Title: Mnemonic introspection in macaques is dependent on dorsolateral prefrontal
cortex but not orbitofrontal cortex

Abbreviated Title: Mnemonic introspection is dlPFC not OFC dependent

Author names:

Sze Chai Kwok [1,2,3], Yudian Cai [1] and Mark J. Buckley [4]

Author affiliations:

[1] *Shanghai Key Laboratory of Brain Functional Genomics, Key Laboratory of Brain*
Functional Genomics Ministry of Education, School of Psychology and Cognitive Science, East
China Normal University, Shanghai 200062, China
[2] *Shanghai Key Laboratory of Magnetic Resonance, East China Normal University, Shanghai*
200062, China
[3] *NYU-ECNU Institute of Brain and Cognitive Science at NYU Shanghai, Shanghai 200062,*
China
[4] *Department of Experimental Psychology, University of Oxford, Oxford OX1 3SR, United*
Kingdom

Corresponding author's email address: sze-chai.kwok@st-hughs.oxon.org

Number of pages in ms: 37; Number of figures: 7; Number of tables: 0

Number of words in Abstract = 245; Number of words in Introduction = 634; Number of words
in Discussion = 1553.

Conflict of Interest Statement: The authors are not aware of any conflicts of interest.

Acknowledgements: This work is supported by the Ministry of Education of PRC Humanities
and Social Sciences Research grant 16YJC190006, STCSM Natural Science Foundation of
Shanghai 16ZR1410200, Fundamental Research Funds for the Central Universities
2018ECNU-HWFW007, NYU Shanghai and NYU-ECNU Institute of Brain and Cognitive
Science at NYU Shanghai (S.C.K.); and the U.K. Medical Research Council (MRC) project
grants G0300817 and MR/K005480/1 (M.J.B.). We acknowledge the support from Dr Keiji
Tanaka and Dr Farshad Mansouri in the Laboratory for Cognitive Brain Mapping at RIKEN
Brain Science Institute, Japan, who helped enable a part of the project.

ABSTRACT

The human prefrontal cortex (PFC) has been associated more with meta-perceptual as

opposed to meta-memory decisions from correlational neuroimaging investigations. Recently,

metacognitive abilities have also been shown to be causally dependent upon anterior and

dorsal PFC in non-human primate lesion studies. Two studies, utilizing post-decision

wagering paradigms and reversible inactivation, challenged this meta-perceptual versus

meta-memory notion and showed that dorsal and anterior prefrontal areas are associated with

metamemory for experienced objects and awareness of ignorance respectively. Causal

investigations are important but scarce; nothing is known, for example, about the causal

contributions of prefrontal sub-regions to spatial metamemory. Here, we investigated the

effects of dorsal versus ventral PFC lesions on two-alternative forced choice spatial

discrimination tasks in male macaque monkeys. Importantly, we were rigorous in approach

and applied three independent but complementary indices used to quantify individual animals'

metacognitive ability ("type II sensitivity") namely *meta-d′, d′* measures, and Phi coefficient

(Φ). Our results were consistent across indices: while neither lesions to superior dorsolateral

PFC (sdlPFC) nor orbitofrontal cortex (OFC) impaired spatial recognition performance, only

monkeys with sdlPFC lesions were impaired in meta-accuracy. Together with the observation

that the same OFC lesioned monkeys were impaired in updating rule-value in a Wisconsin

Card Sorting Test analog, we therefore document a functional double-dissociation between

these two PFC regions. Out study presents important causal evidence that other dimensions,

namely domain-specific processing (e.g., spatial versus non-spatial metamemory), also need

considerations in understanding the functional specialization in the neural underpinnings of

introspection.

Significance Statement

This study demonstrates macaque monkeys' meta-cognitive capability of introspecting its own memory success is causally dependent on intact superior dorsolateral prefrontal cortices (PFC) but not the orbitofrontal cortices. Combining neurosurgical techniques on monkeys and state-of-the-art measures of metacognition, we affirm a critical role of the PFC in supporting spatial meta-recognition memory and delineate functional specificity within PFC for distinct elements of metacognition.

INTRODUCTION

Metacognition refers to awareness of one's own cognition (e.g., knowledge of one's

accuracy, knowing what one knows, or indeed knowing when one does not know). Human

neuroimaging has generally associated prefrontal cortex (PFC) more with meta-judgements

based on perceptual as opposed to memory decisions (Fleming et al., 2012; Morales et al.,

2018), backed up by structural neuroimaging measures (Fleming et al., 2010), and

neuropsychology (Rounis et al., 2010; Shekhar and Rahnev, 2018). Nonetheless, two recent

studies showed that neural activations in dorsal PFC and anterior PFC in the macaque brain are

associated with metamemory of experienced object recognition (Miyamoto et al., 2017) and

awareness of ignorance of experience of objects respectively (Miyamoto et al., 2018),

implying a more complex functional architecture. Indeed, pharmacological intervention

delineated three subareas supporting metamemory, one for temporally remote items in dorsal

area 9 (or 9/46d), a more posterior one for more recent items in area 6, and a third for awareness

of ignorance in the most anterior part of PFC, namely area 10 (frontopolar cortex).

Recent human neuroimaging associates frontopolar cortex with metacognitive control

processes whereas other metacognitive processes underlying decision-making per se are

associated with dissociable neural systems more posteriorly within PFC (Qiu et al., 2018). The

orbitofrontal cortex (OFC) is one such hub associated with value-based decision-making

(Buckley et al., 2009; Noonan et al., 2010; Baltz et al., 2018), inferring the consequences of

potential behavior (Schuck et al., 2016), and decision confidence (Kepecs et al., 2008), such

that lesions to the OFC might affect decision confidence without affecting first-order task

performance (Lak et al., 2014). Since confidence estimation is a fundamental component of

decision-making, and the OFC has been implicated in goal directed decisions that require the

evaluation of predicted outcomes (Rudebeck and Murray, 2014), OFC might be causally

required to support the computation of some dissociable elements of metacognition (Kepecs et

al., 2008; Lak et al., 2014). Moreover, connections differ, for example, posterior parietal areas

of the brain involved in egocentric spatial processing have more robust connections with dorsal

PFC whereas temporal lobe areas implicated in object-identity processing have more robust

connection with ventral PFC (Yeterian et al., 2012). In light of the above and findings that

second-order metacognitive processes could be separated from confidence per se (Dotan et al.,

2018), we investigated the causal roles of one dorsal (sdlPFC) and one ventral (OFC) subregion

of PFC in spatial recognition memory and hypothesized that the sdlPFC but not OFC will be

causally required for accurate spatial memory introspection. Specifically, we contrasted the

first-order memory and second-order metamemory performances of macaques with superior

dorsolateral PFC (sdlPFC) lesion (i.e., lateral area 9) (n=3), or with OFC-lesioned monkeys

(n=3), to unoperated controls (n=7) (FIG. 1) in two delayed-matching-to-position spatial

recognition tasks (FIG. 2). On the basis of a wide-ranging PFC lesion study literature review in

macaque monkeys, we expected that neither lesion would impair first-order spatial recognition

per se, thereby making any changes on second-order recognition easier to discern and interpret.

Cognizant of confidence reporting not being available in our task-design, we employed

response time as a proxy (Dotan et al., 2018) for confidence but as this is a rough proxy we

accordingly adopted a rigorous analytical approach and applied three quite different but

complementary indices used to quantify individual animals' metacognitive ability ("type II

sensitivity") namely *meta-d',* and *d'* measures, and Phi coefficient (Φ). We found that sdlPFC

but not OFC lesioned monkeys were impaired in meta-accuracy in a high spatial memory

demand task variant without showing any impairments in spatial recognition performance itself.

We further established that these putative metacognitive deficits were specific to spatial

recognition memory rather than to other confounds such as rule-learning, reward evaluation, or

general representation of task information using analyses of existing data from the same

sdlPFC-lesioned monkeys when previously tested on a Wisconsin Card Sorting Test analog

(Buckley et al., 2009).

RESULTS

Meta-deficits in dlPFC-lesioned group in spatial recognition

Consistently with the two meta-indices, we revealed a significant main effect of "Group" in the spatial-variant task with a one-way ANOVA on SDT *meta-d'/d'*: $F(2, 9) = 5.464$, $P = 0.028$, $\eta^2 = 0.548$, and given our specific predictions for an impairment in the sdlPFC group we ran post-hoc tests for the comparisons: CON vs. sdlPFC, Dunnet $P = 0.034$, CON vs. OFC, $P = 0.968$ one-tailed; and on hierarchical-model *meta-d'/d'*: $F(2, 9) = 6.524$, $P = 0.018$, $\eta^2 = 0.594$, post hoc test: CON vs. sdlPFC, Dunnet $P = 0.020$, CON vs. OFC, $P = 0.964$ one-tailed.

Given the relatively small sample size, we additionally ran non-parametric tests and revealed the same pattern for both SDT *meta-d'/d'*, Kruskal-Wallis test across monkeys in three groups; $\chi^2(2) = 5.154$, $P = 0.076$, Dunn's post hoc test: CON vs. sdlPFC, $P = 0.075$, CON vs. OFC, $P = 0.120$; for hierarchical-model *meta-d'/d'*, Kruskal-Wallis test across monkeys; $\chi^2(2) = 6.436$, $P = 0.040$, Dunn's post hoc test: CON vs. sdlPFC, $P = 0.034$, CON vs. OFC, $P = 0.148$. The sdlPFC monkeys were impaired in meta-accuracy in spatially demanding recognition, whereas the OFC group did not show any meta-deficit in either of the tasks. In contrast, in the temporal-variant task, we found no main effect of Group in meta-accuracy, *meta-d'/d'*: $F(2, 10) = 0.773$, $P = 0.487$, $\eta^2 = 0.134$, Kruskal-Wallis test across monkeys; $\chi^2(2) = 0.195$, P = 0.907; hierarchical-model *meta-d'/d'*: $F(2, 10) = 0.500$, $P = 0.621$, $\eta^2 = 0.141$, Kruskal-Wallis test across monkeys; $\chi^2(2) = 1.92$, $P = 0.902$.

To add credibility to these results, we replicated these findings with the Phi coefficient (Φ) in both tasks, that is in the spatial-variant, Group effect: $F(2, 9) = 4.904$, $P = 0.036$, $\eta^2 = 0.521$, post hoc test: CON vs. sdlPFC, Dunnet $P = 0.026$, CON vs. OFC, $P = 0.904$ one-tailed, and in the temporal-variant, Group effect: $F(2, 10) = 0.822$, $P = 0.467$, $\eta^2 = 0.091$. The non-parametric tests confirm the same pattern for the spatial-variant, Kruskal-Wallis test across monkeys in three groups; $\chi^2(2) = 6.064$, $P = 0.048$; Dunn's post hoc test: CON vs. sdlPFC, $P = 0.029$,

CON vs. OFC, $P = 0.198$, but not the temporal-variant, Kruskal-Wallis test across monkeys;

$\chi^2(2) = 1.513$, $P = 0.469$. These results using three indices convergently revealed severe

impairment in metamemory of recognition in the sdlPFC lesioned monkeys (but not in OFC

group), confirming that metacognitive ability was impaired in the spatially-demanding spatial

recognition task, but not the temporally-demanding task (FIG. 3A-C).

These meta-indices in principle refer to how meaningful a subject's confidence is in

distinguishing between correct and incorrect responses. We accordingly ran two mixed-design

repeated-measures ANOVAs on percentage correct with "Group" as a between-subject

variable and "Confidence" as a within-subjects variable for the two tasks separately and

obtained a significant interaction with the spatial-variant task $F(2, 9) = 5.416$, $P = 0.029$, but

not with the temporal-variant task $F(2, 10) = 0.355$, $P = 0.710$. Percentage correct in

high-confidence trials is usually higher than low-confidence trials, $P < 0.01$ for both CON and

OFC monkeys in both tasks, but such effects were disrupted in the sdlPFC monkeys in the

spatial-variant task $P = 0.696$, indicating the sdlPFC monkeys were unable to keep track of the

efficacy of confidence during memory judgement (FIG. 3D-E). Correspondingly, one-way

ANOVAs having "Group" as a between-subjects variable on *meta-d'* (a sensitivity measure

quantifying the ability to discriminate between correct and incorrect judgments) for the two

tasks separately also revealed that a significant main effect of "Group" in the spatial-variant

task on SDT *meta-d'*: $F(2, 9) = 5.701$, $P = 0.025$, $\eta^2 = 0.559$, post hoc test: CON vs. sdlPFC,

Dunnet $P = 0.015$, CON vs. OFC, $P = 0.867$ one-tailed, but not in the temporal-variant $F(2, 10)$

$= 0.558$, $P = 0.589$, $\eta^2 = 0.100$. Given the relatively small sample per group, we additionally

ran the statistics using nonparametric tests. These again revealed a significant group effect in

the spatial-variant, Kruskal-Wallis test across all monkeys; $\chi^2(2) = 5.154$, $P = 0.025$, Dunn's

post hoc test: CON vs. sdlPFC, $P = 0.025$, CON vs. OFC, $P = 0.257$, but not in the

temporal-variant, $\chi^2(2) = 1.438$, $P = 0.487$.

In order to ascertain that these lesion effects were task-specific (Task:

spatial-variant/temporal-variant), we ran three separate mixed-design repeated-measures

ANOVAs considering only the CON and sdlPFC groups with "Group" as a between-subjects

variable and "Task" as a within-subjects variable and confirmed a marginally significant "Task

$\times$ Group" interaction on SDT *meta-d'/d'*: $F(1, 7) = 4.194$, $P = 0.080$ and on hierarchical-model

meta-d'/d': $F(1, 7) = 4.599$, $P = 0.069$, as well as a slightly weaker effect on Phi coefficient (Φ):

$F(1, 7) = 1.939$, $P = 0.206$.

sdlPFC and OFC lesions did not result in recognition impairment

Given that metacognition is quantified by the correspondence between confidence and

type I task performance, it is theoretically important to establish that the task (first-order)

performances were matched between the groups in order to argue for the presence of a true

difference in metacognition caused by the sdlPFC lesion. Despite the deficits in metamemory

accuracy in the sdlPFC group, importantly, we established that there were not any memory

deficits in their type I performance. In two mixed-design repeat-measures ANOVAs we entered

the percentage correct or RT with one between-subjects factor "Group" and one within-subjects

factor "Condition" and found neither a main effect nor interaction effects with "Group" in the

two tasks, temporal-variant: % correct: $F(2, 10) = 0.284$, $P = 0.759$, RT: $F(2, 10) = 0.932$, $P =$

0.425, no "Group $\times$ Condition" interaction all Ps > 0.05; spatial-variant: % correct: $F(2, 9) =$

3.868 $P = 0.061$, RT: $F(2, 9) = 0.794$, $P = 0.481$, no "Group $\times$ Condition" interactions all Ps >

0.05. The ANOVAs also showed that performance decreased with Delay, % correct: $F(4, 40) =$

26.964, $P < 0.001$, RT: $F(4, 40) = 9.918$, $P < 0.001$, and with Separation, % correct: $F(3, 27) =$

33.960, $P < 0.001$, RT: $F(3, 27) = 0.121$ $P = 0.105$ for all three groups (FIG. 4). These analyses

point to the fact that neither sdlPFC nor OFC lesion resulted in any type 1 recognition memory

impairment.

Meta-deficits could not be explained away by speed-accuracy trade off

Here we have utilized reaction time as a proxy for memory decision confidence (Dotan et

al., 2018). The meta-deficit effects observed here might be confounded by some

speed-and-accuracy trade-off strategy adopted by the monkeys towards maximizing the

time-spent per unit of reward (correct) ratio. Speed-and-accuracy trade-off taps into the

monitoring of the current state of mind as regards the uncertainty properties of the judgement

(Bogacz et al., 2010), whereas RT-indexed metacognition – defined as an introspective

evaluation process – taps into the higher-level function. We thus analyzed the ratio between

percentage correct and RT for each monkey for both tasks. In two mixed-design

repeat-measures ANOVAs we entered the inverse efficiency (RT / % correct) with one

between-subjects factor "Group" and one within-subjects factor "Condition". We found neither

a main effect nor interaction effects with "Group" in the two tasks, all Ps > 0.05 (FIG. 5). We

conclude that the putative lesion-related meta-deficits were not resultant from any

speed-and-accuracy trade off.

No meta-memory deficit following sdlPFC lesion in short-term abstract rule memory in

Wisconsin Card Sorting Test (WCST)

While the sdlPFC monkeys were impaired in metamemory for spatial recognition, we have

not been able to ascribe the effects specifically to spatial recognition per se. Is this deficit

uniquely ascribable to the metamemory in temporo-spatial recognition, or more generally to the

metamemory of learning abstract rules, or other higher cognitive processes? Considering

performance supporting WCST demands multi-processes such as memory and acquisition of

abstract rules, as well as reward-value evaluation, we thus analyzed some extant data obtained

from WCST to test for metacognitive deficits specifically in the sdlPFC monkeys. In contrast to

the spatial recognition task, no meta-deficits were found with WCST in the sdlPFC group in

comparison with the CON group in one-way ANOVAs on SDT *meta-d'/d'*: $F(1, 10) = 0.677$, P

$= 0.430$, $\eta^2 = 0.063$, Kruskal-Wallis test across monkeys; $\chi^2(2) = 0.419$, $P = 0.518$;

hierarchical-model *meta-d'/d'*: $F(1, 10) = 0.018$, $P = 0.896$, $\eta^2 = 0.002$, Kruskal-Wallis test

across monkeys; $\chi^2(2) = 0.077$, $P = 0.782$; and Phi coefficient (Φ): $F(1, 10) = 1.132$, $P =$

0.312, $\eta^2 = 0.102$, Kruskal-Wallis test across monkeys; $\chi^2(2) = 0.009$, $P = 0.926$ (FIG. 6).

These analyses confirm that meta-deficits caused by sdlPFC lesion were highly specific for

spatial recognition and ruled out the explanation that such meta-deficits were attributable to

processes involved in the maintenance of abstract rules or general representation of knowledge.

DISCUSSION

As expected, neither sdlPFC nor OFC aspirative lesions impaired first-order spatial

recognition memory performance; yet importantly, and consistent with our hypotheses, sdlPFC

lesions selectively impaired second-order meta-recognition within a recognition memory

paradigm taxing recent spatial memory. No such change in metacognitive ability was observed

after OFC lesions both affirming a critical functional role of the sdlPFC in supporting spatial

metamemory and showing evidence for functional specificity within prefrontal cortex (PFC)

for elements of metacognition. Our findings are robust because we assessed multiple measures

of metamemory (both SDT and hierarchical model *meta-d'/d'* and Phi coefficient (Φ)) and

found consistent significance across measures. Our findings are important because they provide

causal evidence, of which there is currently very little, towards refining our understanding of

functional specialization within primate PFC underpinning introspection during memory

recognition.

Critical role of dlPFC in meta-recognition memory

Whilst the role of dlPFC in meta-evaluation of visual perception has been relatively well

established, by evidence observed in functional activation during post-decision evaluation

(Ochsner et al., 2005; Wan et al., 2016), structural and connectivity profiles (Fleming et al.,

2010; Fleming and Dolan, 2012), and neuromodulation studies (Meiron and Lavidor, 2014),

the neural basis of meta-recognition memory within this region remains largely unknown.

Previously, combined lesions to mid-dlPFC lesion (areas 46 and 9/46) and superior part of the

mid-dlPFC (lateral area 9) were found not to affect standard first-order recognition memory

maintenance for recently presented objects (in contrast to lesions to ventrolateral PFC and OFC

which do impair first-order object recognition (Bachevalier and Mishkin, 1986; Kowalska et al.,

1991)). The mid dlPFC lesions did however severely impaired executive processes of

monitoring visual working memory information in a self-ordered version of the task (Petrides,

2000); moreover, more restricted lesions within lateral area 9 (similar to our sdlPFC lesions)

were sufficient to impair the self-ordered task. Together with findings that patients with more

diffuse frontal lobe pathology exhibit impaired feeling-of-knowing in the absence of amnesia

(Janowsky et al., 1989), our pattern of results corroborate the extant evidence that a dissociation

between type 1 vs. type 2 performance in recognition is associated with dlPFC damage. This

dissociation also aligns with the recent distinction stipulating the dlPFC's putative role in

metacognitive control as opposed to decision-making per se (Qiu et al., 2018).

Importantly, and in light of recent evidence from the macaque showing that there exist

dissociated networks within PFC underlying metacognition (Miyamoto et al., 2017)(Miyamoto

et al, 2018), our results significantly extend the causal evidence for understanding the

functional neuroanatomy of metacognition in PFC in several key ways. Firstly, we extend

causal evidence for metacognition into the spatial domain. Secondly, we dissociate the

functional neuroanatomy of first-order spatial recognition memory from second-order spatial

meta-recognition within PFC. Circumscribed muscimol injections to different localities within

mid-dlPFC area impair visually-guided saccades in a visuospatial working memory task in a

topographical manner (Sawaguchi and Iba, 2001); further evidence that this dlPFC region but

not the more superior dlPFC region impairs first-order spatial recognition comes from previous

surgical lesions in monkeys (Levy and Goldman-Rakic, 1999). Our sdlPFC lesion fails to

encroach much on this first-order region and accordingly do not impair first-order spatial

recognition; at the same time our sdlPFC lesions impair meta-recognition in the same task,

therefore demonstrating functional dissociation between first-order and second-order spatial

recognition within PFC. Taken together with Miyamoto et al. (2017; 2018), our study confirms

that sdlPFC neither contributes exclusively to object nor spatial meta-recognition, rather it

contributes to both. Interestingly we found dissociation between meta-recognition deficits after

sdlPFC lesions in a spatially demanding but not temporally demanding version of our task. The

main difference between these two versions is that in the former the spatial difficulty is

intentionally varied between trials; given it is a spatial recognition task we postulate efficacious

metacognitive monitoring of performance will therefore be in flux and continually challenged

necessitating a dynamic signal from sdlPFC input into the wider neural system in support of

metacognitive computation. By contrast, in the temporal-variant the spatial difficulty is

constant across trials so any metacognitive evaluation signal from sdlPFC would likely be a

less important parameter to the system and either absent or less liable to be disrupted by the

sdlPFC lesion accordingly, in accordance with our behavioural observations.

OFC represents value/confidence but does not supports self-introspection

At first glance, OFC is an interpreter of specific values especially in terms of sorting and

representation of inferred information (Wallis, 2007; Jones et al., 2012); could it be the

cornerstone of meta-appraisal towards one's own memory performance? Despite proposals that

the OFC is a key part of continuous decision-making under uncertainty (Kennerley et al., 2011),

related to the explicit manifestation of decision confidence (Kepecs et al., 2008; Lak et al., 2014)

and various aspects of decision-making (Izquierdo, 2017), its role in metacognition in the

present experimental context appears to be none as evident in the total absence of

meta-impairment in the OFC group. One likely possibility is that value-assignment

(Padoa-Schioppa and Assad, 2006), valuation of inferred information (Jones et al., 2012), and

decision-confidence (Lak et al., 2014) per se are fundamentally distinct/dissociable from

metamemory introspection. Indeed, meta-decision processes – as measured by the SDT and

hierarchical-model *meta-d'* here – are in principle "bias-free" and are immune to any bias due to

"confidence" (Fleming and Lau, 2014), suggesting that the computation performed by the OFC

underlying confidence signals (Kepecs et al., 2008; Lak et al., 2014) does not necessarily

equate to the same neurobiological prerequisite for meta-cognitive computation. Relatedly, in

our tasks there was no explicit requirement for reporting confidence, in which case the memory

response need not to be bound with any explicit value valuation processes (Rolls and

Grabenhorst, 2008) or reward-based updating (Buckley et al., 2009). The introspection

following memory decision was thus based entirely on some self-generated space, without any

feedback or input exerted externally on their decision confidence or monitoring of degree of

uncertainty (Kennerley et al., 2011). This task feature discrepancy might explain the lack of

meta-awareness deficits even when the OFC was absent.

An alternative but not mutually exclusive explanation, given that frontal vs. parietal cortex

neural correlates of metacognition are somewhat domain-specific (McCurdy et al., 2013), is

that OFC's contribution to metacognition might analogously be domain-specific. Indeed,

previous studies tapping into the OFC role in meta-related processes were all on perceptual

decisions, such as odor discrimination judgement (Kepecs et al., 2008; Lak et al., 2014),

whereas at present the tasks in question concern mnemonic decisions; some recent work on

humans has similarly evinced such specificity for perceptual vs. memory related metacognition

(Fleming et al., 2014; Ye et al., 2018).

Applications and limitations of reaction-time (RT) approach

We categorized trials into fast-response trials (high confidence) and slow-response trials

(low confidence) according to the median of RT (Dotan et al., 2018). A similar method was

used in humans, suggesting fast-response and slow-response are supported by distinct

mechanisms (Novikov et al., 2017) and processes (van de Vijver et al., 2011; Coomans et al.,

2016). While recognizing that other possibilities exist, for example fast trials and slow trials

might also relate to different levels of attention besides uncertainty (Novikov et al., 2017),

certainty is nonetheless informed by RT (Kiani et al., 2014) and all considered process are

presumably related to self-monitoring. In our approach we focus on four kinds of trials:

correct/high confidence (fast RT), incorrect/low confidence (slow RT), correct/low confidence

(slow RT), incorrect/high confidence (fast RT); accordingly, our analyses may be considered to

be particularly relevant to implicit metacognitive accuracy of the decision-making system. The

advantage of RT-indexed metacognition should not be overlooked too, namely that it does not

suffer from any training-induced associations (such as environmental cue associations,

behavioral cue associations, response competition) (Hampton, 2009), which could contaminate

true introspection.

If, using implicit metacognitive measurement, we cannot assert that the effect is purely

represented by metacognition, then the RT in type I task may indeed be entangled with

metacognitive reaction time, leaving us with a mixed read-out of the directional certainty (Caio

M. Moreira and Kagan., 2018). Considering the performance on type I tasks in sdlPFC lesioned

monkeys did not differ from that in CON monkeys, we infer neither attention (Novikov et al.,

2017) nor motor function are affected by sdlPFC lesions. Moreover, the result of

speed-accuracy trade off analyses confirmed that changed cognitive strategy cannot account for

the deficits in the sdlPFC group in the spatial task.

A wider theoretical implication afforded by the current study is that this same group of

sdlPFC monkeys, despite their impaired memory self-appraisal in the

delayed-matching-to-position task, were completely intact in all aspects of a rule-guided

memory Wisconsin Card Sorting Test analog (Buckley et al., 2009). This constitutes a stark

contrast to the OFC-lesioned monkeys, who were impaired in updating rule-value

representation in the WCST analog (Buckley et al., 2009), but not in their introspection in the

present delayed-matching-to-position tasks. These results taken together constitute a double

dissociation within PFC between dorsolateral and ventromedial PFC regions in differentially

supporting two related, yet perhaps distinct, higher-order processes, providing compelling

evidence suggestive of functional specialization of dual supervisory, self-monitoring abilities

between dorsolateral vs. ventral parts of the PFC.

MATERIALS AND METHODS

Animals. Data were acquired from 16 adult macaque monkeys (eleven *Macaca mulatta*, three

Macaca fuscata, and two *Macaca fascicularis*): Three monkeys had orbitofrontal lesion (OFC,

consisting of two *M. mulatta* and one *M. fuscata*), another 3 monkeys had superior dorsolateral

prefrontal cortex lesion (sdlPFC, consisting of one *M. mulatta* and two *M. fuscata*), 7 served as

unoperated controls (CON, consisting of five *M. mulatta* and two *M. fascicularis*) for the main

tasks, and 3 further CON were included only for the WCST analog analysis. All but six

macaque monkeys were trained, operated and tested in Oxford, UK, the other six (three

Macaca fuscata and three *Macaca mulatta*) in RIKEN Brain Science Institute, Wako, Japan.

All animal training, surgery and experimental procedures were the same in both laboratories.

Those conducted in the UK were licensed in compliance with the UK Animals (Scientific

Procedures) Act 1986, and those in Japan were done in accordance with the guidelines of the

Japanese Physiological Society and approved by RIKEN's Animal Experiment Committee.

Surgery. The operations were performed in sterile conditions (under gaseous general

anesthesia, and with pre-, peri-, and post-operative analgesia) with the aid of an operating

microscope and the same surgeon performed all operations in both laboratories. Detailed

description of the surgical procedures has been reported previously (Buckley et al., 2009).

The intended extent of the sdlPFC lesion was designed to include the cortex on the

dorsolateral aspect of the PFC extending up to midline (i.e., lateral area 9 and the dorsal

portions of areas 46 and 9/46) but excluding ventrally situated dlPFC cortex; the lesion

excluded posteriorly located premotor areas 8A, 8Bd, and 8Bv, nor did it extend anteriorly into

area 10.

The intended extent of the OFC lesion included at its lateral extent, the cortex in the medial

bank of the lateral orbital sulcus; the lesion included all of the cortex between the medial and

lateral orbital sulci, and also extended medially until the lateral bank of the rostral sulcus. The

anterior extent of the lesion was an imaginary line drawn between the anterior tips of the lateral

and medial orbital sulci, and the posterior extent was an imaginary line drawn just anterior to

the posterior tips of these two sulci. The intended lesion therefore included areas 11, 13 and 14

of the orbital surface and did not extend posteriorly into the agranular insula.

Histology. After the conclusion of the experiments the animals with ablations were sedated,

deeply anesthetized, and then perfused through the heart with saline solution (0.9%), which

was followed by formol saline solution (10% formalin in 0.9% saline solution). The brains

were blocked in the coronal stereotaxic plane posterior to the lunate sulcus, removed from the

skull, allowed to sink in sucrose formalin solution (30% sucrose,10% formalin), and sectioned

coronally at 50μm on a freezing 10 microtomes. Every 10^{th} section through the temporal lobe

was stained with cresyl violet and mounted. When referring to cytoarchiteturally defined

regions in the lesion description below we have adopted the nomenclature and conventions of

Petrides and Pandya (Petrides and Pandya, 1999) and have reconstructed lesion extents on

standard drawings based upon those provided by the Laboratory of Neuropsychology at NIMH.

All three of the sdlPFC lesions were as intended. None of the OFC lesioned animals

sustained any bilateral damage outside the area of the intended region; two animals sustained

extremely slight unilateral damage beyond the intended lateral boundary of the lesion OFC2

and OFC3; in all three animals the lesions did not extend as far medially as intended. Detailed

photomicrographs of stained coronal sections through the intended sdlPFC and OFC lesions for

all animals and reconstructions of precise lesion extents in individual animals have been

published previously (Buckley et al, 2009); as the lesions were largely as intended, FIG.2

illustrates summary drawings of the intended extent of the lesions.

Metamemory quantification. Three different but complimentary indices were used to

quantify individual animals' metacognitive ability ("type II sensitivity"), as defined as the

ability to accurately link confidence with performance. Here, we calculated the *meta-d′/d′*, a

metric for estimating the metacognitive efficiency (level of metacognition given a particular

level of performance or signal processing capacity), which enables a model-based approach to

the computation of type II sensitivity that is independent of response bias and type I sensitivity

(*d′*) on the primary task (Rounis et al., 2010). We also computed metacognitive efficiency using

a hierarchical Bayesian estimation method, which can avoid edge-correction confounds and

enhance statistical power (Fleming and Daw, 2017). Both *meta-d′* and *d′* measures assume that

the variance of the internal response takes a Gaussian distribution, and that the distributions

associated with the two type 1 responses respectively are of equal variance. To ensure our

results were not due to any idiosyncratic violation of the assumptions of signal detection theory

(SDT), we additionally calculated the Phi coefficient (Φ), which does not make these

parametric assumptions (Fleming and Lau, 2014), and supplemented the statistical analyses

using nonparametric tests (Kruskal-Wallis). The correlations among the three meta-cognitive

metrics are very high (FIG. 7).

Signal detection theoretic and hierarchical Bayesian estimation meta-index (meta-d′/d′): Using

a Type II SDT toolbox (Maniscalco and Lau, 2012), which has been extensively used for

evaluation of metacognitive ability (Baird et al., 2013; Fleming et al., 2014), it is possible to

compute a measure of metacognitive accuracy that is unconfounded by type I performance

directly from the empirical type II receiver operating characteristic (ROC) curve. The type II

ROC curve reflects the relationship between the accuracy of type I and the observer's

confidence rating. This approach exploits the link between type I and type II SDT models to

express observed type II sensitivity at the level of the type I SDT model (termed *meta-d′*).

Maximum likelihood estimation is used to determine the parameter values of the type I SDT

model that provide the best fit to the observed type II data. A measure of metacognitive ability

that controls for differences in type I sensitivity is then calculated by taking the ratio of *meta-d'*

and the type I sensitivity parameter *d'*: meta efficiency, computed as *meta-d'/d'*. The most

straightforward approach to computing meta efficiency involves an equal variance SDT model

in which the variances of internal distributions of evidence for "target" and "foil" in the type I

model are assumed to be equal. We thus quantified metacognitive sensitivity with the

SDT-based measure *meta-d'*. Based on type II signal detection theory, meta-efficiency (in

terms of *meta-d'/d'*) reflects how much information, in signal-to-noise units, provides a

response-bias free measure of how well confidence ratings track task accuracy. The toolbox for

the SDT-based *meta-d'/d'* estimation was available at

http://www.columbia.edu/~bsm2105/type2sdt/. Of note, the standard type II SDT toolbox is

designed for 2AFC tasks, in which S1 and S2 are always constant in the left or right, but our

target and foil are randomly presented at the screen and we only recorded the separation of the

two probes and did not track the specific position of the target and the foil. Since the target and

the foil were presented randomly on the screen, and the SDT algorithm only requires the

distribution of those four kinds of trials, we divided the number of those trials equally to S1

trials and S2 trials in a random manner considering the animals would not have any preference

to any given side/location of the screen. In addition, we have also replicated the analyses using

a variant of metacognitive efficiency (H-*meta-d'*) with a hierarchical Bayesian estimation

method (https://github.com/smfleming/HMeta-d), which can avoid edge-correction confounds

and enhance statistical power (Fleming and Daw, 2017).

Phi coefficient (Φ): In order to ensure our results were not due to any idiosyncratic violation of

the parametric assumptions of SDT, we additionally calculated the phi coefficient index, which

does not make the SDT assumptions. The phi coefficient is a contingency index of preference

for optimal choice (Kornell et al., 2007; Middlebrooks and Sommer, 2011) and was calculated

according to the following formula using the number of trials classified in each case [n(case)]:

$$\text{phi coefficient } (\Phi) = \frac{n(Correct\ High) \times n(Incorrect\ Low) - n(Correct\ Low) \times n(Incorrect\ High)}{\sqrt{n(Correct) \times n(Incorrect) \times n(High) \times n(Low)}}$$

.

The Φ coefficient evaluates how optimally each trial was assigned for high or low confidence

based on performance in the preceding cognitive judgment, reflecting the correlation between

the two binary variables. Note that despite differences in their mathematical assumptions, the

three meta-cognitive metrics are highly correlated with each other (FIG. S1).

For the computation for SDT *meta-d'/d'*, hierarchical-model *meta-d'/d'*, and Phi coefficient,

four types of trials and their distribution are required. The computation performed here are

based on the premise that confidence is computed in a retrospective manner (Pleskac and

Busemeyer, 2010). Using a summary of the decision process, trials that are responded fast are

judged as more of higher certainty (Kiani et al., 2014). Following this logic, we accordingly

used trial-specific reaction times (RT) as a proxy for confidence (Dotan et al., 2018). We

collapsed all trials per monkey and classified the trials within-monkeys by the median of all RT

crossing with correct/incorrect responses into four kinds: correct/high confidence (fast RT),

incorrect/low confidence (slow RT), correct/low confidence (slow RT), incorrect/high

confidence (fast RT). For each of the final analyses, each monkey had one single value for the

measurement of meta-ability. It should be noted that all the statistical analyses on

meta-cognition were performed within-subjects and thus the numerally (but not statistically

significant) faster RT for the sdlPFC monkeys would not affect our main findings based on

confidence for trial classification.

Behavioral tasks and pre-analysis. *Spatial recognition tasks (delayed-matching-to-position,*

or DMP). A temporally-demanding DMP task and a spatially-demanding DMP task were

performed by the monkeys. In both variants, each trial consisted of an encoding phase in which

a spatial position ("sample") was indicated by a red cross. After the monkey touched the sample,

a blue square ("distractor") appeared in the center of an imaginary (i.e., invisible) circle whose

circumference transected the centre of the red cross. A touch to the blue square initiated a

variable delay interval (i.e., the manipulation of delay for the temporal-variant) and then a

choice phase consisting of two identical red crosses in different positions, both located on the

circumference of the aforementioned imaginary circle (hence equidistant from the blue square

distractor just touched), albeit one positioned in the same (i.e., spatial-match) position as the

first red cross and the other (i.e., non-match) positioned some angle (with respect to the centre

of the imaginary circle) away from the spatial-match along the invisible circumference. From

trial to trial we could vary the angle of separation along the circumference to allow for easy

trials (i.e., large angle and accordingly large spatial separation) and harder trials (i.e., smaller

angles and accordingly smaller spatial separation) (cf. the manipulation of separation between

two probes for the spatial-variant). As mentioned above, one of the crosses appeared in the

same position as the sample (target; S+) and the other one in a different position (foil; S-). A

touch to the S+ resulted in a delivery of a reward pellet as reward (see section on Apparatus

below), removed the S-, and the S+ remained alone for a further 1 s for positive feedback. The

screen would then be blanked for an ITI of 6 s before the next trial. A touch to the S- removed

both S+ and S- from the screen and the screen would be blanked for an ITI of 12 s. There was

no time constraint imposed on responses made to the choices and therefore there were no

missed trials. No repetition correction routines were implemented following an error response;

each trial was new and independent of the outcome of the preceding trial. In terms of sizes of

visual stimuli, the sample subtended a visual angle of 9 □ in task acquisition and the

temporal-variant task, or 6.8 □ in the spatial-variant task; the distractor subtended a visual

angle of 4.6 □ in all tasks (FIG. 2).

In the temporal-variant DMP, there were five trial-types with differing delay intervals (either 1,

2, 4, 8, or 16 s) between the distractor and probes. Trials within a session were divided into five

trial-types with differing intervals of delay between the distractor and probes. The trial-type

order was randomized within each successive set of five trials (with one trial of each trial-type

per set) so that the delay changed unpredictably from one trial to another. The two probe

choices were separated by a visual angle of 21.7 □. In the spatial-variant DMP, the separation

between two red crosses varied across trials; there were four different trial-types with differing

spatial separations (visual angles of either 4.8°, 8.6°, 15.2°, or 21.7°; equivalent to 5, 9, 16, or

23 cm on screen) between probe choices. Delays were fixed at 1 s for the spatial-variant DMP.

In the final testing, each animal accrued 200 rewards to complete the temporal-variant (across

two daily sessions) and 150 rewards (one session) to complete the spatial-variant. Since the

animals accrued varying numbers of errors to complete the tasks, the mean total numbers of

trials were 271.2 and 209.8 trials (averaged across groups) respectively for the two tasks. The

analysis on metamemory was done by collapsing all trials across sessions and thus the learning

trend of these data was not considered. One CON did not complete the spatial-variant so only

12 monkeys were analyzed in the spatial-variant task.

For the formal analyses, trials with RT longer than 20 s and shorter than 100 ms in the memory

judgement were discarded (< 0.5% of all trials). We also set a stringent selection of good trials

on which the monkeys were attentive is crucial for the metacognition analysis. To this end, we

set a requirement for the monkeys to touch the distractor before the memory task so as to ensure

they were not distracted and/or less willing/ready prior to initiating the memory judgement.

Trials with touch-distractor times longer than 1000 ms (15.0% and 20.2% trials discarded

respectively for temporal-variant task and spatial-variant task) were not included for analysis.

There were no differences in touch-distractor times between the groups in either of the tasks

following this trial removal procedure, with a one-way ANOVA on temporal-variant: $F(2, 10)$

$= 1.27, P = 0.322$; and on spatial-variant: $F(2, 9) = 0.800, P = 0.479$. The results did not differ

if we chose to use other touch-distractor times cut-off criteria of either 800 ms, 900 ms, 1100

ms, or 1200 ms.

Wisconsin Card Sorting Test (WCST) analog. Given that the DMP tasks involve multiple

processes which might confound our main results, we therefore analyzed extant data obtained

from a WCST analog, which is a validated rule-guided task taxing multi-processes such as

perception (involved in matching stimuli), memory and acquisition of abstract rules, and

reward-value evaluation (Buckley et al., 2009; Mansouri et al., 2015). We accordingly used

some WCST data to rule out that the putative meta-deficits observed here were not attributable

to these perceptual and reward-value evaluation processes. The WCST analog paradigm is

summarized as follows: on each trial, a randomly selected sample (a square, a circle, or a cross

of different colors) is displayed alone on the center of the touch screen, and when the sample is

touched, three additional choice items immediately appear (one matching in color, one

matching in shape, and one not matching in either dimension), with their positions randomly

chosen. If the animal's choice is correct (i.e., the animal selects the choice item that matches

according to the currently reinforced rule, which changes unannounced every time the animal

attains 85% in 20 consecutive trials), then a reward pellet is delivered, and the correct choice

remains on the screen for 1 s to provide visual feedback; if the animal makes an incorrect choice,

then no reward is given, and the stimuli are removed and replaced by an error signal (white

circle), which is presented on the screen for 1 s instead.

We analyzed WCST data from 12 monkey data points (9 CON vs. 3 sdlPFC). Six out of the

nine CON monkey data points here were from the pre-lesion data of the six lesioned monkeys

(3 sdlPFC and 3 OFC). We included 3,000 trials (acquired from ten 300-trial daily sessions) per

monkey data point. We collapsed all trials and classified the trials into four types of trials for

the computation for the meta-efficiency and Phi coefficient. Since the type II SDT toolbox was

designed for 2AFC tasks, and the WCST task contained three stimuli, we ran three separate sets

of computation, each one discarding only either the bottom, left, or right choice, for each of the

three meta-indices. For each monkey, we then computed the mean of these three values as his

meta-score to enter into the meta-indices calculation. In this analysis, we did not include the

OFC monkeys because the OFC monkeys were severely impaired in the WCST type I task, thus

making any analyses on meta-ability invalid (their chance level implies they did not know how

to make correct judgments, violating the prerequisite for the meta-assessment of their

judgement).

Preliminary training. All monkeys completed preliminary training and task acquisition before

performing the two main tasks and WCST described above. We conducted the spatial-variant

task immediately after the temporal-variant task without any additional training. The monkeys

performed one session per day, 6–7 d per week. For the lesioned animals, the task was

administered post-operatively (on average 22 months post-lesion). For the two DMP tasks,

during task acquisition the monkeys were trained until they reached ≥ 90% performance level

within a 100-reward session. All trials in this stage consisted of a short delay interval (1 s), and

a wide separation between choice positions (21.7°, or 23 cm) to make the trials "easy" to

acquire. Upon reaching criterion, the three groups were not different in the number of errors

accrued, $F < 1$ and number of rewards received, $F < 1$, indicating that the groups of lesioned

monkeys learned to perform these spatial recognition problems as well as controls.

Apparatus. The tasks were performed in an automated test apparatus. The subject sat,

unrestrained, in a wheeled transport cage fixed in position in front of a touch-sensitive screen

on which the stimuli could be displayed. The animals could reach out between the horizontal or

vertical bars (spaced approx. 45 mm apart) at the front of the transport cage to touch the screen.

An automated pellet delivery system delivered banana flavored reward pellets (190 mg

supplied by Noyes Company Inc. and Neuroscience Inc.) into a food well (approx. 80 mm in

diameter) positioned beneath and to one side of the screen, in response to correct choices made

by the subject to the touch screen. Pellet delivery was accompanied by an audible click. A

spring-loaded lunchbox (length 200 mm, width 100 mm, height 100 mm) was positioned

beneath and to one side of the subject; this opened immediately with a loud crack on

completion of the testing session and contained the subject's daily diet of wet monkey chow,

primate pellets, nuts, raisins, and a slice of apple, banana, and orange (water was provided in

the home cage *ad libitum*). An infrared camera allowed the subject to be observed while it was

engaged in the task. The entire apparatus was housed in an experimental cubicle that was dark

apart from the background illumination from the touch screen. A computer, with a

millisecond accuracy timer-card to record reaction times, controlled the experiment and data

acquisition. Identical software controlled the tasks in both laboratories to ensure that the tasks

were replicated exactly.

Figure 1. (A) The intended OFC lesion extent shown on a ventral drawing of the macaque
brain (far left) together with reconstructions, from histological stained sections and
photomicrographs, of the actual lesion of the three OFC lesioned animals; (B) The intended
sdlPFC lesion extent shown on lateral drawings of the macaque brain (far left) together with
reconstructions, from histological stained sections and photomicrographs, of the actual lesion
of the three sdlPFC lesioned animals. Detailed photomicrographs of stained coronal sections
through the intended sdlPFC and OFC lesions have already been published previously for each
of the individual animals and are available elsewhere (Buckley et al., 2009).

Figure 2. Delayed-matching-to-position tasks. Each trial consisted of a sample (red cross)

stage, a distractor (blue square) stage, a delay, and then a probe/choice (2 red crosses) stage.

Temporally-taxing DMP: five levels of delay interval between distractor and probes (either 1, 2,

4, 8, or 16 s); spatially-taxing variant DMP: four levels of separation between two red crosses in

probe (visual angles of either 4.8°, 8.6°, 15.2°, or 21.7° which are equivalent to 23, 16, 9, and 5

cm on screen; all delay fixed at 1s). S+ denotes the target, S- the foil. The grey dotted circle in

the figure is invisible to the animal and is just shown to illustrate the two choices are always

equidistant to the distractor to obviate proximity bias.

Figure 3. Differential deficits in meta-indices and accuracy x confidence interaction in

sdlPFC group in spatially-demanding recognition. Meta performance for the 3 monkey

groups (OFC, sdlPFC, CON). Metacognitive accuracy in sdlPFC group was lower than CON

group for spatial-variant task, but not for the temporal-variant task: (A) SDT *meta-d'/d'*, (B)

hierarchical-model *meta-d'/d'*, and (C) Phi coefficient (Φ). Horizontal axes represent the two

spatial recognition tasks (temporal-variant; spatial-variant). Vertical axes represent the three

meta indices. (D – E) Accuracy in high-confidence trials is usually higher than low-confidence

trials (for both CON and OFC monkeys in both tasks) but such effects were disrupted in the

sdlPFC monkeys especially in the spatial-variant task. ⊗ indicates significant "Group x

Confidence" interaction $P < 0.05$, * $P < 0.05$. Colored dots depict individuals. Error bars

indicate 90% confidence interval around the estimate computed by a bootstrapping procedure.

Figure 4. sdlPFC and OFC lesions did not result in recognition impairment. Memory task

performance was intact in both tasks: (A) temporal-variant percentage correct, (B)

temporal-variant reaction time, (C) spatial-variant percentage correct, and (D) spatial-variant

reaction time. Error bars indicate 90% confidence interval around the estimate computed by a

bootstrapping procedure.

Figure 5. Meta-memory deficits could not be explained away by speed-accuracy trade off.
Inverse efficiency (reaction time / percentage correct) shows no main effect of "Group" in (A)
temporal-variant task and (B) spatial-variant task. Error bars indicate 90% confidence interval
around the estimate computed by a bootstrapping procedure.

Figure 6. No meta-memory deficit following dlPFC lesion in WCST analog. Meta

performance for the 2 monkey groups (sdlPFC, CON) in the two spatial recognition tasks

(temporal-variant; spatial-variant) and WCST analog. Vertical axes represent the three meta

indices: (A) SDT *meta-d'/d'*, (B) hierarchical-model *meta-d'/d'* and (C) Phi coefficient (Φ).

Metacognitive accuracy in sdlPFC group was lower than CON group for spatial-variant task

(see also FIG. 3A-C), but not for WCST analog or temporal-variant task. Error bars indicate 90%

confidence interval around the estimate computed by a bootstrapping procedure. * $P < 0.05$.

Colored dots depict individual monkeys.

Figure 7. Strong correlations among the three meta-cognitive metrics. Pearson correlations

computed among the three indices were all statistically significant (all *P*s < 0.001). Colored

dots depict individual data points collapsed across monkey groups, with each monkey shown

twice (temporal- and spatial-variants).

655   REFERENCES

656   Bachevalier J, Mishkin M (1986) Visual recognition impairment follows
657   ventromedial but not dorsolateral prefrontal lesions in monkeys. Behavioural
658   Brain Research 20:249-261.
659   Baird B, Smallwood J, Gorgolewski KJ, Margulies DS (2013) Medial and lateral
660   networks in anterior prefrontal cortex support metacognitive ability for
661   memory and perception. The Journal of Neuroscience 33:16657-16665.
662   Baltz ET, Yalcinbas EA, Renteria R, Gremel CM (2018) Orbital frontal cortex
663   updates state-induced value change for decision-making. eLife 7:e35988.
664   Bogacz R, Wagenmakers E-J, Forstmann BU, Nieuwenhuis S (2010) The neural basis
665   of the speed–accuracy tradeoff. Trends in Neurosciences 33:10-16.
666   Buckley MJ, Mansouri FA, Hoda H, Mahboubi M, Browning PGF, Kwok SC,
667   Phillips A, Tanaka K (2009) Dissociable components of rule-guided behavior
668   depend on distinct medial and prefrontal regions. Science 325:52-58.
669   Caio M. Moreira MRKKMW, Kagan. I (2018) Post-decision wagering after
670   perceptual judgments reveals bi-directional

671   certainty readouts. Cgonition 176:40-52.
672   Coomans F, Hofman A, Brinkhuis M, van der Maas HL, Maris G (2016)
673   Distinguishing Fast and Slow Processes in Accuracy - Response Time Data.
674   PLoS One 11:e0155149.
675   Dotan D, Meyniel F, Dehaene S (2018) On-line confidence monitoring during
676   decision making. Cognition 171:112-121.
677   Fleming SM, Dolan RJ (2012) The neural basis of metacognitive ability. Phil Trans R
678   Soc B 367:1338-1349.
679   Fleming SM, Lau H (2014) How to measure metacognition. Frontiers in Human
680   Neuroscience 8:443.
681   Fleming SM, Daw ND (2017) Self-Evaluation of Decision-Making: A General
682   Bayesian Framework for Metacognitive Computation. Psychological Review
683   124:91-114.
684   Fleming SM, Huijgen J, Dolan RJ (2012) Prefrontal contributions to metacognition in
685   perceptual decision making. The Journal of neuroscience : the official journal
686   of the Society for Neuroscience 32:6117-6125.
687   Fleming SM, Ryu J, Golfinos JG, Blackmon KE (2014) Domain-specific impairment
688   in metacognitive accuracy following anterior prefrontal lesions. Brain
689   137:2811-2822.
690   Fleming SM, Weil RS, Nagy Z, Dolan RJ, Rees G (2010) Relating introspective
691   accuracy to individual differences in brain structure. Science 329:1541-1543.
692   Hampton RR (2009) Multiple demonstrations of metacognition in nonhumans:
693   Converging evidence or multiple mechanisms? Comp Cogn Behav Rev
694   4:17-28.

Izquierdo A (2017) Functional heterogeneity within rat orbitofrontal cortex in reward
learning and decision making. The Journal of Neuroscience 37:10529-10540.
Janowsky JS, Shimamura AP, Squire LR (1989) Memory and metamemory:
Comparisons between patients with frontal lobe lesions and amnesic patients.
Psychobiology 17:3-11.
Jones JL, Esber GR, McDannald MA, Gruber AJ, Hernandez A, Mirenzi A,
Schoenbaum G (2012) Orbitofrontal cortex supports behavior and learning
using inferred but not cached values. Science 338:953-956.
Kennerley SW, Behrens TE, Wallis JD (2011) Double dissociation of value
computations in orbitofrontal and anterior cingulate neurons. Nature
Neuroscience 14:1581-1589.
Kepecs A, Uchida N, Zariwala HA, Mainen ZF (2008) Neural correlates, computation
and behavioural impact of decision confidence. Nature 455:227-231.
Kiani R, Corthell L, Shadlen MN (2014) Choice certainty is informed by both
evidence and decision time. Neuron 84:1329-1342.
Kornell N, Son LK, Terrace HS (2007) Transfer of Metacognitive Skills and Hint
Seeking in Monkeys. Psychological Science 18:64-71.
Kowalska DM, Bachevalier J, Mishkin M (1991) The role of the inferior prefrontal
convexity in performance of delayed nonmatching-to-sample.
Neuropsychologia 29:583-600.
Lak A, Costa GM, Romberg E, Koulakov AA, Mainen ZF, Kepecs A (2014)
Orbitofrontal cortex is required for optimal waiting based on decision
confidence. Neuron 84:190-201.
Levy R, Goldman-Rakic PS (1999) Association of Storage and Processing Functions
in the Dorsolateral Prefrontal Cortex of the Nonhuman Primate. The Journal
of Neuroscience 19:5149-5158.
Maniscalco B, Lau H (2012) A signal detection theoretic approach for estimating
metacognitive sensitivity from confidence ratings. Consciousness and
Cognition 21:422-430.
Mansouri FA, Buckley MJ, Mahboubi M, Tanaka K (2015) Behavioral consequences
of selective damage to frontal pole and posterior cingulate cortices.
Proceedings of the National Academy of Sciences 112:E3940-E3949.
McCurdy LY, Maniscalco B, Metcalfe J, Liu KY, de Lange FP, Lau H (2013)
Anatomical coupling between distinct metacognitive systems for memory and
visual perception. The Journal of Neuroscience 33:1897-1906.
Meiron O, Lavidor M (2014) Prefrontal oscillatory stimulation modulates access to
cognitive control references in retrospective metacognitive commentary.
Clinical Neurophysiology 125:77-82.
Middlebrooks PG, Sommer MA (2011) Metacognition in monkeys during an
oculomotor task. Journal of Experimental Psychology: Learning Memory and
Cognition 37:325-337.

Miyamoto K, Setsuie R, Osada T, Miyashita Y (2018) Reversible Silencing of the
Frontopolar Cortex Selectively Impairs Metacognitive Judgment on
Non-experience in Primates. Neuron 97:980-989.
Miyamoto K, Osada T, Setsuie R, Takeda M, Tamura K, Adachi Y, Miyashita Y
(2017) Causal neural network of metamemory for retrospection in primates.
Science 355:188-193.
Morales J, Lau H, Fleming SM (2018) Domain-General and Domain-Specific
Patterns of Activity Supporting Metacognition in Human Prefrontal Cortex.
The Journal of Neuroscience 38:3534-3546.
Noonan M, Walton M, Behrens T, Sallet J, Buckley M, Rushworth M (2010) Separate
value comparison and learning mechanisms in macaque medial and lateral
orbitofrontal cortex. Proceedings of the National Academy of Sciences
107:20547-20552.
Novikov NA, Nurislamova YM, Zhozhikashvili NA, Kalenkovich EE, Lapina AA,
Chernyshev BV (2017) Slow and Fast Responses: Two Mechanisms of Trial
Outcome Processing Revealed by EEG Oscillations. Frontiers in Human
Neuroscience 11.
Ochsner KN, Beer JS, Robertson ER, Cooper JC, Gabrieli JD, Kihsltrom JF,
D'Esposito M (2005) The neural correlates of direct and reflected
self-knowledge. Neuroimage 28:797-814.
Padoa-Schioppa C, Assad JA (2006) Neurons in the orbitofrontal cortex encode
economic value. Nature 441:223-226.
Petrides M (2000) Dissociable roles of mid-dorsolateral prefrontal and anterior
inferotemporal cortex in visual working memory. The Journal of Neuroscience
20:7496-7503.
Petrides M, Pandya D (1999) Dorsolateral prefrontal cortex: comparative
cytoarchitectonic analysis in the human and the macaque brain and
corticocortical connection patterns. European Journal of Neuroscience
11:1011-1036.
Pleskac TJ, Busemeyer JR (2010) Two-stage dynamic signal detection: A theory of
choice, decision time, and confidence. Psychological Review 117:864-901.
Qiu L, Su J, Ni Y, Bai Y, Zhang X, Li X, Wan X (2018) The neural system of
metacognition accompanying decision-making in the prefrontal cortex. PLoS
biology 16:e2004037.
Rolls ET, Grabenhorst F (2008) The orbitofrontal cortex and beyond: from affect to
decision-making. Progress in Neurobiology 86:216-244.
Rounis E, Maniscalco B, Rothwell JC, Passingham RE, Lau H (2010) Theta-burst
transcranial magnetic stimulation to the prefrontal cortex impairs
metacognitive visual awareness. Cognitive Neuroscience 1:165-175.
Rudebeck PH, Murray EA (2014) The orbitofrontal oracle: cortical mechanisms for
the prediction and evaluation of specific behavioral outcomes. Neuron
84:1143-1156.

Sawaguchi T, Iba M (2001) Prefrontal cortical representation of visuospatial working
memory in monkeys examined by local inactivation with muscimol. Journal of
Neurophysiology 86:2041-2053.
Schuck NW, Cai MB, Wilson RC, Niv Y (2016) Human orbitofrontal cortex
represents a cognitive map of state space. Neuron 91:1402-1412.
Shekhar M, Rahnev D (2018) Distinguishing the roles of dorsolateral and anterior
PFC in visual metacognition. The Journal of Neuroscience 38:5078-5087.
van de Vijver I, Ridderinkhof KR, Cohen MX (2011) Frontal oscillatory dynamics
predict feedback learning and action adjustment. J Cogn Neurosci
23:4106-4121.
Wallis JD (2007) Orbitofrontal cortex and its contribution to decision-making. Annual
Review of Neuroscience 30:31-56.
Wan X, Cheng K, Tanaka K (2016) The neural system of postdecision evaluation in
rostral frontal cortex during problem-solving tasks. eNeuro 3:1-24.
Ye Q, Zou F, Lau H, Hu Y, Kwok SC (2018) Causal evidence for mnemonic
metacognition in human precuneus. The Journal of Neuroscience
38:6379-6387.
Yeterian EH, Pandya DN, Tomaiuolo F, Petrides M (2012) The cortical connectivity
of the prefrontal cortex in the monkey brain. Cortex 48:58-81.

802 **Author Contributions**: S.C.K. and Y.C. and M.J.B. conceived the research. S.C.K. and M.J.B.

803 designed and performed experiments. S.C.K. and Y.C. analyzed data. S.C.K. and M.J.B. wrote

804 the manuscript with input from Y.C.

806 **Conflict of Interest**: The authors declare no competing financial interests.