Sarah Atherton[2], Ulf Jondelius[1],[2]

A taxonomic review and revisions of Microstomidae (Platyhelminthes:

Macrostomorpha)

1. Department of Zoology, Systematics and Evolution, Stockholms Universitet,

SE-106 91 Stockholm

2. Department of Zoology, Naturhistoriska riksmuseet

Box 50007, SE-104 05 Stockholm

Abstract

Microstomidae (Platyhelminthes: Macrostomorpha) diversity has been almost entirely

ignored within recent years, likely due to inconsistent and often old taxonomic

literature and a general rarity of sexually mature collected specimens. Herein, we

reconstruct the phylogenetic relationships of the group using both previously

published and new 18S and CO1 gene sequences. We present some taxonomic

revisions of Microstomidae and further describe 8 new species of *Microstomum* based

on both molecular and morphological evidence. Finally, we briefly review the

morphological taxonomy of each species and provide a key to aid in future research

and identification that is not dependent on reproductive morphology. Our goal is to

clarify the taxonomy and facilitate future research into an otherwise very understudied

group of tiny (but important) flatworms.

Introduction

Macrostomorpha DOE, 1986 is a group of free-living Platyhelminthes found

worldwide in aquatic and semi-aquatic habitats (Rieger 2001). They are very small in

size, typically only 1-2 mm, but are nevertheless important and often abundant

members of their communities, feeding on diatoms and other microorganisms or even

other small invertebrates (Ax 1995; Carrasco 2012; Atherton & Jondelius 2018b).

While the deep splits within Macrostomorpha remain ambiguous, current

phylogenetic hypotheses (Littlewood et al. 1999; Litvaitis & Rohde, 1999; Janssen et

al. 2015), place Microstomidae as sister to Dolichomacrostomidae within the group

Dolichomicrostomida.

Microstomidae LUTHER 1907 are characterised by ciliated sensory pits at the

frontal end, an intestine extending anteriorly past the mouth, and the ability to

reproduce asexually through fissioning. Currently, Microstomidae is unevenly divided

between three genera: *Myozonella* BEKLEMISHEV, 1955, with only a single species

Myozonella microstomoides BEKLEMISHEV, 1955 collected from a village in western

Russia and characterized by a muscular ring surrounding the anterior intestine;

Alaurina BUSCH 1851, with three currently accepted species all with an elongated and

unciliated (apart from sensory cilia) anterior proboscis; and the highly diverse and

speciose genus *Microstomum* ØRSTED 1843.

Species of Microstomidae have been considered difficult to distinguish

through traditional morphological methods. Their small size, relatively few

morphological characters, and high amounts of intraspecific variation (Luther 1960;

Bauchhens 1971; Heitkamp 1982; Atherton & Jondelius 2018b) all present taxonomic

challenges, especially as sexually mature specimens are generally rare. Furthermore,

the taxonomic literature may sometimes lack detail and is often old and/or difficult to

access, such that even the otherwise-reliable WORMS database

(http://www.marinespecies.org/aphia.php?p=taxdetails&id=142210) includes

numerous mistakes (e.g. recognition of *M. hanatum*, still listing *M. giganteum* as

synonymous to *M. lineare*). Taken together, it is unsurprising that while there has

been some recent studies of species diversity within Macrostomorpha in general (e.g.,

Schockaert 2014; Sun et al. 2014; Janssen et al. 2015; Fang et al. 2016; Reyes &

Brusa 2017; Lin et al. 2017a, b;), taxonomic research within Microstomidae from

even the last 50 years has been restricted to a very few articles (Faubel 1974, 1984;

Rogozin 2015; Atherton & Jondelius 2018 a,b).

Species of Microstomidae primarily reproduce through asexual fission, but

their life cycle may also include short periods of sexual reproduction occurring

primarily in the autumn (Bauchhenss 1971; Heitkamp 1982). During this time,

individual zooids will develop male and female sexual structures, including for the

male complex: single or bilateral testes, vasa deferentia, and a male copulatory

apparatus with vesicula seminalis, stylet and antrum masculinum; and for the female

complex: a single ovary and a female antrum. Species of Microstomidae may be

sexually mature for as little as two weeks a year or less (Bauchhenss 1971; Faubel

1974) and thus full or partial sexual anatomy is currently only known for roughly half

of the nominal species. Since species identification of Macrostomorpha has often

been based heavily or, in some cases, entirely on the morphology of the reproductive

organs (e.g. Nasonov 1935; Ax & Armonies 1987; Faubel & Cameron 2001), some

authors have argued that any species with unknown sexual anatomy should be

instantly considered species dubiae (Faubel 1984).

For Microstomidae with limited morphological data available, analysis of

nucleotide sequence data will be an indispensable tool when solving taxonomic

problems. DNA taxonomy is not without its own pitfalls (e.g. sensitivity to sampling,

overlap between intra and interspecific variation; DeSalle et al. 2005; Meier et al.

2006; Sauer & Hausdorf 2012), but has nevertheless been demonstrated as a reliable

and reproducible method to identify lineages without depending solely on

morphological characters that may be sensitive to environmental conditions or access

to sexually mature specimens (Pérez-Ponce de León & Poulin 2016; Fonseca et al.

2017). A number of different approaches to sequence-based species discovery, or

DNA-taxonomy, have been developed, including haplotype networks based on

statistical parsimony (Templeton *et al.* 1992), Bayesian species delineation (Yang &

Rannala 2010; Zhang *et al.* 2013), and maximum likelihood approaches with the

General Mixed Yule-Coalescent model (Pons *et al.* 2006).

DNA taxonomy may be more effective than traditional morphological

methods in discerning species, as re-examination of morphological taxa often reveals

hidden biodiversity in morphologically inseparable, i.e. cryptic species (Fiser et al.

2018). Successful DNA-taxonomy requires use of a molecular marker with an

appropriate level of variation, or, preferably, more than one such marker (Fontaneto et

al. 2015). Recent molecular investigations into the widespread taxon *Microstomum*

lineare revealed an assemblage of four genetically divergent but morphologically

similar species, some of which were found in the same sample (Atherton & Jondelius

2018b). The presence of multiple distinct but cryptic lineages in single locations

potentially could have much larger implications for Microstomidae research,

especially since studies in the past have often been performed on specimens collected

from environmental samples (e.g. Bauchhenss 1971; Heitkamp 1982; Reuter &

Palmberg 1989; Gutafsson et al. 1995).

In this paper, we reconstruct the phylogenetic relationships of *Microstomum*

using both previously published and new 18S and CO1 nucleotide sequences. Our

dataset comprises sequences from 336 specimens representing 15 previously

described nominal species and eight new species. The new taxa are diagnosed on the

basis of DNA-taxonomy as well as morphological characters that distinguish them

from all other known *Microstomum* species. We also review the morphological

taxonomy of the species within Microstomidae and provide a key to aid in future

research and identification. Our goals are to simplify the taxonomy, demonstrate that

asexual *Microstomum* specimens can often be identified, and to facilitate future

research into an otherwise very understudied group of microscopic flatworms that are

highly abundant in both limnic and marine habitats.

Methods

Sediments and aquatic vegetation were collected from fresh and marine

waters. Collection details for each specimen, including sampling date and GPS

coordinates, are listed in Table S1. Samples were transported back to laboratories and

processed according to type. Animals were extracted from marine sediments

following a standard anesthetization-decantation technique (Martens 1984) within 48

hours of collection. Freshwater sediments were first allowed to stagnate for 24-72

hours before the water was either directly removed into a glass petri dish or was first

filtered through a 63 µm sieve and backwashed to the petri dish. Vegetation was

simply washed through a fine mesh sieve.

Following extraction, animals were manually isolated under a Nikon SMZ

1500 stereomicroscope, transferred to a glass slide and identified with either a Nikon

Eclipse 80i compound microscope equipped with DIC (differential interference

contrast) or a Leitz LaborLux S compound microscope. Light micrographs and digital

videos were captured with a Cannon EOS 5D Mark III digital camera, and

measurements were taken with an ocular micrometer. Following documentation,

individual specimens were fixed in 95% ethanol and transported to Naturhistoriska

riksmuseet in Stockholm for DNA extraction and analysis.

DNA was extracted from whole animals using the DNeasy Blood & Tissue kit

(Qiagen, Valencia, CA) following the manufacturer's instructions. PCR amplification

was performed using 0.2 ml PuReTaq Ready-To-Go PCR Beads (GE Healthcare)

with 5 pmol each forward and reverse primers and 2 µl DNA. Table S2 lists primer

pairs and protocols used for amplification and sequencing. Products were viewed on

1-2% agarose gels, purified using ExoSAP-IT enzymes (Exonuclease and Shrimp

Alkaline Phosphatase; GE Healthcare) and sent for commercial sequencing to

Macrogen Europe (Netherlands) for commercial sequencing.

Amplification was performed on the complete nuclear 18S gene as well as a

~650 bp segment of the mitochondrial cytochrome oxidase c subunit 1 (CO1) loci

pertaining to the "Folmer region" (Folmer et al. 1994). The 18S ribosomal gene is

currently the most commonly used molecular marker for flatworms on GenBank

(8895 entries as of 5 July 2018), while CO1 gene sequences are known to be highly

variable in Platyhelminthes (e.g. Larsson et al. 2008; Telford et al. 2000; Atherton &

Jondelius 2018b) and other meiofauna (e.g. Verheye et al. 2016; Alvarez-Campos et

al. 2017; Kieneke & Nikoukar 2017; Michaloudi et al. 2017; Sahraean et al. 2017;

Zhao et al. 2018) and are thus applicable in identifying species and assessing

intraspecific diversity.

Two catenulid species, *Stenostomum leucops* and *Parcatenula* sp., and one

polyclad species, *Hoploplana californica*, were selected as outgroup taxa. Catenulida

is considered the sister group of all Rhabditophora, and Polycladida is the closest

relative of the Macrostomorpha (Egger et al. 2015; Laumer et al. 2015). Outgroup

sequences as well as available species of Macrostomorpha were downloaded from

GenBank and combined with new sequences for analysis. Table S1 list accession

numbers for all specimens used in subsequent analyses

Digital morphological vouchers (pictures and video) as well as collection data

of four previously undescribed *Microstomum* species were downloaded from the

Dryad Digital Repository (https://datadryad.org/resource/doi:10.5061/dryad.b5908)

and the Macrostomorpha Taxonomy and Phylogeny website

(http://www.macrostomorpha.info/). Further information on the collection and

processing of these species can be found in Janssen et al. (2015).

Sequence assembly and alignment was performed in Mega v 7.0.21 (Kumar et

al. 2016), and trace files were manually edited. Alignments for each marker were

generated using MUSCLE (Edgar 2004) with the default settings. CO1 sequences

were translated to amino acids using the flatworm mitochondrial genetic code

(Telford et al. 2000), manually checked for stop codons and reading frame shifts,

aligned and then reverted back to the original nucleotides for analysis. Maximum

likelihood (ML) analysis was performed on each marker individually along with

concatenated datasets in IQTree v1.6.4. The best fitting model was determined using

the ModelFinder algorithm (Kalyaanamoorthy et al. 2017) implemented in IQTree.

We performed 1000 ultrafast bootstrap replicates (Nguyen et al. 2015) in IQTree,

applying edge-proportional partitions with best-fit models independently selected for

each gene in the concatenated datasets.

Species of Microstomidae were inferred using the Bayesian implementation of

the Poisson Tree Processor (bPTP) in the web-based interface (http://species.h-

its.org/) with the default parameters (Zhang et al. 2013). Input trees were generated

from ML analyses of concatenated datasets that included Microstomidae sequences

and two sequences each of *Myozonaria bistylifera*, *Myozonaria fissipara* and

otherwise unidentified Myozonariinae downloaded from GenBank as outgroups

(Table S1). All bPTP tests were re-run up to three times with the input tree pruned so

that each prospective species included a maximum of five randomly selected

sequences in order to ensure that results were not influenced by an imbalance in the

number of specimens in any given clade. Following recommended guidelines,

outgroups were not included in any bPTP analysis.

Results

The full-concatenated alignment had a total of 336 specimens and total length

of 2348 bp. The K3Pu+F+G4 substitution model was selected as the best fit for the

CO1 gene sets and the TIM2e+I substitution model was selected for the 18S gene

sets. Tree topologies were consistent across all individual gene and concatenated

analyses.

Fig. 1 summarizes the results of the entire concatenated phylogeny (see also

Fig. S1). Results were generally consistent with Janssen et al. (2015), with maximum

(1.00) support for the two major subgroupings Macrostomidae and

Dolichomicrostomida. Results found further high (>0.90) support for: a monophyletic

Macrostomidae consisting of *Macrostomum* and *Psammomacrostomum*;

Myozonariinae as sister to Dolichomacrostominae, including the genera

Paramalostomum, *Austromacrostomum*, *Cylidromacrostomum* and

Dolichomacrostomum; and a monophyletic Microstomidae, which in this analysis

included *Alaurina composita* nested within *Microstomum*.

bPTP results were consistent across analyses, and a total of twenty species

were identified within the *Microstomum* clade (Fig. 2). Twelve of these were

previously described: *Alaurina composita* METSCHNIKOV 1865, *Microstomum artoisi*

ATHERTON & JONDELIUS 2018, *M. bispiralis* STIREWALT 1937*, M. crildensis* FAUBEL

1984*, M. edmondi* ATHERTON & JONDELIUS 2018*, M. laurae* ATHERTON & JONDELIUS

2018*, M. lineare* (MÜLLER 1773)*, M. papillosum* (GRAFF 1882)*, M. rubromaculatum*

(GRAFF 1882), *M. septentrionale* (SABUSSOW 1900), *M. tchaikovskyi* ATHERTON &

JONDELIUS 2018, and *M. zicklerorum* ATHERTON & JONDELIUS 2018. The other eight

species are new and are described below.

We obtained high support for four main clades within *Microstomum* (Figs. 1,

2), composed of: 1. all five freshwater species included in the analysis (*M. artoisi, M.*

bispiralis, M. lineare, M. tchaikovskyi, M zicklerorum); 2. six species (*M. afzelli* sp.

nov., *M. crildensis*, *M. inexcultus*, *M. marisrubri* sp. nov., *M. papillosum, M. schultei*

sp. nov.); 3. five species (*M. curinigalletti* sp. nov., *M. edmondi*, *M. lotti* sp. nov., *M.*

rubromaculatum, *M. weberi* sp. nov); and 4. *Alaurina composita* grouping with *M.*

laurae, *M. septentrionale* and *M. westbladi* nom.

Unfortunately, nucleotide sequences from two species of *Alaurina* and one

species of *Myozonella* were unavailable due to lack of specimens; however, results

from our phylogenetic analyses found all specimens of *Alaurina composita* nested

within the *Microstomum* clade (Figs. 1, 2), and thus we herein propose to synonymise

Alaurina with *Microstomum* and transfer its three species to *Microstomum*.

Nomenclature Acts

Synonymization of Alaurina *and amended diagnosis of* Microstomum *SCHMIDT 1848*

243   **Gen. Microstomum SCHMIDT 1848**

Microstomidae with ciliary pits at frontal end and preoral intestine. Two

excretion openings in the anterior body. Rhabdites and ingested nematocysts present

in some species. Asexual reproduction through fissioning as well as sexual

reproduction. Sexual anatomy including one or two testes beside or behind the

intestine; one or two ovaries consisting of one or more lobes and containing a central

oocyte and a peripheral layer of nutrient cells. Body size of solitary individual 0.6-

4mm, with chains comprising up to 18 zooids and 15 mm.

Type species: *Microstomum lineare* (Müller, 1773).

43 species: 33 marine, 9 limnic, and 1 inhabiting both brackish (up to 7‰) and

freshwaters.

Synonymization of Microstomum hamatum *WESTBLAD 1953 and* Microstomum

lucidum *(FUHRMANN, 1896)*

M. lucidum and *M. hamatum* are morphological identical accepting the

ambiguously-defined "dark greyish ± pigment spots" in *M. hamatum* (Westblad

1953). Both species have bodies colored yellow due to the intestine, with identical

body shapes and sizes and adhesive papillae on the posterior ends. Both have

rhabdites sparsely present over the entire body, concentrated at the posterior end and a

preoral intestine extending to the brain. Moreover, both species inhabit the Northern

Atlantic in overlapping habitats, in mud on opposite sides of the English Channel:

Baie de la Forêt near Concarneau for *M. lucidum*, and the harbor near Plymouth for

M. hamatum. Unfortunately, Furhmann provided no images of *M. lucidum* apart from

a drawing of a single rhabdite bundle and was unable to collect specimens with sexual

anatomy. Nevertheless, the similarities in morphology and ecology suggest it is highly

likely that these represent the same species if—as we have interpreted it—the "±" of

Westblad's (1953) description means that some individuals may not have dark grey

spots. Regardless, body pigmentation is known to be highly variable in this genus

(Stirewalt 1937; Luther 1960), and Westblad (1953) gave no explanation on how the

two species should be distinguished. In this case, *Microstomum hamatum* WESTBLAD

1953 must be considered a junior synonym for *Microstomum lucidum* FUHRMANN

1896.

Species Descriptions

279   Order Macrostomida KARLING, 1940

280   Family Microstomidae LUTHER, 1907

281   Genus *Microstomum* SCHMIDT, 1848

283   ***Microstomum afzelii* sp. nov.**

284   Fig. 3

Diagnosis

Marine *Microstomum* with an almost elliptical body shape; body length 0.8 mm, 2

zooids. Colorless and without pigmented eyespots. Anterior end smoothly rounded.

Posterior end slightly more tapered but also with rounded tip. Ciliary pits small and

shallow; may be difficult to discern in some specimens. Few adhesive papillae present

on anterior and posterior ends of body. Rhabdites sparsely present throughout body.

Nematocysts particularly concentrated at posterior end. Preoral intestine extending

anterior to brain. Reproductive system unknown. GenBank accession number for

partial CO1 sequence XXX; for 18S XXX.

Etymology

Named for Lars Fredrik Afzelius, the first director of the Tjärnö Marine Laboratory,

in appreciation of his contributions to the marine scientific community.

Material examined

Holotype: asexual specimen, Sweden, Stömstad, Saltö, 58°52'29''N 11°08'41'' E; 18

June 2016; 10 cm depth, marine, eulittoral sand, S. Atherton leg. (SMNH-Type-XXX;

GenBank Accession CO1 XXX, 18S XXX)

Additional materials: asexual specimen; same as type (SMNH-XXX; GenBank

Accession CO1 XXX, 18S XXX)

Description

Microstomum with a total body length of 0.8 mm, 2 zooids. Body colorless

and without pigmented eyespots. Body shape almost elliptical, with widest point

occurring approximately at midpoint or slightly anterior, lightly decreasing in width

to a rounded anterior end, and a slightly more pronounced tapering to the rounded

posterior end. Ration of body width:length ~1:2.5 in slightly compressed animal.

Ciliary pits weakly present, may be difficult to distinguish in some specimens.

Small and shallow, located at level of posterior brain.

Epidermis uniformly covered with cilia. A few longer sensory cilia present on

posterior end. Nematocysts present, highly concentrated at posterior. Rhabdites

present, 15-20 µl long, sparsely scattered across entire body. Approximately 6-8

adhesive papillae, 10-15 µm long, on the anterior end; a further 10-12 located at the

posterior rim.

Mouth circular, 50 µm in diameter, surrounded by cilia. Pharynx also ciliated,

138 µm long, surrounded by relatively few pharyngeal glands. Preoral intestine

extending anterior to brain. Intestine filled with lightly orange-brown food.

Reproductive system unknown.

Notes

There are currently 23 species of *Microstomum* that have neither pigmented

eyespots nor distinct body pigmentation. Of these, *M. afzelii* sp. nov. can generally be

easily separated by its body shape, especially the rounded body ends, preoral intestine

extending anterior to the brain and the presence of the adhesive papillae on the

anterior end. The other species of *Microstomum* without very distinct and

characteristic pigmentation can specifically be separated from *M. afzelii* sp. nov. in

the following ways:

M. album (ATTEMS 1896) – anterior proboscis present; more numerous posterior

adhesive papillae, presence of adhesive papillae on midbody; more numerous

rhabdites; short preoral intestine

M. bispiralis STIREWALT 1937 – posterior end pointed; preoral intestine short;

adhesive papillae and rhabdites absent; limnic habitat; see also Figs. 1, 2

M. breviceps MARCUS 1951 – very distinct spatulate posterior end; biparte pharynx

M. canum (FUHRMANN 1894) – longer, more pointed anterior end; posterior end

pointed; ciliary pits deep; adhesive papillae and rhabdites absent; limnic habitat

M. crildensis FAUBEL 1984– short preoral intestine; anterior adhesive papillae absent;

rhabdites absent; distinct constriction of body width at level of brain; see also Figs. 1,

2

M. davenporti GRAFF 1911– anterior adhesive papillae absent; more numerous

anterior rhabdites; biparte pharynx

M. edmondi ATHERTON & JONDELIUS 2018 – posterior end blunt; deep bottle-shaped

ciliary pits; anterior adhesive papillae absent; more numerous rhabdites; see also Figs.

1, 2

M. giganteum HALLEZ 1878 – very large body size; preoral intestine short; adhesive

papillae and rhabdites absent; limnic habitat

M. inexcultus sp. nov. – rhabdites absent; anterior adhesive papillae absent; more

numerous posterior adhesive papillae; see also Figs. 1, 2

M. jenseni RIEDEL, 1932 – posterior end tail-like; preoral intestine only reaching to

level of brain; anterior adhesive papillae absent

M. lineare (MÜLLER 1773) – often with anterior red eyestripes; posterior end tail-like;

adhesive papillae appearing absent or very small; rhabdites absent; limnic or brackish

water habitat; see also Figs. 1, 2

M. lotti sp. nov. – distinct spatulate tail-plate; very large ciliated pits; anterior crown

of adhesive papillae; small preoral intestine; see also Figs. 1, 2

M. lucidum (FUHRMANN 1896) – preoral intestine extending only to level of brain;

anterior adhesive papillae absent

M. marisrubri sp. nov. – rhabdites absent; more numerous posterior adhesive papillae;

anterior end longer; preoral intestine small; see also Figs. 1, 2

M. mundum GRAFF 1905 – characteristic lateral bulges in intestine; preoral intestine

extending only to level of brain; posterior end tail-like; anterior adhesive papillae

absent

M. papillosum (GRAFF 1882) – more numerous rhabdites at anterior end; adhesive

papillae scattered across entire body; see also Figs. 1, 2

M. rhabdotum MARCUS 1950 – biparte pharynx; paired stripes of rhabdite bundles at

ciliary pits; preoral intestine extending only to level of brain

M. spiculifer FAUBEL 1974 – anterior adhesive papillae absent; more numerous

posterior rhabdites; longer preoral intestine reaching almost to very tip of body.

M. trichotum MARCUS 1950 – long and tapered anterior end; more numerous rhabdites

concentrated at anterior; anterior adhesive papillae absent; preoral intestine extending

only to level of brain

M. ulum MARCUS 1950 – preoral intestine short; deep ciliary pits; distinct spatulate

tail plate; more numerous rhabdites concentrated at posterior end

M. weberi sp. nov. – bottle-shaped ciliary pits; anterior crown of adhesive papillae;

see also Figs. 1, 2

M. westbladi nom. nov. – more numerous rhabdites highly concentrated at anterior

end; adhesive papillae scattered across entire body; characteristic shape of field of

adhesive papillae at anterior end; see also Figs. 1, 2

M. zicklerorum ATHERTON & JONDELIUS 2018 – individuals may have anterior red

eyestripes; posterior end tail-like; adhesive papillae and rhabdites absent; limnic

habitat; see also Figs. 1, 2

388   ***Microstomum curinigalletti* sp. nov.**

389   Fig. 4

Diagnosis

Marine *Microstomum* with a body length of 1.4 mm, 2 zooids. Anterior body

end colored ochre to dark yellow-orange. Body otherwise colorless without

pigmented eyespots. Body shape with a conically rounded anterior end and wide,

blunt posterior end. Numerous adhesive papillae present at posterior end. Rhabdites

present but few, scattered across the body. Ciliary pits large and bottle-shaped,

located anterior to pharynx. Preoral intestine short. Reproductive system unknown.

GenBank accession number for partial CO1 sequence XXXX; for 18S XXX.

Etymology

This species is dedicated with many thanks to Dr. Marco Curini-Galletti, who found

the first specimens of these animals.

Material examined

Holotype: Asexual specimen; Sweden, Fiskebäckskil, Kristineberg Sven Lovén

Center for Marine Research, 58°14′59″ N, 11°26′45″ E, 20 Aug. 2015, marine,

sublittoral phytal on algae, M. Curini-Galletti leg. (SMNH-Type-XXXX; GenBank

Accession 18S XXX)

Additional materials: 5 asexual; same as holotype (SMNH-XXXX; GenBank

Accession 18S XXX)

Description

Microstomum with a total body length to 1.4 mm, 2 zooids. Anterior body end

colored ochre to dark yellow-orange. Body otherwise colorless and reflecting

intestines. Pigmented eyespots absent. Body shape with a conically rounded anterior

end; posterior end blunt and almost as wide as rest of body. Ratio of width generally

consistent across body, width:length is 1:4 in slightly compressed animal.

Ciliary pits large, clearly distinguishable anterior to pharynx; bottle shaped,

extending at a 45° angle posterior into the body ~65 µm; opening 30 µm long.

Epidermis uniformly covered with cilia. Nematocysts present. Few rhabdites,

to 25 µm long, scattered throughout the body. Numerous adhesive papillae, 12-15 µm

long, present at posterior tip.

Mouth small and circular surrounded by cilia. Pharynx surrounded by normal

ring of pharyngeal glands. Preoral intestine very short, hardly extending past mouth.

Reproductive system unknown

Notes

In body shape and size and, especially, size, shape and location of ciliary pits

(Fig. 4C), *Microstomum curinigalletti* sp. nov. is most similar to *M. edmondi*. The two

species also share habitats and distributions and were collected in the same samples.

M. curinigalletti sp. nov., however, can easily be distinguished from *M. edmondi*

through the yellow pigmentation of the anterior end and the shorter preoral intestine.

Results from the phylogenetic and bPTP analyses also clearly separated the two

species (Figs. 1, 2).

436   ***Microstomum inexcultus* sp. nov.**

437   Fig. 5

Diagnosis

Marine *Microstomum* with a body length of 0.85, 2 zooids. Colorless and without

pigmented eyespots. Body shape with a conically pointed anterior end and a tapered

posterior end with a blunt tip. Numerous adhesive papillae present at posterior end.

Rhabdites absent. Ciliary pits clearly present. Preoral intestine long, stretching almost

to anterior body end. Reproductive system unknown. GenBank accession number for

partial CO1 sequence XXXX; for 18S XXX.

Etymology

This species named for its very simple appearance without rhabdites (*inexcultus*,

Latin: unadorned, unembellished)

Material examined

Holotype: Asexual specimen; Sweden, Stömstad, Saltö, 58°52'29''N 11°08'41'' E;

18 Jun. 2016; 10 cm depth, marine, eulittoral sand, S. Atherton leg. (SMNH-Type-

XXXX; GenBank Accession CO1 XXX, 18S XXX)

Additional materials: 5 asexual specimens, same as holotype; 2 asexual specimens

Sweden, 58°25'24''N 11°39'34'' E; 15 Jun. 2015; 1 m depth, marine; S. Atherton

leg. (SMNH-XXXX; GenBank Accession CO1 XXX, 18S XXX)

Description

Microstomum with a total body length to 0.85 mm, 2 zooids. Body colorless

and without pigmented eyespots. Body shape with a conically pointed anterior end

and a tapered posterior end with a blunt tip. Maximum body width just posterior to

pharynx; ratio of maximum width:length is ~1:3 in slightly compressed animal.

Ciliary pits clearly distinguishable at level of anterior pharynx 170 µm from

anterior body end; roundish to cup-shaped, extending at a slight angle into the body

27 µm; opening 20 µl long.

Epidermis uniformly covered with cilia. Very few nematocysts present,

primarily at body ends. Rhabdites absent. Numerous adhesive papillae, ~15 long,

present at posterior end and along up to ~⅕ of the posterior body; may also be present

at zone of division in very well developed zooids.

Mouth elliptical, 45 µm long, surrounded by cilia. Pharynx 125 µm long,

surrounded by normal ring of pharyngeal glands. Preoral intestine extending well

anterior to brain, reaching almost to anterior body tip.

Reproductive system unknown

Notes

Microstomum inexcultus sp. nov. can be distinguished from all other species of

Microstomum through a combination of the following characters: absent of

pigmentation or eyespots; preoral brain very long and extending almost to anterior

body tip (Fig. 5C); rhabdites absent. The absence of rhabdites in particular

distinguished this species, since only 11 other species of *Microstomum* do not have

rhabdites recorded; these can be distinguished from *M. inexcultus* sp. nov. by the

following:

M. artoisi ATHERTON & JONDELIUS 2018 – anterior red eyestripes present; preoral

intestine short; posterior end tail-like; adhesive papillae absent; limnic habitat; see

also Figs. 1, 2

M. bispiralis STIREWALT 2018 – preoral intestine short; posterior end pointed;

adhesive papillae absent; limnic habitat; see also Figs. 1, 2

M. caerulescens (SCHMARDA, 1859) – body blue; adhesive papillae absent; limnic

habitat

M. canum (FUHRMANN 1894) – posterior end tail-like; ciliary pits deep; adhesive

papillae absent; limnic habitat

M. crildensis FAUBEL, 1984 – more rounded anterior; constriction of body width at

level of brain; short preoral intestine; see also Figs. 1, 2.

M. giganteum HALLEZ 1878 – very large body size; preoral intestine short; adhesive

papillae absent; limnic habitat

M. lineare (MÜLLER, 1773) – anterior red eyestripes often present; posterior end tail-

like; adhesive papillae appearing very small or absent; limnic or brackish water

habitat; see also Figs. 1, 2

M. marisrubri sp. nov. – longer frontal end; small preoral intestine; anterior adhesive

papillae present; see also Figs. 1, 2.

M. punctatum DORNER 1902 – body brown-yellow with black spots; preoral intestine

short; posterior end tail-like; limnic habitat

M. tchaikovskyi ATHERTON & JONDELIUS 2018 – anterior red eyestripes present;

preoral intestine short; posterior end tail-like; adhesive papillae absent; limnic habitat;

see also Figs. 1, 2

M. zicklerorum ATHERTON & JONDELIUS 2018 – anterior red eyestripes may be

present; preoral intestine short; posterior end tail-like; adhesive papillae absent;

limnic habitat; see also Figs. 1, 2

512   ***Microstomum lotti* sp. nov.**

513   Fig. 6

Diagnosis

Marine *Microstomum* with a body length of 0.8, 2 zooid. Body colorless without

pigmented eyespots. Anterior end conical; posterior end forming a spatulate tail-plate

set off by a constriction. Adhesive papillae present along posterior half of the body

and very numerous at tail plate. An additional row of adhesive papillae present

ventrally at the frontal region ~30 µm from anterior tip. Ciliary pits rounded. Preoral

intestine short. Rhabdites present. Reproductive system unknown. GenBank accession

for 18S KP730483, KP730493.

Etymology

This species is named in honor of Christian Lott, who was a co-organizer of the 2010

BIOSAND workshop, where the sample was collected.

Material examined

Holotype: Asexual specimen; Italy, Sant Andrea Bay, 42°48′31′′ N, 10°08′30′′ E, 26

April 2010, marine (MTP LS 660; GenBank Accession 18S KP730483)

Other materials: same as holotype (MTP LS 635; GenBank Accession 18S

KP730493)

Description

Microstomum with a total body length to 0.8 mm, 2 zooids. Body colorless

and without pigmented eyespots. Anterior end conically pointed with row of adhesive

papillae at the frontal end, ~30 µm from anterior tip. Posterior end forming a distinct

tail-plate separated by a constriction of the body width. Width to length ration is

approximately 1:5.

Ciliary pits large, present posterior to brain; each forming a roundish

indentation angled slightly anteriorly; extending inward 21 µm with 16 µm long

openings.

Epidermis uniformly covered with cilia. Rhabdites bundles scattered

throughout the body, ~12-17 µm. In addition to those at the frontal end, numerous

adhesive papillae, 10 µm long, are present at the tail-plate and less frequently at the

margins to approximately halfway up the body.

Mouth circular and surrounded by cilia. Pharynx 220 µm long, surrounded by

normal ring of pharyngeal glands. Preoral intestine reaching to level halfway between

pharynx and brain.

Reproductive system unknown.

Notes

The only species of *Microstomum* with distinct spatulate tail-plates are *M. breviceps*

and *M. ulum*. *M. lotti* sp. nov. differs from *M. breviceps* by the row of adhesive

papillae on the frontal end (Fig. 8C) and by the small preoral intestine. *M. lotti* sp.

nov. is more similar to *M. ulum*, although the two species still can be differentiated by

the field of rhabdites at the frontal end of *M. ulum* and the row of adhesive papillae at

the frontal end of *M. lotti* sp. nov.

560   ***Microstomum marisrubri* sp. nov.**

561   Fig. 7

Diagnosis

Marine *Microstomum* with a body length of 0.65, 1 zooid. Body colorless without

pigmented eyespots. Anterior end large, with a rounded tip. Posterior end blunt.

Adhesive papillae sparsely present over entire body but extremely numerous at

posterior ⅕ of body. Ciliary pits present, cup shaped, located at level between anterior

pharynx and brain. Preoral intestine short, reaching just anterior to pharynx.

Rhabdites absent, but with many nematocysts intensely concentrated at posterior end.

Male stylet a curved tube tapering distally to a slightly enlarged end. GenBank

accession number for partial CO1 sequence KP730580; for 18S KP730505.

Etymology

This species is named after the collection location *(maris,* Latin: sea; *rubri* Latin: red)

Material examined

Holotype: □; Egypt, Mangrove Bay, 25°52'15″ N, 34°25'04″ E, 11 January 2009,

marine (lactophenol preparation NHMUK-2015.1.30.22; digital morphological

voucher MTP LS 524; GenBank Accession CO1 KP730580, 18S KP730505)

Description

Microstomum with a total body length to 0.65 mm, 2 zooids. Body colorless

and without pigmented eyespots. Anterior end long, measuring 95 µm from tip to

anterior brain, with a widely rounded tip. Body width slightly decreasing from frontal

end to ciliary pits, slightly increasing toward the middle of the animal and then

decreasing again to a blunt posterior end. Animal measures approximately

80:70:120:45 µm at frontal end: ciliary pits: body middle: posterior end.

Ciliary pits present posterior to brain; each forming a roundish indentation

angled at approximately 45° into the body; extending 10 µm with 10 µm long

openings.

Epidermis uniformly covered with cilia. Rhabdites absent. Very many

nematocysts present throughout the animal and especially intensely concentrated at

the posterior end. Numerous adhesive papillae, ~10 µm long, present along ~⅕ of the

posterior body and also sparsely occurring anteriorly.

594   Mouth surrounded by cilia. Pharynx 95 µm long, surrounded by normal ring

595   of pharyngeal glands. Preoral intestine short, terminating just posterior to level of

596   brain.

597   Male reproductive system with single developing testis connected by short vas

deferens to a rounded vesicula granulorum. Male stylet, 63 µm long, a slightly

tapering tube, 9 µm wide at base and ~3 µm at narrowest part of distal end; curving a

little less than 90°, with a slight widening at the distal tip to ~4.5 µm. Genital pores

separate. Male pore 18 µm in diameter.

Notes

The stylet and reproductive system of *Microstomum marisrubri* sp. nov. (Fig. 7 E) is

very similar to *M. papillosum*, although it is smaller (62 µm compared to 80 µm,

respectively). Other differences between the two species can be found in the size of

the preoral intestine, which *for M. papillsum* extends past the brain, and the presence

of numerous rhabdites at the anterior end. As discussed above, few species of

Microstomum lack rhabdites, and these can be distinguished:

M. artoisi ATHERTON & JONDELIUS 2018 – anterior red eyestripes present; posterior

end tail-like; adhesive papillae absent; limnic habitat; see also Figs. 1, 2

M. bispiralis STIREWALT, 1937 – posterior body pointed; adhesive papillae absent;

limnic habitat; see also Figs. 1, 2.

M. caerulescens (SCHMARDA, 1859) – body blue; preoral intestine extending past

brain; adhesive papillae absent; limnic habitat

M. canum (Fuhrmann 1894) – posterior end tail-like; adhesive papillae absent;

limnic habitat

M. crildensis Faubel, 1984 – shape and size of the male stylet; lack of anterior

adhesive papillae and fewer posterior adhesive papillae; see also Figs. 1, 2.

M. giganteum Hallez 1878 – very large body size; posterior end a thin tail; adhesive

papillae absent; limnic habitat

M. inexcultus sp. nov. – smaller frontal end; large preoral intestine; lack of anterior

adhesive papillae; see also Figs. 1, 2.

M. lineare (Müller, 1773) – anterior red eyestripes often present; posterior end tail-

like; adhesive papillae very small or appearing absent; limnic or brackish water

habitat; see also Figs. 1, 2

M. punctatum Dorner 1902 – body brown-yellow with black spots; posterior end

tail-like; limnic habitat

M. tchaikovskyi Atherton & Jondelius 2018 – anterior red eyestripes present;

posterior end tail-like; adhesive papillae absent; limnic habitat; see also Figs. 1, 2

M. zicklerorum Atherton & Jondelius 2018 – anterior red eyestripes may be

present; posterior end tail-like; adhesive papillae absent; limnic habitat; see also Figs.

1, 2

637   ***Microstomum schultei* sp. nov.**

638   Fig. 8

Diagnosis

Marine *Microstomum* with a body length of 1.5 mm, 1 zooid. Body appears light

reddish-brown without pigmented eyespots. Anterior end rounded and set apart from

the rest of the body by a slight constriction at the ciliary pits. Posterior end conically

blunted. Adhesive papillae and rhabdites concentrated at anterior and posterior ends.

Ciliary pits present, cup shaped. Preoral intestine short. Reproductive system with

curved male stylet, tapering from a wider base to a narrow tube with a beveled tip.

GenBank accession number for 18S KP730494.

Etymology

This species is named in honor of Gregor Schulte, who was a field assistant during the

2010 BIOSAND workshop and aided in sample collection.

Material examined

Holotype: □; Italy, Fetovaia Bay, 42°43′36′′ N, 10°09′33′′ E, 05 May 2010, marine

(lactophenol preparation NHMUK-2015.1.30.21; digital morphological voucher MTP

LS 700; GenBank Accession 18S KP730494)

Description

Microstomum with a total body length to 1.5 mm, 1 zooid. Body light reddish-

brown, without pigmented eyespots. Anterior end rounded and set apart from the rest

of the body by a slight constriction at the ciliary pits. Posterior end conically blunted.

Ratio of maximum width:length is ~1:5 in slightly compressed animal.

Ciliary pits present at level just posterior to brain, 170 μm from anterior body

end; roundish to cup-shaped, extending into the body 30 μm with 15 μm long

openings.

Epidermis uniformly covered with cilia. Rhabdites present throughout the

body, but especially concentrated at the anterior and posterior ends. Numerous short

adhesive papillae, ~10 µm long, present, heavily concentrated at posterior end along

~⅙ of the body and also anterior to brain.

Mouth circular, 40 µm long, surrounded by cilia. Pharynx 185 µm long,

surrounded by normal ring of pharyngeal glands. Preoral intestine short.

Reproductive with single medial ovary and caudally developing eggs. Male

system with vesicula granulorum connected to male stylet, 60 µm long, curved and

tapering from a wider base to form a narrow tube with a beveled tip. Stylet base 20

µm wide, distal tube ~4 µm wide, opening ~5 µm long. Sperm ~70 µm long, with a

rounded head and a single flagellum with lateral setae.

Notes

There are a few species of *Microstomum* with a similarly shaped male stylet (Fig.

6D). *M. schultei* sp. nov. can be separated from each of these through:

M. crildensis FAUBEL, 1984 – stylet much smaller (32 µm instead of 60 µm), with a

more narrow base; rhabdites absent; anterior adhesive papillae absent; body colorless;

see also Figs. 1, 2

M. gabriellae MARCUS, 1950 – stylet even more curved, tracing an entire half circle

180°; terminal end not beveled; preoral intestine large, extending past brain; red

eyespot present

M. septentrionale (SABUSSOW 1900)– stylet without beveled terminal end; preoral

intestine reaching to level of anterior brain; body color with dorsal yellow

pigmentation; see also Figs. 1, 2.

In addition, *M. schultei* sp. nov can generally be distinguished from the other species

of *Microstomum* through a combination of the presence of adhesive papillae at the

anterior and posterior ends and a short preoral intestine (defined here as a preoral

intestine that does not extend to at least the level of posterior brain; Fig. 6). *M.*

schultei sp. nov. can be separated from the other species of *Microstomum* with these

characters by the following ways:

M. album (ATTEMS 1896) – body colorless or white; anterior proboscis present;

ciliary pits small

M. crildensis FAUBEL, 1984 – see above

M. curinigalletti sp. nov. – frontal end colored ochre; ciliary pits large and bottle

shaped; anterior adhesive papillae absent; fewer rhabdites; see also Figs. 1, 2

M. lotti sp. nov. – posterior end spatulate; anterior crown of adhesive papillae; body

colorless; see also Figs. 1, 2

M. marisrubri sp. nov. – longer frontal end; more numerous posterior adhesive

papillae; rhabdites absent; body colorless; see also Figs. 1, 2

M. punctatum DORNER 1902 – body brown-yellow with black spots; pointed anterior

and tail-like posterior ends; ciliary pits small and shallow; rhabdites absent; limnic

habitat

M. rubromaculatum (GRAFF, 1882) - red pigmentation stripe at anterior end present;

widely blunt posterior end; see also Figs. 1, 2

M. ulum MARCUS 1950 – pointed anterior end; distinct spatulate tail-plate at posterior

end; more rhabdites concentrated at anterior end; body colorless

***Microstomum weberi* sp. nov.**

Fig. 9

Diagnosis

Marine *Microstomum* with a body length of 0.65 mm, 2 zooids. Body colorless

without pigmented eyespots. Anterior end conical; posterior end blunt. Adhesive

papillae occur at posterior end and in a ventral row at the frontal end. Ciliary pits

bottle-shaped. Preoral intestine extending to anterior brain or just beyond. Large

rhabdites bundles present at anterior and posterior ends. Reproductive system

unknown. GenBank accession number for partial CO1 sequence KP730576; for 18S

KP730487.

Etymology

This species is named in honor of Miriam Weber, who collected the sample and was

co-organizer of the 2010 BIOSAND workshop

Material examined

Holotype: Asexual specimen; Italy, Pianoso, 42°34′29″ N, 10°03′59″ E, 30 April

2010, marine (MTP LS 666; GenBank Accession CO1 KP730576, 18S KP730487)

Description

        Body length 0.65 mm, 2 zooids. Body colorless and without pigmented

eyespots. Anterior end conically rounded. Body width smaller at anterior end and

increases perceptively just after ciliary pits. Posterior tapering to a wide blunt end.

        Ciliary pits bottle-shaped, present at level of anterior pharynx; extending 13

μm into the body at slight posteriorly-directed angle; with 12 μm long openings.

Epidermis uniformly covered with cilia. Large rhabdites bundles, to 20 µm,

present only at anterior and posterior ends. A row of adhesive papillae at the frontal

end, situated ventrally 25 µm from anterior tip. Additional adhesive papillae, ~7-10

µm long, also present at posterior end, but are otherwise absent from the body

margins.

Mouth eliptical and surrounded by cilia. Pharynx 105 µm long, surrounded by

normal ring of pharyngeal glands. Preoral intestine reaching just past anterior brain.

Reproductive system unknown.

Notes

This is only the second species of *Microstomum* with a ventral row of adhesive

tubules on the frontal end (Fig. 9C), the other being *M. lotti* sp. nov. *M. weberi* sp.

nov. can be distinguished by general body shape and size, through the bottle-shaped

ciliary pits (Fig. 9C), the longer preoral intestine and by the large rhabdite bundles

present only near the body ends. Results from the phylogenetic and bPTP analyses

also clearly separated the two species (Figs. 1, 2).

Microstomum westbladi **nom. nov.**

Fig. 10

Synonym

Microstomum papillosum GRAFF 1882, sensu Westblad 1953, pg 406-7

Material examined

Holotype: Asexual, sectioned specimen; Norway, Herdla; 15 July 1934; 5-10 m

depth, marine amphioxus sand, E. Westblad leg. (SMNH-Type-XXX74)

Additional materials: 1 asexual, sectioned, same as holotype (SMNH-94979); 1

asexual specimen, Sweden, Fiskebäckskil, 58°15'05''N 11°27'52'' E, 26 Aug. 2015,

marine, eulittoral sand, S. Atherton leg. (SMNH-XXXX; GenBank Accession CO1

XXX, 18S XXX)

Westblad, 1953 – Norway, Herdla, Kværnepoll current; Sweden, Gullmarfjord

This paper: Sweden, Fiskebäckskil, 58°15'05''N 11°27'52'' E

Diagnosis

Marine *Microstomum* with strap shaped body, to 2.0 mm long and 5 zooids.

Colorless and without pigmented eyespots. Anterior end rounded. Posterior tapered to

a blunt end. Ciliary pits small. Numerous rhabdites intensely concentrated at frontal

end and present throughout body. Numerous adhesive papillae present at posterior

end and more sparsely anterior. Characteristic grouping of adhesive papillae covering

the anterior tip: field longest at both lateral sides and curving upwards toward the

middle. Preoral intestine extending to brain. Reproductive system including male

stylet, 60 µm long, and shaped as a smooth curve 180° curve, with a maximum width

~⅙ of the way to the distal end, decreasing proximally and also distally to a point.

GenBank accession number for partial CO1 sequence XXXX; for 18S XXX.

Etymology

This species is named in honor of Dr. Einar Westblad, who first described these

animals.

Materials examined

2 asexual; Sweden, Malmön, 58°20'18'' N 11°20'26'' E; 13 Aug. 2015; marine, 1.5

m depth; S. Atherton & Y. Jondelius leg. (SMNH-XXXX; GenBank Accession CO1

XXX, 18S XXX)

Description

Microstomum with a total body length to 2.0 mm, 5 zooids. Body colorless

and without pigmented eyespots. Body strap-shaped with a bluntly rounded anterior

end, an almost equal width along entire body until a slight decreasing in width to a

bluntly rounded posterior end.

Ciliary pits small and shallow cups, located at level of anterior pharynx.

Epidermis uniformly covered with cilia. Nematocysts not seen or recorded.

Rhabdites present, ~15-30 µm long, throughout entire body, but intensely

concentrated at anterior end. Numerous adhesive papillae, ~12-15 µm long, present at

posterior end and more sparsely anterior. Further adhesive papillae present at the

anterior tip grouped in a very distinct pattern: the field covers the anterior tip, toupee

like, i.e. longest at both lateral sides and curving upwards toward the middle.

Mouth elliptical surrounded by cilia. Pharynx 110 µm long, surrounded by

normal ring of pharyngeal glands. Preoral intestine extending to brain.

Reproductive system mostly unknown. Following Westblad (1953) stylet

recorded as 60 µm long, shaped as a smooth curve 180° curve, with the width first

increasing along the most proximal ~$\frac{1}{6}$ of the stylet and then smoothly decreasing in

width to a pointed, terminal opening.

Notes

Microstomum westbladi nom. nov. can be readily distinguished from *M.*

papillosum by the characteristic pattern of the anterior adhesive papillae (Fig. 10B,

C), the very high concentration of rhabdites in the anterior end (Fig. 10D) and the

shape of the stylet, which decreases in width toward the proximal end as well as

distally. In addition, results of the phylogenetic and bPTP analyses clearly separate

the two species (Figs. 1, 2).

M. westbladi nom. nov. is most similar to *M. septentrionale*, particularly since

both species are of similar size and shape, have a very high concentration of rhabdites

in the anterior end, and are known to have overlapping habitats. However, *M.*

westbladi nom. nov. can be distinguished morphologically by its colorless body

without dark or orange-yellow in the anterior and through the characteristic pattern of

anterior adhesive papillae. Additionally, the stylet shape of *M. westbladi* nom. nov.

differs by decreasing in width toward the proximal end. *M. septentrionale* and *M.*

westbladi nom. nov. are closely related sister taxa (Figs. 1, 2).

Discussion

Phylogenetic Analyses and DNA Taxonomy

The use of nucleotide sequences to delineate species and capture biodiversity

is an important advancement in Microstomidae taxonomy. As noted in the

introduction, the animals may have relatively few morphological characters on which

to base descriptions, and thus identification from morphology alone may require high

amounts of effort, specimens and specialized taxonomic expertise. Further, many

morphological features are highly variable depending on environmental conditions

(Stirewalt 1937; Bauchhenss 1971; Heitkamp 1982). Laboratory studies of

Microstomum lineare have demonstrated that body shape and size varies based on

food availability and sexual maturity; body color ranges from colorless, white,

yellow, grey, reddish or brownish based on light, food type and habitat; and even

amount of pigmentation in the characteristic red eyespots can range from large bright

red slashes at the frontal end to non-existence depending on light intensity (Luther

1960; Bauchhenss 1971; Heitkamp 1982; Atherton & Jondelius 2018b). DNA

taxonomy provides a robust, reproducible and relatively easy method of identification

that can also account for the hidden biodiversity of cryptic species (Pérez-Ponce de

León & Poulin 2016; Fonseca et al. 2017).

The results of the bPTP analyses on the concatenated 18S and CO1 datasets

(Fig. 2) were able to reliably separate the species of Microstomidae and supported the

more traditional morphological delineation. Support for four species (*M. crildensis,*

M. laurae, M. lotti, M. septentrionale) was consistent but low (0.49-0.68) and is

unsurprising given the overall low intraspecific genetic diversity within each

Microstomum species. Support was moderate to high (≥0.79) for all other species.

Within the Microstomidae, our results found all limnic species (*M. artoisi, M.*

bispiralis, M. lineare, M. tchaikovskyi, M. zicklerorum) grouped together in a single

clade with high support (0.91), indicating a single instance of freshwater invasion

from marine habitats. The transition of marine to freshwaters within a lineage is a

common occurrence for many different animal taxa (Lee & Bell 1999, Davis et al.

2012) and has occurred independently at least five other times within the

Platyhelminthes, including within Macrostomidae (Schockaert et al. 2008). Currently,

nine species of Microstomidae have been reported exclusively from freshwaters while

one (*M. lineare*) inhabits both limnic and brackish waters to 7‰. Further broad

sampling efforts in different parts of the world will be required to illuminate the

origin and distribution patterns of freshwater Microstomidae.

Morphological Taxonomy

Morphological identification of flatworms in general is often dependent on the

sexual anatomy, and species identification of Macrostomorpha specifically is heavily

or entirely based on the morphology of the reproductive organs (e.g. Nasonov 1935;

Ax & Armonies 1987; Faubel & Cameron 2001). Some authors have argued that

species of Microstomidae with unknown sexual anatomy are incompletely described

and should not be considered valid (Faubel 1984). Unfortunately, specimens of

Microstomidae are most often immature when collected due to very short seasons of

sexual reproduction, and thus scientists are often reluctant to describe new species

(e.g. Yanoviak 2001; Delp 2002; Janssen et al. 2015). Ignoring or invalidating

immature Microstomidae leads to large parts of biodiversity left unexplored in a

group of animals that are ubiquitous, abundant (e.g. Martens & Schockaert 1981,

Reise 1984) and ecologically important (Zeppilli et al. 2015) and prevents the

discovered taxa from being included in future research or conservation efforts (Jörger

& Schrödl 2013). Further, invalidating asexually reproducing immature species

assumes that all species must have sexual anatomy, an assumption that is not

necessarily legitimate. Instances of obligate asexuality are widespread throughout the

animal kingdom (reviewed in Schön et al. 2009), and animal taxa already possessing

the ability to reproduce asexually can give rise to asexual lineages without sexual

organs, as has been demonstrated in e.g. aphids (Simon et al. 2002), ostracods (Butlin

et al. 1998) as well as numerous other meio- and macrofaunal taxa (Simon et al.

2003).

In order to facilitate future research, we created a key to Microstomidae that is

not dependent on the reproductive system so that species may be distinguished even

when immature. As the taxonomic literature is often old and may be difficult to

acquire, we further summarize the morphological and molecular taxonomy of all

currently accepted species within Microstomidae. Important distinguishing

morphological characters include: shape of anterior end (blunt, conical, proboscis,

rounded; Fig. 11); shape of posterior end (blunt, pointed, rounded, spatulate, tail-like;

Fig. 12); size and shape of ciliary pits (rounded, bottle-shaped; Fig. 13); length of

preoral intestine (defined as long if extending anterior to the brain, medium if

reaching to level of brain, and short if terminating posterior to level of brain; Fig. 14);

body pigmentation; presence, location and color of pigmented eyespots; presence and

location of rhabdites; presence, pattern and location of adhesive papillae; as well as

other characters specific to individual species (e.g. the lateral intestinal bulges of *M.*

mundum; the biparte pharynx of *M. breviceps*; Fig. 15). Table S3 summarizes the

morphological characters for each species.

Taxonomic Summary

911   **Fam. Microstomidae Luther, 1907**

912   Macrostomida with intestine extending anterior to the mouth (preoral

913   intestine). Ciliated sensory pits present at frontal end, typically around level of brain.

914 Frontal glands present surrounding pharynx. Asexual reproduction through fissioning

915 occurring. Marine, brackish and freshwater habitats; two genera.

916   Type: *Microstomum* SCHMIDT 1848

918   **Gen. *Microstomum* SCHMIDT 1848**

Microstomidae with ciliary pits at frontal end and preoral intestine. Two

excretion openings in the anterior body. Rhabdites and nematocysts present in some

species. Asexual reproduction through fissioning as well as sexual reproduction

present. Sexual anatomy including one or two testes beside or behind the intestine;

one or two ovaries consisting of one or more lobes and containing a central oocyte

and a peripheral layer of adipose cells. Body size of solitary individual 0.6-4mm, with

chains comprising up to 18 zooids and 15 mm.

Type species: *Microstomum lineare* (Müller, 1773).

43 species: 33 marine, 9 limnic, and 1 inhabiting both brackish (up to 7‰) and

freshwaters.

Species Key:

1.	Pigmented Eyes Present	2.
	Pigmented Eyes Absent	13.
2.	Pigmented eyes red	3.
	Pigmented eyes dark, not red	9.
3.	Adhesive papillae present at the posterior end of the animal	4.
	Posterior adhesive papillae not present	7.

4.	Adhesive papillae present at the anterior end of the animal	5.
	Anterior adhesive papillae not present	6.
5.	Eyespots marginal dots close to anterior tip; body color reddish; preoral intestine to level of brain	*M. gabriellae*
	Field of red pigmentation forming broad transverse band at anterior end; preoral intestine small, not reaching to brain	*M. rubromaculatum*
6.	Paired red eyespots as small dots located at lateral margins of the body at level of brain, posterior to small ciliary pits	*M. laurae*
	Single medial eyespot, oval shaped, located anterior to the brain; intestine with lateral lobules and bulges	*M. groenlandicum*
7.	Eyespots present as red stripes at dorso-anterior end	*M. lineare* complex**
	Eyespot not so	8.
8.	Body color yellow to orange; eyespots at level of ciliary pits; rhabdites present concentrated at body ends	*M. spiriferum*
	Body yellow-green with two small eyespots at frontal margins; posterior end with stiff cilia twice as long as locomotory cilia; rhabdites present across body	*M. bioculatum*
9.	Anterior end enlarged to form a proboscis with papillae and sensory cilia	10.
	Without anterior proboscis	11.
10.	Sensory cilia even spaced along lateral margins of body; eyespots posterior to mouth	*M. proliferum*

	Without evenly spaced lateral sensory cilia; eyespots present just anterior to mouth	*M. compositum*
11.	Preoral intestine short, not quite reaching level of brain; ciliary pits small; without adhesive papillae	*M. dermophthalmum*
	Not so	12.
12.	Eyes with distinct crescent lens present at frontal end; posterior end tapering and rounded; marine species	*M. melanophthalmum*
	Eyes with distinct crescent lens present anterior to ciliary pits; posterior end tail-like; freshwater species	*M. paradii*
13.	Adhesive papillae present at the anterior end of the animal	14.
	Anterior adhesive papillae not present	28.
14.	Posterior end forming a spatulate tail plate set off by a constriction in the body width	15.
	Posterior end not so	17.
15.	Ciliary pits small and shallow; preoral intestine reaching anterior to brain; pharynx biparte with longer anterior cilia and shorter posterior	*M. breviceps*
	Not so	16.
16.	Adhesive papillae on frontal end situated in a row ~30 μm from the anterior tip of the animal	*M. lotti*
	Rhabdites heavily concentrated at anterior end above pharynx	*M. ulum*
17.	Preoral intestine short, terminating posterior to level of brain	18.

	Preoral intestine extending to brain or anterior	22.
18.	Rhabdites present	19.
	Rhabdites absent	21.
19.	Anterior end enlarged to form a proboscis, non ciliated but with sensory bristles	M. album
	Anterior end not so	20.
20.	Anterior end bluntly rounded; field of red pigmentation forming broad transverse band at anterior end	M. rubromaculatum
	Body evenly light reddish-brown; anterior end rounded and set apart by slight constriction at ciliary pits	M. schultei
21.	Body color brownish yellow with numerous anterior black dots; anterior end pointed and separated by distinct constriction at ciliary pits	M. punctatum
	Body colorless; anterior end long and rounded; with numerous adhesive papillae on posterior end	M. marisrubri
22.	Preoral intestine extending anterior to brain	23.
	Preoral intestine terminating at level of brain	25.
23.	Ciliary pits clearly discernable and bottle-shaped; adhesive papillae present in a row ventrally, ~25 μm from the anterior tip	M. weberi
	Not so	24.
24.	Rhabdites very numerous at frontal end; adhesive papillae scattered along margins of entire body	M. papillosum
	Rhabdites only sparsely present; few adhesive papillae present on anterior and posterior ends only	M. afzelii

25.	Rhabdites present throughout body and extremely numerous at anterior end	26.
	Rhabdites not present or present at anterior ends only	27.
26.	Body colorless; field of adhesive papillae present at anterior tip of animal, longest at lateral sides and sloping upwards medially	M. westbladi
	Body yellow with darker yellow to orange frontal end; adhesive papillae sparsely scattered over entire body	M. septentrionale
27.	Adhesive papillae present at both body ends; pharynx biparte with longer anterior and shorter posterior cilia; paired stripes of rhabdite bundles on dorsolateral sides	M. rhabdotum
28.	Rhabdites present	29.
	Rhabdites absent	37.
29.	Posterior end tail-like	30.
	Posterior end rounded or blunt, not tail-like	31.
30.	Rhammite glands colored red; cilliary pits deep and narrow	M. jenseni
	Intestine with lateral bulges and smaller lobules	M. mundum
31.	Ciliary pits large and very distinct, bottle shaped	32.
	Ciliary pits not so	33.
32.	With ochre frontal end; preoral intestine very short, hardly extending past mouth	M. curinigalettii
	Body colorless; preoral gut extending anterior to brain	M. edmondi

33.	Preoral intestine extending anterior to brain	34.
	Preoral intestine terminating at level of brain	36.
34.	Pharynx divided into two parts by annular fold with longer anterior and shorter posterior cilia	*M. davenporti*
	Pharynx not so	35.
35.	Body red with yellow rhammite glands	*M. mortenseni*
	Body colorless; rhabdites concentrated at anterior end	*M. spiculifer*
36.	Anterior end long and tapered, resembling a proboscis; body white	*M. trichotum*
	Body colorless and may have ± dark grey pigment spots in some individuals; anterior end bluntly rounded	*M. lucidum*
37.	Preoral intestine extending anterior to brain	38.
	Preoral intestine terminating at or before level of brain	40.
38.	Adhesive papillae present at posterior tip and along up to ~1/5 of the posterior body	*M. inexcultus*
	Adhesive papillae absent	39.
39.	Ciliary pits deep; anterior end conically pointed; posterior end tail-like with long sensory cilia	*M. canum*
	Body ends end blunt; bluish color	*M. caerulescens*
40.	Rounded posterior end with adhesive papillae	*M. crildensis*
	Posterior adhesive papillae not present	41.

41.	Preoral intestine reaching to level of brain	*M. lineare*
	Preoral intestine terminating before level of brain	42.
42.	Animal comparatively large, single zooid to 4 mm, 18.5 mm maximum for 8 zooids; posterior end rounded or with thin-tail	*M. giganteum*
	Maximum body length 4 mm, 8 zooids; body transparent grey to light yellow with tapered pointed posterior end	*M. bispiralis*

931
932   ***M. lineare* complex refers to the species that are distinguished from *M. lineare*
933   primarily on the basis of DNA information, specifically *M. artoisi*, *M. tchaikovskyi*,
934   *M. zicklerorum* (Atherton & Jondelius, 2018b)
935

List of Currently Recognized Microstomum *Species in Alphabetical Order*

1. *Microstomum album* (ATTEM 1896)

Synonyms

Alaurina alba Attems 1896, pg 56

Type: Not recorded

Type Locality: North Sea, Helgoland

Habitat: Marine, pelagic, phytal

Brinkmann 1905 – Denmark, Frederikshavn in Kattegat

Graff 1913 – France, Gulf of Lion, Banyuls sur Mer

Body length of solitary animal to 1.6 mm, chains to 2.5 mm, 4 zooids. Body

colorless or white, displaying yellow-orange intestine. Pigmented eyes absent.

Anterior end smoothly tapering to conically pointed proboscis, non-ciliated but with

tactile bristles and a ring of papillae. Posterior end rounded. Small ciliary pits present

slightly anterior to mouth. Rhabdites present, numerous over entire body. Preoral

intestine short. Adhesive papillae sparsely present along entire mid to posterior body

and highly concentrated at posterior end.

Reproductive system with single testis and ovary. Stylet arched projecting into

antrum masculinum. True prostatic vesicle present in addition to seminal vesicle

present. Female system with ovary and short duct. Gonopores separate.

2. *Microstomum afzelii* sp. nov.; Fig. 3

Type: Asexual, MTP LS 700; GenBank accession 18S KP730494

Type Locality: Sweden, Saltö, 58°52'29''N 11°08'41'' E, 10 cm

Habitat: Marine, eulittoral sand

GenBank Accession:

18S – KP730494

Body length 0.8 mm, 2 zooids. Body colorless, without pigmented eyespots.

Body shape elliptical with rounded anterior and posterior ends. Ciliary pits weakly

present, small and difficult to discern. Adhesive papillae and rhabdites sparsely

present. Nematocysts particularly concentrated at posterior end. Preoral intestine long,

extending anterior to brain.

Reproductive system unknown.

3. *Microstomum artoisi* ATHERTON & JONDELIUS 2018; Fig. 17D

Type: Asexual, SMNH-Type-9022; GenBank accession 18S MH221276, CO1

MH221384, ITS MH221478

Type locality: Belgium, Hoge Kemper Park, 51°00'03''N 5°40'44''E, 10 cm depth

Habitat: Freshwater phytal, 5-10 cm depth

GenBank Accession:

18S – MH221275 – MH221276

CO1 – MH221383 – MH221384

ITS – MH221477 – MH221478

Molecular Diagnosis:

COI gene region with reference to GenBank Accession MH221288: 39-A; 42-

G; 57-A; 81-C; 114-A; 180-T; 195-C; 207-C; 216-G; 234-C; 240-G; 321-G; 390-C;

409-A; 612-T

ITS1, 5.8S, ITS-2 gene region with reference to GenBank Accession

MH221390: 56-A; 383-T; 839-A; 961-T

Morphological Description:

Body length to approximately 3.5 mm, maximum 6 zooids. Body color

greyish or slightly tinged pinkish clear. Anterior end abruptly conical, posterior end

tail-like without adhesive papillae. Sensory bristles present and ciliary pits present.

Two red eyestripes present at dorso-anterior end. Amount of eyespot pigmentation in

some individuals very bright, almost merging medially to form a dorsal. Preoral

intestine short.

Reproductive system not seen.

1004

1005 Notes: See notes of *M. lineare*.

1006

1007

1008 **4. *Microstomum bioculatum* FAUBEL, 1984**

1009 Type: Not deposited

1010 Type locality: North Sea, Sylt Island, off Keitum

1011 Habitat: Marine, eulittoral and superlittoral lentic upper mudflats and salt meadows.

1012

1013   Body length to 1.4 mm, maximum 4 zooids. Body yellow-green with two very

1014 small red eyespots at frontal margins. Ends rounded; posterior with long, stiff cilia

1015 that are twice the length of the locomotory cilia. Rhabdites present. Ciliated pits,

1016 preoral intestine and adhesive papillae all apparently absent.

1017   Reproductive system not seen.

1018

Notes: The absence of the preoral intestine and, especially, ciliated pits is extremely

unusual, as both are defining characters of the genus. Additionally, there seems to be

some confusion regarding the preoral intestine, since the original written description

by Faubel (1984) specifically precludes its presence ("Ein Kopfdarm und typische

Einschnürungen des Körpers in Höhe des Gehirns fehlen"), but Faubel's original

drawing of the species distinctly includes a preoral intestine that extends anterior to

the brain.

1026

1027

1028   **5. *Microstomum bispiralis* STIREWALT, 1937; Figs. 16, 17E**

1029   Type: Deposition not recorded

1030   Type locality: near University of Virginia, Virginia, USA

1031   Habitat: Freshwater pools

1032

1033   This paper: Sweden, Finland

1034

1035   GenBank Accession:

1036   18S – XXX

1037   CO1 – XXX

1038

Maximum 8 zooids, 4 most common, with body length to 4 mm. Following

laboratory cultures, individuals can be transparent-grey to bright yellow. Pigmented

eyespots absent. Anterior end short conically pointed. Posterior end tapered to a point.

Ciliary pits present anterior to mouth, each shaped as a tube extending posteriorly at

45° angle and ending in an enlarged spherical pocket. Pharynx ciliated, and occupies

⅓ to ½ length of anterior zooid. Preoral intestine short. Adhesive papillae and

rhabdites absent. Nematocysts may be present in some specimens.

Nervous system and cerebral ganglion as typical for the genus; consisting of

two anterior ganglia connected by a short and wide transverse commissure. A

dorsolateral and a ventrolateral nerve extend posteriorly from each ganglion.

Ventrolateral nerves are connected by a commissure ventral to the pharynx. One

nerve extends laterally from each ganglion to the ciliary pit and another from each

ganglion to branch profusely in the anterior end of the body.

Protanderous hermaphrodites, with male formation occurring in laboratory

cultures during January and February. Male genital system consisting of two compact

and spherical testes, one developing earlier than the other. Vasa deferentia extending

posteriorly from each testis to empty dorsolaterally into the vesicula seminalis.

Vesicula seminalis shaped as a small sphere, located ventral and posterior to the

intestine; glandular and surrounded by a heavy muscular sheath of the copulatory

organ. Copulatory stylet a smooth spiral sharply bent in two planes, 120 µm long,

widened at the base and drawn out distally to a sharp point.

Female reproductive system includes a single ovary lying slightly anterior to

the middle of the zooid, ventral to the intestine. Numerous single-celled glands open

to the vagina, which opens to a female pore at the beginning of the posterior ⅓ of the

zooid.

Somatic chromosome number 16 (n=8)

1065

Notes: This is the first time that *M. bispiralis* is reported from Sweden or anywhere

else outside of the original location. The specimens collected in Finland and Sweden

matched the original descriptions, including the male stylet and other sexual anatomy

that was found in three individuals (Fig. 16C). The very great distance between the

locations does suggest the possibility of cryptic species or much larger dispersal

abilities than has previously been shown in any species of *Microstomum* (see

Atherton & Jondelius 2018b), although true support for either scenario must be

deferred until DNA sequence data can be attained from specimens collected at the

type locality.

The absence of adhesive papillae and rhabdites combined with a short preoral

intestine is a unique set of characters that separates this species from nearly all of the

other currently accepted *Microstomum* species. *M. bispiralis* is most similar to *M.*

lineare and *M. lineare*-like or cryptic species (including *M. artoisi, M. giganteum, M.*

tchaikovskyi, and *M. zicklerorum*), which in part is due to the very wide

morphological variability reported for *M. lineare*. However, *M. bispiralis* is typically

smaller with a slightly higher body width:length ratio, has a shorter preoral intestine,

and always and entirely lack the pigmented eyestripes. Further, all species within the

M. lineare complex are separated through 18S and CO1 gene evidence (see Figs. 1,

2), while the shape and size of the male stylet is different between *M. bispiralis, M.*

giganteum and *M. lineare*.

1086

1087

1088 **6. *Microstomum breviceps* MARCUS, 1951**

1089 Lectotype: Asexual, sectioned, SMNH-Type-XXXX; E. Marcus leg.**

1090 Type locality: Brazil, São Sebastião

1091 Habitat: Marine, between algae, mainly *Sargassum stenophyllum*

1092

Maximum 3 zooids, 0.7 mm. Color and pigmented eyes unknown as worms

were only described based on preserved state. Anterior end rounded, posterior

spatulate. Adhesive papillae present anteroventrally above level of the brain, at the

posterior tip and in the zone of division in mature zooids. Ciliated pits small.

Rhabdites occur throughout the body. Preoral intestine extends anterior to brain.

Mouth circular. Pharynx divided into two parts by an annular fold, with longer cilia in

the anterior part and shorter cilia in the posterior. Glands around the pharynx may be

pale pink in color. Intestine unciliated.

Reproductive organs not seen.

1102

Notes: *Microstomum breviceps* is similar to *M. davenporti* in that both have similar

patterns of adhesive papillae, a long preoral intestine and a characteristic divided

pharynx with long and short cilia and with an annular fold. *M. breviceps* can be

separated from *M. davenporti* by its body shape: the anterior end of *M. breviceps* is

truly round instead of tapering blunt, and the posterior spatulate tail-plate is very

distinct from the rest of the body. Other differences can be found in the high

concentration of rhabdites surrounding the oval mouth in *M. davenporti*, and the very

large distances separating their locations (São Sebastião, Brazil and Woods Hole

Massachusetts, USA for *M. breviceps* and *M. davenporti*, respectively).

**A type specimen was not previously designated for *M. breviceps*. However, two

original specimens were deposited at the Swedish Museum of Natural History, and

thus, following the rules of the International Code of Zoological Nomenclature, we

hereby designate one sectioned asexual specimen (formerly SMNH108136, now

SMNH-Type-XXXX) as lectotype and one whole mount asexual specimen

(SMNH108137) as paralectotype for *Microstomum breviceps* Marcus, 1951.

1118

1119

7. *Microstomum caerulescens* (Schmarda, 1859)

Synonyms:

Strongylostomum caerulescens - Schmarda 1859

Strongylostomum coerulescens - Parádi 1881, pg 311; Diesing 1862, pg 235; Graff

1882, pg 253

Typhlomicrostomum coerulescens - Diesing 1862

Microstomum coerulescens - Graff 1882, pg 253

1127

1128   Type: Not recorded

1129   Type locality: Jamaica, Freshwater lakes near Kingston

1130   Habitat: Freshwater lakes

1131

1132   Body small, 0.67 mm; two zooids. Anterior and posterior ends blunt, almost

1133   flat. Bluish color; eyespots not recorded. Preoral intestine extends to close to the

1134   anterior end. Adhesive papillae, rhabdites and ciliary pits not recorded.

1135   Reproductive system not observed.

1136

Notes: Ciliary pits were not described for this species, nor were they included in

Schmarda's (1859) drawing. However, the original description occurred at a very

early time period, and positively identifying ciliary pits may be difficult even with

modern equipment when they are small and/or shallow. The importance of the

character for this genus (as first argued by Uljanin 1870; Graff, 1882) suggests that

the presence or absence of ciliary pits for this species should remain in question until

they can be re-examined. Even without absolute knowledge of the adhesive papillae,

ciliary pits, and reproductive system, the combination of blue body color, absence of

pigmented eyespots, large pre-oral intestine, blunt body ends and freshwater habitat is

enough to separate this species from all other currently described species of

Microstomum.

1148

1149

1150   **8. *Microstomum canum* (FUHRMANN 1894)**

1151   Synonyms:

1152   *Microstoma canum* Fuhrmann 1894, pg 232

1153

1154   Type: Not recorded

1155   Type locality: Switzerland, Basel, Augustinerholzbach

1156   Habitat: Freshwater

1157

1158   Graff 1913 – France

1159

1160   Body length to 2.0 mm, 8 zooids. Body grey. Eyes absent. Anterior end long

1161   and tapering to a point. Posterior end pointed, with long cilia but without adhesive

1162   papillae. Ciliary pits deep, present at level of mouth, surrounded by long cilia. Preoral

1163   intestine extends anterior to brain. Rhabdites and nematocysts not seen.

1164   Reproductive system unknown.

1165

Notes: Hofsten (1912) and Meixner (1915) postulated that *M. canum* is a synonym of

M. lineare based on variability of the latter. While the wide morphological variations

of *M. lineare* undoubtedly creates difficulty in absolutely ruling out synonymy,

potential differences between the two species exist in the extension of the preoral

intestine anterior to the brain, a more posterior position of the ciliary pits and,

perhaps, a more pointed anterior end (based Furhmann's original drawing) for *M.*

canum.

1173

1174

9. *Microstomum compositum* (METSCHNIKOV 1865); Fig. 18F

Synonyms:

Alaurina composita Metschnikov 1865, pg 178

Alauretta viridirostrum Mereschkowsky 1879, pg 35

Alaurina viridirostrum Graff 1882, pg 261

Alaurina claparedii Graff 1882, pg 262

Dinophilis simplex Verrill 1892, pg 485

1182

1183   Type: deposition not recorded

1184   Type Locality: Germany, Helgoland, Coast of Skye-Hebrides

1185   Habitat: Marine pelagic or planktonic

1186

Brinkmann 1905 – Denmark, Skagens Rev, Horns Rev

Gamble 1893 – United Kingdom, Island of Skye

Graff 1913 – United States, Rhode Isalnd, Newport

Hofker 1930 – Netherlands, Den Helder, IJsselmeer

This paper: Sweden, Fiskebäckskil, Kristineberg Sven Lovén Center for Marine

Research, 58°14′59′′ N, 11°26′45′′

1193

1194   GenBank Sequences

1195   18S – XXX

1196   CO1 – XXX

1197

Body length of solitary animal to 0.9 mm, chains to 3.0 mm. Maximum 10

zooids, 4 most common. Body color light yellow with greenish anterior (proboscis).

Black pigmented eyespots present. Anterior end sharply decreasing in width to form

distinct proboscis, with sensory cilia, papillae and numerous circular folds, and

denoted by paired transverse tufts of cilia at base. Posterior tip with thin tail and

terminal sensory cilia. Preoral intestine extending anterior to brain. Intestine non-

ciliated, often with diatoms and protozoa. Rhabdites may be present in the anterior

end.

Male reproductive system including a single testis, which may be bilobed in

some specimens. Stylet weakly arched connecting to seminal vesicle.

Female system unknown.

1209

1210   Notes: Brinkmann (1905, pg 66) argued that both *Alaurina viridirostrum* and *A.*

1211   *claparedii* were identical to this species, stating that the very few differences were

1212   due to errors derived from outdated methods of investigation and/or a lack of

1213   sufficient specimens.

1214

1215

1216   **10. *Microstomum crildensis* FAUBEL, 1984; Fig. 18E**

1217   Type: Deposition not recorded

1218   Type locality: Germany, Jade Busen, Crildumer Siel

1219   Habitat: Marine, eulittoral

1220

1221   This paper: Sweden, Fiskebäckskil, Kristineberg Sven Lovén Center for Marine

1222   Research, 58°14′59″ N, 11°26′45″; Stömstad, Saltö, 58°52′29′′N 11°08′41′′ E

1223

1224   GenBank Accession:

1225   18S – XXX

1226   CO1 – XXX

1227

Body to 1.5 mm, maximum 4 zooids. Solitary animal 0.9 mm. Body

colorless; pigmented eyes absent. Ends rounded, posterior with adhesive papillae.

Constriction at level of brain at ciliary pits. Preoral intestine short, not quite reaching

level of brain. Rhabdites and nematocysts absent.

Single testis, located at the very caudal end of the animal. Vesicula

granulorum connected to a 32 μm long stylet, shaped as a curved tube of

approximately consistent width and with distal terminal opening. Single ovary

present.

1236

Notes: Three specimens identified as *Microstomum crildensis* were collected from

Swedish Skagerrak coast. Unfortunately, none of these specimens displayed sexual

organs, and so the reproductive system—most importantly the male stylet—could not

be used to corroborate identification. However, in every other respect, these

specimens matched the original description and drawings, including: overall body

shape including a body constriction at the ciliary pits, presence adhesive tubes on the

posterior end only, absence of rhabdites, and small preoral intestine. The habitat—

eulittoral muddy sediments of the North Sea—was also consistent.

Currently, the only other species of *Microstomum* with posterior adhesive

papillae that entirely lacks rhabdites are *M. inexcultus* sp. nov., *M. marisrubri* sp. nov.

and *M. punctatum.* M. crildensis can be distinguished from each of these species

through:

1249

M. inexcultus sp. nov. – long preoral intestine; constriction of the anterior body width

near the ciliary pits; see also Figs. 1, 2

M. marisrubri. sp. nov. – shape and size of the male stylet; presence of anterior

adhesive papillae and more numerous posterior adhesive papillae; see also Figs. 1, 2

M. punctatum – brown-yellow body coloring with anterior black spots; tail-like

posterior end; limnic habitat

1256

1257

1258 **11. *Microstomum curinigalletti* sp. nov.; Fig. 4**

1259 Type: Asexual specimen, SMNH-XXXX; GenBank Accession XXX

1260 Type locality: Sweden, Fiskebäckskil, 58°14′59″ N, 11°26′45″ E, 20 cm depth

1261 Habitat: Marine, sublittoral phytal on algae

1262

1263 GenBank Accession:

1264   18S – XXX

1265   CO1 – XXX

1266

Body length 1.4 mm, 2 zooids. Anterior body end ochre to dark yellow-

orange. Body otherwise colorless without pigmented eyespots. Anterior end conically

rounded; posterior end blunt with numerous adhesive papillae. Ciliary pits large and

bottle-shaped, located anterior to pharynx. Preoral intestine very short. Few rhabdites

scattered across the body. Nematocysts present.

Reproductive system unknown.

1273

1274

1275   **12. *Microstomum davenporti* GRAFF, 1911**

1276   Type: Deposition not recorded

1277   Type locality: Woods Hole, Massachusetts, USA and Stamford, Connecticut, USA

1278   Habitat: Marine, on *Ulva* sp. and *Sargaassum* sp.

1279

Body length to 1.5 mm; maximum 4 zooids. Body colorless or whiteish with a

yellow-ocher intestine. Eyes absent. Anterior conically pointed; posterior end broad

with numerous adhesive papillae. Ciliary pits present but very shallow. Rhabdites

present, 12 µm, highly concentrated at the anterior end and around the oval mouth.

Preoral intestine long. Pharynx divided into two parts by an annular fold, with longer

cilia in the anterior part and shorter cilia in the posterior

Reproductive system not seen.

1287

1288

1289   **13. *Microstomum dermophthalmum* RIEDEL, 1932**

1290   Type: Deposition not recorded.

1291   Type locality: Engelskmandshavn, Godhavn, Greenland

1292   Habitat: Marine, coarse sand and sandy mud, 1-15 m depth

1293

1294   Steinböck 1938—West Greenland, sand and mud from 5-10 m depth; Iceland, low

1295   tide zone, coarse sand and algae, outer harbor of Ísafjör∂ur.

1296   Riedl 1956 – Adriatic sea

1297

Small, body length to 0.32 mm in solitary animal. Anterior end short conical;

posterior end tapering to a rounded tip. Body colorless. Marginal eyes present very

near the anterior end. Ciliary pits small and indistinct, at level posterior to marginal

eyes. Rhabdites concentrated at anterior and posterior ends of the body. Adhesive

papillae not recorded. Preoral intestine not quite reaching level of brain. Pharynx with

5.4 µm long cilia. Intestine unciliated.

Paired testes, sausage-shaped, obliquely lateral and posterior to ovary.

Vesicula granulorum 20 µm long, 13 µm wide, strongly muscled. Male stylet strait

tube with distal toothed thickening; small ~10-12 µm long (based on drawing).

Antrum masculine small, spherical. Genital pores separate.

1308

1309

1310   **14. *Microstomum edmondi* ATHERTON & JONDELIUS 2018; Fig. 18A**

1311   Type: ♂, SMNH-Type-8890; GenBank accession MF185704

1312   Type Location: Sweden, Fiskebäckskil, 58°14'52''N, 11°27'05''E, 25 cm

1313   Habitat: Marine benthic, 15-25 cm depth

1314

1315   GenBank Accession:

1316   18S – XXX-XXX

1317   CO1 – MF185700 – MF185711

1318

Body length to 1.7 mm, maximum 4 zooids. Conical pointed anterior end;

blunt, almost flat posterior end with numerous 5-8 µm long adhesive papillae along

rim. Ciliary pits large, bottle shaped. Body clear. Pigmented eyes absent. Dense field

of cilia clearly covering epidermis. Preoral intestine extending anterior to brain.

Mouth distinctly encircled by glands. Large rhabdite bundles to 45 µm long, primarily

in the posterior end.

Protandrous hermaphrodite. Male reproductive system with single large testis.

Vesicula seminalis circular to elliptical, 98 µm long, and containing the ends of

numerous prostate glands in the distal part. Stylet approximately 67 µm long; shaped

as an elongate, narrow funnel, slightly curved in one plane with a short, arched tip.

Female reproductive system with single ovary and gonopore. Eggs develop caudally.

1330

1331

1332   **15. *Microstomum gabriellae* MARCUS, 1950**

1333   Lectotype: Asexual, whole mount, SMNH-Type-XXX; E. Marcus leg.**

1334   Type location: Brazil, São Sebastião

1335   Habitat: Marine, between algae, mainly *Sargassum stenophyllum*

1336

Body length to 3 mm; maximum 7 zooids. Color reddish, with anterior

marginal paired red eyespots. Anterior end short and conically rounded; posterior end

rounded. Ciliated pits are thin and deep, located at level of posterior brain. Adhesive

papillae present at the posterior of each well-developed zooid and more sparsely

along the body. Rhabdites present, scarce and throughout the body. Nematocysts not

seen. Pharyngeal nerve ring present, connected to brain by two short nerves. Preoral

intestine reaching to level of brain.

Reproductive system based on original drawings (Marcus, 1950). Male system

with paired round testes connecting separately to vesicular seminalis by vas deferens.

Stylet approximately 45-50 µm long, shaped as a tube with near consistent width

throughout, broadly curved 180°.

1348

**A type specimen was not previously designated for *M. gabriellae*. However, nine

original specimens were deposited at the Swedish Museum of Natural History, and

thus, following the rules of the International Code of Zoological Nomenclature, we

hereby designate one whole-mount asexual specimen (formerly SMNH108140, now

SMNH-Type-XXXX) as lectotype. The other eight originally deposited specimens

(SMNH108138-9, SMNH108141-6) are thus designated as paralectotypes.

1355

1356

1357   **16. *Microstomum giganteum* HALLEZ 1878**

1358   Type: Not recorded

1359   Type Location: France, Lille (?)

1360   Habitat: Freshwater in Europe, far less common than *M. lineare*

1361

Hallez 1878, 1879 – France, Lille, Wimereux

Keller 1894 – Switzerland

Zabusov 1894 – Russia, Kazan

Zykov 1897 – Russia, in the viscidity of Moscow

Volz 1898 – Switzerland

Dorner 1902 – Germany, Königsberg

Markov 1904 – Russia, Kharkiv region, Udy River

Graff 1909 – Germany

1913 – Scotland, Denmark

Nasanov 1919 – Kirov oblast

Southern 1936 – Ireland, Clare, shore of Lough Atorick

Kostenko 1988 – Ukraine, Kiev, Okrung Reservoirs

Rogozin 2015 – Lipovka River falling into Lake Ilmen, 55°00′42″ N, 60°09′20″ E;

Lake Bol'shoe Miassovo, 55°10′39″ N, 60°17′31″ E

1376

1377   Body large, single zooid to 4 mm, max length 18.5 mm for 8 zooids. Body

1378   whitish, grey or colorless with brown-red intestine. Pigmented eyespots absent.

1379   Anterior end rounded; posterior ends rounded with a thin tail. Adhesive papillae

1380   absent. Preoral intestine extending to anterior pharynx or very slightly past.

1381   Nematocysts present.

1382   Male copulatory organ with an oblong seminal vesicle, 150-180 µm long.

1383   Stylet with three parts: 85 µm distal part inserted in a 30 µm mid part which is

1384   inserted in turn into a 40 µm long proximal part. Total length of the stylet to 180 µm,

1385   shaped as a spiral with 2.5 full turns. Caudal ovary with three follicles with triangular

1386   ovicels and 4-5 abortive eggs.

1387

Notes: There was much discussion in the literature about whether *Microstomum*

giganetum should be considered a "safe species" or a larger variant of *M. lineare*. The

species name was reduced to a synonym by Luther (1960), who considered it to

represent a pure line of *M. lineare* that had developed over a long period of asexual

reproduction. *M. giganteum* eventually regained its independence after Kostenko

(1988) carried out a comparative study of both forms and found several

morphological differences, most importantly in the different structures of the

reproductive system and male stylet, without intermediate or transitional forms.

M. giganteum differs from *M. lineare* through its larger size (8 zooid body

length of ≥8.5 mm compared with 4-5 mm, respectively); lack of elongated tail,

adhesive papillae and pigmented eyespots; pharyngeal glands restricted to oral cavity;

and reproductive system including larger male stylet with three parts forming greater

number of spiral turns (2.5 versus 1.5, respectively).

1401

1402

1403   **17. *Microstomum groenlandicum* (LEVINSEN, 1879)**

1404   Synonyms:

1405   *Microstoma groenlandicum* Levinsen 1879, pg 449

1406

1407   Type: Deposition not recorded

1408   Type Locality: Davis Strain, Disko Island, Greenland

1409   Habitat: Marine, among *Ulva* sp.

1410

Gamble 1893 – United Kingdom, English Channel

Graff 1882 – Greenland, Baffins Bay

Graff 1905, 1913 – Norway, Bergen

Meixner 1938 – United Kingdom, English Chanel

Southern 1912 – Ireland, Clare Island

Southern 1936 – Ireland, Blacksod Bay, New Harbor

1417

Body length to 2 mm, with maximum 8 zoods. Body colorless, yellow or

reddish. A single medial red eyespot anterior to the brain, oval shaped. Anterior tip of

the body conically pointed. Posterior end long, tail-like with many adhesive papillae.

Rhabdites present throughout the body, concentrated at the anterior end. Small ciliary

pits located at level between the brain and the mouth. Preoral intestine extending to

the brain. Intestine with lateral lobules and bulges.

Testes not seen. Male stylet narrow, cylindrical and weakly spiraling, ending

with a small, spoon-shaped thickening. Single medial ovary.

1426

1427

1428 **18. *Microstomum inexcultus* sp. nov.; Fig. 5**

1429 Type: Asexual specimen; SMNH-Type-XXXX; GenBank Accession XXX

1430 Type Locality: Sweden, Saltö, 58°52'29''N 11°08'41'' E,10 cm depth,

1431 Habitat: Marine, eulittoral sand

1432

1433 GenBank Accession:

1434   18S – XXX

1435   CO1 – XXX

1436

1437   Body length 0.85, 2 zooids. Colorless and without pigmented eyespots.

1438 Anterior end pointed; posterior tapered to a blunt tip with numerous adhesive papillae.

1439 Rhabdites absent. Ciliary pits clearly present. Preoral intestine long, stretching almost

1440 to anterior body end.

1441   Reproductive system unknown.

1442

1443

1444 **19. *Microstomum jenseni* RIEDEL, 1932**

1445 Synonyms:

1446   *Microstomum tortipenis* Steinböck, 1938

1447

1448   Type: Deposition not recorded

1449   Type locality: Engelskmandshavn, Godhavn, Disko Island, Davis Strain

1450   Habitat: Marine benthic, to 300 m

1451

1452   Steinböck 1933 – Iceland, Isafjördur, fine sand, shallow water

1453   Westblad 1953, 1954 – Scandinavian west coast, Gullmar Fjord, 60-80 m depth, mud;

1454   Herdla, Kværnepollen, 40 m, mud

1455   Faubel 1974, 1977 – Germany, North Frisian Islands, Sylt

1456   Faubel and Warwick 2005 – United Kingdom, Scilly, St. Martin and surrounding

1457

Body length around 0.6 mm. Colorless or whiteish with red-yellow intestine.

Anterior conically pointed; posterior end long and tail-like with adhesive papillae.

Rhabdites evenly distributed over body. Rhammite glands colored red at anterior end.

Ciliary pits deep and narrow. Preoral intestine extending to level of brain. Intestine

lightly ciliated or unciliated, with minotian cells. Nematocysts present.

Paired testes rounded to club-shaped. Paired vasa deferentia separately

connected to vesicula granulorum. Male stylet 80 µm, shaped as an elongate weak

spiral. Ovary unpaired

1466

1467

1468   **20. *Microstomum laurae* ATHERTON & JONDELIUS 2018; Fig. 18B**

1469   Type: Asexual, SMNH-Type-8903, GenBank accession MF185712

1470   Type location: Strömstad, Saltö, 58°52'29''N, 11°08'41''E, 10 cm depth

1471   Habitat: Marine eulittoral

1472

1473   GenBank Accession

1474   18S – XXX

1475   CO1 – MF185712-MF185713

1476

Body length to 760 µm, 2 zooids. Body colorless and reflective of orange-red

intestine. Anterior and posterior ends rounded. Posterior rim with approximately

twenty adhesive papillae. Paired red eyespots 43 µm long, located in the lateral

margins at level of the brain. Ciliary pits very small and shallow, directly below

eyespots. Rhabdites concentrated in the anterior end above the pharynx. Nematocysts

present. Preoral intestine extending anterior to brain.

Reproductive system unknown.

1484

1485

21. *Microstomum lineare* (MÜLLER, 1773); Fig. 17A

Synonyms:

Fasciola linearis – Müller, 1773, pg 67

Planaria linearis – Müller, 1776, pg 223

Planaria vulgaris – Fabricius, 1826, pg 18

Derostoma flavicans – Ehrenberg 1831

Derostoma lineare – Duges, 1828, pg 141

Microstoma lineare – Örsted 1843, pg 566

Smigrostoma littoral – Örsted 1845, pg 417

Anotocelis linearis – Diesing, 1862, pg 237

Anotocelis flavicans – Diesing, 1862, pg 242

Microstomum caudatum – Leidy 1852, pg 350

Microstomum inerme – Zacharias, 1894, pg 83

Microstomum littorale – Diesing 1861, pg 242

1500

1501   Type: No deposition recorded

1502   Type Locality: Unknown

1503   Habitat: Freshwater and brackish water to 7‰ salinity; littoral benthal or phytal; to

1504   55+ m depth

1505

An Der Lan 1939, 1961 –Macedonia, Lake Ohrid, springs, marshy banks, mud, to 33

m depth

1962 – Hungary, Danube

Atherton & Jondelius 2018b – Sweden, Finland, Belgium, benthic or phytal

Ax 1957 – Germany, Elbe, littoral

Bauchhenss 1971 – Germany, Franconia, Nuremberg and surrounding areas

Beklemishev 1921 – Russia, Saint Petersburg

Braccini & Leal-Zanchet 1913 – Brazil

Braun 1885 – Latvia, Estonia

Brusa 2006 – Argentina, Buenos Aires, Atalaya and Punta Piedras, benthic and phytal

of rivers

Chodorowski 1959 – Poland, Harsz Lake

Dorner 1902 – Poland (Prussia)

Ferguson et al. 1939 – United States, Virginia, Mountain Lake Biological Station

Fuhrmann 1894 – Switzerland, Basel

Fulinski & Szynal 1932 – Ukraine, Poland, Lviv, Gródek

Gamo & Norena 1998 – Spain, Iberian Peninsula

Gieysztor 1926 – Poland, Warsaw, Drozdowice, Zacisze

Graff 1911 – United States, Canada, Lake Ontario, on algae in puddles

1913 – United States, New Jersey, Michigan, Pennsylvania; Denmark

Hartog 1977 – Ireland, Cork, Bennet's Lough

Heitkamp 1979, 1982 – Germany, Lower Saxony, ponds between Göttingen and

Weser

Hofsten 1907 – Switzerland, Bern

1911 – Switzerland

Kaiser 1967, 1969 – Germany, Mühlhausen

1974 – Germany, Thuringia

Kolasa 1979, 1983 – Poland, Lake Zbe□chy, surrounding rivers and streams

Kolasa et al. 1987 – United States, New York, streams around Millbrook

Korgina, 2011 – Russia, Uvod' Reservoir

Kosler 1962 – Germany, Hiddensee, benthic

Kraus 1965 – Austria, Neusiedlersee, Seewinkel National Park

Luther 1960 – Sweden, Denmark, Norway, lakes, rivers, Baltic sea, vegetation, sand,

mud, to 30+ m

Mack-Fira□ 1968 – Romania, Scurtu, Filipoiu Canal

1973, 1974 – Romania, Lagoon Complex Razelm-Sinoë

Meixner 1915 – Austria, Lunzer See

Müller & Faubel 1993 – Germany, Elbe Estuary

Nasonov 1919a, b, 1924 – Russia, Saint Petersburg and surrounding

1925 – Russia, Kola Peninsula

1546   1929 – Japan

1547   Noreña 1995 – Argentina, Paraná River

1548   Noreña et al. 1999 – Spain, Extramadura

1549   2006 – Peru, Loreto, Pacaya-Samiria National Reserve

1550   2007 – Spain, Iberian Peninsula, Galician and Cantabrian coast

Okugawa 1953 – Japan

Papi 1952 – Italy, Pisa and surrounding areas, artificial ditches, still or slow waters

Petrovna & Vladimirovich, 2013 – Russia, Tartarstan, Kazan, Kaban Lake

Pörner 1966 – Germany, Jena and surrounding

Reisinger 1933 – Indonesia, Sumatra

Reisinger & Steinböck 1925 – Austria, Wörthersee

Rixen 1961 – Germany, Schleswig-Holstein, Lakes and ponds, phytal region, detritus-

rich sand

1559   1968 – Germany, Bodensee, benthic and phytal regions, to 50 m

Rogozin, 2016 – Russia, Miass, Il'menskoye

Ročník 2007 – Czechia, lednické rybníky ponds

Sabussow 1900 – Russia, Solowetzki Island

Schwank 1976 – Germany, Bondensee

Seifert 1939 – Germany, Hiddensee

Sibiriakova 1929 – Russia, Angara

Southern 1936 – Ireland, Cork, Tipperary, Wicklow

Steinböck 1933 – Faeroe Islands

1568   1949, 1951a, b – Italy, Lago Maggiore, to 54 m

Tokinov & Berdnik, 2016 – Russia, Volga, Blue Lakes

Tu 1934 – China, Beijing, Tsing Hus University, campus pond

Urban-Malinga 2011 – Poland, Puck Bay, Hel Marine Station, 7 m depth

Valkanov 1926 - Bulgaria

Vara & Leal-Zanchet 2013 – Brazil, Rio Grande do Sul, in irrigated rice fields

Woodworth 1896 – United States, Michigan

Young 1970 – United Kingdom

1973 – United Kingdom, Wales

Zykov 1902 – Russia

1578

GenBank Accession

28S – KC869844, specimen from United States; KP730557, KP730520,

specimens from Finland; AJ270172 specimen from United Kingdom

18S – MH221198-257, MH221271-4, specimens from Sweden; KP730484,

MH221258-70 specimens from Finland; MH221235-7, specimens from Belgium;

D85092, specimen from Japan; KC869791, specimen from United States; U70082

collection location not listed

16S – KC869750, specimen from United States

ITS – MH221390-463, MH221475-6, specimens from Sweden; MH221464-

474, specimens from Finland; MH221428-30, specimens from Belgium

CO1 – MH221368-80, KP730567, specimens from Finland; MF185697-9,

MH221288-367, MH221381-2 specimens from Sweden; MH221329-31, specimens

from Belgium; AY228756, AJ405979-80, specimens from United Kingdom

Ef1a – AF288068, collection location not listed

1593

Body length of solitary animal 0.4-1 mm; chains to 8 mm, 18 zooids

maximum. Anterior end conically pointed. Posterior end tail-like. Body color

variable: clear, grey, yellow, pink or brownish. Pigmented eyespots present as paired

red stripes at dorso-anterior end. Amount of pigmentation highly variable and may be

absent in some individuals based on light intensity.

Sensory bristles present. Cilia covering body, shorter in mouth. Ciliary pits at

level of brain. Intestine ciliated and often including food remnants, including

Cladocera, chironomid larvae, Copepoda, hydroids, Nematoda and other small

animals. Nematocysts occuring throughout body.

Following TEM an SEM observations by Reuter (1978), groups of 6-8

adhesive papillae, each to 4 µm long, present in row at ventrolaterally at frontal end.

Nervous system includes brain, three pairs of nerve chords and pharyngeal

ring system. Protonephridia with bilateral stems and pores in anterior body.

Male reproductive system with bilateral testes, circular or eliptical vesicular

seminalis with prostate glands and separate gonopore. Male stylet spiraled, to 200 µm

long. Female system with single ovary, ciliated oviduct and ventral pore.

Proterandrous development.

1611

Notes: *Microstomum lineare* is an extremely common and abundant species in

brackish and freshwater habitats worldwide and can be found February through late

November or early December (Bauchhenss, 1971; Heitkamp, 1982). Lifecycle

includes 4 generations per year, with sexual reproduction occurring primarily in late

summer and fall. *M. lineare* purportedly has a global distribution and extremely wide

tolerance for differing environmental conditions, including salinity (freshwater to

7‰; Karling 1974), temperature (to 47°C; Graff 1913), calcium (to 50 mg/L; Young

1973), oxygen and pH (Kaiser 1969; Heitkamp 1979, 1982).

Positive identification of *M. lineare* should be cautiously done since several

other species of *Microstomum* are either very similar or identical in appearance to *M.*

lineare. In part this is due to the very wide morphological variability reported for the

species itself. Three species, *M. artoisi, M. tchaikovskyi,* and *M. zicklerorum*, were

separated from *M. lineare* primarily through DNA evidence based on the

mitochondrial cytochrome c oxidase subunit I and nuclear internal transcribed spacers

I and II and 5.8S rDNA gene loci, and thus have few or no morphologically

distinguishing characters (Atherton & Jondelius 2018b). A fourth, *M. giganteum*,

though separable based on morphological evidence, can easily be misidentified and

has historically been challenged as a "safe species". However, *M. giganteum* is a

larger species than *M. lineare* (8 zooid body length of ≥8.5 mm compared with 4-5

mm, respectively), without an elongated tail, adhesive papillae or pigmented eyespots,

with a clearly visible esophagus, with pharyngeal glands restricted to the oral cavity

only and with a three-part stylet (Kostenko 1988; Rogozin 2015).

1634

1635

1636   **22. *Microstomum lotti* sp. nov.; Fig. 6**

1637   Type: MTP LS 660

1638   Type Locality: Italy, Sant Andrea Bay, 42°48'31'' N, 10°08'30'' E

1639   Habitat: Marine

1640

1641   GenBank Accession:

1642   18S – KP730483, KP730493

1643

Body length 0.8 mm, 2 zooid. Colorless and without pigmented eyespots.

Anterior end conical; posterior end a spatulate tail plate. Adhesive papillae present in

a row ventrally at the frontal region ~30 μm from anterior end, along posterior half of

the body and very numerous at tail plate. Ciliary pits large and circular. Preoral

intestine short. Rhabdites present.

Reproductive system unknown.

1650

1651

1652   **23. *Microstomum lucidum* (FUHRMANN, 1896)**

1653   Synonyms:

1654   *Microstoma lucidum* Fuhrmann 1896, pg 1011

1655   *Microstomum hamatum* Westblad 1953

1656   *Microstomum hanatum* WORMS (accessed 10 Aug. 2018)

1657

1658   Type: deposition not recorded

1659   Type location: English Channel, Baie de la Forêt

1660   Habitat: Marine sublittoral

1661

1662   Steinböck 1931 – Faroes, Vaagfjord, Suderø, mud 10 m

1663   Meixner 1938 – United Kingdom, Plymouth, Wembury

1664   Westblad 1953, 1954 – Norway, Bergen University, black mud; England, Plymouth,

1665   40 m black mud; Blyth, 10 m, black mud.

1666   Karling 1966 – Norway, Herdla, Oöy and Blomöy

1667

Body length to 1.5 mm, 4 zooids. Colorless and transparent with yellowish

intestines. After Westblad (1953) may have (±) dark grey pigment spots in some

individuals. Eyes absent. Ends bluntly rounded. Posterior end with adhesive papillae.

Rhabdites present all over the body. Ciliary pits weakly visible. Preoral intestine

extending to brain.

One or two testes, with only one being large and well developed, connecting

via a single short vas deferens to the large vesicula seminalis. Stylet 60-70 µm, funnel

shaped, proximally slightly thickened and curved like a hook distally. Antrum

masculinum narrow.

Female system with unpaired ovary.

1678

1679

1680   **24. *Microstomum marisrubri* sp. nov.; Fig. 7**

1681   Type: MTP LS 524

1682   Type Locality: Egypt, Mangrove Bay, 25°52'15'' N, 34°25'04'' E

1683   Habitat: Marine

1684

1685   GenBank Accession:

1686   18S – KP730505

1687   CO1 – KP730580

1688

Body length 0.65, 1 zooid. Body colorless, without pigmented eyespots. Body

anterior to brain long, with rounded end. Posterior end blunt with numerous adhesive

papillae. Adhesive papillae sparsely present over entire body. Ciliary pits present at

level between anterior pharynx and brain. Preoral intestine short. Rhabdites absent.

Nematocysts intensely concentrated at posterior end.

Male reproductive system including a single testis connected by short vas

deferens to a rounded vesicula granulorum. Male stylet 63 µm long, shaped as a

slightly tapering curved tube with a slight widening at the distal tip. Genital pores

separate.

1698

1699

1700   **25. *Microstomum melanophthalmum* STEINBÖCK 1933**

1701   Type: deposition not recorded

1702   Type location: Croatia, Rovinj, S. Andrea

1703   Habitat: Marine, sublittoral to 7 m depth

1704

1705   Riedl, 1956 , 1959 – Croatia, Isola Saint Andrea; Italy, Capo di Sorrento, Sicily, Porto

1706   Venere; France, Banyuls-sur-mer

1707

Body length to 2.0 mm, to two zooids. Clear with yellow intestine. Black or

dark brown eyes with distinct crescent lens near the frontal end. Anterior tip blunt.

Posterior end conically rounded with adhesive tubes. Ciliary pits present just anterior

to brain. Preoral intestine large, extending to well above brain. Rhabdites present.

Male reproductive system with seminal vesicular attached to stylet, 30 µm

long, shaped as a sharply decreasing flat funnel connected to a tube hooked at an

angle of 90° with a pointed end and terminal opening. Female system not seen.

1715

1716

1717   **26. *Microstomum mortenseni* RIEDEL 1932**

1718   Type: deposition not recorded

1719   Type location: Greenland, Disco Bay

1720   Habitat: Marine, sublittoral benthal to 300 m depth

1721

Body to 1.2 mm in solitary zooid. Based on Riedel's (1932) original drawings,

anterior end conically rounded and posterior end rounded and wide. Ciliary pits

present at frontal end. Red color with yellow tinged rhammite glands. Pigmented

eyespots absent. Rhabdites present at lateral parts of body. Preoral intestine long,

anterior to brain. Intestine non-ciliated. Adhesive papillae not recorded.

Reproductive incompletely known. Ovary single with medial female

gonopore. S shaped stylet.

1729

1730

1731   **27. *Microstomum mundum* GRAFF 1905**

1732   Type: Deposition not recorded.

1733   Type locality: Black Sea, Sevastopol Bay, St. Georges Monastery

1734   Habitat: Marine, benthic in the sand

1735

Body to 2 mm, 8 zooids. Colorless without pigmented eyespots. Anterior end

conical. Adhesive papillae present in the posterior quarter of each well-developed

zooid. Ciliary pits present at level of brain. Rhabdites present throughout the body,

16-20 µm long. Nematocysts present. Intestine, including preoral intestine, with

characteristic lateral bulges and smaller lobules. Preoral intestine extending to a level

halfway between the ciliary pits and anterior tip.

Reproductive system unknown.

1743

1744

1745   **--*Microstomum ornatum* ULJANIN 1870**

1746   Species dubiae by Westblad (1953) based on its incomplete description, perfunctory

1747   drawings, the posterior positioning of the ovary, which is unique among species of

1748   *Microstomum*, and the anterior direction of egg development, which seems to conflict

1749   with the posterior position of the female duct.

1750

1751

1752   **28. *Microstomum papillosum* GRAFF 1882; Fig. 18D**

1753   Synonyms:

1754   *Microstoma papillosum* – Claparède 1861, p152

1755

1756   Type: Deposition not recorded

1757   Type Locality: Sartor Oë, Norway

1758   Habitat: Marine, littoral to sublitteral.

1759

1760   Armonies 1987 - North Sea, Sartorø near Bergen

1761   2017 – Germany, Sylt, North Sea

Ax 1959 – Black Sea, Bosporus, Marmara Sea

Böhmig 1889

Faubel 1974, 1977 – Germany, Sylt, North Sea

Faubel & Warwick 2005 – United Kingdom, Scilly, St. Martin and surrounding

Hellwig 1987

Micoletzky 1910 – Gulf of Trieste, Croatia, Italy

Riedl 1956 – Italy, Capo di Sorrento, Sicily, Porto Venere; France, Banyuls-sur-mer

Steinböck 1938 – Iceland, Isafjördur, benthic and in algae

Wehrenberg & Reise 1985 – Germany, Sylt, North Sea

1771

1772   GenBank Accession

1773   28S – FJ715316, AF021330 specimens from Germany

1774   18S – FJ715296, specimen from Germany

1775   16S – KP730453, specimen from Germany

1776   CO1 – KP730570, specimen from Germany

1777

Body length to 1 mm, 5 zooids. Body shape with rounded anterior and a

slightly tapered, rounded posterior. Body colorless except for reddish-brown intestine.

Pigmented eyespots absent. Shallow ciliary pits present at level of anterior pharynx.

Adhesive papillae scattered across the whole body but most numerous at anterior and

posterior ends. Rhabdites present, especially at anterior end. Preoral intestine long,

extending almost to anterior end.

Male genital system consisting of unpaired testis and vesicula seminalis.

Stylet shaped as a curved tube, tapering to a thin distal terminal opening, 80 µm long.

Female system including single ovary.

1787

1788   Notes: *Microstomum papillosum* has a somewhat similar stylet shape as *M. gabriellae*

1789   and *M. crildensis*, although its stylet is larger (80 µm compared to 50 and 32 µm,

1790   respectively) and tapers to a thinner tube distally. Other differences include, for *M.*

1791 *crildensis*, the size of the preoral intestine and the ciliary pits and the absence of

1792 rhabdites; and for *M. gabriellae*, the red body coloring and red eyespots.

1793

1794

1795 **29. *Microstomum paradii* (Duges 1828)**

1796 Synonyms:

1797 *Derostoma squalus* – Duges 1828, pg 142

1798

1799 Type: Deposition not recorded

1800 Type locality: Romania, Transylvania, Cluj-Napoca

1801 Habitat: Freshwater

1802

1803 Graff 1913 – France, Montpellier

1804

1805   Body to 2.5 mm, 2 zooids. Anterior end rounded; posterior tail-like with

1806 adhesive tubules. Colorless with light brown intestine. Black eyespots with conical

1807 lenses present anterior to ciliary pits. Rhabdites sparsely present throughout the body.

1808 Preoral intestine long.

1809   Reproductive system not seen.

1810

1811

1812 **30. *Microstomum proliferum* (Busch 1851)**

1813 Synonyms:

1814 *Alaurina prolifera* Busch 1851, pg 114

1815

1816   Types: deposition not recorded

1817   Type locality: United Kingdom, Skye Islands

1818   Habitat: Marine, pelagic

1819

1820   Fewkes 1883 – USA, Newport

1821   Graff 1913 – USA, New Jersey, Newport; Spain, Malaga

1822

Body to 2.2 mm, 2 zooids. Pigmented eyes present lateral and posterior to

mouth. Anterior end with conically pointed proboscis, non-ciliated, with many

circular folds and denoted at the base by a tuft of cilia on each side. Sensory bristles

present evenly spaced along lateral margins of the body, twice as long as the cilia, and

single even longer bristle present at posterior end. Adhesive papillae present along

entire body.

1829

1830   Notes: Brinkmann (1905) argued that this species is identical to *Alaurina composita*

1831   since any differences were either errors or derived from outdated methods of

1832   investigation. Graff (1913) argued against this, saying the two species differ in the

1833   sensory cilia and the location of the pigmented eyespots.

1834

1835

1836   **31. *Microstomum punctatum* DORNER 1902**

1837   Types: Deposition not recorded

1838   Type location: Linkenen Pond

1839   Habitat: Freshwater pond, muddy with plants

1840

1841 Kaiser 1974 – Germany, Mühlhausen, Bad Langensalza

1842 Wegelin 1966 – Germany, Halle, Elbe and surrounding

1843

Body length 1 mm, 4 zooids maximum. Anterior end spade-like, pointed and

separated from the posterior by a distinct constriction at level of the mouth. Posterior

end tail-like. Body color brownish yellow, with numerous black dots at the anterior.

Ciliary pits weak, at level of mouth. Adhesive papillae very sparsely present.

Rhabdites absent. Pharynx very large, ¼ to ⅓ body size. Preoral intestine extremely

small, without extending to anterior end of pharynx.

Single ovary. Male system unknown.

1851

1852

1853 **32. *Microstomum rhabdotum* MARCUS 1950**

1854 Type: Deposition not recorded

1855 Type location: Brazil, São Sebastião

1856 Habitat: Marine, between *Sargassum stenophyllum*

1857

Body length to 0.7 mm, 3 zooids. Rounded anterior and posterior ends.

Colorless and without pigmented eyespots. Adhesive papillae present at anterior,

mouth, dividing zones, posterior. Biparte pharynx separated by annular fold with

larger anterior cilia than posterior. Paired stripes of rhabdite bundles on the

dorsolateral sides, joining at level of ciliary pits; median dorsal and ventral areas free

of rhabdites; Preoral intestine reaches to anterior border of brain.

Reproductive system not seen.

1865

1866

1867 **33. *Microstomum rubromaculatum* (GRAFF 1882); Fig. 18C**

1868 Synonyms:

1869 *Microstoma rubromaculatum* Graff 1882, pg 251

1870

1871 Type: deposition not recorded

1872 Type location: Gulf of Naples, Tyrrhenian Sea

1873 Habitat: Marine, sublittoral phytal in *Sargassum* spp, benthic on shell sand.

1874

Atherton & Jondelius 2018a – Sweden, Fiskebäckskil

Graff 1913 – France, Concarneau

Steinböck 1931 – Faroes, Vaagfjord, Suderø, mud 10 m

Westblad, 1934 – Norway, Herdla

Southern 1936 – Ireland, Dublin, Malahide Inlet, New Harbor, Galaway, 1-2 m

Steinböck 1938 – Iceland, Ísafjör∂ur, fine sand and algae, 1-2 m

Meixner 1938 – Enlgand, Wembury

Westblad 1953 – Sweden, Gullmar Fjord, Fiskebäckskil, 1-2 m

1883

1884 GenBank Accession:

1885   18S – XXX

1886   CO1 – MF185684-MF185696, specimens from Sweden

1887

Body length to 2 mm; maximum of 8 zooids. Very large red eyespot

pigmentation, merging dorsally to form a broad transverse band at the anterior of the

animal. Otherwise colorless or slightly pinkish reflective of intestines. Body ends

bluntly rounded. Adhesive tubules present and concentrated at anterior and posterior

ends. Rhabdites present throughout animal. Ciliary pits present with long cilia.

Preoral intestine small.

Reproductive system based on specimens collected from Fiskebäckskil,

Sweden (Atherton & Jondelius, 2018a). Male system with paired testes, copulatory

apparatus, antrum masculinum and separate gonopore. Stylet 75-112 µm long, shaped

as a single wide spiral bent around a 90° angle, terminating with a fingerlike hook.

Female system with a single ovary situated mid-body, antrum, and a separate

gonopore. Eggs develop caudally.

1900

1901 Notes: The amounts of eyespot pigmentation may be greatly variable, with

1902 pigmentation ranging from a bright circular band around the anterior end to two

1903 clearly distinct separate spots. Steinböck (1938) found one animal with a single large

1904 pigment spot in the middle that became thinner towards the margins of the body.

1905

1906

1907 **34. *Microstomum schultei* sp. nov.; Fig. 8**

1908 Type: MTP LS 700

1909 Type Locality: Italy, Fetovia Bay, 42°43'36'' N, 10°09'33'' E

1910 Habitat: Marine

1911

1912 GenBank Accession:

1913   18S – KP730494

1914

1915   Body length 1.5 mm, 1 zooid. Body light reddish-brown without pigmented

1916   eyespots. Anterior end rounded, with a small constriction at the ciliary pits. Posterior

1917   end conically blunted. Adhesive papillae and rhabdites present, concentrated at body

1918   ends. Ciliary pits present posterior to level of brain. Preoral intestine short.

1919   Reproductive system with curved male stylet, tapering from a wider base to a

1920   narrow tube with a beveled tip.

1921

1922

1923   **35. *Microstomum septentrionale* (Sabussow 1900); Fig. 18G**

1924   Synonyms:

1925   *Microstoma ornatum* Sabussow 1897, pg 14

1926   *Microstoma septentrionale* Sabussow 1900, pg 15, 81, 181

1927

1928   Type: Deposition not recorded

1929   Type location: Russia, Kola Peninsula

1930   Habitat: Marine, littoral among algae

1931

1932   Karling 1934 – Iceland, Herdla Kværnepollen

1933   Westblad 1953 – Sweden, Gullmarfjord, Gåsövik

1934   This paper – Sweden, Fiskebäckskil, 58°14′59″ N, 11°26′45″ E

1935

1936   GenBank Accession:

1937   18S – XXX

1938   CO1 – XXX

1939

1940   Maximum body length 1 mm, 2 zooids. Body translucent yellow with bright

1941   yellow intestines. Some specimens may have darker yellow or orange yellow in the

1942   anterior end. Pigmented eyespots absent. Anterior end rounded with a tapering

1943   rounded posterior end. Adhesive papillae sparsely scattered over the entire body.

1944   Preoral intestine reaching to level of anterior brain. Rhabdites present throughout the

1945   body, and especially concentrated in anterior end. Ciliary pits present at level of brain.

1946   Single ovary and bilobed testis. Copulatory organ a curved tube, with a wide

1947   base and tapering distally to a point. Stylet length to 68 µm. Antrum masculinum

1948   present just anterior to stylet, large and ciliated. Separate ventral male and female

1949   pores.

1950

1951

1952   **36. *Microstomum spiculifer* FAUBEL 1974**

1953   Type: Sagittal section series; Deposition numbers not listed

1954   Type location: North Sea, North Frisian Uslands, Sylt Island, List

1955   Habitat: Marine, eulittoral, semi-exposed sandy beach

1956

1957   Body length to 1.5 mm long, maximum 4 zooids. Body color clear with

1958   reddish brown intestine. Eyes absent. Ends rounded, posterior with multiple adhesive

1959   papillae. Preoral intestine reaching to level of brain or slightly anterior. Ciliated pits

1960   present at brain. Rhabdites present across body, concentrated at anterior.

1961   Testis unpaired, located laterally and posterior to male copulatory apparatus.

1962   Masculine antrum elongated. Stylet strait, ca. to 55 µm long, decreasing from a width

1963   of 7-8 µm to a distal point. Ovary unpaired. Male and female pores separate.

1964

1965

1966   **37. *Microstomum spiriferum* WESTBLAD 1953**

1967   Lectotype: ⬜, whole mount, SMNH-Type-XXXX; E. Westblad leg.**

1968   Type Locality: Sweden, Gullmarsfjorden, Gullmarsvik

1969   Habitat: Marine, mud, sublittoral between 10-60 m depth

1970

1971   Westblad 1953 – Sweden west coast, 10 m ; Norway, Tromsö, Ramfjord, 10-25 m;

1972   United Kingdom, Scotland, Millport, 30-35 m

1973

1974   Body to 1.5 mm, generally composed of 2 zooids. Color pale reddish-yellow

1975   due to rich pigment cells in dorsal side. Red eyespots located at level of ciliary pits,

1976   may extend dorsally in some animals. Anterior and posterior ends rounded. Rhabdites

1977   present, concentrated in anterior and posterior ends. Adhesive papillae absent. Preoral

1978   intestine extending to level of ciliary pits.

1979   Testis singular, small, connecting via a seminal duct to the vesicula seminalis

1980   and posterior penis stylet. Penis stylet large, 140 μm, with 5-6 dexiotropal torsions.

1981   With two female ducts and orifices, both connected to the ovary. The anterior acts as

1982   a ciliated oviduct and is separated from the ovary by a sphincter. The posterior acts as

1983   a female copulatory organ and may contain spermatozoids in the proximal end.

1984

1985   **A type specimen was not previously designated for *M. spiriferum*. However,

1986   fourteen original specimens were deposited at the Swedish Museum of Natural

1987   History, and thus, following the rules of the International Code of Zoological

1988   Nomenclature, we hereby designate one whole mount hermaphroditic specimen

1989   (formerly SMNH94968, now SMNH-Type-XXXX) as lectotype for *Microstomum*

1990   *spiriferum* WESTBLAD 1953. The remaining thirteen specimens (SMNH94966-7,

1991   SMNH94969-73, SMNH97839-44) are thus designated as paralectotypes.

1992

1993

1994   **38. *Microstomum tchaikovskyi* ATHERTON & JONDELIUS 2018; Fig. 17C**

1995   Type: Asexual, SMNH-Type-9025; GenBank accession 18S MH221277, CO1

1996   MH221385, ITS MH221479

1997   Type Locality: Sweden, Sollentuna, 59°26'26''N 18°00'03''E, 15 cm depth

1998   Habitat: Freshwaters of Sweden and Finland, phytal to 20 cm depth

1999

2000   GenBank Accession:

2001   18S – MH221277, from Sweden; MH221278, from Finland

2002   CO1 – MH221385, from Sweden; MH221386, from Finland

2003   ITS – MH221479, from Sweden; MH221480, from Finland

2004

2005   Molecular Diagnosis:

2006   COI gene region with reference to GenBank Accession MH221288: 6-T; 15-

2007   A; T-16; 18-A; 30-G; 54-A; 69-A; 84-A; 180-G; 186-G; 228-A; 246-C; 279-A; 310-

2008   C; 312-T; 318-A; 321-A; 333-A; 351-A; 435-T; 486-G; 540-A; 603-A; 609-1; 625-C;

2009   627-T

2010   ITS1, 5.8S, ITS-2 gene region with reference to GenBank Accession

2011   MH221390: 121-T; 218-A; 288-T; 353-G; 358-G; 394-G; 497-T; 533-A; 730-G; 759-

2012   C; 768-C; 787-A; 800-T; 845-T; 887-T; 999-T

2013

2014   Morphological Description:

2015   Body length approximately 4 mm, 6 zooids. Body clear with darker intestines.

2016 Preoral intestine short. Anterior end abruptly tapering, posterior end tail-like without

2017 adhesive papillae. Sensory bristles present. Ciliary pits present at the level of anterior

2018 pharynx. Red eye pigmentation in two stripes, located dorsally at anterior end. May

2019 be weak in some individuals. Nematocysts few, scattered across the body.

2020   Reproductive system not recorded.

2021

2022 Notes: See notes of *M. lineare*.

2023

2024 **39. *Microstomum trichotum* Marcus 1950**

2025 Lectotype: ♂, whole mount, SMNH-Type-XXXX, E. Marcus leg.**

2026 Type Locality: Brazil, São Sebastião

2027 Habitat: Marine, among calcareous algae incl. Corallinaceae and *Jania rubens*

2028

2029   Body length small, approximately 0.5 mm, 3 zooids. Body color white,

2030 without pigmented eyespots. Anterior end tapered and long, resembling the proboscis

2031 characteristic of *Alaurina* and densely ciliated, posterior end rounded with ca. 20

2032 adhesive papillae. Preoral intestine to anterior brain. Ciliated pits small. Anterior

2033 rhammite tract absent. Rhabdites present throughout body.

2034   Male reproductive system with single testis. Stylet shaped as a curved funnel,

2035 approximately 12 µm long. Female system not seen.

2036

2037 **A type specimen was not previously designated for *M. trichotum*. However, four

2038 original specimens were deposited at the Swedish Museum of Natural History, and

2039 thus, following the rules of the International Code of Zoological Nomenclature, we

2040 hereby designate one whole mount male specimen (formerly SMNH108151, now

2041 SMNH-Type-XXXX) as lectotype for *Microstomum trichotum* MARCUS 1950. The

2042 remaining three specimens (SMNH108149-50, SMNH10852) are thus designated as

2043 paralectotypes.

2044

2045

2046 **40. *Microstomum ulum* MARCUS 1950**

2047 Lectotype: Asexual, whole mount, SMNH-Type-XXXX, Leg. E. Marcus

2048 Type locality: Brazil, São Sebastião

2049 Habitat: Marine, sublittoral in coarse sand

2050

2051 Body size to 1.1 mm, 4 zooids. Colorless and without pigmented eyespots.

2052 Anterior end conically pointed, posterior separated by a constriction, forming a

2053 distinctly spatulate tail-plate. Adhesive papillae present. Rhabdites present throughout

2054 the body, but heavily concentrated at anterior end above the pharynx. Deep ciliary pits

2055 present at level of brain. Preoral intestine short.

2056 Reproductive system not seen.

2057

2058 **A type specimen was not previously designated for *M. ulum.* However, three

2059 original specimens were deposited at the Swedish Museum of Natural History, and

2060 thus, following the rules of the International Code of Zoological Nomenclature, we

2061 hereby designate one whole mount asexual specimen (formerly SMNH108153, now

2062 SMNH-Type-XXXX) as lectotype for *Microstomum ulum* MARCUS 1950, while the

2063 remaining two specimens (SMNH108154-5) are thus designated as paralectotypes.

2064

2065

2066 **41. *Microstomum weberi* sp. nov.; Fig. 9**

2067 Type: MTP LS 666

2068 Type Locality: Italy, Pianosa, 42°34'29'' N, 10°03'59'' E

2069 Habitat: Marine

2070

2071 GenBank Accession:

2072   18S – KP730487

2073   CO1 – KP730576

2074

2075   Body length of 0.65 mm, 2 zooid. Colorless and without pigmented eyespots.

2076 Anterior end conical; posterior end widely blunt. Adhesive papillae present at

2077 posterior end and in a ventral row at the frontal end 25 µm from anterior tip. Ciliary

2078 pits bottle-shaped. Preoral intestine extending to anterior brain or just beyond. Large

2079 rhabdites bundles present at anterior and posterior ends.

2080   Reproductive system unknown.

2081

2082

2083 **42. *Microstomum westbladi* nom. nov.; Fig. 10**

2084 Synonym:

2085 *Microstomum papillosum* GRAFF 1882, sensu Westblad 1953, pg 406-7

2086

2087 Type: Asexual, sectioned, SMNH-Type-XXXX

2088 Type Locality: Norway, Herdla

2089 Habitat: Marine, sublittoral sand and mud

2090   GenBank Accession:

2091   18S – XXX

2092   CO1 – XXX

2093

2094   Westblad, 1953 – Norway, Herdla, Kværnepoll current; Sweden, Gullmarfjord

2095   This paper – Sweden, Malmön, 58°20'18'' N 11°20'26'' E

2096

Body length to 2.0 mm, 5 zooids. Colorless and without pigmented eyespots.

Anterior and posterior ends bluntly rounded; posterior with numerous adhesive

papillae. Ciliary pits small. Rhabdites numerous and intensely concentrated at frontal

end. Adhesive papillae present sparsely along body entire length, with characteristic

grouping present at anterior tip: field longest at both lateral sides and curving upwards

toward the middle. Preoral intestine extending to brain.

Reproductive system including male stylet, 60 µm long, and shaped as a

smooth curve 180° curve, with a maximum width ~⅙ of the way to the distal end,

decreasing proximally and also distally to a point.

2106

2107

2108   **43. *Microstomum zicklerorum* ATHERTON & JONDELIUS 2018; Fig. 17B**

2109   Type: Asexual, SMNH-Type-9026; GenBank accession 18S MH221281, CO1

2110   MH221389, ITS MH221483

2111   Type Locality: USA, Massachusetts, North Chelmsford 42°38'13''N 71°23'30''W

2112   Habitat: Freshwater benthic or phytal; to 25 cm depth

2113

2114   GenBank Accession:

2115   18S – MH221279 – MH221280

2116   CO1 – MH221387 – MH221389

2117   ITS – MH221481 – MH221483

2118

Molecular Diagnosis:

COI gene region with reference to GenBank Accession MH221288: 24-A; 30-

T; 123-T; 165-A; 186-A; 192-G; 201-G; 267-C; 300-C; 318-G; 324-G; 330-A; 336-C;

387-C; 501-A; 525-G; 555-G; 558-A; 588-A; 609-G; 630-C

ITS1, 5.8S, ITS-2 gene region with reference to GenBank Accession

MH221390: 197-A; 317-A; 507-T; 829-A; 879-A; 888-C; 1000-A

2125

Morphological Description:

Body length to 1.5 mm, 2 zooids. Body clear and reflective of intestines. Red

eye pigmentation extremely weak or not present. Anterior end conical, posterior end

tail-like without adhesive papillae. Sensory bristles present. Ciliary pits present with

circular openings lateral at the level of mid-pharynx. Preoral intestine short. Mouth

and pharynx large leading to strait intestine. Intestine containing desmid algae.

Nematocysts present throughout the body.

Reproductive system not recorded.

2134

2135   Notes: See notes of *M. lineare.*

2136

2137

2138   **Gen. Myozonella BEKLEMISHEV 1955**

2139   Microstomidae with ciliary pits at frontal end and intestine extending anterior

2140 to mouth opening (preoral intestine). Muscle ring present surrounding intestine.

2141 Asexual reproduction through fissioning present; sexual reproduction unknown.

2142 Freshwater, one species.

2143   Type species: *Myozonella microstomoides* BEKLEMISHEV 1955

2144

2145 **1. *Myozonella microstomoides* BEKLEMISHEV 1955**

2146 Type: Deposition not recorded

2147 Type locality: Russia, Near Bryansk in town of Sudimir

2148 Habitat: Freshwater, phytal in *Utriculara* sp and *Calopteryx* sp.

2149

Body size to 1.0 mm, 2 zooids. Colorless and without pigmented eyespots.

Anterior end conically pointed, posterior with thin tail. Adhesive papillae and

rhabdites absent. Ciliary pits present dorsolaterally, narrow and deep. Preoral intestine

extending to level of ciliary pits. Muscle ring present surrounding intestines just

posterior to pharynx.

Reproductive system not seen.

2156

2157

Acknowledgements

We would like to thank many people for their valuable help, insight and support: Dr.

Tom Artois, Edmond Atherton, Laura Atherton, Adam Bates, Dr. Marco Curini-

Galletti, Dr. Rick Hochberg, Ylva Jondelius and the entire Zickler family. We are

grateful to the staffs of the Stensoffa Research Station, Abisko Scientific Research

station, and Umeå Marina Forskningcentrum. We are also particularly indebted to Dr.

Lukas Schärer for the collection, documentation and sequencing of four of the species

included in this manuscript and for generally providing insights and instruction on

Macrostomorpha taxonomy. Many thanks also to Cassandra Vion-Bailly for creating

and providing all drawings in this manuscript. This research was supported by a grant

from the Swedish Taxonomy Initiative (no. DHA 2014-151 4.3) to UJ.

2169

2170

Work Cited

Álvarez-Campos P, Giribet G, Riesgo A, 2017. The *Syllis gracilis* species complex: a

molecular approach to a difficult taxonomic problem (Annelida,

Syllidae). Mol Phylo Evol. 2017;109:138-50.

An Der Lan H. Zur rhabdocoelen Turbellarienfauna des Ochrid-Sees (Balkan). Sber

Akad Wiss Math Nat Kl Wien Abt I. 1939;148:195-254.

An Der Lan H. Zur Ökologie und Populations dynamik rhabdocoeler Turbellarien im

Ohrid-See. Arhiv Biol Nauka Archs Sci Biol. 1961;13:59-66.

An Der Lan H. Zur Turbellarienfauna der Donau. Arch Hydrobiol. 1962;27 Suppl

1:3-27.

Armonies W. Freilebende Plathelminthen in supralitoralen Salzwiesen der Nordsee:

Ökologie einer borealen Brackwasser-Lebensgemeinschaft. Microfauna

Marina. 1987;3:81-156.

Armonies W. Long-term change of meiofaunal species composition in a sandy beach,

with description of 7 new species of Platyhelminthes. Helg Mar Res.

2017;71:12.

Atherton S, Jondelius U. *Microstomum* (Platyhelminthes, Macrostomorpha,

Microstomidae) from the Swedish west coast: two new species and a

population description. Eur J Taxon. 2018a;398.

Atherton S, Jondelius U. Wide distributions and cryptic diversity within a

Microstomum (Platyhelminthes) species complex. Zool Script. 2018b;1-13.

Attems C. Beitrag zur Kenntnis der rhabdocoelen Turbellarien Helgolands. Wiss

Meeresunters. 1896;2:219–32.

Ax P, Armonies W. Amphiatlantic identities in the composition of the boreal brackish

water community of Platyhelminthes. A comparison between the Canadian

and European Atlantic coast. Microfauna Mar. 1987;3:7-80.

Ax P. Die Einwanderung mariner Elemente der Mikrofauna in das limnische

Mesopsammal der Elbe. Verhandl Deutsch Zool Ges Hamburg. 1956:427-35.

Ax P. Zur Systematik, Ökologie und Tiergeographie der Turbellarienfauna in den

ponto-kaspischen Brackwassermeeren. Zool Jb Syst. 1959;87:43-184.

Ax P. Brackish-water Plathelminthes from the Faroe Islands. In: Biology of

Turbellaria and some Related Flatworms. Dordrecht: Springer. 1995. p. 45-7.

Bauchhenss J. Die Kleinturbellarien Frankens. Ein Beitrag zur Systematik und

Ökologie der Turbellarien excl. Tricladida in Süddeutschland. Int Revue ges

Hydrobiol. 1971;56:609-66.

Beklemishev V. Materialy po sistematike i faunistike turbellarii Vostochnoi Rossii.

Isvest Ross Akad Nauk. 1921;15:631-56.

Beklemishev V. [A clear case of parallelism of development in turbellaria: muscle

muff of the intestine in the group of Macrostomida]. Dokl Akad Nauk SSSR.

1955;104:789-791. Russian.

Böhmig L. *Microstoma papillosum*. Zool Anz. 1889;12:479-83.

Braccini J, Leal-Zanchet A. Turbellarian assemblages in freshwater lagoons in

southern Brazil. Invert Biol. 2013;132:305-14.

Braun M. Über die Turbellarien Livlands. Zool Anz. 1885;8:696-9.

Brinkmann A. Studier over Danmarks rhabdoele og acoele Turbellarier. Vidensk

Medd Dansk Naturhist Foren. 1905;1:1-159.

Brusa F. Macrostomida (Platyhelminthes: Rhabditophora) from Argentina with the

description of two new species of *Macrostomum* and of stylet ultrastrucutre.

Zool Sci. 2006;23:853-62.

Busch W. Beobachtungen über Anatomie und Entwicklung einiger wirbellosen

Seethiere. Berlin: Hirschwald; 1851. 143 pp.

Butlin R, Schön I, Martens K. Asexual reproduction in nonmarine ostracods.

Heredity. 1998;81:473-80

Carrasco N, Perissinotto R. Development of a halotolerant community in the St. Lucia

Estuary (South Africa) during a hypersaline phase. PloS One. 2012;7:e29927.

Chodorowski A. Ecological differentiation of turbellarians in Harsz Lake. Polsk Arch

Hydrobiol. 1959;6:33-73.

Claparède E. Recherches anatomique sur les Annélides, Turbellariés, Opalines et

Gregarines observés dans les Hebrides. Mem Soc Geneve Georg. 1861;96.

Davis A, Unmack P, Pusey B, Johnson J, Pearson R. Marine-freshwater transitions

are associated with the evolution of dietary diversification in terapontid

grunters (Teleostei: Terapontidae). J Evol Biol. 2012;25:1163-79.

Delp A. Flatworm predation on juvenile freshwater mussels. Masters thesis Graduate

College of Southwest Missouri State University. 2002. 31 pp.

DeSalle R, Egen M, Siddall M. The unholy trinity: Taxonomy, species delimitation

and DNA barcoding. Philos Trans R Soc Lond Biol Sci. 2005;360:1905-16.

Diesing K. Revision der Turbellarien. Abtheilung: Rhabdocoelen. Sber Akad Wiss

Wien, Math-nat Kl. 1862;45:191-318.

Dorner G. Darstellung der Turbellarienfauna der Binnengewässer Ostpreußens. Schr

Phys Ökon Ges Königsberg I Pr. 1902;43:1-58.

Dugès A. Recherches sur l'organisation et les moeurs des planariees. Ann Sci Nat Ser

1. 1828;15:139-82.

Edgar R. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucl Acids Res. 2004;32:1792-7.

Egger E, Lapraz F, Tomiczek B, Müller S, Dessimoz C, Girstmair J et al. A

transcriptomic-phylogenomic analysis of the evolutionary relationships of

flatworms. Cur Biol. 2015;25:1347–53.

Ehrenberg C. Phytozoa, Turbellaria africana et asiatica. In: Hemprich, F. &

Ehrenberg, C. (eds.), "Symbolae Physicae." Animalia evertebrata exclusis

insectis recensuit Dr. CG Ehrenberg. Series Prima cum Tabularum Decade

Prima. Berolini: Berlin. 1831; folia a-d.

Fabricius O. Fortsaettelse af nye zoologiske bidrag. VI. Nogle lidet bekjendte of

tildels nye flad-orme (Planariae). Kong Danske Vid Selsk Naturvid og

Mathem Afhandlingar, 2. Deel Kjöbenhavn. 1826. 34 pp.

Fang C, Wang L, Zang Y, Wang A. Two new species of brackish-water

Macrostomum (Platyhelminthes, Macrostomida) from southern China.

Zootaxa. 2016;4170 (2): 298–310.

Faubel A, Cameron B. Platyhelminthes from salt marshes of Coomera river

southeastern Queensland, Australia. Mem Qld Mus. 2001;46:511-9.

Faubel A, Warwick R. The marine flora and fauna of the Isles of Scilly: Free-living

Plathelminthes ('Turbellaria'). J Nat Hist. 2005;31:1-45.

Faubel A. Macrostomida (Turbellaria) von einem Sandstrand der Nordseeinsel Sylt.

Mikrofauna Meeresboden. 1974;45:339-70.

Faubel A. The distribution of Acoela and Macrostomida (Turbellaria) in the littoral of

the North Frisian Islands Sylt, Römö, Jordsand and Amrum (North Sea).

Sencken-bergiana Marit. 1977;9:59-74.

Faubel A. Experimentelle Untersuchungen zur Wirkung von Rohoel und

Rohoel/Tensid-Gemischen im Ökosystem Wattenmeer. X. Turbellaria.

Senckenbergiana Marit. 1984;16:153-70.

2270

Ferguson F, Stirewalt M, Brown T, Hayes W. Studies on the turbellarian fauna of the

Mountain Lake Biological Station. I. Ecology and Distribution. J Elisha

Mitchell Sci Soc. 1939;55:274-88.

Feweks J. Note on *Alaurina prolifera* Busch. Amer Naturalist XVII. 1883;668-9.

Fišer C, Robinson C, Malard F. Cryptic species as a window into the paradigm shift

of the species concept. Mol Ecol. 2018;27(3):613-35.

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification

of mitochondrial cytochrome c oxidase subunit I from diverse metazoan

invertebrates. Mol Mar Biol Biotech. 1994;3:294-9.

Fonseca G, Fontaneto D, Maikon D. Addressing biodiversity shortfalls in meiofauna.

J Exp Mar Biol Ecol. 2017;502:26-38.

Fontaneto D, Flot J.-F, Tang, C. Q. Guidelines for DNA taxonomy, with a focus on

the meiofauna. Marine Biodiversity. 2015. 45(3), 433–451.

http://doi.org/10.1007/s12526-015-0319-7

Fuhrmann O. Die Turbellarien der Umgebung von Basel. Rev Suisse Zool.

1894;2:215-92.

Fuhrmann O. Nouveaux Rhabdocoelides marins de la Baie de Concarneau. Comp

Rend Soc Biol Année. 1896;1011-3.

Fuhrmann O. Neue Turbellarien der Bucht von Concarneau. Zool Anz. 1898;21:252-

6.

Fulinski B, Szynal E. Über die Turbellarienfauna aus der Umgebung von

Grzymaloew (Podolien). Kosmos (Lwow), Ser A, Rozprawy. 1932;57:177-

217.

Gamble F. Contributions to a knowledge of British marine Turbellaria. Quart J

Microsc Sci Lond. 1893;34:433-528.

Gamo J, Noreña C, Janssen C. Old and new records of turbellarians from the central

areas of Spain. Hydrobiologia. 1998;383:299-305.

Gieysztor M. Über die Rhabdocoelidenfauna der Umgebung von Warschau. Bull

Intern Acad Polon Ser B, Sci Nat. 1926;617-71.

Graff L. Monographie der Turbellarien: I. Rhabdocoelida. Verlag Wilhelm

Engelmann: Leipzig. 1882. 442 pp.

Graff L. Turbellaria I. Acoela. In: Schultz F, editors. Das Tierreich Vol.

23. Berlin: Friedlande. 1905;28-9.

Graff L. Turbellaria, Strudelwürmer. I. Allgemeines und Rhabdocoelida. Herausgeg.

von A. Brauer. Heft 19, Berlin. 1909;59–142

Graff L. Acoela, Rhabdocoela, Alloeocoela des Ostens der Vereinigten Staaten von

America. Mit Nachträgen zu den marinen Turbellarien Orotavas und der

Küsten Europas. Z Wiss Zool. 1911;99:1-108.

Graff L. Turbellaria II. Rhabdocoelida. Das Tierreich. 1913;35:1-484.

Gutafsson M, Eriksson K, Hydén A. Never ending growth and a growth factor. I.

Immunocytochemical evidence for the presence of epidermal growth factor in

a tapeworm. Hydrobiologia 1995;305:229-33

Hallez P. Contributions à l'histoire naturelle des Turbellariès - Travaux de l'Institut

Zoologique de Lille et de la Station Maritime de Wimereux. Theses, Fac Sci

Paris. 1878. 213 pp.

Hallez P. Contribution a 1'histoire naturelle des Turbellariés. Trav Inst Zool Lille Stat

marit Wimereux. 1879;2:1–213.

Hartog J. The marginal tentacles of *Rhodactis sanctithomae* (Corallimorpharia) and

the sweeper tentacles of *Montastrea cavernosa* (Scleractinia); their cnidom

and possible function. Proc 3rd int Coral Reef Symp. 1977;1:463–79.

Heitkamp U. Zur Systematik, Biologie und Ökologie der Turbellarien periodischer

und perennierender Kleingewässer Südniedersachsens. Habilitationsschrift

Göttingen. 1979. 311 pp.

Heitkamp U. Untersuchungen zur Biologie, Ökologie und Systematik limnischer

Turbellarien periodischer und perennierender Kleingewässer

Südniedersachsens. Arch Hydrobiol. 1982;64 Suppl: 65-188.

Hellwig M. Ökologie freilebender Plathelminthen im Grenzraum Watt-Salzwiese

lenitischer Gezeitenküsten. Microfauna Mar. 1987;3:157-248.

Hofker J. Faunistische Beobachtungen in der Zuidersee während der Trockenlegung.

Z Morph Ökol Tiere. 1930;18:189-216.

Hofsten N. Studien über Turbellarien aus dem Berner Oberland. Z wiss Zool.

1907;85:391-654.

Hofsten N. Neue Beobachtungen über die Rhabdocölen und Allöocölen der Schweiz.

Zool Bijdr Uppsala. 1911;1:1-84.

Hofsten N. Revision der schweizerischen Rhabdocoelen und Alloeocoelen. Rev

Suisse Zool. 1912;20:543-687.

Janssen T, Vizoso D, Schulte G, Littlewood D, Waeschenbach A, Schärer L. The first

multi-gene phylogeny of the Macrostomorpha sheds light on the evolution of

sexual and asexual reproduction in basal Platyhelminthes. Mol Phylog Evol.

2015;92:82-107.

Jörger K, Schrödl M. How to describe a cryptic species? Practical challenges of

molecular taxonomy. Front Zool. 2013;10:59.

Kaiser H. Zur Turbellarienfauna zweier Moorgebiete in der Umgebung von

Mühlhausen/Thür. Abh Ber Naturk-Mus Gotha. 1967;4:9-32.

Kaiser H. Die Turbellarienfauna der limnokrenen Erdfallquellen und ihrer Abflüsse

im Raume Mühlhausen – Bad Langensalza. Abh Ber Naturk-Mus Gotha.

1969;5:33-56.

Kaiser H. Die Turbellarienfauna in salzhaltigen Gewässern und Quellregionen

Nordwest-Thüringens. Limnologica. 1974;9:1-62.

Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nature methods.

2017 May 8;14(6):587.

Karling T. Ein Beitrag zur Kenntnis der Nemertinen des nnischen Meerbusens. Mem

Soc Fauna Flora Fenn. 1934;10:76–90.

Karling T. Zur Morphologie und Systematik der Alloeocoela cumulata und

Rhabdocoela Lecithophora (Turbellaria). Acta Zool Fenn. 1940;26:1-260.

Karling T. On nematocysts and similar structure in turbellarians. Acta Zool Fenn.

1966;116:3-28.

Karling T. Turbellarian fauna of the Baltic proper. Identification, ecology and

biogeography. Fauna fenn. 1974;27:1-101

Keller R. Pädagogisch-psychometrische Studien. Biol Centralblatt Bd XIV. 1894;2.

Kieneke A, Nikoukar H. Integrative morphological and molecular investigation of

Turbanella hyalina Schultze, 1853 (Gastrotricha: Macrodasyida), including a

redescription of the species. Zool Anz. 2017;267:168-86.

Kolasa J. Ecological and faunistical characteristics of Turbellaria in the eutrophic

Lake Zbe☐chy. Acta Hydrobiol. 1979;21:435-59.

Kolasa J. On the origin of the stream interstitial microturbellaria. Pol Arch Hydrobiol.

1983;29:405-13

Kolasa J, Strayer J, Bannon-O'Donnell E. Microturbellarians from interstitial waters,

streams, and springs in southeastern New York. J N Am Benthol Soc.

1987;6:125-32.

Korgina Y. New and rare (for the Volga basin) members of the order

Lecithoepitheliata (Turbellaria). Inl Wat Biol. 2011;4:12-20.

Kosler A. Die Bodenfauna der Gewässer um Hiddensee. Wiss Z Ernst-Moritz-Arndt-

Univ Greifswald. 1962;11:57-64.

Kostenko A. On the species autonomy of the turbellarian *Microstomum giganteum*

(Turbellaria, Microstomidae). Vestn Zool. 1988;2:59-61

Kraus H. Zur Turbellarienfauna des Seewinkels im Neusiedlerseegebiet. Wiss Arb

Burgenland. 1965;32:60-115.

Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis

Version 7.0 for Bigger Datasets. Mol Biol Evol. 2016;33:1870-4.

Larsson K, Ahmadxadeh A, Jondelius U. DNA taxonomy of Swedish Catenulida

(Platyhelminthes) and a phylogenetic framework for catenulid classification.

Org Divers Evol. 2008;8:399-412.

Laumer C, Bekkouche N, Kerbl A, Goetz F, Neves R, Sørensen M, et al. Spiralian

phylogeny informs the evolution of microscopic lineages. Cur

Biol. 2015;25:2000-6.

Lee C, Bell M. Causes and consequences of recent freshwater invasions by saltwater

animals. Trends Ecol Evol. 1999;14:284-8.

Leidy J. Corrections and additions to former papers on Helminthology. Proc Acad Nat

Sci Phila. 1852;5:288-9

Levinsen G. Bidrag til kundskab om Groenlands turbellariefauna. Vidensk Medd

Dansk Naturhist Foren. 1879;3:165-204.

Lin Y, Zhuang J, Chen L, Zhong L, Wang A. First record of Microstomidae

(Platyhelminthes: Macrostomorpha) from China, and the phylogeny of one

new species of *Microstomum*. Sichuan J Zool. 2017a;36(4):441–9.

Lin Y, Fang W, Xin F, Zhang L, Zhang Y, Wang A. 2017b. Two new species and the

molecular phylogeny of eight species of *Macrostomum* (Platyhelminthes:

Macrostomorpha) from southern China. Zootaxa. 2017b;4337:423-9.

Littlewood D, Rohde K, Clough K. The interrelationships of all major groups of

Platyhelminthes: Phylogenetic evidence from morphology and molecules. Biol

J Linn Soc. 1999;66:75-114

Litvaitis M, Rohde K. A molecular test of platyhelminth phylogeny: Inferences from

partial 28S rDNA sequences. Invertebr Biol. 1999;118:42-56.

Luther A. Ueber die systematische Stellung der Rhabdocoelenfamilie Catenulidae.

Zool. Anz. 1907;31:718-23.

Luther A. Die Turbellarien Ostfennoskandiens I. Acoela, Catenulida, Macrostomida,

Lecithoepitheliata, Prolecithophora, Proseriata. Fauna Fenn. 1960;7:1-155.

Mack-Fira☐ V. Rhabdocoelen aus dem Überschwemmungsgebiet der Donau. Limnol

Berichte X Jubiläumstagung Donauforschung, Bulgarien, 10–20 Okt 1966.

1968;251-8.

Mack-Fira☐ V. Turbellaries du complexe lagunaire Razelm-Sinoe (Mer Noire). Rapp

Comm int Mer Médit. 1973;21:633-5.

Mack-Fira□ V. The turbellarian fauna of the Romanian littoral waters of the Black

Sea and its annexes. In: Riser N, Morse M, eds. Biology of the Turbellaria.

McGraw Hill Book Co.:New York. 1974;248-90.

Marcus E. Turbellaria Brasileiros (8). Bolm Fac Filos Ciênc Letr Univ S Paulo Zool.

1950;17:5-191.

Marcus E. 1951: Turbellaria Brasileiros (9). Bolm Fac Filos Ciênc Letr Univ S Paulo

Zool. 1951;16:5-215.

Markov M. [An outline of the fauna of the turbellarians of the Kharkiv and Poltava

provinces.] Uchen App Kharkiv University Ser Natural. Science. 1904;80.

Russian.

Martens P, Schockaert E. Sand dwelling Turbellaria from the Netherlands delta area.

Hydrobiologia. 1986;84:113-27.

Martens P. Comparison of three different extraction methods for Turbellaria. Mar

Ecol Prog Ser. 1984;14:229-34.

Meier R, Shiyang K, Vaidya G, Ng P. DNA barcoding and taxonomy in diptera: A

tale of high intraspecific variability and low identification success. System Biol.

2006;55:715-28.

Meixner J. Zur Turbellarienfauna der Ostalpen, insonderheit des Lunzer Seengebietes.

Zool Jb Syst. 1915;38:459-588.

Meixner J. Turbellaria (Strudelwürmer) I. Allgemeiner Teil. In: Grimpe G, Wagler E,

Remane A, Eds. Die Tierwelt der Nord- und Ostsee, VIb. Leipzig:

Akademischer Verlag. 1938. 146 pp.

Mereschkowsky K. [On some new Turbellaria of the White Sea]. Trudi d St

Petersburger Ges d Naturforscher. 1879;9:35-55. Russian.

Metschnikov E. [About the history of Rhabdocoels]. Ann Mag Nat Hist 3 Ser XVII.

1865. 57 pp. Russian.

Michaloudi E, Mills S, Papakostas S, Stelzer C, Triantafyllidis A, Kappas I, et al.

Morphological and taxonomic demarcation of *Brachionus asplanchnoidis*

Charin within the *Brachionus plicatilis* cryptic species complex (Rotifera,

Monogononta). Hydrobiologia. 2017;796(1):19-37.

Micoletzky H. Die Turbellarienfauna des Golfes von Triest. Arb Zool Inst Univ Wien.

1910;18:167-82.

Müller D, Faubel A. The 'Turbellaria' of the River Elbe Estuary. A faunistic analysis

of oligohaline and limnic areas. Arch Hydrobiol. 1993;75 Suppl:363-96.

Müller O. Vermium terrestrium et fluviatilium, seu animalium infusoriorum,

helminthicorum et testaceorum, non marinorum succincta historia. Havniae et

Lipsiae. 1773. 80 pp.

2451

Müller O. Zoologiae danicae prodromus, seu animalium Daniae et Norvegiae

indigenarum characteres, nomina et synonyma imprimmis popularium.

Havniae. 1776;221-82.

Nasonov N. Contributions à la faune des Turbellaria de la Russie. I. Bull Acad Sci

Russie. 1919a;13:619-46.

Nasonov N. Contributions à la faune des Turbellaria de la Russie. IV. Bull Acad Sci

Russie. 1919b;13:1179-97.

Nasonov N. Les trait généraux de la distribution géographique des Turbellaria

Rhabdocoelida dans la Russie d'Europe. Bull Acad Sci Russie. 1924;18:327-

52.

Nasonov N. Sur la faune des Turbellaria de la péninsule de Kola. Bull Acad Sci

Russie. 1925;53-74.

Nasonov N. Zur Fauna der Turbellaria Rhabdocoelida der japanischen

Süßwasserbecken. Comptes Rendus Acad Sc URSS. 1929;423-8.

Nasonov N. Über den Heliotropismus der Turbellaria Rhabdocoelida des Baikalsees.

Trudy Lab. Eksper. Zool Morf Zhivot Akad Nauk SSSR 1935;4:195-204.

Nguyen L, Schmidt H, Haeseler A, Minh B. IQ-TREE: A Fast and Effective

Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol

Biol Evol. 2015;32:268–74.

Noreña C. Studies on the taxonomy and ecology of the Turbellaria (Plathelminthes) in

the floodplain of the Paraná river (Argentina). II. Taxonomy and ecology of

the Turbellaria. Arch Hydrobiol. 1995; 107 Suppl:211-62.

Noreña C, Campo A, Real M. Taxonomy and morphology of limnic

microturbellarians (Plathelminthes) in Extremadura (Spain). Hydrobiologia

1999;397:21-8.

Noreña C, Damborenea C, Brusa F, Escobedo M. Free-living Platyhelminthes of the

Pacaya-Samiria National Reserve, a Peruvian Amazon floodplain. Zootaxa.

2006;1313.

Noreña C, Damborenea C, Faubel A, Brusa F. Composition of meiobenthonic

Platyhelminthes from brackish environments of the Galician and Cantabrian

coasts of Spain with the description of a new species

of Djeziraia (Polycystididae, Kalyptorhynchia). J Nat Hist. 2007;41:29-32.

Okugawa K. A Monograph of Turbellaria (Acoela, Rhabdocoela, Alioeocoera and

Tricladida) of Japan and its adjacent regions. Kyoto Gakugei Univ Ser B.

1953;3:20-43.

Oersted A. Forsög til en ny classification af planarierne (Planaria Dugès) grundet paa

mikroskopisk-anatomiske undersögelser. Naturh Tidsskr. 1843;4:519-81.

Oersted A. Fortegnelse over Dyr, samlede Christianiafjord ved Dröbak fra 21-24 Juli

1844. Naturh Tidsskr. 1845;2.

Papi F. Note faunistiche sui Turbellari dell'Italia centrale. Monit Zool Ital. 1952;60:3-

13.

Paradi K. Die in der Umgebung von Klaussenburg gefundenen rhabdocoelen

Turbellarien. Med-natunwiss Anz. 1881;6:161-74.

Pérez-Ponce de León, G., Poulin R. Taxonomic distribution of cryptic diversity

among metazoans: Not so homogeneous after all. Biol Let. 2016;12(8),

20160371.

Petrovna T, Vladimirovich B. Ciliated worms (Platyhelminthes, "Turbellaria") of the

Kaban Lake System (Kazan). Sci Not Kazan Univ, Ser Nat Sci. 2013;195-9.

Russian.

Pons J, Barraclough T, Gomez-Zurita J, Cardoso A, Duran D, Hazell S, et al.

Sequence-based species delimitation for the DNA taxonomy of undescribed

insects. System Biol. 2006;55:595-609.

Pörner H. Die rhabdocoeliden Turbellarien der Gewässer von Jena und Umgebung.

Limnologica. 1966;4:27-44.

Reise, K. Free-living Platyhelminthes (Turbellaria) of a marine sand flat: An

ecological study. Microfauna Mar. 1984;1:1-62.

Reisinger E, Steinböck O. Zur Turbellarienfauna des Wörthersees. Carinthia II.

1925;114/115:109-14.

Reisinger E. Turbellarien der deutschen limnologischen Sunda-Expedition. Arch

Hydrobiologie. 1933;4 Suppl 12:239-62.

Reuter M. Scanning and transmission electron microscopic observations on surface

structures of three turbellarian species. Zool Scr. 1978;7:5-11.

Reuter M, Palmberg I. Development and differentiation of neuronal subsets in

asexually reproducing *Microstomum lineare*. Immunocyctochemistry of 5-HT,

RF-amide and SCP$_B$. Histochemistry. 1989;91:123-31.

Reyes J, Brusa F. Species of *Macrostomum* (Macrostomorpha: Macrostomidae) from

the coastal region of Lima, Peru, with comments on *M. rostratum* Papi, 1951.

Zootaxa. 2017;4362:246-58.

Riedel G. Ergebnisse einer von E. Reisinger und O. Steinböck mit Hilfe des Rask-

Örsted-Fonds durchgeführten zoologischen Reise in Grönland 1926. 3.

Macrostomida. Vidensk Medd Dansk Naturhist Foren. 1932;94:33-90.

Riedl R. Zur Kenntnis der Turbellarien adriatischer Schlammböden sowie ihre

geographische und faunistische Verbreitung. Thalassia Jugosl. 1956;1:69-184.

Riedl R. Turbellarien aus submarinen Höhlen. 1. Archoophora. Ergebnisse der Öster-

reichischen Tyrrhenia-Expedition 1952, Teil VII. Pubbl Staz Zool Napoli.

1959;Suppl 30:178-208.

Rieger R. Phylogenetic systematics of the Macrostomorpha. In: Littlewood D. & Bray

R, Eds. Interrelationships of the Platyhelminthes. New York :Taylor & Francis

Inc. 2001;28-38.

Rixen J. Kleinturbellarien aus dem Litoral der Binnengewässer Schleswig-Holsteins.

Arch Hydrobiol. 1961;57:464-538.

Rixen J. Beitrag zur Kenntnis der Turbellarienfauna des Bodenseegebietes. Arch

Hydrobiol. 1968;64:335-65.

Ročník L. [Zoobenthos of the Lednické rybníky ponds]. Acta Univ Agric Silvic

Mendel Brun. 2007;2:85–94. Russian.

Rogozin A. New species of turbellaria in the fauna of the Urals. Biol Bull.

2015;42:755-8.

Rogozin A. Fauna turbellaria of the Ilmensky Reserve (South Urals). Act Prob

Human Nat Sci. 2016;8:44-7.

Sabussow H. Turbellarien-Studien 1. Ueber den Bau der männlichen

Geschlechtsorgane von *Stenostoma leucops* O. Schm. Zool Jahrb Anat.

1897;10:47-54

Sabussow H. Beobachtungen iiber die Turbellerien der Inseln von Solowetzk. Trudi

Kazan Univ. 1900. 34 pp.

Sahraean N, Van Campenhout J, Rigaux A, Mosallanejad H, Leliaert F, Moens T.

Lack of population genetic structure in the marine nematodes *Ptycholaimellus*

pandispiculatus and *Terschellingia longicaudata* in beaches of the Persian

Gulf. Iran Mar Ecol. 2017;38(3)1-13.

Sauer J, Hausdorf B. A comparison of DNA-based methods for delimiting species in a

Cretan land snail radiation reveals shortcomings of exclusively molecular

taxonomy. Cladistics. 2012;28:300-16.

Schmarda L. Neue wirbellose Thiere beobachtet und gesammelt auf einer Reise um

die Erde 1853 bis 1857. Turbellarien, Rotatorien und Anneliden. Wilhelm

Engelmann, Leipzig. 1859. 66 pp.

Schmidt E. Die rhabdocoelen Strudelwürmer (Turbellaria Rhabdocoela) des süssen

Wassers. Verlag von Friedrich Mauke, Jena. 1848. 65 pp.

Schockaert E. Marine Macrostomorpha (Platyhelminthes, Rhabditophora) from the

Algarve (Southern Portugal). Zootaxa. 2015;3872:577–90.

Schön I, Martens K, Dijk P. Lost sex. The evolutionary biology of parthenogenesis.

The Netherlands: Springer. 2009. 615 pp.

Schwank P. Quantitative Untersuchungen an litoralen Turbellarien des Bodensees. J

Ges Naturkde Württ. 1976;131:163-81.

Seifert R. Die Zusammensetzung der Bodenfauna der Hiddenseer Boddengewässer.

Mitt Naturwiss Verein Neuvorpommern Rügen. 1939;67:1-33.

Sibiriakova O. La faune Turbellaria Rhabdocoelida du fleuve Angara. Russ

Hydriobiol Z. 1929;8:237-50.

Simon J, Rispe C, Sunnucks. Ecology and evolution of sex in aphids. Trends Ecol

Evol. 2002;17:34-9.

Simon J, Delmotte F, Rispe C, Crease T. Phylogenetic relationships between

parthenogens and their sexual relatives: the possible routes to parthenogenesis

in animals. Biol J Linn Soc. 2003;79:151-63.

Southern R. Platyhelmia. Clare Ireland Survey pt. 56. Proc Roy Irish Acad.

1912;31:1-18.

Southern R. Turbellaria of Ireland. Proc Roy Irish Acad Sect B. 1936;43:43-72.

Steinböck O. Marine Turbellaria. VIII. In: Jensen A, Lundbeck W, Eds. The Zoology

of the Faeroes. 1931. 26

pp.

Steinböck O. Zur Turbellarienfauna der Südalpen, zugleich ein Beitrag zur

geographischen Verbreitung der Süßwasserturbellarien. Zoogeographica.

1933;1:209-62.

Steinböck O. Marine Turbellaria. Zool Iceland II. 1938;9:1-26.

Steinböck O. Zur Turbellarienfauna des Lagio Maggiore und des Lago di Como.

Mem Ist Ital Idrobiol. 1949;5:231-54.

2585

Steinböck O. Turbellarienstudien am Lago Maggiore. I. Mem Ist Ital Idrobiol.

1951a;6:137-64.

Steinböck O. Turbellarienstudien am Lago Maggiore. II. Mem Ist Ital Idrobiol.

1951b;6:165-76.

Stirewalt M. *Microstomum bispiralis* n. sp. Zool Anz. 1937;119:314-21.

Sun T, Zhang L, Wang A, Zhang Y. Three new species of freshwater *Macrostomum*

(Platyhelminthes, Macrostomida) from southern China. Zootaxa

2014;4012:120–34.

Telford M, Herniou E, Russell R, Littlewood D. Changes in mitochondrial genetic

codes as phylogenetic characters: two examples from the flatworms. Proc Natl

Acad Sci USA. 2000;97:1359-64.

Templeton A, Crandall K, Sing C. A cladistic analysis of phenotypic associations

with haplotypes inferred from restriction endonuclease mapping and DNA-

sequence data III. Cladogram estimation. Genetics. 1992;132:619-33.

Tokinov R, Berdnik S. [Microturbellaria (Platyhelminthes: Catenulida,

Rhabditophora) Nature Reserve "Blue Lakes" (Middle Volga Region)]. Russ J

Appl Ecol. 2016;123:15-8. Russian.

Tu T. Notes on some turbellarians from the Tsing Hua Campus. Sci Rep Nat Tsing

Hua Univ. 1934;1:191-205.

Uljanin W. Turbellarien der Bucht von Sewastopol. Arb 2 Vers Russ Naturf Moskau.

1870;2.

Urban-Malinga B. Free-living interstitial Platyhelminthes of the Baltic Sea: diversity

and abundance. Polish J Ecol. 2011;3:623-9.

Valkanov A. Beitrag zur Süsswasser. Fauna Bulgariens. Arb Bulg Naturforsch Ges.

1926;12:183-6.

Vara D, Leal-Zanchet A. Turbelários límnicos (Platyhelminthes) em ecossistemas de

arroz irrigado de Planície Costeira do sul do Brasil. Biota Neotropica.

2013;13:241-50.

Verheye M, Backeljau T, d'Acoz C. Looking beneath the tip of the iceberg:

Diversification of the genus *Epimeria* on the Antarctic shelf (Crustacea,

Amphipoda). Polar Biol. 2016;39(5):925–45.

Verrill A. Marine planarians of New England. Trans Connect Acad Arts Sci.

1892;8:458-520.

Volz W. Über neue Turbellarien aus der Schweiz. Zool Anz. 1898;21:605-12.

Wegelin R. Beitrag zur Kenntnis der Grundwasserfauna des Saale-Elbe-

Einzuggebietes. Zool Jb Syst. 1966;93:1-117.

Wehrenberg C, Reise K. Artenspektrum und Abundanz freilebender Plathelminthes in

sublitoralen Saenden der Nordsee bei Sylt. Microfauna Mar. 1985;2:163-80.

Westblad E. Marine Macrostomida (Turbellaria) from Scandinavia and England. Ark

Zool. 1953;4:391-408.

Westblad E. Some Hydroidea and Turbellaria from western Norway. Årbok Univ

Bergen 1954, Naturvit Rekke 1954;10:1-22.

Woodworth W. Report on the Turbellaria collected by the Michigan State Fish

Commission during the summers 1893 and 1894. Bull Mus Comp Zool

Harvard College 1896;29:239-43.

Yang Z, Rannala B. Bayesian species delimitation using multilocus sequence data.

Proc Nat Acad Sci USA, 2010;107:9264-9.

Yanoviak S. The macrofauna of water-filled tree holes on Barro Colorado Island,

Panama. Biotropica 2001;33:110-20.

Young J. British and Irish freshwater Microturbellaria: Historical records, new

records and a key for their identification. Arch Hydrobiol. 1970;87:210-41.

Young J. The occurrence of Microturbellaria in some British lakes of diverse

chemical content. Arch Hydrobiol. 1973;72:202-24.

Zabusov I. Microstomidae O. Schm. in the vicinities of Kazan, Tr. Kazan. Obshch

Estestvoisp Kazan Univ. 1894;27:1–36.

Zacharias O. Faunistische Mittheilungen. Forschungsber Biol Stat Plön Th 2. Berlin

1894.

Zeppilli D, Sarrazin J, Leduc D, Arbizu P, Fontaneto D, Fontanier C, et al. Is the

meiofauna a good indicator for climate change and anthropogenic impacts?

Mar Biodivers, 2015;45:505-35.

Zhang J, Kapli P, Pavlidis P, Stamatakis A. A general species delimitation method

with applications to phylogenetic placements. Bioinformatics, 2013;29: 2869-

76.

Zhao D, Kong L, Yu H, Li Q. Cryptic genetic diversity of *Neverita didyma* in the

coast of China revealed by phylogeographic analysis: implications for

management and conservation. Conserv Gen. 2018;19(2):275-82.

Zykov W. Beiträge zur Turbellarienfauna Russlands I. Zool Anz. 1897;20:450–2.

Zykov W. Beiträge zur Turbellarienfauna Russlands II. Zool Anz. 1902;25:478-80.

2654

2655

2656   Figure and Supplimental Captions

2657

2658   **Fig. 1: Concatenated 18S and CO1 gene tree summary.** Percent bootstrap values

2659   are given at each node.

2660

2661   **Fig. 2: bPTP analysis of 18S and CO1 gene tree**. A maximum of five specimens

2662   were chosen for each species. Specimens are listed with the isolate label in

2663   parentheses. At each node, percent bootstrap values (black above) and bPTP (red

2664   below) values are given. Transitions from blue to red lines highlight species according

2665   to the results of the bPTP analysis.

2666

2667   **Fig. 3:** *Microstomum afzelii* **sp. nov.** A. Drawing of ventral view. B.

2668   Micropictograph of side view. ap adhesive papillae; b brain; c ciliary pits; i intestine;

2669   m mouth; n nematocysts; ph pharynx; r rhabdites

2670

2671   **Fig. 4:** *Microstomum curinigalletti* **sp. nov.** A. Drawing of whole body. B.

2672   Micropictograph of whole body C. Anterior end with focus on the ciliary pits. D.

2673   Posterior end. ap adhesive papillae; b brain; c ciliary pits; i intestine; m mouth; ph

2674   pharynx; r rhabdites

2675

2676   **Fig. 5:** *Microstomum inexcultus* **sp. nov.** A. Drawing of whole body. B.

2677   Micropictograph of whole body C. Anterior end. D. Posterior end with focus on

2678   adhesive papillae. ap adhesive papillae; b brain; c ciliary pits; i intestine; m mouth; n

2679   nematocysts; ph pharynx

2680

2681   **Fig. 6:** *Microstomum lotti* **sp. nov.** A. Drawing of whole body. B. Micropictograph of

2682   whole body C. Anterior end with focus on adhesive papillae. D. Anterior end with

2683   focus on ciliary pits E. Posterior end with focus on adhesive papillae. ap adhesive

2684   papillae; b brain; c ciliary pits; i intestine; m mouth; ph pharynx; r rhabdites

2685

2686   **Fig. 7:** *Microstomum marisrubri* **sp. nov.** A. Drawing of whole body. B.

2687   Micropictograph of whole body C. Posterior end with focus on the nematocysts. D.

2688   Posterior end with focus on the adhesive papillae. E. Male stylet. ap adhesive

2689   papillae; b brain; c ciliary pits; i intestine; m mouth; mp male pore; n nematocyst; ph

2690   pharynx; r rhabdites; s male stylet; t testis; vg vesicula granulorum

2691

2692   **Fig. 8:** *Microstomum schultei* **sp. nov.** A. Drawing of whole body. B.

2693   Micropictograph of whole body C. Anterior end with focus on the ciliary pits. D.

2694   Male stylet. E. Sperm. ap adhesive papillae; b brain; c ciliary pits; i intestine; m

2695   mouth; o ovary; ph pharynx; r rhabdites; s male stylet; vg vesicula granulorum

2696

2697   **Fig. 9:** *Microstomum weberi* **sp. nov.** A. Drawing of whole body. B.

2698   Micropictograph of whole body C. Anterior end side view. D. Anterior end with focus

2699   on adhesive papillae. ap adhesive papillae; b brain; c ciliary pits; i intestine; m mouth;

2700   ph pharynx; r rhabdites

2701

2702   **Fig. 10:** *Microstomum westbladi* **nom. nov.** A. Drawing of whole body. B.

2703   Micropictograph of whole body C. Anterior end focus on adhesive papillae. D.

2704   Anterior end with focus on rhabdite field. ap adhesive papillae; b brain; c ciliary pits;

2705   i intestine; m mouth; ph pharynx; r rhabdites; s male stylet

2706

Fig. 11: Different shapes of the anterior ends within *Microstomum*. A. Drawing

depicting the elongated frontal end of *M. trichotum*. B. Drawing depicting an anterior

proboscis. C. Micropictograph of the proboscis of *M. compositum*. D. Drawing

depicting a conical anterior. E. Micropictograph of the conical anterior of *M. lineare*.

F. Micropictograph of the conical anterior of *M. bispiralis*. G. Drawing depicting a

blunt anterior. H. Drawing depicting a rounded anterior. I. Micropictograph depicting

the rounded anterior of *M. septentrionale*. b brain; ph pharynx

2714

Fig. 12: Different shapes of the posterior ends within *Microstomum*. A. Drawing

depicting a rounded posterior. B. Drawing depicting a blunt posterior. C.

Micropictograph of the blunt posterior of *M. edmondi*. D. Drawing depicting a thin-

tail posterior of e.g. *M. giganteum*. E. Drawing depicting a spatulate posterior. F.

Micropictograph of the spatulate posterior of *M. lotti*. G. Drawing depicting a tail-like

posterior. H. Drawing depicting a pointed posterior. I. Micropictograph depicting the

pointed posterior of *M. bispiralis*.

2722

2723 **Fig. 13: Different shapes of the ciliary pits within *Microstomum*.** A. Drawing

2724 depicting bottle-shaped ciliary pits. B. Micropictograph of the bottle-shaped ciliary

2725 pits of *M. edmondi*. C. Drawing of rounded ciliary pits. D. Micropictograph of the

2726 rounded ciliary pits of *M. rubromaculatum*. Arrows indicate ciliary pits; b brain; ph

2727 pharynx

2728

Fig. 14: Different lengths of the preoral intestine within *Microstomum*. A.

Drawing depicting a long preoral intestine that extends anterior to the brain. B.

Micropictograph of the long preoral intestine of *M. inexcultus* sp. nov. C. Drawing

depicting a medium preoral intestine that terminates at the level of the brain. D.

Micropictograph of the medium preoral intestine of *M. septentrionale*. E. Drawing

depicting a short preoral intestine that terminates posterior to the brain. D.

Micropictograph of the short preoral intestine of *M. bispiralis*. b brain; i intestine; m

mouth; ph pharynx

2737

2738 **Fig. 15: Different shapes of the pharynx within *Microstomum*.** A. Drawing

2739 depicting the characteristic biparte pharynx of e.g. *M. breviceps*. B. Drawing

2740 depicting a typically-shaped pharynx of *Microstomum* C. Micropictograph of the

2741 typically-shaped pharynx of *M. bispiralis*. b brain; i intestine; m mouth; ph pharynx

2742

2743 **Fig. 16: *Microstomum bispiralis*; specimen from Sweden.** A. Micropictograph of

2744 whole body B. Anterior end. C. Posterior end with focus on sexual anatomy. b brain;

2745 c ciliary pits; i intestine; mp male pore; n nematocyst; ph pharynx; s male stylet; t

2746 testis; vg vesicula granulorum

2747

2748 **Fig. 17: Examples of limnic *Microstomum*.** A. *M. lineare* B. *M. zicklerorum* C. *M.*

2749 *tchaikovskyi* D. *M. artoisi* E. *M. bispiralis*. Bar = 200 µm

2750

2751 **Fig. 18: Examples of marine *Microstomum*.** A. *M. edmondi* B. *M. laurae* C. *M.*

2752 *rubromaculatum* D. *M. papillosum* E. *M. crildensis* F. *M. compositum* G. *M.*

2753 *septentrionale.* Bar = 100 µm

2754

2755 **Fig. S1: Concatenated 18S and CO1 gene tree with all specimens**

2756

2757 **Table S1. Collecting information for the specimens used in this study**. Collecting

2758 location, specific coordinates and dates are given for each specimen, where available.

2759 In addition, the GenBank Accession Numbers for 18S, and CO1 sequences and

2760 references are listed.

2761

2762   **Table S2: Primer sequences, references and protocols for amplification of 18S**

2763   **and CO1 sequences.**

2764

2765   **Table S3: A summary of the morphological characters for each species**.

A

ap

b

c

m

ph

i

r

50 μm

B

m

ph

r

i

ap

50 μm

C

ap

c

m

i

25 μm

D

ap

25 μm