Modified TCA/acetone precipitation of proteins for proteomic analysis

Liangjie Niu[§], Hang Zhang[§], Hui Liu, Xiaolin Wu, Wei Wang

State Key Laboratory of Wheat and Maize Crop Science, Collaborative Innovation Center of Henan Grain Crops,

College of Life Sciences, Henan Agricultural University, Zhengzhou, China

§The co-first authors.

*Corresponding author at: College of Life Sciences, Henan Agricultural University, Zhengzhou 450002, China. E-

mail address: wangwei@henau.edu.cn (W. Wang).

Abstract

Protein extracts obtained from cells or tissues often need to remove interfering substances for preparing high-quality protein samples in proteomic analysis. A number of protein extraction methods have been applied to various biological samples, especially TCA/acetone precipitation and phenol extraction. TCA/acetone precipitation, as a common method, is thought to minimize protein degradation and proteases activity as well as reduce contaminants like salts and polyphenols. However, the TCA/acetone precipitation relies on the completely pulverizing and repeatedly rinse of tissue powder to remove the interfering substances, which is laborious and time-consuming. In addition, a prolonged incubation in TCA/acetone or acetone can lead to the modifications and degradation of proteins, and the precipitated proteins are more difficult to re-dissolve. We have described a modified TCA/acetone precipitation of plant proteins for proteomic analysis. Proteins of cells or tissues were extracted using SDS-containing buffer, precipitated with equal volume of 20% TCA/acetone, and washed with acetone. Compared to classical TCA/acetone precipitation, this protocol generates comparable yields, spot numbers, and proteome profiling, but takes less time (ca. 45 min), thus avoiding excess protein modification and degradation after extended-period incubation in TCA/acetone or acetone. The modified TCA/acetone precipitation method is simple, fast, and suitable for various plant tissues in proteomic analysis.

Keywords: Protein extraction, TCA/acetone precipitation, Removal of interfering substances, SDS sample buffer; 2DE; MALDI-TOF mass spectrometry

Background

Protein extracts obtained from cells or tissues often contain interfering substances, which must be removed for preparing high-quality protein samples [1]. In particular, plant tissues contain a diverse group of secondary compounds, such as phenolics, lipids, pigments, organic acids and carbohydrates, which greatly interfere with protein extraction and proteomic analysis [2]. Thus, sample quality is critical for the coverage, reliability, and throughput of proteomic analysis, protein extraction in proteomics remains a challenge, even though advanced detection approaches (especially LC-MS/MS) can greatly enhance the sensitivity and reliability of protein identification [3].We are always interested in developing protein extraction protocols for 2DE-based proteomics. In last decade, Wang and the colleagues have established many such protocols as integrating TCA/acetone precipitation with phenol extraction [4-8]. A protein extraction protocol that can be universally applied to various biological samples with minimal optimization remains essential in current proteomics.

A number of methods are available for concentrating dilute protein solutions and simultaneously removing interfering substances, especially TCA/acetone precipitation and phenol extraction [9-11]. TCA/acetone precipitation is a common method for precipitation and concentration of total proteins, initially developed by Damerval et al. [12] and then modified for use in various tissues [10, 13-19].

TCA/acetone precipitation is thought to minimize protein degradation and proteases activity as well as reduce contaminants such as salts or polyphenols [20]. During acetone/TCA precipitation, organic-soluble substances are rinsed out, leaving proteins and other insoluble substances in the precipitate, and proteins are extracted using a buffer of choice [2, 5, 9]. The success of TCA/acetone precipitation relies on the completely pulverizing and repeatedly rinse of tissue powder to remove the interfering substances, which is laborious and time-consuming.

However, a prolonged incubation of tissue powder in TCA/acetone or acetone can lead to the

modifications of proteins by acetone and the proportion of modified peptide increased over time,

thus affecting the outcome of MS/MS analysis [21]. Moreover, long exposure to the acidic pH in

TCA/acetone probably causes protein degradation [6, 22]. Alternatively, protein extract can be

precipitated using aqueous 10% TCA [23], but the TCA precipitated proteins are more difficult to

dissolve and need using NaOH to increase their solubilization [24]. Therefore, aqueous TCA

precipitation is not commonly used like TCA/acetone precipitation in proteomic analysis.

To overcome the limitations of aqueous TCA precipitation and TCA/acetone precipitation, we

here report a modified, rapid method of TCA/acetone precipitation of plant proteins for proteomic

analysis. We systematically compared the modified and classical TCA/acetone precipitation

regarding protein yields and proteome profiles.

Methods

Materials

Maize (*Zea mays* L.) seeds were soaked in water for 2 h to soften seed coats and endosperms, and

the embryos were manually dissected and used for protein extraction. Leaves and roots (root tips,

ca. 1 cm) of maize seedlings were sampled and used for protein extraction.

Escherichia coli BL21 was cultivated at 37°C for 12 h with good aeration until 0.8 optical density.

The cells were collected by centrifugation (1,000 g, 3 min, 4°C), rinsed with phosphate-buffered

saline thrice, and used for protein extraction.

Modified TCA/ Acetone precipitation of proteins

The modified protocol is designed for running in eppendorf tubes within 1 h, and can be reliably

adapted to big volumes. It includes protein extraction, precipitation, and dissolving. The detailed

steps were given in Fig. 1. Maize tissues and *E. coli* cells were used for evaluating the protocol.

Organic solvents were pre-cooled at -20°C and were supplemented with 5 mM dithiothreitol (DTT)

before use. The subsequent steps were carried out at 4°C unless otherwise indicated.

Protein extraction. Maize embryos (0.2g), leaves (0.4g), and roots (0.4g) and *E. coli* cells (0.2g)

were homogenized in a pre-cooled mortar (interior diameter 5 cm)on ice in 2.0 ml of the

extraction buffercontaining1% SDS, 0.1 M Tris-HCl (pH 6.8), 2 mM EDTA-Na$_2$, 20 mM DTT and 2

mM PMSF (adding before use). The homogenate was transferred into eppendorf tubes and

centrifuged at 15,000 g for 5 min. Then, the supernatant (protein extract) was pipetted into fresh

tubes.

Protein precipitation. Add cold 20% TCA/acetone into the protein extract (1: 1, v/v, with a final

10% TCA/50% acetone), place the mixture on ice for 5 min, centrifuge the mixture (15000g, 3 min),

and discard the supernatant. Wash protein precipitate with 80% acetone, followed by

centrifugation as above. Repeat the wash step once or more.

Protein dissolving. Air-dry the protein precipitates shortly (1- 3 min), and dissolve in a buffer of

choice for SDS-PAGE, IEF or iTRAQ analysis. Notably, do not over-dry the precipitates, otherwise

they will be more difficult to resolubilize.

As a parallel experiment, the classical TCA/ acetone method was exactly done according to

previously described [10].

Protein assay

For SDS-PAGE, protein precipitates were dissolved in a SDS-containing buffer (0.5% SDS, 50mM

Tris-HCl, pH 6.8, and 20mM DTT). Protein concentration was determined by the Bradford assay

(Bio-Rad, Hercules, CA) [25]. Prior to SDS-PAGE, protein extracts were mixed with appropriate

volume 4 x SDS sample buffer [26]. For 2DE, protein precipitates were dissolved in the 2DE

rehydration solution without IPG buffer to avoid its interference [27], and protein concentrations

were determined by the Bradford Assay. After then, the IPG buffer was supplemented into protein

samples to 0.5% concentration.

SDS-PAGE

SDS-PAGE was run using 12.5% gel [28] and protein was visualized using Coomassie brilliant blue (CBB) G250.

2-DE and MS/MS

Isoelectric focusing (IEF) was performed using 11-cm linear IPG strips (pH 4–7, Bio-Rad). Approximately 600 μg of proteins in 200 μl of the rehydration solution was loaded by passive rehydration with the PROTEAN IEF system (Bio-Rad) for 12 h at 20°C. IEF and subsequent SDS-PAGE, and gel staining were performed as previously described [28].Digital 2-DE images were processed and analyzed using PDQUEST 8.0 software (Bio-Rad). Protein samples were analyzed by 2DE in three biological replicates.

The spots with at least2-fold quantitative variations in abundance among maize embryos, leaves and roots by two methods respectively were selected for mass spectrometry (MS) analysis. One-way ANOVA was performed based on three biological replications.

The selected protein spots were extracted, digested and analyzed by the MALDI-TOF/TOF analyzer (AB SCIEX TOF/TOF-5800, USA) as described previously [29]. MALDI-TOF/TOF spectra were acquired in the positive ion mode and automatically submitted to Mascot 2.2 (http://www.matrixscience.com) for identification against NCBInr database (version Feb 17, 2017; species, *Zea mays*, 279566 sequences). Only significant scores defined by Mascot probability analysis greater than "identity" were considered for assigning protein identity. All of the positive protein identification scores were significant (p< 0.05).

Bioinformatics Analysis

The unidentified proteins were searched by BLAST using Universal Protein (http://www.uniprot.org/) or National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov/). Grand average of hydropathicity (GRAVY) analyses were

performed using ProtParam tool (http://web.expasy.org/protparam/). Subcellular localization

information was predicted using the online predictor Plant-mPLoc

(http://www.csbio.sjtu.edu.cn/bioinf/plant-multi/).

Results

The development of the modified TCA and acetone precipitation

Rather than preparing tissue powder by extensively TCA/acetone rinse in the classical method, we

directly extracted proteins in cells or tissues and then precipitated proteins in the extracts with

equal volume of 20% TCA/acetone. Various aqueous buffers can be used for protein extraction,

but the composition of the extraction buffers will greatly affect the profiles of the extracted

proteome. The Laemmli's SDS buffer [24] was used for protein extraction, because SDS-containing

buffer can enhance protein extraction and solubility, especially under heating.

The ratio of TCA and acetone was optimized in initial tests (Fig. 2). We compared the effect of

10% (w/v) TCA in various concentrations (0-80%, v/v) of aqueous acetone and the effect of various

TCA concentrations in aqueous 50% acetone on protein extraction and separation. The presence

of TCA could improve protein resolution by SDS-PAGE, but no significant differences were

observed among 5-20% TCA (Fig. 2A). At a fixed 10% TCA, protein patterns were quite similar

among different acetone ratio, but the background was clearer with the increased acetone ratio

(Fig. 2B). Based on the quality of protein gels, we choose the low ratio combination of 10% TCA/50%

acetone (final concentration) for further testing.

Different from aqueous TCA precipitation, proteins precipitated by 10% TCA/50% acetone are

easy to dissolve. After incubated on ice for 10 min, protein precipitates are recovered by

centrifugation, and washed with 80% acetone thrice to remove residual TCA in the precipitated

protein. Finally, the air-dried protein precipitates are dissolved in a buffer of choice for SDS-PAGE,

IEF or iTRAQ analysis.

Comparison of the performance in both methods

We made a comprehensive comparison of protein yields and resolving in 2-DE by the modified method and the classical method. Regarding protein yields, the former was slightly higher, but did not significantly differ from the classical method (t-test, $p < 0.05$). Similar results were obtained by comparison of spot numbers in 2D gels between both methods (Table 1). Compared with other separation methods (e.g. SDS-PAGE, LC), 2DE analysis can display visible results regarding the quality of protein samples.

For all materials tested, 2D gels obtained with the two methods were generally comparable regarding the number, abundance and distribution of protein spots, without profound deviations (Fig.3-5); however, several protein spots exhibited at least 2-fold differences in abundance. For example, spots 1, 2 and 4 were more abundant in maize roots by the modified method, whereas the classical method resulted in more abundant spot 3. Of the 11 differential abundance proteins (DAPs) selected for MS/MS identification, nine were identified with MS/MS analysis (Table 2). In particular, the modified method selectively depleted globulin-1 in maize embryos (Fig. 4). Our recent studies showed that globulin-1 (also known as vicilin) was the most abundant storage proteins in maize embryos [11, 30], and selective depletion of globulin-1 improved proteome profiling of maize embryos [11]. In addition, the modified protocol also produced good 2DE maps in *E. coli* cells (Fig. 6).

It needed to note that the aqueous TCA precipitation cause severely-denatured proteins that were very difficult to dissolve, and was rarely used in proteomic analysis. Thus, we did not compare it with the modified method in the present study.

Discussion

The classical TCA/acetone precipitation applies a strategy of removal of interfering substances before protein extraction, involving incubation for extended periods (from 45 min to overnight) in

TCA/acetone and between the rinsing steps[10,12,13,15,20]. These steps can lead to the

modifications of proteins by acetone [21] or possible protein degradation after long exposure to

harsh TCA/acetone [6, 23, 31], thus affecting the outcome of MS/MS analysis.

In contrast, the modified method here is using a strategy of removing interfering substances

after protein extraction, taking less time to avoid protein modifications by acetone, but producing

similar or better results regarding protein yields, 2DE spot numbers and proteome profiling. The

resultant protein precipitates are easy to wash using acetone compared to tissue powder in the

classical TCA/acetone precipitation method.

Previously, Wang et al. [29] observed that in oil seed protein extraction uses 10% TCA/ acetone,

rather than aqueous TCA, because the former resulted in protein precipitates easily to dissolve in

SDS buffer or 2-D rehydration buffer. The combination of TCA and acetone is more effective than

either TCA or acetone alone to precipitate proteins [17, 32]. It is noted that some proteins were

preferably extracted by the modified or the classical method, but the rationale remains open to

question. Recently, we reported a chloroform-assisted phenol extraction method for depletion of

abundant storage protein (globulin-1) in monocot seeds (maize) and in dicot (soybean and pea)

seeds [8]. The modified method exhibited also highly efficient to deplete globulin-1, suggesting

another application of the modified method in proteomic analysis.

In the modified method, final protein pellets were dissolved in the same 2DE buffer as in the

classical method, so protein profiles highly depends on the extraction efficiency in the SDS buffer

and precipitation efficiency by 20% TCA/acetone. We observed some repeatedly, significant

differences in abundance of several DAPs between both methods. There were specific DAPs

associated with each method in different samples. However, the rationale behind this

phenomenon remains unclear. We tried to analyze the hydropathicity, physicochemical property,

and Subcellular compartments of these DAPs (Table 2); however, no a clearly conclusion could be

drawn. Reasonably, different extract methods can produce protein profiles with substantial or

subtle differences [33], but these inherent differences were yet difficult to explain, as discussed

at a previous study [15].

To summarize, the greatest advantage of the modified method is simple and fast. Despite the

similarity in steps to aqueous TCA precipitation, but it overcame the latter's drawback that TCA-

precipitated proteins are difficult to dissolve. Moreover, the modified method uses equal volume

20% TCA/acetone to precipitate proteins, and can handle bigger volume of protein extracts than

acetone precipitation in a microtube. Because the modified method precipitates proteins in

aqueous extracts, it is expected to be universally applicable for various plant tissues in proteomic

analysis.

Declarations

Authors' contributions

LJN and WW drafted the manuscript. HZ and LN performed the experiments. All authors contributed the experimental design and data analysis.

Acknowledgements

Financial support for this study was provided by National Natural Science Foundation of China (http://www.nsfc.gov.cn/, grant no. 31771700) and the Program for Innovative Research Team (in Science and Technology) in University of Henan Province, china (http://www.haedu.gov.cn/, grant no. 15IRTSTHN015).

Competing interests

The authors declare that they have no competing interests.

References

1. Link AJ, LaBaer J. Trichloroacetic acid (TCA) precipitation of proteins. Cold Spring Harb Protoc.

2011; 2011: 993-4.

2. Wang W, Tai F, Chen S. Optimizing protein extraction from plant tissues for enhanced

proteomics analysis. J Sep Sci. 2008; 31:2032-9.

3. Niu LJ, Yuan HY, Gong FP, Wu XL, Wang W, Protein extraction methods shape much of the

extracted proteomes, front. Plant Sci. 2018 (online)

4. Wang W, Scali M, Vignani R, Spadafora A, Sensi E, Mazzuca S, et al. Protein extraction for two-

dimensional electrophoresis from olive leaf, a plant tissue containing high levels of interfering

compounds. Electrophoresis. 2003; 24:2369-75.

5. Wu X, Xiong E, Wang W, Scali M, Cresti M. Universal sample preparation method integrating

trichloroacetic acid/acetone precipitation with phenol extraction for crop proteomic analysis.

Nat Protoc. 2014; 9:362-74.

6. Wang W, Vignani R, Scali M, Cresti M. A universal and rapid protocol for protein extraction

from recalcitrant plant tissues for proteomic analysis. Electrophoresis. 2006; 27:2782-6.

7. Wang W, Wu X, Xiong E, Tai F. Improving gel-based proteome analysis of soluble protein

extracts by heat prefractionation. Proteomics. 2012; 12:938-43.

8. Xiong E, Wu X, Yang L, Gong F, Tai F, Wang W. Chloroform-assisted phenol extraction improving

proteome profiling of maize embryos through selective depletion of high-abundance storage

proteins. PLoS One. 2014; 9:e112724.

9. Méchin V, Damerval C, Zivy M. Total protein extraction with TCA-acetone. Methods Mol Biol.

2007; 355:1-8.

10. Isaacson T, Damasceno CM, Saravanan RS, He Y, Catalá C, Saladié M, et al. Sample extraction

techniques for enhanced proteomic analysis of plant tissues. Nat Protoc. 2006; 1:769-74.

11. Wu X, Gong F, Wang W. Protein extraction from plant tissues for 2DE and its application in

proteomic analysis. Proteomics. 2014; 14:645-58.

12. Damerval C, Vienne DD, Zivy M, Thiellement H. Technical improvements in two-dimensional

electrophoresis increase the level of genetic variation detected in wheat-seedling proteins.

Electrophoresis. 1986; 7: 52–4.

13. Granier F. Extraction of plant proteins for two-dimensional electrophoresis. Electrophoresis.

1988; 9:712-8.

14. Saravanan RS, Rose JK. A critical evaluation of sample extraction techniques for enhanced

proteomic analysis of recalcitrant plant tissues. Proteomics. 2004; 4:2522-32.

15. Carpentier SC, Witters E, Laukens K, Deckers P, Swennen R, Panis B. Preparation of protein

extracts from recalcitrant plant tissues: an evaluation of different methods for two-

dimensional gel electrophoresis analysis. Proteomics. 2005; 5:2497-507.

16. Maldonado AM, Echevarría-Zomeño S, Jean-Baptiste S, Hernández M, Jorrín-Novo JV.

Evaluation of three different protocols of protein extraction for *Arabidopsis thaliana* leaf

proteome analysis by two-dimensional electrophoresis. J Proteomics. 2008; 71:461-72.

17. Fic E, Kedracka-Krok S, Jankowska U, Pirog A, Dziedzicka-Wasylewska M. Comparison of

protein precipitation methods for various rat brain structures prior to proteomic analysis.

Electrophoresis. 2010; 31:3573-9.

18. Zadražnik T, Hollung K, Egge-Jacobsen W, Meglič V, Šušta..r-Vozlič J. Differential proteomic

analysis of drought stress response in leaves of common bean (*Phaseolus vulgaris* L.). J

Proteomics. 2013; 78:254-72.

19. Hao R, Adoligbe C, Jiang B, Zhao X, Gui L, Qu K, et al. An optimized trichloroacetic

acid/acetone precipitation method for two-dimensional gel electrophoresis analysis of

qinchuan cattle *Longissimus dorsi* muscle containing high proportion of marbling. PLoS One.

267 2015; 10:e0124723.

268 20. Shaw MM, Riederer BM. Sample preparation for two-dimensional gel electrophoresis.

269 Proteomics. 2003; 3:1408-17.

270 21. Simpson DM, Beynon RJ. Acetone precipitation of proteins and the modification of peptides.

271 J Proteome Res. 2010; 9:444-50.

272 22. Alias N, Aizat WM, Amin NDM, Muhammad N, Noor NM. A simple protein extraction method

273 for proteomic analysis of mahogany (*Swietenia macrophylla*) embryos. Plant Omics. 2017;

274 10:176-82.

275 23. Balbuena TS, Silveira V, Junqueira M, Dias LL, Santa-Catarina C, Shevchenko A, et al. Changes

276 in the 2-DE protein profile during zygotic embryogenesis in the Brazilian Pine (*Araucaria*

277 *angustifolia*). J Proteomics. 2009; 72:337-52.

278 24. Nandakumar MP, Shen J, Raman B, Marten MR. Solubilization of trichloroacetic acid (TCA)

279 precipitated microbial proteins via NaOH for two-dimensional electrophoresis. J Proteome

280 Res. 2003; 2:89-93.

281 25. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of

282 protein utilizing the principle of protein-dye binding. Anal Biochem. 1976; 72:248-54.

283 26. Laemmli UK. Cleavage of structural proteins during the assembly of the head of

284 bacteriophage T4. Nature. 1970; 227:680-5.

285 27. Wang, N., Cao, D., Gong, F. P., Ku, L. X., Chen, Y. H., and Wang, W. Differences in properties

286 and proteomes of the midribs contribute to the size of the leaf angle in two near-isogenic

287 maize lines. J. Proteomics. 2015; 128, 113-22.

288 28. Wu, X., Gong, F., Yang, L., Hu, X., Tai, F., and Wang, W. Proteomic analysis reveals differential

289 accumulation of small heat shock proteins and late embryogenesis abundant proteins

290 between ABA-deficient mutant vp5 seeds and wild-type Vp5 seeds in maize. Front. Plant Sci.

2014; 5, 801.

29. Wang W, Vignani R, Scali M, Sensi E, Tiberi P, Cresti M. Removal of lipid contaminants by

organic solvents from oilseed protein extract prior to electrophoresis. Anal Biochem. 2004;

329:139-41.

30. Ning F, Wu X, Zhang H, Wu Z, Niu L, Yang H, et al. Accumulation Profiles of Embryonic Salt-

Soluble Proteins in Maize Hybrids and Parental Lines Indicate Matroclinous Inheritance: A

Proteomic Analysis. Front Plant Sci. 2017; 8:1824.

31. Harder A, Wildgruber R, Nawrocki A, Fey SJ, Larsen PM, Görg A. Comparison of yeast cell

protein solubilization procedures for two-dimensional electrophoresis. Electrophoresis. 1999;

20(4-5):826-9.

32. Görg A, Obermaier C, Boguth G, Csordas A, Diaz JJ, Madjar JJ. Very alkaline immobilized pH

gradients for two-dimensional electrophoresis of ribosomal and nuclear proteins. Plant J.

2004;39(5):715-33.

33. Rose JK, Bashir S, Giovannoni JJ, Jahn MM, Saravanan RS. Tackling the plant proteome:

practical approaches, hurdles and experimental tools. Electrophoresis. 1997; 18:328-37.

307
308
309
310 **Figure 1.** The comparison of the steps between the modified and the classical TCA/acetone

311 precipitation methods. The latter was done according to Isaacson *et al.* [13]. The SDS extraction

312 buffer contains 1% (w/v) SDS, 0.1 M Tris-HCl (pH 6.8), 2 mM EDTA-Na$_2$, 20 mM DTT and 2 mM

313 PMSF (adding before use). All organic solvents are pre-chilled at -20℃ and contain 5 mM DTT

314 (adding before use).

Figure 2 Optimization of TCA/acetone ratio used in the modified method. Equal amounts (ca. 30 μg) of maize root proteins were analyzed by SDS-PAGE (12.5% resolving gel). Protein was stained using CBB. A, final acetone concentration was 50% (v/v), but TCA concentration varied from 0-20% (w/v) in the aqueous mixture. B, final TCA concentration was 10% (w/v), but acetone concentration varied from 0-80% (v/v) in the aqueous mixture.

Figure 3 Comparison of 2DE protein profiles of maize root proteins extracted using two methods. Figures were from two independent experiments. *Left panel*: the modified TCA/acetone precipitation. *Right panel*: the classical TCA/acetone precipitation. Spots with increased abundance were indicated in red. About 800 μg of proteins was resolved in pH 4-7 (linear) strip by IEF and then in 12.5% gel by SDS-PAGE. Proteins were visualized using CBB. Differential abundance spots (numbered, >2-fold) were subject to MS/MS analysis.

Figure 4 Comparison of 2DE profiles of maize embryo proteins extracted using two methods.

Shown were two independent experiments. Left panel: the modified TCA/acetone precipitation.

Right panel: the classical TCA/acetone precipitation. About 800 µg of proteins were resolved in

pH 4-7 (linear) strip by IEF and then in 12.5% gel by SDS-PAGE. Protein was visualized using CBB.

Differential abundance spots (numbered, >2-fold) were subject to MS/MS analysis.

339
340
341 **Figure 5** Comparison of 2DE profiles of maize leaf proteins extracted using two methods. Shown

342   were two independent experiments. Left panel: the modified TCA/acetone precipitation. Right

343   panel: the classical TCA/acetone precipitation. About 800 μg of proteins were resolved in pH 4-

344   7 (linear) strip by IEF and then in 12.5% gel by SDS-PAGE. Protein was visualized using CBB.

345   Differential abundance spots (numbered, >2-fold) were subject to MS/MS analysis.

Figure 6 Comparison of 2DE profiles of E. coli proteins extracted using two methods. Left panel:

the modified TCA/acetone precipitation. Right panel: the classical TCA/acetone precipitation. E.

coli BL21 was cultivated at 37°C for 12 h with good aeration until optical density of about 0.8.

Cells were collected by centrifugation and washed with phosphate-buffered saline (PBS).

Proteins were extracted by using two methods. About 800 μg of proteins were resolved in pH 4-

7 (linear) strip by IEF and then in 12.5% gel by SDS-PAGE. Protein was visualized using CBB.

355 **Table 1** Comparison of protein yield and spot number in 2DE between both methods

Maize tissues	The modified method		The classical method	
	Yield [a]	Spot No [b]	Yield [a]	Spot No [b]
Roots	4.82±0.07	738±11	4.70±0.08	743±10
Leaves	4.13±0.11	310±9	4.06±0.23	314±4
Embryos	5.80±0.13	314±12	5.56±0.38	304±18

361 Notes: a, Protein yields were expressed as μg/mg fresh weight. b, Average spot number detected in 2D gels

362 using PDQUEST software (version 8.0, Bio-Rad). All data were from at least three independent experiments.

363 The corresponding data from the two methods did not significantly differ at $p < 0.05$ according to t-test.

Table 2 The identification of the differential extracted proteins between the two methods in maize

Protein name	Accession	MW(kDa)/p*I*	Protein score	Protein Score C. I. (%)	Coverage (%)	GRAVY	Subcellular localization	Matched Peptides
Calreticulin-2, partial (spot 1)	ONM21219	41.11/4.81	240	100	47.6	-0.870	Endoplasmic reticulum	FEDGWESR;NPNYQGKWK;QTGSIYEHWDILPPK;WKAPMID NPDFK; FYAISAEYPEFSNK;YIGIELWQVK;KDDNMAGEWNHTSGK; SGTLFDNIIITDDPALAK;WNGDAEDKGIQTSEDYR;KPEGYD DIPKEIPDPDAK;KPEDWDDKEYIPDPEDK;KPEDWDDEEDG EWTAPTIPNPEYK;
Fructokinase-1 (spot 2)	AAP42805	34.67/4.87	86	99.937	39.0	0.139	Chloroplast	MLAAILR;EALWPSR;EFMFYR;LGDDEFGR;DFHGAVPSFK;T AHLRAMEIAK;FANACGAITTTK;DNGVDDGGVVFDSGAR;L GGGAAFVGKLGDDEFGR;AAVFHYGSISLIAEPCR;VSEVELE FLTGIDSVEDDVVMK
Glycine-rich RNA-binding protein 2(spot 3)	ACG28116	15.48/6.1	182	100	34.6	-0.487	Nucleus	NITVNEAQSR;GGGYGNSDGNWR;RDGGGGYGGGGGGYGG GGGYGGGGGGYGGGNR
Elongation factor 1-β (spot 4)	ONM59608	15.67/4.38	134	100	13.2	-0.360	Cell membrane	LDEYLLTR;KLDEYLLTR;LSGITAEGQGVK;WFNHIDALVR
Glutathione transferase (spot 5)	CAA73369	25.10/6.21	132	100	43.8	-0.174	Cytoplasm	GDGQAQAR;NLYPPEK;LYDCGTR;AEMVEILR;KLYDCGTR; VYDFVCGMK;FWADYVDKK;ECPRLAAWAK;GLAYEYEQD LGNK;QGLQLLDFWVSPFGQR;KQGLQLLDFWVSPGQR;QG LQLLDFWVSPFGQRCR;GLAYEYLEQDLGNKSELLLR
Lea14-A (spot 6)	AMY96568	16.08/5.64	514	100	96.7	0.024	Chloroplast. Golgi apparatus. Nucleus.	DGATLAGRVDVR;DWDIDYEMR;SGELKLPTLSSIF;DAGRD WDIDYEMR;LDVPVKVPYDFLVSLAK;TVASGTVPDPGSLAG DGATTR;VGLTVDLPVVGKLTLPLTK;NPYSHAIPVCEVTYTL R;LANIQKPEAELADVTVGHVGR;SAGRTVASGTVPDPGSLA GDGATTR;VDVRNPYSHAIPVCEVTYTLR;GFVADKLANIQK PEAELADVTVGHVGR
Sedoheptulose-1,7-bisphosphatase (spot 11)	ACG31345	41.79/6.8	101	99.998	19.0	-0.136	Chloroplast	EKYTLR;LLICMGEAMR;FEETLYGSSR;GIFTNVTSPTAK;YT GGMVPDVNQIIVK;LTGVTGGDQVAAAMGIYGPR

Note: For MS/MS analysis, differential abundance spots (> 2 folds) were extracted, in-gel digested (trypsin, 37°C, 20 h), and analyzed by the MALDI-TOF/TOF analyzer (AB SCIEX
TOF/TOF-5800, USA). MALDI-TOF/TOF spectra were acquired in the positive ion mode and automatically submitted to Mascot 2.2 (http://www.matrixscience.com) for
identification against NCBInr database (version Feb 17, 2017; species, Zea mays, 279566 sequences). The search parameters were as follows: type of search: combined (MS +
MS/MS); enzyme: trypsin; dynamical modifications: oxidation (M); fixed modifications: carbamidomethyl (C); mass values: monoisotopic; protein mass: unrestricted; peptide
mass tolerance: ±100 ppm; fragment mass tolerance: ±0.4 Da; peptide charge state: 1+; max missed cleavages: 1. Unambiguous identification was judged by the number of
matched peptide sequences, sequence coverage, Mascot score and the quality of MS/MS spectra. All of the positive protein identification scores were significant (p<0.05).