KLF4 binding during reprogramming is involved in 3D architectural

rewiring and transcriptional regulation of enhancer hubs

Dafne Campigli Di Giammartino[1]*, Yiyuan Liu[1]*, Andreas Kloetgen[2]*, Alexander

Polyzos[1], Daleum Kim[1], Matthias Stadtfeld[3], Aristotelis Tsirigos[2,4,5]# and Effie

Apostolou[1]#

[1]Sanford I. Weill Department of Medicine, Sandra and Edward Meyer Cancer
Center, Weill Cornell Medicine, New York, NY 10021 USA

[2] Department of Pathology, NYU School of Medicine, New York, NY, 10016, USA

[3]Skirball Institute of Biomolecular Medicine, Department of Cell Biology and
Helen L. and Martin S. Kimmel Center for Biology and Medicine, NYU School of
Medicine, New York, NY 10016, USA

[4]Laura and Isaac Perlmutter Cancer Center and Helen L. and Martin S. Kimmel
Center for Stem Cell Biology, NYU School of Medicine, New York, NY, 10016,
USA

[5]Applied Bioinformatics Laboratories, NYU School of Medicine, New York, NY,
10016, USA

* These authors contributed equally

#Corresponding Authors

Lead Contact: efa2001@med.cornell.edu

SUMMARY

Cell fate transitions are accompanied by global transcriptional, epigenetic and topological changes driven by transcription factors (TFs), as is strikingly exemplified in somatic cell reprogramming to pluripotent stem cells (PSCs) by OCT4, KLF4, SOX2 and cMYC. How TFs orchestrate the complex molecular changes around their targets in a temporal manner remains largely elusive. Here, using KLF4 as a paradigm, we provide the first TF-centric view of chromatin reorganization during reprogramming and its association to enhancer rewiring and gene regulation. We captured the enhancer connectomes in fibroblasts and PSCs by H3K27ac HiChIP and identified complex 3D enhancer hubs that were strongly correlated with cell-type specific gene expression and coregulation. KLF4-centric conformational analysis at different reprogramming stages revealed that KLF4 is involved in the dissociation of fibroblast-specific and the establishment of PSC-specific enhancer loops concomitantly with repression or activation of linked genes. Moreover, KLF4 occupancy was significantly enriched within highly connected enhancers, suggesting a role in the formation and regulation of complex enhancer hubs. Indeed, disruption of a single KLF4 binding site from a PSC-specific enhancer was sufficient to reduce expression of multiple genes within the enhancer hub, partly by impairing long-range contact. Our study provides an integrative view of the intricate activities of a master regulator during a controlled cell fate transition and offers novel insights into the order and nature of molecular events that follow TF binding.

INTRODUCTION

The identity of each cell type is determined by a unique gene expression

program as well as characteristic epigenetic landscape and three-dimensional

(3D) chromatin topology. All of these features are under the control and constant

supervision of a network of critical transcription factors (TF), known as master

regulators of cell identity[1,2]. Although, the ability of master regulators to maintain

or change cell identity is well appreciated, the underlying mechanisms remain

poorly understood.

Somatic cell reprogramming into induced pluripotent stem cells (iPSCs) by

the so-called Yamanaka factors OCT4, KLF4, SOX2 and cMYC (OKSM) offers a

tractable system to study the molecular mechanisms of cell fate determination

and the roles and activities of each reprogramming TF[3,4]. Research over the last

decade started dissecting on a genome-wide level the transcriptional and

epigenetic changes that result in successful erasure of somatic identity and

establishment of pluripotency[5]. Distinct or synergistic roles of the reprogramming

TFs as well as specific direct and indirect mechanisms for coordinating these

molecular changes have been proposed[6-10]. In addition to the transcriptional and

epigenetic changes, recent studies utilizing targeted or global chromatin

conformation capture techniques revealed that the 3D chromatin topology differs

between somatic and pluripotent stem cells (PSCs) and is largely reset during

reprogramming[11-16]. However, the principles of chromatin reorganization during

iPSC generation, the association with enhancer and gene activity and the

involvement of TFs in these processes have only started to be explored.

The potential role of reprogramming TFs in 3D chromatin organization is

currently based mostly on computational analyses of 4C or Hi-C datasets, which

reveal a strong enrichment of OKS binding around long-range interactions in

PSCs or during reprogramming[11-13,15]. The architectural function of KLF4 is

further supported by experimental evidence, as it has been shown that KLF4

depletion abrogates specific loops around target loci in the context of PSCs [14],

and also in human epidermal keratinocytes[17], while depletion of the related factor

KLF1 disrupts selected long-range interactions in the context of erythropoiesis

[18,19]. These findings establish a link between reprogramming TF binding and

chromatin architecture and suggest that OKS, and particularly KLF4, may

actively orchestrate long-range chromatin interactions during reprogramming in

order to establish and maintain the pluripotent transcription program. To test this

possibility in a direct, genome-wide and dynamic manner during the course of

reprogramming, we used KLF4 as a paradigm to capture the TF-centric

topological reorganization and determine the relationships with epigenetic and

transcriptional changes. To do so, we used a well-characterized reprogramming

system[20,21] to apply genome-wide assays that map KLF4 binding (ChIP-seq),

chromatin accessibility (ATAC-seq), enhancer and gene activity (H3K27ac ChIP-

seq and RNA-seq), enhancer connectivity (H3K27ac HiChIP) as well as KLF4-

centric chromatin looping (KLF4 HiChIP) at different stages during acquisition of

pluripotency (Fig.1a top). Integrative analysis of our results generated a

reference map of stage-specific chromatin changes around KLF4 bound loci and

established strong links with enhancer rewiring and concordant transcriptional

changes. Genetic manipulation of KLF4 binding from a PSC enhancer further

supported the ability of KLF4 to function both as a transcriptional regulator and a

chromatin organizer. In conclusion, our study offers novel insights into the

intricate roles of a master regulator during cell fate transition.

Results

KLF4 binding induces chromatin opening and precedes enhancer and gene

activation

To determine the molecular changes around KLF4 targets during iPSC formation,

we first mapped its genome-wide binding at different stages of reprogramming

using "reprogrammable" mouse embryonic fibroblasts (MEFs) (Rosa26-

M2rtTA/Col1a1-OKSM) induced with doxycycline (dox)[20,21] in the presence of

ascorbic acid. Under these conditions the resulting iPSCs are molecularly and

functionally indistinguishable from embryonic stem cells[22,23] and, thus, the term

PSCs will be used throughout the text to describe either cell type. For our earliest

time point, we collected bulk populations on day 3 after dox treatment, whereas

at later stages, on day 6 and day 9, we sorted SSEA1 positive cells to enrich for

cells that are on the right trajectory towards induced pluripotency[21,24] (Fig.1a

bottom and Supplementary Fig.1a). Finally, we used PSCs as a reference point

for established pluripotency. ChIP-seq analysis showed a highly dynamic pattern

of KLF4 occupancy during reprogramming with two major categories of binding

sites: (i) transiently detected in reprogramming intermediate populations, but

absent in PSCs (Transient KLF4 targets) and (ii) PSC targets, which represent

the actual KLF4 binding repertoire once stem cell identity is acquired (Fig.1b and

Supplementary Table 1). Among the PSC KLF4 binding sites, 30% were already

present on day 3 (Early KLF4 targets), while the rest were either gradually

established during reprogramming (Mid KLF4 targets) or detected only in

established PSCs (Late KLF4 targets). To gain insights into the nature and

potential function of each category of KLF4 targets we performed genomic

annotation based on their chromatin state classification introduced by Chronis et

al [6] as well as Gene Ontology (GO) analysis using the GREAT tool [25] (Fig.1c).

Early KLF4-targets mostly enriched for promoters of genes involved in regulation

of metabolic processes and cell junction organization, in agreement with the

previously reported early role of KLF4 in regulating these processes[21]. On the

other hand, Mid and Late KLF4 targets included an increasing number of

pluripotency-associated enhancers and enriched for stem cell maintenance

genes, including many master regulators of pluripotency, such as *Sox2*, *Nanog*,

Esrrb and *Klf4*. Finally, Transient KLF4 targets enriched for transient and partial-

iPSC enhancers of genes involved in negative regulation of cell cycle, apoptosis

and various signaling pathways associated with differentiation, such as TGF-beta

signaling (Fig.1c). The different nature of stage-specific KLF4 binding sites was

also supported by motif analysis, which revealed distinct classes of TFs enriched

among Early (e.g. cMYC and YY1), Mid (e.g.ELF5 and RFX), Late (e.g. ESRRB

and NR5A2) and Transient (e.g.FOSL1 and JUNB) KLF4 genomic targets

(Supplementary Fig.1b).

The differential ability of KLF4 to bind Early or Late targets prompted us to

investigate the chromatin accessibility in MEFs and its dynamic changes during

reprogramming (Fig.1d and Supplemental Fig.1c). Integration of ATAC-seq and

KLF4 ChIP-seq datasets revealed that ~60% of the Early KLF4 binding sites

were already open in MEFs, suggesting that preexisting accessibility could partly

explain the differential binding behavior of KLF4 during reprogramming (Fig.1d

and Supplementary Fig.1d) . In contrast, the majority (>70%) of mid, late and

transient KLF4 targets were "closed" in MEFs and gained accessibility at later

reprogramming stages (Fig.1d and Supplementary Fig.1d) in concordance with

the kinetics of KLF4 binding. This result is in agreement with the proposed

pioneer activity of KLF4[7,26] and/or with synergistic binding with the other

reprogramming factors[6].

KLF4 has been proposed to function both as an activator and repressor of

gene expression[9,21]. To assess the impact of KLF4 binding on enhancer and

gene activity, we first performed ChIP-seq for H3K27 acetylation in MEFs, PSCs

and intermediate reprogramming stages to map potential changes in enhancer

usage during iPSC generation (Supplementary Fig.1e). Our analysis

demostrated an acquisition of H3K27 acetylation concurrently with or

subsequently to KLF4 binding (Fig.1e and Supplementary Fig.1f), suggesting that

KLF4 might predominantly function as an activator. In further support of this

notion, less than 5% of decommisioned MEF enhancers were targeted by KLF4,

whereas about 35% of the total acquired PSC enhancers and almost the entirety

of (so-called) superenhancers (SE)[27] were bound by KLF4 concomitantly or prior

to H3K27 acetylation (Fig.1f, Fig.1g and Supplementary Table 2). Moreover,

RNA-seq analysis (Supplementary Fig.1g and Supplementary Table 3) of genes

that are linked to Early, Mid, Late or Transient KLF4 ChIP-seq peaks showed a

strong trend for upregulation, rather than downregulation, at the respective

stages of reprogramming (Supplementary Fig.1h). Representative examples of

ChIP-seq and ATAC-seq tracks as well as RNA-seq results for each category of

KLF4 targets are shown in Figure 1h.

In conclusion, our data revealed a stage-specific KLF4 binding during

reprogramming, which progressively targets PSC-associated enhancers and

eliminates transient, PSC-unrelated occupancy. KLF4 binding appeared to

depend only minimally on preexisting chromatin accessibility, had strong effects

on chromatin opening and accordingly on enhancer and gene activation.

Enhancers are extensively rewired during reprogramming and form

complex interactomes in established pluripotent stem cells.

Previous studies utilizing targeted (4C-seq) or global (Hi-C) chromatin

conformation assays have demonstrated that chromatin topology around specific

genomic loci or in large scale, such as compartments and Topologically

Associated Domains (TADs), is largely cell-type specific and drastically

reorganized during reprogramming[11,13-16,28]. However, the small-scale, cell type-

specific regulatory loops, such as enhancer-promoter interactions, were

insufficiently captured in these studies due to technical limitations. Here, we

performed H3K27ac HiChIP[29] in MEFs and PSCs in order to generate high-

resolution contact maps around active enhancers and promoters and

characterize the degree of architectural reorganization during reprogramming.

After calling high-confidence interactions using HiC-bench tool[30] (see methods),

we further refined our set by considering only loops that overlapped with

H3K27ac ChIP-seq peaks in at least one anchor (Supplementary Fig.2a).

Differential looping analyses (pvalue<0.01. LogFC>2) revealed about 50,000

contacts that were enriched either in MEFs or in PSCs (Fig.2a, Supplementary

Fig.2b and Supplementary Table 4). By applying stringent statistic criteria

(pvalue>0.5, logFC<0.5) , we also identified a group of ~6,000 constant H3K27ac

contacts that show similar interaction strength in both MEFs and PSCs.

Enrichment analysis using the LOLA tool[31], revealed that MEF-specific loops

were enriched for binding of somatic TFs, such as Runx2 and Junb, whereas

PSC-specific loops were strongly associated with pluripotency TFs, including

OKS and NANOG as well as transcriptional co-activators, such as CBP and

Mediator subunits (Supplementary Fig.2c).

In agreement with previous publications[32], H3K27ac HiChIP detected a

large number of Enhancer-Enhancer (E-E), Enhancer-Promoter (E-P) and

Promoter-Promoter (P-P) contacts (Supplementary Fig.2d). Interestingly, we

observed that genomic regions that were detected as anchors in PSC-specific

H3K27ac loops were involved in significantly higher number of long-range

contacts (higher connectivity) than the ones detected in MEF-specific and

constant loops (Fig. 2b). Together, these results document the dramatic 3D

genomic reorganization of enhancers between MEFs and PSCs and reveal that

cell type-specific enhancer connectomes are characterized by different degrees

of complexity.

3D-organized enhancer cliques and enhancer hubs are associated with

cell-type specific gene expression and coregulation

The presence of complex enhancer-promoter interactions, such as "enhancer

cliques" (multiple enhancers contacting the same promoter) or "enhancer hubs"

(multiple promoters communicating with the same enhancer) (Supplementary

Table 5) prompted us to examine their potential cell-type specific function in

regulating gene expression. Integration of RNA-seq data overall showed a

significant correlation of MEF-specific or PSC-specific H3K27ac loops with

increased expression of associated genes in the respective cell type (Fig. 2c and

Supplementary Fig.2e). Interestingly, the strength of this correlation and the

levels of cell type-specific expression were higher when we focused on genes

within enhancer cliques (at least two enhancer contacts) and increased with the

number of enhancer contacts per gene (Fig.2c). These results support that

multiple enhancers connected to the same gene in a cell-type specific manner

are more likely to ensure robust and tight transcriptional activation.

We then focused on enhancers that establish connections with multiple

gene promoters hypothesizing that they may coregulate these genes in the

context of 3D regulatory hubs (enhancer hubs). To test this hypothesis, we

selected all protein-coding genes that participate in such hubs (2 or more genes

contacting the same enhancer) and are expressed (>1TPM) in either MEFs or

PSCs and performed pair-wise comparisons among genes within hubs to

calculate the percentage of co-regulation or anti-regulation. For control groups

we used random gene pairs (from the same pool of genes as the test group)

either of similar linear distance with our test group (global random) or within the

same TADs[33](TAD-matched random). The results show a significant

overrepresentation of coregulated gene pairs within enhancer hubs compared to

our control groups (Fig.2d). This analysis revealed 217 gene pairs that reside

within PSC-enhancer hubs and become concordantly upregulated during

reprogramming (Supplementary Table 5). As an example, we found that the

previously described Zfp42 SE also contacts and potentially co-regulates the

distal *Triml2* gene (Fig.2e). *Triml2* , similarly to *Zfp42,* is activated specifically in

PSCs, suggesting a potential function in the maintenance of stem cell identity. A

more complex hub was revealed around a SE enhancer located 60kb

downstream of *Nanog,* which appeared to form PSC-specific contacts with

multiple pluripotency-associated genes, including the proximal (~25kb) Slc2a3

and the distal (~450kb) Phc1 (Supplementary Fig.2f). Therefore, our analyses

revealed that many PSC SE and typical enhancers (TE) that were previously

assigned to only one gene[27,34], formed hubs with multiple genes and not

necessarily the most proximal ones (enhancer skipping) (Supplementary Fig.2g)

in a PSC-specific manner. Importantly, Gene ontology analysis of the newly

discovered SE and TE target genes enriched for stem cell maintenance and cell

cycle categories, supporting their functional relevance (Supplementary Fig.2g).

To better understand the biological basis of low or high enhancer

connectivity, we ranked PSC H3K27ac anchors based on the number of contacts

they establish and looked for features that may contribute to their distinct looping

behaviour. Using available ChIP-seq datasets, we compared binding intensities

at regions with the lowest (only one contact, n=3,594) and highest connectivity

(equal or more than four contacts, n=1434) and found that binding of pluripotency

TFs, including KLF4, was significantly stronger ($p<10^{-16}$) in highly connected

enhancers, similarly to Mediator subunits Med1 and Med12 (Fig.2f). However,

well-described architectural proteins, such as CTCF and cohesin[35] or the recently

described enhancer-promoter mediator YY1[36,37], did not show significantly

increased binding within highly connected enhancers (Fig. 2f). The involvement

of KLF4 in PSC-specific H3K27ac interactions was further corroborated by the

fact that about 80% of the H3K27ac PSC-specific loops were occupied by KLF4

with almost 50% being Early or Mid KLF4 targets during reprogramming (Fig.2g

and Supplementary Fig.2h). These results establish a link between KLF4 binding

and enhancer-promoter looping in established PSCs and during reprogramming,

supporting a potential architectural function for KLF4 in addition to its

transcriptional and pioneering activity.

Dynamic KLF4-centered chromatin reorganization during reprogramming

associates with enhancer rewiring and transcriptional changes of target

genes.

To capture the KLF4-centric chromatin reorganization during iPSC formation, we

performed HiChIP in early (day 3) and mid (day 6) stages of reprogramming as

well as in PSC using KLF4 antibody (Supplementary Fig.3a). Principle

component analysis (PCA) on all high-confidence interactions distinguished

primarily KLF4 from H3K27ac HiChIP data (Supplementary Fig.3b) regardless of

cell type or stage, demonstrating the different nature of chromatin contacts that

each antibody captures. PCA2 and PCA3 showed a clear separation of the

different reprogramming stages, suggesting dynamic reorganization of KLF4-

bound contacts (Fig.3a). Differential looping analysis generated 4 clusters of

differential KLF4-centered interactions (Fig.3b and Supplementary Table 6): two

clusters of gained loops, detected either in mid or late reprogramming stages

and two clusters of lost loops detected only in early or mid stages. Integration of

KLF4 ChIP-seq data revealed a strong correlation of stage-specifc KLF4-looping

with the expected stage-specific KLF4 binding (Fig.3c). Specifically, the majority

of gained loops overlapped with mid and late KLF4 binding, while lost loops had

predominantly transient KLF4 occupancy. Interestingly, in about 25-30% of KLF4

loops, KLF4 binding was detected prior to loop formation or persisted after loop

loss (Fig.3c), suggesting that KLF4 binding may be causal but not always

sufficient for looping formation or detection.

298   To gain insights into the role and nature of the different KLF4-centered

loop clusters we investigated the expression changes of associated genes during

reprogramming. Fig.3d shows that the relative percentages of up- and down-

regulated genes are different among the distinct loop clusters. Lost KLF4

contacts mostly associate with gene repression, while gained KLF4 loops with

gene activation during reprogramming (Fig.3d). Furthermore, integration of

H3K27ac HiChIP data from MEFs and PSCs, showed that most of the gained

KLF4 loops were also detected in PSC-enriched H3K27ac HiChIP contacts, while

lost KLF4 loops mostly enriched for MEF enhancer contacts (Fig.3e). Examples

of such loops and genes are shown in Figure 3f, which illustrates the normalized

KLF4 and H3K27 HiChIP signal around *Mycn* and *Ets1* in the format of virtual 4C

along with the respective ChIP-seq tracks during reprogramming (Fig.3g). *Mycn*

is a pluripotency-associated gene activated during reprogramming (Fig.3h),

concomitantly with the establishment of KLF4-mediated loops with a distal

enhancer. In contrast, *Ets1* is downregulated during reprogramming (Fig.3h) in

concordance with the gradual loss of a pre-existing H3K27ac-loop in MEFs that

was abolished in PSCs (Fig. 3f). Importantly, comparison of published Hi-C

datasets with our H3K27ac HiChIP results validated the cell-type specific nature

of these loops (Supplementary Fig.3c). Moreover, independent 3C-qPCR

experiments validated that the *Mycn*-enhancer loop was gradually established

during reprogramming, while the *Ets1*-enhancer contact frequency was

significantly decreased by day 6 (Fig.3i). Together, these observations suggest

that stage-specific KLF4 binding is involved both in the formation/activation of

PSC enhancer loops and abrogation/repression of prexisting somatic enhancer

loops.

KLF4 binding is involved in different types of chromatin loops

The previous results establish strong links between KLF4-centered chromatin

reorganization during reprogramming with enhancer looping as well as gene

expression changes, but they do not explain all observed long-range contacts. As

shown in Fig.3d, a high percentage (~50%) of genes within dynamic KLF4 loops

did not significantly change their expression during reprogramming. In

accordance, a large proportion of both gained and lost KLF4-centered

interactions appeared to be either constitutive enhancer loops (similar H3K27ac

HiChIP signal in both MEF and PSC) or independent of H3K27ac looping

(Fig.3e). These data demonstrate that looping around KLF4-bound loci might not

always result in changes in enhancer and/or gene activity.

The fact that KLF4 participates in H3K27ac-dependent and independent

chromatin contacts prompted us to investigate the potentially distinct looping

mechanisms and architectural partners of KLF4 for each loop category. To do so,

we first clustered all PSC KLF4 binding sites based on their presence within

H3K27ac-dependent or independent loops (KLF4 loops that were also detected

or not in H3K27ac HiChIP) (Supplementary Table 7). To increase stringency,

KLF4 binding sites present in both anchors of a loop (bi-anchored sites) were

considered. As a control group, KLF4 target sites that were detected neither in

KLF4 nor in H3K27ac HiChIP contacts were used (non-anchored sites).

Importantly, KLF4 binding strength was similar in all categories (Supplementary

Fig.3d), suggesting that the differential looping probability around KLF4 target

sites may depend on the presence or absence of other factors. Enrichment

analysis using published ChIP-seq datasets showed that KLF4 binding sites

within H3K27ac-dependent loops enriched for active enhancer features such as

binding of pluripotency TFs (ESRRB, NANOG, SOX2 and POU5F1), YY1 as well

as POLR2A and Mediator subunits (Supplementary Fig.3e and Supplementary

Table 8). In addition, H3K27ac-dependent KLF4 loops were associated with

occupancy/activity of specific histone modulators such as BRD4, p300 and

KDM2B or KDM6B, suggesting a potential architectural function for such

complexes. On the other hand, H3K27ac-independent KLF4 anchors

preferentially enriched for binding of classic architectural factors, such as

Cohesin and CTCF (Supplementary Fig.3e), but also of PRC1 and PRC2

components, which have been reported to mediate looping among repressed or

bivalent genes in PSCs[13,38]. Given that many of the enriched factors directly

interact with KLF4 protein, these data suggest that KLF4 might establish

chromatin loops with distinct properties and functions by recruiting different

architectural cofactors.

Disruption of KLF4 binding site from a pluripotency-associated enhancer interferes with looping and decreases expression of multiple genes within the hub.

To examine the functional significance of KLF4 in long-range chromatin interactions and associated gene expression, we decided to focus on the *Tbx3* locus, which was detected to contact multiple enhancers specifically in PSCs by both KLF4 and H3K27ac HiChIP and validated using published Hi-C data (Fig.4a and Supplementary Fig.4a). 3C-qPCR experiments during reprogramming confirmed the establishment of a contact between *Tbx3* promoter with a distal (-90kb) enhancer at late stages of reprogramming and concomitantly with gene activation (Fig.4b and 4c). To experimentally test the relevance of this putative enhancer for *Tbx3* gene regulation, we utilized CRISPR/Cas9 technology to genetically delete it in PSCs using a previous characterized proximal enhancer as a reference[39] (Fig.4d, Supplementary Fig.4b and Supplementary Table 9). RT-pPCR analysis of homozygous knock-out (KO) clones (Supplementary Fig.4c) demostrated that the transcriptional levels of *Tbx3* were severely impaired upon disruption of either enhancer (Dis-KO and Pro-KO), with the distal enhancer showing a stronger effect (Fig.4e). Interestingly, the RNA levels of the surrounding non-coding genes, *Gm1603* and *Aw549542,* which were part of the same enhancer hub with *Tbx3* (Fig.4a and Supplementary Fig.4d), were also reduced (Fig 4f). Importantly, mutation of just one KLF4 binding motif within the distal enhancer (DisKLF4[mut]) (Fig 4d and Supplementary Fig.4b and e), also resulted in a significant downregulation of *Tbx3*, *Gm1603* and *Aw549542* RNA levels (Fig. 4e), supporting a critical role of KLF4 binding in the regulatory activity

of this enhancer. In addition, and consistently with the transcriptional

downregulation, the long-range contact between *Tbx3* and the distal enhancer

was specifically and significantly reduced upon mutation of the KLF4 motif as

shown by 3C-qPCR (Fig.4f), while the interaction of *Tbx3* with the proximal

enhancer or a KLF4-independent contact in chromosome 7 remained unaffected

(Fig.4f). These results provide direct evidence for the role of KLF4 both as

transcriptional regulator and chromatin organizer within enhancer hubs.

Discussion

Here, we describe the dynamic binding of KLF4 and its effects on chromatin

accessibility, enhancer activity and connectivity, gene expression and 3D

chromatin organization during iPSC reprogramming. Our data demostrated that

KLF4 initial binding mostly occured in previously inaccessible chromatin regions

and coincided with chromatin opening. This binding behavior is likely the result of

its documented pioneering function[7,26] as well as cooperativity with other

reprogramming TFs[6]. KLF4 binding was predominantly linked to gene

upregulation rather than downregulation, suggesting that a previously reported

repressive activity of this TF[6,9,21] may be manifested at earlier timepoints[6] or by

indirect mechanisms, for example through activation of the suppressor SAP30,

as it has been recently shown in a different reprogramming system[40]. Finally,

KLF4 binding often occurred prior to enhancer and gene activation (early KLF4

binding-late H3K27 acetylation), suggesting a causality, but also the requirement

of additional factors, such as ESRRB or NANOG[6], which are reactivated only

later in reprogramming (not shown).

Previous studies utilizing 4C, 5C or Hi-C have reported that 3D genomic

architecture is drastically reorganized between somatic and pluripotent cells[11,13-

16]. However, the principles and mechanisms of such reorganization remain

largely elusive. A recent study performed Hi-C analysis at different stages of B

cell reprogramming and offered important insights into the kinetics and degree of

chromatin reorganization on the level of chromosome compartments and TADs[28].

However, only a minor fraction of cell-type specific loops were detected. Here,

we chose to apply H3K27ac HiChIP analysis, which has been recently shown to

have significantly higher discovery rate for cell-type specific loops, compared to

Hi-C and Capture Hi-C methods[29,32]. Indeed, our data revealed a dramatically

rewired enhancer connectome between MEFs and PSCs generating a reference

map of conformational changes, relevant for cell identity. Our analyses unraveled

cell-type specific H3K27ac interactomes that included simple and complex P-P,

E-E and E-P contacts. Interestingly, genes that communicated with multiple

enhancers (enhancer cliques[32]) in a cell-type specific manner appeared to be

more strongly upregulated in the respective cell-type, which supports the

possibility that genes critical for cell identity are under the control of more than

one enhancers ensuring increased specificity and robustness. In parallel, our

analysis identified a number of cell-type specific 3D enhancer hubs, including

many SE-hubs, which contained two or more coregulated genes. This provides

genome-wide evidence that complex 3D enhancer hubs may play a functional

role in coordinating gene regulation during cell fate transitions and may therefore

enable identification of novel candidate genes and enhancers critical for the

maintenace or establishment of specific cellular identities.

Encouraged by previous studies reporting potential architectural functions

for various KLF protein members[14,18,19], we went on to capture for the first time

the KLF4-centric chromatin reorganization around KLF4-binding sites during

iPSC formation. Integrative computational analysis established strong links

between KLF4-bound stage-specific contacts with cell-type specific enhancer

looping and concordant gene expression changes. Specifically, our analyses

showed that KLF4 binding associated with *de novo* formation of PSC-specific

enhancer loops during reprogramming promoting activation of linked genes. On

the other hand, transient KLF4 binding on preexisting enhancer loops in somatic

cells was linked to their gradual abrogation and silencing. This phenomenon of

TF "hijacking" on preestablished loops has been previously reported during

differentiation and reprogramming[28,41], but mostly as a trigger for looping

stablization and gene activation and less for loop destabilization and

transcriptional repression.

Finally, our results revealed that KLF4 binding occurs within distinct

categories of chromatin contacts: (i) loops enriched in active enhancer features,

such as H3K27ac, Mediator and pluripotency TFs, (ii) structural H3K27ac-

independent loops, which were established by strong architecural factors such as

Cohesin and CTCF and (iii) potentially repressive or paused loops that we

associated with PRC1/2 components. The role of KLF4 binding in the formation,

stabilization or functionality of these distinct loop categories remains to be shown.

In support of a critical role of KLF4 in maintaining enhancer-promoter

communication and activity, we showed that disruption of a single KLF4 binding

site from a newly identified PSC-specific enhancer, decreased expression of

multiple genes within the enhancer hub, partly by reducing looping frequency.

In conclusion, our study provided the first TF-centric chromatin

reorganization analysis during a cell fate transition and established mechanistic

and functional links with enhancer rewiring and gene regulation. The integrative

analysis of KLF4 occupancy and looping data with changes in chromatin

accessibility and enhancer and gene activity offered novel insights on the

multiple activities of a master regulator during somatic cell reprogramming. In

addition, our approach identified complex PSC-specific 3D enhancer hubs and

provided evidence for their role in gene coregulation during reprogramming.

Importantly, KLF4 binding was strongly enriched in high connectivity hubs,

suggesting a potential regulatory and architectural fuction for this TF, which was

further supported by genetic engineering experiments. Given the reported role of

KLF4 in multiple cell fate decisions, such as keratinocyte and monocyte

differentiation, and its context-dependent function as a tumor-suppressor or

oncogene[42], it will be interesting to identify and interrogate the significance of

KLF4-mediated 3D hubs in different cellular contexts. Future work will be critical

to (i) delineate whether chromatin looping around TF-binding sites occurs by

active tethering through recruited architectural cofactors or by "self-organization"

through liquid-phase separation as a result of local epigenetic alterations and (ii)

determine the functional significance of TF-mediated 3D chromatin hubs in cell

fate transitions.

FIGURES AND FIGURE LEGENDS

Figure 1

Figure 1. Dynamic KLF4 binding during reprogramming and association

with chromatin accessibility and enhancer activity. (a) Schematic illustration

of the experimental system and strategy. (**b**) Tornado plots of KLF4 ChIP-seq

signals at different reprogramming stages clustered in four different categories:

Early, Mid, Late and Transient KLF4 binding. ChIP-seq signals were centered

1kb upstream/downstream of peak centers. **(c)** Left: Pie charts of functional

classification of KLF4 Early, Mid, Late and Transient peaks (based on[6]) (piPSC,

partial iPSCs); Right: GREAT gene ontology analysis of Early, Mid, Late and

Transient KLF4 target sites. (**d-e**) Line plots showing the percentages of KLF4

Early, Mid, Late and Transient targets that overlapped with (d) ATAC-seq peaks

or (e) H3K27ac ChIP-seq peaks at each reprogramming stage (**f**) Tornado plot of

H3K27ac ChIP-seq signal at each reprogramming stage 2.5Kb upstream or

downstream of peak centers. **g)** Bar plots showing overlap of KLF4 binding with

either lost (MEF and d3) or gained (d9 and PSC) H3K27ac peaks (top) or with

typical PSC enhancers (TE) versus superenhancers (SE) as characterized by

[27](bottom). **h)** Examples of specific KLF4 targets with respective IGV tracks for

KLF4 ChIP-seq, ATAC-seq, H3K27ac ChIP-seq at each reprogramming stage.

Genomic coordinates are shown in parenthesis and the signal values on the right

side. The transcriptional changes of the depicted genes during reprogramming

are shown at the bottom, expressed as transcripts per million (TPM).

Figure 2

Figure 2. Detection and characterization of H3K27ac-associated loops in

MEFs and PSCs by H3K27ac HiChIP analysis. (a) Heatmap of differential

loops detected by H3K27ac HiChIP in MEF versus PSC. Differential loops were

called by average logFC > 2 and p-value < 0.01, constant loops were called by

average logFC > -0.5 & < 0.5 and p-value > 0.5. Heatmap shows row-wise z-

score of normalized interaction strength per replicate. **(b)** Connectivity of

H3K27ac differential or constant loops measured as number of high-confidence

contacts around each H3K27Ac HiChIP fragment. Asterisks indicate significant

difference with p<0.001 **(c)** RNA expression changes between MEFs and PSCs

of genes that were involved in at least two (left) or at least five (right) MEF-

specific, PSC-specific or constant H3K27ac loops. All protein-coding genes were

used as control. Asterisks indicate significant difference (p<0.001) as calculated

by paired t-test. **(d)** Barplot indicating the percentages of gene pairs within

enhancer hubs (enhancers contacting 2 or more gene promoters) that become

transcriptionally co-regulated (both up or both down) or anti-regulated (one up

one down) between MEFs and PSCs. Global Random or TAD-matched gene

pairs were used as controls (see also Methods). Genes that remain unchanged

(FC<1.5, p>0.5) were not considered in this analysis. Significance is indicated by

asterisks and was calculated by Fisher's exact test **(e)** Example of a newly

identified enhancer hub in PSCs. Normalized HiChIP signals around the

viewpoint are illustrated as virtual 4C plots. H3K27ac ChIP-seq and RNA-seq

IGV tracks for each reprogramming stage are shown below to highlight

concordance with H3K27ac HiChIP data and coordinated upregulation of target

genes. **(f)** Violin plots showing ChIP-seq intensity under peaks located within

PSC-specific enhancers with either low connectivity (only one contact, n=3594)

or high connectivity (more than 4 loops, n=1434). In addition to KLF4 (in-house

ChIP-seq data), we examined other pluripotency TFs as well as known

architectural factors using public ChIP-seq datasets in PSCs. Signal comparison

between low and high connectivity was done by Wilcoxon signed-rank test and

the p-values are reported. **(g)** Pie chart showing overlap of H3K27ac PSC-

specific loops with KLF4 Early, Mid, Late and Transient ChIP-seq peaks**.**

Figure 3

Figure 3. Capturing the KLF4-mediated chromatin reorganization during

reprogramming and its association to enhancer and gene activity a) PCA

analysis of high-confidence interactions of H3K27ac and KLF4 HiChIP samples

(see also Supplementary figures 3a and 3b). **(b)** Heatmap of differential KLF4

HiChIP analysis depicting 4 distinct clusters grouped into gained or lost loops.

Heatmap shows row-wise z-score of the normalized interaction strength for KLF4

HiChIP replicates. **(c)** Pie charts representing overlap between gained or lost

KLF4 HiChIP loops with Early, Mid, Late and Transient KLF4 ChIP-seq peaks.

(d) Stacked barplot indicating the percentages of genes within gained or lost

KLF4 loops that become upregulated or downregulated (logFC > 1.5, FDR < 0.01

in PSC vs day3) during reprogramming. **(e)** Stacked barplot showing the

percentage of gained or lost KLF4 loops that were also detected by H3K27ac

HiChIP analysis and were enriched in either MEFs or in PSC or in both (constant

loops). Note that about 25-30% of the loops were not detected by H3K27ac

(H3K27ac-independent KLF4 loops) (see also Supplementary figure 3d,e). **(f)**

Examples of KLF4-mediated loops that are either acquired (*Mycn*) or lost (*Ets1*)

during reprogramming. Normalized KLF4 and H3K27ac HiChIP signals are

illustrated in a virtual 4C format around the viewpoints (*Mycn* and *Ets1* gene

promoters). The respective ChIP-seq IGV tracks are shown in **(g)**, while the RNA

changes during reprogramming expressed as transcripts per million (TPM) are

shown in **(h)**. **(i)** 3C-qPCR validation of the relative interaction frequency between

Mycn and Ets1 promoters and the respective distal enhancers as depicted in (f)

at different reprogramming stages. Unpaired one-tailed Student's t-test was used

to determine significance relative to the MEF time point (value indicated on the

top of each bar). Error bars indicate standard deviation in *n=2* biological

replicates.

Figure 4

Figure 4. Disruption of KLF4 binding site from a distal *Tbx3* enhancer

abrogates looping and decreases *Tbx3* expression in PSCs. (a) Normalized

KLF4 and H3K27ac HiChIP signals are illustrated as virtual 4C line plots around

the *Tbx3* promoter. The respective ChIP-seq IGV tracks are shown at the bottom.

(b) *Tbx3* RNA changes during reprogramming expressed as transcripts per

million (TPM) **(c)** 3C-qPCR validation of the relative interaction frequency

between the *Tbx3* promoter and the most distal enhancer as depicted in (a) at

different reprogramming stages. Unpaired one-tailed Student's t-test was used to

determine significance relative to MEF (value indicated on the top of each bar).

Error bars indicate standard deviation in *n=2* biological replicates. **(d)** Schematic

diagram showing CRISPR-mediated manipulations of the *Tbx3* distal (Dis) and

proximal (Prox) enhancers. KO: knockout, KLF4mut: Mutation or indel of a KLF4

binding motif within the distal enhancer. **(e)** RT-qPCR showing expression

changes of *Tbx3* and nearby genes in either wild-type (w.t) PSCs or in CRISPR-

Cas9 engineered PSC carrying homozygous deletions of the distal *Tbx3*

enhancer (Dis-KO), the proximal (Prox-KO) or the KLF4mut. Unpaired one-tailed

Student's t-test was used to determine significance relative to w.t. (value

indicated on the top of each bar). Error bars indicate standard deviation from n=2

biological replicates **(f)** 3C-qPCR analysis to detect changes in looping formation

between the *Tbx3* promoter and either the distal or proximal enhancer following

mutation of the KLF4 binding site within the distal enhancer. Unpaired one-tailed

Student's t-test was used to determine significance relative to w.t. (value

indicated on the top of each bar). Error bars indicate standard deviation from n=2

biological replicates.

METHODS

Cell lines, culture conditions and reprogramming experiments
Mouse ES V6.5 were cultured on irradiated feeder cells in KO-DMEM media
(Invitrogen) supplemented with 15% heat-inactivated fetal bovine serum,
GlutaMAX, penicillin-streptomycin, non-essential amino acids, β-mercaptoethanol

and 1000 U/ml LIF, with or without the presence of 2i (1uM MEKinhibitor
(Stemgent 04-0006) and 3uM GSK3 inhibitor (Stemgent 04-0004)).
Mouse embryonic fibroblasts (MEFs) were isolated from a "reprogrammable"
mouse harboring a polycystronic OKSM cassette in the Col1a1 locus and M2rtTA
in the Rosa26 locus[20]. Cells were reprogrammed in the presence of 1ug/ml
doxycycline and 50ug/ml ascorbic acid and cultured in ES medium as described
above. Cells were collected at the indicated time points.

MACS and FACS
For isolating the SSEA1 positive cells from reprogramming intermediates at day6
and day9 we used magnetic microbeads conjugated to anti-SSEA1 antibody
(MACS Miltenyi Biotec #130-094-530) as per manufacture instructions. SSEA
positive and negative fractions were then stained for FACS analysis with an anti-
Thy1 antibody conjugated to pacific blue fluorophore (ebioscience # 48-0902-82)
and anti-SSEA antibody conjugated to APC fluorophore (biolegend #125608).
611
Generation, selection and validation of KO
gRNAs were cloned into the px458 vector using the BbsI restriction enzyme[43].
0.3 million ESC cells (V6.5) were transfected using 2ug of Left-*Tbx3*-plasmid and
2ug of Right-Tbx3-plasmid (for *Tbx3* enhancer deletions) or 4ug of *Tbx3*-
KLF4mut-vector (mutation of KLF4 binding site within *Tbx3* distal enhancer).
DNA was pre-mixed with 50ul media with no additions, and in a separate tube
10ul of Lipofectamine 2000 (Invitrogen #11668019) was pre-mixed with 50ul
media with no addition. After 5 minutes the two tubes were combined and
incubated at room temperature for 20 more minutes. Cells were then added to
the solution and plated on a gelatinized 12 well. 48hrs post-transfection, GFP
positive single cells were sorted by FACS into 96 well plates. Genotyping of the
single cell colonies was performed using a three-primer strategy (for deletions) or
by surveyor with T7 (for in-del mutation). Two colonies with homozygous
mutations (or w.t. colonies as control) were expanded and used for RT-qPCR
and 3C experiments. All gRNA and primers are described in Supplementary
Table 9.
628
3C-qPCR
For each sample 1 to 2 million cells were lysed in 300ul of lysis buffer (10mM
Tris-HCl pH 8.0, 10mM NaCl, 0.2% Igepal CA630 with protease inhibitors) and
incubated on ice for 20 mins. Cells were centrifuged 2500g for 5min at 4°C and
pellet washed once in lysis buffer. Pellets were resuspended in 50ul of 0.5% SDS
and incubated at 65°C for 10 mins. 145ul of water and 25ul of 10% triton were
added to the samples and incubated 15mins at 37°C. 100 Units of MboI
restriction enzyme and 25ul of NEB buffer 2 were added and incubated over
night at 37°C with rotation. Next day the enzyme was inactivated at 65°C for 20
mins. The ligation reaction was carried out over night at 16°C by adding 120ul of
NEB T4 ligase buffer, 100ul of 10% Triton, 6ul of 20mg/l BSA, 100ul of 10mM
ATP and 5ul of T4 ligase (NEB #M0202).The day after, 50ul of 20mg/ml
proteinase K and 120ul of 10% SDS were added and the samples were
incubated over night at 65°C. Lastly, 10ul of 10mg/ml RNAse was added and

samples incubated 1 hour at 37°C. Following phenol chloroform purification the
DNA was precipitated using 1.6 volumes of 100% ethanol and 0.1 volume of 3M
sodium acetate. After incubation at -80C for 1 hour samples were spun for
15mins at 4°C at 16000rpm. Pellets were washed twice with 70% ethanol and
dissolved in 100ul of 10mM Tris pH8. Qbit was used to measure sample
concentrations and 100ng of material was used to amplify the desired regions by
qPCR. All primer sequences can be found in Supplementary table 9.
650
ChIP-seq
ChIP-seq was performend as previously described[44]. Specifically, cells were
crosslinked in 1% formaldehyde at RT for 10 minutes and quenched with 125mM
glycine for 5 mins at RT. 50 million cells were used for KLF4 ChIPs and 10
million for H3K27acetylation ChIP. Cell pellets were washed twice in PBS and
resuspended in 400ul lysis buffer (10mM Tris pH8, 1mM EDTA, 0.5% SDS) per
20 million cells. Cells were sonicated in a bioruptor device (30 cycles 30sec
on/off, high setting) and spun down 10 minutes at 4°C at maximum speed.
Supernatants were diluted 5 times with dilution buffer (0.01%SDS, 1.1%
triton,1.2mM EDTA,16.7mM Tris pH8, 167mM NaCl) and incubated with the
respective antibody (2-3ug/10M cells) (KLF4 R&D #3158, H3K27ac ab4729)
O/N with rotation at 4°C. Next day, protein G Dynabeads (ThermoScientific)
preblocked with BSA protein (100ng per 10ul Dynabeads) were added (10ul
blocked Dynabeads per 10 million cells) and incubated for 2-3 hours at 4°C.
Beads were immobilized on a magnet and washed twice in low salt buffer (0.1%
SDS,1% triton, 2mM EDTA, 150mM NaCl, 20mM Tris pH8), twice in high salt
buffer (0.1% SDS,1% triton, 2mM EDTA, 500mM NaCl, 20mM Tris pH8), twice in
LiCl buffer (0.25M LiCl, 1% NP40, 1% deoxycholic acid (sodium salt), 1mM
EDTA, 10mM Tris pH8) and once in TE. DNA was then eluted from the beads by
incubating with 150ul elution buffer (1% SDS, 100mM NaHCO3) for 20 minutes
at 65°C (vortexing every 10min). Supernatants were collected and reverse-
crosslinked by incubation at 65°C O/N in presence of proteinase K. After RNase
A treatment for 1hr at 37°C, DNA was purified using the minElute kit (Qiagen). 6-
10ng of immunoprecipitated material was used for ChIP-seq library preparation
using the KAPA Hyper prep kit (KAPA Biosystems). Libraries were sequenced on
an Illumina HiSeq 2500 platform on SE50 mode.
677
ATAC-seq
ATAC-seq was performed as previously described[45]. In brief, a total of 50,000
cells were washed once with 50 µL of cold PBS and resuspended in 50 µL lysis
buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.2% (v/v) IGEPAL
CA-630). The suspension of nuclei was then centrifuged for 10 min at 800 g at
4°C, followed by the addition of 50 µL transposition reaction mix (25 µL TD buffer,
2.5 µL Tn5 transposase and 22.5 µL nuclease-free H_2O) using reagents from the
Nextera DNA library Preparation Kit (Illumina #FC-121-103). Samples were then
incubated at 37°C for 30min. DNA was isolated using a ZYMO Kit (#D4014).
ATAC-seq libraries were first subjected to 5 cycles of pre-amplification. To
determine the suitable number of cycles required for the second round of PCR
the library was assessed by quantitative PCR as described in Buenrostro et al

and the library was then PCR amplified for the appropriate number of cycles
using Nextera primers. Samples were subject to a dual size selection (0.55x-
1.5x) using SPRI beads (Beckman Coulter #B23317). Finally, the ATAC libraries
were sequenced on a HiSeq 2500 platform on PE50 mode.
694
RNA-seq
Total RNA was prepared with TRIZOL (Life technologies #15596018) following
manufacturer's instructions. Libraries were generated by the Weill Cornell
Genomics core facility using the Illumina TruSeq stranded mRNA library
preparation kit (#20020594) and sequenced on an Illumina HiSeq4000 platform
on SE50 mode.
701
HiChIP
HiChIPs were performed as previously described[29] with some modifications. In
brief, up to 15 million crosslinked cells (for KLF4 HiChIPs two samples of 15
million cells were combined at the end, for each sample replicate) were
resuspended in 500 µL of ice-cold Hi-C lysis buffer (10 mM Tris-HCl pH 7.5, 10
mM NaCl, 0.2% NP-40, 1× protease inhibitors) and rotated at 4°C for 30 min.
Nuclei were pelleted and washed once with 500 µL of ice-cold Hi-C lysis buffer.
Pellet was then resuspended in 100 µL of 0.5% SDS and incubated at 62°C for
10 min. 285 µL of water and 50 µL of 10% Triton X-100 were added, and
samples were rotated at 37°C for 15 min. 50 µL of NEB Buffer 2 and 15 µL of
25 U/µL MboI restriction enzyme (NEB, R0147) were then added, and sample
was rotated at 37°C for 2 h. MboI was then heat inactivated at 62°C for 20 min.
We added 52 µL of incorporation master mix: 37.5 µL of 0.4 mM biotin–dATP
(Thermo Fisher, 19524016); 4.5 µL of a dCTP, dGTP, and dTTP mix at 10 mM
each; and 10 µL of 5 U/µL DNA Polymerase I, Large (Klenow) Fragment (NEB,
M0210). The reactions were then rotated at 37°C for 1 h. 948 µL of ligation
master mix was then added: 150 µL of 10× NEB T4 DNA ligase buffer with 10
mM ATP (NEB, B0202), 125 µL of 10% Triton X-100, 3 µL of 50 mg/mL BSA
(Thermo Fisher, AM2616), 10 µL of 400 U/µL T4 DNA Ligase (NEB, M0202),
and 660 µL of water. The reactions were then rotated at room temperature for
4 h. After proximity ligation, the nuclei were pelleted and the supernatant was
removed. The nuclear pellet was brought up to 880 µL in Nuclear Lysis Buffer
(50 mM Tris-HCl pH 7.5, 10 mM EDTA, 0.5% SDS, 1× Roche protease
inhibitors, 11697498001), and sonicated with a Bioruptor 300 (Diagenode) for 8
cycles of 30sec each, on a medium setting. Clarified samples were transferred
to Eppendorf tubes and diluted five times with ChIP Dilution Buffer (0.01% SDS,
1.1% Triton X-100, 1.2 mM EDTA, 16.7 mM Tris-HCl pH 7.5, 167 mM NaCl).
Cells were precleared with 30 µL of Protein G dynabeads (Life technology
#10004D) in rotation at 4°C for 1 h. Supernatants were transferred into fresh
tubes and antibody was added (8 µg of KLF4 antibody or 3ug H3K27Ac
antibody for 15 million cells) and incubated overnight at 4°C. The next day 30
µL of Protein G dynabeads were added to samples and rotated at 4°C for 2 h.
After bead capture, beads were washed three times each with low-salt wash
buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 7.5, 150
mM NaCl), high-salt wash buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20

mM Tris-HCl pH 7.5, 500 mM NaCl), and LiCl wash buffer (10 mM Tris-HCl pH
7.5, 250 mM LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM EDTA, make
fresh). Samples were eluted with 150 µL of DNA elution buffer (50 mM sodium
bicarbonate pH 8.0, 1% SDS, freshly made) and incubated at 37°C for 30 min
with rotation. Supernatant was transferred to a fresh tube and elution repeated
with another 150 µL elution buffer. 5 µL of Proteinase K (20mg/ml) (Thermo
Fisher) were added to the 300 µL reaction and samples were incubated
overnight in 65°C. Samples were purified with DNA Clean and Concentrator
columns (Zymo Research) and eluted in 10 µL of water. Post-ChIP DNA was
quantified by Qubit (Thermo Fisher) to estimate the amount of Tn5 (Illumina)
needed to generate libraries at the correct size distribution (see below). 5 µL of
Streptavidin C-1 beads (Thermo Fisher) were washed with Tween Wash Buffer
(5 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 M NaCl, 0.05% Tween-20) then
resuspended in 10 µL of 2× biotin binding buffer (10 mM Tris-HCl pH 7.5, 1
mM EDTA, 2 M NaCl). Beads were added to the samples and incubated at
room temperature for 15 min with shaking. After capture, beads were washed
twice by adding 500 µL of Tween Wash Buffer and incubated at 55°C for 2 min
with shaking. Samples were then washed in 100 µL of 1× TD Buffer (2× TD
Buffer is 20 mM Tris-HCl pH 7.5, 10 mM magnesium chloride, 20%
dimethylformamide). After washes, beads were resuspended in 25 µL of 2× TD
Buffer, Tn5 (for 50 ng of post-ChIP DNA we used 2.5 µL of Tn5), and water to
50 µL. Tn5 amount was adjusted linearly for different amounts of post-ChIP
DNA, with a maximum amount of 4 µL of Tn5. Samples were incubated at
55°C with interval shaking for 10 min. After removing the supernatant 50 mM
EDTA was added to samples and incubated with interval shaking at 50°C for
30 min. Beads were then washed two times each in 50 mM EDTA then Tween
Wash Buffer at 55°C for 2 min. Lastly, beads were washed in 10 mM Tris
before PCR amplification. Beads were resuspended in 25 µL of Phusion HF 2×
(New England Biosciences), 1 µL of each Nextera Ad1_noMX and Nextera
Ad2.X at 12.5 µM, and 23 µL of water. The following PCR program was
performed: 72 °C for 5 min, 98 °C for 1 min, then cycle at 98 °C for 15 s, 63 °C
for 30 s, and 72 °C for 1 min (cycle number was estimated based on the
amount of material from the post-ChIP Qubit (approximately 50 ng was run in
six cycles, while 25 ng was run in seven, 12.5 ng was run in eight, etc.). Size
selection was performed using two-sided size selection with the Ampure XP
beads. After PCR, libraries were placed on a magnet and eluted into new tubes.
25 µL of Ampure XP beads were added, and the supernatant was kept to
capture fragments less than 700 bp. Supernatant was transferred to a new
tube, and 15 µL of fresh beads was added to capture fragments greater than
300 bp. After size selection, libraries were quantified with Qbit and sent for
Bioanalyzer to check for the quality and final size of the library. Libraries were
sequenced on a HiSeq 2500 platform on PE75 mode.
779
ATAC-seq data analysis
Mapping, peak calling and peak processing. Paired-end reads were aligned to
mm10 (bowtie2 version 2.3.2, --no-unal --local --very-sensitive-local --no-
discordant --no-mixed --contain --overlap --dovetail -I 10 -X 2000), and

mitochondrial DNA alignments were excluded. Fragments marked as positional
duplicates (sambamba version 0.6.6) or overlapping with mouseENCODE
blacklisted genomic regions (liftOver to mm10;[46]) were filtered out. Read ends
were adjusted for Tn5 transposase offsets. Peaks were called at $p<10^{-5}$ (MACS
version 2.1.1) per replicate, and only common peaks between two independent
biological replicates were retained for further analysis.
790
ChIP-seq data analysis
Mapping, peak calling and peak processing. Study and published ChIP-seq
reads were trimmed for adapters with parameters of maximum error rate of 0.4
and minimum length of 15 bp (cutadapt version 1.8.1), and low-quality ends
(sickle version 1.33), respectively. Alignment to the mouse reference genome
version mm10 (GRCm38.p4) was performed using standard parameters,
permitting a maximum of one mismatch in seed alignment (bowtie2 version 2.3.2).
Reads marked as positional duplicates (sambamba version 0.6.6) or overlapping
with mouseENCODE blacklisted genomic regions (liftOver to mm10;[46]) were
filtered out. Study ChIP-seq peaks (enrichment of signals over background
determined by input samples) were called at p<0.01 (MACS version 2.1.1) per
biological replicate, and peaks detected in more than half of biological replicates
were retained for further analysis. Published ChIP-seq replicates were merged,
and peaks were called at $p<10^{-5}$ using input samples where applicable.
805
806   **Overlap analysis of ChIP-seq peaks for chromatin states of reprogramming**
807   **cell types.** Chromatin states (1 kb resolution) during reprogramming were
808   retrieved from ref [6], and cis-regulatory elements were annotated from chromatin
809   states as in the original publication. The assignment of ChIP-seq peaks to cis-
810   regulatory elements was determined by the largest degree of overlap in bp.
811
ChIP-seq gene ontology (GO) enrichment analysis. Spatial proximity of ChIP-
seq peaks to transcript start sites (TSSs) and enriched GOs were uncovered
utilizing the GREAT (version 3.0.0) web application. We selected the "basal plus
extension rule" for the association of gene ontology annotations with regulatory
domains (customized setting: 5 kb upstream and 1 kb downstream of TSSs, and
further extended both directions by 250 kb). Enrichment of ontology annotations
was assessed by the binomial test of ChIP-seq peak-overlaps with annotated
regulatory regions.
820
821   **PSC typical enhancers and super-enhancers.** Coordinates of typical- and
822   super-enhancers in mESCs were ascertained from ref[34], lifted over from mm9 to
823   mm10, and gene symbols featured in the GENCODE reference (version M6)
824   were retained.
825
Overlaps of KLF4 binding with early lost or late gained H3K27ac peaks
during reprogramming or at typical- and super-enhancers. The maximum
permitted distance between KLF4 binding detected in day3, 6 and 9 with PSC
and H3K27ac peaks or enhancers in ref[34] was 250 bp. Where H3K27ac peaks or

enhancers overlapped with KLF4 sites of different stages, the earliest stage was
prioritized (Fig. 1g).
832

Motif analysis
For each KLF4 cluster we generated a random background (by shuffling the
peaks randomly throughout the genome) to test motif enrichment within each
cluster. Analysis of the KLF4 clusters was performed with the use of HOMER and
'findMotifsGenome.pl' command with the following parameters: '-bg random.bed -
size 200 -len 15'. Only motifs with p-value≤1e-5 were considered significant.Two
heatmaps with the z-transformed '-log10(p-value)' and z-transformed 'motif
frequency' of selected motifs for each cluster are presented in Supplementary
Fig.1d.
842

PCA analysis for ATAC-seq, RNA-seq and ChIP-seq experiments:
We first merged all accessible regions / H3K27ac peak detected from ATAC-seq
/ H3K27ac ChIP-seq in any reprogramming stage using bedtools v2.25.0. Then,
we calculated the coverage of reads for each merged accessible region and
H3K27ac peak for each replicate independently. For the RNA-seq data, we
calculated the coverage for each exon and only exons with at least 1 read
covering every single base of the exon were used for downstream analysis. PCA
analysis was performed with R and PCA plots were generated with 'ggplot2'
library. In each PCA plot, we present the variability captured by the first two PCs
(PC1 and PC2).
853

854 **RNA-seq data analysis**
855 Expression of genes was quantified in transcripts per kilobase million (TPM)
856 using quasi mapping (Salmon, version 0.8.2) to GENCODE (version M6, mm10)
857 reference gene annotation. Salmon provides alignment-free transcript
858 quantification information in a single step[47].
859

Line plots for gene expression analysis
We plotted the median expression levels of all protein coding genes with their
corresponding 95% confidence interval (CI) that are bound by KLF4 in a distance
less than 50 kb from their corresponding transcription start site (TSS). For each
KLF4 cluster we calculated the closest (≤50 kb) TSS to each KLF4 binding site
and plotted the median expression levels (TPM) of all genes annotated in each
KLF4 cluster with the use of R.
867

Preprocessing of HiChIP / public Hi-C datasets
HiChIP datasets and public Hi-C datasets were uniformly pre-processed with the
HiC-bench platform[30], which is outlined in short in the following. First, all paired-
end sequencing reads were aligned against the mouse genome version mm10
with bowtie2 version 2.2.3[48] (specific settings: --very-sensitive-local --local).
Read-filtering was conducted by the GenomicTools[49] gtools-hic filter command
(integrated in HiC-bench), which discards multi-mapped reads ("multihit"), read-
pairs with only one mappable read ("single sided"), duplicated read-pairs
("ds.duplicate"), read-pairs with a low mapping quality of MAPQ < 20, read-pairs

resulting from self-ligated fragments and short-range interactions resulting from
read-pairs aligning within 10kb (together called "ds.filtered"). The percentage of
accepted intra-chromosomal read-pairs ("ds.accecpted intra") was high across all
HiChIP replicates and conditions and was consistently above 35% (Figure S3a).
In order to create counts-matrices per chromosome in a binned fashion, we set
the bin size to 10kb for all datasets. For all the HiChIP sample and chromosome
matrices, the trajectories of each matrix bin to both anchors were overlaid with
the ChIP-Seq signal of the respective matching sample, requiring a minimal
overlap of 1bp between a HiChIP-bin and a ChIP-peak. Only loops of which at
least one anchor was supported by a ChIP-peak were kept for further analyses.
Next, we applied sequencing-depth normalization (called "scaling") and distance
normalization as recently described[50].
High-confidence loops were identified by requiring at least 1 read-pair per loop
across all replicates of a single condition in order to maintain a signal that is
replicable. Next, all loops were ordered by average loop strength per cell-type
and antibody and the top 50k / 100k interactions were used for downstream
analyses for KLF4 and H3K27ac, respectively.
894
Principal component analysis for HiChIP samples
Principal component analysis (PCA) as shown in Figures 3a and S3b was
performed on all available replicates on the high-confidence loops. Therefore, for
each detected high-confidence loop from any sample, the per replicate
normalized loop strength was extracted before filtering for top 50k / 100k loops in
order to also integrate lowly detected interactions in the analysis. PCA was
performed using the prcomp function of R (version 3.3.0; scale=TRUE,
center=TRUE)
903
904　**HiChIP/Hi-C heatmaps**
905　All HiChIP and Hi-C heatmaps were generated with the image.plot function of R
906　the package fields (version 8.10), using the normalized interaction matrices.
907　Color scale was set the same across all conditions but separately for Hi-C and
908　HiChIP samples.
909
Differential loop analysis
Differential looping analysis was performed on each loop independently by
applying a two-sided t-test on the normalized counts between any pairwise
comparisons: ES-KLF4 vs d3-KLF4, ES-KLF4 vs d6-KLF4, d3-KLF4 vs d6-KLF4
and ES-H3K27ac vs MEF-H3K27ac. In order to estimate the change in loop
strength, we calculated the log fold-change (logFC) between the average
normalized loop-strength per condition for the same pairwise comparisons after
adding a pseudo-count of 1 to each replicate and loop. For constant H3K27ac
loops, we selected loops with p-value > 0.5 and absolute logFC < 0.5 for the
pairwise comparison of H3K27ac. MEF/ES-specific H3K27ac loops were
selected by p-value < 0.01 and logFC > 2 / logFC < -2 taken from the ES
H3K27ac vs MEF H3K27ac comparison, respectively. Mid and late established
KLF4 loops were selected by applying p-value < 0.01 and logFC > 2 in the
pairwise comparisons of ES-KLF4 vs d3-KLF4 and d6-KLF4 vs d3-KLF4 (mid)

and ES-KLF4 vs d3-KLF4 and ES-KLF4 vs d6-KLF4 (late). Transient and mid-
lost KLF4 loops were selected by applying a p-value < 0.01 and logFC < -2 in the
pairwise comparisons of PSC-KLF4 vs d6-KLF4 and PSC-KLF4 vs d3-KLF4
(transient) and PSC-KLF4 vs d3-KLF4 and d6-KLF4 vs d3-KLF4 (mid-lost). All
calculations were performed in R version 3.3.0, using the native t.test function.
929
Annotation of H3K27ac HiChIP loop anchors as promoter or enhancers.
H3K27ac HiChIP loop anchors were overlapped with transcription start sites
(TSSs) of GENCODE (version M6) genes. Presence of one or more TSSs was
considered a promoter HiChIP anchor, and the absence of any TSS but
presence of at least one H3K27ac ChIP-seq constitutes an enhancer HiChIP
anchor. In estimating connectivity, all HiChIP anchors, either promoter, enhancer
or otherwise desolate, were considered.
937
RNA expression integration (enhancer cliques)
For RNA expression integration, we overlapped all canonical TSSs of protein-
coding genes (transcript support level/TSL = 1) downloaded from Ensembl
Genes V85 for the mouse genome mm10 with all loop anchors. Because the
TSS is a 1bp position in the genome, each gene was uniquely assigned to one
bin, however, multiple TSSs per gene with a TSL=1 mapping to different bins are
possible. Next, we filtered genes by occurrence of differential loop clusters that
were obtained from the HiChIP experiments and have TPM > 1 in at least one
reprogramming stage, and analyzed the expression patterns of such genes
throughout reprogramming. For H3K27ac HiChIP data integration, we assigned
genes to MEF/PSC-specific loops if their TSSs were found in >= 1, >= 2 or >= 5
MEF/PSC-specific loops but in none of the other (Figures 2c and S2e). Genes
contained in anchors of constant loops were solely filtered by having at least two
constant loop anchors. To further validate expression changes based on
differential looping, we applied a two-sided t-test between genes logFCs of
constant H3K27ac loops versus genes with MEF/PSC-specific loops. As a
negative control, we compared logFCs of genes with constant loops versus all
annotated protein-coding genes.
956
Co-regulation of gene expression by H3K27ac HiChIP (enhancer hubs). In
this analysis, promoter anchors of enhancer-mediated loops were filtered for
protein-coding genes that have an expression TPM > 1 in MEF or PSC.
Enhancer fragments that contact two to ten promoter fragments in PSC specific
H3K27ac loops were selected. Genes were paired across different promoter
fragments connecting to the same enhancer anchor (later on called hub), and
repeated gene pairs were removed from the overall pool. Owing to lower
promoter connectivity of enhancer hubs in MEF-specific or constant H3K27ac
HiChIP loops, the constraint of promoter contacts was reduced to five or three for
enhancer hubs in MEF-specific or constant loops. Gene pairs were considered
co-expressed, if both genes were up-regulated in PSC compared to MEF ($p<10^{-2}$
and fold change threshold of 1.5 in R limma package version 3.32.10) or vice
versa for down-regulation. Or otherwise, at least one gene in a pair unchanged
between MEF and PSC constitute unchanged gene pairs. Enrichment of co-

regulated gene pairs in the genuine data compared to either gene pairs in
random shuffled distance matched window or from within TADs was statistically
assessed using Fisher's exact test (Hi-C data from Pelham-Webb et al,
unpublished results). Topologically associated domains (TADs) were called from
normalized Hi-C matrices in PSCs processed at 40kb resolution using a recently
published software[51]. The insulation window parameter was set to 500 kb.
977

Overlap between H3K27ac PSC-specific loops and KLF4 clusters
979 Overlap between any of the KLF4 peaks with any of the HiChIP anchors
980 (H3K27aC or KLF4 loops) was performed with the use of bedtools v2.25.0.
981

ChIP-seq feature enrichment at lowly or highly connected H3K27ac HiChIP
enhancer anchors. H3K27ac HiChIP anchors were selected for low (contacting
only one other anchor; N = 3,594) or high connectivity (contacting four or more
anchors; N = 1,434), and the low connectivity anchors were sub-sampled to
match numbers of the high connectivity class. Average signals of ChIP-seq
features (reads per kilobase per million/RPKM) per base under peaks or the
fraction of peaks that are within anchors were statistically (Wilcoxon signed-rank)
tested for disparity between low and high connectivity anchors (Fig.2g). Publicly
available ChIP-seq data was used for OCT4, SOX2, MED1, SMC1, CTCF and
YY1:
992

	Study Accession	Sample Accession
OCT4	GSE44288	SRR713340
SOX2	GSE44288	SRR713341
MED1	GSE22557	SRR058987|SRR058988
SMC1	GSE22557	SRR058981|SRR058982|SRR058995|SRR058996
CTCF	https://www.enco deproject.org	https://www.encodeproject.org/files/ENCFF001LIY/ @@download/ENCFF001LIY.fastq.gz
YY1	GSE99518	SRR5634080

KLF4 peaks involved in KLF4 and H3K27ac looping.
Anchors of high-confidence PSC KLF4 loops and of PSC-specific H3K27ac loops
were stratified by KLF4 binding. . KLF4 peaks were overlapped with anchors of
loops of KLF4 and H3K27ac, separately. In these separate analyses, the
viewpoint was placed at this anchor to see if any of the partner anchors has
KLF4 binding. If so, these were called "bi-anchor" for the viewpoint anchor.
Thereby KLF4 peaks in ChIP-seq carried the identity of the viewpoint anchor. A
small number of KLF4 peaks may have conflict identities, due to partial
overlapping with multiple anchors, and thereby removed. A KLF4 peak is called
'no-anchor' if it does not overlap with any anchors in neither KLF4 nor H3K27ac
HiChIP loops.
1005

KLF4 looping involved in RNA expression changes
1007 To estimate the effect of KLF4 associated looping on changes in RNA expression,
1008 we followed a similar approach as for 'enhancer cliques'. After selecting

1009	expressed genes within anchors of each KLF4 loop cluster, we further filtered for
1010	differentially expressed genes between PSC and day3 (FDR < 0.01; logFC > 1.5
1011	(upregulated) or logFC < -1.5 (downregulated)). Information on differential
1012	expression was derived by edgeR (version 3.14.0) with subsequent multiple
1013	testing correction. Genes determined as 'no change' were selected by applying
1014	FDR > 0.5 and absolute logFC < 0.5. All remaining genes were discarded from
1015	the analysis.
1016	
1017	**LOLA enrichment analysis**
1018	The identified differential loops were subjected for an enrichment analysis of
1019	further transcription factor bindings and other DNA binding proteins. First, the
1020	anchors of each differential loop were mapped back to the original ChIP-peaks,
1021	because the 10kb stretches of the bins would give too many false positive
1022	findings. Each anchor that was overlapping an actual ChIP-peak by at least 1bp
1023	was subjected for further analysis. Since two anchors can theoretically overlap
1024	with a single ChIP-peak using this approach, the resulting list was collapsed and
1025	only unique ChIP-peaks were kept. Next, we applied LOLA version 1.4.0 [31]
1026	against a database of analyzed ChIP-Seq datasets taken from LOLA Region
1027	Databases (regionDB) for mm10 (for Figure S3e we used Codex and encode
1028	TFBSmm10 databases). We excluded all ChIP-Seq datasets that were marked
1029	as treated with any agent and had less than 3,000 peaks in total (5,000 peaks for
1030	Figure S3e). As a universe for LOLA, we used only unique ChIP-peaks of the
1031	union of all ChIP-Seq peaks for the respective antibody across all reprogramming
1032	stages.
1033	
1034	**Virtual 4C**
1035	Virtual 4C was performed to identify interaction signals of gene promoters or
1036	enhancers with their genomic vicinity. For this approach, we used the filtered
1037	HiChIP read-pairs as described above before binning and normalization of each
1038	replicate. We extracted all read pairs for which a read mate maps within +/- 10kb
1039	around the virtual viewpoint. Next, we defined successive overlapping windows
1040	for each chromosome at a 10kb resolution, and all adjacent windows are
1041	overlapping by 95% of their length (i.e. 9.5kb, or a shift of 500bp between
1042	adjacent windows). We then counted the second mapped read mate in all
1043	overlapping bins. Thus, each read-pair accounts for +1 in exactly 19 bins,
1044	however, the overlap of bins achieves a smoothed signal. Read counts for all
1045	bins were normalized to total sequencing depth of the respective replicate by
1046	edgeR version 3.14.0 to calculate counts-per-million (CPM) per bin. For
1047	visualization, the average of the normalized virtual 4C-signal across replicates of
1048	a single condition was calculated. We have used the following viewpoints:
1049	
1050	*Zfp42/Triml2* locus chr8:43,321,380-43,321,381
1051	*Mycn* locus chr12:12,941,173-12,941,174
1052	*Ets1* locus chr9:32,696,322-32,696,323
1053	*Tbx3* locus chr5:119,670,608-119,670,609 (promoter) / chr5:119,585,186-
1054	119,585,187 (Dist enhancer)
1055	

ACKNOWLEDGEMENTS

We thank all members from the Apostolou and Tsirigos groups for critical reading of the manuscript. We would also like to acknowledge Olivier Elemento and Ashley Doane for input with initial computational analyses and WCM Epigenomics and Genomics facilities for support on next-generation sequencing. D.C.D.G is a New York Stem Cell Foundation postdoctoral fellow and also supported by the Family-Friendly Postdoctoral Initiative at Weill Cornell Medicine. This work was partly funded by the NIH Director's New Innovator Award (1DP2DA043813), the Edward Jr Mallinckrodt Foundation and the Tri-Institutional Stem Cell Initiative by the Starr Foundation.

AUTHOR CONTRIBUTIONS

E.A conceived, designed and supervised the study with input from A.T and M.S. A.T. supervised the computational analyses. D.C.D.G designed and performed most of the experiments with help from D.K and E.A. Y.L analyzed all ChIP-seq, RNA-seq and ATAC-seq datasets. A.K analyzed all HiChIP and Hi-C data. Y.L, A.K and A.P.P performed integrative analyses. D.K contributed to the CRISPR/Cas9 experiments. E.A and D.C.G wrote the manuscript with input from all authors.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

References

Young, R. A. Control of the embryonic stem cell state. *Cell* **144**, 940-954, doi:10.1016/j.cell.2011.01.032 (2011).
Natoli, G. Maintaining cell identity through global control of genomic organization. *Immunity* **33**, 12-24, doi:10.1016/j.immuni.2010.07.006 (2010).
Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**, 663-676 (2006).
Graf, T. Historical origins of transdifferentiation and reprogramming. *Cell Stem Cell* **9**, 504-516, doi:10.1016/j.stem.2011.11.012 (2011).
Apostolou, E. & Hochedlinger, K. Chromatin dynamics during cellular reprogramming. *Nature* **502**, 462-471, doi:10.1038/nature12749 (2013).
Chronis, C. *et al.* Cooperative Binding of Transcription Factors Orchestrates Reprogramming. *Cell* **168**, 442-459 e420, doi:10.1016/j.cell.2016.12.016 (2017).
Soufi, A., Donahue, G. & Zaret, K. S. Facilitators and impediments of the pluripotency reprogramming factors' initial engagement with the genome. *Cell* **151**, 994-1004, doi:10.1016/j.cell.2012.09.045 (2012).

8   Chen, J. *et al.* Hierarchical Oct4 Binding in Concert with Primed
Epigenetic Rearrangements during Somatic Cell Reprogramming. *Cell Rep* **14**,
1540-1554, doi:10.1016/j.celrep.2016.01.013 (2016).
9   Sridharan, R. *et al.* Role of the murine reprogramming factors in the
induction of pluripotency. *Cell* **136**, 364-377, doi:S0092-8674(09)00007-5 [pii]
10.1016/j.cell.2009.01.001 (2009).
10   Knaupp, A. S. *et al.* Transient and Permanent Reconfiguration of
Chromatin and Transcription Factor Occupancy Drive Reprogramming. *Cell Stem*
Cell **21**, 834-845 e836, doi:10.1016/j.stem.2017.11.007 (2017).
11   Apostolou, E. *et al.* Genome-wide Chromatin Interactions of the Nanog
Locus in Pluripotency, Differentiation, and Reprogramming. *Cell stem cell* **12**,
699-712, doi:S1934-5909(13)00149-5 [pii]10.1016/j.stem.2013.04.013 (2013).
12   de Wit, E. *et al.* The pluripotent genome in three dimensions is shaped
around pluripotency factors. *Nature* **501**, 227-231, doi:10.1038/nature12420
(2013).
13   Denholtz, M. *et al.* Long-range chromatin contacts in embryonic stem cells
reveal a role for pluripotency factors and polycomb proteins in genome
organization. *Cell Stem Cell* **13**, 602-616, doi:10.1016/j.stem.2013.08.013 (2013).
14   Wei, Z. *et al.* Klf4 Organizes Long-Range Chromosomal Interactions with
the Oct4 Locus in Reprogramming and Pluripotency. *Cell stem cell*, doi:S1934-
5909(13)00203-8 [pii]10.1016/j.stem.2013.05.010 (2013).
15   Krijger, P. H. *et al.* Cell-of-Origin-Specific 3D Genome Structure Acquired
during Somatic Cell Reprogramming. *Cell Stem Cell* **18**, 597-610,
doi:10.1016/j.stem.2016.01.007 (2016).
16   Beagan, J. A. *et al.* Local Genome Topology Can Exhibit an Incompletely
Rewired 3D-Folding State during Somatic Cell Reprogramming. *Cell Stem Cell*
18, 611-624, doi:10.1016/j.stem.2016.04.004 (2016).
17   Rubin, A. J. *et al.* Lineage-specific dynamic and pre-established enhancer-
promoter contacts cooperate in terminal differentiation. *Nat Genet* **49**, 1522-1528,
doi:10.1038/ng.3935 (2017).
18   Drissen, R. *et al.* The active spatial organization of the beta-globin locus
requires the transcription factor EKLF. *Genes Dev* **18**, 2485-2490,
doi:10.1101/gad.317004 (2004).
19   Schoenfelder, S. *et al.* Preferential associations between co-regulated
genes reveal a transcriptional interactome in erythroid cells. *Nat Genet* **42**, 53-61,
doi:10.1038/ng.496 (2010).
20   Stadtfeld, M., Maherali, N., Borkent, M. & Hochedlinger, K. A
reprogrammable mouse strain from gene-targeted embryonic stem cells. *Nat*
Methods **7**, 53-55, doi:10.1038/nmeth.1409 (2010).
21   Polo, J. M. *et al.* A molecular roadmap of reprogramming somatic cells
into iPS cells. *Cell* **151**, 1617-1632, doi:10.1016/j.cell.2012.11.039 (2012).
22   Stadtfeld, M. *et al.* Ascorbic acid prevents loss of Dlk1-Dio3 imprinting and
facilitates generation of all-iPS cell mice from terminally differentiated B cells. *Nat*
Genet **44**, 398-405, S391-392, doi:10.1038/ng.1110 (2012).
23   Stadtfeld, M. *et al.* Aberrant silencing of imprinted genes on chromosome
12qF1 in mouse induced pluripotent stem cells. *Nature* **465**, 175-181,
doi:10.1038/nature09017 (2010).

24   Stadtfeld, M., Maherali, N., Breault, D. T. & Hochedlinger, K. Defining
molecular cornerstones during fibroblast to iPS cell reprogramming in mouse.
Cell Stem Cell **2**, 230-240, doi:10.1016/j.stem.2008.02.001 (2008).
25   McLean, C. Y. *et al.* GREAT improves functional interpretation of cis-
regulatory regions. *Nat Biotechnol* **28**, 495-501, doi:10.1038/nbt.1630 (2010).
26   Soufi, A. *et al.* Pioneer transcription factors target partial DNA motifs on
nucleosomes to initiate reprogramming. *Cell* **161**, 555-568,
doi:10.1016/j.cell.2015.03.017 (2015).
27   Whyte, W. A. *et al.* Master transcription factors and mediator establish
super-enhancers at key cell identity genes. *Cell* **153**, 307-319,
doi:10.1016/j.cell.2013.03.035 (2013).
28   Stadhouders, R. *et al.* Transcription factors orchestrate dynamic interplay
between genome topology and gene regulation during cell reprogramming. *Nat*
Genet, doi:10.1038/s41588-017-0030-7 (2018).
29   Mumbach, M. R. *et al.* HiChIP: efficient and sensitive analysis of protein-
directed genome architecture. *Nat Methods* **13**, 919-922,
doi:10.1038/nmeth.3999 (2016).
30   Lazaris, C., Kelly, S., Ntziachristos, P., Aifantis, I. & Tsirigos, A. HiC-
bench: comprehensive and reproducible Hi-C data analysis designed for
parameter exploration and benchmarking. *BMC Genomics* **18**, 22,
doi:10.1186/s12864-016-3387-6 (2017).
31   Sheffield, N. C. & Bock, C. LOLA: enrichment analysis for genomic region
sets and regulatory elements in R and Bioconductor. *Bioinformatics* **32**, 587-589,
doi:10.1093/bioinformatics/btv612 (2016).
32   Mumbach, M. R. *et al.* Enhancer connectome in primary human cells
identifies target genes of disease-associated DNA elements. *Nat Genet* **49**,
1602-1612, doi:10.1038/ng.3963 (2017).
33   Dixon, J. R. *et al.* Topological domains in mammalian genomes identified
by analysis of chromatin interactions. *Nature* **485**, 376-380,
doi:10.1038/nature11082 (2012).
34   Dowen, J. M. *et al.* Control of cell identity genes occurs in insulated
neighborhoods in mammalian chromosomes. *Cell* **159**, 374-387,
doi:10.1016/j.cell.2014.09.030 (2014).
35   Gomez-Diaz, E. & Corces, V. G. Architectural proteins: regulators of 3D
genome organization in cell fate. *Trends Cell Biol* **24**, 703-711,
doi:10.1016/j.tcb.2014.08.003 (2014).
36   Weintraub, A. S. *et al.* YY1 Is a Structural Regulator of Enhancer-
Promoter Loops. *Cell* **171**, 1573-1588 e1528, doi:10.1016/j.cell.2017.11.008
(2017).
37   Beagan, J. A. *et al.* YY1 and CTCF orchestrate a 3D chromatin looping
switch during early neural lineage commitment. *Genome Res* **27**, 1139-1152,
doi:10.1101/gr.215160.116 (2017).
38   Schoenfelder, S. *et al.* Polycomb repressive complex PRC1 spatially
constrains the mouse embryonic stem cell genome. *Nat Genet* **47**, 1179-1186,
doi:10.1038/ng.3393 (2015).

39   Kearns, N. A. *et al.* Functional annotation of native enhancers with a
Cas9-histone demethylase fusion. *Nat Methods* **12**, 401-403,
doi:10.1038/nmeth.3325 (2015).
40   Li, D. *et al.* Chromatin Accessibility Dynamics during iPSC
Reprogramming. *Cell Stem Cell* **21**, 819-833 e816,
doi:10.1016/j.stem.2017.10.012 (2017).
41   Samstein, R. M. *et al.* Foxp3 exploits a pre-existent enhancer landscape
for regulatory T cell lineage specification. *Cell* **151**, 153-166,
doi:10.1016/j.cell.2012.06.053 (2012).
42   Ghaleb, A. M. & Yang, V. W. Kruppel-like factor 4 (KLF4): What we
currently know. *Gene* **611**, 27-37, doi:10.1016/j.gene.2017.02.025 (2017).
43   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system.
Nat Protoc **8**, 2281-2308, doi:10.1038/nprot.2013.143 (2013).
44   Liu, Y. *et al.* Widespread Mitotic Bookmarking by Histone Marks and
Transcription Factors in Pluripotent Stem Cells. *Cell Rep* **19**, 1283-1293,
doi:10.1016/j.celrep.2017.04.067 (2017).
45   Buenrostro, J. D., Wu, B., Chang, H. Y. & Greenleaf, W. J. ATAC-seq: A
Method for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol*
109, 21 29 21-29, doi:10.1002/0471142727.mb2129s109 (2015).
46   Consortium, E. P. An integrated encyclopedia of DNA elements in the
human genome. *Nature* **489**, 57-74, doi:10.1038/nature11247 (2012).
47   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon
provides fast and bias-aware quantification of transcript expression. *Nat Methods*
14, 417-419, doi:10.1038/nmeth.4197 (2017).
48   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.
Nat Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).
49   Tsirigos, A., Haiminen, N., Bilal, E. & Utro, F. GenomicTools: a
computational platform for developing high-throughput analytics in genomics.
Bioinformatics **28**, 282-283, doi:10.1093/bioinformatics/btr646 (2012).
50   Gong, Y. *et al.* Stratification of TAD boundaries reveals preferential
insulation of super-enhancers by strong boundaries. *Nat Commun* **9**, 542,
doi:10.1038/s41467-018-03017-1 (2018).
51   Crane, E. *et al.* Condensin-driven remodelling of X chromosome topology
during dosage compensation. *Nature* **523**, 240-244, doi:10.1038/nature14450
(2015).
1222

1223