A model-based high throughput method for fecundity estimation in fruit fly

studies.

Enoch Ng'oma[1*], Elizabeth G. King[1], and Kevin M. Middleton[2]

[1]Division of Biological Sciences, University of Missouri, Columbia, MO 65211

[2]Department of Pathology and Anatomical Sciences, University of Missouri School of

Medicine, Columbia, MO 65212

*Corresponding author:

Enoch Ng'oma

401 Tucker Hall

Division of Biological Sciences

University of Missouri, Columbia, MO 65211

Tel.: 573-882-8562

E-mail: ngomae@missouri.edu

Abstract

The ability to quantify fecundity is critically important to a wide range of experimental applications, particularly in widely-used model organisms such as *Drosophila melanogaster.* However, the standard method of hand-counting eggs is time consuming and limits the feasibility of large-scale experiments. We develop a predictive model to automate the counting of eggs from images of eggs removed from the media surface and washed onto dark filter paper. A cross-validation approach demonstrates our method performs well, with a correlation between predicted and hand counted values of 0.88. We show how this method can be applied to a large data set where egg densities vary widely.

Keywords: image analysis; fecundity; *Drosophila melanogaster*; quantitative biological imaging; cross-validation

Introduction

Reproductive output, with its close tie to Darwinian fitness, is potentially the most important of an individual's phenotypes and thus a critical phenotype to be able measure in many different experiments. While counting the number of eggs or offspring produced by an individual is simple in concept, in practice it is often quite challenging. In many insect systems, including the widely used model organism *Drosophila melanogaster (D. melanogaster)*, females produce a large number of eggs over their lifetime, and the most common method of quantifying lifetime fecundity is by hand counting eggs soon after they are laid. These features limit the ability to quantify lifetime

fecundity for large numbers of experimental replicates. Thus, the development of

methods that allow the automation of egg counting while maintaining accuracy have the

potential to expand the set of biological questions that can be investigated.

As image analysis techniques in general have gotten more sophisticated, there is

a growing interest in applying these techniques to different types of biological data[1].

There are several challenges associated with applying image analysis methods to

fecundity data in *D. melanogaster*. Female flies lay eggs on the food media, which for

most standard recipes[2] is close in color to the eggs and does not provide a high

contrast background. This challenge has been addressed in the past by providing egg

laying media that is higher contrast (e.g., transparent media[3] or the addition of charcoal

to the media[4]), or by washing the eggs off of the surface of the food and filtering them

through black filter paper[5–7]. Irregularities on the surface of the food, the presence of

dust or other particulate matter, and the clumping of many eggs together are all issues

that have the potential to affect any automated image analysis. Existing methods for

automating egg counting in *Drosophila* have developed methods to identify individual

eggs on the food surface and then quantify the total number present[3,4]. While these

methods perform well for some applications, their performance declines when eggs are

at high densities with a lot of clumping, or when there are many other similarly sized

non-egg objects in the image. In addition, because both rely on images of the food

surface, for recipes that produce lightly colored food, which may be necessary in studies

with a specific diet manipulation, it becomes more difficult for these methods to

accurately quantify eggs.

We sought to develop a method for robustly predicting egg counts applicable to a

wide range of applications, with a minimal impact of the challenges described above.

Our method first produces images with a high contrast between eggs and the

background and then takes advantage of the simple relationship between the area of

light-colored pixels and the number of eggs present to develop a predictive model that

can be applied to a large set of images. This method has the flexibility to be applied to a

wide range of egg densities and can be applied to many experimental contexts.

Methods

Fecundity samples

The set of samples used to develop and validate our egg counting method were

part of a quantitative genetic study employing a half-sibling, split family design, with

families split over three different dietary conditions using flies derived from the

Drosophila Synthetic Population Resource[8]. All vials began with 15-24 female flies and

6 male flies with fecundity estimated once per week over a 24-hour period for the entire

lifetime of the flies. Females were provided with fresh media each Monday and following

a 24-hour egg laying period, these egg vials were collected the following Tuesday and

stored at -20C until processing. We continued this process each week until all females

within a vial had died. To visualize eggs, we modified a protocol developed by Rose and

colleagues[5–7]. We added 2.5 mL of 50% bleach to each vial (frozen or thawed) and

gently swirled for about 2 min 50 s on an ORBITRON Rotator 1 (Boekel Scientific, USA)

to separate eggs from the media. Eggs were flushed from each vial onto a black filter

disc fit into a custom-built vacuum apparatus with a brief spray of embryo solution (12%

Triton X-100, SIGMA-ALDRICH, USA) followed by a spray of water to rinse. As a quality

check, flask filtrate was re-filtered to a new disc periodically, and processed vials

inspected under a dissecting microscope. The filter discs were prepared from black

landscape fabric (DEWITT Weed Barrier Pro, www.dewittcompany.com) to a radius of

3.65 cm.

Discs bearing eggs were photographed using a Canon EOS Rebel TSi (Canon

Inc., Japan) camera vertically attached to a steel clamp using the following settings:

exposure time 1/25 seconds, aperture F5.6, and ISO 100. Egg-bearing discs were

mounted on a black stage placed within a cubical photo studio 35cm x 35cm x 35cm

constructed with PVC frame and enclosed by a white fabric (Fig. 1). The entire setup

was placed in a dark room lit only by two facing white light lamps, 23W, 1200 lumens

(model # LBP23PAR2/5K, UTILITECH, China) clamped to the stage outside the studio

box to allow for soft uniform lighting across the imaging platform. Image files were

immediately inspected for quality with ImageJ[9].

Estimation of egg counts

We developed a predictive model to estimate the counts of eggs from the images

of the filter discs. To train the model and assess its performance, eggs on 345 discs

were manually counted using the *Multi-point* tool in ImageJ[9]. To ensure that all possible

imaging conditions were available for model training, manually counted discs were

selected across the whole lifespan of flies and across all three diet treatments. Some

background reflectivity of the discs and small debris present meant that some non-zero

white area was present in almost all images, so our training set includes an over-

representation of images with relatively few eggs, to prevent upward bias in the

fecundity predictions.

Each image was manually cropped to be square at the edges of the disc, and the

background outside the disc was set to black to eliminate any reflectivity or debris

occurring outside of the disc using GNU Image Manipulation Program[10]. Subsequently,

the image was binarized by thresholding 8-bit grayscale values using the opencv

package[11] in Python[12] (v. 3.6; https://github.com/EGKingLab/fecundity_estimation).

From this thresholded image, the area of white pixels was summed

(https://github.com/EGKingLab/fecundity_estimation). Because images were not all the

same size due to slight variations in camera zoom, we included size of the image as a

covariate in the linear model predicting egg count from white pixel area and image

dimension:

$$Egg\ Count = \beta_0 + \beta_1 \sqrt{Area} + \beta_2\ ImageSize$$

where *Area* is the sum of the white pixels and *Image Size* is the linear dimension of one

edge of the image in pixels. We use $\sqrt{Area}$ because pixel area is predicted to scale as a

square of egg count.

The optimal grayscale threshold value was determined by cross-validation using

R[13] (v. 3.5 https://github.com/EGKingLab/fecundity_estimation). We tested threshold

values between 30 and 150 (for 8-bit images with 256 levels of gray), across a range of

training/testing data splits (0.5 to 0.9) and a range of proportions of data (0.5 to 1.0).

The latter was used to test the adequacy of our data. Each model was fit 1,000 times at

each combination of threshold, train/test split, and data proportion, with the root mean

squared difference between predicted count and manual count (RMSd) retained for

each iteration (https://github.com/EGKingLab/fecundity_estimation). The optimal

threshold value was determined from the mean RMSd for the 1,000 iterations at each

combination (Fig. 2). That threshold was 53 for the training set, using 90% of the data

each iteration for training, and only required using 50% of the total data in any one

iteration. The resulting predictive model performed well for our training set (Fig. 3a). The

trained model was then used to predict egg counts for all 3,768 images that were part of

this dataset (Fig. 3b). Images with negative predicted egg counts were set to 0. All

images were also visually inspected, and those determined to be of poor quality, with

excessive background reflectivity, were counted manually and were not included in the

training or prediction set.

Data and software availability

Our raw data, including our fecundity estimates, and all egg image files, have

been uploaded to Zenodo and may be retrieved at

http://doi.org/10.5281/zenodo.1285237. All scripts to reproduce all of our analyses are

available at GitHub: https://github.com/EGKingLab/fecundity_estimation. Code is

available to directly download egg image files from the Zenodo and reproduce our

analyses.

Results

We developed a high throughput method for obtaining egg counts by creating a

predictive model to obtain estimated egg counts from images of eggs that were

separated from the food surface and filtered onto a dark fabric disc by taking advantage

of the simple relationship between the number of eggs present and the amount of white

area on a thresholded image of a disc containing *D. melanogaster* eggs (Fig. 2a). We

used a set of hand counted images to optimize the model and show that it performs well

(Fig 3a).

We tested several parameters to produce an optimal prediction model. First, by

using different proportions of our total dataset of hand counted images, we showed our

hand counted set was sufficiently large to train the predictive model and achieve a high

predictive ability. Using only 50% of the hand counted data, we achieved similar error

rates in the resulting model, demonstrating our dataset was sufficiently large to saturate

the model (Fig 2c,d). We also tested a range of thresholds to identify the value that

would remove the most background noise without obscuring actual eggs (Fig 2c,d). To

assess the robusticity of our model across different parameters, we employed cross

validation to ensure we tested the ability of the model to predict egg counts for new data

(data not used to fit the model). For each iteration, 90% of the data was used to train the

model and we assessed the ability of this model to predict the values of the remaining

10% of the data. Our best performing model was able to predict new data well, with a

mean absolute error of ~177 eggs and a high correlation (r = 0.89 between the

predicted egg count and the manual egg counts. We note that with a 24-hour egg laying

period for a vial with several females, this error rate represents a relatively small

number of eggs. After identifying the optimal threshold value, we fit a model using our

full dataset and found similar error rates when comparing the predicted values from this

model to the hand counted images (r = 0.88, mean absolute error = 128 eggs; Fig. 3a).

We were then able to use this model to predict egg counts for our entire dataset of

3,768 images (Fig. 3b).

Discussion

The method of washing eggs from the surface of the food onto black filter paper has been used previously in studies quantifying fecundity in *D. melanogaster*[5–7]. While this is more time consuming than imaging the surface of the food directly, this approach has several advantages. First, removing eggs from the surface of the food typically results in less clumped eggs and removes surface irregularities such as bubbles in the food. Second, it allows for eggs to be imaged on a high contrast background no matter what food recipe is used. This feature is particularly important when diet is a manipulated variable in an experiment and/or it is undesirable to add charcoal or some other darkening agent to the food. Finally, any imaging (including on the food surface) provides a permanent record of the phenotype while hand-counting and discarding vials does not allow for validation of the phenotyping or later error checking. In previous work using this method, eggs washed onto black filter paper were still manually counted. Here, we have shown how this process can be automated, providing a significant time savings, e.g., avoiding manual counts of ~84,000 eggs in this study.

Other automated egg counting methods rely on identifying individual objects in an image as eggs, including those of Waithe et al.[3] and Nouhaud et al.[4] It is not surprising then that both methods perform more poorly when egg densities are high and multiple eggs clump together. The method we have developed is able to handle a wide range of densities because the egg count is estimated directly from the area of white pixels, removing the need to identify eggs as individual objects. The method described here is susceptible to debris on the filter paper surface (e.g., food particles, etc.), because any lightly colored object will be included in the total thresholded area. We included a large

number of low egg count discs to help calibrate the average amount of background

debris on our discs. Nevertheless, it is important when applying this method to make

sure minimal amounts of debris are present. This may be achieved either by careful

egg-washing and/or pre-cleaning discs before imaging (and recording eggs that lie on

the debris being removed).

Overall, we present an approach to automating egg counting that can be optimized

and tailored to a wide variety of situations and applications. We use a cross-validation,

train/test approach to identify the model parameters that allow us to optimally predict

new egg count values. The cross-validation approach shows a high correspondence

between predicted and hand-counted egg counts for new data (data not used to form

the predictive model), showing our high performance is not inflated due to overfitting to

our hand-counted set. This approach could potentially be applied to any images with

enough contrast to threshold the background from the objects of interest and may be

especially applicable to situations where it is challenging to separate individual objects.

Acknowledgements

Larry Cabral provided with valuable ideas through personal communication to

establish our fecundity protocol. Elizabeth Lopresti, Michael Reed, Osvaldo Enriquez,

Anna Perinchery and Kyla Winford helped with fly husbandry, time-intensive

experimental set-ups, collection and entry of data. This work was supported by NIH

grant R01 GM117135 to Elizabeth G. King, The University of Missouri, and a University

of Missouri Research Board Grant.

References

1. Nketia TA, Sailem H, Rohde G, Machiraju R, Rittscher J. Analysis of live cell images: Methods, tools and opportunities. Methods 2017; 115:65–79.

2. Bloomington Drosophila Stock Center. Fly Food Recipes. 2017 accessed 2018 Aug 1; Available from: https://bdsc.indiana.edu/information/recipes/index.html

3. Waithe D, Rennert P, Brostow G, Piper MDW. QuantiFly: Robust Trainable Software for Automated Drosophila Egg Counting. PLOS ONE 2015; 10:e0127659.

4. Nouhaud P, Mallard F, Poupardin R, Barghi N, Schlötterer C. High-throughput fecundity measurements in Drosophila. Sci Rep 2018; 8.

5. Rose Lab Data. Fecundity: Animated Introduction. 2012; Available from: https://www.youtube.com/watch?v=xrEBNB8bUwo&sns=em

6. Rose Lab Data. Fecundity: Live Demonstration. 2012; Available from: https://www.youtube.com/watch?v=uqFZ4ZTC6ZY&sns=em

7. Burke MK, Barter TT, Cabral LG, Kezos JN, Phillips MA, Rutledge GA, Phung KH, Chen RH, Nguyen HD, Mueller LD, et al. Rapid divergence and convergence of life-history in experimentally evolved *Drosophila melanogaster*. Evolution 2016; 70:2085–98.

8. Ng'oma E, Fidelis W, Middleton K, King EG. The evolutionary potential of diet-dependent effects on lifespan and fecundity in a multi-parental population of *Drosophila melanogaster*. bioRxiv 2018; :343947.

9. Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, Eliceiri

KW. ImageJ2: ImageJ for the next generation of scientific image data. BMC

Bioinformatics 2017; 18:529.

10. The GIMP Team. GNU Image Manipulation Program. 2015. Available from:

http://www.gimp.org

11. Bradski, G. The OpenCV Library. Dr Dobb's Journal of Software Tools 2000.

12. Python Software Foundation. Python Language Reference. Available from:

http://www.python.org/

13. R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing. 2018; Vienna, Austria. Available from:

http://www.R-project.org/

Figure 1:

Figure 2:

Figure 3

Figure 1: Image acquisition set up. Egg discs were imaged using a Canon EOS Rebel

TS*i* camera vertically attached to a steel clamp using the following settings: exposure

time 1/25 seconds, aperture F5.6, and ISO 100. The discs were mounted on a black

stage placed within a cubical photo studio 35cm x 35cm x 35cm constructed with PVC

frame and enclosed by a white fabric. The entire setup is placed in a dark room lit only

by two facing white light lamps, clamped to the stage outside the studio box to allow for

soft uniform lighting across the imaging platform.

Figure 2: Performance of the model predicting egg counts from images. **a.** A

representative raw image of eggs separated from fly media and filtered onto black filter

fabric. **b.** The same image thresholded at the optimal threshold level identified by the

model. Due to the time required for optimization of threshold value, we carried this

process out in two steps. **c.** Coarse optimization was carried out for every 5 grayscale

values between 30 and 150, which revealed a single minimum in that range, falling

between 50 and 60. **d**. Fine optimization was then carried for integer values between 50

and 60, resulting in a single optimal value of 53. Colors represent the proportion of data

used for each model fit. Even at the optimal value, the difference in minimum (at 50% of

data) and maximum (at 100% of data) in root mean squared difference was ~2 eggs.

Figure 3: The performance of the predictive model. **a.** The square root of image area is

on the x-axis and the egg count is on the y-axis. True, hand-counted values are shown

in red and the predicted values from the model are shown in blue. Pairs of values

(hand-counted and predicted) for individual images are connected by a red line. **b.**

Distribution of predicted values for all images in the data set. Inflated zero counts results

from assigning a value of 0 to any predicted values <0.