Liquid droplet germ granules require assembly and localized regulators for mRNA

repression

Scott Takeo Aoki[1], Tina R Lynch[1], Sarah L Crittenden[1,2], Craig A Bingman[1], Marvin Wickens[1],

Judith Kimble[1,2,*]

[1] Department of Biochemistry and [2] Howard Hughes Medical Institute, University of Wisconsin-

Madison, Madison, WI, USA

* For correspondence:

Judith Kimble

HHMI/Department of Biochemistry

University of Wisconsin-Madison

433 Babcock Drive

Madison, WI 53706-1544

Tel 608-262-6188

Fax 608-265-5820

jekimble@wisc.edu

Keywords: RNA-protein granule, germ granule, mRNA repression, scaffold protein, Argonaute

Abbreviations: RNA, ribonucleic acid; UTR, untranslated region; GFP, green fluorescent

protein; CRISPR, Clustered Regularly Interspaced Short Palindromic Repeats; RNAi,

ribonucleic acid interference

Running title: P-granule assembly not sufficient for mRNA repression

Summary:

Cytoplasmic RNA-protein (RNP) granules have diverse biophysical properties, from liquid to

solid, and play enigmatic roles in RNA metabolism. Nematode P-granules are paradigmatic

liquid droplet granules and central to germ cell development. Here we analyze a key P-granule

scaffolding protein, called PGL, to investigate the functional relationship between P-granule

assembly and function. Using a protein-RNA tethering assay, we find that reporter mRNA

expression is repressed when recruited to PGL granules. We determine the crystal structure of

the PGL N-terminal region to 1.5 Å, discover its dimerization and identify key residues at the

dimer interface. *In vivo* mutations of those interface residues prevent P-granule assembly, de-

repress PGL-tethered mRNA and reduce fertility. Therefore, PGL dimerization lies at the heart

of both P-granule assembly and function. Finally, we identify the P-granule-associated

Argonaute WAGO-1 as crucial for repression of PGL-tethered mRNA. We conclude that P-

granule function requires both assembly and localized regulators.

Introduction

RNA-protein (RNP) granules, otherwise known as biomolecular condensates [1], are ubiquitous

non-membrane bound "organelles". Some RNP granules exist in a solid-like state with little

component exchange [2], while others behave as liquid-like droplets with components dynamically

diffusing in and out of the granule [3, 4]. Intriguingly, mRNA regulators localizing to liquid granules

do not rely on granule association for their activities [5]. Therefore, despite intense interest and an

ever-expanding literature, the functional relationship between granule assembly and mRNA

regulation remains a mystery.

Here, we investigate the functional relationship between RNP granule assembly and function in

C. elegans germline P-granules. P-granules are paradigmatic RNP granules [6, 7] with striking

similarities to germ granules in *Drosophila* and vertebrates, both in subcellular location and

composition [8]. They are crucial for fertility and totipotency [9], and some of their components can

display liquid droplet behavior [6, 10, 11]. In adult germ cells, P-granules localize to the cytoplasmic

face of nuclear pores (**Figure 1A**) and contain both untranslated mRNAs [12] and numerous RNA

regulatory proteins [13]. Yet their molecular function is poorly defined. RNAi-mediated gene

repression is seated there [14] and loss of P-granules correlates with aberrant upregulation of

somatic transcripts [15, 16, 17]. Yet it remains unclear whether mRNA repression depends on

granule scaffold assembly or on localized regulators in granules. A definitive understanding of

this fundamental question requires manipulation of mRNA localization as well as manipulation of

granule formation without eliminating pivotal assembly proteins. The former is made possible

with a protein-mRNA tethering assay [18, 19, 20], but the latter requires a deeper understanding of

P-granule scaffolding proteins and the molecular basis of their assembly.

Two PGL proteins, PGL-1 and PGL-3, are the key scaffolding proteins required for P-granule

assembly [21, 22]. Genetic removal of PGL proteins causes mislocalization of P-granule proteins [23],

aberrant expression of spermatogenic and somatic mRNAs [15, 16, 17], and temperature-dependent

sterility [21, 22]. PGL proteins self-assemble into granules, both *in vitro* using purified recombinant

protein [24] and in intestinal nematode cells or mammalian cells in culture when expressed on

their own [25, 26]. Artificial PGL granules display liquid droplet behavior [6, 11, 24], indicating that PGL

protein alone is sufficient to recapitulate the biophysical properties of P-granules in nematode

cells [10].

PGL assembly into granules was poorly understood prior to this work though progress had been

made. PGL-1 and PGL-3 are close paralogs [22] with the same architecture (**Figure 1B**). Many

RNP granule assembly proteins rely on low complexity sequences for multivalent-multivalent,

low affinity interactions [1]. The only low complexity sequences in PGL are RGG repeats at the C-

terminus, which mediate PGL RNA binding and are dispensable for granule formation [24, 25].

Assays for granule assembly in mammalian cells implicated the conserved N-terminal region of

PGL as critical [25], and structural studies identified a central dimerization domain (DD) within that

conserved region (**Figure 1B**) [27]. Yet higher-ordered self-assembly demands additional PGL-

PGL contacts.

In this work, we employ a tethering assay to manipulate mRNA localization in and out of

granules, structural analyses to identify a new PGL dimerization domain and incisive mutational

intervention to discover the role of dimerization in P-granule assembly and mRNA regulation.

Our findings provide evidence that repression of mRNA expression in P-granules requires both

assembly and localized regulators and hence makes a major advance in understanding the

functional relationship between RNP granule assembly and function.

Results

Tethered PGL represses a reporter mRNA in nematode germ cells

Prior studies have suggested that P-granules regulate mRNA expression (see Introduction). To

test this notion directly, we relied on a protein-mRNA tethering assay (**Figure 1C**) [18, 19], widely

used to investigate RNA regulatory proteins [20]. Our assay examined the expression of mRNAs

to which the PGL-1 protein was tethered via λN22, a short peptide that binds with high affinity

and specificity to the boxB RNA hairpin [18]. This assay has been used successfully to identify the

functions of a variety of RNA binding proteins in several organisms, including nematodes [18, 28].

For the reporter, we inserted three boxBs into the 3'UTR of an established GFP-histone

transgene that is ubiquitously expressed throughout the germline [29] (**Figure 1C**, Methods). To

tether PGL to the GFP reporter mRNA via boxB, we generated PGL::SNAP::λN22 with

sequential CRISPR gene editing in an internal, non-conserved protein region of PGL-1 (**Figures**

1C and **S1A,** see Methods). The SNAP tag [30] is used to visualize subcellular localization and

λN22 provides tethering. PGL::SNAP::λN22 homozyotes were sterile (100%, n=94), but could

be maintained and tested as a fertile heterozygote (PGL-1::SNAP::λN22/+). Given that *pgl-1* null

homozygotes are fertile [21], the fertility defects from the addition of λN22 to PGL-1 is likely not

due to defective protein function. Regardless, the logic of our tethering strategy is simple: if

tethered PGL-1 localizes to granules and represses GFP expression as predicted, this assay

provides a powerful entrée into fundamental questions about granule function.

To evaluate mRNA regulation, we assayed reporter GFP fluorescence in both living animals and

fixed, extruded gonads; the former facilitated scoring many samples and the latter permitted

scoring subcellular localization of both PGL protein via SNAP and *gfp* RNA with single molecule

fluorescence *in situ* hybridization (smFISH). In controls carrying PGL-1::SNAP without λN22,

GFP fluorescence was robust (**Figure 1D,F**), but GFP fluorescence was absent in animals with

PGL-1::SNAP::λN22 (**Figure 1E,G**). Essentially the same result was found in fixed germlines:

PGL-1::SNAP without λN22 germlines expressed GFP (100% gonads, n=35) (**Figure 1H** and

S2C,N), but PGL-1::SNAP with λN22 had GFP was faintly detected in only a few gonads (5%

gonads, n=39) (**Figure 1I** and **S2G,N**). Importantly, the PGL-1::SNAP and PGL-1::SNAP::λN22

proteins both assembled into cytoplasmic granules at the nuclear periphery (**Figure 1J,K** and

S2D,H), similar to untagged PGL-1 and PGL-3 reported previously [21, 22]. The SNAP signal was

less for PGL-1::SNAP::λN22, perhaps because animals were heterozygous. We conclude that

tethered PGL-1 localizes to perinuclear granules and represses expression of the reporter

mRNA.

We next asked if the reporter RNA localized with PGL in perinuclear granules. To this end, we

used smFISH to detect *gfp* RNAs and SNAP staining to detect PGL (**Figure S2A**). Control germ

cells expressing PGL-1::SNAP without λN22 possessed *gfp* RNAs in both nuclear and

cytoplasmic puncta (**Figure S2B-E**). We interpret nuclear puncta as nascent transcripts at

active transcription sites and cytoplasmic puncta as mRNAs, based on a previous study [31]. GFP

fluorescence was robust (**Figure S2C,K**), and PGL-1::SNAP localized to perinuclear granules

(**Figure S2D**), as in **Figure 1**. Germ cells expressing PGL-1::SNAP::λN22 lacked robust GFP

fluorescence (**Figure S2G,K**) and PGL-1::SNAP::λN22 localized to perinuclear granules (**Figure**

S2H), also as in **Figure 1**. The cytoplasmic RNA puncta were less diffuse with tethered PGL-1

than in the control and frequently colocalized with PGL-1::SNAP::λN22 in perinuclear granules

(**Figure S2F-I,O**; see **Figure S3** for additional images). Imaging of the rachis, the shared

cytoplasmic space in the germline, did not reveal substantial aggregates of reporter RNA as

observed with mRNA in other studies [32]. The presence of *gfp* cytoplasmic transcripts (**Figure**

S2F) demonstrates that the reporter was not transcriptionally silenced, despite the lack of GFP

fluorescence. In summary, reporter RNA localized with PGL-1 when tethered and PGL-1

tethering repressed reporter protein expression. These results support the idea that P-granules

are sites of post-transcriptional repression.

Identification of the PGL N-terminal Dimerization Domain

To test the significance of P-granule assembly to mRNA repression, we sought to perturb PGL-

1 assembly into granules. An emerging principle is that multivalent-multivalent interactions drive

granule formation (e.g. protein with at least two multimerization domains) [1, 33]. Because PGLs

are key assembly proteins for P-granules and can form granules on their own [6, 11, 24, 25], we

reasoned that PGLs use multiple self-interactions to drive granule assembly. We previously

identified one dimerization domain (DD) centrally in PGL [27], but DD missense mutations grossly

affected protein stability. We postulated the existence of another PGL multimerization domain

that might be more amenable to manipulation and again turned to structural analyses.

The region N-terminal to DD has high sequence conservation (**Figure S1B**), which implies a

critical role in PGL function. Our initial efforts to express trypsin-mapped recombinant protein

fragments [27] of this N-terminal region proved unfruitful. However, addition of amino acids

disordered in the DD crystal structures [27] permitted robust expression sufficient for biochemical

and structural characterization (**Figure S4A,B**, see Methods for more details). Henceforth, we

refer to this stable protein fragment as the N-terminal dimerization domain (NtDD) (**Figure 1B**).

We determined the *C. japonica* PGL-1 NtDD crystal structure to 1.5 Å (**Figure 2**, see **Table S1**

for statistics, see Methods for details on crystallization and structure determination). The NtDD

had a novel fold consisting of 11 alpha helices and a single N-terminal beta strand (**Figure 2B**).

The asymmetric unit (ASU) was composed of four NtDD domains (**Figure 2A**), which were

structurally similar (RMSD 0.219 - 0.254, chains B-D aligned to A), except for minor differences

in termini and internal loops. While each ASU possessed two pairs of identical interfaces

(**Figure 2A**), one of these interface pairs consisted of a network of conserved amino acid side

chains making extensive salt bridges and hydrogen bonds (**Figure 3A-C** and **Figure S4C-E**).

The complexity and conservation of these interactions suggested biological relevance. We next

tested for dimerization *in vitro*. Recombinant PGL-3 NtDD formed a dimer on a sizing column

combined with multi-angle light scattering (SEC-MALS, **Figure 3D,E**). To ask if the conserved

interface in the NtDD crystal structure might be its dimerization interface, we used our structural

model and *in silico* prediction [34] to design missense mutations predicted to disrupt the interface.

These analyses yielded two distinct mutants: R123E with a single mutated residue and K126E

K129E with two mutated residues. Both NtDD interface mutants formed monomers rather than

dimers in solution (**Figure 3D,E**). We conclude that the dimers observed in the crystal structure

represent the NtDD dimer detected in solution. Therefore, PGL proteins possess two

dimerization domains. We renamed the original DD to Central DD (CDD) for clarity (**Figure 1B**).

PGL NtDD dimerization is critical for granule formation

To assess the role of NtDD dimerization in PGL granule self-assembly, we used an assay in

mammalian cells where expression of wild-type PGL-1 tagged with GST was sufficient for

assembly into granules [25]. Similar to that report, wild-type PGL-1 tagged with GFP also formed

large cytoplasmic granules when expressed in mammalian cells (**Figure 3F,G**), while GFP

alone was diffuse (**Figure 3H**). However, PGL-1::GFP mutated to either K126E K129E or

R123E no longer self-assembled into granules (**Figure 3I,J**). We conclude that NtDD

dimerization is essential for granule formation in mammalian cells.

To assess the role of NtDD dimerization in nematode germ cells, we used CRISPR to introduce

the dimerization defective mutations into SNAP-tagged PGL-1 (**Figure 4A**, **Figure S1A**,

Methods). We first asked about effects on fertility (**Figure 4B**). Most wild-type PGL-1 (no SNAP)

and PGL-1::SNAP animals were fertile at 20°C and 25°C (**Figure 4B**). In contrast, most *pgl-1*

null mutants were fertile at 20° but few were fertile at 25°C (**Figure 4B**), as reported previously

[21, 22]. Fertility of the NtDD dimerization mutants, K126E K129E and R123E, by contrast, was

sharply reduced at both 20° and 25°; many mutant worms were sterile at 20°C and most were

sterile at 25°C (**Figure 4B**). Fertility was therefore impacted more severely in dimerization

mutants than *pgl-1* null mutants (**Figure 4B**), suggesting that a PGL mutant protein incapable of

NtDD dimerization functions as a dominant-negative. The NtDD dimerization mutants had

smaller than normal germlines and many lacked oocytes (**Figure S5B-D**), which are defects

typical of *pgl-1* and *pgl-1 pgl-3* null mutants [21, 22]. We conclude that PGL-1 NtDD dimerization is

critical for fertility.

To investigate the role of PGL NtDD dimerization in granule assembly, we compared the

subcellular localization of wild-type PGL-1::SNAP to NtDD dimerization-defective PGL-1::SNAP

mutant proteins (**Figure 4C-F**). Wild-type PGL-1::SNAP assembled into cytoplasmic granules at

the nuclear periphery (**Figure 4C**), but the K126E K129E and R123E mutant proteins were

largely diffuse in both fertile (**Figure 4E,F**) and sterile (**Figure S5E,F**) germlines. Protein

expression levels may affect the propensity of PGL-1 to form granules but were difficult to

compare due to variability in germline size. Mutant PGL-1 fluorescent intensity was above

background (**Figure S6A-E**), albeit less than wild-type PGL-1. Immunoblots also revealed only a

modest difference in protein expression between PGL-1 wild-type and mutant protein (**Figure**

S6F). Despite these differences, the mutant PGLs were capable of forming small perinuclear

granules in a variable number of germ cells (**Figure 4E,F** and **Figure S5E,F**). In addition, for

each mutant, we found a single germline (1 of 59 for K126E K129E; 1 of 54 for R123E) with

PGL-1 perinuclear granules in all germ cells (**Figure S5G,H**). Therefore, both PGL-1 mutant

proteins are capable of incorporating into granules, but do so much more inefficiently than their

wild-type counterparts (**Figure 5C**).

We next asked why the fertility defects of PGL-1 NtDD dimerization mutants were more severe

than a *pgl-1* null mutant. The likely explanation was interference with assembly of other P-

granule components into granules. Normally, PGL-1 interacts with PGL-3 [22], and both PGL-1

and PGL-3 rely on GLH-1 or GLH-4 Vasa helicases to localize to the nuclear periphery in adult

germ cells [26, 35]. In contrast, GLH proteins can assemble at the nuclear pore independently of

PGLs [22]. We postulated that PGL-1 assembly mutants might interfere with PGL-3 assembly into

granules but not affect GLH-1. To test this idea, we epitope-tagged endogenous *pgl-3* and *glh-1*

(see Methods) and compared localization of PGL-3::V5 and GLH-1::Myc in germ cells also

expressing either wild-type PGL-1::SNAP or the dimerization defective mutant PGL-1(K126E

K129E)::SNAP. In the presence of wild-type PGL-1::SNAP, all three proteins, PGL-1, PGL-3

and GLH-1, co-localized to granules at the nuclear periphery (**Figure 4G-K**), as previously

observed for untagged proteins [21, 22, 36]. By contrast, a dimerization-defective mutant protein,

PGL-1(K126E K129E)::SNAP, was diffuse rather than granular, and wild-type PGL-3 became

similarly diffuse (**Figure 4L-N**). GLH-1, however, was still capable of localizing to granules at the

nuclear periphery (**Figure 4O-P**), similar to previous reports [22]. Although these GLH-1 granules

appeared smaller than normal, their formation was seen around virtually all germline nuclei

(**Figure 4P**). We conclude that NtDD dimerization of PGL-1 is crucial for assembly of both PGL-

1 and PGL-3 into granules and that this likely explains the severe fertility defects of PGL-1

dimerization mutants.

PGL granule assembly is required for PGL-mediated mRNA repression

The identification of assembly-defective PGL-1 proteins coupled with our tethering assay

(**Figure 5A**) allowed us to test the relationship between granule assembly and mRNA

repression. We introduced the K126E K129E mutation into PGL-1::SNAP::λN22 and tested for

reporter expression. PGL-1(K126E K129E)::SNAP::λN22 homozygotes were fertile (21%, n=96)

to an extent comparable to PGL-1(K126E K129E)::SNAP without λN22 (**Figure 4B**). While

control PGL-1::SNAP::λN22 repressed the reporter (**Figure 5D,F**), assembly-defective PGL-1

was not repressive and the vast majority of germ cells expressed GFP (**Figure 5E,G and**

S2K,N). The PGL-1(K126E K129E)::SNAP::λN22 mutant protein was diffuse and non-granular

compared to wild-type PGL-1::SNAP (**Figure 5H,I**), similar to PGL-1(K126E K129E)::SNAP

without λN22 (**Figure 4E**). By smFISH, *gfp* RNA signal in PGL-1(K126E K129E)::SNAP::λN22

mutant germlines was observed robustly throughout the cytoplasm (**Figure S2J-M**). Formally,

the PGL-1 interface residues might affect granule assembly and mRNA repression

independently but the simplest explanation is that PGL-1 must assemble into granules to

repress mRNA.

PGL-mediated mRNA repression relies on WAGO-1, a cytoplasmic Argonaute

PGL tethering provides a simple assay for identification of additional factors needed for P-

granule mRNA repression (**Figure S7A**). We depleted candidate P-granule-associated RNA

regulators with RNAi and sought GFP reporter de-repression in PGL-1::SNAP::λN22 worms. In

this candidate screen, knockdown of the cytoplasmic Argonaute WAGO-1 had a dramatic effect

(**Figure S7B**). RNAi against other candidates, by contrast, had either no or a minor effect on

GFP repression (**Figure S7B**). The RNAi screen highlighted WAGO-1 as a key factor in PGL-

mediated mRNA repression, a finding consistent with previous studies showing that WAGO-1

localizes to P-granules and regulates gene expression [37, 38].

To further investigate *wago-1*, we first inserted an epitope-tag at the endogenous locus (**Figure**

S7C, Methods). WAGO-1::3xV5 co-localized with PGL-1(wt)::SNAP in perinuclear granules

(**Figure 6A**), consistent with a previous report that WAGO-1 resides in P-granules [37]. We next

asked if WAGO-1 association with perinuclear granules was dependent on PGL assembly. In

animals expressing assembly-defective PGL-1(R123E)::SNAP, which fails to form granules

efficiently (**Figure 4E,F**), WAGO-1 was diffuse in about half the germlines (**Figure 6B**), but

granular in the other half (**Figure 6C**). Therefore, WAGO-1 can assemble into P-granules

independently of PGL-1. To investigate if PGL assembly is independent of WAGO-1, we

generated an internal deletion that creates a frameshift and fails to express WAGO-1 protein

(**Figure 6D**, **Figure S7C**, Methods). PGL-1::SNAP localized to perinuclear granules in the

absence of WAGO-1 (**Figure 6D).** In sum, PGL-1 and WAGO-1 can assemble into granules

independently of each other.

We finally explored the role of WAGO-1 in mRNA repression within granules (**Figure 7A**). To

this end, we tested PGL-1::SNAP::λN22 for its ability to repress reporter RNA in the presence or

absence of WAGO-1. We again conducted our assays in living animals to ensure a large

sample size (**Figure 7B-E**) and fixed extruded gonads to visualize PGL in addition to GFP

fluorescence (**Figure 7F-I**). The reporter was repressed with wild-type WAGO-1 (**Figure 7D,F**),

as expected, but de-repressed in the *wago-1* null mutant (**Figure 7E,G**), consistent with *wago-1*

RNAi (**Figure S7B**). The de-repression was seen in ~70% of germlines when assayed in living

animals (**Figure 7E**) and in ~90% of fixed gonads (**Figure 7G**). This more penetrant de-

repression in fixed gonads might reflect higher sensitivity or lower sample size. Loss of *wago-1*

had no observable effect on fluorescent reporter expression (**Figure S8**), arguing against

varying reporter mRNA levels causing the increase in fluorescence. Regardless, the

percentages demonstrate strong but incomplete de-repression, a result consistent with WAGO-

1 functioning redundantly with other Argonautes [37, 38, 39]. As expected, PGL-1::SNAP::λN22

assembles into granules, with or without WAGO-1 (**Figure 7H,I**). Moreover, by smFISH, the *gfp*

reporter mRNA colocalizes with PGL-1 both in animals with wild-type WAGO-1 (**Figure S9A-**

D,I) and in the *wago-1* null mutant (**Figures S9E-I** and **S10**). We conclude that WAGO-1 is a

major regulator of mRNA repression in P-granules. These results also provide direct evidence

that granule formation alone is not sufficient for mRNA regulation (see **Figure 8**, Discussion).

Discussion

This work investigates the functional relationship between assembly of a paradigmatic liquid

droplet, the *C. elegans* P-granule, and the activities of its regulatory components. Our analyses

make three key advances. First, we discover that PGL-1 dimerization of the N-terminal domain

is required for its assembly into P-granules. Second, we demonstrate that PGL-mediated mRNA

repression relies on PGL granule assembly. Third, we find that mRNA repression by P-granules

employs the activity of at least one P-granule constituent, the Argonaute WAGO-1. Together,

these advances support a model that PGL assembly into granules is necessary for its biological

function but not sufficient to repress the expression of localized mRNAs (**Figure 8**). Below we

discuss these advances and their implications for RNP granules more generally.

Dimerization drives PGL protein assembly into granules

An emerging principle of RNP granule assembly is that multivalent macromolecular interactions

drive granule formation [1, 33]. This work extends that principle to nematode P-granules and their

primary assembly proteins, the PGLs (see Introduction). A previous study reported that PGL

proteins possess a dimerization domain (DD) in their central region [27], and here we report the

discovery of a second PGL DD in the N-terminal region (NtDD). Thus, PGLs possess two

protein folds that confer multivalency. Based on insights from the NtDD structure, we designed

two distinct PGL-1 mutant proteins that are unable to dimerize *in vitro* and are severely

compromised *in vivo* for assembly into P-granules and fertility. These results provide evidence

that PGL dimerization is crucial for P-granule assembly and for P-granule biological function.

Although attempts to disrupt dimerization via the central DD led to protein instability and hence

could not test its biological significance [27], we include both N-terminal and central DDs in our

model for how PGLs mediate assembly into P-granules (**Figure 8A**). A critical future direction is

to investigate how each dimerization domain contributes to higher order and likely oligomeric

assembly.

Self-assembling, multivalent proteins have been identified for several RNP granules. Examples

include TDP-43 in nuclear granules [40], Oskar and Vasa for *Drosophila* polar granules [41, 42, 43];

EDC3 and LSM4 for P-bodies [44]; and MEG-3 and MEG-4 for embryonic P-granules [6, 45, 46].

These various assembly proteins rely on a combination of multimerization domains and low

complexity, intrinsically disordered sequences to facilitate granule formation [6, 44, 45, 47, 48, 49]. A

leading hypothesis of liquid droplet assembly invokes reliance on multiple, weak interactions

among RNP constituents [1, 33]. Our work highlights the idea that interactions between structured

regions are a driving force *in vivo* for assembly of liquid droplet granules as well as their

biological function (see below). Recombinant PGL proteins make liquid droplets on their own *in*

vitro [6, 11, 24], suggesting that PGLs possess regions responsible for low affinity contacts in the

full-length protein. We suggest that PGL uses both dimerization domains as well as additional

low affinity contacts to facilitate liquid droplet formation. Alternatively, the DDs may be subject to

post-translational modifications that modulate their affinity *in vivo*. PGL did not require its C-

terminal RGG repeats to form granules in mammalian culture [25], but RGG repeats in other

assembly proteins enhance granule formation [49]. The PGL RGG repeats may instead be

needed to trigger robust granule assembly with RNA [24] or impart liquid droplet properties

associated with PGL in nematodes [10]. Regardless, the discovery that PGL proteins are

multivalent with two dimerization domains is an essential advance for understanding the

assembly of this paradigmatic liquid droplet.

PGL-mediated mRNA repression relies on recruitment into P-granules

Our study employs a protein-RNA tethering assay to provide direct evidence that mRNA

expression is repressed when the mRNAs are recruited to P-granules. Three lines of indirect

evidence had previously led to this idea: P-granules contain repressed mRNAs [12];

spermatogenic and somatic transcripts are aberrantly expressed in the absence of P-granules

[15, 16, 17]; and a variety of RNA regulatory factors co-localize in P-granules, including inhibitory

RNA binding proteins, like the Pumilio homolog FBF-2 [32], and RNA regulatory enzymes, like the

Argonaut/Piwi PRG-1 [50] and the deadenylase PARN-1 [51]. Despite these clues, the significance

of P-granule assembly for biological function remained enigmatic. Our discovery of key amino

acids required for PGL assembly into granules (see above) allowed us to compare RNA

regulation when coupled to assembly-competent or assembly-defective PGL. We found that a

reporter mRNA tethered to assembly-competent PGL protein localized with PGL to P-granules

and was repressed, but a reporter tethered to an assembly-defective PGL did not localize to

granules and was expressed. Together, these results suggest that localizing mRNAs into P-

granules leads to their repression (**Figure 8A,B**).

Argonaute WAGO-1 promotes repression of mRNA expression in P-granules

Expression of mRNAs localized within an RNP granule might be repressed by either of two

broad mechanisms. Granule localization might recruit RNAs to form interactions with negative-

acting factors (e.g. RNA turnover machinery) or it might sequester RNAs away from positive-

acting factors (e.g. translational machinery). Our results suggest that localization with

assembled PGL and key repressors is the primary mechanism (**Figure 8**). Previous work

identified an *in vitro* RNase activity for PGL DD [27]. However, PGL-tethering of RNA, and

subsequent recruitment into granules, is not sufficient for repression on its own but also relies

on a P-granule-localized Argonaute, WAGO-1 (**Figure 8C**). Previous studies reported that

WAGO-1 is P-granule-associated and represses transcripts that have been primed by the

piRNA pathway as part of the secondary RNAi response [38]. Our tethering results broaden the

role of WAGO-1 to repress any RNA associated with the P-granule scaffold. Previous studies

also reported that WAGO-1 is functionally redundant with other Argonautes [37, 38, 39]. We suspect

this redundancy may explain why some, not all, *wago-1* null mutant germlines repressed PGL-

tethered reporter mRNAs. The specific mechanism for mRNA repression, mRNA turnover or

translational repression, is still unclear, although we favor RNA turnover driven by PGL's RNase

activity and the Argonaute. Additional work must be done to understand the relative importance

and individual roles of these Argonautes and PGL in mRNA repression.

Other cytoplasmic RNP granules have been proposed to repress mRNAs. Granule formation of

a yeast amyloid-like RNA binding protein correlates with translational inhibition of transcripts

critical for gametogenesis [52], and P-bodies contain mRNAs that are translationally repressed in

cells [53]. Our work extends this idea further by showing the dependence of other enzymatic

factors in liquid droplet granules for mRNA repression. Our model proposes that loss of PGL

dimerization prevents P-granule assembly and hence compromises the coalescence of RNAs

with WAGO-1 and other P-granule-associated repressors (**Figure 8B**), while loss of WAGO-1

relieves granule-recruited mRNAs from repression (**Figure 8C**). Therefore, the scaffolding

function of PGL is important but not sufficient for P-granule function. We also propose that,

without WAGO-1, mRNAs that are normally destined for P-granule association can exist in

either of two states. Those remaining in the granule are not expressed due either to the

presence of other RNA repressors or the absence of ribosomes, but those diffusing out of the

liquid droplet granule reach the translational machinery for expression (**Figure 8C**). By

extension, we propose that other RNP granules with liquid droplet properties may require both

scaffolding proteins and active enzymatic factors to regulate their mRNAs.

This work adds to the emerging theme that granules play a general role in regulating the

function of their components. For example, stress granules sequester the mTORC1 protein

complex to block activation of mTOR signaling [54], mammalian cells can trap hormones and

melanin in amyloid-like aggregates to prevent active signaling [55, 56], and liquid droplet granule

formation of cGAS with cytosolic DNA triggers its enzymatic activity *in vitro* and cell culture [57].

The functional relationship between PGL and WAGO-1 is unknown. PGL may simply recruit or

retain mRNAs in P-granules to be regulated by WAGO-1. A more enticing model is that PGL

forms a higher ordered, RNP complex with mRNA that enhances WAGO-1's biochemical

activity. Further studies that pair insights into mechanisms of granule assembly with direct *in*

vivo assays of regulation will be pivotal moving forward to decipher the mechanistic function of

other RNP granules in their biological context.

Materials and Methods

Protein expression and purification

We previously used *C. elegans* PGL-3 recombinant protein and limited proteolysis to identify a central

dimerization domain (CDD) [27]. While we could express CDD efficiently we could not express recombinant

protein that was N-terminal to the cleavage site (PGL-3 amino acid residues 205-206). We tried moving

the six-histidine purification tag to the N- and C- termini, shortened the protein regions used for

expression, and tried several different orthologs with little success. The insight came after aligning protein

sequences of several Caenorhabditid sp. and studying the CDD domain boundary (**Figure S1B**).

Protease cleavage occurred in a conserved portion of the N-terminal region and this region was

disordered in our DD crystal structures. After inclusion of this region (PGL-3 amino acid residues 205-

212), we could express and purify recombinant N-terminal protein from *C. elegans* PGL-1 and its

orthologs. We refer to this region as the N-terminal domain (NtDD).

This study used primarily *C. elegans* PGL-3 and *C. japonica* PGL-1 recombinant NtDD proteins. The *C.*

elegans PGL-3 coding region was PCR amplified from cDNA. A codon-optimized (*E. coli*) version of *C.*

japonica PGL-1 NtDD was ordered as a gBlock (IDT, Coralville, IA). We included a six-histidine tag at the

C-terminus that was removed later with carboxypeptidase A [58]. Constructs were cloned into a pET21a

vector (MilliporeSigma, Burlington, MA) using Gibson Assembly cloning [59], and plasmids transformed into

Rosetta2 cells (MilliporeSigma, Burlington, MA). Cultures were grown at 37°C with shaking (225 rpm) until

~0.8 OD, cooled for 30-60 minutes, and induced with a final concentration of 0.1 mM IPTG. Cultures were

then grown at 16°C with shaking (160 rpm) for 16-18 hours, collected, and bacterial pellets frozen until

use. Selenomethionine-incorporated *C. japonica* protein was expressed in SelenoMethionine Medium

Complete (Molecular Dimensions, Suffolk, UK), and grown, induced, and collected in a similar manner.

Bacterial pellets were defrosted on ice and reconstituted in lysis buffer (20 mM sodium phosphate pH 7.4,

300 mM NaCl, 10 mM imidazole, 5 mM beta-mercaptoethanol (BME)) with protease inhibitors

(cOmplete™ EDTA-free, Roche, Indianapolis, IN). Lysozyme (Sigma-Aldrich, St. Louis, MO) was added

at 50 μg/ml and incubated on ice for 20 minutes prior to lysis in a french press. Samples were spun at low

(3220 x g, 4°C, 20 minutes) and high speed (10,000 x g, 20°C, 10 minutes), then incubated with 1.5 ml

NiNTA beads (Thermo Fisher Scientific, Waltham, MA) for 1 hour at 4°C with rotation. Sample

supernatant was separated by gravity flow, washed twice with lysis buffer, and eluted using lysis buffer

with increasing imidazole concentrations (20, 40, 60, 80, 100, 250 mM). Eluted samples were checked for

protein via Bradford assay (Bio-Rad, Hercules, CA), and dialyzed overnight in HN buffer (20 mM HEPES

pH 7.4, 100 mM NaCl). The dialyzed samples were concentrated with a Centriprep 10K concentrator

(Millipore), calcium added to 1 mM $CaCl_2$, and the histidine tag removed with carboxypeptidase A bound

to agarose (Sigma, St. Louis, MO) at a ratio of 10 protein:1 enzyme (w/w). Samples were incubated at

room temperature (~20°C) for 45-90 minutes with rotation prior to supernatant elution by centrifugation in

microflow columns (Thermo Fisher Scientific, Waltham, MA). Samples were run on a S200 sizing column

(GE Healthcare, Chicago, IL) in HNT buffer (20 mM HEPES pH 7.4, 100 mM NaCl, 0.5 mM TCEP pH

7.4). Fractions containing recombinant protein were collected, concentrated in an Amicon 10K

concentrator (MilliporeSigma, Burlington, MA), and protein concentration estimated by A280. Samples

were frozen in liquid nitrogen or used immediately.

Crystallization and structure determination

C. elegans PGL-1, *C. elegans* PGL-3, and *C. japonica* PGL-1 NTD recombinant protein were screened in

crystallization conditions using 400 nl hanging and sitting drop 96-well trays set up with the Mosquito

(TTP Labtech, Cambridge, MA) in 20°C. Several conditions produced labile crystal plates. Data was

collected to 4 Å from *C. elegans* PGL-1 crystal plates, determined to have a very large unit cell (86 Å x 86

Å x 460 Å) and P6 point group, and eventually determined to have perfect merohedral twinning. *C.*

japonica PGL-1 also crystallized as large (60-150 Å) rhomboid crystals in 40-45% PEG 400 at low (Na

Citrate pH 5.5-6.0) and physiologic pH (imidazole pH 7.5-8.0). Crystals grown in citrate or imidazole both

diffracted well, but we used imidazole (100 mM imidazole pH 7.5, 45% PEG 400, 1 mM TCEP pH 7.4)

due to its higher reproducibility for large crystals and its modestly better resolution. The crystals did not

require additional cryo-protection due to the high PEG 400. We eventually collected a full data set to 1.5

Å in space group C2.

PGL-1 NtDD was a novel domain. Novelty and translational pseudosymmetry precluded us from using

any model for molecular replacement. Trial heavy atom soaks also proved unfruitful, and the *C. japonica*

PGL-1 NtDD has just two methonines past the start codon, making selenomethionine phasing

challenging. To boost anomalous signal, we mutated two non-conserved isoleucines to methionines

(I63M, I212M). This methionine mutant provided phases to 3.6 Å by single anomalous dispersion (SAD)

that we used to build a 1.6 Å model of the mutant protein (PDB ID: **5W4D**). We used this model for

molecular replacement into the wild-type data set to build a complete 1.5 Å model (PDB ID: **5W4A**). Data

and model statistics are in **Table S2**. Model coordinates and data are available at RCSB (www.rcsb.org).

Size exclusion chromatography with multi-angle laser light scattering (SEC-MALS)

Molecular weights of *C. elegans* PGL-3 NtDD wild type and mutant recombinant protein were determined

by conducting SEC-MALS experiments using Agilent Technologies 1260 LC HPLS system (Agilent

Technologies, Santa Clara, CA) equipped with Dawn® Heleos™II 18-angle MALS light scattering

detector, Optilab® T-rEX™ (refractometer with EXtended range) refractive index detector, WyattQELS™

quasi-elastic (dynamic) light scattering (QELS) detector and ASTRA software (all four from Wyatt

Technology Europe GmbH, Dernbach, Germany). A total of 500 µL (1 mg/mL) of the samples in HNT

buffer (20 mM HEPES pH 7.5, 100 mM NaCl, 0.5 mM TCEP pH 7.4) were injected and run on a Superdex

75 10/300 GL column (GE Healthcare) pre-equilibrated with the same buffer, at a flow rate of 0.5 mL/min

at 20°C. Lysozyme (Sigma-Aldrich, St. Louis, MO) was used as a control.

Mammalian cell culture maintenance, transfection and imaging

Full length PGL-1 was cloned into a pcDNA 3.1 vector (Thermo Fisher Scientific, Waltham, MA) with a C-

terminal eGFP and OLLAS epitope linker. Mutations to PGL-1 were created using Gibson Assembly

cloning [59]. Chinese Hamster Ovary (CHO) cells (ATCC, Manassas, VA) were propagated according to

distributor's recommendations. Briefly, cells were grown in F-12K Medium (Thermo Fisher Scientific,

Waltham, MA) with 10% fetal bovine serum (Gibco), and split with Trypsin 0.25% (Gibco) every 2-3 days.

Cells were grown to 70% confluence and transfected with TransIT-CHO Transfection Kit (Mirus Bio LLC,

Madison, WI). Transfected cells were split the following day and grown in Ibitreat 15 u-Slide 8 well slides

(Ibidi, Madison, WI) overnight. Hoechst stain (Invitrogen, Carlsbad, CA) was added to wells prior to

imaging by confocal microscopy for GFP and Hoechst fluorescence, and transmitted light. Well dilutions

were chosen based on adequate cell spacing to discern each cell, and 25 fields of view were taken based

on the highest concentration of GFP-positive cells. Experiments were repeated four times with similar

results. During image collection, we observed a single example of a granule-like blob in the PGL-

1::OLLAS::GFP K126E K129E. The cell appeared unhealthy, and thus the granule may be an artifact of

cell death, but we included it in our study for completeness.

Worm maintenance, CRISPR mutagenesis, fertility and imaging

Frozen strains:

N2 Bristol

JK5687: *pgl-1(q894)[PGL-1::SNAP] IV*

JK5902: *pgl-1(q975)[PGL-1(R123E)::SNAP] IV*

JK6158: *wago-1 q1087[WAGO-1::3xV5]; pgl-1(q894)[PGL-1::SNAP] IV*

JK6159: *wago-1 q1089[WAGO-1(null deletion)::3xV5]; pgl-1(q894)[PGL-1::SNAP] IV*

JK6157: *wago-1 q1087[WAGO-1::3xV5]; pgl-1(q975)[PGL-1(R123E)::SNAP] IV*

JK5898: *glh-1(q858)[GLH-1::3xMYC] I; pgl-1(q894)[PGL-1::SNAP] IV; pgl-3(q861)[PGL-3::3xV5] V*

JK5970: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q894)[PGL-1::SNAP] IV*

JK5873: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1::SNAP::λN22]/nT1[qIs51](IV;V)*

JK5874: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1::SNAP::λN22]/nT1[qIs51](IV;V)*

JK6149: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1::SNAP::λN22]/nT1[qIs51](IV;V)*

JK6150: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1:SNAP:λN22]/nT1[qIs51](IV;V)*

JK6147: *wago-1 q1089[WAGO-1(null deletion)::3xV5]; qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1::SNAP::λN22]/nT1[qIs51](IV;V)*

JK6148: *wago-1 q1089[WAGO-1(null deletion)::3xV5]; qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q994)[PGL-1::SNAP::λN22]/nT1[qIs51](IV;V)*

JK6367: *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II*

JK6368: *wago-1 q1089[WAGO-1(null deletion)::3xV5]; qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II*

Worm strains that could not be frozen:

1. *pgl-1(q960)[PGL-1(K126E K129E)::SNAP] IV*

2. *glh-1(q858)[GLH-1::3xMYC] I; pgl-1(q960)[PGL-1(K126E K129E)::SNAP] IV; pgl-3(q861)[PGL-3::3xV5]*

V

3. *qSi375[(mex-5 promoter::eGFP::linker::his-58::3xboxb::tbb-2 3'UTR) *weSi2] II; pgl-1(q1053)[PGL-*

1(K126E K129E)::SNAP::λN22]/nT1[qIs51](IV;V)

C. elegans were maintained as previously reported [60]. For CRISPR-Cas9 mutagenesis, a Cas9 protein

co-conversion approach was used [61]. Briefly, worms were injected with a target CRISPR-Cas9 RNA

(crRNA) or a plasmid expressing a Cas9-scaffold with tandem target sequence RNA (sgRNA) to a gene

of interest [61], a target crRNA to *dpy-10* or *unc-58*, a scaffolding tracrRNA (IDT), recombinant Cas9 protein

[62], a *dpy-10/unc-58* repair DNA oligo that inserted a dominant mutation [61], and an epitope tag/missense

mutant repair oligo or PCR product. See below for a Table of guide RNAs and repair templates used. F1s

with the co-injection marker phenotype were additionally screened by a combination of PCR without or

with restriction enzyme digest to identify those with the repair of interest. In JK5687, a SNAP tag [30] was

inserted between PGL-1 amino acids G713 and G714 in N2 worms. A 3xMYC tag was added to the N-

terminus of GLH-1 between G17 and F18. A 3xV5 tag was added in the C-terminal region of PGL-3

between residues G627 and S628. A 3xV5 tag was added to the C-terminus of WAGO-1 between

residues E914 and A915. To generate the *wago-1* null allele, a WAGO-1::3xV5 allele was mutated so that

648 base pairs were deleted from the N-terminus and proper coding frame shifted to add premature stop

codons (**Figure S7C**). The *wago-1* null allele was confirmed by staining and imaging (**Figure 6D**). F2s

were PCR screened to identify homozygous SNAP alleles and the PCR product sequenced to confirm

proper repair. Three worm strains were too infertile to freeze. All worms were outcrossed at least twice

with N2, with the exception of *glh-1(q858)[GLH-1::3xMYC] I; pgl-1(q960)[PGL-1(K126E K129E)::SNAP]*

IV; pgl-3(q861)[PGL-3::3xV5] V that was backcrossed with JK5898.

Worms were singled into the peripheral wells of a 24-well plate that contained NGM agar and OP50

bacteria. Worms were allowed to propagate for 5 days at 20°C or 25°C, and then scored for progeny and

gravid progeny. We report the progeny numbers here.

CRISPR-Cas9 guide RNAs and repair oligos

Name	Type	Strain targeted	mutation	Sequence
CRISPR-Cas9 guide RNAs:				
glh-1 sgRNA 1	CRISPR-Cas9 sgRNA plasmid	N2	3xmyc	target sequence: TCCACTACCGAATCCAGTTT
pgl-3 sgRNAin1	CRISPR-Cas9 sgRNA plasmid	N2	3xV5	target sequence: GCAACGGAACGTCTGGAAG
pgl-1 crRNA 1	CRISPR-Cas9 RNA	N2	SNAP	target sequence: cccaccagttcagcttatgg
pgl-1 crRNA 5	CRISPR-Cas9 RNA	JK5687, JK5898, JK5874	K126E K129E	target sequence: gtcttcagtcttcttcagct
pgl-1 crRNA 8	CRISPR-Cas9 RNA	JK5687	R123E	target sequence: cttcttcagcttggccttac
SNAP crRNA 1	CRISPR-Cas9 RNA	JK5687	λN22	target sequence: CCTGGGCTGGGTCCTGCAGG
wago-1 crRNA 1	CRISPR-Cas9 RNA	N2	3xV5	target sequence: TGCGACTCCTTTGCTTCAAT
wago-1 crRNA 2	CRISPR-Cas9 RNA		null deletion	target sequence: ATTGGAGTCATAGCTCCTGG

Name	Type	Strain targeted	mutation	Sequence	Enzyme screen
wago-1 crRNA 3	CRISPR-Cas9 RNA		null deletion	target sequence: AAATCTGGGCAAGCGTACTG	

CRISPR-Cas9 repair oligos

Name	Type	Strain targeted	mutation	Sequence	Enzyme screen
DNA repair template:					
glh-1 3xmyc repair 1	ssDNA repair oligo	N2	3xmyc	ttccaccggtttattttgattaaaaactttatttcagCgAAAACTGG AAACGAACAGAAGCTTATTTCCGAGGAAGACC TCGCCGGAGAGCAAAAGCTCATCTCTGAAGAG GATCTTGGAGCCGAACAGAAGCTTATCTCTGA AGAAGACCTCGGAGGATTCGGTAGTGGAGGC GGTTTCGGTGGTGGTAACAATGGAG	n/a
pgl-3 3xV5 repair 1	ssDNA repair oligo	N2	3xV5	agtttgccagcagcaacggaacCtcCggaCgaggcGGAAAG CCAATCCCAAACCCACTCCTCGGACTCGACTC CACCGGAGGAAAGCCAATCCCAAACCCACTCC TCGGACTCGACTCCACCATCGGAAAGCCAATC CCAAACCCACTCCTCGGACTCGACTCCACCGG Atcttatggaggtggtcgcggtggcgatcgt	n/a

				ggattcggtcaatttgctcccaccagttcagcttatggaAGTGGC GGTATGGACAAAGACTGCGAAATGAAGCGCAC CACCCTGGATAGCCCTCTGGGCAAGCTGGAAC TGTCTGGGTGCGAACAGGGCCTGCACCGTATC ATCTTCCTGGGCAAAGGAACATCTGCCGCCGA CGCCGTGGAAGTGCCTGCCCCAGCCGCCGTG CTGGGCGGACCAGAGCCACTGATGCAGGCCA CCGCCTGGCTCAACGCCTACTTTCACCAGCCT GAGGCCATCGAGGAGTTCCCTGTGCCAGCCCT GCACCACCCAGTGTTCCAGCAGGAGAGCTTTA CCCGCCAGGTGCTGTGGAAACTGCTGAAAGTG GTGAAGTTCGGAGAGGTCATCAGCTACAGCCA CCTGGCCGCCCTGGCCGGCAATCCCGCCGCC ACCGCCGCCGTGAAAACCGCCCTGAGCGGAA ATCCCGTGCCCATTCTGATCCCCTGCCACCGG GTGGTGCAGGGCGACCTGGACGTGGGGGGCT ACGAGGGCGGGCTCGCCGTGAAAGAGTGGCT GCTGGCCCACGAGGGCCACAGACTGGGCAAG	
pgl-1 SNAP	PCR product	JK5687	SNAP	CCTGGGCTGGGTCCTGCAGGCGGATCCggagga ggtggtcgcggaggatatggcggtggagaccgtg	n/a
pgl-1 K126E K129E repair	ssDNA repair oligo	JK5687, JK5898, JK5874	K126E K129E	tcgatgacgacaagaagctcggaatgctcgcccgtaaggcTGag ctgaagGagactgaagacgctaagattcttcaagctcttctcaaagt	BlpI
pgl-1 R123E repair	ssDNA repair oligo	JK5687	R123E	tttctgtcatcgatgacgacaagaagctcggaatgctcgcTGAGa aggccaagctgaagaagactgaagacgctaagattcttc	HpyAV

				ctccaccgccatatcctccgcgaccacctcctccGGATCCGTT GGCGGCCTTCCATTGGGCTTGCTTCTCGGCAC GACGCTCACGACGACGGGTACGGGCGTTTCC	
PGL-1 SNAP lambda repair	ssDNA repair oligo	JK5687	λN22	AGAAGAGCCTGCAGGACCCAGCCCAGGCTTG CCCAGTCTGTGGCCCTCGT	n/a
wago-1 3xV5 repair	ssDNA repair oligo	N2	3xV5	ATT TAT TTA TAT TTT GCA GGA CTA AAG GAC CAA TTG AAG GAA AGC CAA TCC CAA ACC CAC TCC TCG GAC TCG ACT CCA CCG GAG GAA AGC CAA TCC CAA ACC CAC TCC TCG GAC TCG ACT CCA CCA TCG GAA AGC CAA TCC CAA ACC CAC TCC TCG GAC TCG ACT CCA CCG GAG CAA AGG AGT CGC AAG GCG AGC GCC TC	n/a
wago-1 deletion repair	ssDNA repair oligo		null deletion	CCA ATG CCA CCA GTC ACT GCT CCA GGA TCA AAA CAT TGA AGT AAA CAA TCG AGA ATA CAC	n/a

Immunoblot

To check PGL-1::SNAP protein expression, worms were boiled in 5x sample buffer (250 mM Tris pH 6.8,

25 mM EDTA pH 8.0, 25% glycerol, 5% SDS, 500 mM beta-mercaptoethanol), run on an SDS-PAGE gel,

and transferred to PVDF. The blot was blocked in 5% dehydrated milk in PBS-T, incubated overnight with

anti-SNAP antibody (NEB, Ipswich, MA), washed and probed with goat-anti-rabbit horseradish peroxidase

secondary antibody (Invitrogen, Carlsbad, CA), and developed with ECL substrate (Thermo Fisher

Scientific, Waltham, MA) and film (Kodak, Rochester, NY). Radiographs were scanned, contrast adjusted

and cropped (Photoshop, Adobe Creative Cloud) as shown.

Fluorescent imaging

To analyze GFP reporter expression, L4 larvae were propagated for approximately 24 hours to adulthood

at 20°C, placed in M9 with 0.1 mM levamisole on a glass slide with a cover slip, imaged at 10x

magnification on a compound microscope and counted for the presence or absence of GFP fluorescence

in their germlines. Numbers represent totals from two separate experiments. The reporter images of live

worms were taken of worms treated in a similar manner and visualized on a Leica SP8 scanning laser

confocal microscope.

For confocal imaging, germlines were extruded, fixed with 1% paraformaldehyde (Electron Microscopy

Sciences, Hatfield, PA) and permeabilized with 0.5% Triton-X as previously described [63]. Germlines were

incubated with 1 µg/ml primary antibodies overnight [anti-MYC (JAC6 (rat), Bio-Rad, Hercules, CA); anti-

V5 (sv5-Pk1 (mouse), Bio-Rad, Hercules, CA)] or 30 nM SNAP JF 549 ligand [64] for 1 hour, stained with

fluorophore-labeled secondary antibodies (Alexa 488 Donkey anti-Mouse, Alexa 555 Donkey anti-Mouse,

Alexa 647 Donkey anti-Mouse, Alexa 488 Goat anti-Rabbit; Invitrogen, Carlsbad, CA) and DAPI

(Invitrogen, Carlsbad, CA), washed and mounted in Vectashield (Vector Laboratories, Burlingame, CA).

Quantitation of GFP and SNAP fluorescence was performed in ImageJ. Graph made in Excel.

Quantitation of GFP and SNAP fluorescence were quantitated in summed intensity projections of confocal

stacks in ImageJ, similar to Crittenden *et al.* [65]. Briefly, a line (40 pixel width) was drawn along the

germline axis. Pixel intensity was measured using plot profile, and intensities averaged and plotted in

Excel.

Single molecule fluorescence *in situ* hybrididation (smFISH)

For smFISH, gonads were extruded, fixed, and hybridized with single molecule FISH probes as described

[31]. The *gfp* exon probe set contains 38 unique oligonucleotides labeled with CAL Fluor Red 610. Briefly,

probes were dissolved in RNase-free TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) to create a 250

µM probe stock. Mid-L4 stage animals were grown on OP50 for 24 hours, then dissected in PBS+0.1%

Tween-20 + 0.25 mM levamisole. Animals were fixed in 4% paraformaldehyde for 20 minutes, incubated

at room temperature in PBS-T (PBS + 0.1% Tween-20) for 10-25 minutes, and equilibrated in smFISH

wash buffer (30 mM sodium citrate pH 7.0, 300 mM NaCl, 1% formamide, 0.1% Tween-20, DEPC water)

for 10-16 minutes. Samples were then incubated in hybridization buffer (30 mM sodium citrate pH 7.0,

300 mM NaCl, 1% formamide, 10% dextran sulfate w/v, DEPC water) plus 0.5 µM smFISH probe at 37°C

for 26-44 hours. 30 nM SNAP 549 ligand was added during the smFISH wash buffer + DAPI wash;

samples were washed at 37°C for approximately 60 minutes. Finally, samples were resuspended in 12 μL

Antifade Prolong Gold mounting medium (Thermo Fisher Scientific, Waltham, MA), mounted on glass

slides, and cured in a dark drawer for at least 24 hours before imaging.

Confocal imaging for smFISH and protein fluorescence

Samples were imaged using a Leica SP8 scanning laser confocal microscope, taking 0.3 μm (smFISH

experiments) or 1 μm (other non-smFISH confocal imaging experiments) slices in sequence. Maximum

intensity partial stack projections were generated and brightness adjusted using ImageJ [66]. All images

were treated equally in ImageJ and Photoshop, with the exception of the transmitted light images.

Imaging experiments were repeated at least twice with similar results, with the exception of PGL-1(K126E

K129E)::SNAP::λN22 worms.

smFISH image analysis

To quantify *gfp* smFISH signal localization between samples, *gfp* smFISH signal and PGL-1::SNAP signal

were each detected then colocalized using Imaris version 9.3.1. The same batch settings were run on all

directly-compared images. In the 3D view for each image, detection for both *gfp* and PGL-1::SNAP

surfaces was checked. If necessary, the edit tool was used to select and delete *gfp* and/or PGL-1::SNAP

surfaces that were detected outside the imaged germline (e.g. on nearby intestine tissue). The *gfp*

surfaces object was selected and the total *gfp* signal was pulled from the statistics tab (Detailed tab,

select Average Values from dropdown, "Intensity Sum Ch=4 Img=1" row and "Sum" column) and stored in

an excel file. The Surface-surface coloc was run from Surface-Surface coloc XTension. The *gfp* and PGL-

1::SNAP channels were selected and the "no smoothing" option was selected. The same statistics tab

navigation was used to pull GFP intensity from the ColocSurface objects that were created by the

XTension. Total GFP intensity in the colocalized surfaces was divided by total GFP intensity in the

germline to calculate how much GFP signal is located in the PGL granules. Some *gfp* and PGL::SNAP

co-localization values were calculated as over 100%. Although colocalization was performed using the

Imaris user interface, the Surface-Surface Colocalization XTension ran an available MATLAB algorithm.

We observed that the Surface-Surface colocalization created a new set of colocalization surfaces, which

are differently shaped and larger than the surfaces written with Imaris algorithms to detect *gfp* or

PGL:SNAP surfaces. Therefore, the total *gfp* intensity in the coloc surfaces was sometimes larger than in

the *gfp* surfaces.

PGL granule staining is sensitive to physical perturbation and some germlines lacked PGL-1::SNAP

signal entirely or almost entirely. GFP smFISH and PGL-1::SNAP signal overlap was therefore calculated

from the top 66% of images in terms of PGL-1::SNAP Intensity (Detailed tab, select Average Values from

dropdown, "Intensity Sum Ch=5 Img=1" row and "Sum" column). Excluding the low PGL images

increased median signal overlap from 26% to 34% in PGL-1::SNAP and from 41% to 57% in PGL-

1::SNAP::λN22 (**Figure S2O**). Excluding the low PGL images increased median signal overlap from

113% to 121% in PGL-1::SNAP::λN22 with WAGO-1 and from 74% to 81% in PGL-1::SNAP::λN22

without WAGO-1 (**Figure S9I**).

Each of the two independent samples were run through a two-tailed t test to calculate significance. We

used the online calculator at http://vassarstats.net.

Imaris batch detection settings

	GFP smFISH channel: Figure S2L	PGL:SNAP channel: Figure S2L	GFP smFISH channel: Figure S9I	PGL:SNAP channel: Figure S9I
[Algorithm]				
Enable Region of Interest	false	false	false	false
Enable Region Growing	true	true	true	true
Enable Tracking	false	false	false	false
[Source Channel]				

Source Channel Index	4	5	4	5
Enable Smooth	true	true	true	true
Surface Grain Size	0.090233 (default -- 2x pixel size)	0.090233 (default -- 2x pixel size)	0.090241 (default)	0.090241 (default)
Enable Eliminate Background	true	true	true	true
Diameter of largest sphere	0.28000 um*	0.60000 um*	0.28000 um	0.60000 um
[Threshold]				
Enable Automatic Threshold	false	false	false	false
Manual Threshold value	2.0241**	5.66575**	3.08326	15.9772
Active Threshold	true	true	true	true
Enable Automatic Threshold B	true	true	true	true
Manual Threshold Value B	19.1165**	14.5472**	97.9303	67.4159
Active Threshold B	false**	false**	false	false
Region Growing Estimated Diameter	0.30000 um*	0.40000 um*	0.30000 um	0.40000 um

[Classify Seed Points]				
"Quality"	above 1.2526*	above 1.9952*	above 3.5559	above 15.052
[Classify Surfaces]				
"Number of Voxels Img = 1"	above 28.000*	above 28.000*	above 28.000	above 28.000

*Values were determined by trial and error, using values from measurements where possible.

**Manual threshold initially set by turning on Upper Threshold (threshold B) -- this changes the value for

the automatic lower threshold. Set the lower threshold to lock that value, then turn off the upper threshold.

This makes the Value B stored, although it isn't actually in use (active = false). Note: the same technique

would provide different values for each image, so the 2.0241 was arbitrarily set on the image settings.

We used a Dell Precision 5820 with a 64-bit Windows 10 Education operating system, an Intel(R)

Xeon(R) W-1245 CPU @3.70GHz processor, and 128 GB of RAM. Imarisx64 9.3.1 and

ImarisFileConverterx64 9.3.1 were installed along with MATLAB R2018b. The Surface-Surface

Colocalization XTension was downloaded and installed from the Bitplane XTension File Exchange

at http://open.bitplane.com/tabid/235/Default.aspx?id=111.

References

1. Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of
    cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298 (2017).
2. Woodruff JB, Hyman AA, Boke E. Organization and function of non-dynamic biomolecular
    condensates. *Trends Biochem Sci* **43**, 81-94 (2018).

3.   Wu H, Fuxreiter M. The structure and dynamics of higher-order assemblies: amyloids,

signalosomes, and granules. *Cell* **165**, 1055-1066 (2016).

4.   Hyman AA, Weber CA, Julicher F. Liquid-liquid phase separation in biology. *Annu Rev*

Cell Dev Biol **30**, 39-58 (2014).

5.   Buchan JR. mRNP granules. Assembly, function, and connections with disease. *RNA Biol*

11, 1019-1030 (2014).

6.   Putnam A, Cassani M, Smith J, Seydoux G. A gel phase promotes condensation of liquid

P granules in *Caenorhabditis elegans* embryos. *Nat Struct Mol Biol* **26**, 220-226 (2019).

7.   Marnik EA, Updike DL. Membraneless organelles: P granules in *Caenorhabditis elegans.*

Traffic **20**, 373-379 (2019).

8.   Voronina E, Seydoux G, Sassone-Corsi P, Nagamori I. RNA granules in germ cells. *Cold*

Spring Harb Perspect Biol **3**, a002774 (2011).

9.   Strome S, Updike D. Specifying and protecting germ cell fate. *Nat Rev Mol Cell Biol* **16**,

406-416 (2015).

10. Brangwynne CP*, et al.* Germline P granules are liquid droplets that localize by controlled

dissolution/condensation. *Science* **324**, 1729-1732 (2009).

11. Zhang G, Wang Z, Du Z, Zhang H. mTOR regulates phase separation of PGL granules to

modulate their autophagic degradation. *Cell* **174**, 1492-1506 e1422 (2018).

12. Schisa JA, Pitt JN, Priess JR. Analysis of RNA associated with P granules in germ cells

of *C. elegans* adults. *Development* **128**, 1287-1298 (2001).

13. Updike D, Strome S. P granule assembly and function in *Caenorhabditis elegans* germ

cells. *J Androl* **31**, 53-60 (2010).

14. Grishok A. Biology and mechanisms of short RNAs in *Caenorhabditis elegans*. *Adv Genet*

83, 1-69 (2013).

15. Updike DL, Knutson AK, Egelhofer TA, Campbell AC, Strome S. Germ-granule

components prevent somatic development in the *C. elegans* germline. *Curr Biol* **24**, 970-

975 (2014).

16. Campbell AC, Updike DL. CSR-1 and P granules suppress sperm-specific transcription in

the *C. elegans* germline. *Development* **142**, 1745-1755 (2015).

17. Knutson AK, Egelhofer T, Rechtsteiner A, Strome S. Germ granules prevent accumulation

of somatic transcripts in the adult *Caenorhabditis elegans* germline. *Genetics* **206**, 163-

178 (2017).

18. Baron-Benhamou J, Gehring NH, Kulozik AE, Hentze MW. Using the λN peptide to tether

proteins to RNAs. *Methods Mol Biol* **257**, 135-154 (2004).

19. Coller J, Wickens M. Tethered function assays using 3' untranslated regions. *Methods* **26**,

142-150 (2002).

20. Bos TJ, Nussbacher JK, Aigner S, Yeo GW. Tethered function assays as tools to elucidate

the molecular roles of RNA-binding proteins. *Adv Exp Med Biol* **907**, 61-88 (2016).

21. Kawasaki I, Shim Y-H, Kirchner J, Kaminker J, Wood WB, Strome S. PGL-1, a predicted

RNA-binding component of germ granules, is essential for fertility in *C. elegans*. *Cell* **94**,

635-645 (1998).

22. Kawasaki I*, et al.* The PGL family proteins associate with germ granules and function

redundantly in *Caenorhabditis elegans* germline development. *Genetics* **167**, 645-661

(2004).

23. Amiri A*, et al.* An isoform of eIF4E is a component of germ granules and is required for

spermatogenesis in *C. elegans*. *Development* **128**, 3899-3912 (2001).

24. Saha S*, et al.* Polar positioning of phase-separated liquid compartments in cells regulated

by an mRNA competition mechanism. *Cell* **166**, 1572-1584 e1516 (2016).

25. Hanazawa M, Yonetani M, Sugimoto A. PGL proteins self associate and bind RNPs to

mediate germ granule assembly in *C. elegans*. *J Cell Biol* **192**, 929-937 (2011).

26. Updike DL, Hachey SJ, Kreher J, Strome S. P granules extend the nuclear pore complex

environment in the *C. elegans* germ line. *J Cell Biol* **192**, 939-948 (2011).

27. Aoki ST, Kershner AM, Bingman CA, Wickens M, Kimble J. PGL germ granule assembly

protein is a base-specific, single-stranded RNase. *Proc Natl Acad Sci USA* **113**, 1279-

1284 (2016).

28. Wedeles CJ, Wu MZ, Claycomb JM. Protection of germline gene expression by the *C.*

elegans Argonaute CSR-1. *Dev Cell* **27**, 664-671 (2013).

29. Zeiser E, Frøkjær-Jensen C, Jorgensen E, Ahringer J. MosSCI and gateway compatible

plasmid toolkit for constitutive and inducible expression of transgenes in the *C. elegans*

germline. *PLoS One* **6**, e20082 (2011).

30. Keppler A, Gendreizig S, Gronemeyer T, Pick H, Vogel H, Johnsson K. A general method

for the covalent labeling of fusion proteins with small molecules *in vivo*. *Nat Biotechnol* **21**,

86-89 (2003).

31. Lee C, Sorensen EB, Lynch TR, Kimble J. *C. elegans* GLP-1/Notch activates transcription

in a probability gradient across the germline stem cell pool. *Elife* **5**, e18370 (2016).

32. Voronina E, Paix A, Seydoux G. The P granule component PGL-1 promotes the

localization and silencing activity of the PUF protein FBF-2 in germline stem cells.

Development **139**, 3732-3740 (2012).

33. Bergeron-Sandoval LP, Safaee N, Michnick SW. Mechanisms and consequences of

macromolecular phase separation. *Cell* **165**, 1067-1079 (2016).

34. Kortemme T, Kim DE, Baker D. Computational alanine scanning of protein-protein

interfaces. *Sci STKE* **2004**, pl2 (2004).

35. Spike C*, et al.* Genetic analysis of the *Caenorhabditis elegans* GLH family of P-granule

proteins. *Genetics* **178**, 1973-1987 (2008).

36. Kuznicki KA, Smith PA, Leung-Chiu WMA, Estevez AO, Scott HC, Bennett KL.

Combinatorial RNA interference indicates GLH-4 can compensate for GLH- 1; these two

P granule components are critical for fertility in *C. elegans*. *Development* **127**, 2907-2916 (2000).

37. Gu W*, et al.* Distinct Argonaute-mediated 22G-RNA pathways direct genome surveillance in the *C. elegans* germline. *Mol Cell* **36**, 231-244 (2009).

38. Shirayama M*, et al.* piRNAs initiate an epigenetic memory of nonself RNA in the *C. elegans* germline. *Cell* **150**, 65-77 (2012).

39. Yigit E*, et al.* Analysis of the *C. elegans* Argonaute family reveals that distinct Argonautes act sequentially during RNAi. *Cell* **127**, 747-757 (2006).

40. Wang A*, et al.* A single N-terminal phosphomimic disrupts TDP-43 polymerization, phase separation, and RNA splicing. *EMBO J* **37**, (2018).

41. Vanzo NF, Ephrussi A. Oskar anchoring restricts pole plasm formation to the posterior of the *Drosophila* oocyte. *Development* **129**, 3705-3714 (2002).

42. Breitwieser W, Markussen FH, Horstmann H, Ephrussi A. Oskar protein interaction with Vasa represents an essential step in polar granule assembly. *Genes Dev* **10**, 2179-2188 (1996).

43. Markussen FH, Michon AM, Breitwieser W, Ephrussi A. Translational control of *oskar* generates Short OSK, the isoform that induces pole plasma assembly. *Development* **121**, 3723-3732 (1995).

44. Decker CJ, Teixeira D, Parker R. Edc3p and a glutamine/asparagine-rich domain of Lsm4p function in processing body assembly in *Saccharomyces cerevisiae*. *J Cell Biol* **179**, 437-449 (2007).

45. Wang JT*, et al.* Regulation of RNA granule dynamics by phosphorylation of serine-rich, intrinsically disordered proteins in *C. elegans*. *Elife* **3**, e04591 (2014).

46. Smith J, Calidas D, Schmidt H, Lu T, Rasoloson D, Seydoux G. Spatial patterning of P granules by RNA-induced phase separation of the intrinsically-disordered protein MEG-3. *Elife* **5**, e21337 (2016).

47. Ling SH, Decker CJ, Walsh MA, She M, Parker R, Song H. Crystal structure of human

Edc3 and its functional implications. *Mol Cell Biol* **28**, 5965-5976 (2008).

48. Jeske M*, et al.* The crystal structure of the *Drosophila* germline inducer Oskar identifies

two domains with distinct Vasa helicase- and RNA-binding activities. *Cell Rep* **12**, 587-

598 (2015).

49. Nott TJ*, et al.* Phase transition of a disordered nuage protein generates environmentally

responsive membraneless organelles. *Mol Cell* **57**, 936-947 (2015).

50. Batista PJ*, et al.* PRG-1 and 21U-RNAs interact to form the piRNA complex required for

fertility in *C. elegans*. *Mol Cell* **31**, 67-78 (2008).

51. Tang W, Tu S, Lee HC, Weng Z, Mello CC. The RNase PARN-1 trims piRNA 3' ends to

promote transcriptome surveillance in *C. elegans*. *Cell* **164**, 974-984 (2016).

52. Berchowitz LE*, et al.* Regulated formation of an amyloid-like translational repressor

governs gametogenesis. *Cell* **163**, 406-418 (2015).

53. Hubstenberger A*, et al.* P-body purification reveals the condensation of repressed mRNA

regulons. *Mol Cell* **68**, 144-157 e145 (2017).

54. Wippich F, Bodenmiller B, Trajkovska MG, Wanka S, Aebersold R, Pelkmans L. Dual

specificity kinase DYRK3 couples stress granule condensation/dissolution to mTORC1

signaling. *Cell* **152**, 791-805 (2013).

55. Maji SK*, et al.* Functional amyloids as natural storage of peptide hormones in pituitary

secretory granules. *Science* **325**, 328-332 (2009).

56. Fowler DM, Koulov AV, Alory-Jost C, Marks MS, Balch WE, Kelly JW. Functional amyloid

formation within mammalian tissue. *PLoS Biol* **4**, e6 (2006).

57. Du M, Chen ZJ. DNA-induced liquid phase condensation of cGAS activates innate

immune signaling. *Science* **361**, 704-709 (2018).

58. Arnau J, Lauritzen C, Petersen GE, Pedersen J. Current strategies for the use of affinity

tags and tag removal for the purification of recombinant proteins. *Protein Expr Purif* **48**, 1-

13 (2006).

59. Gibson DG. Enzymatic assembly of overlapping DNA fragments. *Methods Enzymol* **498**,

349-361 (2011).

60. Brenner S. The genetics of *Caenorhabditis elegans*. *Genetics* **77**, 71-94 (1974).

61. Arribere JA, Bell RT, Fu BX, Artiles KL, Hartman PS, Fire AZ. Efficient marker-free

recovery of custom genetic modifications with CRISPR/Cas9 in *Caenorhabditis elegans*.

Genetics **198**, 837-846 (2014).

62. Paix A, Folkmann A, Rasoloson D, Seydoux G. High efficiency, homology-directed

genome editing in *Caenorhabditis elegans* using CRISPR-Cas9 ribonucleoprotein

complexes. *Genetics* **201**, 47-54 (2015).

63. Crittenden SL, Seidel HS, Kimble J. Analysis of the *C. elegans* germline stem cell pool.

Methods Mol Biol **1463**, 1-33 (2017).

64. Grimm JB*, et al.* A general method to improve fluorophores for live-cell and single-

molecule microscopy. *Nat Methods* **12**, 244-250, 243 p following 250 (2015).

65. Crittenden SL, Lee C, Mohanty I, Battula S, Knobel K, Kimble J. Sexual dimorphism of

niche architecture and regulation of the *C. elegans* germline stem cell pool. *Mol Biol Cell*

30, mbcE19030164 (2019).

66. Schindelin J, Rueden CT, Hiner MC, Eliceiri KW. The ImageJ ecosystem: An open

platform for biomedical image analysis. *Mol Reprod Dev* **82**, 518-529 (2015).

67. Magis C*, et al.* T-Coffee: Tree-based consistency objective function for alignment

evaluation. *Methods Mol Biol* **1079**, 117-129 (2014).

Acknowledgements

The authors thank M. Cox for equipment; J. Claycomb for plasmids and strains; L. Lavis for

supplying the SNAP JF 549 ligand; T. Hoang, S. Strome, D. Updike, and members of the

Kimble and Wickens labs for helpful discussions. Use of the LS-CAT Sector 21 was supported

by the Michigan Economic Development Corporation and the Michigan Technology Tri-Corridor

(Grant 085P1000817). The Keck Biophysics Facility (SEC-MALS) is supported in part by NCI

CCSG P30 CA060553 grant awarded to the Robert H Lurie Comprehensive Cancer Center.

STA was supported by K99HD081208. CAB was supported by NIH grants GM094584,

GM094622 and GM098248. MW was supported by NIH grant GM50942. JK is an Investigator of

the Howard Hughes Medical Institute.

Author contributions

STA conceived and performed experiments, analyzed data and wrote the paper. TRL and SLC

performed experiments, analyzed data and helped write the paper. CAB performed experiments

and analyzed data. MW analyzed data and helped write the paper. JK conceived experiments,

analyzed data and wrote the paper.

Competing interests: The authors declare no competing interests.

Figure Legends

Figure 1. PGL-tethering represses an mRNA reporter *in vivo*

(A) Left, *C. elegans* adult hermaphrodite possesses two gonadal arms with proliferating germ

cells at one end (asterisk) and differentiating gametes at the other. Gonads make sperm (blue)

first and then oocytes (pink). Right, P-granules (magenta) reside at the nuclear periphery of all

germ cells until late oogenesis. (B) Linear diagram of *C. elegans* PGL-1. (C) Protein-mRNA

tethering assay. The reporter mRNA encodes GFP-histone H2B and harbors three boxb

hairpins in its 3'UTR; a ubiquitous germline promoter drives expression (see Methods). λN22

peptide (light blue) is inserted into PGL-1 with a SNAP tag (magenta). Binding of PGL-

1::SNAP::λN22 to boxB hairpins recruits reporter mRNA. (D-G) GFP reporter expression in

germ cells of live animals. (D,E) Brightfield image. (F,G) GFP fluorescence (green); auto

fluorescence (red). n, number of animals scored for GFP expression. "%", germlines with

detectable GFP. Scale bar, 10 μm, in D applies to D-G. (H-K) Representative images in fixed

gonads. (H,I) GFP fluorescence. (J,K) SNAP staining (magenta) and DAPI (cyan). n, number of

germlines scored for GFP expression. Scale bar, 10 μm, in (H) applies to images. **Figure 1** and

Figure 5 results were performed in parallel, and thus results from (E,G) are the same reported

in **Figure 5B,D**.

Figure 2. Crystal structure of PGL NtDD

(A) Crystal structure of *C. japonica* PGL-1 NtDD to 1.5 Å. See **Table S1** for crystal structure

data and model statistics. NtDD has four copies per asymmetric unit (ASU). Copies in yellow,

gold, tan, and brown. Arrows indicate two pairs of subunit interfaces in the ASU. Red arrows

highlight the interface relying on conserved amino acids (see text). (B) Enlarged image of a

single NtDD.

Figure 3. NTD dimerization and its role in PGL self-assembly

(A) Structural model of the NtDD dimer. (B-C) Enlargement of dimer interface (red box in A).

PGL-1 amino acids (B) K126 and K129, and (C) R123 interact with apposing subunit side

chains. Residue labels in yellow or gold to indicate their representative subunits. (D,E) Size

exclusion chromatography and multi-angle light scattering (SEC-MALS) of recombinant PGL-3

(D,E) NtDD wild type, (D) K126E K129E and (E) R123E proteins. A280 UV absorbance (left y

axis) was normalized to the maximum value. Molecular weight (MW, right y axis) for MALS in

daltons (Da). Wild-type protein (blue) measured the approximate size of a dimer, while both

mutant proteins (red) measured approximately as monomers. (F) Diagram of *C. elegans* PGL-1

C-terminally tagged with GFP. (G-J) Representative images of (G) GFP-tagged PGL-1, (H) GFP

alone, and GFP-tagged PGL-1 (I) K126E K129E and (J) R123E mutants expressed in Chinese

Hamster Ovary (CHO) cells. Cell cultures were imaged live, and GFP-positive cells counted for

the presence or absence of granules. Images show the majority result (percentages noted

above image). Scale bar, 10 μm.

Figure 4. NtDD dimerization is critical for fertility and P-granule formation in nematodes

(A) Sites of SNAP tag insertion and missense mutations in *C. elegans* PGL-1. (B) Fertility of

SNAP-tagged PGL-1 animals. Percentages were obtained after scoring individuals for

production of larval progeny after 5 days at either 20°C or 25°C. (C-P) Extruded adult germlines,

fixed, stained and imaged in same region of meiotic pachytene (see **Figure S5A**). (C-F)

Representative images of SNAP staining to visualize PGL-1 expression and granule formation.

All images are partial z-stacks to maximize visualization of P-granules. Images were taken from

germlines containing embryos; similar images were obtained from germlines lacking embryos

(**Figure S5E,F**). (C) PGL-1::SNAP localizes to granules around nuclei (n=49). (D) Control, wild-

type animal lacking SNAP tag shows virtually no background staining (n=20). (E) PGL-1(K126E

K129E)::SNAP is diffuse (n=38). (F) PGL-1(R123E)::SNAP is diffuse (n=24). (G-P)

Representative images showing localization of three P-granule components in germ cells

expressing either (G-K) PGL-1::SNAP (n=20) or (L-P) PGL-1(K126E K129E)::SNAP (n=14).

(G,L) DNA (DAPI); (H,M) SNAP (PGL-1::SNAP or mutant); (I,N) V5 (PGL-3); (J,O) MYC (GLH-

1); (K,P) Merge. Scale bar, 10 µm for all images, except 2.5-fold enlargements of nuclei in

boxes placed outside main images.

Figure 5. PGL assembly is required for tethered mRNA reporter repression

(A) Protein-mRNA tethering assay and PGL assembly. To test the necessity of granule

formation for mRNA repression, NtDD assembly mutations were added to PGL-1::SNAP::λN22

and germlines observed for GFP reporter expression. (B-E) GFP reporter expression in germ

cells of live animals. (B,C) brightfield image. (D,E) GFP fluorescence (green); auto fluorescence

(red). n, number of animals scored for GFP expression. Scale bar, 10 µm, in B applies to B-E

images. (F-I) Representative images of PGL granule formation, seen by SNAP staining

(magenta) and GFP fluorescence (green) in fixed gonads. n, number of germlines scored for

GFP expression. "%", germlines with detectable GFP. Scale bar, 10 µm, in F applies to F-I

images. **Figure 1** and **Figure 5** results were performed in parallel, and thus results from B,D are

the same reported in **Figure 1E,G**.

Figure 6. PGL-1 and WAGO-1 assemble independently in P-granules

(A-D) Representative images showing localization of PGL-1 and WAGO-1 in germ cells

expressing (A) PGL-1::SNAP, WAGO-1::3xV5 (n=33). (B) PGL-1(R123E)::SNAP, WAGO-

1::3xV5 without WAGO-1 puncta (11 of 22 germlines). (C) PGL-1(R123E)::SNAP, WAGO-

1::3xV5 with WAGO-1 puncta (11 of 22 germlines). (D) PGL-1::SNAP, WAGO-1(null)::3xV5

(n=29). DNA (DAPI); SNAP (PGL-1::SNAP or mutant); V5 (WAGO-1::3xV5 wild-type or null).

Scale bar, 10 µm for all images, except 2.5-fold enlargements placed outside main images.

Figure 7. Argonaute WAGO-1 is required for PGL-tethered mRNA reporter repression

(A) Protein-mRNA tethering assay and WAGO-1. To test the necessity of WAGO-1 for mRNA

repression, PGL-1::SNAP::λN22 germlines were analyzed for GFP reporter expression in the

presence or absence of WAGO-1. (B-E) GFP reporter expression in germ cells of live animals

with (B,D) wild-type *wago-1* or (C,E) *wago-1* null. (B,C) Brightfield image. (D,E) GFP

fluorescence (green); auto fluorescence (red). n, number of animals scored for GFP expression.

Scale bar, 10 µm, in B applies to B-E images. (F-I) Representative images of fixed germlines

with (F,H) wild-type *wago-1* or (G,I) *wago-1* null. (F,G) GFP fluorescence. (H,I) DNA (DAPI) and

PGL granule formation seen by SNAP staining. n, number of germlines scored for GFP

expression. Scale bar, 10 µm, in F applies to images F-I.

Figure 8. Model of P-granule assembly and mRNA repression

(A) P-granules assemble at the nuclear pore with assembly-competent PGL protein. PGL is

shown as a dimer for simplicity but multivalent PGLs likely form an oligomeric protein-RNA

network. WAGO-1 binds to RNA, as expected for an Argonaute, and WAGO-associated RNAs

are repressed through mRNA turnover or translational repression (purple). (B) When PGL NtDD

cannot dimerize, PGL fails to assemble into P-granules at the nuclear periphery. WAGO-1

assembles independently into P-granules. PGL-associated, non-granular mRNAs are available

for translation (ribosomes, black). (C) In the absence of cytoplasmic Argonaute WAGO-1, PGL

proteins assemble into P-granules normally with its associated mRNA, but loss of WAGO-1

perturbs repression of P-granule-localized transcripts. PGL's liquid droplet properties permit

diffusion of some transcripts into the cytoplasm for translation. RNAs in the granule are not

translated, either because of repressors or the lack of ribosomes. See text for further

Discussion.

Supplemental Tables and Figures

Table S1. *C. japonica* PGL-1 NtDD crystal structure data and model statistics

Figure S1. PGL sequence alignment and the *pgl-1* locus

(A) *pgl-1* primary transcript. 5' and 3' UTRs are grey, exons are white, numbered 1-8 and

separated by introns. Sites of *pgl-1* mutations are labeled, including location of SNAP tag

(magenta) and λN22 fusion (blue). (B) Sequence alignment of PGL NtDD domain in *C. elegans*

(*Ce*), *C. japonica* (*Cj*), *C. brenneri* (*Cbn*), *C. briggsae* (*Cbr*), *C. remanei* (*Cr*). Alignment and

conservation (cons.) determined by T-Coffee [67]. Starred residues (*) are identical. Period (.) and

colon (:) residues are similar. Residues participating in salt bridges only are in orange. Residues

participating in hydrogen bonds only are in yellow. Residues forming both hydrogen bonds and

salt bridges are highlighted in red. *C. elegans* PGL-1 missense mutations and their allele

numbers are labeled. Dashed lines mark the newly-annotated end of PGL-1 NtDD domain and

start of PGL-1 CDD domain.

Figure S2. GFP reporter transcripts localize to P-granules when tethered

(A) Single molecule fluorescent *in situ* hybridization (smFISH) in nematode germ cells to

visualize RNA expression and localization. (B-M) Gonads were extruded from animals harboring

the *gfp* reporter and (B-E) PGL-1::SNAP (n=30); (F-I) PGL-1::SNAP::λN22 (n=27); (J-M) PGL-1

(K126E K129E)::SNAP::λN22 (n=10). Gonads were fixed and imaged for *gfp* RNA using (B,F,J)

smFISH; (C,G,K) GFP protein fluorescence; (D,H,L) DNA (DAPI) and SNAP. The smFISH, DNA

and SNAP images are merged in E,I,M. White arrows mark examples of intranuclear puncta;

black arrows mark examples of cytoplasmic puncta. Scale bar, 5 μm, for all images, except for

2.5-fold enlarged images in inset. For germline location, see **Figure S5A**. (N) Quantification of

GFP signal in confocal images for the GFP reporter expressed with PGL-1::SNAP (n=23), PGL-

1::SNAP::λN22 (n=24), PGL-1 (K126E K129E)::SNAP::λN22 (n=10). A silenced GFP reporter

(n=25) served as a negative control. Mean GFP fluorescent signal and standard deviation

reported in Arbitrary Units. (O) smFISH and PGL-1::SNAP signal colocalization in germlines

expressing the GFP reporter and either PGL-1::SNAP or PGL-1::SNAP::λN22. Reported as the

(Sum of smFISH GFP Intensity in SNAP:smFISH colocalized binned signal)/(Total smFISH GFP

binned signal). Box plots represent the first and third quartiles, black line is median, whiskers to

min and max values. Co-localization greater in germlines expressing PGL-1::SNAP::λN22

(p<0.01). See Methods for more details.

Figure S3. GFP reporter transcripts localize to P-granules when tethered

(A-F) Six additional examples of germlines harboring PGL-1::SNAP::λN22 (n=27). Gonads were

fixed and imaged for *gfp* RNA using smFISH, DNA (DAPI) and SNAP. The three are merged in

images on right. White arrows mark examples of intranuclear puncta; black arrows mark

examples of cytoplasmic puncta. Scale bar, 5 µm, for all images, except for 2.5-fold enlarged

images in inset. For germline location, see **Figure S5A**.

Figure S4. Supplemental biochemical and structural analyses of PGL NtDD

(A,B) Coomassie-stained polyacrylamide gel of recombinant PGL NtDD wild-type and mutant

protein. Ladder marker sizes labeled in kilodaltons (kDa) on right. (A) Recombinant *C. japonica*

(Cj) PGL-1 NtDD protein used for crystallization. Recombinant *C. elegans* PGL-3 NTD protein

included for comparison. (B) Wild-type and mutant *C. elegans* PGL-3 NtDD recombinant

proteins used for biochemical characterization. (C) Tables of predicted hydrogen bonds and salt

bridges at the NtDD dimerization interface. Amino acid numbers correspond to *C. japonica* PGL-

1 NtDD. (D,E) Surface representation of the NtDD dimerization interface. (D) Amino acids

colored by identity (red) and similarity (pink). (E) Amino acids at dimerization interface (purple).

Figure S5. Supplemental images of PGL-1 dimerization mutants

(A) Schematic of an adult hermaphrodite germline. An asterisk marks proliferating germ cells

here and in images B-D. The germline produces oocytes at this stage; sperm were made earlier

and stored in the spermatheca (not shown). Red box marks region imaged. (B-D)

Representative brightfield images of extruded gonads from worms grown at 20°C. Scale bar, 10

μm. (B) Wild-type PGL-1::SNAP gonads are of normal size and produce oocytes and embryos.

(C) Representative image of PGL-1(K126E K129E)::SNAP sterile gonads, which are small and

produce no gametes. (D) PGL-1(R123E)::SNAP sterile gonads are also small and produce no

gametes. (E-H) Representative partial z-projection stacks of SNAP and DNA stained germlines.

Scale bar, 10 μm, applies for all images. (E,F) PGL-1::SNAP mutant protein is expressed in

sterile gonads (no embryos observed). (E) PGL-1(K126E K129E)::SNAP, n=21 gonads imaged.

(F) PGL-1(R123E)::SNAP, n=30 gonads imaged. (G,H) In single rare gonads, PGL-1::SNAP

mutants were seen to assemble into granules in all germ cells.

1001

Figure S6. PGL-1 protein expression. (A-D) Examples of SNAP staining in (A) PGL-1::SNAP,

(B) wild-type N2 negative control, (C) PGL-1(K126E K129E)::SNAP and (D) PGL-

1(R123E)::SNAP. (E) Fluorescent signal (y-axis, arbitrary units) from SNAP-stained PGL-

1::SNAP, N2, PGL-1(K126E K129E)::SNAP and PGL-1(R123E)::SNAP germlines were

measured from the beginning of the germline (start of progenitor zone, **Figure S5A**) to the

meiotic pachytene region (x-axis, μM). Solid line represents the mean with black showing the

standard error. (F) Immunoblot of embryo-containing adult, N2, PGL-1::SNAP or PGL-

1(R123E)::SNAP expressing worms. Samples were separated by SDS-PAGE and probed with

SNAP and actin antibodies. Experiment was repeated twice with similar results.

1011

Figure S7. Targeted RNAi screen reveals WAGO-1 as a potential candidate in repressing PGL-

tethered mRNA transcripts

(A) Candidate gene RNAi screening strategy. PGL-1::SNAP::λN22 tethered to GFP reporter

mRNAs repress GFP expression. We propagated larval worms on bacteria expressing RNAi for

specific P-granule-associated enzymatic factors. After 5 days (one generation), young adult

worm progeny were observed for GFP nuclei in their germlines, indicating tethering de-

repression. (B) Candidate genes and their effect on GFP expression. Sequences of candidate

genomic regions reported. Results were repeated at least three times and summarized here. (C)

wago-1 (R06C7.1.1) primary transcript. 5' and 3' UTRs are grey, exons are white, numbered 1-9

and separated by introns. Site of 3xV5 insertion and deleted region of null allele noted.

1022

Figure S8. GFP reporter expression in the presence or absence of WAGO-1. (A-D) Examples

of GFP fluorescence in (A) positive control *mut-2(-)*; GFP reporter, (B) negative control wild-type

N2, (C) GFP reporter, and (D) *wago-1(-)*; GFP reporter. *mut-2* prevents reporter silencing. (E)

GFP Fluorescent signal (y-axis, arbitrary units) from germlines of described genotypes were

measured from the beginning of the germline (start of progenitor zone, **Figure S5A**) to the

meiotic pachytene region (x-axis, µM). The greater than 100% values are due to the use of both

Imaris and Matlab for quantitation (see Methods). Solid line represents the mean with black

showing the standard error. Note that the GFP reporter expressed similarly with or without

WAGO-1.

1032

Figure S9. PGL-1-tethered GFP reporter transcripts localize to granules in the presence or

absence of WAGO-1

Gonads were extruded from *gfp* reporter, PGL-1::SNAP::λN22 animals harboring (A-D) *wago-1*

wild-type (n=22); (E-H) *wago-1* null (n=21). Gonads were fixed and imaged for (A,E) *gfp* RNA

using smFISH, (B,F) GFP protein fluorescence, (C,G) DNA (DAPI) and SNAP. *gfp* RNA

smFISH, DNA and SNAP merged in D, H. White arrows mark examples of intranuclear puncta;

black arrows mark examples of cytoplasmic puncta. Scale bar, 5 µm, for all images, except for

2.5-fold enlarged images in inset. For germline location, see **Figure S5A**. (I) smFISH and PGL-

1::SNAP signal colocalization in germlines expressing the GFP reporter, PGL-1::SNAP::λN22 in

the presence or absence of WAGO-1. Reported as the (Sum of smFISH GFP Intensity in

SNAP:smFISH colocalized binned signal)/(Total smFISH GFP binned signal). Box plots

represent the first and third quartiles, black line is median, whiskers to min and max values. Co-

localization similar in germlines expressing PGL-1::SNAP::λN22 with or without WAGO-1

(p=0.039). See Methods for more details.

1047

Figure S10. Additional images of PGL-1-tethered GFP reporter transcripts without WAGO-1

(A-F) Six additional examples of germlines harboring PGL-1::SNAP::λN22, *wago-1* null imaged

for *gfp* RNA, DNA, and SNAP. *gfp* RNA, DNA and SNAP merged in right images. White arrows

mark examples of intranuclear puncta; black arrows mark examples of cytoplasmic puncta.

Scale bar, 5 μm, for all images, except for 2.5-fold enlarged images in inset. For germline

location, see **Figure S5A**.

A

B

N-term. Dimerization Domain (NtDD)	Central Dimerization Domain (CDD)	C-region	RGG

1   217 223   449   664   730

C PGL-1::SNAP::λN22

SNAP λN22

Reporter

GFP histone 3xboxb AAA → expression?
(GFP fluorescence in nuclei)

PGL-1::SNAP | PGL-1::SNAP::λN22/ +

D germline / intestine | **E** germline / intestine

F GFP auto 100% GFP (n=219) | **G** GFP auto 0% GFP (n=111)

H GFP 100% GFP n=35 | **I** GFP 5% GFP n=39

J DNA SNAP | **K** DNA SNAP

A residues at dimerization interface
*** NtDD | CDD | C-region | SNAP | RGG

B

Protein	allele	20°C		25°C	
		% fertile	n	% fertile	n
PGL-1(wt)	+	99	96	98	93
PGL-1(wt)::SNAP	q894	99	93	91	93
PGL-1 null	bn102	96	92	18	96
PGL-1(K126E K129E)::SNAP	q960	28	93	0	92
PGL-1(R123E)::SNAP	q975	78	95	3	96

PGL-1(wt)::SNAP | PGL-1(wt) (no SNAP) | PGL-1(K126E K129E):: SNAP | PGL-1(R123E):: SNAP

C SNAP / DNA / DNA SNAP
D SNAP / DNA / DNA SNAP
E SNAP / DNA / DNA SNAP
F SNAP / DNA / DNA SNAP

PGL-1(wt)::SNAP

G DNA
H SNAP (PGL-1)
I V5 (PGL-3)
J MYC (GLH-1)
K SNAP V5 MYC

PGL-1(K126E K129E)::SNAP

L DNA
M SNAP (PGL-1)
N V5 (PGL-3)
O MYC (GLH-1)
P SNAP V5 MYC

A PGL-1(dimerization defective)::SNAP::λN22

K126E K129E   λN22

Reporter

GFP histone 3xboxb —AAA → GFP?

PGL-1::SNAP::λN22/ +   PGL-1(K126E K129E):: SNAP::λN22

B germline intestine   C germline intestine

D GFP auto   0% GFP (n=111)   E GFP auto   98% GFP (n=95)

F GFP   5% GFP (n=39)   G GFP   100% GFP (n=22)

H DNA SNAP   I DNA SNAP

A PGL-1::SNAP, WAGO-1::3xV5 (n=33)

B PGL-1(R123E)::SNAP , WAGO-1::3xV5 (50%, n=22)

C PGL-1(R123E)::SNAP , WAGO-1::3xV5 (50%, n=22)

D PGL-1::SNAP, WAGO-1(null)::3xV5 (n=29)

Table S1. *C. japonica* NtDD data collection and refinement statistics.

	NtDD, Selenomethionine (5W4D)	NtDD, wild-type (5W4A)
Wavelength	0.9786	0.984
Resolution range	48.47 - 1.599 (1.656 - 1.599)	30.36 - 1.5 (1.554 - 1.5)
Space group	C 1 2 1	C 1 2 1
Unit cell	133.3 94.8 72.5 90 91.4 90	132.77 94.67 72.95 90 90.756 90
Total reflections	880332 (81451)	2176741 (194944)
Unique reflections	115340 (11239)	143729 (14301)
Multiplicity	7.6 (7.2)	15.1 (13.6)
Completeness (%)	97.06 (95.16)	99.77 (99.33)
Mean I/sigma(I)	24.44 (2.30)	16.91 (1.85)
Wilson B-factor	22.04	22.44
R-merge	0.0469 (0.8775)	0.08108 (1.282)
R-meas	0.05038 (0.945)	0.08353 (1.332)
R-pim	0.01827 (0.348)	0.01984 (0.357)
CC1/2	0.999 (0.74)	0.997 (0.662)
CC*	1 (0.922)	0.999 (0.893)
Reflections used in refinement	115302 (11238)	143649 (14297)
Reflections used for R-free	1424 (129)	1468 (151)
R-work	0.1607 (0.2590)	0.1681 (0.3038)
R-free	0.1925 (0.2707)	0.2039 (0.3232)
CC(work)	0.962 (0.864)	0.967 (0.802)
CC(free)	0.939 (0.819)	0.974 (0.794)
Number of non-	7593	7785

hydrogen atoms		
macromolecules	6768	6813
ligands	168	136
solvent	657	836
Protein residues	846	853
RMS(bonds)	0.010	0.010
RMS(angles)	1.00	0.99
Ramachandran favored (%)	98.68	98.10
Ramachandran allowed (%)	1.32	1.90
Ramachandran outliers (%)	0.00	0.00
Rotamer outliers (%)	1.33	0.66
Clashscore	2.29	1.86
Average B-factor	30.58	30.21
macromolecules	29.29	28.97
ligands	53.64	51.75
solvent	37.95	36.82
Number of TLS groups	1	1

Statistics for the highest-resolution shell are shown in parentheses.

A Germ cell

smFISH — nascent transcripts — AAA

smFISH — mature mRNAs — AAA

nucleus

cytoplasm

→ RNA localization?

PGL-1::SNAP

B smFISH to reporter RNA

C GFP

D DNA SNAP

E DNA SNAP smFISH

PGL-1::SNAP::λN22/ +

F smFISH to reporter RNA

G GFP

H DNA SNAP

I DNA SNAP smFISH

PGL-1::SNAP::λN22 K126E K129E

J smFISH to reporter RNA

K GFP

L DNA SNAP

M DNA SNAP smFISH

N

GFP fluorescence Arbitrary Units

- PGL-1::SNAP
- neg
- PGL-1::SNAP::λN22
- PGL-1(K126E K129E):: SNAP::λN22

O

smFISH-SNAP overlap (%)

PGL-1::SNAP

PGL-1::SNAP::λN22

PGL-1::SNAP::λN22/ +, six additional germlines

C

H-bonds		salt bridges	
subunit 1	subunit 2	subunit 1	subunit 2
A:R128	B:T158	A:R128	B:E189
A:R128	B:E189	A:K131	B:E162
A:K131	B:E162	A:K134	B:D167
A:L132	B:N196	A:K134	B:D169
A:K134	B:D167	A:E162	B:K131
A:K134	B:D169	A:D167	B:K134
A:T158	B:R128	A:D169	B:K134
A:E162	B:K131	A:E189	B:R128
A:D167	B:K134		
A:D169	B:K134		
A:E189	B:R128		
A:N196	B:L132		

A

progenitor zone

meiotic pachytene

maturing oocytes

cross section

B PGL-1::SNAP

newly fertilized embryo

maturing oocytes

C PGL-1::SNAP K126E K129E

gut

severely underproliferated germline with no oocytes or embryos

D PGL-1::SNAP R123E

gut

gut

smaller than normal germline with no oocytes or embryos

PGL-1::SNAP staining in most gonads (sterile)

PGL-1::SNAP granules in rare gonads

E K126E K129E

SNAP
DNA
DNA SNAP

F R123E

SNAP
DNA
DNA SNAP

G K126E K129E

SNAP
DNA
DNA SNAP

H R123E

SNAP
DNA
DNA SNAP

A PGL-1::SNAP

DNA

SNAP

B wild-type (N2)

DNA

SNAP

C PGL-1::SNAP K126E K129E

DNA

SNAP

D PGL-1::SNAP R123E

DNA

SNAP

E

— wild-type (N2)
— PGL-1::SNAP
— PGL-1::SNAP K126E K129E
— PGL-1::SNAP R123E

F

A

PGL-1::SNAP::λN22   λN22

Reporter

● GFP histone 3xboxb ... AAA

RNAi against P-granule associated,
RNA regulatory genes

↓

expression?
(GFP fluorescence in nuclei)

B

Gene	Sequence	Genome region (start...end)	GFP %	n
empty	–	–	0	204
csr-1	F20D12.1	CAGTACGCATTTCCTCGTGA...TTTACAGTCTTGCGTGCTGG	0	46
drh-3	D2005.5	CGCTCAACGTGTCAGTGATT...TCAAATGCTTCCATTGCAAG	0	59
ego-1	F26A3.3	ATCCGATCCGAACCATTACA...GGCATTTGGGATGAAAATTG	2.0	49
ekl-1	F22D6.6	ATGCAAAGCCAATTCCAGTC...GATTCAGGCGGTCGTGATAT	0	64
mut-2	K04F10.6	GCCTGGCACCAATGTAGTTT...CGTCCTCGTGATCCCTTAAA	0	70
parn-1	K10C8.1	TTCCGCCCAAATATTTCAAG...ATCAAGTCTTCATGGACGGC	0	46
pgl-1	ZK381.4	AATTTGTTCAAGGAATCAAC...GGAGAAGGTGTTACTGTAAAAGCG	0	64
prg-1	D2030.6	AGTGTGCTTTGCGATCTCCT...GACCACTTCCAGATGCCATT	1.4	69
rde-1	K08H10.7	CGACATCTGTTTCAGCAGGA...GATTGCGCTCCTCTGTTTTC	1.9	54
wago-1	**R06C7.1**	GGGAAGAGATTGACGTTGGA...AATCGGCTACAAACAAACCG	**43.6**	**94**
wago-4	F58G1.1	CATTGACGGTTGGTATTCAAAAT...TCAACAAGACCAATGTGAAGTTG	1.4	71

C *wago-1* (R06C7.1.1)

q1089 (**deletion and frameshift**)

q1087 (**3xV5**)

1 2 3 4 5 6 7 8 9

500 bp

A *mut-2 (-)*; GFP reporter

DNA

GFP

B wild-type (N2)

DNA

GFP

C GFP reporter

DNA

GFP

D *wago-1 (-)*; GFP reporter

DNA

GFP

E

— wild-type (N2)
— *mut-2 (-)*; GFP reporter
— GFP reporter
— *wago-1 (-)*; GFP reporter

μM

PGL-1::SNAP::λN22, WAGO-1 wild type

PGL-1::SNAP::λN22, WAGO-1 null

I

PGL-1::SNAP::λN22, WAGO-1 null, six additional germlines