Title: The genetics of the mood disorder spectrum: genome-wide association

analyses of over 185,000 cases and 439,000 controls

Authors: Jonathan R. I. Coleman[1;2], Héléna A. Gaspar[1;2], Julien Bryois[3], Bipolar

Disorder Working Group of the Psychiatric Genomics Consortium[4], Major Depressive

Disorder Working Group of the Psychiatric Genomics Consortium[4], Gerome Breen[1;2]

Affiliations:

1. Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry,

8   Psychology and Neuroscience, King's College London, London, United Kingdom

2. NIHR Maudsley Biomedical Research Centre, King's College London,

London, United Kingdom

3. Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,

Stockholm, Sweden

4. Full consortium authorship listed in Article Information

Correspondence to: Gerome Breen, Social, Genetic and Developmental Psychiatry

Centre, Institute of Psychiatry, Psychology and Neuroscience, London, SE5 8AF,

UK, gerome.breen@kcl.ac.uk, +442078480409

Short title: GWAS of the mood disorder spectrum

Keywords: major depressive disorder; bipolar disorder; mood disorders; affective

disorders; genome-wide association study; genetic correlation

Abstract: 240 words

Main text: 4272 words (including citations and headers)

Tables: 1

Figures: 6

Supplementary Materials: 2 (Materials and Figures, Tables)

Abstract

Background

Mood disorders (including major depressive disorder and bipolar disorder) affect 10-20% of the population. They range from brief, mild episodes to severe, incapacitating conditions that markedly impact lives. Despite their diagnostic distinction, multiple approaches have shown considerable sharing of risk factors across the mood disorders.

Methods

To clarify their shared molecular genetic basis, and to highlight disorder-specific associations, we meta-analysed data from the latest Psychiatric Genomics Consortium (PGC) genome-wide association studies of major depression (including data from 23andMe) and bipolar disorder, and an additional major depressive disorder cohort from UK Biobank (total: 185,285 cases, 439,741 controls; non-overlapping N = 609,424).

Results

Seventy-three loci reached genome-wide significance in the meta-analysis, including 15 that are novel for mood disorders. More genome-wide significant loci from the PGC analysis of major depression than bipolar disorder reached genome-wide significance. Genetic correlations revealed that type 2 bipolar disorder correlates strongly with recurrent and single episode major depressive disorder. Systems biology analyses highlight both similarities and differences between the mood disorders, particularly in the mouse brain cell types implicated by the expression patterns of associated genes. The mood disorders also differ in their genetic

correlation with educational attainment – positive in bipolar disorder but negative in

major depressive disorder.

<u>Conclusions</u>

The mood disorders share several genetic associations, and can be combined

effectively to increase variant discovery. However, we demonstrate several

differences between these disorders. Analysing subtypes of major depressive

disorder and bipolar disorder provides evidence for a genetic mood disorders

spectrum.

<u>Introduction</u>

Mood disorders affect 10-20% of the global population across their lifetime,

ranging from brief, mild episodes to severe, incapacitating conditions that markedly

impact lives (1–4). Major depressive disorder and bipolar disorder are the most

common forms and have been grouped together since the third edition of the

Diagnostic and Statistical Manual of Mental Disorders (DSM-III) (5). Although given

dedicated chapters in DSM5, they remain grouped as mood disorders in the

International Classification of Disorders (ICD11) (6, 7).

Depressive episodes are common to major depressive disorder and type 2

bipolar disorder, and are usually present in type 1 bipolar disorder (7). The bipolar

disorders are distinguished from major depressive disorder by the presence of mania

in type 1 and hypomania in type 2 (7). However, these distinctions are not absolute –

some individuals with major depressive disorder may later develop bipolar disorder,

and some endorse (hypo)manic symptoms (8–10). Following their first depressive

episode, a non-remitting individual might develop recurrent major depressive

disorder or bipolar disorder. Treatment guidelines for these disorders differ (11, 12).

Identifying shared and distinct genetic associations for major depressive disorder

and bipolar disorder could aid our understanding of these diagnostic trajectories.

Twin studies suggest that 35-45% of variance in risk for major depressive

disorder and 65-70% of the variance in bipolar disorder risk is accounted for by

additive genetic factors (13). These genetic components are partially shared, with a

twin genetic correlation (r_g) of ~65%, and common variant based r_g derived from the

results of genome-wide association studies (GWAS) of 30-35% (14–17).

Considerable progress has been made in identifying specific genetic variants that

underlie genetic risk. Recently, the Psychiatric Genomics Consortium (PGC)

published a GWAS of bipolar disorder, including over 20,000 cases, with 30 genomic

loci reaching genome-wide significance (16). They also performed a GWAS of major

depression, including over 135,000 individuals with major depressive disorder and

other definitions of depression, with 44 loci reaching genome-wide significance (15).

The PGC GWAS of major depression has since been combined with a broad

depression GWAS (Supplementary Note).

GWAS have identified statistical associations with major depressive disorder

and with bipolar disorder individually, but have not explored the genetic aspects of

the relationship between these disorders. In addition, both major depressive disorder

and bipolar disorder exhibit considerable clinical heterogeneity and can be separated

into subtypes. For example, the DSM5 includes categories for bipolar disorder type 1

and type 2, and for single episode and recurrent major depressive disorder (7). We

use the PGC analyses of major depression and bipolar disorder, along with analyses

of formally-defined major depressive disorder from UK Biobank, to explore two aims

(18, 19). Firstly, we seek to identify shared and distinct mood disorder genetics by

combining studies of major depressive disorder and bipolar disorder. We then

explore the genetic relationship of mood disorders to traits from the wider GWAS

literature. Secondly, we assess the overall genetic similarities and differences of

bipolar disorder subtypes (from the PGC) and major depressive disorder subtypes

(from UK Biobank), through comparing their genetic correlations and polygenic risk

scores from GWAS.

<u>Materials and Methods</u>

<u>Participants</u>

Our primary aim was to combine analyses of bipolar disorder and major

depression to examine the shared and distinct genetics of these disorders. Summary

statistics were derived from participants of Western European ancestries. Full

descriptions of each study and their composite cohorts are provided in each paper

(15, 16, 19). Brief descriptions are provided in the Supplementary Methods. Except

where otherwise specified, summary statistics are available (or will be made

available) at https://www.med.unc.edu/pgc/results-and-downloads.

Major depression data were drawn from the full cohort (PGC MDD: 135,458

cases, 344,901 controls) from (15). This included data from 23andMe (20), access to

which requires a Data Transfer Agreement; consequently, the data analysed here

differ from the summary statistics available at the link above. Data for bipolar

disorder were drawn from the discovery analysis previously reported (PGC BD:

20,352 cases, 31,358 controls), not including replication results (16).

Secondly, we wished to examine genetic correlations between mood disorder

subtypes. Summary statistics were available for the primary bipolar disorder

subtypes, type 1 bipolar disorder (BD1: 14,879 cases, 30,992 controls) and type 2

bipolar disorder (BD2: 3,421 cases, 22,155 controls), and for schizoaffective bipolar

disorder (SAB: 977 cases, 8,690 controls), a mood disorder including psychotic

symptoms. Controls are shared across these subtype analyses.

Subtype GWAS are not yet available from PGC MDD. As such, a major

depressive disorder cohort was derived from the online mental health questionnaire

in the UK Biobank (UKB MDD: 29,475 cases, 63,482 controls; Resource 22 on

http://biobank.ctsu.ox.ac.uk) (18). The definition of major depressive disorder in this

cohort is based on DSM-5, as described in full elsewhere (18), and in Supplementary

Table 1 (7). We defined three major depressive disorder subtypes for analysis.

Individuals meeting criteria for major depressive disorder were classed as recurrent

cases if they reported multiple depressed periods across their lifetime (rMDD, N =

17,451), and single-episode cases otherwise (sMDD, N = 12,024, Supplementary

Table 1). Individuals reporting depressive symptoms, but not meeting case criteria,

were excluded from the main analysis but used as a "sub-threshold depression"

subtype to examine the continuity of genetic associations with major depressive

disorder below clinical thresholds (subMDD, N = 21,596). All subtypes were

analysed with the full set of controls. Details on the quality control and analysis of the

UK Biobank phenotypes is provided in the Supplementary Methods.

Meta-analysis of GWAS data

We meta-analysed PGC MDD and UKB MDD to obtain a single major

depressive disorder GWAS (combined MDD). We meta-analysed combined MDD

with PGC BD, comparing mood disorder cases to controls (MOOD). Further meta-

analyses were performed between PGC MDD and each bipolar disorder subtype and

major depressive disorder subtype to assess the relative increase in variant

discovery when adding different mood disorder definitions to PGC MDD

(Supplementary Results).

Summary statistics were limited to common variants (MAF > 0.05;

Supplementary Methods) either genotyped or imputed with high confidence (INFO

score > 0.6) in all studies. Controls were shared between PGC MDD and PGC BD,

and (due to the inclusion of summary data in PGC MDD) the extent of this overlap

was unknown. Meta-analyses were therefore performed in METACARPA, which

controls for sample overlap of unknown extent between studies using the variance-

covariance matrix of the observed effect sizes at each variant, weighted by the

sample sizes (21, 22). METACARPA adjusted adequately for known overlap

between cohorts (Supplementary Methods). For later analyses (particularly linkage

disequilibrium score regression) we used as the sample size a "non-overlapping N"

estimated for each meta-analysis (Supplementary Methods). The definition,

annotation and visualisation of each meta-analysis is described in the

Supplementary Materials.

Sensitivity analysis using down-sampled PGC MDD

Results from MOOD showed greater similarity to PGC MDD than to PGC BD.

Cross-trait meta-analyses may be biased if the power of the composite analyses

differs substantially (23, 24). The mean chi-square of combined MDD [1.7] exceeded

that of PGC BD [1.39], suggesting this bias may affect our results (Supplementary

Table 2). We therefore repeated our analyses, meta-analysing UKB MDD with

summary statistics for PGC MDD that did not include participants from 23andMe nor

the UK Biobank (mean chi-square = 1.35). All analyses were performed on the full

and the down-sampled analyses, with the exception of GSMR analyses. Full results

of the down-sampled analyses are described in the Supplementary Materials.

Estimation of SNP-based heritability captured by common variants and genetic

correlations with published GWAS

The SNP-based heritability captured by common variants was assessed using

linkage disequilibrium score regression (LDSC) for each meta-analysed set of data

(25). SNP-based heritability estimates were transformed to the liability scale,

assuming population prevalences of 15% for combined MDD, 1% for PGC BD, and

16% for MOOD, and lower and upper bounds of these prevalences for comparison

(Supplementary Methods). LDSC separates genome-wide inflation into components

due to polygenicity and confounding (25). Inflation not due to polygenicity was

quantified as (intercept-1)/(mean observed chi-square-1) (26). Genetic correlations

were calculated in LDSC between each analysis and 414 traits curated from

published GWAS. Local estimates of SNP-based heritability and genetic covariance

were obtained using HESS v0.5.3b (Supplementary Methods and Results) (27, 28).

Genetic correlations between subtype analyses

To assess the structure of genetic correlations within the mood disorders,

SNP-based heritabilities and genetic correlations were calculated in LDSC between

bipolar disorder subtypes (BD1, BD2, SAB), and major depressive disorder subtypes

(rMDD, sMDD, subMDD). Putative differences between genetic correlations were

identified using a z-test (p < 0.05), and formally tested by applying a block-jackknife,

with Bonferroni correction for significance (p < 8.3×10^{-4}; Supplementary Methods).

Differences between the genetic correlations of PGC MDD and each bipolar disorder

subtype, and between PGC BD and each major depressive disorder subtype were

also tested (Bonferroni correction for significance, p < 0.0083). Genetic correlations

were hierarchically clustered using the gplots package in R v1.4.1 (29, 30).

Hierarchical clustering was performed using just the subtypes, and including results

from six external GWAS relevant to mood disorders (Supplementary Methods). To

validate our conclusion of a genetic mood disorder spectrum, we performed principal

component analysis of the genetic correlation matrix including the six external

GWAS (Supplementary Methods and Results).

Association of PGC BD polygenic risk scores with major depressive disorder

subtypes

Polygenic risk score analyses were performed using PRSice2 to assess

whether rMDD was genetically more similar to PGC BD than were sMDD or subMDD

(Supplementary Methods) (36).

Gene-wise, gene-set, and tissue and single-cell enrichment analyses

For all analyses, gene-wise p-values were calculated as the aggregate of the

mean and smallest p-value of SNPs annotated to Ensembl gene locations using

MAGMA v1.06 (Supplementary Methods and Results) (37). Gene set analysis was

performed in MAGMA (Supplementary Methods and Results). Further analyses were

performed to assess the enrichment of associated genes with expression-specificity

profiles from tissues (Genotype-Tissue Expression project, version 7) and broadly-

defined ("level 1") and narrowly-defined ("level 2") mouse brain cell-types (38, 39).

Analyses were performed in MAGMA following previously described methods with

minor modifications, with Bonferroni-correction for significance (Supplementary

Methods) (38). Similar analyses can be performed in LDSC-SEG – we report

MAGMA results, which reflect specificity of expression across the range, whereas

LDSC-SEG compares the top 10% of the range with the remainder (40). Results

using LDSC are included in the Supplementary Tables.

Mendelian randomisation (GSMR)

Bidirectional Mendelian randomisation analyses were performed using the

GSMR option in GCTA to allow exploratory inference of the causal direction of

known relationships between mood disorder traits and other traits (41, 42).

Specifically, the relationship between the mood disorder analyses (MOOD, combined

MDD, PGC BD) and schizophrenia, intelligence, educational attainment, body mass

index, and coronary artery disease were explored (Supplementary Methods) (32,

43–46). These traits were previously examined in the PGC major depression GWAS

– we additionally tested intelligence following the results of our genetic correlation

analyses (15).

Conditional and reversed-effect analyses

Additional analyses were performed to identify shared and distinct mood

disorder loci, using mtCOJO, an extension of GSMR (Supplementary Methods) (41,

42). Analyses were performed on combined MDD conditional on PGC BD, and on

PGC BD conditional on combined MDD (Supplementary Results). To identify loci

with opposite directions of effect between combined MDD and PGC BD, the MOOD

meta-analysis was repeated with reversed direction of effects for PGC BD

(Supplementary Methods and Results).

Results

Evidence for confounding in meta-analyses

Meta-analysis results were assessed for genome-wide inflation of test

statistics using LDSC (25). The LDSC intercept was significantly >1 in most cases

(1.00-1.06), which has previously been interpreted as confounding (Supplementary

Table 2). However, such inflation can occur in large cohorts without confounding

(47). Estimates of inflation not due to polygenicity were small in all meta-analyses (4-

7%, Supplementary Table 2).

Combined MOOD meta-analysis

We meta-analysed the PGC MDD, PGC BD and the UKB MDD cohorts

(MOOD, cases = 185,285, controls = 439,741, non-overlapping N = 609,424). 73 loci

reached genome-wide significance, of which 55 were also seen in the meta-analysis

of PGC MDD and UKB MDD (combined MDD, Supplementary Table 3,

Supplementary Figures 1 and 2). Results are summarised in Table 1: 39 of the 44

PGC MDD loci reached genome-wide significance in MOOD (Supplementary Table

3, Supplementary Figures 1-8). In comparison, only four of the 19 PGC BD loci

reached genome-wide significance in MOOD. MOOD loci overlapped considerably

with previous studies of depression and depressive symptoms (51/73) (20, 23, 48–

52), bipolar disorder (3/73) (53–56), neuroticism (32/73) (23, 57–59), and

schizophrenia (15/73) (32, 60), although participants overlap between MOOD and

many of these studies. Locus 52 (chromosome 12) passed genome-wide

significance in a previous meta-analysis of broad depression and bipolar disorder,

although the two other loci from this study did not replicate (51). Six of the 73

associations are entirely novel (p > 5x10^{-8} in previous studies of all phenotypes;

Table 1, Supplementary Table 4).

The down-sampled MOOD (cases = 95,481, controls = 287,932, non-

overlapping N = 280,214) showed increased similarity to PGC BD compared to

MOOD, but remained more similar to PGC MDD. Nineteen loci reached genome-

wide significance in down-sampled MOOD, including nine (20%) from PGC MDD,

compared with two (11%) reported in PGC BD (Supplementary Table 3). 17/19 loci

were also observed in MOOD. Of the two loci not observed in MOOD, one passed

genome-wide significance in PGC BD.

Locus	Chr	BP	Index SNP	A1	A2	OR	SE	p	Previous report
1	1	37192741	rs1002656	T	C	0.97	0.005	2.71×10^{-11}	DO, N
2	1	72837239	rs7531118	T	C	0.96	0.004	1.05×10^{-16}	D, DO, S, O
4	1	80795989	rs6667297	A	G	0.97	0.005	5.86×10^{-11}	D, DO
5	1	90796053	rs4261101	A	G	0.97	0.005	1.78×10^{-8}	D
6	1	175913828	rs10913112	T	C	0.97	0.005	1.46×10^{-10}	DO, O
7	1	177370033	rs16851203	T	C	0.96	0.007	2.38×10^{-9}	DO, S, O
9	2	22582968	rs61533748	T	C	0.97	0.004	3.84×10^{-11}	DO, N
10	2	57987593	rs11682175	T	C	0.97	0.004	2.18×10^{-11}	D, DO, BS, N, S, O
11	2	157111313	rs1226412	T	C	1.03	0.005	1.27×10^{-8}	D, DO, N, O
12	2	198807015	rs1518367	A	T	0.97	0.005	1.18×10^{-8}	BS, S, O
13	3	108148557	rs1531188	T	C	0.96	0.006	1.61×10^{-9}	O
14	3	158107180	rs7430565	A	G	0.97	0.004	2.30×10^{-11}	D, DO, N, O
16	4	42047778	rs34215985	C	G	0.97	0.006	1.72×10^{-10}	D, DO, N
17	5	77709430	rs4529173	T	C	0.97	0.005	4.29×10^{-9}	O
18	5	88002653	rs447801	T	C	1.03	0.004	2.29×10^{-10}	D, DO, N, O
19	5	92995013	rs71639293	A	G	1.03	0.005	5.85×10^{-9}	DO, N
20	5	103904226	rs12658032	A	G	1.04	0.005	2.19×10^{-16}	D, DO, N, O
21	5	106603482	rs55993664	A	C	0.97	0.006	1.87×10^{-8}	NOVEL LOCUS
22	5	124251883	rs116755193	T	C	0.97	0.005	1.47×10^{-10}	D, O
23	5	164523472	rs11135349	A	C	0.97	0.004	2.96×10^{-11}	D, DO, N
24	5	166992078	rs4869056	A	G	0.97	0.005	5.21×10^{-9}	D
25	6	28673998	rs145410455	A	G	0.94	0.007	7.17×10^{-18}	D, DO, BO, BS, DS, N, S, O
26	6	101339400	rs7771570	T	C	0.97	0.004	9.68×10^{-10}	DO, N, O
27	6	105365891	rs1933802	C	G	0.98	0.004	1.05×10^{-8}	DO, S, O
28	7	12267221	rs4721057	A	G	0.97	0.004	7.31×10^{-11}	D, DO, N, O
29	7	24826589	rs79879286	C	G	1.04	0.006	1.97×10^{-11}	B, BS, DO, S
30	7	82514089	rs34866621	T	C	1.03	0.005	2.21×10^{-8}	DO, O
31	7	109099919	rs58104186	A	G	1.03	0.004	7.12×10^{-9}	D, DO

34	9	11379630	rs10959753	T	C	0.96	0.005	1.45×10^{-13}	D, DO, N, O
35	9	37207269	rs4526442	T	C	0.96	0.006	7.97×10^{-11}	DO, O
36	9	81413414	rs11137850	A	G	1.03	0.005	1.25×10^{-8}	NOVEL LOCUS
38	9	119733380	rs10759881	A	C	1.03	0.005	8.56×10^{-9}	D, DO
40	9	122664468	rs10818400	T	G	0.98	0.004	1.29×10^{-8}	N
41	9	126682068	rs7029033	T	C	1.04	0.008	2.61×10^{-8}	D, DO, O
42	10	104684544	rs78821730	A	G	0.96	0.007	2.95×10^{-8}	N, BS, S, O
43	10	106563924	rs61867293	T	C	0.96	0.005	5.64×10^{-12}	D, DO, N, O
44	11	16293680	rs977509	T	C	0.97	0.005	1.19×10^{-8}	DO, N, O
45	11	31850105	rs1806153	T	G	1.03	0.005	2.81×10^{-9}	D, DO, N, O
46	11	32765866	rs143864773	T	C	1.04	0.008	1.70×10^{-8}	NOVEL LOCUS
47	11	61557803	rs102275	T	C	0.97	0.005	5.04×10^{-11}	B, DO, BO, O
48	11	63632673	rs10792422	T	G	0.98	0.004	2.18×10^{-8}	O
49	11	88743208	rs4753209	A	T	0.97	0.004	4.15×10^{-9}	DO, N, O
50	11	99268617	rs1504721	A	C	0.98	0.004	2.24×10^{-8}	O
51	11	113392994	rs2514218	T	C	0.97	0.005	3.22×10^{-10}	DO, BS, N, S, O
52	12	2344644	rs769087	A	G	1.03	0.005	3.27×10^{-8}	B, BD, BO, DS, BS, S, O
53	12	23947737	rs4074723	A	C	0.97	0.004	3.18×10^{-9}	D, DO, N, O
54	12	121186246	rs58235352	A	G	0.95	0.009	1.64×10^{-10}	DO, O
55	12	121907336	rs7962128	A	G	1.02	0.004	3.63×10^{-8}	NOVEL LOCUS
56	13	44327799	rs4143229	A	C	0.95	0.008	2.73×10^{-10}	D
57	13	53625781	rs12552	A	G	1.04	0.004	1.25×10^{-23}	D, DO, O
58	14	42074726	rs61990288	A	G	0.97	0.004	2.29×10^{-10}	D, DO, O
60	14	64686207	rs915057	A	G	0.98	0.004	1.92×10^{-8}	D, DO, O
61	14	75130235	rs1045430	T	G	0.97	0.004	9.83×10^{-11}	D, DO, N, O
62	14	104017953	rs10149470	A	G	0.97	0.004	1.15×10^{-10}	D, DS, DO, BS, S, O
63	15	36355868	rs1828385	A	C	0.97	0.004	1.15×10^{-8}	NOVEL LOCUS
64	15	37643831	rs8037355	T	C	0.97	0.004	4.09×10^{-15}	D, DO, O
65	16	6310645	rs8063603	A	G	0.97	0.005	5.36×10^{-11}	D, DO

66	16	7667332	rs11077206	C	G	1.03	0.004	5.49×10^{-10}	D, DO, N, O
67	16	13038723	rs12935276	T	G	0.97	0.005	4.75×10^{-10}	D, DO, N, O
68	16	13750257	rs7403810	T	G	1.03	0.005	7.52×10^{-11}	DO, BS, S, O
69	16	72214276	rs11643192	A	C	1.03	0.004	1.46×10^{-11}	D, O
70	17	27363750	rs75581564	A	G	1.04	0.006	2.47×10^{-10}	D, DO, O
71	18	31349072	rs4534926	C	G	1.03	0.004	9.14×10^{-9}	DO, N
72	18	36883737	rs62099069	A	T	0.97	0.004	9.52×10^{-10}	D, O
73	18	42260348	rs117763335	T	C	0.97	0.005	1.33×10^{-8}	O
74	18	50614732	rs11663393	A	G	1.03	0.004	1.56×10^{-10}	D, DO, N, O
75	18	52517906	rs1833288	A	G	1.03	0.005	4.54×10^{-8}	D, DS, DO, N, S, O
76	18	53101598	rs12958048	A	G	1.04	0.005	4.86×10^{-14}	D, DO, BS, N, S, O
77	19	30939989	rs33431	T	C	1.02	0.004	4.04×10^{-8}	DO, O
78	20	45841052	rs910187	A	G	0.97	0.005	3.09×10^{-9}	DO, O
79	22	41621714	rs2179744	A	G	1.03	0.005	3.83×10^{-12}	D, B, DO, BS, N, S, O
80	22	42815358	rs7288411	A	G	1.03	0.005	3.86×10^{-8}	NOVEL LOCUS
81	22	50679436	rs113872034	A	G	0.96	0.006	1.10×10^{-9}	O

Table 1: Loci genome-wide significant ($p < 5 \times 10^{-8}$) in the MOOD meta-analysis.
Locus – shared locus number for annotation (Supplementary Table 3), Chr – chromosome,
BP – base position, A1 – effect allele, A2 – non-effect allele, Previous report – locus
previously implicated in PGC MDD (D), PGC BD (B), previous combined studies of bipolar
disorder and major depressive disorder (BD), other studies of major depressive disorder or
depressive symptoms (DO), other studies of bipolar disorder (BO), previous combined
studies of bipolar disorder and schizophrenia (BS), previous combined studies of major
depressive disorder and schizophrenia (DS), neuroticism (N), schizophrenia (S), or other
studies (O – see Supplementary Table 4).

SNP-based heritability and genetic correlations

The estimate of SNP-based heritability for MOOD (8.8%) was closer to PGC

MDD (9%) than to PGC BD (17-23%) (15, 16). Significant genetic correlations

between MOOD and other traits included psychiatric and behavioural, reproductive,

cardiometabolic, and sociodemographic traits (Figure 1, Supplementary Table 5).

Genetic correlations with psychiatric and behavioural traits are consistently observed

across psychiatric traits (17, 61). The genetic correlation with educational attainment

differs, being negative in combined MDD, but positive in PGC BD (Supplementary

Table 6). The genetic correlation (r_g) between MOOD and educational attainment

was -0.058 (p=0.004), intermediate between the results of combined MDD and of

PGC BD. Notably, the genetic correlation with intelligence (IQ) was not significant in

combined MDD, PGC BD, nor MOOD (p>1.27x10^{-4}). However, sensitivity analyses

(see below), indicated that including 23andMe in the PGC MDD sample obscured a

negative genetic correlation of MDD with IQ.

The SNP-based heritability of down-sampled MOOD from LDSC was 11%,

closer to PGC MDD than to PGC BD (Supplementary Table 2). Genetic correlations

varied (Supplementary Tables 5 and 7) with some more similar to PGC BD

(schizophrenia: down-sampled rg = 0.61, combined MDD rg = 0.35, PGC BD rg =

0.7), and others more similar to combined MDD (ADHD: down-sampled rg = 0.48,

combined MDD rg = 0.45, PGC BD rg = 0.14). The genetic correlation with IQ was

significant (rg = -0.13, p = 5x10^{-7}), because the excluded 23andMe depression

cohort has a positive genetic correlation with IQ (rg = 0.06, p = 0.01). The greater

genetic correlation of MOOD with combined MDD (0.98) compared to PGC BD

(0.55) persisted when comparing down-sampled MOOD to combined MDD (0.85)

and PGC BD (0.75; Supplementary Table 6).

Figure 1a: Selected genetic correlations of psychiatric traits with the main meta-analysis (MOOD), the separate mood disorder analyses (combined MDD and PGC BD), and the down-sampled analyses (down-sampled MOOD, down-sampled MDD). Full genetic correlation results are provided in Supplementary Table 5.

Figure 1b: Selected genetic correlations of other traits with the main meta-analysis (MOOD), the separate mood disorder analyses (combined MDD and PGC BD), and the down-sampled analyses (down-sampled MOOD, down-sampled MDD). Full genetic correlation results are provided in Supplementary Table 5.

Relationship between mood disorder subtypes

Analyses were performed using GWAS data from subtypes of bipolar disorder

(BD1, BD2, SAB) and major depressive disorder (rMDD, sMDD, subMDD). SNP-

based heritability for the subtypes ranged from subMDD and sMDD (8%), through

BD2 and rMDD (10% and 12%, respectively) to BD1 and SAB (22% and 29%

respectively, Figure 2, Supplementary Table 2).

Figure 2: SNP-based heritability estimates for the subtypes of bipolar disorder and
subtypes of major depressive disorder. Points = SNP-based heritability estimates.
Lines = 95% confidence intervals. Full SNP-based heritability results are provided in
Supplementary Table 2.

7   The major depressive disorder subtypes were strongly and significantly

8   genetically correlated (r_g = 0.9-0.94, $p_{rg = 0}$ < 8.3x10^{-4}). These correlations did not

9   differ significantly from 1 (all $p_{rg = 1}$ > 0.3), nor from each other (all $p_{\Delta rg = 0}$ > 0.5, Figure

10   2, Supplementary Table 8). BD1 and SAB were strongly correlated (r_g = 0.77, $p_{rg = 0}$ =

6×10^{-13}, $p_{rg=1} = 0.03$), as were BD1 and BD2 ($r_g = 0.86$, $p_{rg=0} = 3 \times 10^{-16}$, $p_{rg=1} = 0.2$).

However, BD2 was not significantly correlated with SAB ($r_g = 0.22$, $p_{rg=0} = 0.02$).

In hierarchical clustering, BD2 clustered with the major depressive disorder

subtypes rather than the bipolar disorder subtypes. The strength of correlation

between BD2 and BD1 did not differ from that between BD2 and rMDD ($r_g = 0.68$, p_{rg}

$_{=0} = 3 \times 10^{-8}$, $p_{rg=1} = 0.01$), following multiple testing correction ($\Delta r_g = 0.18$, $p = 0.02$).

Overall, these results suggest a spectrum of genetic relationships between major

depressive disorder and bipolar disorder, with BD2 bridging the two disorders (Figure

3; Supplementary Figure 9). This spectrum remained when six external phenotypes

were added, and was supported by results from principal component analysis

(Supplementary Results, Supplementary Figure 10).

Figure 3: Genetic correlations across the mood disorder spectrum. Labelled arrows show genetic correlations significantly different from 0. Solid arrows represent genetic correlations not significantly different from 1 (p < 0.00333, Bonferroni correction for 15 tests). Full results are provided in Supplementary Table 8.

Polygenic risk score analyses showed that individuals with high polygenic risk

scores for PGC BD were more likely to report rMDD than sMDD, and more likely to

report sMDD than subMDD (Supplementary Results).

Tissue and cell-type specificity analyses

The results of gene-wise and gene set analyses are described in the

Supplementary Results. The tissue-specificity of associated genes differed minimally

between the analyses (Supplementary Table 9). All brain regions were significantly

enriched in all analyses, and the pituitary was also enriched in combined MDD and

PGC BD ($p < 9.43 \times 10^{-4}$, Bonferroni correction for 53 regions, Supplementary Table

9). Results from down-sampled MOOD and down-sampled MDD were generally

consistent with the main analyses, except spinal cord was not enriched in either, nor

was the cordate in the down-sampled MDD analysis.

In contrast, cell-type enrichments differed between combined MDD and PGC

BD (Figure 4, Supplementary Tables 10 and 11). Genes associated with PGC BD

were enriched for expression in pyramidal cells from the CA1 region of the

hippocampus and the somatosensory cortex, and in striatal interneurons. None of

these enrichments were significant in combined MDD. Genes only associated with

combined MDD were significantly enriched for expression in neuroblasts and

dopaminergic neurons from adult mice. Further cell-types (dopaminergic

neuroblasts; dopaminergic, GABAergic and midbrain nucleus neurons from

embryonic mice; interneurons; and medium spiny neurons) were enriched for both

combined MDD and PGC BD, but the rank and strength of enrichment differed, most

notably for medium spiny neurons. The general pattern of differences persisted when

comparing PGC BD with down-sampled MDD, although genes associated with

down-sampled MDD were not enriched for expression in adult dopaminergic

neurons, embryonic midbrain nucleus neurons, interneurons, nor medium spiny

neurons (Supplementary Figure 11).

Figure 4: Cell-type expression specificity of genes associated with bipolar disorder (PGC BIP, left) and major depressive disorder (combined MDD, right). Black vertical lines = significant enrichment (p < 2x10^{-3}, Bonferroni correction for 24 cell types). See Supplementary Table 10 for full results.

Shared and distinct relationships with mood disorders and inferred causality

Bidirectional Mendelian randomisation was used to investigate previously-

described relationships between mood disorder phenotypes (combined MDD, PGC

BD) and external traits: schizophrenia, educational attainment, IQ, body mass index

(BMI) and coronary artery disease (CAD; Figure 5, Supplementary Table 12).

Associations with PGC BD should be interpreted cautiously, as only 19 loci reached

1   genome-wide significance, several of which were removed as potentially pleiotropic

2   in the analyses below.

Figure 5: GSMR results from analyses with the main meta-analysis (MOOD), and the major depression and bipolar disorder analyses (combined MDD, PGC BD). External traits are coronary artery disease (CAD), educational attainment (EDU), body mass index (BMI), and schizophrenia (SCZ). Betas are on the scale of the outcome GWAS (logit for binary traits, phenotype scale for continuous). * $p < 0.004$ (Bonferroni correction for two-way comparisons with six external traits). For figure data, including the number of non-pleiotropic SNPs included in each instrument, see Supplementary Table 12.

A positive bidirectional relationship was observed between combined MDD

and PGC BD, and between schizophrenia and both combined MDD and PGC BD.

This is consistent with psychiatric disorders acting as causal risk factors for the

development of further psychiatric disorders, or being correlated with other causal

risk factors, including (but not limited to) the observed shared genetic basis.

The relationship with educational years differed between the mood disorders

– there was a negative bidirectional relationship between educational years and

combined MDD, but a positive bidirectional relationship with PGC BD (albeit with

only nominal significance from PGC BD to educational years). In contrast, no

significant relationship was observed between mood phenotypes and IQ. This is

consistent with differing causal roles of education (or correlates of education) on the

mood disorders, with a weaker reciprocal effect of the mood disorders altering the

length of education.

A positive association was seen from BMI to combined MDD, but not from

combined MDD to BMI. In contrast, only a nominally significant negative relationship

was seen from PGC BD to BMI. A positive association was observed from combined

MDD to CAD; no relationship was observed between CAD and PGC BD.

Discussion

We identified 73 genetic loci by meta-analysing cohorts of major depression

and bipolar disorder, including 15 loci novel to mood disorders. Our overall mood

disorders meta-analysis results (MOOD) have more in common with our major

depressive disorder analysis (combined MDD) than our bipolar disorder analysis

(PGC BD). Partly, this results from the greater power of the major depressive

disorder analysis compared to the bipolar disorder analysis. Nevertheless, genetic

associations from our sensitivity analysis with equivalently powered cohorts (using

down-sampled MDD in place of combined MDD) still showed a greater overall

similarity to those from major depressive disorder rather than bipolar disorder.

This may reflect a complex genetic architecture in bipolar disorder, wherein

one set of variants may be associated more with manic symptoms and another set

with depressive symptoms. Variants associated more with mania (or psychosis) may

have higher effect sizes, detectable at current bipolar disorder GWAS sample sizes,

and may not be strongly associated with major depressive disorder. This could

contribute to the observed higher heritability of bipolar disorder compared to major

depressive disorder, and agrees with reports that most of the genetic variance for

mania is not shared with depression (13, 14). In this case, meta-analysis of bipolar

disorder and major depressive disorder cohorts would support variants associated

more with depression, but not those associated more with mania. This is consistent

with our findings, and with depressive symptoms being both the unifying feature of

the mood disorders and the core feature of major depressive disorder.

We assessed genetic correlations between mood disorder subtypes. We

observed high, consistent correlations between major depressive disorder subtypes,

including sub-threshold depression. Bipolar disorder type 2 showed greater genetic

similarity to major depressive disorder compared to type 1. In this, we build on

similar findings from polygenic risk scores analyses (16, 56). Individuals with high

polygenic risk scores for PGC BD were more likely to report recurrent than single-

episode major depressive disorder. However, the genetic correlation of PGC BD with

recurrent major depressive disorder was not significantly greater than that with

single-episode major depressive disorder. This might reflect the difference in power

between these methods. We also examined the genetic correlations between mood

disorder subtypes in the context of relevant external traits (Supplementary Results).

Our subtype analyses support a genetic mood spectrum consisting of the

schizophrenia-like bipolar disorder type 1 and schizoaffective disorder at one pole,

and the depressive disorders at the other, with bipolar disorder type 2 occupying an

intermediate position.

Conditional and reversed-effect analyses (Supplementary Results) suggest

that few of the loci we identified are disorder-specific. However, our results highlight

some differences between the genetics of the mood disorders. The expression

specificity of associated genes in mouse brain cell types differed between bipolar

disorder and major depressive disorder analyses. Cell-types more associated with

bipolar disorder (pyramidal neurons and striatal interneurons) were also enriched in

analyses of schizophrenia (38). Cell-types more associated in major depressive

disorder (neuroblasts, adult dopaminergic neurons, embryonic GABAergic neurons)

had weaker enrichments in schizophrenia, but were enriched in analyses of

neuroticism (57). The higher rank of the enrichment of serotonergic neurons with

major depressive disorder compared to bipolar disorder is striking given the use of

drugs targeting the serotonergic system in the treatment of depression (63).

Nevertheless, cell-type enrichment analyses are still novel, and require cautious

interpretation, especially given the use of non-human reference data (38, 64).

We explored potential causal relationships between the mood disorders and

other traits using Mendelian randomisation. The interpretation of these analyses is

challenging, especially for complex traits, when the ascertainment of cases varies,

and when there are relatively few (< 20) variants used as instruments (for example,

in the PGC BD and down-sampled analyses presented) (41, 67, 68). Major

depressive disorder and bipolar disorder demonstrate considerable heterogeneity

(as our subtype analyses show for bipolar disorder types 1 and 2), potentially

confounding the results of Mendelian randomisation. That said, our analyses are

consistent with a bidirectional influence of educational attainment on risk for mood

disorders (and vice versa), with different directions of effect in the two mood

disorders. We found no significant relationship between IQ and either mood disorder.

We also find results consistent with major depressive disorder increasing the risk for

coronary artery disease in a relatively well powered analysis. This mirrors

epidemiological findings, although the mechanism remains unclear (69).

Despite the presence of depressive episodes, the mood disorders are

diagnostically distinct. This is reflected in their differing epidemiology – for example,

more women than men suffer from major depressive disorder, whereas diagnoses of

bipolar disorder are roughly equal between the sexes (3). Differences in our genetic

results between major depressive disorder and bipolar disorder may result from

epidemiological heterogeneity, rather than distinct biological mechanisms (70).

Deeper phenotyping of GWAS datasets is ongoing, and will enable the effect of

confounding factors such as sex to be estimated in future studies (71).

We extend previous findings showing genetic continuity across the mood

disorders (15–17, 56). Combined analyses of major depressive disorder and bipolar

disorder may increase variant discovery, as well as the discovery of shared and

distinct neurobiological gene sets and cell types. Our results also indicate some

genetic differences between major depressive disorder and bipolar disorder,

including opposite bidirectional relationships of each mood disorder with educational

attainment, a possible influence of major depressive disorder on coronary artery

disease risk and differing mouse brain cell types implicated by the enrichment

patterns of associated genes in each disorder. Finally, our data are consistent with

the existence of a genetic mood disorder spectrum with separate clusters for bipolar

disorder type 1 and depressive disorders, linked by bipolar disorder type 2, and with

depression as the common symptom. The mood disorders have a partially genetic

aetiology that is partly shared. The identification of specific sets of genetic variants

differentially associated with depression and with mania remains an aim for future

research.

Acknowledgements

This paper has previously been made available as a preprint on bioRxiv at

https://www.biorxiv.org/content/10.1101/383331v1.

We are deeply indebted to the investigators who comprise the PGC, and to the

hundreds of thousands of subjects who have shared their life experiences with PGC

investigators. This study represents independent research partly funded by the

National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) at

South London and Maudsley NHS Foundation Trust and King's College London. The

views expressed are those of the author(s) and not necessarily those of the NHS,

the NIHR or the Department of Health and Social Care. High performance computing

facilities were funded with capital equipment grants from the GSTT Charity

(TR130505) and Maudsley Charity (980). The PGC has received major funding from

the US National Institute of Mental Health (NIMH) and the US National Institute of

Drug Abuse (NIDA) of the US National Institutes of Health (NIH; U01 MH109528 to

PFS, U01MH109514 to MCO, and U01 MH1095320 to A Agrawal). We acknowledge

the continued support of the NL Genetic Cluster Computer

(http://www.geneticcluster.org/) hosted by SURFsara in the management and

curation of PGC data, with funding from Scientific Organization Netherlands (480-05-

003 to DP). Central analysis of PGC data was funded by UK Medical Research

Council (MRC) Centre and Program Grants (G0801418, G0800509 to PAH, MCO,

MJO) and grants from the Australian National Health and Medical Research Council

(NHMRC; 1078901,108788 to NRW). GB, JRIC, HG, CL were supported in part by

the NIHR as part of the Maudsley BRC. DP is funded by the Dutch Brain Foundation

and the VU University Amsterdam Netherlands. PFS gratefully acknowledges

support from the Swedish Research Council (Vetenskapsrådet, award D0886501).

Acknowledgements and funding for individual cohorts follows. BD_TRS: This work

was funded by the Deutsche Forschungsgemeinschaft (DFG, grants FOR2107

DA1151/5-1, SFB-TRR58, and Project C09 to UD) and the Interdisciplinary Center

for Clinical Research (IZKF) of the medical faculty of Münster (grant Dan3/012/17 to

UD). BiGS: Research was funded by the NIMH (Chicago: R01 MH103368 to ESG,

NIMH: R01 MH061613 and ZIA MH002843 to FJM, Pittsburgh: MH63480 to VN,

UCSD: MH078151, MH081804, MH59567 to JK). FJM was supported by the NIMH

Intramural Research Program, NIH, DHHS. BOMA-Australia: Funding was supplied

by the Australian NHMRC (1037196, 1066177, and 1063960 to JMF, 1103623 to

SEM, 1037196 to PBM, 1078399 to GWM, 1037196 to PRS). JMF would like to

thank Janette M O'Neil and Betty C Lynch for their support. BOMA-Germany I,

BOMA-Germany II, BOMA-Germany III, PsyCourse, and Münster MDD Cohort: This

work was supported by the German Ministry for Education and Research (BMBF)

through the Integrated Network IntegraMent (Integrated Understanding of Causes

and Mechanisms in Mental Disorders), under the auspices of the e:Med program

(01ZX1314A/01ZX1614A to MMN and SC, 01ZX1314G/01ZX1614G to MR,

01ZX1314K to TGS) and through grants NGFNplus MooDS (Systematic

Investigation of the Molecular Causes of Major Mood Disorders and Schizophrenia;

grant 01GS08144, 01GS08147 to MMN, MR and SC). This work was also supported

by the DFG (NO246/10-1 to MMN [FOR 2107], RI 908/11-1 to MR [FOR 2107], WI

3429/3-1 to SHW, SCHU 1603/4-1, SCHU 1603/5-1 [KFO 241] and SCHU 1603/7-1

[PsyCourse] to TGS), the Swiss National Science Foundation (SNSF, grant 156791

to SC) and the European Union (N Health-F2-2008-222963 to BTB and VA). MMN is

supported through the Excellence Cluster ImmunoSensation. TGS is supported by

an unrestricted grant from the Dr. Lisa-Oehler Foundation. AJF received support

from the BONFOR Programme of the University of Bonn, Germany. MH was

supported by the DFG. BOMA-Romania: The work was supported by Unitatea

Executiva Pentru Finantarea Invatamantului Superior a Cercetarii (89/2012 to MG-

S). Bulgarian Trios: Recruitment was funded by the Janssen Research Foundation,

and genotyping was funded by multiple grants to the Stanley Center for Psychiatric

Research at the Broad Institute from the Stanley Medical Research Institute, The

Merck Genome Research Foundation, and the Herman Foundation to GK. CoFaMS

– Adelaide: Research was funded by the Australian NHMRC (APP1060524 to BTB).

CONVERGE: Research was funded by the Wellcome Trust (WT090532/Z/09/Z,

WT083573/Z/07/Z and WT089269/Z/09/Z to J Flint) and the NIMH (MH100549 to

KSK). Danish RADIANT: Research was funded by Højteknologifonden (0001-2009-2

to TW) and the Lundbeck Foundation, (R24-A3242 to TW). deCODE: Research was

funded by FP7-People-2011-IAPP grant agreement PsychDPC, (286213 to KS), and

NIDA (R01 DA017932 to KS, R01 DA034076 to TT). The authors are thankful to the

participants and staff at the Patient Recruitment Center. Edinburgh: Genotyping was

conducted at the Genetics Core Laboratory at the Clinical Research Facility

(University of Edinburgh). Research was funded by the Wellcome Trust

(104036/Z/14/Z to AMM, T-KC, and DJP). DJM is supported by an NRS Clinical

Fellowship funded by the CSO. EGCUT: Research was funded by European Union

Project, (EstRC-IUT20-60, No. 2014-2020.4.01.15- 0012, 692145 to AM). Fran: This

research was supported by Foundation FondaMental, Créteil, France and by the

Investissements d'Avenir Programs managed by the ANR (ANR-11-IDEX-0004-02

and ANR-10-COHO-10-01 to ML). GenPOD/Newmeds: Research was funded by

MRC (G0200243 to GL and MCO), EU 6th Framework, (LSHB-CT-2003-503428 to

RH), IMI-JU, (15008 to GL). GenScot: Research was funded by the UK Chief

Scientist Office (CZD/16/6 to DJP) and the Scottish Funding Council (HR03006 to

DJP). We are grateful to all the families who took part, the general practitioners and

the Scottish School of Primary Care for their help in recruiting them, and the whole

Generation Scotland team, which includes interviewers, computer and laboratory

technicians, clerical workers, research scientists, volunteers, managers,

receptionists, healthcare assistants and nurses. Genotyping was conducted at the

Genetics Core Laboratory at the Clinical Research Facility (University of Edinburgh).

GERA: Participants in the Genetic Epidemiology Research on Adult Health and

Aging Study are part of the Kaiser Permanente Research Program on Genes,

Environment, and Health, supported by the NIA, NIMH, OD, (RC2 AG036607 to CS,

NRisch) and the Wayne and Gladys Valley Foundation, The Ellison Medical

Foundation, the Robert Wood Johnson Foundation, and the Kaiser Permanente

Regional and National Community Benefit Programs. GSK_Munich: We thank all

participants in the GSK-Munich study. We thank numerous people at GSK and Max-

Planck Institute, BKH Augsburg and Klinikum Ingolstadt in Germany who contributed

to this project. Halifax: Halifax data were obtained with support from the Canadian

Institutes of Health Research to MA. Harvard i2b2: Research funded by NIMH (R01

MH085542 to JWS, R01 MH086026 to RHP). iPSYCH: The iPSYCH (The Lundbeck

Foundation Initiative for Integrative Psychiatric Research) team acknowledges

funding from The Lundbeck Foundation (grant no R102-A9118 and R155-2014-

1724, R129-A3973 and R24- A3243), the Stanley Medical Research Institute, the

European Research Council (294838), the Novo Nordisk Foundation for supporting

the Danish National Biobank resource, the Capital Region of Denmark, (R144-

A5327), and grants from Aarhus and Copenhagen Universities and University

Hospitals, including support to the iSEQ Center, the GenomeDK HPC facility, and

the CIRRAU Center. All funding was to the iPSYCH PIs: TW, ADB, OM, MN, DH,

and PBM. Janssen: Funded by Janssen Research & Development, LLC. We are

grateful to the study volunteers for participating in the research studies and to the

clinicians and support staff for enabling patient recruitment and blood sample

collection. We thank the staff in the former Neuroscience Biomarkers of Janssen

Research & Development for laboratory and operational support (e.g., biobanking,

processing, plating, and sample de-identification), and to the staff at Illumina for

genotyping Janssen DNA samples. MARS/BiDirect: This work was funded by the

Max Planck Society, by the Max Planck Excellence Foundation, and by a grant from

the German Federal Ministry for Education and Research (BMBF) in the National

Genome Research Network framework (NGFN2 and NGFN-Plus, FKZ 01GS0481),

and by the BMBF Program FKZ01ES0811. We acknowledge all study participants.

We thank numerous people at Max-Planck Institute, and all study sites in Germany

and Switzerland who contributed to this project. Controls were from the Dortmund

Health Study which was supported by the German Migraine & Headache Society,

and by unrestricted grants to the University of Münster from Almirall, Astra Zeneca,

Berlin Chemie, Boehringer, Boots Health Care, Glaxo-Smith-Kline, Janssen Cilag,

McNeil Pharma, MSD Sharp & Dohme, and Pfizer. Blood collection was funded by

the Institute of Epidemiology and Social Medicine, University of Münster. Genotyping

was supported by the German Ministry of Research and Education (BMBF grant

01ER0816, 01ER1506 to KB). Mayo Bipolar Disorder Biobank: Research was

funded by grants from the Marriot Foundation and the Mayo Clinic Center for

Individualized Medicine to JMB and MF. Michigan (NIMH/Pritzker Neuropsychiatric

Disorders Research Consortium): Research was funded by NIMH (R01

MH09414501A1, MH105653 to MB). We thank the participants who donated their

time and DNA to make this study possible. We thank members of the NIMH Human

Genetics Initiative and the University of Michigan Prechter Bipolar DNA Repository

for generously providing phenotype data and DNA samples. Many of the authors are

members of the Pritzker Neuropsychiatric Disorders Research Consortium which is

supported by the Pritzker Neuropsychiatric Disorders Research Fund L.L.C. A

shared intellectual property agreement exists between this philanthropic fund and the

University of Michigan, Stanford University, the Weill Medical College of Cornell

University, HudsonAlpha Institute of Biotechnology, the Universities of California at

Davis, and at Irvine, to encourage the development of appropriate findings for

research and clinical applications. Mount Sinai: This work was funded in part by a

NARSAD Young Investigator award to EAS, and by NIH (R01MH106531,

R01MH109536 to PS and EAS). NeuRA-CASSI-Australia: This work was funded by

the NSW Ministry of Health, Office of Health and Medical Research, and by the

NHRMC (568807 to CSW and TWW). CSW was a recipient of NHMRC Fellowships

(#1117079, #1021970). NeuRA-IGP-Australia: Research was funded by the NHMRC

(630471, 1061875, 1081603 to MJG. NESDA: Research was funded by Nederlandse

Organisatie voor Wetenschappelijk (NOW; ZonMW Geestkracht grant to PWJHP).

Norway: Research was funded by the Vetenskapsrådet to IA, the Western Norway

Regional Health Authority to KJO, the Research Council of Norway (#421716 to IM,

#249711, #248778, #223273, and #217776 to OAA), the South-East Norway

Regional Health Authority (#2012-132 and #2012-131 to OAA, #2016-064 to OBS,

#2017-004 to OAA and OBS, #2013-088, #2014-102, and #2011-085 to IM), and the

KG Jebsen Stiftelsen to OAA. TE was funded by The South-East Norway Regional

Health Authority (#2015-078) and a research grant from Mrs. Throne-Holst. NTR:

Research was funded by NWO (480-15-001/674 to DIB). Pfizer: Research was

funded by the EU Innovative Medicine Initiative Joint Undertaking (115008.5).

PsyColaus: PsyCoLaus/CoLaus received additional support from research grants

from GlaxoSmithKline and the Faculty of Biology and Medicine of Lausanne, and the

SNSF (3200B0–105993, 3200B0- 118308, 33CSCO-122661, 33CS30-139468,

33CS30- 148401 to MP). QIMR: We thank the twins and their families for their willing

participation in our studies. Research was funded by NHMRC (941177, 971232,

3399450 and 443011 to NGM) and NIAAA (AA07535, AA07728, and AA10249 to

ACH). RADIANT: Research was funded by MRC (G0701420 to GB and CML,

G0901245 to GB) and NIMH (U01 MH109528 to GB). Rotterdam Study: The

Rotterdam Study is funded by Erasmus Medical Center and Erasmus University, and

NWO (175.010.2005.011, 911-03- 012 to AGU). SHIP-LEGEND/TREND: SHIP is

part of the Community Medicine Research net of the University of Greifswald which

is funded by the DFG (GR 1912/5-1 to HJG), Federal Ministry of Education and

Research (grants 01ZZ9603, 01ZZ0103, and 01ZZ0403), the Ministry of Cultural

Affairs, and the Social Ministry of the Federal State of Mecklenburg-West

Pomerania. Genotyping in SHIP was funded by Siemens Healthineers and the

Federal State of Mecklenburg-West Pomerania. Genotyping in SHIP-TREND-0 was

supported by the Federal Ministry of Education and Research (grant 03ZIK012).

Span2: Research was funded by Instituto de Salud Carlos III (PI12/01139,

PI14/01700, PI15/01789, PI16/01505), and cofinanced by the European Regional

Development Fund (ERDF), Agència de Gestió d'Ajuts Universitaris i de Recerca-

AGAUR, Generalitat de Catalunya (2014SGR1357), Departament de Salut,

Generalitat de Catalunya, Spain, and a NARSAD Young Investigator Grant from the

Brain & Behavior Research Foundation. This project has also received funding from

the European Union's Horizon 2020 Research and Innovation Programme under the

grant agreements No 667302 and 643051. CSM is a recipient of a Sara Borrell

contract (CD15/00199) and a mobility grant (MV16/00039) from the Instituto de

Salud Carlos III, Ministerio de Economía, Industria y Competitividad, Spain. MR is a

recipient of a Miguel de Servet contract (CP09/00119 and CPII15/00023) from the

Instituto de Salud Carlos III, Ministerio de Economía, Industria y Competitividad,

Spain. STAR*D: Research was funded by NIMH (R01 MH-072802 to SPH). The

authors appreciate the efforts of the STAR*D investigator team for acquiring,

compiling, and sharing the STAR*D clinical data set. SUNY DMC: Research was

funded by NIMH (R01MH085542 to CP, MTP, JAK, and HM). SWEBIC: Research

was funded by NIMH (MH077139), the Vetenskapsrådet (K2014-62X-14647-12-51

and K2010-61P-21568-01-4), the Swedish foundation for Strategic Research (KF10-

0039) and the Stanley Center for Psychiatric Research, Broad Institute from a grant

from Stanley Medical Research Institute, all to ML. We are deeply grateful for the

participation of all subjects contributing to this research, and to the collection team

that worked to recruit them. We also wish to thank the Swedish National Quality

Register for Bipolar Disorders: BipoläR. Sweden: This work was funded by the

Vetenskapsrådet (to MS and CL), the Stockholm County Council (to MS, CL, LB, LF,

and UÖ) and the Söderström Foundation (to LB). TwinGene: Research was funded

by GenomeEUtwin, (EU/QLRT-2001-01254; QLG2-CT-2002-01254 to NLP), Heart

and Lung Foundation (20070481 to PKM), SFF and Vetenskapsrådet, (M-2005-1112

to U de Faire).We thank the Karolinska Institutet for infrastructural support of the

Swedish Twin Registry. UCL: Research was funded by the MRC (G1000708 to AM).

UCLA-Utrecht (Los Angeles): Research was funded by NIMH (R01MH090553,

U01MH105578 to NBF, RAO, LMOL, and APSO). UK - BDRN: Research was

funded by MRC Centre and Program Grants (G0801418, G0800509 to MCO and

MJO), the Wellcome Trust (078901 to NC, IJ, LAJ), the Stanley Medical Research

Institute (5710002223-01 to NC, IJ, LAJ), and a European Commission Marie Curie

Fellowship (623932 to ADF). BDRN would like to acknowledge the research

participants who continue to give their time to participate in our research. UK

Biobank: This research has been carried out under application numbers 4844, 6818,

and 16577, funded by the National Institute for Health Research under its Biomedical

Research Centres funding initiative (to GB) and the Wellcome Trust (04036/Z/14/Z to

AMM). UNIBO / University of Barcelona, Hospital Clinic, IDIBAPS, CIBERSAM: EV

thanks the support of the Spanish Ministry of Economy and Competitiveness

(PI15/00283 to EV) integrated into the Plan Nacional de I+D+I y cofinanciado por el

ISCIII-Subdirección General de Evaluación y el Fondo Europeo de Desarrollo

Regional (FEDER); CIBERSAM; and the Comissionat per a Universitats i Recerca

del DIUE de la Generalitat de Catalunya to the Bipolar Disorders Group (2014 SGR

398). USC: Research funded by NIH (R01MH085542 to JLS). WTCCC: The principal

funder of this project was the Wellcome Trust to NC and AHY. For the 1958 Birth

Cohort, venous blood collection was funded by the UK MRC. AHY is funded by the

National Institute for Health Research (NIHR) Biomedical Research Centre at South

London and Maudsley NHS Foundation Trust and King's College London. 23andMe:

We thank the 23andMe research participants included in the analysis, all of whom

provided informed consent and participated in the research online according to a

human subjects protocol approved by an external AAHRPP-accredited institutional

review board (Ethical & Independent Review Services), and the employees of

23andMe for making this work possible. 23andMe acknowledges the-invaluable

contributions of Michelle Agee, Babak Alipanahi, Adam Auton, Robert K. Bell,

Katarzyna Bryc, Sarah L. Elson, Pierre Fontanillas, Nicholas A. Furlotte, David A.

Hinds, Bethann S. Hromatka, Karen E. Huber, Aaron Kleinman, Nadia K. Litterman,

Matthew H. McIntyre, Joanna L. Mountain, Carrie A.M. Northover, Steven J. Pitts, J.

Fah Sathirapongsasuti, Olga V. Sazonova, Janie F. Shelton, Suyash Shringarpure,

Chao Tian, Joyce Y. Tung, Vladimir Vacic, and Catherine H. Wilson.

<u>Disclosures</u>

OA Andreassen has received speaker fees from Lundbeck. ATF Beekman is on

speaker's bureaus for Lundbeck and GlaxoSmithKline. G Breen reports consultancy

and speaker fees from Eli Lilly, Otsuka and Illumina and grant funding from Eli Lilly.

G Crawford is a cofounder of Element Genomics. E Domenici was formerly an

employee of Hoffmann–La Roche and a consultant to Roche and Pierre-Fabre. J

Nurnberger is an investigator for Janssen and was an investigator for Assurex. SA

Paciga is an employee of Pfizer. JA Quiroz was formerly an employee of Hoffmann–

La Roche. S Steinberg, H Stefansson, K Stefansson and TE Thorgeirsson are

employed by deCODE Genetics/Amgen. PF Sullivan reports the following potentially

competing financial interests. Current: Lundbeck (advisory committee, grant

recipient). Past three years: Pfizer (scientific advisory board), Element Genomics

(consultation fee), and Roche (speaker reimbursement). AH Young has given paid

lectures and is on advisory boards for the following companies with drugs used in

affective and related disorders: Astrazenaca, Eli Lilly, Janssen, Lundbeck, Sunovion,

Servier, Livanova. AH Young is Lead Investigator for Embolden Study

(Astrazenaca), BCI Neuroplasticity study and Aripiprazole Mania Study, which are

investigator-initiated studies from Astrazenaca, Eli Lilly, Lundbeck, and Wyeth. All

other authors declare no financial interests or potential conflicts of interest.

Article Information

The Bipolar Disorder and Major Depressive Disorder Working Groups of the

Psychiatric Genomics Consortium are collaborative co-authors for this article. The

individual authors are (numbers refer to affiliations listed in the Supplement): Enda

M. Byrne[4], Andreas J. Forstner[5;6;7;8;9], Peter A. Holmans[10], Christiaan A. de Leeuw[11],

Manuel Mattheisen[12;13;14;15;16], Andrew McQuillin[17], Jennifer M. Whitehead Pavlides[18],

Tune H. Pers[19;20], Stephan Ripke[21;22;23], Eli A. Stahl[19;24;25], Stacy Steinberg[26], Vassily

Trubetskoy[22], Maciej Trzaskowski[4], Yunpeng Wang[27;28], Liam Abbott[21], Abdel

Abdellaoui[29], Mark J. Adams[30], Annelie Nordin Adolfsson[31], Esben Agerbo[16;32;33],

Huda Akil[34], Diego Albani[35], Ney Alliey-Rodriguez[36], Thomas D. Als[12;13;16], Till F. M.

Andlauer[37;38], Adebayo Anjorin[39], Verneri Antilla[23], Sandra Van der Auwera[40],

Swapnil Awasthi[22], Silviu-Alin Bacanu[41], Judith A Badner[42], Marie Bækvad-

Hansen[16;43], Jack D. Barchas[44], Nicholas Bass[17], Michael Bauer[45], Aartjan T. F.

Beekman[46], Richard Belliveau[21], Sarah E. Bergen[3], Tim B. Bigdeli[41;47], Elisabeth B.

Binder[37;48], Erlend Bøen[49], Marco Boks[50], James Boocock[51], Monika Budde[52],

William Bunney[53], Margit Burmeister[54], Henriette N. Buttenschøn[3;12;55], Jonas

Bybjerg-Grauholm[16;43], William Byerley[56], Na Cai[57;58], Miquel Casas[59;60;61;62], Enrique

Castelao[63], Felecia Cerrato[21], Pablo Cervantes[64], Kimberly Chambert[21], Alexander

W. Charney[25], Danfeng Chen[21], Jane Hvarregaard Christensen[12;13;55], Claire

Churchhouse[21;23], David St Clair[65], Toni-Kim Clarke[30], Lucía Colodro-Conde[66],

William Coryell[67], Baptiste Couvy-Duchesne[18;68], David W. Craig[69], Gregory E.

Crawford[70;71], Cristiana Cruceanu[37;64], Piotr M. Czerski[72], Anders M. Dale[73;74;75;76],

Gail Davies[77], Ian J. Deary[77], Franziska Degenhardt[7;8], Jurgen Del-Favero[78], J

Raymond DePaulo[79], Eske M. Derks[66], Nese Direk[80;81], Srdjan Djurovic[82;83], Amanda

L. Dobbyn[24;25], Conor V. Dolan[29], Ashley Dumont[21], Erin C. Dunn[21;84;85], Thalia C.

Eley[1], Torbjørn Elvsåshagen[86;87], Valentina Escott-Price[10], Chun Chieh Fan[76], Hilary K. Finucane[88;89], Sascha B. Fischer[5;9], Matthew Flickinger[90], Jerome C. Foo[91], Tatiana M. Foroud[92], Liz Forty[10], Josef Frank[91], Christine Fraser[10], Nelson B. Freimer[93], Louise Frisén[94;95;96], Katrin Gade[52;97], Diane Gage[21], Julie Garnham[98], Claudia Giambartolomei[51], Fernando S. Goes[99], Jaqueline Goldstein[21], Scott D. Gordon[66], Katherine Gordon-Smith[100], Elaine K. Green[101], Melissa J. Green[102], Tiffany A. Greenwood[75], Jakob Grove[12;13;16;103], Weihua Guan[104], Lynsey S. Hall[30;105], Marian L. Hamshere[10], Christine Søholm Hansen[16;43], Thomas F. Hansen[16;106;107], Martin Hautzinger[108], Urs Heilbronner[52], Albert M. van Hemert[109], Stefan Herms[5;7;8;9], Ian B. Hickie[110], Maria Hipolito[111], Per Hoffmann[5;7;8;9], Dominic Holland[73;112], Georg Homuth[113], Carsten Horn[114], Jouke-Jan Hottenga[29], Laura Huckins[24;25], Marcus Ising[15], Stéphane Jamain[116;117], Rick Jansen[46], Jessica S. Johnson[24;25], Simone de Jong[1;2], Eric Jorgenson[118], Anders Juréus[3], Radhika Kandaswamy[1], Robert Karlsson[3], James L. Kennedy[119;120;121;122], Farnush Farhadi Hassan Kiadeh[123], Sarah Kittel-Schneider[124], James A. Knowles[125;126], Manolis Kogevinas[127], Isaac S. Kohane[128;129;130], Anna C. Koller[7;8], Julia Kraft[22], Warren W. Kretzschmar[131], Jesper Krogh[132], Ralph Kupka[46;133], Zoltán Kutalik[134;135], Catharina Lavebratt[94], Jacob Lawrence[136], William B. Lawson[111], Markus Leber[137], Phil H. Lee[21;23;138], Shawn E. Levy[139], Jun Z. Li[140], Yihan Li[131], Penelope A. Lind[66], Chunyu Liu[141], Loes M. Olde Loohuis[93], Anna Maaser[7;8], Donald J. MacIntyre[142;143], Dean F. MacKinnon[99], Pamela B. Mahon[79;144], Wolfgang Maier[145], Robert M. Maier[18], Jonathan Marchini[146], Lina Martinsson[95], Hamdi Mbarek[29], Steve McCarroll[21;147], Patrick McGrath[148], Peter McGuffin[1], Melvin G. McInnis[149], James D. McKay[150], Helena Medeiros[126], Sarah E. Medland[66], Divya Mehta[18;151], Fan Meng[34;149], Christel M. Middeldorp[29;152;153], Evelin Mihailov[154], Yuri Milaneschi[46], Lili Milani[154], Saira Saeed Mirza[80], Francis M.

Mondimore[99], Grant W. Montgomery[4], Derek W. Morris[155;156], Sara Mostafavi[157;158],

Thomas W Mühleisen[5;159], Niamh Mullins[1], Matthias Nauck[160;161], Bernard Ng[158],

Hoang Nguyen[24;25], Caroline M. Nievergelt[75;162], Michel G. Nivard[29], Evaristus A.

Nwulia[111], Dale R. Nyholt[163], Claire O'Donovan[98], Paul F. O'Reilly[1], Anil P. S. Ori[93],

Lilijana Oruc[164], Urban Ösby[165], Hogni Oskarsson[166], Jodie N. Painter[66], José

Guzman Parra[167], Carsten Bøcker Pedersen[16;32;33], Marianne Giørtz Pedersen[16;32;33],

Amy Perry[100], Roseann E. Peterson[41;168], Erik Pettersson[3], Wouter J. Peyrot[46],

Andrea Pfennig[45], Giorgio Pistis[63], Shaun M. Purcell[25;144], Jorge A. Quiroz[169], Per

Qvist[12;13;55], Eline J. Regeer[170], Andreas Reif[124], Céline S. Reinbold[5;9], John P.

Rice[171], Brien P. Riley[41], Fabio Rivas[167], Margarita Rivera[1;172], Panos

Roussos[24;25;173], Douglas M. Ruderfer[174], Euijung Ryu[175], Cristina Sánchez-

Mora[59;60;62], Alan F. Schatzberg[176], William A. Scheftner[177], Robert Schoevers[178],

Nicholas J. Schork[179], Eva C. Schulte[52;180], Tatyana Shehktman[75], Ling Shen[118],

Jianxin Shi[181], Paul D. Shilling[75], Stanley I. Shyn[182], Engilbert Sigurdsson[183], Claire

Slaney[98], Olav B. Smeland[73;184;185], Johannes H. Smit[46], Daniel J. Smith[186], Janet L.

Sobell[187], Anne T. Spijker[188], Michael Steffens[189], John S. Strauss[121;190], Fabian

Streit[91], Jana Strohmaier[91], Szabolcs Szelinger[191], Katherine E. Tansey[192], Henning

Teismann[193], Alexander Teumer[194], Robert C Thompson[149], Wesley

Thompson[55;75;87;107], Pippa A. Thomson[195], Thorgeir E. Thorgeirsson[26], Matthew

Traylor[196], Jens Treutlein[91], André G. Uitterlinden[197], Daniel Umbricht[198], Helmut

Vedder[199], Alexander Viktorin[3], Peter M. Visscher[4;18], Weiqing Wang[24;25], Stanley J.

Watson[149], Bradley T. Webb[168], Cynthia Shannon Weickert[102;200], Thomas W.

Weickert[102;200], Shantel Marie Weinsheimer[55;107], Jürgen Wellmann[193], Gonneke

Willemsen[29], Stephanie H. Witt[91], Yang Wu[4], Hualin S. Xi[201], Wei Xu[202;203], Jian

Yang[4;18], Allan H. Young[204], Peter Zandi[205], Peng Zhang[206], Futao Zhang[4], Sebastian

Zollner[149], Rolf Adolfsson[31], Ingrid Agartz[14;49;207], Martin Alda[98;208], Volker Arolt[209],

Lena Backlund[95], Bernhard T. Baune[210], Frank Bellivier[211;212;213;214], Klaus Berger[193],

Wade H. Berrettini[215], Joanna M. Biernacka[175], Douglas H. R. Blackwood[30], Michael

Boehnke[90], Dorret I. Boomsma[29], Aiden Corvin[156], Nicholas Craddock[10], Mark J.

Daly[21;23], Udo Dannlowski[209], Enrico Domenici[216], Katharina Domschke[217], Tõnu

Esko[19;147;154;218], Bruno Etain[211;213;214;219], Mark Frye[220], Janice M. Fullerton[200;221],

Elliot S. Gershon[36;222], EJC de Geus[29;223], Michael Gill[156], Fernando Goes[79], Hans J.

Grabe[40], Maria Grigoroiu-Serbanescu[224], Steven P. Hamilton[225], Joanna Hauser[72],

Caroline Hayward[226], Andrew C. Heath[171], David M. Hougaard[16;43], Christina M.

Hultman[3], Ian Jones[10], Lisa A. Jones[100], René S. Kahn[25;50], Kenneth S. Kendler[41],

George Kirov[10], Stefan Kloiber[115;121;190], Mikael Landén[3;227], Marion Leboyer[117;211;228],

Glyn Lewis[17], Qingqin S. Li[229], Jolanta Lissowska[230], Susanne Lucae[115], Pamela A.

F. Madden[119], Patrik K. Magnusson[3], Nicholas G. Martin[66;231], Fermin Mayoral[167],

Susan L. McElroy[232], Andrew M. McIntosh[30;77], Francis J. McMahon[233], Ingrid

Melle[234;235], Andres Metspalu[154;236], Philip B. Mitchell[102], Gunnar Morken[237;238], Ole

Mors[16;239], Preben Bo Mortensen[12;16;32;33], Bertram Müller-Myhsok[37;240;241], Richard

M. Myers[139], Benjamin M. Neale[19;21;23], Vishwajit Nimgaonkar[242], Merete

Nordentoft[16;243], Markus M. Nöthen[7;8], Michael C. O'Donovan[10], Ketil J.

Oedegaard[244;245], Michael J. Owen[10], Sara A. Paciga[246], Carlos Pato[126;247], Michele

T. Pato[126], Nancy L. Pedersen[3], Brenda W. J. H. Penninx[46], Roy H. Perlis[248;249],

David J. Porteous[195], Danielle Posthuma[11;250], James B. Potash[79], Martin Preisig[63],

Josep Antoni Ramos-Quiroga[59;60;61;62], Marta Ribasés[59;60;62], Marcella Rietschel[91],

Guy A. Rouleau[251;252], Catherine Schaefer[118], Martin Schalling[94], Peter R.

Schofield[200;221], Thomas G. Schulze[52;79;91;97;233], Alessandro Serretti[253], Jordan W.

Smoller[21;84;85], Hreinn Stefansson[26], Kari Stefansson[26;254], Eystein Stordal[255;256],

Henning Tiemeier[80;257;258], Gustavo Turecki[259], Rudolf Uher[98], Arne E. Vaaler[260],

Eduard Vieta[261], John B. Vincent[190], Henry Völzke[194], Myrna M. Weissman[148;262],

Thomas Werge[16;107;263], Ole A. Andreassen[184;185], Anders D. Børglum[12;13;16], Sven

Cichon[5;7;9;159], Howard J. Edenberg[264], Arianna Di Florio[10;265], John Kelsoe[75],

Douglas F. Levinson[176], Cathryn M. Lewis[1;2;266], John I. Nurnberger[92;267], Roel A.

Ophoff[50;51;93], Laura J. Scott[90], Pamela Sklar[24;25†], Patrick F. Sullivan[3;265;268], Naomi

R. Wray[4;18].

References

1. Craddock N, Owen MJ (2010): The Kraepelinian dichotomy--going, going... but still not gone. *Br J Psychiatry*. 196: 92–95.

2. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005): Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 62: 593–602.

3. Weissman MM, Bland RC, Canino GJ, Faravelli C, Greenwald S, Hwu HG, *et al.* (1996): Cross-national epidemiology of major depression and bipolar disorder. *JAMA*. 276: 293–299.

4. Steel Z, Marnane C, Iranpour C, Chey T, Jackson JW, Patel V, Silove D (2014): The global prevalence of common mental disorders: a systematic review and meta-analysis 1980-2013. *Int J Epidemiol*. . doi: 10.1093/ije/dyu038.

5. Spitzer RL, Md KK, Williams JBW (1980): Diagnostic and Statistical Manual of Mental Disorders, Third Edition. *AMERICAN PSYCHIATRIC ASSOCIATION*. .

6. Reed GM, First MB, Kogan CS, Hyman SE, Gureje O, Gaebel W, *et al.* (2019): Innovations and changes in the ICD-11 classification of mental, behavioural and neurodevelopmental disorders. *World Psychiatry*. 18: 3–19.

7. American Psychiatric Association (2013): *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub.

8. Cassano GB, Rucci P, Frank E, Fagiolini A, Dell'Osso L, Shear MK, Kupfer DJ (2004): The mood spectrum in unipolar and bipolar disorder: arguments for a unitary approach. *Am J Psychiatry*. 161: 1264–1269.

9. Fiedorowicz JG, Endicott J, Leon AC, Solomon DA, Keller MB, Coryell WH (2011): Subthreshold hypomanic symptoms in progression from unipolar major depression to bipolar disorder. *Am J Psychiatry*. 168: 40–48.

10. Ratheesh A, Davey C, Hetrick S, Alvarez-Jimenez M, Voutier C, Bechdolf A, *et al.* (2017): A systematic review and meta-analysis of prospective transition from major depression to bipolar disorder. *Acta Psychiatr Scand.* 135: 273–284.

11. National Collaborating Centre for Mental Health (UK) (2012): *Common Mental Health Disorders: Identification and Pathways to Care.* Leicester (UK): British Psychological Society.

12. National Collaborating Centre for Mental Health (UK) (2018): *Bipolar Disorder: The NICE Guideline on the Assessment and Management of Bipolar Disorder in Adults, Children and Young People in Primary and Secondary Care.* Leicester (UK): British Psychological Society.

13. Polderman TJC, Benyamin B, de Leeuw CA, Sullivan PF, van Bochoven A, Visscher PM, Posthuma D (2015): Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nat Genet.* 47: 702–709.

14. McGuffin P, Rijsdijk F, Andrew M, Sham P, Katz R, Cardno A (2003): The heritability of bipolar affective disorder and the genetic relationship to unipolar depression. *Arch Gen Psychiatry.* 60: 497–502.

15. Wray NR, Ripke S, Mattheisen M, Trzaskowski M, Byrne EM, Abdellaoui A, *et al.* (2018): Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet.* 50: 668–681.

16. Stahl EA, Breen G, Forstner AJ, McQuillin A, Ripke S, Trubetskoy V, *et al.* (2019): Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat Genet.* 173062.

17. The Brainstorm Consortium, Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, *et al.* (2018): Analysis of shared heritability in common disorders of the brain. *Science.* 360: eaap8757.

18. Davis KAS, Coleman JRI, Adams M, Allen N, Breen G, Cullen B, *et al.* (2018): Mental health in UK Biobank: development, implementation and results from an online questionnaire completed by 157 366 participants. *BJPsych Open*. 4: 83–90.

19. Coleman JRI, Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium, UK Biobank Mental Health Consortium, Eley TC, Breen G (2018, January 12): Genome-wide gene-environment analyses of depression and reported lifetime traumatic experiences in UK Biobank. *bioRxiv*. .

20. Hyde CL, Nagle MW, Tian C, Chen X, Paciga SA, Wendland JR, *et al.* (2016): Identification of 15 genetic loci associated with risk of major depression in individuals of European descent. *Nat Genet*. 48: 1031–1036.

21. Gilly A, Tachmazidou I, Zeggini E (2015): Meta-analysis of summary statistics from quantitative trait association studies with unknown sample overlap. *GENETIC EPIDEMIOLOGY*. (Vol. 39), WILEY-BLACKWELL 111 RIVER ST, HOBOKEN 07030-5774, NJ USA, pp 552–553.

22. Southam L, Gilly A, Süveges D, Farmaki A-E, Schwartzentruber J, Tachmazidou I, *et al.* (2017): Whole genome sequencing and imputation in isolated populations identify genetic associations with medically-relevant complex traits. *Nat Commun*. 8: 15606.

23. Turley P, Walters RK, Maghzian O, Okbay A, Lee JJ, Fontana MA, *et al.* (2018): Multi-trait analysis of genome-wide association summary statistics using MTAG. *Nat Genet*. 118810.

24. Hill WD (2018): Comment on "Large-Scale Cognitive GWAS Meta-Analysis Reveals Tissue-Specific Neural Expression and Potential Nootropic Drug Targets" by Lam et al. *Twin Res Hum Genet*. 21: 84–88.

25. Bulik-Sullivan BK, Loh P-R, Finucane HK, Ripke S, Yang J, Schizophrenia Working Group of the Psychiatric Genomics Consortium, *et al.* (2015): LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat Genet.* 47: 291–295.

26. Nievergelt CM, Maihofer AX, Klengel T, Atkinson EG, Chen C-Y, Choi KW, *et al.* (2018, November 1): Largest genome-wide association study for PTSD identifies genetic risk loci in European and African ancestries and implicates novel biological pathways. *bioRxiv.* .

27. Shi H, Kichaev G, Pasaniuc B (2016): Contrasting the Genetic Architecture of 30 Complex Traits from Summary Association Data. *Am J Hum Genet.* 99: 139–153.

28. Shi H, Mancuso N, Spendlove S, Pasaniuc B (2017): Local Genetic Correlation Gives Insights into the Shared Genetic Architecture of Complex Traits. *Am J Hum Genet.* 101: 737–751.

29. Team RC (2015): R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2013. *Document freely available on the internet at: http://www r-project org.* .

30. Warnes GR, Bolker B, Bonebakker L, Gentleman R, Liaw WH, Lumley T, *et al.* (2016): gplots: various R programming tools for plotting data, version 3.0. 1. . Retrieved from https://CRAN.R-project.org/package=gplots.

31. Purves KL, Coleman JRI, Rayner C, Hettema JM, Deckert J, McIntosh AM, *et al.* (2017, October 16): The Common Genetic Architecture of Anxiety Disorders. *bioRxiv.* bioRxiv.

32. Schizophrenia Working Group of the Psychiatric Genomics Consortium (2014): Biological insights from 108 schizophrenia-associated genetic loci. *Nature.* 511:

421–427.

33. Demontis D, Walters RK, Martin J, Mattheisen M, Als TD, Agerbo E, *et al.* (2019): Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat Genet.* 51: 63–75.

34. Baselmans BML, Bartels M (2018): A genetic perspective on the relationship between eudaimonic -and hedonic well-being. *Sci Rep.* 8: 14610.

35. Baselmans BML, Jansen R, Ip HF, van Dongen J, Abdellaoui A, van de Weijer MP, *et al.* (2019): Multivariate genome-wide analyses of the well-being spectrum. *Nat Genet.* . doi: 10.1038/s41588-018-0320-8.

36. Choi SW, O'Reilly PF (2019): PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience.* 8. doi: 10.1093/gigascience/giz082.

37. de Leeuw CA, Mooij JM, Heskes T, Posthuma D (2015): MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol.* 11: e1004219.

38. Skene NG, Bryois J, Bakken TE, Breen G, Crowley JJ, Gaspar HA, *et al.* (2018): Genetic identification of brain cell types underlying schizophrenia. *Nat Genet.* 50: 825–833.

39. GTEx Consortium, Laboratory, Data Analysis &Coordinating Center (LDACC)—Analysis Working Group, Statistical Methods groups—Analysis Working Group, Enhancing GTEx (eGTEx) groups, NIH Common Fund, NIH/NCI, *et al.* (2017): Genetic effects on gene expression across human tissues. *Nature.* 550: 204–213.

40. Finucane HK, Reshef YA, Anttila V, Slowikowski K, Gusev A, Byrnes A, *et al.* (2018): Heritability enrichment of specifically expressed genes identifies disease-relevant tissues and cell types. *Nat Genet.* 50: 621–629.

41. Zhu Z, Zheng Z, Zhang F, Wu Y, Trzaskowski M, Maier R, *et al.* (2018): Causal

associations between risk factors and common diseases inferred from GWAS summary data. *Nat Commun.* 9: 224.

42. Yang J, Lee SH, Goddard ME, Visscher PM (2011): GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet.* 88: 76–82.

43. Savage JE, Jansen PR, Stringer S, Watanabe K, Bryois J, de Leeuw CA, *et al.* (2018): Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet.* . doi: 10.1038/s41588-018-0152-6.

44. Lee JJ, Wedow R, Okbay A, Kong E, Maghzian O, Zacher M, *et al.* (2018): Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet.* . doi: 10.1038/s41588-018-0147-3.

45. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, *et al.* (2015): Genetic studies of body mass index yield new insights for obesity biology. *Nature.* 518: 197–206.

46. Nelson CP, Goel A, Butterworth AS, Kanoni S, Webb TR, Marouli E, *et al.* (2017): Association analyses based on false discovery rate implicate new loci for coronary artery disease. *Nat Genet.* 49: 1385–1391.

47. Loh P-R, Kichaev G, Gazal S, Schoech AP, Price AL (2018): Mixed-model association for biobank-scale datasets. *Nat Genet.* 50: 906–908.

48. Howard DM, Adams MJ, Shirali M, Clarke T-K, Marioni RE, Davies G, *et al.* (2018): Genome-wide association study of depression phenotypes in UK Biobank identifies variants in excitatory synaptic pathways. *Nat Commun.* 9: 1470.

49. Li X, Luo Z, Gu C, Hall LS, McIntosh AM, Zeng Y, *et al.* (2018): Common

variants on 6q16.2, 12q24.31 and 16p13.3 are associated with major depressive disorder. *Neuropsychopharmacology*. 43: 2146–2153.

50. Okbay A, Baselmans BML, De Neve J-E, Turley P, Nivard MG, Fontana MA, *et al.* (2016): Genetic variants associated with subjective well-being, depressive symptoms, and neuroticism identified through genome-wide analyses. *Nat Genet*. 48: 624–633.

51. Amare AT, Vaez A, Hsu Y-H, Direk N, Kamali Z, Howard DM, *et al.* (2019): Bivariate genome-wide association analyses of the broad depression phenotype combined with major depressive disorder, bipolar disorder or schizophrenia reveal eight novel genetic loci for depression. *Mol Psychiatry*. 1.

52. Howard DM, Adams MJ, Clarke T-K, Hafferty JD, Gibson J, Shirali M, *et al.* (2019): Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci*. . doi: 10.1038/s41593-018-0326-7.

53. Jiang Y, Zhang H (2011): Propensity score-based nonparametric test revealing genetic variants underlying bipolar disorder. *Genet Epidemiol*. 35: 125–132.

54. Ikeda M, Takahashi A, Kamatani Y, Okahisa Y, Kunugi H, Mori N, *et al.* (2018): A genome-wide association study identifies two novel susceptibility loci and trans population polygenicity associated with bipolar disorder. *Mol Psychiatry*. 23: 639–647.

55. Psychiatric GWAS Consortium Bipolar Disorder Working Group (2011): Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat Genet*. 43: 977–983.

56. Charney AW, Ruderfer DM, Stahl EA, Moran JL, Chambert K, Belliveau RA, *et al.* (2017): Evidence for genetic heterogeneity between clinical subtypes of

bipolar disorder. *Transl Psychiatry*. 7: e993.

57. Nagel M, Jansen PR, Stringer S, Watanabe K, de Leeuw CA, Bryois J, *et al.* (2018): Meta-analysis of genome-wide association studies for neuroticism in 449,484 individuals identifies novel genetic loci and pathways. *Nat Genet.* . doi: 10.1038/s41588-018-0151-7.

58. Smith DJ, Escott-Price V, Davies G, Bailey MES, Colodro-Conde L, Ward J, *et al.* (2016): Genome-wide analysis of over 106 000 individuals identifies 9 neuroticism-associated loci. *Mol Psychiatry*. 21: 749–757.

59. Luciano M, Hagenaars SP, Davies G, Hill WD, Clarke T-K, Shirali M, *et al.* (2018): Association analysis in over 329,000 individuals identifies 116 independent variants influencing neuroticism. *Nat Genet*. 50: 6–11.

60. Li Z, Chen J, Yu H, He L, Xu Y, Zhang D, *et al.* (2017): Genome-wide association analysis identifies 30 new susceptibility loci for schizophrenia. *Nat Genet*. 49: 1576–1583.

61. Cross-Disorder Group of the Psychiatric Genomics Consortium (2013): Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*. 381: 1371.

62. Baselmans BML, van de Weijer MP, Abdellaoui A, Vink JM, Hottenga JJ, Willemsen G, *et al.* (2019): A Genetic Investigation of the Well-Being Spectrum. *Behav Genet*. 49: 286–297.

63. National Institute for Healthcare and Excellence (2009): *Depression in adults: recognition and management: Clinical guideline [CG90]*. . Retrieved from https://www.nice.org.uk/guidance/cg90.

64. Coleman JRI, Bryois J, Gaspar HA, Jansen PR, Savage JE, Skene N, *et al.* (2019): Biological annotation of genetic loci associated with intelligence in a

meta-analysis of 87,740 individuals. *Mol Psychiatry*. 24: 182–197.

65. Jansen PR, Watanabe K, Stringer S, Skene N, Bryois J, Hammerschlag AR, *et al.* (2019): Genome-wide analysis of insomnia in 1,331,010 individuals identifies new risk loci and functional pathways. *Nat Genet*. 51: 394–403.

66. Peyrot WJ, Lee SH, Milaneschi Y, Abdellaoui A, Byrne EM, Esko T, *et al.* (2015): The association between lower educational attainment and depression owing to shared genetic effects? Results in ~25,000 subjects. *Mol Psychiatry*. 20: 735–743.

67. Verbanck M, Chen C-Y, Neale B, Do R (2018): Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet*. 50: 693–698.

68. Hemani G, Bowden J, Davey Smith G (2018): Evaluating the potential role of pleiotropy in Mendelian randomization studies. *Hum Mol Genet*. . doi: 10.1093/hmg/ddy163.

69. Carney RM, Freedland KE (2017): Depression and coronary heart disease. *Nat Rev Cardiol*. 14: 145–155.

70. Weissbrod O, Flint J, Rosset S (2018): Estimating SNP-Based Heritability and Genetic Correlation in Case-Control Studies Directly and with Summary Statistics. *Am J Hum Genet*. 103: 89–99.

71. Sullivan PF, Agrawal A, Bulik CM, Andreassen OA, Børglum AD, Breen G, *et al.* (2018): Psychiatric Genomics: An Update and an Agenda. *Am J Psychiatry*. 175: 15–27.

Data availability

GWAS results from analyses including 23andMe are restricted by a data transfer agreement with 23andMe. For these analyses, LD-independent sets of 10,000 SNPs will be made available via the Psychiatric Genetics Consortium (https://www.med.unc.edu/pgc/results-and-downloads). Summary statistics not including 23andMe will be made available via the Psychiatric Genetics Consortium (https://www.med.unc.edu/pgc/results-and-downloads).