Manipulation of the social and spatial organization of the vector of

Chagas disease, *Triatoma infestans*, by the parasite *Trypanosoma*

cruzi

Short title: Behavioral manipulation of Chagas disease vector by the parasite

Stéphanie Depickère[1,#a,#b,*], Gonzalo Marcelo Ramírez-Ávila[2], Jean-Louis Deneubourg[3]

[1] Laboratorio Entomología Médica, INLASA, Rafael Zubieta N∘1889, Miraflores, La Paz, Bolivia

[2] Instituto de Investigaciones Físicas, Universidad Mayor de San Andrés, La Paz, Bolivia

[3] CENOLI, Université Libre de Bruxelles, CP 231 Bld du Triomphe, Brussels, Belgium

[#a] Current address: Instituto de Investigaciones Físicas, Universidad Mayor de San Andrés, La Paz,

Bolivia

[#b] Current address: The Abdus Salam, International Centre for Theoretical Physics (ICTP), Trieste,

Italy

24　Corresponding autor: Stéphanie Depickère E-mail: stephanie.depickere@gmail.com

Abstract

Insects of Triatominae subfamily are vectors of the parasite *Trypanosoma cruzi*, the etiological agent
of Chagas disease. Some species of Triatominae, like *Triatoma infestans* in the Southern Cone of
South America, live in the human neighborhood, aggregating in wall or roof cracks during the day and
going out to feed on animal or human blood at night. Except for their feeding specialization, these
insects share this cycle of activities where social interaction alternate with mainly individual foraging
activity with many other arthropods species. The understanding of mechanisms leading to group
cohesion is essential for understanding the spatial organization of the insects and the parasite
dispersion. Experiments with adults of *T. infestans* were conducted to study how sex and infection by
T. cruzi affect their aggregation and geotaxis behavior. Thanks to video recordings, lasting 150 min for
each experiment, the position of the insects was recorded. We show that the insects presented a high
negative geotaxis and aggregative behavior. We demonstrate that males have a higher negative
geotaxis and a higher aggregation level than females and that the infection by *T. cruzi* strengthens both
behaviors. The aggregation level and the negative geotaxis were stronger in infected insects than in
non-infected ones, the difference between sexes being maintained. The question of the influence of the
geotaxis on the bug social behavior is also asked, as all the hitherto published works about aggregation
were done in horizontal devices. The importance of these results is discussed in term of parasitic
manipulation, dispersion of the vector and strategy of its monitoring.

Author summary

Chagas disease, one of the major neglected diseases in Latin America, is caused by a parasite
transmitted by feces of hematophagous insects which live in or near the human dwellings, resting in
wall cracks during the day, and searching for food at night. Spatial organization of these insects is not
known. Thus, we conducted experiments to analyze aggregation (the act of gathering) and geotaxis
(move according to gravity) in adult insects, and to study the influence of the parasite on these two
behaviors. We demonstrate that males exhibit a higher aggregation and a stronger negative geotaxis

(moving upwards) than females and that the infection strengthens both behaviors in both sexes. These

results suggesting a non-homogeneous distribution of individuals on dwellings' walls are

epidemiologically important for the control and the monitoring of insects. They also open a new line

of research to study how these behaviors are influenced by factors like the heterogeneities of the

environment (presence of shelters), or the variability between individuals (different individual stages

together, and presence of infection).

Keywords:

Aggregation, gregariousness, Triatominae, individual behavior, geotaxis, sex differences, parasitic

manipulation, Chagas disease, *Triatoma infestans, Trypanosoma cruzi*, South America, Bolivia

Introduction

Chagas disease is one of the most important neglected tropical diseases with 6-7 million people who are estimated to be infected, and 20% of the population who are at risk [1–6]. This vector-borne disease is caused by the parasite *Trypanosoma cruzi* (Kinetoplastida: Trypanosomatidae) and is transmitted by contact with infected feces/ urine of hematophagous insects of the Triatominae subfamily (Hemiptera: Reduviidae). Currently, 149 extant species have been described worldwide, and all of them are considered as potential vectors [7]. Most of them live in sylvatic habitats, and only a dozen of species are regarded as vectors of major epidemiological importance due to their capacity to live in the surrounding of the human dwellings where they find stable shelters and food abundance (8, 9,10). *Triatoma infestans*, the subject of our study, is the main vector in the Southern Cone of South America [9]. Except for their feeding specialization, the domiciliary species share similar lifestyle and cycle of activities with many gregarious arthropods including other anthropic species like cockroaches [8,10–13]. During the daytime, they assembled in dark and sheltered places such as cracks in the walls or roof, or behind objects hanging on walls. At night, they leave their shelter to actively seek a host upon which to feed and then, they come back to a resting place to digest. The digestive phase can last from some days to several weeks according to the blood meal size, the individual and the environmental conditions [8,14].

The control strategy for Chagas disease relies mainly on the control of the domestic vectors through chemical control [1]. Face to the increase of the insecticide resistance exhibited by these insects, and to the reinvasion of dwellings by residual or sylvatic population of triatomines [15–18]; it is necessary to study the behaviors leading to a better understanding of their distribution and their dispersion. In this perspective, aggregation and geotaxis are key behaviors; knowing them better and understanding how the parasite dispersion may influence them is fundamental. Indeed, aggregation is a widespread behavior [19–23] that results from a response of individuals to environmental heterogeneity, and from social interactions involving attractions between individuals [24–30]. The social interactions maintain the group cohesion and the associated adaptive values of group living

(help in the reproduction [31,32], defense [23,33], and feeding [32,34]). In triatomines, protection

against predation is usually evoked as the main benefice of clustering, but surviving might also be

enhanced thanks to protection against hydric loss, and to a higher probability of coprophagy, symbiont

exchange, and of sex encounters, as it was showed for other insects [35–39]. Aggregation in

triatomines was studied with a focus towards the substances that mediate it, and on the factors that

modulate the aggregative response [40–48]. All these works analyzed nymphal instars behavior

response; in adults very few is known except that they can aggregate around feces [47]. Geotaxis, also

called gravitaxis, is a crucial behavior involved in insect orientation [49]. Animals can exhibit

locomotion that is gravitationally directed vertically down or up (positive or negative geotaxis,

respectively). Almost nothing is known about triatomine geotaxis, except that *T. infestans* tend to be

more concentrated in the upper half of the walls in houses or chicken houses [36,50]. Moreover, to our

knowledge, no studies were conducted to analyze the synergy or conflict between gregariousness and

geotaxis of triatomines.

It is well-known that parasites can modify physiological, behavioral, and/or morphological

traits of their hosts to increase their fitness, even if it is at the cost of the host fitness [51]. This usually

means that infected hosts will behave in ways that facilitate the transmission of the parasite [52,53].

Literature about the effects and a possible manipulation of triatomines behavior by *T. cruzi* is

relatively sparse, covering only six species: *Mepraia spinolai, Panstrongylus megistus, Rhodnius*

prolixus, Triatoma brasiliensis, Triatoma dimidiata and *T. infestans*. Authors have been especially

interested in the parasite's effects on four groups of the host's behavior: life-history trait, feeding,

defecation, and dispersion/ locomotion. It seems that *T. cruzi* increases the development time and

biting rate, and decreases the longevity and defecation time in *M. spinolai* [54,55], but no change was

observed in *P. megistus* [56], *R. prolixus* [57], *T. dimidiata* [58], *T. infestans* [59], and almost no

change in *T. brasiliensis* [60]. Reproduction was decreased by *T. cruzi* in *T. brasiliensis* [60]. The

dispersion was higher in infected females of *T. dimidiata* than in non-infected females; no effect was

found in males [61]. The locomotory activity of *R. prolixus* was decreased by infection: the total

number of movements was 20% less than that observed in non-infected insects [62]. The time to find a

host for an infected *M. spinolai* was almost twice as fast as for a non-infected insect [55]. In conclusion, modification of the triatomine traits seems to be species-dependent, age-dependent, sex-dependent, and even environment/ physiology-dependent.

In this work, experiments were conducted to study aggregation and geotaxis in adults of *T. infestans* and to analyze the effect of the infection by *T. cruzi*. Our hypotheses, based on the literature, were that these two behaviors –gregariousness and geotaxis - are strongly intertwined and are increased in infected individuals. We show that both sexes exhibit a high clustering, and a high negative geotaxis, males demonstrating a higher response than females. Interestingly, the infection by *T. cruzi* strengthens both behaviors significantly in both sexes.

Methods

T. infestans specimens were collected in dwellings from Yacuiba Municipality, Department of Tarija, in the southern region of Bolivia. The infection rate of the insects was determined by analyzing drops of feces under a light microscope. Bugs were then separated into two groups infected/ non-infected. Insects were reared at 26±1°C, 60±15% RH, 12:12 night:dark cycle, in plastic pots containing a folded piece of paper. They were fed on hens once every two weeks. Four conditions were then studied: non-infected males, non-infected females, infected males and infected females (abbreviated as NegMal, NegFem, PosMal, and PosFem in Figures and Tables). Sixteen experiments were carried out with non-infected males and females respectively, and twelve and nine experiments with infected males and females respectively. Due to experimental problems, five experiments in both sexes using non-infected insects included some infected insects, with a proportion of the infected population being less than 20%. These experiments were kept and analyzed in the same way.

Setup and Methods

A glass aquarium was used (50 x 20 x 50 cm) to avoid escaping of *T. infestans* which is unable

to climb on glass walls. As Triatominae prefer to rest on a vertical surface, the insects were allowed to

climb on one of the vertical surfaces of this aquarium (50 x 50 cm) thanks to a paper sheet (43 x

44 cm) [36,50]. The glass setup was washed, and the paper changed at the end of each experiment. It

was illuminated by a centered 60W light bulb, placed at 50 cm behind the wall covered by the paper

sheet. The paper guaranteed a homogeneous illumination of the setup. A video camera placed in front

of the setup recorded the bug activity for 150 min. A 1 m high polystyrene wall surrounded the setup

to isolate it. Experiments were conducted in a quiet and dark room to avoid any disturbance.

At the beginning of the experiment, 10 bugs (8-10 days of starvation) were dropped on the

bottom of the setup. They explored their environment rapidly and climbed on the wall. From the

recordings, a snapshot was extracted at 1 min, 5 min and then every 5 min up to 150 min (31 snapshots

in total). A processing program allowed us to record the spatial position of the thorax of each bug on

each snapshot. With these spatial coordinates, the inter-individual distances were computed. As the

length of an adult bug is on average 2.5 cm, we fixed the maximum distance of aggregation between

two individuals to 4 cm due to a tactile (legs or antennae) or visual perception.

Indexes and statistics

Several indexes of position and aggregation were calculated using processing programs: 1) the

number of individuals on the paper sheet; 2) the number of aggregated individuals; 3) the number and

the size of the clusters; 4) the spatial stability of the individual (% of individuals that were found at

time t+1 in a circle of 10mm in radius centered on the coordinate of the insect at time t); 5) the spatial

stability of the biggest cluster (study of the distance between the centroid of the biggest cluster at time

t+1 and the centroid of the biggest cluster at time t). Finally, the individual position of the insects in

the setup at the end of each experiment was analyzed, recording the vertical orientation of the bugs

(position 0: head towards the top), to put forward a privileged position. A vertical orientation was

defined as inside an angle of ±30° to the vertical, head oriented towards the top or the bottom. Outside

this range, the insect is not considered in a vertical position anymore. The structure of the clusters was

also compared between infected males and infected females, conditions where bigger clusters

emerged. Each cluster was considered as an undirected network where each node was an individual.

Links between nodes were established when the distance between them was less or equal to 4 cm (the

threshold for considering aggregation). The cluster K-density (ratio of the number of edges divided by

the number of possible edges) was compared for clusters with a size greater than three individuals.

The comparisons between conditions were made thanks to Anderson-Darling k-sample test

[63]. In case of obtaining a $p<0.05$, an Anderson-Darling all-pairs comparison test was performed.

These statistics were calculated thanks to the functions adKSampleTest() and adAllPairsTest() of the

PMCMRplus package of R [64,65]. Circular statistics were carried out with Oriana 4.02 (Kovach

Computing services). Uniformity of data was tested using Rao's test, mean comparisons using V-test,

and distribution comparisons using Mardia-Watson-Wheeler pairwise test. The structure of the clusters

was analyzed thanks to *igraph* package in R [66]. The linear regression was done thanks to lm()

function, and the comparison of the slopes of regression thanks to *lsmeans* package of R, using Least-

squared means (predicted marginal means) [67].

Results

Negative geotaxis: spatial distribution and total population

The bugs quickly climbed on the wall and stayed there, showing a high negative geotaxis; and

after an exploratory phase, the insects began to cluster and rest (Fig 1). After 10 min, more than 80%

of the individuals were on the wall (90% after 25 min) for the four conditions; and this proportion

remained constant until the end of the experiment where no statistical difference was detected between

the four conditions (Fig 2). The bugs were mostly located in the upper half of the setup; the median

vertical position reached a plateau value (stationary state) after 15 min with a value greater than 35 cm

for the four conditions (Fig 3). Their vertical distributions revealed a statistical difference between

sexes, e.g., males were located higher than females, and also between non-infected and infected

individuals, e.g., infected males were higher than non-infected males (Fig 4). These trends were also

found inside the top strip of 4 cm (40-44 cm), a zone corresponding to 10% of the total area of the

setup and where 56% (80%) of the non-infected (infected) males and 33% (44%) of the non-infected

(infected) females were located (Fig 5).

Fig 1. Examples of insect's positions at 150 min in the setup for the four conditions

Fig 2. Median fraction of bugs on the wall. Number of experiments: 16, 16, 9, 12 for NegFem

(pink), NegMal (cyan), PosFem (red) and PosMal (blue) respectively. Inserted figure: boxplot

distribution of bugs on the wall at 150 min. Anderson-Darling test at 150 min between the 4

conditions: TkN = -0.14, p = 0.47

Fig 3. Evolution of the median vertical position (quantiles 25%-75%) of the non-infected and

infected insects on the wall during the experimental time. Median position at 150 min: NegFem

(pink): 36.7 cm, PosFem (red): 39.0 cm, NegMal (cyan): 40.9 cm, PosMal (blue): 41.8 cm

Fig 4. Boxplot distribution of the vertical position of the individuals at 150 min. Number of

observations: non-infected females: 145; non-infected males: 151; infected females: 90; infected

males: 117. Anderson-Darling k-sample test between the four conditions for all individuals: TkN =

24.4, p < 0.001. Results of Anderson-Darling all-pairs comparison tests are shown at the top of the

figure (conditions with different letters correspond to conditions statistically different at p < 0.001).

Fig 5. Boxplot distribution of the median fraction of individuals at the top of the setup (40-44cm)

at 150 min. Number of observations: 16, 16, 9, 12 for NegFem, NegMal, PosFem, and PosMal

respectively. Anderson-Darling k-sample test between the four conditions for all individuals: TkN =

8.3, p < 0.001. Results of Anderson-Darling all-pairs comparison tests are shown at the top of the

figure (conditions with different letters correspond to conditions statistically different at p < 0.03)

At the end of the experiment, more than 80% of the individuals had a vertical orientation from which around 70-80% had the head turned towards the top of the setup ($\pm 30°$). The four conditions showed that individuals are not uniformly distributed (Rao's test < 0.01), but rather centered on 0 (V-tests < 0.001, Fig 6). When the distributions of individual orientations were compared, no difference appeared between sexes. Interestingly, the infection affected the orientation of the males which demonstrated a higher proportion of insects with the head towards the bottom when infected (Fig 6).

Fig 6. Vertical orientation of insects for the four conditions. Wedge's angles of 60°, frequencies are shown as radius of the wedge. 0° represents the head to the top. The dark line shows the mean direction and its length, and the standard deviation of the distribution. Number of observations: NegFem: 145, NegMal: 154, PosFem: 90 and PosMal: 117. Rao's tests gave p < 0.01 for the four conditions. V-tests (testing the null hypothesis of uniformity against non-uniform distribution with a mean of 0) gave p < 0.001 for the four conditions. Mardia-Watson-Wheeler pairwise tests between the four conditions: NegFem/ NegMal: p = 0.271, NegFem/ PosFem: p = 0.156, NegFem/ PosMal: p = 0.001, NegMal/ PosFem: p = 0.015, NegMal/ PosMal: p = 0.003, PosFem/ PosMal: p = 0.049

To summarize, without being infected, both sexes showed a high negative geotaxis that was higher for males than for females (Fig 4). Most of the insects of both sexes were oriented the head toward the top (Fig 6). The infection by *T. cruzi* strengthened this bug's geotaxis, especially for males.

Global Clustering

The median aggregated bug fraction increased up to reach a plateau around 30 min, gathering around 40% and 60% of non-infected and infected females respectively, and 70% and 90% of non-infected and infected males respectively (Fig 7). A statistical difference was observed at 150 min (end of the experiment) between sexes (males showed a higher aggregated fraction than females), and between infected conditions (infected bugs with a higher aggregated level than non-infected ones) (Figs 7 and 8). Insects in all conditions tended to gather in one or two clusters. The biggest cluster

assembled 40% (70%) of the aggregated non-infected (infected) population in males, and 30% (40%)

of the aggregated non-infected (infected) population in females (Fig 7). No difference was observed

between sexes ($p = 0.047$ and $p = 0.22$ for non-infected and infected bugs), neither between infection

conditions (Fig 7). When the structure of the clusters are compared between infected sexes, clusters of

infected males looked more compact, with a significantly smaller distance between aggregated

individuals, and they also looked denser, with a higher K-density (Fig 9 and 10).

Fig 7. Dynamics of aggregation for the four conditions: NegFem (pink), NegMal (cyan), PosFem

(red) and PosMal (blue). A: median fraction of aggregated individuals; B: median size of the biggest

cluster; C: spatial stability of the biggest cluster. Anderson-Darling k-sample test at 150 min between

the four conditions: (A) TkN = 7.4, $p < 0.001$, (B) TkN = 5.1, $p = 0.001$, and (C) TkN = -0.4, $p =$

0.573. Results of Anderson-Darling all-pairs comparison test are shown with different letters

corresponding to conditions statistically different at $p < 0.05$: (A) NegFem: a, NegMal: b, PosFem: b,

PosMal: c; (B) NegFem: a, NegMal:b, PosFem: ab, PosMal: b

Fig 8. Relative frequency of experiments according to the aggregated fraction of individuals for

the four conditions

Fig 9. Structure of the clusters in infected conditions. Boxplot of the size of edges (or links)

between aggregated bugs (left) and K-density of the clusters of size > 3 individuals (right). Anderson-

Darling k-sample test for size of the edges: TkN = 9.1, $p < 0.001$ (72 and 255 observations for PosFem

and PosMal respectively); and for the K-density: TkN = 4.7, $p = 0.004$ (17 and 21 observations for

PosFem and PosMal respectively)

Fig 10. Example of clusters of 10 bugs in infected conditions. Nodes represent the insect positions

on the wall. Links between nodes are present only when the distance is less or equal to 4 cm (the

aggregative distance)

At the end of the experiments, individuals were very stable in space: the median fraction of

individuals that moved less than 1 cm was greater than 60% for the four conditions, and no difference

between sexes and infection condition was observed (Fig 11). The biggest cluster also showed strong

spatial stability, with no statistical difference between the four conditions (Fig 7).

Fig 11. Boxplot of stable individuals (move < 10 mm between two snapshots) for the four

conditions. Anderson-Darling k-sample test between the four conditions: TkN = 2.9, p = 0.01. Results

of Anderson-Darling all-pairs comparison tests are shown at the top of the figure (conditions with

different letters correspond to conditions statistically different at p < 0.005)

In order to verify that the fraction of aggregated individuals was not directly due to the method

of calculating this fraction and the increase of the bugs density at the top of the setup, 20,000

repetitions of groups of N simulations were performed ($N = 16$ for NegFem and NegMal, $N = 9$ for

PosFem, and $N = 12$ for MalPos). For each simulation, 10 points were vertically distributed following

the experimental vertical distribution of the bugs at 150 min (Fig 12), and homogeneously horizontally

distributed. For each group of simulations, the mean fraction of aggregated individuals was calculated

for each repetition. The mean aggregated fractions found in the simulation were 0.26, 0.42, 0.36, and

0.72 for NegFem, NegMal, PosFem and PosMal respectively, revealing that an increase of the

geotaxis leads to a rise in the observed aggregation level. However, the probability of observing a

mean aggregated fraction higher or equal to the corresponding experimental one was p < 0.0001 for all

the conditions, demonstrating that the observed phenomenon involves an active aggregation due to the

inter-attraction between individuals.

Fig 12. Vertical distribution of insects on the wall for the four conditions. Fraction of bugs by

horizontal strips of 5 mm, from 0 to 440 mm, corresponding to the height of the setup for non-infected

(females: pink, males: cyan) and infected bugs (females: red, males: blue)

Fig 13 shows that the fraction of aggregated individuals in a strip of 0.5cm was proportional to

the fraction of the population settled in this strip. The slope of the regression line was the lowest for

the NegFem condition, and the highest for the PosMal condition, being intermediate and similar for

the two other conditions. The slopes of the linear regression were compared computing a model

including the interaction between the total number of bugs and the conditions: a significant interaction

was found ($F_{3, 348} = 30.04$, $p < 0.001$), giving a slope equal to 0.61 for NegFem, 0.85 for NegMal, 0.84

for PosFem, and 0.92 for PosMal. The comparison of the slopes between the four conditions showed

that NegFem's slope was lower than the three other conditions' slopes ($p < 0.001$), MalPos' slope was

higher than the three other conditions' slopes ($p < 0.02$), and MalNeg's slope was not different from

PosFem's slope ($p = 0.98$). These results demonstrate that, for the same density, the aggregation was

higher for males than for females and for infected insects than for non-infected ones.

Fig 13. Fraction of aggregated individuals according to the population present in a horizontal

strip of 5mm high for the four conditions. Linear regressions: NegFem (pink): y = 0.6096 x - 0.0019

(SE of slope: 0.0294), $R^2 = 0.83$, $p < 0.001$; NegMal (cyan): y = 0.8492 x - 0.0015 (SE of slope:

0.0169), $R^2 = 0.97$, $p < 0.001$; PosFem (red): y = 0.8403 x - 0.0017 (SE of slope: 0.0268), $R^2 = 0.92$,

$p < 0.001$; PosMal (blue): y = 0.9168 x - 0.0001 (SE of slope: 0.0075), $R^2 = 0.99$, $p < 0.001$

In conclusion, we observed an active aggregation whose level was significantly greater for

males than for females, and for infected than for non-infected insects. Clusters of infected bugs looked

denser and more compact in males than in females, reinforcing this conclusion.

Discussion

This work represents the first detailed analysis of aggregation and geotaxis in adult males and

females of *T. infestans*, and how both sexes are affected by the presence of *T. cruzi*. As shown before

in nymphal instars [43,68], adults exhibited an active aggregation due to the inter-attraction between

individuals, illustrating the social character of these insects. We show that a stable aggregation

emerges for both sexes, but the fraction of aggregated individuals and the density of clusters are higher

for males than for females. This difference between the genders is maintained under infection by *T.*

cruzi, but the latter reinforces the gregariousness in both sexes.

Our results are in agreement with previous studies. Indeed, a multi-factorial analysis (using

species, development stages and, feces source altogether) of the aggregative response of individuals to

feces shows that the aggregation level was lower (but not statistically different) for females than for

males [47]. It is well established that clustering or reduction of the inter-individual distances of social

and subsocial/ presocial arthropods reduces various stresses and therefore energy consumption

[37,69,70]. Different studies show that clustering reduces water loss [37–39,70]. We hypothesize that

the clustering of *T. infestans* individuals provides a similar benefit. As their weight is lower than

females' one, males could be under higher hydric stress. So they could tend to have a higher gregarious

trend. Moreover, it could be more adaptive for females to be less aggressive to disperse their eggs and

increase their probability of survival. In our experiments, males and females were supposed to be in

similar physiological status due to their comparable period of starvation (8-10 days), but in case of

infection, *T. cruzi* and *T. infestans* compete for nutrients, bug individuals showing a reduced resistance

to starvation when they are infected [71]. It might be speculated that infected bugs were in a stronger

starvation state and therefore were more gregarious to reduce the cost of the different stresses.

We almost know nothing about the distribution of triatomines inside a dwelling. Even if all the

stages are found in the upper part of the walls [36,50], our results suggest that males and females rest

in a stratified manner on the walls, males being above of females. Domestic cockroaches have a

similar way of life than triatomines, except the feeding habits. *Periplaneta americana*, for instance,

shows a preference for vertical areas, and males were vertically positioned above females [72]. Due to

the vertical air current, they can detect the female pheromone easily and oriented themselves towards

them. In triatomines, the sex pheromone is emitted by the metasternal glands of females, inducing

males to move towards the females (positive anemotaxis) [14]. On a wall, due to the difference in

temperature between the bottom and the top (up to 6°C [36]), an air current move towards the top of

the wall could allow the males to feel at some distance the sexual pheromone released by the females.

Most of the individuals showed a vertical orientation with the head towards the top, allowing them to

escape quickly from predators generally located below. Infection by *T. cruzi* enhanced the negative

geotaxis, especially in males, for which it was also observed a higher proportion of individuals with

the head toward the bottom.

It is interesting to note that a part of the experiments with non-infected insects was in fact

composed of a mix of non-infected and infected insects (less than 20%). One or two individuals are

not enough to observe a change at the level of the group. The latter suggests that a high proportion of

infected insects is required to change the group behavior.

Behavioral alterations upon infection are called parasitic manipulation when they are adaptive

for the parasite, altering phenotypic traits of its host in a way that enhances its probability of

transmission. There are some examples where the parasitism affects the geotaxis and the gregarious

behavior of the hosts [73–78]. Is there any advantage to *T. cruzi* to strengthen the negative geotaxis

and the aggregation behavior in males of *T. infestans*? Two hypotheses can be put forward: 1)

increasing the longevity of the vector, far from the predators present on the ground surface; that can

result from both a higher negative geotaxis and a higher aggregation level. As previously mentioned,

as aggregation could reduce the stress for the *T. infestans*, it could also increase the survival of *T.*

cruzi; 2) facilitate the host-finding. It could be easier for them to find a food source being higher on a

vertical surface. Actually, these insects are known to let themselves fall on the host [79]: the common

name of these insects in South of Latin America is "vinchuca" – meaning in Quechua "bug that lets

itself fall," i.e., from roof to reach sleeping people to feed on them. [61] have already reported an

increase of the dispersion on the field of infected females of *T. dimidiata*. Nymphal instars of *R.*

prolixus showed, on the contrary, a reduction of their locomotory activity [62].

A low height device like the setup used in these experiments allowed us to highlight

differences, between sexes, and between infected and non-infected insects. The questions that the

results generate showed that little is known about the spatial distribution of the insects in natural

conditions and how they behave. Our results predict that in a natural/anthropic environment the

percentage of infected insects should increase with the height of the settlement. More experiments

have to be conducted to understand the dispersion and aggregation behaviors of *T. infestans*, both in

the laboratory and on the field. For example one of our hypotheses concerns the influence of the height

of the setup: the higher the setup, the greater would be the spatial segregation between the four

categories. The response of the bugs should be modulated according to factors like the development

stages and the physiological condition of the insects, the bug density, the numbers of available and

suitable shelters, and the infection of the bugs. One important question is the robustness of our results

in front of the presence of shelters (small cavities). Similarly, another question is about how mixed

groups would distribute (both sexes, and/ or both infected and non-infected bugs).

Our results also open a new vision for controlling/ monitoring triatomines on the field, letting

appear a higher risk coming from the upper part of the rooms. More precise studies about bug

distribution on the field should help to improve control and monitoring by trapping [80]. Finally, our

results lead us to propose simple tests easily feasible in the field, based on the geotaxis and the

aggregative behavior of the bugs, to detect infected insects.

Acknowledgments

417   S. Depickère is particularly grateful to T. Chavez, F. Lardeux, Don Hugo and E. Siñani for logistic

418   support.

References

1.   WHO. Chagas disease (American trypanosomiasis) [Internet]. World Health Organization Fact
Sheets. 2018. p. 1.
2.   Clayton J. Chagas disease: pushing through the pipeline. Nature. 2010; 465(7301): S12-15.

3. Rodrigues Coura J, Albajar Viñas P. Chagas disease: a new worldwide challenge. Nature.
2010; 465(7301): S6-7.

4. Rassi AJ, Rassi A, Marin-Neto JA. Chagas disease. Lancet. 2010; 375(9723): 1388–402. DOI:
10.1016/S0140-6736(10)60061-X

5. Lee BY, Bacon KM, Bottazzi ME, Hotez PJ. Global economic burden of Chagas disease: A
computational simulation model. Lancet Infect Dis. 2013; 13(4): 342–8. DOI: 10.1016/S1473-
3099(13)70002-1

6. Hotez PJ, Bottazzi ME, Franco-Paredes C, Ault SK, Periago MR. The neglected tropical
diseases of Latin America and the Caribbean: A review of disease burden and distribution and
a roadmap for control and elimination. PLoS Negl Trop Dis. 2008; 2(9): e300. DOI:
10.1371/journal.pntd.0000300

7. Justi SA, Galvão C. The Evolutionary origin of diversity in Chagas disease vectors. Trends
Parasitol. 2017; 33(1): 42–52. DOI: 10.1016/j.pt.2016.11.002

8. Mullen GR, Durden LA. Medical and Veterinary Entomology. Vol. Second Rev. Oxford:
Academic Press Inc.; 2009. 637 p.

9. Flores-Ferrer A, Marcou O, Waleckx E, Dumonteil E, Gourbière S. Evolutionary ecology of
Chagas disease; what do we know and what do we need? Evol Appl. 2017; 11(4): 470–87.
DOI: 10.1111/eva.12582

10. Gillott C. Entomology. Third Edition. 2005. 834 p.

11. Costa JT. The other insect societies. Belknap Press of Harvard University Press; 2006. 767 p.

12. Crall JD, Souffrant AD, Akandwanaho D, Hescock SD, Callan SE, Coronado WM, et al. Social
context modulates idiosyncrasy of behaviour in the gregarious cockroach *Blaberus discoidalis*.
Anim Behav. 2016; 111: 297–305. DOI: 10.1016/j.anbehav.2015.10.032

13. Varadínová Z, Stejskal V, Frynta D. Patterns of aggregation behaviour in six species of
cockroach: comparing two experimental approaches. Entomol Exp Appl. 2010; 136(2): 184–
90. DOI: 10.1111/j.1570-7458.2010.01016.x

14. Lazzari CR, Pereira MH, Lorenzo MG. Behavioural biology of Chagas disease vectors. Mem
Inst Oswaldo Cruz. 2013; 108(Suppl 1): 34–47. DOI: 10.1590/0074-0276130409

15. Lardeux F, Depickère S, Duchon S, Chavez T. Insecticide resistance of *Triatoma infestans*
(Hemiptera, Reduviidae) vector of Chagas disease in Bolivia. Trop Med Int Heal. 2010; 15(9):
1037–48.

16. Germano MD, Roca Acevedo G, Mougabure Cueto GA, Toloza AC, Vassena C V., Picollo MI.
New Findings of Insecticide Resistance in *Triatoma infestans* (Heteroptera: Reduviidae) from
the Gran Chaco. J Med Entomol. 2010; 47(6): 1077–81. DOI: 10.1603/ME10069

17. Germano MD, Picollo MI, Mougabure-cueto GA. Microgeographical study of insecticide
resistance in *Triatoma infestans* from Argentina. Acta Trop. 2013; 128(3): 561–5. DOI:
10.1016/j.actatropica.2013.08.007

18. Depickère S, Buitrago R, Siñani E, Baune M, Monje M, Lopez R, et al. Susceptibility and
resistance to deltamethrin of wild and domestic populations of *Triatoma infestans* (Reduviidae:
Triatominae) in Bolivia: new discoveries. Mem Inst Oswaldo Cruz. 2012 Dec; 107(8): 1042–7.
DOI: 10.1590/S0074-02762012000800013

19. Allee WC. Animals aggregations: a study in general sociology. Chicago: University of Chicago
Press; 1931. 431 p.

20. Parrish JK, Edelstein-Keshet L. Complexity, pattern, and evolutionary trade-offs in animal
aggregation. Science. 1999; 284(5411): 99–101. DOI: 10.1126/science.284.5411.99

21. Allee WC. Animal aggregations. Q Rev Biol. 1927; 2: 367–98.

22. Camazine S, Deneubourg J-L, Franks NR, Sneyd J, Bonabeau E, Theraulaz G. Self-

Organization in Biological Systems. Princeton, editor. New Jersey: Princeton University Press;
2001. 538 p.

23.   Krause J, Ruxton GD. Living in groups. Oxford: Oxford University Press; 2002. 224 p.

24.   Fraenkel GS, Gunn DL. The Orientation of Animals. Kineses, taxes and compass reactions.
New-York: Dover, New York; 1961. 376 p.

25.   Depickère S, Fresneau D, Deneubourg J-L. A basis for spatial and social patterns in ant
species: dynamics and mechanisms of aggregation. J Insect Behav. 2004; 17(1): 81–97.

26.   Depickère S, Fresneau D, Deneubourg J-L. The influence of red light on the aggregation of two
castes of the ant, *Lasius niger*. J Insect Physiol. 2004; 50(7): 629–35.

27.   Jeanson R, Rivault C, Deneubourg J-L, Blanco S, Fournier R, Jost C, et al. Self-organized
aggregation in cockroaches. Anim Behav. 2005; 69: 169–80.

28.   Lorenzo MG, Lazzari CR. Temperature and relative humidity affect the selection of shelters by
Triatoma infestans, vector of Chagas disease. Acta Trop. 1999; 72: 241–9.

29.   Schilman PE, Lazzari CR. Temperature preference in *Rhodnius prolixus*, effects and possible
consequences. Acta Trop. 2004; 90(1): 115–22.

30.   Depickère S, Fresneau D, Deneubourg J-L. Dynamics of aggregation in *Lasius niger*
(Formicidae): influence of polyethism. Insectes Soc. 2004; 51(1): 81–90.

31.   Lloyd JE. Fireflies of Melanesia: Bioluminescence, mating behavior, and synchronous flashing
(Coleoptera: Lampyridae). Environ Entomol. 1973; 2: 991–1008.

32.   Wilson EO. Sociobiology: the new synthesis. Cambridge, Massachusetts: The Belknap Press of
Harvard University Press; 2000. 697 p.

33.   Millor J, Pham-Delegue M, Deneubourg J-L, Camazine S. Self-organized defensive behavior
in honeybees. Proc Natl Acad Sci USA. 1999; 96(22): 12611–5.

34.   Hölldobler B, Wilson EO. The Ants. Cambridge, Massachussetts: The Belknap Press of
Harvard University Press; 1990.

35.   Beard C Ben, Cordon-Rosales C, Durvasula R V. Bacterial symbionts of the Triatominae and
their potential use in control of Chagas disease transmission. Annu Rev Entomol. 2002; 47:
123–41.

36.   Schofield CJ. The behaviour of Triatominae (Hemiptera: Reduviidae): a review. Bull Entomol
Res. 1979; 69(3): 363–79.

37.   Dambach M, Goehlen B. Aggregation density and longevity correlate with humidity in first-
instar nymphs of the cockroach (*Blattella germanica* L., Dictyoptera). J Insect Physiol. 1999;
45(5): 423–9.

38.   Yoder JA, Grojean NC. Group influence on water conservation in the giant Madagascar
hissing-cockroach, *Gromphadorhina portentosa* (Dictyoptera: Blaberidae). Physiol Entomol.
1997; 22: 79–82.

39.   Yoder JA, Hobbs HH, Hazelton MC. Aggregate protection against dehydration in adult females
of the cave cricket, *Hadenoecus cumberlandicus* (Orthoptera, rhaphidophoridae). J Cave Karst
Stud. 2002; 64(2): 140–4.

40.   Vitta ACR, Lorenzo Figueiras AN, Lazzari CR, Diotaiuti L, Lorenzo MG. Aggregation
mediated by faeces and footprints in *Triatoma pseudomaculata* (Heteroptera: Reduviidae), a
Chagas disease vector. Mem Inst Oswaldo Cruz. 2002; 97(6): 865–7.

41.   Lorenzo Figueiras AN, Lazzari CR. Aggregation behaviour and interspecific responses in
Rhodnius prolixus Stal. Mem Inst Oswaldo Cruz. 2002; 97(4): 569–71.

42.   Falvo ML, Lorenzo Figueiras AN, Manrique G. Spatio-temporal analysis of the role of faecal
depositions in aggregation behaviour of the triatomine *Rhodnius prolixus*. Physiol Entomol.
2015; 41: 24–30. DOI: 10.1111/phen.12120

43. Lorenzo Figueiras AN, Kenigsten A, Lazzari CR. Aggregation in the haematophagous bug
Triatoma infestans: chemical signals and temporal pattern. J Insect Physiol. 1994; 40(4): 311–
6.

44. Lorenzo Figueiras AN, Lazzari CR. Aggregation behaviour and interspecific responses in three
species of Triatominae. Mem Inst Oswaldo Cruz. 1998; 93(1): 133–7.

45. Reisenman CE, Lorenzo Figueiras AN, Giurfa M, Lazzari CR. Interaction of visual and
olfactory cues in the aggregation behaviour of the haematophagous bug *Triatoma infestans*. J
Comp Physiol A. 2000; 186: 961–8.

46. Rocha Pires HH, Lorenzo MG, Diotaiuti L, Lazzari CR, Lorenzo Figueiras AN, Pires HHR, et
al. Aggregation behaviour in *Panstrongylus megistus* and *Triatoma infestans*: inter and
intraspecific responses. Acta Trop. 2002; 81: 47–52.

47. Cruz-Lopez L, Malo EA, Rojas JC. Aggregation pheromone in five species of Triatominae
(Hemiptera: Reduviidae). Mem Inst Oswaldo Cruz. 1993; 88(4): 535–9.

48. Minoli SA, Baraballe S, Lorenzo Figueiras AN. Daily rhythm of aggregation in the
haematophagous bug *Triatoma infestans* (Heteroptera: Reduviidae). Mem Inst Oswaldo Cruz.
2007; 102(4): 449–54. DOI: 10.1590/S0074-02762007005000033

49. Wigglesworth VB. The principles of insect physiology. 7th editio. New York, NY, USA:
Chapman & Hall; 1972. 887 p. DOI: 10.1007/978-94-009-5973-6

50. Wisnivesky-Colli C, Pietrokovsky S, Bottazzi V, Haedo A, Solarz N, Schweigmann N, et al.
Laboratory comparison of feeding success among *Triatoma infestans, T. guasayana,* and *T.
sordida* (Hemiptera : Reduviidae). J Med Entomol. 1995; 32(5): 583–7.

51. Van Houte S, Ros VI, Van Oers MM. Walking with insects: molecular mechanisms behind
parasitic manipulation of host behaviour. Mol Ecol. 2013; 22: 3458–75. DOI:
10.1111/mec.12307

52. Hughes DP, Brodeur J, Thomas F, editors. Host Manipulation by Parasites. Oxford University
Press; 2012. 232 p.

53. Hurd H. Manipulation of medically important insect vectors by their parasites. Annu Rev
Entomol. 2003; 48(1): 141–61. DOI: 10.1146/annurev.ento.48.091801.112722

54. Botto-Mahan C. *Trypanosoma cruzi* induces life-history trait changes in the wild kissing bug
Mepraia spinolai: implications for parasite transmission. Vector-Borne Zoonotic Dis. 2009;
9(5): 505–10. DOI: 10.1089/vbz.2008.0003

55. Botto-Mahan C, Cattan PE, Medel R. Chagas disease parasite induces behavioural changes in
the kissing bug *Mepraia spinolai*. Acta Trop. 2006; 98: 219–23.

56. Lima MM, Borges-Pereira J, Albuquerque Dos Santos JA, Teixeira Pinto Z, Vianna Braga M.
Development and reproduction of *Panstrongylus megistus* (Hemiptera: Reduviidae) infected
with *Trypanosoma cruzi*, under laboratory conditions. Ann Entomol Soc Am. 1992; 85(4):
458–61. DOI: 10.1093/aesa/85.4.458

57. Takano-Lee M, Edman JD. Lack of manipulation of *Rhodnius prolixus* (Hemiptera:
Reduviidae) vector competence by *Trypanosoma cruzi*. J Med Entomol. 2002; 39(1): 44–51.
DOI: 10.1603/0022-2585-39.1.44

58. Zeledón R. El *Triatoma dimidiata* (Latreille, 1811) y su relación con la enfermedad de Chagas.
San José (Costa Rica): UNED; 1981. 146 p.

59. Schaub GA. Developmental time and mortality of larvae of *Triatoma infestans* infected with
Trypanosoma cruzi. Trans R Soc Trop Med Hyg. 1988; 82: 94–7.

60. Oliveira TG, Carvalho-Costa FA, Gomes TF, Sarquis O, Sposina R, Lima MM. Developmental
and reproductive patterns of *Triatoma brasiliensis* infected with *Trypanosoma cruzi* under
laboratory conditions. Mem Inst Oswaldo Cruz. 2010; 105(8): 1057–60.

61. Ramirez-Sierra MJ, Herrera-Aguilar M, Gourbière S, Dumonteil E. Patterns of house

infestation dynamic by non-domiciliated *Triatoma dimidiata* reveal a spatial gradient of
infestation in rural villages and potential insect manipulation by *Trypanosoma cruzi*. Trop Med
Int Heal. 2010; 15(1): 77–86.

62.   Marliére NP, Latorre-Estivalis JM, Lorenzo MG, Carrasco D, Alves-Silva J, Rodrigues J de O,
et al. Trypanosomes modify the behavior of their insect hosts: effects on locomotion and on the
expression of a related gene. PLoS Negl Trop Dis. 2015; 9(8): e0003973. DOI:
10.1371/journal.pntd.0003973

63.   Scholz F, Stephens M. K-Sample Anderson-Darling Tests. J Am Stat Assoc. 1987; 82: 918–24.

64.   R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2016.

65.   Pohlert T. PMCMRplus: calculate pairwise multiple comparisons of mean rank sums extended.
R Packag version 101. 2018; .

66.   Csardi G, Nepusz T. The igraph software package for complex network research. InterJournal.
2006; Complex Sy: 1695.

67.   Lenth R V. Least-Squares Means: the R package lsmeans. J Stat Softw. 2016; 69(1): 1:33.
DOI: 10.18637/jss.v069.i01

68.   Lorenzo Figueiras AN, Lazzari C. Aggregation behavior and interspecific responses in three
species of Triatominae. Mem Inst Oswaldo Cruz. 1998; 93(1): 133–7.

69.   Tanaka S, Wolda H, Denlinger DL. Group size affects the metabolic rate of a tropical beetle.
Physiol Entomol. 1988; 13: 239–41.

70.   Benoit JB, Lopez-Martinez G, Michaud MR, Elnitsky MA, Lee RE, Denlinger DL.
Mechanisms to reduce dehydration stress in larvae of the Antarctic midge, *Belgica antarctica*. J
Insect Physiol. 2007; 53(7): 656–67. DOI: 10.1016/j.jinsphys.2007.04.006

71.   Schaub GA. Does *Trypanosoma cruzi* stress its vectors? Parasitol Today. 1989; 5(6): 185–8.

72.   Silverman JM, Bell WJ. The role of vertical and horizontal object orientation in mate-finding
and predator-avoidance by the american cockroach. Anim Behav. 1979; 27: 652–7.

73.   Hughes DP, Kathirithamby J, Turillazzi S, Beani L. Social wasps desert the colony and
aggregate outside if parasitized: parasite manipulation? Behav Ecol. 2004; 15(6): 1037–43.

74.   Lefèvre T, Thomas F. Behind the scene, something else is pulling the strings: Emphasizing
parasitic manipulation in vector-borne diseases. Infect Genet Evol. 2008; 8: 504–19. DOI:
10.1016/j.meegid.2007.05.008

75.   Jacquin L, Mori Q, Steffen M, Medoc V. Non-specific manipulation of gammarid behaviour by
P . minutus parasite enhances their predation by definitive bird hosts. PLoS One. 2014; 9(7):
e101684. DOI: 10.1371/journal.pone.0101684

76.   Arnal A, Droit A, Elguero E, Ducasse H, Sánchez MI, Levèvre T, et al. Activity level and
aggregation behavior in the crustacean gammarid *Gammarus insensibilis* parasitized by the
manipulative trematode *Microphallus papillorobustus*. Front Ecol Evol. 2015; 3: 109. DOI:
10.3389/fevo.2015.00109

77.   Weinersmith KL, Brown CE, Clingen KB, Jacobsen MC, Topper LB, Hechinger RF.
Euhaplorchis californiensis cercariae exhibit positive phototaxis and negative geotaxis. J
Parasitol. 2018; 104(3): 329–33. DOI: 10.1645/17-80

78.   Thomas F, Adamo S, Moore J. Parasitic manipulation: where are we and where should we go?
Behav Processes. 2005; 68: 185–99. DOI: 10.1016/j.beproc.2004.06.010

79.   Lazzari CR, Lorenzo MG. Exploiting triatomine behaviour: alternative perspectives for their
control. Mem Inst Oswaldo Cruz. 2009; 104(Suppl. I): 65–70.

80.   Rojas de Arias A, Abad-Franch F, Acosta N, López E, González N, Zerba E, et al. Post-control
surveillance of *Triatoma infestans* and *Triatoma sordida* with chemically-baited sticky traps.

PLoS Negl Trop Dis. 2012; 6(9): e1822. DOI: 10.1371/journal.pntd.0001822

Bug fraction on the wall at 150min

NegFem

NegMal

PosFem

PosMal