1   Dissection of complex, fitness-related traits in multiple *Drosophila* mapping
2   populations offers insight into the genetic control of stress resistance
3
4

5   Elizabeth R. Everman, Casey L. McNeil, Jennifer L. Hackett, Clint L. Bain, Stuart J. Macdonald
6
7   Department of Molecular Biosciences, University of Kansas, 1200 Sunnyside Avenue, Lawrence,
8   Kansas 66045, USA.

28 Running title: Genetic dissection of fitness traits
29
30 Keywords: Starvation resistance, triglyceride level, DSPR, DGRP, MPP
31

32 Corresponding author:
33
Stuart Macdonald
Department of Molecular Biosciences
4043 Haworth Hall
1200 Sunnyside Avenue
University of Kansas
Lawrence, KS 66045
40
41 Tel: 785-864-5362
42 Email: sjmac@ku.edu

Abstract

We leverage two complementary *Drosophila melanogaster* mapping panels to genetically dissect starvation resistance, an important fitness trait. Using >1600 genotypes of the multiparental *Drosophila* Synthetic Population Resource (DSPR) we map numerous starvation stress QTL that collectively explain a substantial fraction of trait heritability. QTL effects further allowed us to estimate DSPR founder phenotypes, predictions that were correlated with the actual founder phenotypes. Starvation resistance has been linked to triglyceride level, and while we observe a modest phenotypic correlation between the traits in the DSPR, overlap among the QTL identified for each trait is low. Since we show that DSPR strains with extreme starvation phenotypes also differ in desiccation resistance and activity level, our data imply that multiple physiological mechanisms contribute to starvation variability. We also exploit the *Drosophila* Genetic Reference Panel (DGRP), identifying a number of sequence variants associated with starvation resistance. Consistent with prior work these sites rarely fall within QTL intervals mapped in the DSPR. Two other groups previously measured starvation resistance in the DGRP, offering a unique opportunity to directly compare mapping results across labs. We found strong phenotypic correlations among studies, but extremely low overlap in the sets of genomewide significant sites. Despite this, our analyses revealed that the most highly-associated variants from each study typically showed the same additive effect sign in independent studies, in contrast to otherwise equivalent sets of random variants. This consistency provides evidence for reproducible trait-associated sites in a widely-used mapping panel, and highlights the polygenic nature of starvation resistance.

Introduction

Starvation resistance is a classic quantitative fitness-related trait that is associated with several other phenotypes that influence survival, lifespan, and reproduction (Murphey and Hall 1969; Service and Rose 1985; Da Lage *et al.* 1990; Rose *et al.* 1992; Toda and Kimura 1997; Karan and Parkash 1998; Hoffmann *et al.* 2005b; Sørensen *et al.* 2007; Telonis-Scott *et al.* 2009). In particular, increased starvation resistance is often negatively correlated with fecundity and positively correlated with longevity, energy storage, and other stress response traits (Service *et al.* 1985; Rose *et al.* 1992; Hoffmann and Parsons 1993; Chippindale *et al.* 1993; Harshman *et al.* 1999; Bochdanovits and de Jong 2003; Bubliy and Loeschcke 2005; Sørensen *et al.* 2007; Schwasinger-Schmidt *et al.* 2012). Because of pervasive phenotypic and genetic correlations between starvation resistance and other traits, the evolution of starvation resistance in natural populations is complex and population-specific, and is driven by adaptation to heterogeneous environments, phenotypic plasticity, and extensive pleiotropy (Service and Rose 1985; Hoffmann and Parsons 1991; Chippindale *et al.* 1993; Falconer and Mackay 1996; Toda and Kimura 1997; Karan and Parkash 1998; Harshman *et al.* 1999; Harbison *et al.* 2004; Pijpe *et al.* 2007; Rion and Kawecki 2007; Bauerfeind *et al.* 2014; Colinet *et al.* 2015; Everman and Morgan 2018).

Artificial selection for increased starvation resistance often results in a corresponding increase in lipid levels in *D. melanogaster* (Chippindale *et al.* 1996, 1998; Djawdan *et al.* 1998; Harshman *et al.* 1999; Goenaga *et al.* 2013; Hardy *et al.* 2018), suggesting that energy storage is one important mechanism that contributes to starvation resistance. However, variation in the association between these traits has also been observed (Hoffmann *et al.* 2001; Schwasinger-Schmidt *et al.* 2012). For example, while Chippindale *et al.* (1996) provided evidence of a strong positive correlation between starvation resistance and lipid concentration following 60 generations of selection for starvation resistance, Hoffmann *et al.* (2001) found that total lipid concentration and starvation resistance in isofemale lines derived from natural populations were not correlated. The correlation between starvation resistance and lipid level is likely dependent upon genetic background and the evolutionary history of a population, resulting in across-population variation in the strength and direction of the correlation between these traits. Furthermore, starvation resistance is likely influenced by physiological factors in addition to lipid

level, including metabolic rate (Hoffmann and Parsons 1993; Rion and Kawecki 2007;
Schwasinger-Schmidt *et al.* 2012) and differences in overall activity level.

Similar to lipid levels, selection for starvation resistance often results in a concomitant
change in desiccation resistance (Hoffmann and Parsons 1989; Chippindale *et al.* 1996; Hoffmann
and Harshman 1999; Harshman *et al.* 1999; Hoffmann *et al.* 2001). Nonetheless, surveys of
natural populations in several *Drosophila* species have shown that starvation and desiccation
resistance can also independently vary (Davidson 1990; Karan and Parkash 1998; Karan *et al.*
1998; Chippindale *et al.* 1998; Hoffmann and Harshman 1999; Gilchrist *et al.* 2008; Goenaga *et*
al. 2013). In addition, opposing clines in starvation and desiccation resistance were observed
across tropical and temperate climates in several *Drosophila* species (reviewed in Hoffmann and
Harshman 1999). Given these variable patterns, Karan and Parkash (1998) and Da Lage *et al.*
(1990) suggest that desiccation and starvation resistance are not physiologically constrained.
Rather, they are both directly and indirectly influenced by climate variability and selection on
other correlated traits such as diapause or thermal tolerance in seasonally variable temperate
environments (Schmidt *et al.* 2005; Sørensen *et al.* 2007; Rion and Kawecki 2007; Goenaga *et al.*
2013).

Genetic dissection of starvation resistance using diverse mapping panels can facilitate a
deeper understanding of the evolution of, and mechanisms underlying, starvation resistance and
its phenotypic association with other traits such as lipid level and desiccation resistance. Previous
studies have examined the genetic basis of starvation resistance in *D. melanogaster* using a
combination of selection experiments (Rose *et al.* 1992; Chippindale *et al.* 1996; Harshman *et al.*
1999; Bochdanovits and de Jong 2003; Bubliy and Loeschcke 2005; Schwasinger-Schmidt *et al.*
2012), gene expression studies following exposure to starvation stress (Harbison *et al.* 2005;
Sørensen *et al.* 2007), and genetic mapping (Harbison *et al.* 2004; Mackay *et al.* 2012; Huang *et*
al. 2014; Everman and Morgan 2018). These studies have provided extensive lists of candidate
genes, some of which have been functionally validated (Lin *et al.* 1998; Clancy *et al.* 2001;
Harbison *et al.* 2004, 2005; Nuzhdin *et al.* 2005; Sørensen *et al.* 2007). However, up to this point
few studies have undertaken an examination of the genetic architecture of triglyceride or lipid
content in the same genetically diverse panel used to examine variation in starvation resistance.

Doing so would allow a detailed comparison of quantitative trait loci (QTL) that contribute to

variation in each trait, provide important insight into the similarity of the genetic architectures

of starvation resistance and correlated traits, and facilitate a better understanding of the

coevolutionary dynamics underlying these traits.

In this study, we used two powerful, complementary *D. melanogaster* mapping panels -

the *Drosophila* Synthetic Population Resource (DSPR) and the *Drosophila* Genetic Reference

Panel (DGRP) - to genetically dissect starvation resistance. In addition to mapping loci that

contribute to variation in this trait, we measured and mapped triglyceride level in the DSPR. Our

approach allows us to accomplish three primary objectives. First, by measuring starvation

resistance and triglyceride level in the same mapping panel, we can assess overlap in the loci that

contribute to variation in each trait, thus gaining insight into the level of pleiotropy between

these traits. Second, by measuring starvation resistance in both the DSPR and the DGRP, we are

able to address variation in the genetic architecture of this trait between two populations that

have unique evolutionary histories. Finally, one advantage of a stable reference panel of inbred

lines is the ability to repeatedly measure trait variation on the same set of genotypes. Here, in

addition to our new screen, we leveraged two previous starvation resistance screens using the

DGRP (Mackay *et al.* 2012; Everman and Morgan 2018) to resolve SNP variants with consistent,

across-study phenotypic effects. This across-study comparison of the genetic architecture of

starvation resistance in the DGRP provides both technical insight into the use of genomewide

association studies (GWAS) to understand the genetic basis of complex traits, and biological

insight into the phenotypic effects of loci that contribute to trait variation.

Overall, despite a significant phenotypic correlation, we report limited overlap among

mapped loci that contribute to variation in starvation resistance and triglyceride level in the

DSPR, suggesting that the genetic basis of these traits is largely independent. This highlights the

role other physiological mechanisms, such as activity level and desiccation resistance that we

explore here, may have in influencing starvation resistance. In common with previous studies

using both the DSPR and DGRP, we also find little overlap in the loci associated with starvation

resistance between mapping panels, likely the combined result of unique genetic backgrounds

and evolutionary histories, as well as differences in power to detect causative loci. A comparison

of the genetic architecture of starvation resistance across three DGRP datasets supports this

interpretation. We found that the sign of the additive effects of the most strongly-associated

SNPs were consistent across datasets, suggesting these SNPs likely contribute to variation in

starvation resistance in the DGRP, but have sufficiently small effects that they are regularly not

identified following genomewide multiple testing correction.

Materials and Methods
158
Mapping populations

Drosophila *Synthetic Population Resource*

The DSPR is a multiparental population that consists of two synthetic populations (pA and

pB) that were each established following an intercross of eight highly-inbred founder lines, with

one founder line shared between the two populations (King *et al.* 2012a). Flies were maintained

in pairs of subpopulations (pA1, pA2, pB1, pB2) at high population density for 50 generations

prior to the establishment of > 1600 genotyped recombinant inbred lines (RILs) generated via 25

generations of full-sib inbreeding (King *et al.* 2012a; b). Founder lines for the pA and pB panels

have also been sequenced at 50x coverage, enabling inference of the haplotype structure of each

RIL via a hidden Markov model (described in King *et al.* (2012a).

Drosophila *Genetic Reference Panel*

The DGRP was established from mated females collected from a natural population in

Raleigh, North Carolina, with inbred lines derived from 20 generations of full-sib mating (Mackay

et al. 2012). Each of the 205 DGRP lines have been fully re-sequenced and genotyped allowing

GWA mapping to be carried out in the panel (Mackay *et al.* 2012; Huang *et al.* 2014).

Phenotyping assays and analysis

Large-scale starvation resistance assay

Strains from the DSPR and DGRP were duplicated from stocks, and adults were discarded

following 1-2 days of oviposition to maintain a relatively even egg density across all experimental

vials. In the following generation, experimental flies (2-4 days old) were sorted by sex over light

CO_2 anesthesia and placed in groups of same-sex individuals on new cornmeal-molasses-yeast

media for 1 day until the start of the starvation assay. The assay was initiated by placing flies on

1% agar media that additionally contained propionic acid and benzoic acid preservatives.

Starvation media was made within 24 hours of the initiation of each block of the assay and was

not replaced throughout its duration. In both mapping panels, survival was measured across 2

vial replicates per sex in ~85% of strains, with ~90% of vials containing 10 flies (minimum flies per

vial = 6). Some strains were replicated 3-4 times for a maximum of 40 flies screened per strain.

Vials were barcoded during the screen, blinding experimenters to strain identification number,

and assisting with efficient data collection and analysis.

Death was assessed for each vial twice per day at approximately 0900 and 2100 hrs. The

first assessment of survival was made 24 hours after flies were transferred to starvation media.

Dead flies at this initial assessment point were not included in the analysis as their death may

have resulted from handling during the initial transfer to experimental vials rather than

starvation stress. Vials with flies that had become entangled in the cotton vial plug at any point

in the duration of the assay were also excluded from the analysis. Flies were considered dead if

they were not moving or unable to dislodge themselves from the starvation media. The

phenotype used for mapping was the mean time to death in hours per strain across the vial

replicates.

We screened the DSPR (861 pA1/pA2 and 864 pB1/pB2 RILs) in a series of batches across

a seven-month period in 2010. Each batch included the majority of RILs that belonged to a

particular subpopulation. Starvation resistance was measured in 168 DGRP lines in a single batch

in 2012. Finally, we measured starvation resistance in the 15 DSPR founder lines, using 5 vial

replicates per founder, in one batch. In all cases flies were reared and tested at approximately

23°C, 30-60% humidity, with constant light.

We assessed variation in starvation resistance due to mapping subpopulation and sex in

the DSPR with a two-way ANOVA, including the interaction, and treated subpopulation (pA1, pA2,

pB1, and pB2) and sex as fixed factors. Male and female-specific differences among the four

subpopulations were tested using Tukey's HSD post hoc comparisons with an experiment-wide

α = 0.05. Differences in starvation resistance due to sex among the DGRP lines were analyzed

with a one-way ANOVA, treating sex as a fixed factor.

Desiccation resistance assay

To investigate the correlation between starvation and desiccation resistance, we

measured desiccation resistance in a subset of pA1/pA2 RILs that had very low (17 RILs) or very

high (16 RILs) average female starvation resistance in the large-scale screen. Desiccation

resistance was assessed in females only with two vial replicates per RIL, 92.9% of which contained

10 flies (minimum flies per vial = 8). We placed experimental flies in empty vials plugged with

cotton inside an airtight container. Relative humidity was reduced to close to 0% by adding a

large quantity of Drierite (calcium sulfate) to the chamber. Survival was assessed every hour

following initiation of the experiment, and mean desiccation resistance per RIL was used in all

analyses.

Activity monitor starvation resistance assay

The *Drosophila* Activity Monitoring System (DAM2, TriKinetics, Inc.) was used to measure

starvation resistance and activity level prior to starvation under environmental conditions similar

to those used in previous starvation resistance studies in *D. melanogaster* (Mackay *et al.* 2012;

Everman and Morgan 2018). For this experiment we selected a subset of pA1/pA2 RILs from the

two tails of the distribution of female starvation resistance measured in the large-scale screen

described above, testing 16 flies of each sex from 16 (19) lines with high (low) female starvation

resistance.

One day prior to adding flies to monitor tubes, cornmeal-yeast-dextrose media was

poured into 100mm diameter petri dishes. Once set, polycarbonate activity monitor tubes (5mm

diameter x 65mm length) were filled to approximately 10mm by pushing them into the media.

The food plug in each tube was sealed with paraffin wax. A single fly was aspirated into each

tube, and the tubes were capped with small pieces of Droso-Plugs (Genesee Scientific

Corporation). Flies were allowed to acclimate to the tubes and incubator conditions for 24 hours

at 25°C and ~50% relative humidity with a 12:12 hr photoperiod. Following the acclimation

period, we measured activity for 24 hours. After the pre-starvation period, each fly was tipped to
a second monitor tube containing starvation media (agar with preservative prepared as described
above) and activity was monitored every day until each fly died.

To assess the effect of the different environmental conditions on starvation resistance,
mean resistance per RIL from the DAM starvation assay was compared to mean resistance per
RIL from the large-scale screen using general linear regression. To determine differences in
activity patterns due to starvation resistance rank (high versus low) during the pre-starvation
period, we used a full three-way ANOVA model with interactions, and treated starvation rank,
sex, and the light status of the incubator (light versus dark) as fixed effects. The effect size of the
main effects and interactions were calculated using Cohen's F, which determines the effect size
as a ratio of the between-group and between-replicate standard deviations (R package: sjstats)
(Cohen 1988; Quinn and Keough 2002; Lüdecke 2018).

Triglyceride level assay

We duplicated 311 pA1/pA2 and 628 pB1/pB2 DSPR RILs from stocks to two replicate
vials, allowing parental flies to lay eggs for 1-2 days. Eleven days following the start of egg laying
we collected two sets of 10-12 females from each parental vial, resulting in four collection vials
from each RIL. Flies were aged for three additional days before measuring triglyceride level.

Experimental females from each collection vial were anesthetized using CO_2, and groups
of 5 were arrayed into deep well plates (Axygen, P-DW-11-C) over ice, with each well pre-loaded
with a single glass bead. This resulted in 8 replicate samples of 5 females per RIL. Immediately
after finishing a plate, we added 400µl of cold homogenization buffer (10mM potassium
phosphate monobasic, 1mM EDTA, 0.05% Tween-20) to each well, homogenized for 45sec in a
Mini-BeadBeater-96 (Bio Spec Products, Inc.), and centrifuged for 4min at 2,500g. We then
moved 50µl of the supernatant to a standard PCR plate, incubated the plate in a thermocycler at
70°C for 5min, and then placed the plate on ice for 5min.

During the incubation steps, we added 30µl of homogenization buffer to 92 of the 96
wells of a flat-bottom, polystyrene assay plate (Greiner, 655101), and subsequently added 20µl
of the heat-deactivated fly homogenate to each. The four remaining wells of every assay plate

were dedicated to controls; one blank well contained 50µl of homogenization buffer only, and

three wells contained 5µl of Glycerol Standard Solution (SigmaAldrich, G7793, 2.5mg/ml) along

with 45µl of homogenization buffer.

The assay plate was then inserted into a BioTek Powerwave XS2 instrument pre-heated

to 37°C and read at 540nm (baseline absorbance scan). After the scan, and within 10min, we

added 100µl of Free Glycerol Reagent (SigmaAldrich, F6428) to each well. The plate was then re-

inserted into the instrument, incubated at 37°C for 5min, and read again at 540nm (free glycerol

absorbance scan). After this second scan, and again within 10min, we added 25µl of Triglyceride

Reagent (SigmaAldrich, T2449) to each well. The plate was again incubated at 37°C for 5min in

the machine and read for a third time at 540nm (triglyceride, or final absorbance scan).

For each sample, we obtained the final absorbance for each sample (FA_{sample}) and

calculated the initial absorbance (IA_{sample}) as the free glycerol measurement minus the baseline

measurement. We also generated the average final absorbance for the three standard wells

(FA_{std}) and the initial absorbance for the one blank well (IA_{blank}). We then estimated the true

serum triglyceride level as

$$\left[FA_{sample} - \left(IA_{sample} \times F\right)\right]/\left[FA_{std} - \left(IA_{blank} \times F\right)\right]$$

where F = 0.8. We then multiplied this value by the concentration of the glycerol standard

solution (2.5mg/ml) and used the average value across all 8 replicate samples as the RIL mean

triglyceride level for mapping and analysis. For precise details of the enzyme assay and

triglyceride calculation, see the SigmaAldrich Serum Triglyceride Determination kit (TR0100).

Differences in triglyceride level due to DSPR subpopulation were investigated with a one-way

ANOVA followed by post hoc comparisons using Tukey's HSD (experiment-wide α = 0.05).

Correlations among traits

The relationships between starvation resistance measured in the large-scale starvation

assay and DAM starvation resistance, desiccation resistance, and triglyceride level were assessed

with general linear models using the mean trait value per RIL. Subpopulation (pA1, pA2, pB1,

pB2) was included as a factor in the analysis of starvation resistance and triglyceride content.

Correlations among the three DGRP starvation datasets - that from Mackay et al. (2012),

Everman and Morgan (2018), and the new screen we report here - were examined in a pairwise

manner using line means, accounting for multiple comparisons with a Bonferroni-adjusted alpha

level (three comparisons: α = 0.0167). Any difference in overall mean starvation resistance

measured over experiments was analyzed with a one-way ANOVA, treating study as a fixed

factor.

Heritability

The genetic and phenotypic variances of starvation resistance and triglyceride content for

the pA and pB DSPR panels, and of starvation resistance for the DGRP panel, were estimated with

a linear mixed model using the lme and varcomp functions in R (R package: APE, Paradis *et al.*

2004; R package: nlme, Pinheiro *et al.* 2017). We calculated broad sense heritability as the

proportion of the total variance of the strain-specific response explained by the estimated

genetic variance component (Lynch and Walsh 1998).

QTL mapping in the DSPR

QTL mapping analysis and power using the DSPR panel are discussed in detail in King *et*

al. (2012a,b). Briefly, QTL mapping and peak analysis was performed on starvation resistance and

triglyceride data using the DSPRqtl R package (github.com/egking/DSPRqtl; FlyRILs.org) by

regressing the mean trait response for each RIL on the additive probabilities that each of the

panel founders contributed to the haplotype of the RIL at each mapped position. Significance

thresholds were assigned following 1000 permutations of the data, and positions of putative

causative loci were estimated with 2-LOD support intervals, which approximate 95% confidence

intervals for QTL position in the DSPR (King *et al.* 2012a). Mean starvation resistance varied

between sexes in both the pA and pB panel ($F_{3,3440}$ = 18.317; p < 0.0001; Fig S1; Table S1), and

subpopulation influenced female starvation resistance in the pA panel (Tukey's HSD p < 0.0001;

Fig S1). Therefore, QTL mapping was performed for males and females of each panel separately,

and subpopulation was included as a covariate in the analysis of pA females. Mean female

triglyceride level was similar between the pA1 and pA2 subpopulations (Tukey's HSD p = 0.75; Fig

S2; Table S2) but varied between the pB1 and pB2 subpopulations (Tukey's HSD p < 0.0001; Fig

S2; Table S2), so subpopulation was included as a covariate in QTL analysis of the pB triglyceride

data.

Genomewide association mapping in the DGRP

Variants associated with male and female starvation resistance in the DGRP were

identified using the DGRP2 web-based GWA mapping tool (http://dgrp2.gnets.ncsu.edu), which

takes into account variable *Wolbachia* infection status and large inversions that segregate among

the lines (Mackay *et al.* 2012; Huang *et al.* 2014). We performed GWA on data collected in this

study and reanalyzed data from Mackay *et al.* (2012) and for starvation resistance data obtained

from young flies (5 – 7 days old) from Everman and Morgan (2018). We additionally assigned the

150 DGRP lines that are shared between the three datasets an across-study mean and performed

GWA on this summary measure of starvation resistance.

SNPs associated with starvation resistance were identified within each of the four

datasets following correction for multiple comparisons using FDR (Benjamini and Hochberg 1995)

in R (p.adjust; R Core Team 2017)*.* Since we found no significantly associated SNPs with an FDR

adjusted p-value < 0.05 for any starvation resistance dataset in either sex, we relaxed the

significance threshold to an FDR adjusted p-value < 0.2. As a significance threshold of $P < 10^{-5}$ is

commonly used in the DGRP (e.g. Mackay *et al.* 2012; Morozova *et al.* 2014; Huang *et al.* 2014;

Everman and Morgan 2018), we also present SNPs associated with starvation resistance in each

of the four datasets using this threshold.

There was minimal overlap in the identity of the above-threshold, starvation-associated

variants in each study. Thus, we sought to examine whether the sign of the additive effects of

these sets of variants was preserved across studies. Additive effects were calculated as one-half

the difference in starvation resistance between lines homozygous for the major allele and lines

homozygous for the minor allele (major allele frequency > 0.5), after accounting for *Wolbachia*

infection and TE insertions (Falconer and Mackay 1996; Huang *et al.* 2014). To determine the

proportion of SNPs that are expected by chance to have additive effects of the same sign across

studies, we obtained random samples of 50 SNPs from all of the DGRP SNP calls (> 2 million SNPs)

and calculated the additive effects of the random SNPs across pairs of datasets for each sex. To

account for the possibility that the frequency spectrum of above-threshold (P < 10^{-5}), associated

SNPs is not represented by a set of randomly-selected variants, we stratified the random subsets

of 50 SNPs according to the distribution of allele frequencies of the top 50 SNPs associated with

starvation resistance for each sex in each study. Allele frequency bins used in this stratification

were 0.05 – 0.1, > 0.1 – 0.2, > 0.2 – 0.3, > 0.3 – 0.4, and > 0.4 – 0. 5. The exact stratification for

each sex and dataset is provided in Table S3. This process was repeated 1000 times for each

paired comparison of datasets (6 comparisons total) using an ordinary nonparametric

bootstrapping procedure with the R package boot (Davison and Hinkley 1997; Canty and Ripley

2017). For each iteration, we used a custom R function (see File S2) to calculate the proportion

of the 50 random stratified SNPs that had positive additive effects in both of the datasets being

compared.

Data availability

369   DGRP   data   from   Mackay   *et   al.*   (2012)   are   available   online   from

370   http://dgrp2.gnets.ncsu.edu, and DGRP data from Everman and Morgan (2018) are available

371   from Dryad (DOI: https://doi.org/10.5061/dryad.vq087). All phenotypes collected in this study

372   are available in File S1. R code for the bootstrapping analysis and calculation of additive effects

373   for DGRP starvation resistance is available in File S2.

Results and Discussion

Phenotypic variation in starvation resistance

Starvation resistance in both the DSPR and DGRP was highly variable among strains (Fig

1, Fig 2). In the DSPR, pB RILs tended to have higher starvation resistance than pA RILs across

sexes (Fig S1), and a narrower range of phenotypic variation in mean starvation resistance (Fig

1). These differences likely reflect the different sets of founder lines used to establish the two

sets of RILs. Males and females in both the DSPR and DGRP differed in mean starvation resistance,

with males being less starvation resistant overall (Fig S1, Fig S3, Table S1, Table S4). Nonetheless,

male and female starvation resistance was significantly correlated in the DSPR (pA: R^2 = 53.0%;

pB: R^2 = 57.0%; Fig S4) and DGRP (R^2 = 68.0%; Fig S5). Any sex-specific differences in starvation

resistance are likely influenced by a combination of higher glycogen and triglyceride levels and

larger body size, which are often observed for females relative to males (Chippindale *et al.* 1996;

Toda and Kimura 1997; Schwasinger-Schmidt *et al.* 2012; Goenaga *et al.* 2013). Finally, the broad

sense heritability for starvation resistance measured for both sexes in the DSPR and DGRP was

routinely high (Table 1).

389
390
391
392   ***Table 1.*** *Broad sense heritability by trait, sex, and mapping panel.*

Panel	Trait	Sex	Heritability
DSPR pA	SR	Female	80.0%
DSPR pA	SR	Male	73.1%
DSPR pB	SR	Female	74.4%
DSPR pB	SR	Male	71.5%
DGRP	SR	Female	87.1%
DGRP	SR	Male	87.0%
DSPR pA	TL	Female	77.5%
DSPR pB	TL	Female	82.3%

SR = Starvation Resistance; TL = Triglyceride Level

Figure 1. Variation in mean (±SD) starvation resistance for each sex. A-D shows data for DSPR RIL panels
*pA and pB. E and F show data for the founder lines. In E and F, names of the founder lines are shown on
the x-axis; the founder line AB8 is the founder shared by the two mapping panels.*

Figure 2. *Variation in mean DGRP ($\pm$ SD) starvation resistance in females (orange) and males (red).*

400
401
402

We screened the DSPR and DGRP at 23°C and under constant light conditions, whereas

others have measured starvation resistance in *D. melanogaster* at a higher temperature and with

a light cycle (e.g. Mackay *et al.* (2012) and Everman and Morgan (2018)). Since starvation

resistance is sensitive to the thermal environment (van Herrewege and David 1997; Karan and

Parkash 1998; Karan *et al.* 1998; Hoffmann *et al.* 2005a; Bauerfeind *et al.* 2014) and may vary

under different photoperiods (Sheeba *et al.* 2000; Xu *et al.* 2008; Seay and Thummel 2011), we

sought to re-measure starvation resistance for a subset of DSPR RILs at 25°C and with a 12 hour

: 12 hour light/dark cycle. Overall, starvation resistance of the re-tested RILs was lower in both

sexes compared to that measured in the original large-scale starvation assay (effect of assay:

$F_{1,136} = 31.60$, $p < 0.0001$; Fig S6A). Despite this, starvation resistance in the subset of RILs was

significantly correlated between the two experiments (Females: $\beta = 0.43 \pm 0.04$, $t = 9.7$, $p <$

0.0001, $R^2 = 73.9\%$; Males: $\beta = 0.59 \pm 0.05$, $t = 10.9$, $p < 0.0001$, $R^2 = 78.3\%$; Fig S6B).

We similarly compared starvation resistance phenotypes for the DGRP measured in the

current study with those generated by Mackay *et al.* (2012) and Everman and Morgan (2018). On

average, starvation resistance in the DGRP measured in this study was much higher compared to

estimates from Mackay *et al.* (2012) and Everman and Morgan (2018) ($F_{2,532} = 1457.5$, $p < 0.0001$;

Fig S7). We can only speculate as to the cause of this major difference, but we did use a lower

rearing/testing temperature than these other studies (23°C versus 25°C), our flies were not

exposed to a light cycle, humidity varied across studies, and the fly culture media is likely

different. Notably, we included preservatives in the starvation media we used here, in contrast

to that used in Mackay *et al.* (2012) and Everman and Morgan (2018). The addition of

preservatives may have contributed to the increased starvation resistance we observed via

suppression of fungal and bacterial growth.

Even given the large across-study difference in mean starvation resistance in the DGRP,

we found moderately strong correlations in both sexes among each dataset, ranging from 50.8%

to 64.4% (Fig S8). The high correspondence among these three DGRP datasets, coupled with the

phenotypic correlation between the subset of DSPR strains assayed using two different

approaches (see above), suggests that fundamental aspects of the genetic control of starvation

resistance are generally consistent over experiments, even when environmental conditions such

as temperature are quite different. The differences we observed in starvation resistance between

studies may reflect ecologically-relevant phenotypic plasticity. The temporally variable thermal

environment is a particularly important source of selection for ectothermic organisms (Bell 2010;

Bergland *et al.* 2014). Plastic shifts in starvation resistance in response to temperature can have

important fitness benefits, including seasonal adaptation to fluctuating resource availability as

has been reported in the butterfly *Bicyclus anynana* (Pijpe *et al.* 2007) and following the induction

of diapause in *D. melanogaster* (Schmidt *et al.* 2005; Rion and Kawecki 2007). Collectively, these

previous studies and our data speak to the important influence of both phenotypic plasticity and

genotype on variation in starvation resistance in natural populations.

Desiccation resistance and activity levels in DSPR lines with extreme starvation resistance

Several artificial selection studies for starvation resistance have observed a correlated

change in desiccation resistance, suggesting these stress traits are related (Hoffmann and

Parsons 1989; Chippindale *et al.* 1996; Hoffmann and Harshman 1999; Harshman *et al.* 1999;

Hoffmann *et al.* 2001). We investigated this correlation in the DSPR by measuring female

desiccation resistance in RILs chosen from the two tails of the phenotypic distribution of female

starvation resistance (17 high starvation RILs and 16 low). We found that desiccation and

starvation resistance were significantly correlated (R^2 = 43.8, $F_{1,31}$ = 24.11, p < 0.0001; Fig 3). We

also found that mean desiccation resistance was considerably lower than starvation resistance in

the DSPR (Figures 1 versus 3), replicating previous studies (Da Lage *et al.* 1990; Hoffmann and

Parsons 1993; van Herrewege and David 1997; Karan and Parkash 1998; Karan *et al.* 1998;

Hoffmann *et al.* 2001). This pattern is most likely due to the much higher risk of death due to

water loss compared to risk of death due to lack of food (Fairbanks and Burch 1970; Da Lage *et*

al. 1990; Gibbs 2003). For instance, flies that die due to desiccation still have the majority of their

lipid stores (Da Lage *et al.* 1989), whereas, under sufficiently humid conditions, flies remain alive

until their fat stores are depleted (Da Lage *et al.* 1990; Karan *et al.* 1998).

Figure 3. *Mean starvation and desiccation resistance are correlated in the DSPR ($F_{1,31}$ = 21.11, p < 0.0001).*
Desiccation resistance is presented as RIL means (±SD). Open symbols indicate "low" starvation resistance
RILs; filled symbols indicate "high" starvation resistance RILs, and the dashed vertical line also separates
these RIL classes.

463
464
465

The relationship between starvation and desiccation resistance in this study provides

support for the genetic correlation that has been proposed to exist between these traits.

However, it does not rule out the potential that the covariation observed between starvation and

desiccation resistance may be influenced by genetic variation in one or more other traits that are

related to these resistance phenotypes, such as body size or overall lifespan (Rose *et al.* 1992;

Parsons 1995; Chippindale *et al.* 1996; Toda and Kimura 1997; Karan and Parkash 1998; Hoffmann

and Harshman 1999; Bubliy and Loeschcke 2005). Indeed, instances of positively and negatively

correlated levels of desiccation and starvation resistance in natural and selected populations

have been described, suggesting that these stress traits can vary independently across different

environments and selection regimes (Da Lage *et al.* 1990; van Herrewege and David 1997; Karan

and Parkash 1998; Karan *et al.* 1998; Kennington *et al.* 2001; Harbison *et al.* 2004). Physiological

independence of desiccation and starvation resistance likely contributes to the varying strength

of the correlation between these traits across multiple studies (Hoffmann and Harshman 1999).

A more intensive sampling of the DSPR would be necessary to thoroughly investigate the genetic

correlation between starvation and desiccation resistance.

One physiological mechanism that may increase tolerance to environmental stressors

(such as starvation or desiccation) is a reduction in metabolic rate (Lighton and Bartholomew

1988; Hoffmann and Parsons 1989, 1991; Chippindale *et al.* 1996; Djawdan *et al.* 1997; Marron

et al. 2003; Rion and Kawecki 2007; Schwasinger-Schmidt *et al.* 2012; Slocumb *et al.* 2015).

Indeed, selection for starvation resistance has been shown to lead to a correlated change in

activity level, an indirect proxy for metabolic rate (Hoffmann and Parsons 1993; Schwasinger-

Schmidt *et al.* 2012). Here, we assessed activity of a subset of high (N = 16) and low (N = 19)

starvation resistance RILs to understand how genetic variability in starvation resistance relates

to activity levels under non-stressful conditions. Prior to starvation, males and females differed

in activity level across the light and dark period ($F_{1,132}$ = 16.9, p < 0.0001; Fig 4B; Table S5), and

high starvation resistance RILs were less active than low starvation resistance RILs ($F_{1,132}$ = 12.5,

p < 0.001; Fig 4A; Table S5). The effects of starvation resistance rank (high vs low), sex, and the

light status (light versus dark) on activity were similar in magnitude (Cohen's F: 0.21-0.36; Table

S5), suggesting that these factors contribute similarly to variation in waking activity levels.

Figure 4. *Activity level on regular media for males (red) and females (black) from a subset of high and low female starvation resistance RILs. Panels in each figure indicate the light and dark periods of a 24-hour monitoring period. Activity while awake was influenced by both a sex-by-light interaction ($F_{1,132}$ = 16.9, P < 0.001) and by starvation resistance class (i.e. high or low; $F_{1,132}$ = 12.5, P < 0.001).*

The differences in activity between high and low starvation resistance lines on regular, nutritive media (Figure 4) were preserved under starvation conditions, with the high starvation resistant lines being less active than the low starvation resistant lines throughout the starvation process (Fig S9; Table S6). This pattern aligns with those from previous studies. For instance, Schwasinger-Schmidt *et al.* (2012) found that activity of flies with high starvation resistance was reduced following 15 generations of selection for starvation resistance in both males and females. Slocumb *et al.* (2015) also found that waking activity was reduced in lines selected to have high starvation resistance. Although previous associations between increased starvation tolerance and lower activity levels, metabolic rate, and changes in behavior have been observed (Murphey and Hall 1969; Hoffmann and Parsons 1989; Blows and Hoffman 1993; Djawdan *et al.* 1997; Karan *et al.* 1998; Schwasinger-Schmidt *et al.* 2012; Masek *et al.* 2014), our findings present a novel addition to our understanding of how increased starvation resistance may occur. Behavioral components of energy conservation are likely to play a role in how individuals

compensate for stressful conditions (van Dijk *et al.* 2002; McCue 2010; Masek *et al.* 2014) and

represent an additional facet of the complex nature of phenotypic variability in starvation

resistance.

Phenotypic variation in triglyceride level

Periods of starvation have been shown to significantly reduce triglyceride levels in both

males and females (Schwasinger-Schmidt *et al.* 2012), and others have suggested that fat stores

and starvation resistance may be genetically correlated (Service *et al.* 1985; Rose *et al.* 1992;

Chippindale *et al.* 1996; Harshman *et al.* 1999; Schwasinger-Schmidt *et al.* 2012; Slocumb *et al.*

2015). To investigate the relationship between these traits in the DSPR, we measured mean

female triglyceride level in a subset of the pA and pB DSPR RILs and found substantial phenotypic

variation among RILs (pA: N = 311, 1.95 – 5.32 mg/mL; pB: N = 628, 1.58 – 5.11 mg/mL; Fig 5).

Broad sense heritability estimates for triglyceride level were high for each panel (Table 1). Mean

starvation resistance and triglyceride level were positively correlated in both DSPR panels,

although the correlation in the pA and pB panels was significantly different (Fig 6). Overall,

variation in mean starvation resistance explained 23.7% of the variation observed in mean

triglyceride level among the DSPR RILs across the two mapping panels (R^2 = 23.7%, $F_{3,929}$ = 95.96,

p < 0.0001; Fig 6), suggesting that a proportion of variation in the starvation resistance response

can be explained by variation in triglyceride level in the DSPR.

Figure 5. *Variation in mean DSPR triglyceride level (±SD) for females in the pA panel (A) and pB panel (B).*

Figure 6. *Mean female starvation resistance and female triglyceride level are positively correlated ($F_{3,929}$ =*
95.96, p < 0.0001). The strength of the correlation varied between the two mapping panels (interaction:
$F_{1,929}$ = 9.32, p < 0. 01). Points are colored to indicate subpopulation for each mapping panel, although
subpopulation was not included in the regression analysis.

542
543
544

The correlation between triglyceride level (or total lipid level, depending on the study)

and starvation resistance has been measured in numerous natural and artificially selected

Drosophila populations, and a positive relationship is often described (Chippindale *et al.* 1996;

Djawdan *et al.* 1997; van Herrewege and David 1997; Harshman *et al.* 1999; Schwasinger-Schmidt

et al. 2012; Goenaga *et al.* 2013; Slocumb *et al.* 2015; Hardy *et al.* 2018). For example, in

isofemale lines derived from populations distributed across approximately 14.4 degrees of

latitude, Goenaga *et al.* (2015) found that 12% of the variation in female starvation resistance

was accounted for by lipid content. Similarly, Chippindale *et al.* (1996) found a very strong

positive relationship between total lipids and starvation resistance following extended selection

for increased starvation resistance and suggested that lipid levels may directly determine

starvation resistance. Hoffmann and Harshman (1999) also inferred that lipid content is the

primary influencer of starvation resistance. However, a strong correspondence between lipid

content and starvation resistance is not always observed in strains derived from natural

populations (Da Lage *et al.* 1990; Robinson *et al.* 2000; Schmidt *et al.* 2005). Hoffman *et al.* (2001)

suggested that variation in the strength of the correlation between triglycerides and starvation

resistance may be due to the evolutionary history of the study population. Evolutionary tradeoffs

between increased lipid storage and other aspects of fitness may also influence the correlation

between starvation resistance and lipid levels (Hardy *et al.* 2015, 2018). Furthermore, artificial

selection may increase starvation resistance via mechanisms that preferentially modify lipid

accumulation or metabolism, rather than by impacting energy level (Hoffmann and Parsons 1989;

Blows and Hoffman 1993; Hoffmann *et al.* 2001; Marron *et al.* 2003; Rion and Kawecki 2007). The

relationship observed between triglyceride levels and starvation resistance in our study supports

the hypothesis that triglyceride levels and starvation resistance are physiologically and

potentially genetically related. Equally, it is evident from our DSPR data that triglyceride level

likely influences starvation resistance to a lesser degree than proposed by Chippindale *et al.*

(1996) and Hoffmann and Harshman (1999).

QTL mapping of starvation resistance and triglyceride level in the DSPR

Following characterization of the extensive phenotypic variation for starvation resistance

and triglyceride level in the DSPR, we used QTL mapping to investigate and compare the genetic

architectures of these traits in this multiparental panel that offers high power to map QTL to

relatively modest genetic intervals (see King *et al.* (2012a)).

We mapped 8 QTL for starvation resistance in the pA panel and 7 QTL in the pB panel,

several of which overlapped between sexes (Fig S10). Both sets of QTL explained a substantial

amount of variation in starvation resistance, with individual peaks accounting for 3.7-13.2% of

the variation (Table 2). The total variance explained by QTL in the pA (pB) population was 26.1%

(32.8%) in females and 17.5% (37.9%) in males, assuming QTL are independent and additive

(Table 2). None of the QTL identified in the pA and pB mapping panels overlapped, and since

power to detect 5% QTL is expected to be high in our study (King *et al.* 2012a) and all DSPR

phenotyping was completed within seven months using the same design and environmental

conditions, this likely reflects genetic variation among the different sets of founders used to

establish the two sets of lines.

Table 2. Summary of QTL identified for starvation resistance and triglyceride level in the DSPR, divided by
mapping panel (pA, pB). No QTL were identified for triglyceride level in the pA panel. Genes and functions
associated with starvation resistance are listed in Table S7. In the last column, % genes associated with
starvation resistance was determined by cross-referencing genes within each QTL interval with previously
reported candidates and biological functions reported by FlyBase (Lin et al. *1998; Clancy* et al. *2001;*
FlyBase Curators et al. *2004; Harbison* et al. *2004; Sørensen* et al. *2007; Nuzhdin* et al. *2007).*

Starvation Resistance QTL: pA DSPR panel

QTL	Sex	Peak LOD	Chr	Physical Interval (Mb)	Genetic Interval (cM)	Variance Explained	No. Genes	No. Genes assoc. with SR
SA1	M	7.41	2L	0.34-0.64	0.36-1.02	3.90	52	3
SA2	F	10.85	2R	4.90-5.53	59.39-60.64	6.01	122	10
SA2	M	10.59	2R	4.87-5.58	59.32-60.74	5.52	126	10
SA3	F	8.05	2R	12.85-13.43	80.75-82.81	4.23	110	9
SA4	F	7.03	2R	14.18-14.72	84.14-86.68	3.70	105	7
SA5	F	10.13	3L	3.40-3.92	6.89-8.88	5.29	52	4
SA6	F	13.21	3R	0.79-2.37	47.10-47.52	6.85	244	11
SA7	M	7.66	3R	15.55-15.86	66.52-67.36	4.03	52	4
SA8	M	7.73	3R	21.07-21.48	86.77-88.20	4.06	80	3

Starvation Resistance QTL: pB DSPR panel

QTL	Sex	Peak LOD	Chr	Physical Interval (Mb)	Genetic Interval (cM)	Variance Explained	No. Genes	No. Genes assoc. with SR
SB1	F	7.68	2L	7.02-7.30	23.69-25.20	4.05	33	1
SB2	M	8.35	2L	10.10-10.50	39.24-40.93	4.39	110	13
SB3	F	7.13	2L	12.14-12.65	45.94-46.96	3.77	44	4
SB3	M	7.05	2L	12.13-12.63	45.92-46.93	3.72	45	4
SB4	F	12.28	2R	1.10-2.70	54.93-55.73	6.41	148	10

QTL	Sex	Peak LOD	Chr	Physical Interval (Mb)	Genetic Interval (cM)	Variance Explained	No. Genes	No. Genes assoc. with SR
SB4	M	19.40	2R	2.01-2.69	55.21-55.72	9.91	81	8
SB5	F	18.98	3L	18.94-19.42	45.48-45.75	9.73	54	5
SB5	M	26.40	3L	19.00-19.43	45.51-45.76	13.24	46	3
SB6	F	9.07	3R	8.86-9.20	52.50-53.29	4.77	48	7
SB6	M	12.77	3R	8.57-9.21	52.33-53.30	6.64	90	9
SB7	F	7.72	3R	25.65-26.15	99.12-100.11	4.08	111	8

Triglyceride Level QTL: pB DSPR panel

QTL	Sex	Peak LOD	Chr	Physical Interval (Mb)	Genetic Interval (cM)	Variance Explained	No. Genes	No. Genes assoc. with SR
TB1	F	8.73	3L	3.45-4.09	7.08-9.52	6.22	84	7
TB2	F	7.90	3R	7.18-8.01	50.50-51.54	5.64	118	5
TB3	F	8.45	3R	8.31-8.71	51.96-52.54	6.03	54	8
TB4	F	7.71	X	17.34-17.43	57.17-57.44	5.51	4	0

SR = Starvation Resistance

595
596
597

We mapped four distinct QTL for triglyceride level in the pB population, each of which

accounted for 5.5-6.2% of the variation in this trait (Table 2; Fig S11), in total explaining 23.4% of

the variation in pB. No QTL for triglyceride level were detected in the pA panel, likely due to the

reduced number of pA RILs assessed (pA N = 311; pB N = 628). However, even with this reduced

power, the QTL map for pA suggests that the genetic architecture for triglyceride level is different

between the two mapping panels, as there is no evidence of near-significant peaks in pA within

QTL intervals statistically identified in the pB panel (Fig S11). As has been seen in a number of

studies using the DSPR and other multiparental populations (King *et al.* 2012b; Giraud *et al.* 2014;

Najarro *et al.* 2015), the estimated phenotypic effects of each founder haplotype suggest that

multiple alleles may be present at our starvation and triglyceride QTL, since the effects do not

fall into two clear "high" and "low" classes (Figs S10 and S11).

Given the phenotypic correlation between triglyceride level and starvation resistance in

the DSPR (Fig 6) and similar correlations previously reported in other studies (Chippindale *et al.*

1998; Hoffmann and Harshman 1999), one might predict overlap of QTL associated with these

traits. Indeed, the intervals of two QTL identified for triglyceride level, TB1 and TB3, do overlap

with the female SA5 and male SB6 QTL intervals, respectively, identified for starvation resistance

(Table 2). However, given the complete lack of evidence for QTL for the same trait co-localizing

in both the pA and pB DSPR mapping panels, it is unlikely the variant(s) underlying TB1 (mapped

in the pB population) and SA5 (mapped in the pA population) are the same. The limited overlap

in the QTL intervals associated with starvation and triglycleride level suggests that the genetic

bases of this pair of traits are largely independent.

To investigate the relationship between the two QTL that do overlap within the same

panel (SB6 and TB3), we assessed the influence of haplotype structure at the overlapping QTL on

the positive phenotypic correlation between triglyceride level and starvation resistance (Fig 6).

In this analysis, we first identified the founder haplotype for each RIL at the positions of the

overlapping QTL peaks, then calculated the mean trait response of these haplotype groups (i.e.

the average phenotypes of RILs carrying each of the founder haplotypes). We then assessed the

correlation between haplotype-specific mean triglyceride level and starvation resistance with a

general linear model. After accounting for the haplotype structure at the overlapping peaks, we

found that mean starvation resistance and triglyceride level were significantly correlated ($F_{1,7}$ =

7.72, $p < 0.05$, R^2 = 52.4%; Fig S12), suggesting some pleiotropic variants may be responsible for

this pair of overlapping starvation resistance and triglyceride level QTL.

Across all QTL identified for starvation resistance and triglyceride level in this study,

several genes within the QTL intervals have functions related to these and other correlated traits

(Table 2, Table S7). Of particular interest are the 30 genes that fall within our QTL intervals that

were identified in previous starvation resistance studies (Clancy *et al.* 2001; Harbison *et al.* 2005;

Sørensen *et al.* 2007) (Table S7). Gene ontology analyses performed for each trait and panel also

revealed enrichment of genes within pA QTL related to glutathione metabolic process (6.91-fold

enrichment, FDR corrected P = .000146; Table S8), as well as several categories that were

enriched for pB triglyceride level (Table S8). However, it should be noted that the sets of genes

implicated by QTL mapping in the DSPR (3-244 genes per interval in this study) are extremely

unlikely to all contribute to trait variation, and their presence within QTL intervals cannot alone

be taken as evidence for causality, particularly if the number of genes that actually contribute to

trait variation is high.

Upon examination of the biological processes assigned to genes within the overlapping

starvation resistance and triglyceride intervals (TB3 and SB6), we found several genes that have

either been predicted or experimentally demonstrated to be associated with traits related to

starvation resistance and triglycerides or metabolism (Table S7 and references therein). Genes

that fall within the intervals of the overlapping peaks include those that influence adult lifespan

(e.g. *ry*, *Men*, *Gnmt* (Simonsen *et al.* 2006; Paik *et al.* 2012; Obata and Miura 2015)), lipid

metabolic processes (including *Lip3*, *CG11598*, *CG11608*, *CG18530* (FlyBase Curators *et al.* 2004)),

insulin signaling (e.g. *poly* (Bolukbasi *et al.* 2012)), response to starvation (e.g. *mthl12*, *Gnmt*

(UniProt Curators 2002; Obata *et al.* 2014)), larval feeding behavior (e.g. *Hug* (Melcher and

Pankratz 2005)), circadian rhythm and sleep (e.g. *timeout*, *Men* (Harbison *et al.* 2004; Benna *et*

al. 2010)), and triglyceride homeostasis (*Gnmt* (Obata *et al.* 2014)) (Table S7). These genes are

promising candidates for future studies seeking to examine the functional genetic relationship

between these two traits.

Predicting starvation resistance in the DSPR founders with mapped QTL haplotype effects

We can estimate the effect associated with each founder haplotype at each mapped QTL

in the DSPR. It follows that a combination of the estimates across all QTL can be used to predict

the actual phenotypes of the original founder strains. We measured starvation resistance in the

15 founder lines in addition to the DSPR RILs (Fig 1E, F) to test this prediction. It is likely that the

strength of the correlation between the estimated and actual trait response is influenced by the

number and effect size of each QTL mapped for the trait. To account for differences in the degree

to which starvation resistance is influenced by each QTL, we calculated the predicted mean trait

response for each founder line weighted by the variance explained by each QTL. By weighting the

estimates by the variance, QTL that have smaller effects will also have less influence on the

overall mean predicted trait response for each founder line, while QTL with large effects will

influence the predicted trait response for each founder line to a proportionately larger degree.

As predicted, using a general linear model the weighted mean predicted starvation

resistance of the founders based on QTL effects was significantly correlated with the sex-

averaged mean starvation resistance measured for the founder lines (R^2 = 44.5%, $F_{1,13}$ = 10.43, p

< 0.01; Fig 7). The slope of this relationship is relatively small (β = 0.16 +/- 0.05), suggesting that

while a large component of variation in starvation resistance is clearly genetic (as supported by

heritability estimates for each panel), substantial variation in the phenotype response is

unaccounted for by additive genetic effects at mapped QTL. This unaccounted-for genetic

variation in starvation resistance is likely due to many QTL with very small effects beyond our

power to detect them (King *et al.* 2012a). However, the strength of the correlation between

predicted and actual responses does suggest that QTL identified from the DSPR mapping panels

identify causative loci that influence the level of starvation resistance among the progenitors of

the RILs.

Figure 7. *Estimated starvation resistance weighted by the variance explained by each QTL and actual*
starvation resistance measured for the 15 founder lines of the DSPR were significantly correlated (β = 0.16
$\pm$ 0.05; $F_{1,13}$ = 10.43, p < 0.01). AB8 identifies the founder line shared between the pA and pB mapping
panels. Grey shading indicates the 95% CI of the regression.

685
686
687

Genomewide association mapping in the DGRP

Dissection of a quantitative genetic trait in multiple mapping panels can allow greater

resolution of the genetic architecture, and provide insight into how alleles unique to each panel

contribute to phenotypic variation. To gain this additional understanding we measured starvation

resistance in the DGRP and assessed the genetic architecture of this trait using GWA mapping in

four starvation resistance datasets: data collected in this study, data from Mackay *et al.* (2012),

data from Everman and Morgan (2018), and a consensus, across-study starvation resistance

measure calculated as the mean response across the three starvation datasets (N shared lines =

150). Using an FDR threshold of 20%, between 0 and 12 SNPs were associated with starvation

resistance for each dataset and sex (Table 3; Table S9). Aside from 3 SNPs that overlap between

the across-study mean dataset and the Mackay *et al.* (2012) dataset, none of these above-

threshold SNPs were the same (Table 3; Table S9). Using the more lenient significance threshold

of $P < 10^{-5}$, between 17 and 48 SNPs were associated with starvation resistance for each dataset

and sex (Table 3; Table S9); however, overlap in associated SNPs among datasets was still minimal

(Fig S13). With this significance threshold, our sex-specific GWA results for the Mackay *et al.*

(2012) and Everman and Morgan (2018) datasets matched those reported in the original studies

(Table 3; Table S9). The SNPs identified using the more lenient significance threshold include all

those identified at the FDR threshold of 20%, so all subsequent analyses are performed on the

larger set of associated SNPs, and we acknowledge that these sets may include larger fractions

of false-positive associations. The equivalent FDR for the $P < 10^{-5}$ thresholds are provided in Table

3.

Table 3. *Summary of DGRP GWA results and lines measured in this study, Mackay* et al. *(2012) and*
Everman and Morgan (2018). Text in italics indicates results from the reanalysis of starvation data
previously presented by Mackay et al. *(2012) and Huang* et al. *(2014) and Everman and Morgan (2018).*
Numbers noted with an asterisk indicate results that match those reported in the original studies.

Study	Significance Threshold	Equivalent FDR Female	Male	No. Strains	No. SNPs: Female	Male
This Study	FDR 0.2	-	-	168	2	3
	$P < 10^{-5}$	0.78	0.79	168	23	19
Mackay et al. (2012)	*FDR 0.2*	-	-	*203*	*11*	*0*
	$P < 10^{-5}$	*0.49*	*0.82*	*203*	*39**	*17**
Everman and Morgan (2018)	*FDR 0.2*	-	-	*164*	*0*	*0*
	$P < 10^{-5}$	*0.67*	*0.65*	*164*	*25**	*22**
Across-Study Mean Response	FDR 0.2	-	-	150	12	0
	$P < 10^{-5}$	0.43	0.80	150	48	22

715
716
717

Compared to genes implicated by QTL identified in the DSPR, which include several that

have been previously associated with starvation or related phenotypes (e.g., lifespan or lipid

content), DGRP GWAS hits implicate far fewer *a priori* strong candidate genes. Additionally, we

did not observe any GO enrichment following analysis of SNPs associated with each dataset,

although we acknowledge that the limited number of implicated genes likely compromised the

power of such enrichment analyses. Of the total 127 unique genes associated with starvation

resistance in the DGRP across studies and sexes, only two have been previously identified as

influencing starvation resistance in other mapping populations (*CG30118*, *scaf6*; Table S8;

(Sørensen *et al.* 2007)). More generally, five genes identified through our GWA had biological

processes linked to the determination of adult lifespan (e.g. *cnc* and *Egfr*; Table S8; (Sykiotis and

Bohmann 2008; Kamakura 2011)), and 5 have been previously associated with lipid metabolism

or metabolic processes (e.g. *GlcAT-P* and *Ugt86Dj*; Table S8; (FlyBase Curators *et al.* 2004; Gaudet

et al. 2010)). Given the relative lack of power in the DGRP (Long *et al.* 2014), and our use of a

permissive genomewide threshold, it could be that many of the GWAS associations are incorrect.

Equally, it could be the case that a series of novel pathways are involved in natural variation for

starvation resistance, and that traditional candidates - often identified via mutagenesis screens

rather than through examination of segregating allelic variation - typically do not harbor the

functional natural variants detectable in a GWAS (Mackay *et al.* 2009).

Repeatability in the SNPs associated with of starvation resistance across studies

The public availability of starvation resistance data for the DGRP from multiple studies

provides a novel opportunity to investigate the reliability and repeatability of associations

identified for a classic quantitative trait in the same mapping panel across independent

phenotypic screens (Lithgow *et al.* 2017). Despite the moderately high phenotypic correlation

between starvation resistance measured in the three studies (Fig S8), overlap in the variants

implicated in starvation resistance in the DGRP was very limited. Only one SNP (associated with

male starvation resistance) overlapped between starvation resistance measured in Mackay *et al.*

(2012) and Everman and Morgan (2018), and 2-4 overlapped between the across-study mean

response and data from this study, Mackay *et al.* (2012), and Everman and Morgan (2018) (Fig

S13).

The lack of overlap in SNPs associated with starvation resistance could be due to

differences in screen environments (discussed above), where genotype-by-environment effects

(often pervasive for complex traits (Gurganus *et al.* 1998)) could lead to different sites impacting

variation in different studies. However, it is potentially more likely that the sets of associated

SNPs have real, but extremely small effects on starvation resistance variation, and power in a

GWA panel of modest size is too low to consistently detect them (Boyle *et al.* 2017). If true, one

would predict that - in contrast to random SNPs - the "top SNPs" identified within each starvation

resistance dataset would have additive effects of the same sign across all studies. In essence,

significantly associated SNPs with positive effects on starvation resistance from data collected in

this study would be expected to have positive additive effects on starvation resistance measured

in Mackay *et al.* (2012) and Everman and Morgan (2018) more often than expected by random

chance.

To test this prediction, we first collected the sign of the additive effects of SNPs that

survived the significance threshold ($P < 10^{-5}$) associated with each measure of starvation

resistance in each dataset for both sexes (Table S9), and determined whether these top SNPs had

additive effects of the same sign in every other data set. We then established a null distribution

of SNP additive effect signs across pairs of datasets. This was accomplished by taking samples of

50 SNPs segregating in the DGRP and extracting the sign of the additive effect of each SNP in the

pair of datasets, regardless of the association statistic for that SNP. The proportion of the 50 SNPs

that had a positive additive effect on the trait was recorded for each of 1000 iterations and used

to build a distribution of additive effect shifts between the data collected in this study and that

presented by Mackay *et al.* (2012) and Everman and Morgan (2018) (see Methods). We note that

to compensate for any allele frequency bias in the variants that are actually most associated with

phenotype we ensured that the frequencies of the sampled SNPs were stratified to match the

top 50 SNPs associated with each sex and dataset (see Methods; Table S3).

Finally, we compared the proportion of top SNPs for which the sign stayed the same

across studies to our expected distribution for each pair of datasets (Fig 8). We found that for the

random samples of SNPs, the probability that the additive effects were positive in both datasets

compared was greater than 50% for most comparisons (distribution means ranging from 45.9-

72.1%; Fig 8). This implies that a random SNP is slightly more likely to have the same sign effect

across data sets, which may be explained in part by the phenotypic correlation between the

datasets (Fig S8). Even given this skew, far more top SNPs than expected by chance had additive

effects of the same sign in each of the starvation resistance studies (i.e. a SNP with a positive

effect on starvation resistance in this study had positive effects on starvation resistance obtained

by Mackay *et al.* (2012) and Everman and Morgan (2018); Fig 8; Table S8; Fig S14). This may

suggest that there is phenotypic signal even in SNPs that have very small effects, and that are not

clearly associated with starvation resistance in the GWAS (see Yang *et al.* 2010). Generally, the

consistency of the additive effects of SNPs associated with starvation resistance in the DGRP

calculated across data sets implies that starvation resistance is a highly polygenic trait, with a

large number of QTL with very small effects that influence variation in this trait (Mackay 2004;

Boyle *et al.* 2017).

Figure 8. *Distribution of the proportion of SNPs that are expected to have additive effects of the same sign in pairwise comparisons of data from this study, Mackay* et al. *(2012) and Everman and Morgan (2018). Data shown were obtained for random samples of 50 SNPs with 1000 iterations. In each plot, red indicates female data, blue indicates male data, and corresponding vertical lines and text annotation indicate the mean of the sex-specific samples. The shaded tails represent the upper and lower 95% confidence intervals of each distribution. The triangles in each plot represent the sex-specific observed proportion of top SNPs from each GWA analysis (P < 10^{-5}) that had additive effects that were the same sign across studies (Fig S14). For example, in A, 100% of the SNPs associated with female starvation resistance in this study had additive effects of the same sign when calculated for starvation resistance reported in Mackay* et al. *(2012). The dataset comparison is indicated above each plot.*

QTL associated with starvation resistance across mapping panels

Across all datasets reporting starvation resistance in the DSPR and DGRP, four SNPs

identified using the $P < 10^{-5}$ significance threshold fell within QTL intervals identified for

starvation resistance in the DSPR (Fig S15). In females, QTL SB4 and SA4 overlapped with two

SNPs associated with starvation resistance from Mackay *et al.* (2012), and one SNP associated

with starvation resistance in this study overlapped with QTL SA4. In males, one SNP associated

with starvation resistance from Mackay *et al.* (2012) overlapped with QTL SB3.

The general lack of correspondence among the loci associated with starvation resistance

in each mapping panel does not invalidate either approach as a strategy to uncover functional

variation. It is likely that many genes contribute to variation in this trait with effects that are

either fairly small or that vary in different genetic backgrounds, and we would not expect to

routinely identify such loci. In addition, comparison of the genetic architecture of quantitative

genetic traits across multiple panels is complicated by a number of additional factors. The genetic

structure of the mapping panel (e.g., whether it is a multiparental panel like the DSPR or a

population-based association study panel like the DGRP) influences the analytical strategy, and

the power and resolution of mapping (Long et al. 2014). The complement of alleles present in

the panel, and the frequency with which they segregate, will also affect the ability to identify the

same locus across mapping panels (e.g., King and Long 2017). Therefore, a lack of overlap in the

identified QTL for a complex, highly-polygenic trait between the DSPR and the DGRP is perhaps

not unexpected.

Conclusions

In this study, we described the complex quantitative genetics of starvation resistance in

two large *D. melanogaster* mapping panels that have been thoroughly genomically characterized.

The DSPR and DGRP panels have the advantage of increased, stable genetic diversity and provide

a unique comparison to many previous quantitative genetic studies of starvation resistance that

may be limited by genetic diversity or mapping power. Correlations between starvation

resistance and the additional traits described in this study offer insight into the genetic control

of related stress response traits and provide support for the hypothesis that the genetic

architecture of stress traits varies by population and is dependent upon sex, environment, and

the evolutionary history of the populations studied. The relationships between the traits

analyzed in this study also offer insight into the broader responses of organisms to starvation

stress, given the high conservation of mechanisms related to starvation resistance in diverse

species (Partridge *et al.* 2005; Rion and Kawecki 2007). Here, we have demonstrated that traits

related to survival under starvation conditions, energy storage, activity levels, and survival under

desiccation stress are phenotypically correlated in the DSPR, consistent with previous artificial

selection studies as well as some natural populations. However, we also clearly demonstrate that

starvation resistance and triglyceride level are largely genetically independent traits, indicating

that evolutionary constraint between these two traits is unlikely. We additionally describe the

highly polygenic nature of starvation resistance using the DGRP, leveraging previously published

phenotypes on the same lines to compare the genetic architecture of the trait across three

studies. Our work shows that despite a lack of overlap across studies in the identity of the variants

associated with phenotype at a nominal genomewide threshold, the sign of the additive effects

of such top SNPs are conserved across studies. In turn this suggests that these variants do

contribute to phenotype but have sufficiently small effects that they are not routinely captured

following a severe, genomewide correction for multiple tests. From this, we gain a much more

detailed understanding of the genetic control of starvation resistance in a genetically diverse

panel and provide insight into the utility of across-study and across-panel comparisons.

Acknowledgments

854   We thank Sophia Loschky and Brittny Smith for assistance with data collection, Tony Long and

855   Libby King for analytical assistance, and both John Kelly and Libby King for comments on a

856   previous version of this manuscript. The DGRP collection was purchased from the Bloomington

857   Drosophila Stock Center (NIH P40 OD018537). The project was supported by NIH grant R01

858   OD010974 to S.J.M. and Anthony D. Long.

References

Bauerfeind S. S., V. Kellermann, N. N. Moghadam, V. Loeschcke, and K. Fischer, 2014
Temperature and photoperiod affect stress resistance traits in *Drosophila melanogaster*.
Physiol. Entomol. 39: 237–246. https://doi.org/10.1111/phen.12068

Bell G., 2010 Fluctuating selection: the perpetual renewal of adaptation in variable
environments. Philos. Trans. R. Soc. B Biol. Sci. 365: 87–97.
https://doi.org/10.1098/rstb.2009.0150

Benjamini Y., and Y. Hochberg, 1995 Controlling the false discovery rate: A practical and
powerful approach to multiple testing. J. R. Stat. Soc. 57: 289–300.

Benna C., S. Bonaccorsi, C. Wülbeck, C. Helfrich-Förster, M. Gatti, *et al.*, 2010 *Drosophila
timeless2* is required for chromosome stability and circadian photoreception. Curr. Biol.
20: 346–352. https://doi.org/10.1016/j.cub.2009.12.048

Bergland A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, and D. A. Petrov, 2014 Genomic
evidence of rapid and stable adaptive oscillations over seasonal time scales in
Drosophila, (D. Bolnick, Ed.). PLoS Genet. 10: e1004775.
https://doi.org/10.1371/journal.pgen.1004775

Blows M. W., and A. A. Hoffman, 1993 The genetics of central and marginal populations of
Drosophila serrata. I. Genetic variation for stress resistance and species borders.
Evolution 47: 1255–1270. https://doi.org/10.2307/2409990

Bochdanovits Z., and G. de Jong, 2003 Experimental evolution in *Drosophila melanogaster*:

    interaction of temperature and food quality selection regimes. Evolution 57: 1829–1836.

Bolukbasi E., S. Vass, N. Cobbe, B. Nelson, V. Simossis, *et al.*, 2012 *Drosophila* poly suggests a

    novel role for the Elongator complex in insulin receptor-target of rapamycin signaling.

    Open Biol. 2: 110031–110031. https://doi.org/10.1098/rsob.110031

Boyle E. A., Y. I. Li, and J. K. Pritchard, 2017 An expanded view of complex traits: from

    polygenic to omnigenic. Cell 169: 1177–1186. https://doi.org/10.1016/j.cell.2017.05.038

Bubliy O. A., and V. Loeschcke, 2005 Correlated responses to selection for stress resistance and

    longevity in a laboratory population of *Drosophila melanogaster*. J. Evol. Biol. 18: 789–

    803. https://doi.org/10.1111/j.1420-9101.2005.00928.x

Canty A., and B. Ripley, 2017 boot: Bootstrap R (S-Plus) Functions. R package version 1.3-20

Chippindale A. K., A. M. Leroi, S. B. Kim, and M. R. Rose, 1993 Phenotypic plasticity and

    selection in *Drosophila* life-history evolution. I. Nutrition and the cost of reproduction. J.

    Evol. Biol. 6: 171–193. https://doi.org/10.1046/j.1420-9101.1993.6020171.x

Chippindale A. K., T. J. F. Chu, and M. R. Rose, 1996 Complex trade-offs and the evolution of

    starvation resistance in *Drosophila melanogaster*. Evolution 50: 753–766.

    https://doi.org/10.2307/2410848

Chippindale A. K., A. G. Gibbs, M. Sheik, K. J. Yee, M. Djawdan, *et al.*, 1998 Resource

    acquisition and the evolution of stress resistance in *Drosophila melanogaster*. Evolution

    52: 1342. https://doi.org/10.2307/2411304

Clancy D. J., D. Gems, L. G. Harshmann, S. Oldham, H. Stocker, *et al.*, 2001 Extension of life-span by loss of *CHICO* a *Drosophila* insulin receptor substrate protein. Science 292: 104–106.

Cohen J., 1988 *Statistical power for the behavioral sciences*. Lawrence Erlbaum, Hillsdale, NJ.

Colinet H., T. Chertemps, I. Boulogne, and D. Siaussat, 2015 Age-related decline of abiotic stress tolerance in young *Drosophila melanogaster* adults. Gerontol. Soc. Am. 00: 1–7.

Da Lage J. L., P. Capy, and J. R. David, 1989 Starvation and desiccation tolerance in *Drosophila melanogaster* adults: effects of environmental temperature. J. Insect Physiol. 35: 453–457.

Da Lage J., P. Capy, and J. David, 1990 Starvation and desiccation tolerance in *Drosophila melanogaster*: differences between European, North African and Afrotropical populations. Genet. Sel. Evol. 22: 381–391.

Davidson J. K., 1990 Non-parallel geographic patterns for tolerance to cold and desiccation in *Drosophila melanogaster* and *D. simulans*. Aust. J. Zool. 38: 155–161.

Davison A. C., and D. V. Hinkley, 1997 *Bootstrap methods and their applications*. Cambridge University Press, Cambridge.

Dijk P. van, G. Staaks, and I. Hardewig, 2002 The effect of fasting and refeeding on temperature preference, activity and growth of roach, *Rutilus rutilus*. Oecologia 130: 496–504. https://doi.org/10.1007/s00442-001-0830-3

Djawdan M., M. R. Rose, and T. J. Bradley, 1997 Does selection for stress resistance lower metabolic rate? Ecology 78: 828–837. https://doi.org/10.2307/2266062

Djawdan M., A. K. Chippindale, M. R. Rose, and T. J. Bradley, 1998 Metabolic reserves and evolved stress resistance in *Drosophila melanogaster*. Physiol. Zool. 71: 584–594.

Everman E. R., and T. J. Morgan, 2018 Antagonistic pleiotropy and mutation accumulation contribute to age-related decline in stress response. Evolution 72: 303–317. https://doi.org/10.1111/evo.13408

Fairbanks L. D., and G. E. Burch, 1970 Rate of water loss and water and fat content of adult *Drosophila melanogaster* of different ages. J. Insect Physiol. 16: 429–436.

Falconer D. S., and T. F. C. Mackay, 1996 *Quantitative genetics*. Longman Group, Ltd., Essex, England.

FlyBase Curators, Swiss-Prot Project Members, and InterPro Project Members, 2004 Gene Ontology in FlyBase through association of InterPro record with GO terms

Gaudet P., M. Livstone, and P. Thomas, 2010 Gene Ontology annotation inferences using phylogenetic trees. Go Ref. Genome Proj.

Gibbs A. G., 2003 Evolution of water conservation mechanisms in *Drosophila*. J. Exp. Biol. 206: 1183–1192. https://doi.org/10.1242/jeb.00233

Gilchrist G. W., L. M. Jeffers, B. West, D. G. Folk, J. Suess, *et al.*, 2008 Clinal patterns of desiccation and starvation resistance in ancestral and invading populations of *Drosophila subobscura*. Evol. Appl. 1: 513–523. https://doi.org/10.1111/j.1752-4571.2008.00040.x

Giraud H., C. Lehermeier, E. Bauer, M. Falque, V. Segura, *et al.*, 2014 Linkage disequilibrium with linkage analysis of multiline crosses reveals different multiallelic QTL for hybrid performance in the flint and dent heterotic groups of maize. Genetics 198: 1717–1734. https://doi.org/10.1534/genetics.114.169367

Goenaga J., J. J. Fanara, and E. Hasson, 2013 Latitudinal variation in starvation resistance is explained by lipid content in natural populations of *Drosophila melanogaster*. Evol. Biol. 40: 601–612. https://doi.org/10.1007/s11692-013-9235-6

Gurganus M. C., J. D. Fry, S. V. Nuzhdin, E. G. Pasyukova, R. F. Lyman, *et al.*, 1998 Genotype-environment interaction at quantitative trait loci affecting sensory bristle number in *Drosophila melanogaster*. Genetics 149: 1883–1898.

Harbison S. T., A. H. Yamamoto, J. J. Fanara, K. K. Norga, and T. F. Mackay, 2004 Quantitative trait loci affecting starvation resistance in *Drosophila melanogaster*. Genetics 166: 1807–1823.

Harbison S. T., S. Chang, K. P. Kamdar, and T. F. Mackay, 2005 Quantitative genomics of starvation stress resistance in *Drosophila*. Genome Biol. 6: R36. https://doi.org/10.1186/gb-2005-6-4-r36

Hardy C. M., R. T. Birse, M. J. Wolf, L. Yu, R. Bodmer, *et al.*, 2015 Obesity-associated cardiac dysfunction in starvation-selected *Drosophila melanogaster*. Am. J. Physiol. - Regul. Integr. Comp. Physiol. 309: R658–R667. https://doi.org/10.1152/ajpregu.00160.2015

Hardy C. M., M. K. Burke, L. J. Everett, M. V. Han, K. M. Lantz, *et al.*, 2018 Genome-wide
analysis of starvation-selected *Drosophila melanogaster*—a genetic model of obesity.
Mol. Biol. Evol. 35: 50–65. https://doi.org/10.1093/molbev/msx254

Harshman, Hoffmann, and Clark, 1999 Selection for starvation resistance in *Drosophila
melanogaster*: physiological correlates, enzyme activities and multiple stress responses.
J. Evol. Biol. 12: 370–379. https://doi.org/10.1046/j.1420-9101.1999.00024.x

Herrewege J. van, and J. R. David, 1997 Starvation and desiccation tolerances in *Drosophila*:
comparison of species from different climatic origins. Écoscience 4: 151–157.

Hoffmann A. A., and P. A. Parsons, 1989 Selection for increased desiccation resistance in
Drosophila melanogaster: additive genetic control and correlated responses for other
stresses. Genetics 122: 837–845.

Hoffmann A. A., and P. A. Parsons, 1991 *Evolutionary genetics and environmental stress*.
Oxford University Press, Oxford.

Hoffmann A. A., and P. A. Parsons, 1993 Direct and correlated responses to selection for
desiccation resistance: a comparison of *Drosophila melanogaster* and *D. simulans*. J.
Evol. Biol. 6: 643–657.

Hoffmann A. A., and L. G. Harshman, 1999 Desiccation and starvation resistance in *Drosophila*:
patterns of variation at the species, population and intrapopulation levels. Heredity 83:
637–643.

Hoffmann A. A., R. Hallas, C. Sinclair, and P. Mitrovski, 2001 Levels of variation in stress resistance in *Drosophila* among strains, local populations, and geographic regions: patterns for desiccation, starvation, cold resistance, and associated traits. Evolution 55: 1621–1630.

Hoffmann A. A., J. Shirriffs, and M. Scott, 2005a Relative importance of plastic vs genetic factors in adaptive differentiation: geographical variation for stress resistance in *Drosophila melanogaster* from eastern Australia. Funct. Ecol. 19: 222–227. https://doi.org/10.1111/j.1365-2435.2005.00959.x

Hoffmann A. A., R. Hallas, A. R. Anderson, and M. Telonis-Scott, 2005b Evidence for a robust sex-specific trade-off between cold resistance and starvation resistance in *Drosophila melanogaster*. J. Evol. Biol. 18: 804–810. https://doi.org/10.1111/j.1420-9101.2004.00871.x

Huang W., A. Massouras, Y. Inoue, J. Peiffer, M. Ràmia, *et al.*, 2014 Natural variation in genome architecture among 205 *Drosophila melanogaster* Genetic Reference Panel lines. Genome Res. 24: 1193–1208. https://doi.org/10.1101/gr.171546.113

Kamakura M., 2011 Royalactin induces queen differentiation in honeybees. Nature 473: 478–483. https://doi.org/10.1038/nature10093

Karan D., N. Dahiya, A. K. Munjal, P. Gibert, B. Moreteau, *et al.*, 1998 Desiccation and starvation tolerance of adult drosophila: opposite latitudinal clines in natural populations of three different species. Evolution 52: 825–831. https://doi.org/10.2307/2411276

Karan D., and R. Parkash, 1998 Desiccation tolerance and starvation resistance exhibit opposite latitudinal clines in Indian geographical populations of *Drosophila kikkawai*. Ecol. Entomol. 23: 391–396.

Kennington W. J., A. S. Gilchrist, D. B. Goldstein, and L. Partridge, 2001 The genetic basis of divergence in desiccation and starvation resistance among tropical and temperate populations of *Drosophila melanogaster*. Heredity 87: 363–372.

King E. G., S. J. Macdonald, and A. D. Long, 2012a Properties and power of the *Drosophila* Synthetic Population Resource for the routine dissection of complex traits. Genetics 191: 935–949. https://doi.org/10.1534/genetics.112.138537

King E. G., C. M. Merkes, C. L. McNeil, S. R. Hoofer, S. Sen, *et al.*, 2012b Genetic dissection of a model complex trait using the *Drosophila* Synthetic Population Resource. Genome Res. 22: 1558–1566. https://doi.org/10.1101/gr.134031.111

Lighton J. R. B., and G. A. Bartholomew, 1988 Standard energy metabolism of a desert harvester ant, *Pogonomyrmex rugosus*: Effects of temperature, body mass, group size, and humidity. Proc. Natl. Acad. Sci. U. S. A. 85: 4765–4769.

Lin Y. J., L. Seroude, and S. Benzer, 1998 Extended life-span and stress resistance in the *Drosophila* mutant *methuselah*. Science 282: 943–946.

Lithgow G. J., M. Driscoll, and P. Phillips, 2017 A long journey to reproducible results. Nat. News 548: 387.

Long A. D., S. J. Macdonald, and E. G. King, 2014 Dissecting complex traits using the *Drosophila* Synthetic Population Resource. Trends Genet. 30: 488–495. https://doi.org/10.1016/j.tig.2014.07.009

Lüdecke D., 2018 *sjstats: Statistical functions for regression models*. (Version 0.15.0.9000) https://doi.org/10.5281/zenodo.1284472

Lynch M., and B. Walsh, 1998 *Genetic and Analysis of Quantitative Traits*. Sinauer, Sunderland, MA.

Mackay T., 2004 The genetic architecture of quantitative traits: lessons from *Drosophila*. Curr. Opin. Genet. Dev. 14: 253–257. https://doi.org/10.1016/j.gde.2004.04.003

Mackay T. F. C., E. A. Stone, and J. F. Ayroles, 2009 The genetics of quantitative traits: challenges and prospects. Nat. Rev. Genet. 10: 565–577. https://doi.org/10.1038/nrg2612

Mackay T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, *et al.*, 2012 The *Drosophila melanogaster* Genetic Reference Panel. Nature 482: 173–178. https://doi.org/10.1038/nature10811

Marron M. T., T. A. Markow, K. J. Kain, and A. G. Gibbs, 2003 Effects of starvation and desiccation on energy metabolism in desert and mesic *Drosophila*. J. Insect Physiol. 49: 261–270.

Masek P., L. A. Reynolds, W. L. Bollinger, C. Moody, A. Mehta, *et al.*, 2014 Altered regulation of sleep and feeding contributes to starvation resistance in *Drosophila melanogaster*. J. Exp. Biol. 217: 3122–3132. https://doi.org/10.1242/jeb.103309

McCue M. D., 2010 Starvation physiology: Reviewing the different strategies animals use to survive a common challenge. Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 156: 1–18. https://doi.org/10.1016/j.cbpa.2010.01.002

Melcher C., and M. J. Pankratz, 2005 Candidate gustatory interneurons modulating feeding behavior in the *Drosophila* brain, (M. Bate, Ed.). PLoS Biol. 3: e305. https://doi.org/10.1371/journal.pbio.0030305

Morozova T. V., T. F. C. Mackay, and R. R. H. Anholt, 2014 Genetics and genomics of alcohol sensitivity. Mol. Genet. Genomics 289: 253–269. https://doi.org/10.1007/s00438-013-0808-y

Murphey R. M., and C. F. Hall, 1969 Some correlates of negative geotaxis in *Drosophila melanogaster*. Anim. Behav. 17: 181–185.

Najarro M. A., J. L. Hackett, B. R. Smith, C. A. Highfill, E. G. King, *et al.*, 2015 Identifying loci contributing to natural variation in xenobiotic resistance in *Drosophila*, (N. Singh, Ed.). PLOS Genet. 11: e1005663. https://doi.org/10.1371/journal.pgen.1005663

Nuzhdin S. V., A. A. Khazaeli, and J. W. Curtsinger, 2005 Survival analysis of life span quantitative trait loci in *Drosophila melanogaster*. Genetics 170: 719–731. https://doi.org/10.1534/genetics.104.038331

Nuzhdin S. V., L. G. Harshman, M. Zhou, and K. Harmon, 2007 Genome-enabled hitchhiking mapping identifies QTLs for stress resistance in natural *Drosophila*. Heredity 99: 313–321. https://doi.org/10.1038/sj.hdy.6801003

Obata F., E. Kuranaga, K. Tomioka, M. Ming, A. Takeishi, *et al.*, 2014 Necrosis-driven systemic immune response alters SAM metabolism through the FOXO-GNMT axis. Cell Rep. 7: 821–833. https://doi.org/10.1016/j.celrep.2014.03.046

Obata F., and M. Miura, 2015 Enhancing S-adenosyl-methionine catabolism extends *Drosophila* lifespan. Nat. Commun. 6. https://doi.org/10.1038/ncomms9332

Paik D., Y. G. Jang, Y. E. Lee, Y. N. Lee, R. Yamamoto, *et al.*, 2012 Misexpression screen delineates novel genes controlling *Drosophila* lifespan. Mech. Ageing Dev. 133: 234–245.

Paradis E., J. Claude, and K. Strimmer, 2004 APE: analysis of phylogenetics and evolution in R language. Bioinformatics 20: 289–290.

Parsons P. A., 1995 Inherited stress resistance and longevity: a stress theory of ageing. Heredity 75: 216–221.

Partridge L., M. D. W. Piper, and W. Mair, 2005 Dietary restriction in *Drosophila*. Mech. Ageing Dev. 126: 938–950. https://doi.org/10.1016/j.mad.2005.03.023

Pijpe J., P. M. Brakefield, and B. J. Zwaan, 2007 Phenotypic plasticity of starvation resistance in the butterfly *Bicyclus anynana*. Evol. Ecol. 21: 589–600. https://doi.org/10.1007/s10682-006-9137-5

Pinheiro J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team, 2017 nlme: linear and nonlinear mixed effects models. R Package Version 31-131.

Quinn G. P., and M. J. Keough, 2002 *Experimental design and data analysis for biologists*. Cambridge University Press, New York.

R Core Team, 2017 *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Rion S., and T. J. Kawecki, 2007 Evolutionary biology of starvation resistance: what we have learned from *Drosophila*. J. Evol. Biol. 20: 1655–1664. https://doi.org/10.1111/j.1420-9101.2007.01405.x

Robinson S. J., Zwaan, B J, and Partridge, L., 2000 Starvation resistance and adult body composition in a latitudinal cline of *Drosophila melanogaster*. Evolution 54: 1819–1824.

Rose M. R., L. N. Vu, S. U. Park, and J. L. Graves, Jr, 1992 Selection on stress resistance increases longevity in *Drosophila melanogaster*. Exp. Gerontol. 27: 241–250.

Schmidt P. S., L. Matzkin, M. Ippolito, and W. F. Eanes, 2005 Geographic variation in diapause incidence, life-history traits, and climatic adaptation in *Drosophila melanogaster*. Evolution 59: 1721–1732.

Schwasinger-Schmidt T. E., S. D. Kachman, and L. G. Harshman, 2012 Evolution of starvation resistance in *Drosophila melanogaster*: measurement of direct and correlated responses to artificial selection. J. Evol. Biol. 25: 378–387. https://doi.org/10.1111/j.1420-9101.2011.02428.x

Seay D. J., and C. S. Thummel, 2011 The circadian clock, light, and cryptochrome regulate

    feeding and metabolism in *Drosophila*. J. Biol. Rhythms 26: 497–506.

    https://doi.org/10.1177/0748730411420080

Service P. M., E. W. Hutchinson, M. D. MacKinley, and M. R. Rose, 1985 Resistance to

    environmental stress in *Drosophila melanogaster* selected for postponed senescence.

    Physiol. Zool. 58: 380–389.

Service P. M., and M. R. Rose, 1985 Genetic covariation among life-history components: the

    effect of novel environments. Evolution 39: 943. https://doi.org/10.2307/2408694

Sheeba V., V. K. Sharma, K. Shubha, M. K. Chandrashekaran, and A. Joshi, 2000 The effect of

    different light regimes on adult lifespan in *Drosophila melanogaster* is partly mediated

    through reproductive output. J. Biol. Rhythms 15: 380–392.

Simonsen A., R. C. Cumming, K. Lindmo, V. Galaviz, S. Cheng, *et al.*, 2006 Genetic modifiers

    of the *Drosophila Blue Cheese* gene link defects in lysosomal transport with decreased

    fife span and altered ubiquitinated-protein profiles. Genetics 176: 1283–1297.

    https://doi.org/10.1534/genetics.106.065011

Slocumb M. E., J. M. Regalado, M. Yoshizawa, G. G. Neely, P. Masek, *et al.*, 2015 Enhanced

    sleep is an evolutionarily adaptive response to starvation stress in *Drosophila*. PloS One

    10: e0131275.

Sørensen J. G., M. M. Nielsen, and V. Loeschcke, 2007 Gene expression profile analysis of

    Drosophila melanogaster selected for resistance to environmental stressors. J. Evol. Biol.

    20: 1624–1636. https://doi.org/10.1111/j.1420-9101.2007.01326.x

Sykiotis G. P., and D. Bohmann, 2008 Keap1/Nrf2 signalling regulates oxidative stress tolerance and lifespan *Drosophila*. Dev Cell 14: 76–85.

Telonis-Scott M., R. Hallas, S. W. McKechnie, C. W. Wee, and A. A. Hoffmann, 2009 Selection for cold resistance alters gene transcript levels in *Drosophila melanogaster*. J. Insect Physiol. 55: 549–555.

Toda M. J., and M. T. Kimura, 1997 Life-history traits related to host selection in mycophagous drosophilids. J. Anim. Ecol. 66: 154–166. https://doi.org/10.2307/6018

UniProt Curators, 2002 Manual transfer of experimentally-verified manual GO annotation data to orthologs by curator judgment of sequence similarity

Xu K., X. Zheng, and A. Sehgal, 2008 Regulation of feeding and metabolism by neuronal and peripheral clocks in *Drosophila*. Cell Metab. 8: 289–300. https://doi.org/10.1016/j.cmet.2008.09.006

Yang J., B. Benyamin, B. P. McEvoy, S. Gordon, A. K. Henders, *et al.*, 2010 Common SNPs explain a large proportion of heritability for human height. Nat. Genet. 42: 565–569. https://doi.org/10.1038/ng.608

Dissection of complex, fitness-related traits in multiple *Drosophila* mapping populations offers insight into the genetic control of stress resistance

Elizabeth R. Everman, Casey L. McNeil, Jennifer L. Hackett, Clint L. Bain, Stuart J. Macdonald

Department of Molecular Biosciences, University of Kansas, 1200 Sunnyside Avenue, Lawrence, Kansas 66045, USA.

Supplemental Figures, Tables, and Supplemental File descriptions

Supplemental Figures

Figure S1. *Mean (± 95% CI) starvation resistance per subpopulation and sex. Sex and subpopulation interacted to influence starvation resistance (F$_{3,3440}$ = 18.317; p < 0.0001), though only females from the pA1 and pA2 subpopulations were significantly different within a panel (Tukey's HSD adj. p < 0.0001).*

Figure S2. Mean female triglyceride level per subpopulation ($\pm$ 95% CI). Subpopulation influenced triglyceride level ($F_{3,935}$ = 37.099; $p < 0.0001$), though this was driven by differences between the pB subpopulations (Tukey's HSD adj. $p < 0.0001$)

Figure S3. *Mean starvation resistance ($\pm$ 95% CI) for males and females in the DGRP ($F_{1,334}$ = 118.21, p < 0.0001).*

Figure S4. *Correlation between sex-specific mean starvation resistance for the DSPR pA mapping panel (A) and pB mapping panel (B). Sex-specific responses were significantly correlated for both panels (pA: $F_{1,859}$ = 968.5, P < 0.0001; pB: $F_{1,860}$ = 1138, p < 0.0001). Grey shading around the regression line in both plots indicates the 95% confidence interval.*

Figure S5. *Male and female mean starvation resistance was significantly correlated in the DGRP ($F_{1,334}$ = 118.21, P < 0.0001). Grey shading around the regression line indicates the 95% confidence interval.*

Figure S6. *A. Starvation resistance was significantly higher in the large-scale starvation screen of all DSPR RILs compared to the DAM starvation assay for the selected subset of RILs (Assay: $F_{1,136}$ = 31.60, p < 0.0001). Mean starvation resistance across RIL means is presented ($\pm$ 95% CI). B. Mean starvation resistance measured in the large-scale starvation resistance screen (x-axis) was correlated with mean starvation resistance measured in the DAM assay (y-axis) in the DSPR (Females: β = 0.43 $\pm$ 0.04, t = 9.7, p < 0.0001, R^2 = 73.9%; Males: β = 0.59 $\pm$ 0.05, t = 10.9, p < 0.0001, R^2 = 78.3%). The multiple R^2 value in the plot includes the interaction between starvation resistance measured under different assay conditions with sex.*

Figure S7. *Mean starvation resistance (±95% CI) was significantly higher in this study compared to Mackay* et al. *(2012) and Everman and Morgan (2018) ($F_{2.532}$ = 1457.5, P < 0.0001).*

Figure S8. *Correlation between sex-specific mean starvation resistance in the DGRP panel for 150 lines that overlap between this study, Mackay* et al. *2012, and Everman and Morgan 2018. Red points indicate males and black points indicate females. All comparisons showed that the three independent measures of starvation resistance were significantly correlated (A: $F_{3,296}$ = 52.69, p < 0.0001; B: $F_{3,\,296}$ = 50.66, p < 0.0001; C: $F_{3.296}$ = 78.18, p < 0.0001).*

Figure S9. *Mean activity levels for females (top panel, black points) and males (bottom panel, red points) (± 95% CI) across the sampling periods during starvation in the DAM assay until death. Low starvation resistance RILs (solid line) tended to be more active during starvation compared to high starvation resistance RILs (dashed line) under both light and dark conditions. Light status (dark versus light) influenced the overall activity level of females but did not influence male activity. Data were analyzed with a repeated measures ANOVA; results are presented in Table S6. Similar to the pre-starvation period (Figure 4), waking activity levels of individuals during the DAM starvation experiment were primarily driven by starvation resistance rank in both sexes (females: $F_{1,8} = 14.87$, $p < 0.01$; males: $F_{1,6} = 13.87$, $p < 0.01$; Table S6). Light status did not influence activity between sampling periods for either sex (females: $F_{1,8} = 0.84$, $p = 0.39$; males: $F_{1,6} = 0.23$, $p = 0.65$; Table S6). However, light status did significantly influence activity in females within each sampling period (females: $F_{1,8} = 43.09$, $p < 0.0001$; Table S6), indicating that female activity in both high and low starvation resistance RILs was consistently higher during times when lights were on. Male activity of high and low starvation resistance RILs tended to remain constant under different light conditions. The larger influence of light status on females is consistent with patterns observed in activity during the pre-starvation period (Fig 4).*

Figure S10. *Starvation resistance QTL and estimated allele effects at each QTL. Data are presented as RIL means (±SE) for estimated starvation resistance when the founder haplotype was present in more than 5 RILs.*

Figure S10 continued.

Figure S11. *Triglyceride level QTL and estimated allele effects of founders at each QTL. Data are presented as RIL means (±SE) for estimated triglyceride level when the founder haplotype was present in more than 5 RILs.*

Figure S12. *Triglyceride level and starvation resistance were correlated after accounting for variation due to founder haplotype at the overlapping peaks TB3 and SB6 ($F_{1,7}$ = 7.72, P < 0.05). Data presented are averages for each founder haplotype in the pB panel, including "NA" for RILs that could not be assigned with confidence to a known haplotype. Point size relates the number of RILs per haplotype for the starvation resistance peak (smallest = 1 RIL; largest = 193 RILs); point color relates the number of RILs per haplotype for the triglyceride level peak (black = 1 RIL; lightest blue = 181 RILs). Grey shading around the regression line indicates the 95% confidence interval.*

Figure S13. *Overlap in SNPs associated with starvation resistance for each data set using the P < 10^{-5} significance threshold. Overlap between data sets was minimal. Plot A presents results for females; plot B presents results for males.*

Figure S14. *Additive effects of SNPs associated with starvation resistance in each study with a threshold of P < 10⁻⁵ calculated for starvation resistance measured in every other dataset. Female data is presented in the left column; male data is presented in the right column. SNPs that passed the FDR threshold of 0.2 are highlighted in red. Generally, SNPs had similar effects (of the same +/- sign) on starvation resistance as measured in each experiment.*

***Figure S15.** Manhattan plots of mean starvation resistance in the DGRP with SNPs that were associated with starvation resistance in previous studies and intervals of sex-specific QTL identified for starvation resistance in the DSPR highlighted. Plots are broken up by chromosome arm in A for females and in B for males. In all plots, points highlighted in black indicate SNPs that are associated with starvation resistance in the DGRP from data obtained in this study; red points indicate SNPs associated with starvation resistance in Everman and Morgan 2018; green points indicate SNPs associated with starvation resistance in Mackay et al. 2012; blue points indicate SNPs associated with starvation resistance averaged across the three datasets. A genomewide significance threshold of $P < 10^{-5}$ is shown with the blue line. Yellow shaded boxes and labels correspond to QTL intervals around peaks mapped in the pB DSPR panel; orange shaded boxes correspond to QTL intervals around peaks mapped in the pA DSPR panel. [Figure is provided as an independent file]*

Supplemental Tables

Table S1. *Analysis of variance of mean starvation resistance in the DSPR.*

Source	df	SS	MS	F	P
Subpopulation	3	330782	110261	183.702	< 0.0001
Sex	1	342197	342197	570.121	< 0.0001
Subpopulation x Sex	3	32982	10994	18.317	< 0.0001
Residual	3440	2064747	600		

Table S2. *Analysis of variance of mean triglyceride level in the DSPR.*

Source	df	SS	MS	F	P
Subpopulation	3	46.81	15.6020	37.099	< 0.0001
Residual	935	393.22	0.4206		

Table S3. *Stratification of the top 50 SNPs associated with starvation resistance across five frequency bins.*

Study	Sex	Allele Frequency Bin	No. SNPs
This Study	F	0.05 - 0.1	14
This Study	F	> 0.1 - 0.2	9
This Study	F	> 0.2 - 0.3	13
This Study	F	> 0.3 - 0.4	6
This Study	F	> 0.4 - 0.5	8
This Study	M	0.05 - 0.1	11
This Study	M	> 0.1 - 0.2	12
This Study	M	> 0.2 - 0.3	13
This Study	M	> 0.3 - 0.4	3
This Study	M	> 0.4 - 0.5	11
Mackay *et al.* 2012	F	0.05 - 0.1	28
Mackay *et al.* 2012	F	> 0.1 - 0.2	11
Mackay *et al.* 2012	F	> 0.2 - 0.3	5
Mackay *et al.* 2012	F	> 0.3 - 0.4	2
Mackay *et al.* 2012	F	> 0.4 - 0.5	4
Mackay *et al.* 2012	M	0.05 - 0.1	21
Mackay *et al.* 2012	M	> 0.1 - 0.2	13
Mackay *et al.* 2012	M	> 0.2 - 0.3	7
Mackay *et al.* 2012	M	> 0.3 - 0.4	2
Mackay *et al.* 2012	M	> 0.4 - 0.5	7
Everman and Morgan 2018	F	0.05 - 0.1	11
Everman and Morgan 2018	F	> 0.1 - 0.2	13
Everman and Morgan 2018	F	> 0.2 - 0.3	10
Everman and Morgan 2018	F	> 0.3 - 0.4	8
Everman and Morgan 2018	F	> 0.4 - 0.5	8
Everman and Morgan 2018	M	0.05 - 0.1	18
Everman and Morgan 2018	M	> 0.1 - 0.2	13
Everman and Morgan 2018	M	> 0.2 - 0.3	12
Everman and Morgan 2018	M	> 0.3 - 0.4	4
Everman and Morgan 2018	M	> 0.4 - 0.5	3

Table S4. *Analysis of variance of mean starvation resistance in the DGRP.*

Source	df	SS	MS	F	P
Sex	1	97213	97213	118.21	< 0.0001
Residual	334	274680	822		

Table S5. *Analysis of variance of activity during the 24-hour period prior to the DAM starvation assay with effect sizes (Cohen's F) for each variable.*

Source	df	SS	MS	F value	P	Effect Size
Starvation Resistance Rank (High vs. Low)	1	1.37	1.37	12.48	< 0.001	0.31
Sex	1	0.61	0.61	5.57	< 0.05	0.21
Lights On/Off	1	1.55	1.55	14.11	< 0.001	0.33
Starvation Resistance Rank x Sex	1	0.00	0.00	0.01	0.92	0.01
Starvation Resistance Rank x Lights On/Off	1	0.00	0.00	0.00	0.98	0.00
Sex x Lights On/Off	1	1.85	1.85	16.87	< 0.0001	0.36
Starvation Resistance Rank x Sex x Lights On/Off	1	0.02	0.02	0.21	0.64	0.04
Residuals	132	14.49	0.11	0.00		

Table S6. *Repeated measures analysis of variance across sampling period for activity during the DAM starvation assay for males and females.*

Female Activity During Starvation
Between Periods

Source	df	SS	MS	F value	P
Starvation Resistance Rank (High vs. Low)	1	11.15	11.15	14.19	< 0.01
Lights On/Off	1	0.66	0.66	0.84	0.39
Starvation Resistance Rank (High vs. Low) x Lights On/Off	1	0.48	0.48	0.61	0.46
Residuals	8	6.29	0.79		

Within Periods

Source	df	SS	MS	F value	P
Starvation Resistance Rank (High vs. Low)	1	19.06	19.06	115.58	< 0.0001
Lights On/Off	1	7.10	7.11	43.09	< 0.0001
Starvation Resistance Rank (High vs. Low) x Lights On/Off	1	0.16	0.16	0.99	0.32
Residuals	575	94.81	0.17		

Male Activity During Starvation
Between Periods

Source	df	SS	MS	F value	P
Starvation Resistance Rank (High vs. Low)	1	5.47	5.47	13.79	< 0.01
Lights On/Off	1	0.09	0.09	0.23	0.65
Starvation Resistance Rank (High vs. Low) x Lights On/Off	1	0.67	0.67	1.69	0.24
Residuals	6	2.38	0.40		

Within Periods

Source	df	SS	MS	F value	P
Starvation Resistance Rank (High vs. Low)	1	5.03	5.04	46.70	< 0.0001
Lights On/Off	1	0.32	0.32	2.93	0.09
Starvation Resistance Rank (High vs. Low) x Lights On/Off	1	0.12	0.12	1.14	0.29
Residuals	471	50.78	0.11		

Table S7. *Data from genes mapped to the region under QTL intervals for starvation resistance in the pA and pB DSPR mapping panels and triglyceride level in the pB DSPR mapping panel based on Flybase release version FB2018_1. Highlighted genes indicate those previously identified in QTL mapping studies of starvation resistance. [Table is provided as an independent file]*

Table S8. *Gene ontology analysis of genes that are included within QTL intervals for starvation resistance and triglyceride level. Gene lists used in each analysis included all genes unique to male and female analyses for each trait. GO analysis was performed using the PANTHER Overrepresentation Analysis (Released 2018-05-21) with Fisher's Exact test with FDR multiple test correction. No enrichment was found for starvation resistance in the pB panel or for starvation resistance in the DGRP in any dataset.*

Starvation Resistance GO Analysis: pA DSPR Panel

Category	Fold Enrichment	FDR
glutathione metabolic process (GO:0006749)	6.91	1.46E-04
cellular modified amino acid metabolic process (GO:0006575)	3.85	1.79E-02

Triglyceride Level GO Analysis: pB DSPR Panel

Category	Fold Enrichment	FDR
heat shock-mediated polytene chromosome puffing (GO:0035080)	42.66	4.25E-04
polytene chromosome puffing (GO:0035079)	38.39	3.35E-04
sensory perception of sweet taste (GO:0050916)	27.42	7.71E-04
detection of chemical stimulus involved in sensory perception of taste (GO:0050912)	19.47	7.16E-04
chaperone cofactor-dependent protein refolding (GO:0051085)	16	5.94E-03
'de novo' posttranslational protein folding (GO:0051084)	16	5.19E-03
protein refolding (GO:0042026)	15.36	5.65E-03
'de novo' protein folding (GO:0006458)	13.24	9.68E-03
cellular response to unfolded protein (GO:0034620)	12.8	4.87E-03
response to unfolded protein (GO:0006986)	12.44	4.78E-03
cellular response to topologically incorrect protein (GO:0035967)	9.95	1.06E-02
response to topologically incorrect protein (GO:0035966)	9.74	1.00E-02
cellular response to heat (GO:0034605)	8.93	5.11E-02
chaperone-mediated protein folding (GO:0061077)	8.93	4.79E-02
response to hypoxia (GO:0001666)	7.22	4.89E-02
sensory perception of taste (GO:0050909)	6.65	2.84E-02

Table S9. *Data from GWA, generated from the DGRP Freeze 2.0 pipeline, based on Flybase release version FB2018_1. All SNPs shown passed the $P < 10^{-5}$ significance threshold; highlighted SNPs passed the FDR threshold of 0.2. [Table is provided as an independent file]*

Supplemental File Descriptions
All are provided as independent files

File S1. Data generated in the analysis of starvation resistance, desiccation resistance, triglyceride level, and activity.

File S2. All code associated with the bootstrapping analysis of SNPs associated with starvation resistance measured in the DGRP in this study, Mackay *et al.* (2012), and Everman and Morgan (2018).

File S3. Raw LOD scores for starvation resistance in the DSPR pA panel.

File S4. Raw LOD scores for starvation resistance in the DSPR pB panel.

File S5. Raw LOD scores for triglyceride level in the DSPR pA panel.

File S6. Raw LOD scores for triglyceride level in the DSPR pB panel.

File S7. GWA results for DGRP starvation resistance collected in this study.

File S8. GWA results for DGRP starvation resistance collected in Mackay *et al.* (2012).

File S9. GWA results for DGRP starvation resistance collected in Everman and Morgan (2018).

File S10. GWA results for average DGRP starvation resistance across the three datasets.