1

Transcriptomic and functional analysis of the oosome, a unique form of germ plasm in the wasp *Nasonia vitripennis*

9   Honghu Quan and Jeremy Lynch*

10   University of Illinois at Chicago, Department of Biological Sciences, Chicago IL 60607

11   USA

12   *Author for correspondence

13   Honghu Quan email: hhq2@uic.edu

14   Jeremy Lynch email: jlynch42@uic.edu

Abstract

Background

The oosome is the germline determinant in the wasp *Nasonia vitripennis* and is homologous to the polar granules of *Drosophila*. Despite a common evolutionary origin and developmental role, the oosome is morphologically quite distinct from polar granules. It is a solid sphere that migrates within the cytoplasm before budding out and forming pole cells.

Results

To gain an understanding of both the molecular basis of the novel form of the oosome, and the conserved essential features of germ plasm, we quantified and compared transcript levels between embryo fragments that contained the oosome, and those that did not. The identity of the localized transcripts indicated that *Nasonia* uses different molecules to carry out conserved germ plasm functions. In addition, functional testing of a sample of localized transcripts revealed potentially novel mechanisms of ribonucleoprotein assembly and pole cell cellularization in the wasp.

Conclusions

Our results demonstrate that numerous novel and unexpected molecules have been recruited in order produce the unique characteristics of the oosome and pole cell formation in *Nasonia*. This work will serve as the basis for further investigation into the patterns of germline determinant evolution among insects, the molecular basis of extreme morphology of ribonucleoproteins, and the incorporation of novel components into developmental networks.

Keywords

42 Oosome, Germ plasm, Primordial Germ Cells, Germline, Nasonia, Drosophila, RNAseq, RNA

43 Interference

Background

Germline establishment is a crucial event for sexually reproducing organisms. Germline cells are

special in that they are able to generate all of the cell fates of the soma and to regenerate

themselves. There are two major strategies to specify the germline among animals: zygotic

induction and maternal provision. In zygotic induction, inductive signals from surrounding

tissues drive the establishment of germline fate, usually relatively late in embryogenesis, after

the transition from maternal to zygotic control [1]. In contrast, in the maternal provision mode,

the germ cells are specified by determinants called germ plasm that are synthesized and localized

during oogenesis and are the first group cells formed during embryogenesis. Germline

specification in this mode occurs very early in development, usually prior to the activation of the

zygotic genome [1, 2]. Classical experiments have shown that germ plasm is both necessary and

sufficient to establish the germline fate [3-5].

It is likely that the maternal provision mode of germ plasm evolved multiple times among the

animals, and this reflected in the molecular basis of germ plasm determinants [6-27]. In

vertebrates, germ plasm (where it exists) is dependent on the maternal localization of *bucky ball*

[28]. In insects, the gene products of *oskar* (*osk*) are both necessary and sufficient to induce germ

plasm, and thus, primordial germ cells (PGCs) [29-32]. Downstream of these nucleators is a suite

of highly conserved germline-associated molecules (i.e., Vasa (Vas), Nanos (Nos), Tudor (Tud),

etc.) that are recruited to the germ plasm where they presumably carry out functions ancestral for

the animal germline [30, 33-35].

There are several conserved properties of PGCs that must be conferred to naive cells by

germ plasm. One of these is a period of transcriptional quiescence germ cells undergo after

being specified. This feature makes the composition of a germ plasm even more important since

it means that mRNAs and proteins critical for at least the early germ cell functions need to be

provided in the germ plasm itself. Germ cells also usually become highly migratory later in

development as they seek to colonize the developing gonad. They are often highly enriched for

mitochondria and have specific metabolic needs [36]. Since they carry the genome that will be

passed to future generations, germ cells have enhanced mechanisms to prevent DNA damage and

to reduce the activity of transposable elements [37, 38]. Finally, they must have the capability to

induce pluripotency to their genome, as germline cells will be the source for all cell fates in the

eventual progeny.

Like all developmental processes, there is likely to be variation in the details of these

conserved features of germline determination, whether due to selective or neutral forces.

Pressures that could impact the composition of the germ plasm could be differential activity of

transposable elements in the genome, or a novel path for migration of the PGCs to the gonad, for

example. Novelties in embryogenesis may also drive germ plasm composition. For example, in

the parthogenetic, paedomorphic embryos of the midge *Miastor,* somatic nuclei undergo

significant chromosomal diminution in the early cleavages while the cell that takes up germ

plasm maintains the full complement of chromosomes, indicating that the germ plasm contains a

component that prevents the loss of chromosomes during the early cleavages [39].

Holometabolous insects are an ideal system with which to study how germ plasm evolves,

given that it is ancestral in this clade [35], the unparalleled levels of diversity, and the strong

baseline understanding germ plasm function derived from *Drosophila melanogaster*. Here we

focus on the parasitic wasp *Nasonia vitripennis* as a model to compare to the fruit fly. Like

Drosophila, *Nasonia* depends on Osk, Vas, and Tud to assemble germ plasm [35, 40]. However,

in contrast to the collection of small granules stably associated with the posterior pole that make

up the *Drosophila* germ plasm, the *Nasonia* germ plasm forms a very large, dense organelle

called the oosome (Figure 1). This highly divergent morphology strongly indicates that the

composition of the *Nasonia* oosome may diverge significantly from the polar granules of

Drosophila.

The behaviors of the oosome and the PGCs in *Nasonia* further imply a divergent

composition of the oosome. In freshly laid eggs, the oosome is tightly bound to the ventral-

posterior cortex of the embryo (Fig. 1A). When the zygotic nucleus forms and moves into the

interior of the embryo, the oosome detaches from the cortex and coalesces into a dense,

extremely large structure in the same central column as the nuclei. It migrates anteriorly,

sometimes being found near to 50% egg length, before migrating back to the posterior pole (Fig.

1B-D). As the cleavage nuclei migrate toward the cortex, the oosome flattens into a crescent on

the posterior pole of the embryo while a large bud protrudes from the pole (Fig. 1E). Typically,

two or three nuclei become associated with the bud and the oosome material. The bud pinches

off, and the nuclei rapidly individuate into pole cells (Fig. 1F,G).

This is distinct from pole cell formation in *Drosophila,* where each pole cell forms

individually upon association with a certain critical amount of pole plasm [41]. After dividing a

few times, the pole cells remain stable at the posterior pole of the egg until gastrulation, when

they migrate through the posterior epithelium and coalesce into to two masses, presumably

where the primordial gonads are developing (Fig. 1H). This migration is distinct from

Drosophila pole cells which remain external on the posterior pole as the germ band undergoes

massive extension well after gastrulation and migrate internally through the gut wall [41].

Thus, it is clear that *Nasonia* and *Drosophila* share some fundamental aspects of germline

establishment, but they also have their own diverged features that fit in their own embryogenesis

program. This raises the question of which genes are the core components for the maternal

provision mode and which genes contribute to their own distinct features in germline

development.

To address these questions, we compared the mRNA content of anterior and posterior

halves of the pre-blastoderm stage *Nasonia* embryos in an effort to identify the components

specifically localized to the oosome. We found only a few of mRNAs conserved in both fly polar

granules and the *Nasonia* oosome, such as *osk*, *nos*, and *ovo* (see Supplemental File 1). The rest

are all novel components germ plasm that either lack *Drosophila* homologs or have homologs in

Drosophila that do not play any roles in germline development. We performed functional studies

for a set of localized transcripts, all of which showed roles in the unique features of *Nasonia*

germline development, demonstrating the value of our approach to identify the molecular

sources of novelty among various insect lineages.

Results

RNA-seq analyses of the anterior and posterior poles of the wasp *Nasonia* early embryos

To identify the maternal transcripts in the oosome, we isolated the total RNA separately from

anterior and posterior poles of the pre-blastoderm stage *Nasonia vitripennis* embryos. Six

samples that comprised three sets of each pole were sequenced and subjected to differential

expression analysis, which are described in more detail in the Methods.

Our first attempt yielded encouraging, but mixed results. Several novel posteriorly

enriched factors were identified, and we did recover some known posterior factors (*Nv-osk* and

Nv-nos). However, other known localized factors such as *Nv-cad* and *Nv-dpp* were not found to

be statistically significant. In addition, one of the posterior samples was of poor quality, and

could not be used in the differential expression analyses, reducing the power of our approach.

Since we wanted results as comprehensive as possible, we repeated the experiment using a few

adjustments (see Methods) and obtained much more robust results. While this second experiment

was being prepared, we moved forward with expression and functional analysis of novel

candidates identified in the first. Since a handful of transcripts with confirmed oosome

localization were found only in the first experiment, the remainder of the results will include

these transcripts along with those identified in the second experiment.

All known maternally localized molecules (*Nv-osk*, *nos*, *dpp*, *cad*, *otd1*, and *gt*) were

found with high significance, giving confidence that our experimental design and analysis

approaches were appropriate for our goal of discovering the mRNAs localized to the oosome.

Overall, we found 92 transcripts with apparent significant enrichment at the posterior pole. These

ranged in levels of enrichment from 1.4 to 55 times higher in the posterior fragments compared

to anterior fragments. Our analyses also uncovered anteriorly enriched mRNAs, of which there

were 89, with a range of fold enrichment from 1.4 to 10 times higher at the anterior.

Novel transcripts localized in the anterior half of the *Nasonia* early embryos

While the anterior factors are not the focus of this manuscript, some interesting observations

were made in examining a small sample of candidate mRNAs. Most transcripts are localized in

small domains at the anterior cortex and seem to extend toward the posterior in variable tendrils,

rather than being uniform or graded caps of anterior localization (Fig. 2A1-K1). A notable

exception is the transcript of *Nasonia* homolog of *mex-3,* an RNA binding protein known for

controlling translation of orthologs of the posterior patterning factor *caudal* in the nematode *C.*

elegans and the beetle *Tribolium* [24, 42]. *Nv-mex3* mRNA is localized in a broad domain

extending far toward the posterior of the embryo (Fig. 2L1). *Nv-mex3* expression is highly

dynamic and variable in both the blastoderm and post-gastrular stages (Fig. 2L2-L3, additional

images not shown).

Transient localization is the most common feature of the anteriorly localized transcripts.

Most are ubiquitous or absent by the time the early syncytial blastoderm forms (Fig. 2 A2-H2,

J1-K3, M1-M3), except for *LOC100313502* which persists into the blastoderm stage where it

forms a anterior cap (Fig. 2 I1-I3). Among the anteriorly enriched transcripts, many have

predicted functions that may be relevant to egg activation (e.g. four encoding ion channels), and

anterior-posterior polarity and patterning (e.g. transcription and translation factors, see

Supplemental file 1). There are also a large number of transcripts with no clear homologs outside

of *Nasonia* see (Additional File 1). One of these (*LOC100119982*, Fig. 2G1-G3) is a member of

a novel family of ankyrin domain containing molecules that are specific to Chalcid wasps and

appear to have obtained a broad diversity of expression and potential function during *Nasonia*

development [43].

Finally, a handful of molecules localized at both poles were detected. These include the

known factors *Nv-otd1* [44] and *Nv-TollC* (JAL in preparation)*,* along with *Nv-endoglucanase*

(Fig. 3E), *Nv-insulin-like growth factor (Nv-igf,* Fig. 3C)*, Nv-ucth* (a ubiquitin carboxy terminus

hydrolase, Fig. 2M)) . The fact that these were found despite lower apparent fold differences

between the two embryonic halves, further gave confidence that our analysis was robust enough

to detect even subtle germ plasm localization of the vast majority of mRNAs.

General description of the novel transcripts localized in the posterior pole of the *Nasonia*

early embryos

From the two analyses, we identified 92 candidate transcripts that were statistically enriched at

the posterior half of the *Nasonia* embryos. We then isolated PCR products for 54 of these genes

and made probes to determine their expression patterns during early embryogenesis. We were

able to confirm 47 transcripts that are expressed posteriorly during pre-blastoderm stage. Of the

remaining transcripts, one of them was not successfully cloned, and the rest were successfully

cloned but did not have any specific, localized expression. Interpretation of these potential false

positives (as well as some false negatives) will be discussed later. Among the 28 transcripts with

expression in the posterior half of the embryo, we grouped them into three categories based on

their expression patterns: 1) genes expressed in oosome and with expression maintained in the

PGCs (Fig. 3), 2) mRNAs localized in the oosome but then degraded in the PGCs (Fig. 4), and 3)

mRNAs strongly enriched in the posterior region of the embryo, but not incorporated in the

oosome (Fig. 5).

Highly germline-associated transcripts

We identified 11 transcripts that are localized to the oosome and maintained in the PGCs at pole

cell formation and beyond. We consider these to be the strongest candidates for having important

roles in the specification and function of the PGCs but cannot exclude *a priori* that they have

other (or no) important functions. Six transcripts are maintained throughout the stages of

germline development followed in this manuscript: oosome, pole cells, and migrating germ cells.

They include *Nasonia* homologs of the fly genes *bark beetle* (*Nv-bark*) (Fig. 1), *spt5* (*Nv-spt5*),

tejas (*Nv-tdrd7*), *insulin-like growth factor* (*Nv-igf*), and two transcripts without fly homologs

(*Nasvi2EG001470* and *Nv-endoglucanase*) (Fig. 3A1-E3).

204   Among these, *Nv-bark* is the best germline marker, bearing strong and consistent

germline association throughout early development, including expression in the late embryonic

gonads (Fig. 1). In this respect, it is better than our previously favored marker, *Nv-nos*, which is

downregulated significantly toward the end of PGC migration [40]. Bark is a large

transmembrane protein and is not expressed in the *Drosophila* germline. Its only known role is in

stabilizing tricellular junction in epithelial cells during embryogenesis [45, 46]. It is not clear

how this function is relevant to germline function, and indicates a novel recruitment of this factor

in the wasp.

212   *Nv-spt5* is significantly enriched in the oosome with low levels of ubiquitous expression

213   in the rest of the embryos in pre-blastoderm stage (Fig. 3A1). In the blastoderm and migrating

214   germ cell stages expression is reduced but still enriched in the pole cells while the ubiquitous

215   expression in the embryo persists (Fig. 3A2-A3). Spt5 homologs are involved in regulating RNA

216   polymerase progression during transcription [47], which might indicate that Nv-Spt5 is involved

217   in repressing or otherwise regulating the onset of transcription in the germ cells.

218   *Nv-tdrd7* is present at appreciable levels throughout the bulk cytoplasm and is also

strongly localized in the oosome (Fig. 3B1). This pattern is well reflected in the quantification of

mRNA levels in the two halves of the embryo, which show significant numbers of reads coming

from the anterior half of the embryo. At the blastoderm stage, *Nv-tdrd7* is moderately enriched in

the pole cells and is zygotically expressed in a ventral-posterior patch (Fig. 3B2), which was

detected in our earlier analysis of dorsal-ventral patterning [48]. After gastrulation, *Nv-tdrd7* is

strongly upregulated in a group of cells that are near to where the germ cells migrate, but it is not

clear if they are germ cells (Fig. 3B3).

Nv-igf is initially expressed in a bipolar pattern (similar to *Nv-otd1*[44]), before becoming

specific to the pole cells during the blastoderm stage and the migrating germ cells after

gastrulation (Fig. 3C1-C3).

Nasvi2EG001470 encodes a short peptide of 80 amino acids and was not included in the

most recent annotation of the *Nasonia* genome at NCBI, but was present in OGS 2.0 [49]. A very

similar sequence is annotated in the close relative *Trichomalopsis* (JAL personal observation),

indicating that it is a *bona fide* transcript that is either novel, or very rapidly evolving.

Nasvi2EG001470 is strongly expressed in the oosome and pole cells, while levels markedly

decrease in migrating germ cells (Fig. 3D1-D3).

Besides the expression in the oosome and the pole cells, *Nv-endoglucanase* is initially

expressed at both poles during pre-blastoderm stage and early blastoderm stage (Fig. 3E1-E2).

Later in blastoderm stage, the expression is down-regulated at the anterior pole and become

specific to the pole cells (Fig. 3E3). Proteins of this type are found extensively in Hemimetabola,

Hymenoptera, and Coleoptera, and appear to have been lost at the origin of Lepidoptera and

Diptera clades. It is intriguing that what roles this protein plays during *Nasonia* embryogenesis.

Five transcripts localized to the oosome enter and are maintained in the pole cells but are

then downregulated in the migrating germ cells (Fig. 3F1-J3). This set includes homologs of

Drosophila anillin (*scraps*), *qin, and innexin1 (ogre),* (*Nv-anillin*, *Nv-qin*, and *Nv-innexin1*,

respectively). Anillin is an actin binding protein that localizes to the contractile ring during

cytokinesis [50]. In *Drosophila* Anillin protein is localized in the cleavage furrows when forming

the PGCs [51], but its mRNA is ubiquitous. The early oosome localization of *Nv-anillin* mRNA

suggests that the protein might also play roles in oosome outside of a potential conserved role in

pole cell formation.

Nv-qin encodes a protein containing tudor domains along with an E3 ubiquitin ligase

domain. *qin* is important in processing germline piRNAs, repressing retroelements assembling

the nuage, and proper completion of oogenesis in the fly [52-54]. While *qin* has an important

late role in germline cells, it is only weakly and diffusely expressed during embryogenesis in

Drosophila [55]. Its mammalian homolog *RNF17* is required for production of specific particles

in the germline nuage and for sperm development, but not for early germline specification [56].

Nv-innexin1 encodes a putative gap junction protein whose fly homolog is most well-

known for its role in proper development and function of the nervous system [57]. Other

unexpected roles for innexin proteins have been described and proposed in insects [58], but at the

moment the potential functional significance of the germline localization in *Nasonia* is unclear.

Two transcripts localized to the oosome and preserved in the pole cells have not clear

Drosophila homologs (Fig. 3I1-J3). One of these (*Nv-C16orf87*) encodes a homolog of the

human protein *C16orf87* and is expressed in the posterior region as well as specifically in the

oosome in pre-blastoderm stage (Fig. 3I1-I3). The protein of *Nv-C16orf87* belongs to the

uncharacterized protein family UPF0547, which contains the zinc-ribbon motif, and functions of

this protein and it homologs are not known. It appears that this gene has been lost specifically in

the Brachyceran fly lineage as it is found in beetles, moths, and some mosquitos, but not

Drosophila (JAL personal observation).

Finally, another ankyrin domain encoding transcript is strongly localized to the oosome

and is taken up into pole cells (Fig. 3J1-J2). It later has a complex and dynamic pattern in the

blastoderm stages (Fig. 3J2-J3, additional images not shown). This transcript is a member of the

newly described CLANK (Chalcid Lineage-specific ANKyrin-domain gene) family, of which

there are nearly 200 in the *Nasonia* genome [43]. To differentiate from the others, we name it

Nasonia   vitiripennis   Oosome   CLANK   (Nv-OoCLANK).

Transcripts enriched in the oosome but excluded from pole cells

A set of 12 genes is expressed in the oosome but not transported to the pole cells. We predicted

that these would have germline roles primarily in the oosome itself, or in the early stages of pole

cell formation. We also expect many transcripts in this set will have roles outside of germline

production, such as in embryonic patterning (as already known for *Nv-dpp* and *Nv-cad*) [59, 60].

Examples of what we consider potential embryonic patterning factors include: a CLIP protease

encoding message (*Nv-mp1*) related to fly Melanization Protease and Easter (Fig. 4A1-A3), *Nv-*

kayak (Fig. 4 B1-B3) encoding at transcription factor downstream of JNK signaling[61], and *Nv-*

elbow (Fig. 4C1-C3) encoding a single zinc-finger transcription factor [62]. *Nv-kayak* is later

expressed in a dorsal domain, indicating a conserved role in extraembryonic patterning.

Interestingly, *Drosophila elbow* interacts with *orthodenticle* in specifying the ocelli and in

photoreceptor cell fate determination [63], and an intriguing possibility is that these proteins

work together in posterior patterning, with *Nv-elbow* possibly playing an important role in

differentiating the posterior targets of *otd* from the anterior ones.

Several oosome resident transcripts have suggestive functional annotations. For example,

the *Nasonia coronin* gene (*Nv-coronin,* (Fig. 4D1-D3)) encodes a protein whose homologs are

known to bind and modulate actin, provide links between the actin and microtubule

cytoskeletons, and regulate endo- and exocytosis in several developmental contexts [64, 65]. A

germline role for the *Drosophila* Coronin ortholog has not been observed. *Nasonia milton* (*Nv-*

milton) is another exciting transcript (Fig. 4E1-E3). *Drosophila* Milton is an adaptor protein that

allows mitochondria to be loaded onto, and transported by, microtubule motors [66].

An oosome resident mRNA encodes a protein whose fly ortholog is uncharacterized, but

whose function may be relevant to oosome function. This is the *Nasonia* homolog of *CG4552*

(*Nv-CG4552*), which encodes a protein with a TBC25 domain (Fig. 4F1-F3). Proteins with this

domain interact with Rabs to regulate membrane trafficking and dynamics. Such activities have

been shown to be crucial for Osk function in the fly [67], and *Nv-CG4552* may play a supporting

role in regulating membrane dynamics in the wasp. Another suggestive localized factor does not

have clear orthologs outside of the hymenoptera, but it does have two predicted RNA

Recognition Motifs, therefore we name it *Nv-rrm* (Fig. 4G1-G3). RRM domains bind RNA and

are components of proteins that regulate RNA localization and translation. This novel lineage

specific protein could therefore be involved in the localization of specific RNAs in the oosome,

or the regulation of translation of specific RNAs within it.

Many of the oosome localized transcripts do not have annotations that lead to simple

hypotheses about their roles in specifying the germline. One of these is *Nv-waterproof*, which

encodes a fatty acyl-CoA reductase. *Drosophila waterproof* produces the hydrophobic molecules

that coat the trachaeal tubes during *Drosophila* embryogenesis and is essential for gas filling of

the trachea [68]. The protein's novel role in *Nasonia* germline is not clear and worth to

investigate, as is its early maternal expression in the oosome and later zygotic expression as the

dorsal strip and in the extraembryonic tissue. (Fig. 4H1-H3). *Nv-CG42269* encodes a predicted

organic ion transporter protein whose *Drosophila* homolog (*CG42269*) has no described function

(Fig. 4 I1-I3).

Three oosome localized transcripts have no clear homologs in *Drosophila* or in other

model organisms. *LOC100123551* has a sterile alpha motif (SAM) domain, which might indicate

protein-protein or -RNA interactions (Fig. 4J1-J3). *LOC103315681* contains weak similarity to

the N-terminal domain of Folded-gastrulation proteins (but is not a folded gastrulation ortholog)

(Fig. 4 K1-K3), while *Nasvi2EG022821* has no discernible conserved domains (Fig. 4L1-L3).

The functions of these factors will be the object of future investigation.

Transcripts enriched in the posterior pole but not specifically the oosome

There are five transcripts that are significantly enriched in the posterior region of the embryo, but

do not show significant association with the oosome (Fig. 5). The early embryonic expression for

these transcripts appears as a cap or broad posterior to anterior gradient of mRNA. The

significance of such transcripts to oosome assembly or to germ cell formation is not clear. Two

of this class are known developmental transcription factors: orthologs of Zerknuellt [69], and

Mothers against dpp [70] (*Nv-zen* and *Nv-mad2,* respectively (there are two mad paralogs in

Nasonia). Two transcripts are predicted to encode catalytic enzymes: a choline kinase

homologous to the *Drosophila CG2201* (*Nv-CG2201*), and a homolog of the ADP ribosylation

factor-like 4 protein (*Nv-ARL4*).   Finally, *LOC100680197* and *LOC100677932* have no

identifiable homologs outside of hymenoptera. *LOC100680197* encodes a protein with MYND-

type zinc-fingers and a p27-like domain, while *LOC100677932* has no clear conserved or

functional domains.

Functional analysis by parental RNA interference showed low phenotypic penetrance

While localization of an mRNA to the oosome and pole cells may strongly suggest a function

related to PGC specification, demonstration of any such function is required. We chose a

sampling of five promising molecules for in depth functional analysis (*Nv-bark*, *Nv-anillin*, *Nv-*

rrm, *Nv-coronin*, and *Nv-innexin1*).

We initially tried to apply our parental RNAi (pRNAi) approach [44], but quickly found

that this was not the ideal approach. Most dsRNAs caused partial sterility, with most of the

obtainable eggs being apparently normal escapers. Eventually, we managed to collect embryos

with phenotypes in a very low penetrance (2%-6%) from three of the five genes that we studied

with the dsRNA concentration of 1ug/uL to 2.5ug/uL. These genes are *Nv-rrm*, *Nv-coronin*, and

Nv-innexin1. They all either have no pole cells formed or less pole cells formed with and

disorganized germ plasm residue at the posterior pole of the embryos in blastoderm stage.

Infrequently, *Nv-coronin* knockdown embryos were characterized by pole cells that did not

migrate to the gonad, but instead remained at the pole after gastrulation. (data not shown).

This issue was more serious for *Nv-bark* and *Nv-anillin*. At concentrations from

1.5ug/uL-2.5ug/uL), there were no eggs laid. At lower concentrations (250ng/uL, 500ng/uL and

750ng/uL), the eggs we collected all showed normal development. Since phenotypic embryos

were either completely absent or extremely rare for our genes of interest, it became necessary to

develop a new technique to assess the functions of the novel oosome genes we discovered.

Development of an embryonic injection protocol for *Nasonia* RNAi

To circumvent the low penetrance problem from the pRNAi, we developed a protocol for

embryonic injection of dsRNA followed by fixation and *in situ* hybridization (see details in

Methods and Materials). As a negative control, we injected dsRNA against eGFP to test whether

the physical injection and the dsRNA itself would affect the structure of the oosome and

formation of the pole cells non-specifically. We were happy to find that even in embryos with

obvious physical damage, the oosome and pole cells could form normally (Fig. 6A1-A4).

Of course, at a certain point, damage becomes too severe, leading to the death of the

embryo. We set a stringent criterion for collecting embryos for later analysis by removing those

where yolk leakage exceeded more than 10% of the embryo size (although all embryos with this

amount of damage showed normal germ plasm and germline development) (Fig. 6A2, A4). We

also excluded embryos that showed major morphological changes as compared to uninjected

embryos.

After removing the embryos with obvious major damage both after injections, and when

performing eggshell dissection after the fixation, we were left with about 85% embryos with

viable embryogenesis by the time of imaging after *in situ* hybridization. The same criteria were

also applied to the experimental knockdowns where the percentages were roughly the same as in

the negative control When determining the phenotypes for the five genes, we considered the

disruptions of developmental events to be potential effects of a knockdown when they were only

specific to the knockdown and were never observed in the negative control.

We performed the same procedures when injecting and collecting the embryos as

described for the negative control. The penetrance given by the embryonic RNAi (eRNAi)

knockdowns is higher than the pRNAi knockdown, ranging from 20% to 39% across

experiments. Phenotypes are evident in both the pre-blastoderm and pole cell stages, for *Nv-bark*,

Nv-anillin and *Nv-rrm*. In contrast, the oosome is not affected in pre-blastoderm embryos after

Nv-coronin and *Nv-innexin1* dsRNA injection, with phenotypes becoming evident only after the

pole cells should have formed.

RNAi against three novel germ plasm components unexpectedly disrupts the oosome at

from an early stage

One transcript that particularly captured our attention was *Nv-bark*, as it was the transcript most

strongly and consistently associated with PGC specification over embryonic development.

However, its potential function in the germline is not clear. Since it encodes a transmembrane

protein involved in epithelial junctions in *Drosophila*, we speculated that it might have a role in

mediating adhesion or migration of the pole cells once they were formed. Surprisingly, the

phenotypes produced by knocking down this transcript showed a much earlier requirement for

this transcript. In early embryos (before migration of nuclei to periphery), the oosome has lost

its integrity as a single unit (Fig. 6B1). Instead there are scattered particles of what appears to be

oosome-like material, concentrated around the lateral cortex of the embryo. Later, at the time the

pole cells would normally form, no budding is observed, and some loose aggregates of germ

plasm like material remains attached to the cortex (Fig. 6B2-B3). In some cases, germ plasm

surrounds nuclei in a way similar to what is seen in pole cells (Fig. 6B3, B4). However, these

nuclei remain part of the embryonic syncytium.

Like *bark,* the mRNA of *Drosophila anillin* is not localized to the polar granules [71].

Anillin protein, however, accumulates at the base of pole cells when they are budding in

Drosophia [51]. Since Anillin is a major component of the contractile ring during mitotic

cytokinesis [50] and is enriched at the bud furrow during pole cell formation in *Drosophila* [51],

we predicted that the enrichment of *Nv-anillin* mRNA in the oosome and pole cells would be

related to *Nasonia's* unique way of forming a single large pole bud instead of several small ones,

as occurs in *Drosophila*. Surprisingly, the phenotype of *Nv-anillin* is indistinguishable from that

of *Nv-bark*: The oosome does not form properly, and germ plasm material remains bound to the

posterior cortex of the embryo, leading to the association of the syncytial nuclei at random

posterior locations (Fig. 6C1-C4). The same set of phenotypes is observed when we knock down

Nv-rrm with eRNAi (Fig. 6D1-D4). Thus, three genes with very different predicted functions all

result in the same phenotype.

It is important to note at this point that the common phenotype of the above three

knockdowns is not the same as the complete loss of germ plasm activity. Such phenotypes are

seen for *Nv-osk*, *Nv-vas*, and *Nv-aubergine (Nv-aub)*. When these genes are knocked down,

posterior mRNAs such as *Nv-nos* take on a uniformly graded posterior cap and no enriched

accumulation of germ plasm markers is ever observed at the posterior [35, 40]. This indicates

that these transcripts are involved in the specific form of the oosome, but may not be essential for

the production of germline-like cells.

Finally, despite the fact that the oosome does not form or migrate through the embryo as

in wildtype, a bud like protrusion similar to the one that initiates pole cell formation is regularly

observed (Fig. 6C3, Fig. 1E), and this bud is not associated with the bulk of the germ plasm like

material. This indicates that neither the oosome, nor its remnants, induce the bud, and that bud

formation may be autonomous to the embryo. This has some precedent in *Drosophila*, where the

autonomous ability of the fly embryo to produce pole-cell like structures at both poles of the

embryo is revealed when Arf guanine exchange factor Steppke is reduced [72]. However, in the

fly, normal global repression of pole-cell formation is overcome by germ plasm components

(primarily *germ-cell-less*), while in *Nasonia,* at least the initial budding appears to be germ plasm

independent.

RNAi against *Nv-coronin* and *Nv-innexin1* does not affect the oosome, but disrupts pole cell

formation

We tested the functions of two other transcripts with eRNAi: *Nv-coronin* and *Nv-innexin1*.

Knockdown of both genes left the oosome intact, and able to migrate through the embryonic

cytoplasm normally (Fig. 6E1, F1). However, in both cases, pole cell formation fails, indicating

that these genes have downstream functions that are specific to pole cell formation (Fig. 6E2-E4,

F2-F4). *Nv-coronin*'s functional annotation is consistent with a role in cellularization, as it is

predicted to interact with both the microtubule and actin cytoskeletons, both of which are crucial

for mitosis and cell formation. While Innexin1 is most well known as a crucial component of

gap junctions and a role in the nervous system [57], a role for at least one Innexin in cellurization

has been demonstrated in the beetle *Tribolium* [58], suggesting that the full potential of these

proteins in regulating cellular processes has not been fully explored.

Discussion

RNA-seq analyses

Our results have uncovered an unexpectedly large amount novelty in the mRNA content of the

germline determinant of the wasp *Nasonia vitripennis*. This was achieved using RNAseq

followed by statistical detection of differential expression of mRNAs between the anterior and

posterior poles of the early embryo. The statistical predictions were then extended by *in situ*

hybridization and RNAi of candidate genes.

Our goal was to identify all transcripts specifically localized to the oosome, and it is

worth contemplating how close we came to achieve this goal. We used three different

approaches that varied in the strategy for mapping, quantifying transcripts and assessing

differential expression. These analyses agreed on the vast majority of the genes with putative

significantly different enrichment at the two poles, and most of the disagreement was at the

margin of differential expression that would be likely biologically significant. In addition, all

analyses found genes previously identified to be localized anteriorly or posteriorly. We even

detected all known genes with both strong oosome localization and strong localization at the

anterior pole. We thought these could be missed, because if the anteriorly localized mRNA

population approached the same levels as that in the oosome, the enrichment might be obscured.

For these reasons, we believe that we have uncovered the vast majority of mRNAs

localized to the *Nasonia* oosome. Of course, no approach can guarantee comprehensiveness, and

some molecules may have been missed for several reasons. For example, while the *Nasonia*

genome is well annotated, it is possible that a very few genes are not represented in the genome

assembly or the transcriptome annotations, and thus would not have been assessed. Indeed, we

did identify slightly different sets of genes when using NCBI annotation 102 versus the OGS 2.0

annotation [49], which were created using different approaches to identify and predict genes.

Most of the differences are at the margins of significance and low levels of differential

expression. However, there are a handful of confirmed localized transcripts detected using

annotation 102 that were missed in analyses with OGS 2.0. In addition, at least one transcript

was not annotated in annotation 2.1, but was found in annotation 2.0 and was confirmed to be

localized to the oosome (*Nasvi2EG001470*, Fig 3D). This is likely a false negative annotation in

annotation 102, as a very similar sequence is found in the transcriptome of the very closely

related wasp *Trichomalopsis sarcophagae* [73].

Another issue that can cause false negatives is the large number of *Nasonia* transcripts

that overlap, which can lead to the concatenation of transcripts. If a localized transcript is fused

to a ubiquitous, highly expressed transcript, the signal of localization can be lost. This problem

is largely solved by preventing novel junctions when mapping with tophat2. However, this also

seem to change the calculation of significance for some transcripts, so performing the analysis

with and without novel junctions gives more complete results.

Finally, there are some unknown artifacts which may cause significantly enriched genes

to be missed. A prime example of this is our discovery of *Nv-coronin*. A preliminary attempt at

this experiment resulted in generally poor sequencing results, and a completely unusable

replicate. In general, this analysis gave predictably worse results, where several known localized

genes were not found to be significant. However, all of the molecules whose functions were

analyzed in this manuscript were found to be statistically significantly enriched at the posterior,

including *Nv-coronin*, and we proceeded to clone them to test their localization and function. In

the subsequent analyses based on high quality sequencing results (used as the basis for this

manuscript), *Nv-coronin* was excluded by cuffdiff for statistical testing for unknown reasons,

despite showing similar posterior enrichment that had previously been deemed statistically

significant. Thus, there is a potential for false negatives if this artifact affects several genes.

Very few transcripts in the "not tested" category show a similarly strong posterior enrichment, so

we believe that this effect is also small.

Comparison of the oosome of *Nasonia* to the polar granules of *Drosophila*

Germ plasm in insect embryos must perform several different functions. First, it must be able to

concentrate and arrange the set of proteins and RNAs (e.g. Tud, Vas, *nos* mRNA, Piwi/Aub) that

are associated with germline fate in the posterior of the egg. Second, it must ensure its own

incorporation into the PGCs as they form by interacting with the specialized cytoskeletal

structures that mediate pole cell formation. Germ plasm must also include molecules involved

specialized features of PGCs.   Such functions include the concentration and selection of

mitochondria, repression of transcription, repression of transposons, and guidance of the germ

cell migration to the gonad primordia.

On the other hand, we had already observed that the oosome's morphology and

interaction with the cytoskeleton (during its migration and formation of the pole cells) was quite

distinct from *Drosophila*, as is the way the pole cells migrate into the interior of the embryo.

Therefore, we expected to find a mixture of conservation and novelty when we examined the

mRNA content of the oosome. Indeed, this is what was found, but with a surprisingly strong bias

toward novelty.

In terms of conservation, our analyses found *Nv-nos, Nv-osk* (already known factors

found in both oosome and polar granules) as well as *Nv-ovo*. We also found *Nv-aub* whose

ortholog is localized as protein, but not mRNA, in *Drosophila*. While there are a large number of

polar granule localized mRNAs that are not found in the oosome, we will only discuss a few that

are significant.

mRNA for *polar granule component (pgc)* encodes a small peptide [74], and is strongly

localized to the posterior pole [75]. Polar-granule-component protein has a crucial role in the

global repression of transcription that occurs in pole cells upon their formation, through an

interaction with the transcription elongation factor TEF-b [74]. This repression is a widely

conserved feature of PGCs across animals, which makes it somewhat surprising that *pgc* appears

to be a novelty in the *Drosophila* lineage [75].

Fascinatingly, TEF-b may be a unifying factor underlying germline quiescence, as it

interacts with PIE-1 in the worm *C. elegans* to repress transcription [76]. In line with this, we

have found a highly conserved transcription elongation factor, *Nv-spt5* (Fig. 3 A1-A3) localized

strongly to the oosome and pole cells. In human cells, Spt5 acts as an inhibitor or transcriptional

elongation, until its C-terminal domain is phosphorylated in a TEF-b dependent manner [77]. If

an interaction between *Nasonia* Spt5 and TEF-b does have a role in regulating the cessation of

transcription in *Nasonia* pole cells, it would be strong evidence for TEF-b being a core conserved

component of the germline fate, whose interaction partners and regulators are labile across

lineages.

Another crucial *Drosophila* germ cell factor that is not present in the *Nasonia* oosome is

germ cell less (*gcl*). The Germ-cell-less protein itself is very highly conserved at the sequence

level in *Nasonia,* but the mRNA showed no enrichment in our RNAseq experiments, and we

independently confirmed by *in situ* hybridization that it is expressed uniformly throughout the

early embryo (not shown). Gcl is important for the proper production of pole cells, apparently by

regulating the orientation centrosome separation at the posterior pole, which is required for

efficient pole cell formation and uptake polar granules by the pole cells [78]. At the molecular

level, Gcl seems to act by downregulating torso signaling, to allow the proper conditions for pole

cells to form. The lack of Gcl function in the germline of *Nasonia* is consistent with the lack of

Torso signaling at the termini in the wasp [79], making the need for Gcl redundant. At the

moment, it is not clear whether the use of Gcl in pole cell formation is a recent novelty in

Drosophila, or whether it was present ancestrally, but lost in the Hymenopteran lineage.

A number of transcripts found in the *Nasonia* oosome are good candidates for generating

essential PGC features. For example, a high concentration of mitochondria is a strongly

conserved feature of germ plasm and PGCs across animals [80, 81]. In *Drosophila*, the long Osk

isoform plays an important role in concentrating mitochondria in the pole plasm [82]. But, since

the long form of Osk appears to be a novelty of *Drosophila* and its close relatives, other

molecules should be expected to perform this role in other species. Suggestively, mRNA

encoding a Milton ortholog was found strongly localized to the *Nasonia* oosome. Milton acts an

adaptor that loads mitochondria onto microtubule motors for transport and localization within

and between cells in *Drosophila* [83], and we propose that *Nasonia* Milton plays a role in

enriching mitochondria around the oosome and in the pole cells in the wasp, and perhaps other

insect species that lack the specialization of long Osk isoform.

Another critical function for germ cells is the control of transposable elements, which is

often dependent on Tudor domain containing proteins. mRNAs for two Tudor domain proteins

are present in the oosome, including *Nv-qin* and *Nv-tdrd7*. Neither of them is enriched in the

polar granules or pole cells in *Drosophila*, but both have crucial roles during oogenesis to reduce

the activity of transposable elements [52, 53, 84]. The presence of these additional Tudor

domain encoding transcripts may indicate that either there is an increased activity of transposable

elements in *Nasonia* that requires an earlier response, or perhaps other mechanisms are employed

in *Drosophila* to combat transposon activity in the early PGCs. Further sampling of germ plasm

of other insects should help to resolve these questions.

Germ cells are known to have a distinct metabolic profile from somatic cells, and this

difference is related to their pluripotent stem cell-like properties, and to the requirements of their

migratory properties [85, 86]. Potentially related to this we have found that a *Nasonia* insulin-

like growth factor I mRNA is localized to the oosome. Interestingly, this mRNA encodes a short

protein that shows much stronger similarity to insulin proteins of vertebrates, than it does to any

of the insulin-like molecules of *Drosophila* (DILPs) (JAL personal observation). In addition, an

mRNA encoding a putative organic cation transporter (*Nv-CG42269)* containing a Major

Facilitator Superfamily (MFS) domain is strongly localized to the oosome. Such molecules are

crucial for regulating cellular metabolism and signaling at multiple levels, by controlling the

trafficking of many small organic molecules (including sugars) within and between cells [87, 88].

Lipid metabolism is also uniquely regulated in germ cells, and the identification of the *Nasonia*

homolog of the Acyl-CoA reductase *waterproof* may reflect this [89]. Finally, we surprisingly

found a transcript encoding a protein similar to endoglucanases found in several insect lineages

(but absent from Diptera). Enzymes of this type are broadly defined by their ability to break

down polymers glucose. The substrate and potential role for this enzyme in the germline is not

yet known.

In addition to providing insight into the conserved functions of germ cell components, we

also found several molecules that do not have clear homology outside of the Hymenoptera, or in

some cases outside of *Nasonia* and its closest relatives. This includes a novel RNA recognition

domain containing protein whose function we analyzed in depth (*Nv-rrm* discussed in the

following section). We also found that an mRNA encoding an ankyrin domain protein (*Nv-*

OoCLANK (*LOC100679945*)) that belongs to a family of proteins that underwent a massive

amplification within chalcid wasp lineage, and which appears to have entered the ancestral

chalcid wasp by horizontal transfer [43]. Finally, a handful of transcripts have no identifiable

domains or homologs. The functional relevance of these molecules will be an area of intense

interest in the future.

Unexpected functions of novel oosome components

Early in our analysis we chose a handful of transcripts for functional analysis. These were

chosen based on a combination of criteria that included high enrichment in the oosome, novelty,

and the potential to give phenotypes that we could characterize with the current set of functional

tools available to us in *Nasonia*. Three of these (*Nv-bark*, *Nv-anillin* and *Nv-rrm*) gave an

unexpected phenotype, where oosome-like material was not coalesced into the typical spherical

oosome structure, but rather was scattered in clumps attached to the plasma membrane near the

posterior pole. Eventually, these clumps of germ plasm-like material come into contact with

syncytial nuclei when they migrate to the cortex. However, this remnant material is unable to

induce the pole cell fate. It is important to reiterate that this phenotype is quite distinct from that

seen for genes that have a core role in oosome assembly (*Nv-osk*, *Nv-tud*, *Nv-vas* and *Nv-aub*

(not shown). In these cases, there is no hint of the oosome, and mRNAs normally localized in

the oosome are distributed in homogenous caps at the posterior pole, rather than as discrete

clumps of material [35].

These phenotypes indicate these genes are involved in the coalescence of the oosome into

single entity within the central column of cytoplasm in the embryo, and/or maintenance of

oosome integrity. The molecular bases of such functions are not completely clear at the moment

for any of these three genes. *Nasonia* Bark is a putative transmembrane protein and would be

predicted to be targeted to the membrane. One potential hypothesis is that *Nasonia* Bark is

targeted to the membrane at the posterior pole of the embryo, releases and/or repels oosome

material from adhering to the plasma membrane, forcing the oosome to remain in the bulk

cytoplasm where it concentrated into a large sphere and moved around by strong cytoplasmic

flows that occur during the earliest cleavages. This model would also imply that interaction with

the cortex prevents oosome material from coalescing, leading to the scattered clumps we observe

in *Nv-bark* knockdowns. Mechanistically, this could be related to the ability of Bark to induce

endocytosis [90], a process that is associated with proper anchoring of the germ plasm and the

recruitment of specialized actin binding proteins to the posterior pole in *Drosophila* [91].

An alternative hypothesis is that *Nasonia* Bark produced in the oosome is not secreted but

is instead incorporated as an important structural component of the large, solid form of the

oosome. This is consistent with the structure of the protein, which contains main protein-protein

interaction domains of different types [45, 46]. Testing these hypotheses will require in depth

analysis of the subcellular localization of *Nasonia* Bark during early embryogenesis, and

proteomic analysis of binding partners of *Nasonia* Bark.

The role of *Nv-anillin* in maintaining the stability of the oosome was also surprising and

the molecular basis of the phenotype will require further investigation. Anillin orthologs are well

known as actin binding proteins involved in assembling the contractile ring required to separate

cells in cytokinesis [50]. Anillin also plays a crucial and novel role in the specialized cytokinesis

of the *Drosophila* pole cells [51]. While these known functions might have indicated that

oosome localization of *Nv-anillin* was related to an important role in the specialized polar bud

formed in *Nasonia*, RNAi showed that this protein has an earlier role in oosome

assembly/maintenance (Fig. 6 C1). Similar to *Nasonia* Bark, one possible function of *Nasonia*

Anillin is as a structural component of the oosome, which may or may not be related to its ability

to bind actin and associated proteins. Alternatively, *Nv-anillin* may act to release and/or repel the

oosome from the cortex, as proposed above for *Nv-bark*.

Anillin homologs have known functions which are directly related to the processes of release of

germ plasm from the embryonic plasma membrane. In *Drosophila*, the germ plasm is tightly

bound to the plasma membrane until the nuclei reach the posterior pole of the embryo. The

centrosomes associated with these nuclei mediate detachment of the pole plasm from the cortex

through interactions of the astral microtubules emanating from the centrosomes [92]. This is in

contrast to *Nasonia,* where the oosome detaches from the cortex at about the same time as the

zygotic nucleus begins its first division at the anterior pole. While nuclei are lacking at this time,

numerous centrosomes are present, as they are provided maternally in a process characteristic of

many Hymenopteran embryos [93]. *Nasonia* Anillin could be relevant to a model where astral

microtubules emanating from maternally provided centrosomes detach oosome material from the

cortex, because Anillin homologs have been shown to mediate interactions between the actin

cytoskeleton and cortical and subcortical microtubule arrays in multiple model systems [94, 95].

Again, in depth examination of the subcellular localization and interactions of *Nasonia* Anillin

will be required to completely understand its role in maintaining the oosome.

Based on the presence of only RNA recognition motifs in the protein, we predict that *Nv-*

rrm will have one of two likely roles. One possibility is that it is involved in translational

regulation of key regulators of oosome structure, presumably including *Nasonia* Anillin and

Nasonia Bark. Alternatively, *Nv-rrm* may be important in binding RNA and protein in order to

maintain the structural integrity of the oosome. In addition, neither of these possibilities is

mutually exclusive.

The knockdowns of *Nv-coronin* and *Nv-innexin1* had specific effects only on the

formation of pole cells, while the oosome appeared to remain intact. Coronin is an actin binding

protein associated with the formation of highly concentrated networks of F-actin [65]. It seems

likely that *Nasonia* Coronin has an important role in organizing an actin cytoskeleton

arrangement specialized for the formation of the large polar bud that initiates pole cell formation.

The potential role of *Nv-innexin1* is somewhat more mysterious. Innexins are typically

known as components of the gap junctions that are found in some tightly integrated epithelial

tissues. Such junctions would not be expected of the motile pole cells. Interestingly, Innexin-7 (a

paralogous protein with a similar structure to *Nasonia* Innexin1) in the beetle *Tribolium* has a

novel role in cellularization of the syncytial blastoderm. Such a function for *Nasonia* Innexin1

would explain the failure in pole cell formation we see after RNAi.

Conclusion

This work has revealed numerous unexpected mRNAs that are localized to the germ plasm of the

wasp *Nasonia*. The results have given insights on the potentially ancestral mechanisms used by

germ plasm to accomplish conserved required functions, such as a possible ancestral role for

Milton orthologs to bring mitochondria to the germ plasm, a function replaced by the long Osk

isoform in *Drosophila*. On the other hand, our results have identified numerous components that

are likely to be specific to *Nasonia* and its relatives in the parasitic wasp lineage (e.g., *Nv-*

OoCLANK, Nv-rrm, and the use of Nv-Bark and Nv-Anillin in assembling the oosome). Deeper

analyses of the functions of these molecules in *Nasonia*, and broader sampling of germ plasm in

other holometabolous insects will be required to determine the patterns of evolutionary change in

the germ plasm. Such analyses will be important because the germ plasm is a uniquely powerful

organelle that can rapidly drive naive nuclei into a highly specialized, yet functionally totipotent

state. Understanding how and why such a fundamental substance changes and is even lost in the

course of evolution will provide foundational insights into the mechanisms of cell fate

determination and the interaction of subcellular organelles and their cellular milieu.

Methods

Sample preparation

In order to identify the components of the maternally deposited mRNAs in the oosome located in the posterior half of the parasitoid wasp *Nasonia vitripennis* embryos, we collected and bisected the pre-blastoderm stage embryos (0-2 hours after egg lay at 25 °C) to detect the differential expression levels between the anterior and posterior halves of the embryos.

The embryos were aligned with the anterior pole to the right on the ice-prechilled and heptane glue-coated slide. Then a thin layer of halocarbon oil 700 (Sigma) was applied to cover the embryos. The slide was transferred on the dry ice-prechilled "guillotine" and then was anchored by tightening the screws on each end. After the halocarbon oil 700 was solidified on dry ice, put the guillotine on the dry ice-prechilled stabilizer and transferred it under the dissection microscope. The embryos were positioned to match the slot in the guillotine where the dry ice-prechilled razor blade will be inserted into. After the embryos were cut, the anterior and posterior halves of the embryos were immediately collected and transferred into the two 1.5mL non-stick RNase-Free microfuge tubes (Ambion) with the dry ice-prechilled probe, separately. Three biological replicates were created.

Total RNA was isolated from these six samples for library preparation. In the library preparation upon which this manuscript is based, we used around one microgram of total RNA from each sample, in which 100ng of total RNA was from the *Nasonia* other 900ug was from a distantly related parasitic wasp (*Melittobia digitata* [43]). Libraries were prepared using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (NEB #E7420) in conjunction with NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB #E7490). Libraries were validated and quantified before being pooled and sequenced on an Illumina HiSeq 2000

sequencer with a 100 bp paired-end protocol. Sequence files are available in the NCBI SRA

database under accession SRP156232.

RNA sequencing data analysis

The quality of the sequencing data was determined using FastQC software. The sequences were

processed by Cufflinks package for differential expression detection, using multiple variations on

the default parameters (job files in Supplemental File 2). Briefly, raw reads were aligned to

annotation   102   of   assembly   2.1   of   the   *N.   vitripennis*   genome

(https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Nasonia_vitripennis/102/)   using

TopHat2[96]. These results were either used directly in cuffdiff [97], or were further processed

using stringtie [98] to generate new transcriptome predictions and quantification. Various

normalization parameters were used, and each permutation gave slightly different results. In

addition, we mapped the reads using assembly 1.0 and annotation 2.0 (OGS 2.0) [49]. All of the

commands jobs in this process of analysis are included in the Supplemental File 2. The cuffdiff

results of all permutations of the analysis are provided in Supplemental File 3. The computing

work was done by the High Performance Computing Cluster located at University of Illinois at

Chicago.

Probes and dsRNAs for the chosen genes were generated by the protocol described in

[40]. Primers for generating these templates are provided in Supplemental File 4. Alkaline

phosphatase *in situ* hybridization was performed by the protocol described in [99].

Embryonic RNA interference (eRNAi)

In order to perform eRNAi on the early *Nasonia* embryos to study the germline candidate genes'

functions, we created the following workflow:

Around 30 pre-blastoderm stage embryos (0-1 hour after egg lay at 25 °C) were collected

and quickly aligned vertically on the heptane glue-coated 18mm×18mm coverslip. This coverslip

was transferred and anchored on the ice-prechilled slide by applying a thin layer of water. The

slide was then put in an air tight petri dish with proper amount of desiccant (Drierite with

indicator, 8 mesh, ACROS Organics) pre-chilled at 4 °C. The embryos were dehydrated in the

desiccant at 4 °C for 45 minutes. After dehydration, the embryos were covered with a layer of

halocarbon oil 700 and were ready for microinjection.

The dsRNAs were dissolved in Nuclease-Free Water (Ambion) at the concentration of

1mg/mL and loaded into the Femtotips II Microinjection Capillary (Cat. No. 930000043,

Eppendorf). The constant pressure was set at 500hpa and the injection pressure was set initially

at 250hpa with periodic adjustment as the needle changed over the course of injection. The

process of injection was performed at room temperature and needed to be done as soon as

possible. After injection, the slide was transferred into a paper towel-moisturized petri dish pre-

warmed at 28 °C to incubate the injected embryos for specific developmental stages. The

embryogenesis of these embryos was stopped at pre-blastoderm stage (before the budding),

beginning of blastoderm stage (during budding), and later in blastoderm stage (pole cells formed

and/or after pole cell divisions). To stop the development, the coverslip was put into the heptane

to wash off the halocarbon oil 700 for three minutes, and then transferred into the 37%

formaldehyde-saturated heptane for 2-5 hours fixation in the dark with the embryos facing up.

After fixation, the coverslip was carefully taken out of the fixative, and flipped upside

down to gently press the embryos on a double-sided tape that was taped on a petri dish, so that

all the embryos can be anchored on the tape for dissection. Add about 15mL PBS with 1%

Tween, use the needle (BD PrecisionGlide Needle, 30G × 1) to carefully remove the eggshells

from the embryos. The dissected embryos were then transferred by pipette into the 1.5mL non-

stick RNase-Free microfuge tubes. The embryos were immediately dehydrated by 100%

Methanol and stored at -20 °C.

Before performing fluorescent *in situ* hybridization (FISH) on those eRNAi knocked out

embryos, they need to be rehydrated with a series of Methanol/PBT washes (75%, 50%, 25%).

The protocol for FISH was adapted from [40]. A detailed protocol is available on request.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

Sequencing results can be found in the NCBI SRA database at:

https://www.ncbi.nlm.nih.gov/sra/SRP156232.

Competing Interests

The authors declare no competing interests.

Funding

This work was supported by startup funds from UIC, and NIH grants 1R03HD087476 and

1R03HD078578.

Authors' contributions

JAL and HHQ collaborated on experimental design, analyzed the data and wrote the paper.

HHQ performed the experiments and collected the data.

Acknowledgements

We thank Daniel Pers for performing the combined *Nasonia/ Melittobia* RNAseq libraries, and

the UIC RRC Sequencing core for arranging the sequencing of our samples.

Competing Interests:

The authors declare no competing interests.

References

1.   Extavour CG, Akam M: **Mechanisms of germ cell specification across the metazoans:**
epigenesis and preformation. *Development* 2003, **130**(24):5869-5884.
2.   Extavour CG: **Evolution of the bilaterian germ line: lineage origin and modulation of**
specification mechanisms. *Integr Comp Biol* 2007, **47**(5):770-785.
3.   Hegner RW: **Experiments with Chrysomelid Beetles. III. The Effects of Killing Parts of**
the Eggs of Leptinotarsa decemlineata. *Biological Bulletin* 1911, **20**:237-251.
4.   Illmensee K, and A. P. Mahowald: **Transplantation of posterior polar plasm in**
Drosophila. Induction of germ cells at the anterior pole of the egg. *Proceedings of the*
National Academy of Sciences 1974, **71**(4):1016-1020.
5.   Illmensee K, and A. P. Mahowald: **The autonomous function of germ plasm in a somatic**
region of the Drosophila egg. *Experimental cell research* 1976, **97**(1):127-140.
6.   Xia QY, Zhou, Z. Y., Lu, C., Cheng, D. J., Dai, F. Y., et al.: **A draft sequence for the genome**
of the domesticated silkworm (Bombyx mori). *Science* 2004, **306**:1937-1940.
7.   Nagy L, Riddiford, L., Kiguchi, K. : **Morphogenesis in the Early Embryo of the**
Lepidopteran Bombyx-Mori. *Developmental Biology* 1994, **165**:137-151.
8.   Juhn J, Marinotti, O., Calvo, E., James, A. A.: **Gene structure and expression of nanos**
(nos) and oskar (osk) orthologues of the vector mosquito, Culex quinquefasciatus.
Insect Molecular Biology 2008, **17**:545-552.
9.   Suzuki N, Shimizu, S., Ando, H. : **Early Embryology of the Alderfly, Sialis-Mitsuhashii**
Okamoto (Megaloptera, Sialidae). *International Journal of Insect*
Morphology & Embryology 1981, **10**:409-418.

10. Werren JH, Richards, S., Desjardins, C. A., Niehuis, O., Gadau, J.: **Functional and**
Evolutionary Insights from the Genomes of Three Parasitoid Nasonia Species. *Science,*
327:343-348.
11. Berg GJ, Gassner, G.: **Fine structure of the blastoderm embryo of the pink bollworm,**
Pectinophora Gossypiella (Saunders) (lepidoptera: Gelechiidae). *International Journal*
of Insect Morphology and Embryology 1978, **7**:81–105.
12. Miura T, Braendle, C., Shingleton, A., Sisk, G., Kambhampati, S., Stern, D. L.: **A**
comparison of parthenogenetic and sexual embryogenesis of the pea aphid
Acyrthosiphon pisum (Hemiptera : Aphidoidea). *Journal of Experimental Zoology Part*
B-Molecular and Developmental Evolution 2003, **295B**:59-81.
13. Bull AL: **Stages of Living Embryos in the Jewel Wasp Mormoniella-**
(Nasonia)-Vitripennis-(Walker)(Hymenoptera, Pteromalidae). *Journal of Insect Morphology &*
Embryology 1982, **11**:1-23.
14. Klag J: **Differentiation of primordial germ cells in the embryonic development of**
Thermobia domestica. Pack. (Thysanura): an ultrastructural study. *Journal of*
Embryology and Experimental Morphology 1977, **38**:93-114.
15. Chang C-c, Dearden P, Akam M: **Germ Line Development in the Grasshopper**
Schistocerca gregaria: vasa As a Marker. *Developmental Biology* 2002, **252**(1):100-118.
16. Consortium THGS: **Insights into social insects from the genome of the honeybee Apis**
mellifera. *Nature* 2006, **443**(7114):931-949.
17. Dearden PK: **Germ cell development in the Honeybee (Apis mellifera); vasa and nanos**
expression. *BMC Dev Biol* 2006, **6**:6.
18. Dearden PK, Wilson MJ, Sablan L, Osborne PW, Havler M, McNaughton E, Kimura K,
Milshina NV, Hasselmann M, Gempe T *et al*: **Patterns of conservation and change in**
honey bee developmental genes. *Genome Res* 2006, **16**(11):1376-1384.
19. Juhn J, James AA: **oskar gene expression in the vector mosquitoes, Anopheles gambiae**
and Aedes aegypti. *Insect Mol Biol* 2006, **15**(3):363-372.
20. Nakao H, Hatakeyama M, Lee JM, Shimoda M, Kanda T: **Expression pattern of Bombyx**
vasa-like (BmVLG) protein and its implications in germ cell development. *Dev Genes*
Evol 2006, **216**(2):94-99.
21. Schroder R: **vasa mRNA accumulates at the posterior pole during blastoderm**
formation in the flour beetle Tribolium castaneum. *Dev Genes Evol* 2006, **216**(5):277-
283.
22. Khila A, Abouheif E: **Reproductive constraint is a developmental mechanism that**
maintains social harmony in advanced ant societies. *Proc Natl Acad Sci U S A* 2008,
105(46):17884-17889.
23. Mito T, Nakamura T, Sarashina I, Chang CC, Ogawa S, Ohuchi H, Noji S: **Dynamic**
expression patterns of vasa during embryogenesis in the cricket Gryllus bimaculatus.
Dev Genes Evol 2008, **218**(7):381-387.
24. Tribolium Genome Sequencing C, Richards S, Gibbs RA, Weinstock GM, Brown SJ, Denell
R, Beeman RW, Gibbs R, Beeman RW, Brown SJ *et al*: **The genome of the model beetle**
and pest Tribolium castaneum. *Nature* 2008, **452**(7190):949-955.
25. Consortium TIAG: **Genome Sequence of the Pea Aphid Acyrthosiphon pisum.** *PLoS Biol*
2010.

26.   Khila A, Abouheif E: **Evaluating the role of reproductive constraints in ant social**
evolution. *Philos Trans R Soc Lond B Biol Sci* 2010, **365**(1540):617-630.
27.   Shigenobu S, Bickel RD, Brisson JA, Butts T, Chang CC, Christiaens O, Davis GK, Duncan EJ,
Ferrier DE, Iga M *et al*: **Comprehensive survey of developmental genes in the pea aphid,**
Acyrthosiphon pisum: frequent lineage-specific duplications and losses of
developmental genes. *Insect Mol Biol* 2010, **19 Suppl 2**:47-62.
28.   Bontems F, Stein A, Marlow F, Lyautey J, Gupta T, Mullins MC, Dosch R: **Bucky ball**
organizes germ plasm assembly in zebrafish. *Curr Biol* 2009, **19**(5):414-422.
29.   Lehmann R, Nussleinvolhard C: **Abdominal Segmentation, Pole Cell-Formation, and**
Embryonic Polarity Require the Localized Activity of Oskar, a Maternal Gene in
Drosophila. *Cell* 1986, **47**(1):141-152.
30.   Ephrussi ALKD, L. K.; Lehmann, R.: **oskar organizes the germ plasm and directs**
localization of the posterior determinant nanos. *Cell* 1991, **66**(1):14.
31.   Ephrussi A, Lehmann R: **Induction of Germ-Cell Formation by Oskar.** *Nature* 1992,
358(6385):387-392.
32.   Rongo CG, E. R.; Lehmann, R.: **Localization of oskar RNA regulates oskar translation and**
requires Oskar protein. *Development* 1995, **121**(9):2737-2746.
33.   Breitwieser W, Markussen, F. H., Horstmann, H., Ephrussi, A.: **Oskar protein interaction**
with Vasa represents an essential step in polar granule assembly. *Genes &*
Development 1996, **10**:2179-2188.
34.   Boswell RE, Mahowald AP: **Tudor, a Gene Required for Assembly of the Germ Plasm in**
Drosophila-Melanogaster. *Cell* 1985, **43**(1):97-104.
35.   Lynch JA, Ozuak O, Khila A, Abouheif E, Desplan C, Roth S: **The phylogenetic origin of**
oskar coincided with the origin of maternally provisioned germ plasm and pole cells at
the base of the Holometabola. *PLoS genetics* 2011, **7**(4):e1002029.
36.   Jaglarz MK, Howard, K. R.: **The active migration of Drosophila primordial germ cells.**
Development 1995, **121**:3495-3503.
37.   Wylie A, Jones AE, D'Brot A, Lu WJ, Kurtz P, Moran JV, Rakheja D, Chen KS, Hammer RE,
Comerford SA *et al*: **p53 genes function to restrain mobile elements.** *Genes Dev* 2016,
30(1):64-77.
38.   Tiwari B, Kurtz P, Jones AE, Wylie A, Amatruda JF, Boggupalli DP, Gonsalvez GB, Abrams
JM: **Retrotransposons Mimic Germ Plasm Determinants to Promote Transgenerational**
Inheritance. *Curr Biol* 2017, **27**(19):3010-3016 e3013.
39.   Bantock CR: **Experiments on chromosome elimination in the gall midge, Mayetiola**
destructor. *Journal of Embryology and Experimental Morphology* 1970, **24**:257-289.
40.   Lynch JA, Desplan C: **Novel modes of localization and function of nanos in the wasp**
Nasonia. *Development* 2010, **137**(22):3813-3821.
41.   Santos AC, Lehmann R: **Germ cell specification and migration in Drosophila and beyond.**
Curr Biol 2004, **14**(14):R578-589.
42.   Draper BWM, C. C.; Bowerman, B.; Hardin, J.; Priess, J. R. : **MEX-3 Is a KH Domain**
Protein That Regulates Blastomere Identity in Early C. elegans Embryos. *Cell* 1996,
87:205-216.

43.　Lynch J, Pers D: **Ankyrin domain encoding genes resulting from an ancient horizontal**
transfer are functionally integrated into developmental gene regulatory networks in
the wasp Nasonia. *bioRxiv* 2018.

44.　Lynch JA, Brent AE, Leaf DS, Pultz MA, Desplan C: **Localized maternal orthodenticle**
patterns anterior and posterior in the long germ wasp Nasonia. *Nature* 2006,
439(7077):728-732.

45.　Byri S, Misra T, Syed ZA, Batz T, Shah J, Boril L, Glashauser J, Aegerter-Wilmsen T, Matzat
T, Moussian B *et al*: **The Triple-Repeat Protein Anakonda Controls Epithelial Tricellular**
Junction Formation in Drosophila. *Dev Cell* 2015, **33**(5):535-548.

46.　Hildebrandt A, Pflanz R, Behr M, Tarp T, Riedel D, Schuh R: **Bark beetle controls**
epithelial morphogenesis by septate junction maturation in Drosophila. *Dev Biol* 2015,
400(2):237-247.

47.　Jennings BH, Shah S, Yamaguchi Y, Seki M, Phillips RG, Handa H, Ish-Horowicz D: **Locus-**
specific requirements for Spt5 in transcriptional activation and repression in
Drosophila. *Curr Biol* 2004, **14**(18):1680-1684.

48.　Pers D, Buchta T, Ozuak O, Wolff S, Pietsch JM, Memon MB, Roth S, Lynch JA: **Global**
analysis of dorsoventral patterning in the wasp Nasonia reveals extensive
incorporation of novelty in a regulatory network. *BMC Biol* 2016, **14**:63.

49.　Rago A, Gilbert DG, Choi JH, Sackton TB, Wang X, Kelkar YD, Werren JH, Colbourne JK:
OGS2: genome re-annotation of the jewel wasp Nasonia vitripennis. *BMC Genomics*
2016, **17**:678.

50.　Field CM, Alberts BM: **Anillin, a Contractile Ring Protein That Cycles from the Nucleus**
to the Cell Cortex. *J Cell Biol* 1995, **131**(1):165-178.

51.　Cinalli RM, Lehmann R: **A spindle-independent cleavage pathway controls germ cell**
formation in Drosophila. *Nature cell biology* 2013, **15**(7):839-845.

52.　Anand A, Kai T: **The tudor domain protein kumo is required to assemble the nuage and**
to generate germline piRNAs in Drosophila. *EMBO J* 2012, **31**(4):870-882.

53.　Zhang Z, Koppetsch BS, Wang J, Tipping C, Weng Z, Theurkauf WE, Zamore PD:
Antisense piRNA amplification, but not piRNA production or nuage assembly, requires
the Tudor-domain protein Qin. *EMBO J* 2014, **33**(6):536-539.

54.　Anand A, Kai T: **Response to Zhang et al.** *EMBO J* 2014, **33**(6):540-541.

55.　Lecuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, Cerovina T, Hughes TR,
Tomancak P, Krause HM: **Global analysis of mRNA localization reveals a prominent role**
in organizing cellular architecture and function. *Cell* 2007, **131**(1):174-187.

56.　Pan J, Goodheart M, Chuma S, Nakatsuji N, Page DC, Wang PJ: **RNF17, a component of**
the mammalian germ cell nuage, is essential for spermiogenesis. *Development* 2005,
132(18):4029-4039.

57.　Watanabe T, Kankel, D. R.: **Molecular Cloning and Analysis olf(1)ogre, a Locus of**
Drosophila melanogaster With Prominent Effects on the Postembryonic Development
of the Central Nervous System. *Genetics* 1990, **126**:1033-1044.

58.　van der Zee M, Benton MA, Vazquez-Faci T, Lamers GE, Jacobs CG, Rabouille C:
Innexin7a forms junctions that stabilize the basal membrane during cellularization of
the blastoderm in Tribolium castaneum. *Development* 2015, **142**(12):2173-2183.

59.   Olesnicky EC, Brent AE, Tonnes L, Walker M, Pultz MA, Leaf D, Desplan C: **A caudal**
mRNA gradient controls posterior development in the wasp Nasonia. *Development*
2006, **133**(20):3973-3982.

60.   Ozuak O, Buchta T, Roth S, Lynch JA: **Dorsoventral polarity of the Nasonia embryo**
primarily relies on a BMP gradient formed without input from Toll. *Curr Biol* 2014,
24(20):2393-2398.

61.   Riesgo-Escovar JR, Hafen E: **Common and distinct roles of DFos and DJun during**
Drosophila development. *Science* 1997, **278**(5338):669-672.

62.   Dorfman R, Glazer L, Weihe U, Wernet MF, Shilo BZ: **Elbow and Noc define a family of**
zinc finger proteins controlling morphogenesis of specific tracheal branches.
Development 2002, **129**(15):3585-3596.

63.   Wernet MF, Meier KM, Baumann-Klausener F, Dorfman R, Weihe U, Labhart T, Desplan
C: **Genetic dissection of photoreceptor subtype specification by the Drosophila**
melanogaster zinc finger proteins elbow and no ocelli. *PLoS genetics* 2014,
10(3):e1004210.

64.   Rybakin V, Clemen CS: **Coronin proteins as multifunctional regulators of the**
cytoskeleton and membrane trafficking. *Bioessays* 2005, **27**(6):625-632.

65.   Gandhi M, Goode BL: **Coronin: the double-edged sword of actin dynamics.** *Subcell*
Biochem 2008, **48**:72-87.

66.   Glater EE, Megeath LJ, Stowers RS, Schwarz TL: **Axonal transport of mitochondria**
requires milton to recruit kinesin heavy chain and is light chain independent. *J Cell Biol*
2006, **173**(4):545-557.

67.   Tanaka T, Nakamura A: **The endocytic pathway acts downstream of Oskar in**
Drosophila germ plasm assembly. *Development* 2008, **135**(6):1107-1117.

68.   Jaspers MH, Pflanz R, Riedel D, Kawelke S, Feussner I, Schuh R: **The fatty acyl-CoA**
reductase Waterproof mediates airway clearance in Drosophila. *Dev Biol* 2014,
385(1):23-31.

69.   Buchta T, Ozuak O, Stappert D, Roth S, Lynch JA: **Patterning the dorsal-ventral axis of**
the wasp Nasonia vitripennis. *Dev Biol* 2013, **381**(1):189-202.

70.   Ozuak O, Buchta T, Roth S, Lynch JA: **Ancient and diverged TGF-beta signaling**
components in Nasonia vitripennis. *Dev Genes Evol* 2014, **224**(4-6):223-233.

71.   Tomancak P, Berman BP, Beaton A, Weiszmann R, Kwan E, Hartenstein V, Celniker SE,
Rubin GM: **Global analysis of patterns of gene expression during Drosophila**
embryogenesis. *Genome Biol* 2007, **8**(7):R145.

72.   Lee DM, Wilk R, Hu J, Krause HM, Harris TJ: **Germ Cell Segregation from the Drosophila**
Soma Is Controlled by an Inhibitory Threshold Set by the Arf-GEF Steppke. *Genetics*
2015, **200**(3):863-872.

73.   Martinson EO, Mrinalini, Kelkar YD, Chang CH, Werren JH: **The Evolution of Venom by**
Co-option of Single-Copy Genes. *Curr Biol* 2017, **27**(13):2007-2013 e2008.

74.   Hanyu-Nakamura K, Sonobe-Nojima H, Tanigawa A, Lasko P, Nakamura A: **Drosophila**
Pgc protein inhibits P-TEFb recruitment to chromatin in primordial germ cells. *Nature*
2008, **451**(7179):730-733.

75.   Nakamura A, Amikura R, Mukai M, Kobayashi S, Lasko PF: **Requirement for a noncoding**
RNA in Drosophila polar granules for germ cell establishment. *Science* 1996,
274(5295):2075-2079.

76.   Zhang F, Barboric M, Blackwell TK, Peterlin BM: **A model of repression: CTD analogs and**
PIE-1 inhibit transcriptional elongation by P-TEFb. *Genes Dev* 2003, **17**(6):748-758.

77.   Yamada T, Yamaguchi Y, Inukai N, Okamoto S, Mura T, Handa H: **P-TEFb-mediated**
phosphorylation of hSpt5 C-terminal repeats is critical for processive transcription
elongation. *Mol Cell* 2006, **21**(2):227-237.

78.   Lerit DA, Shebelut CW, Lawlor KJ, Rusan NM, Gavis ER, Schedl P, Deshpande G: **Germ**
Cell-less Promotes Centrosome Segregation to Induce Germ Cell Formation. *Cell Rep*
2017, **18**(4):831-839.

79.   Amarnath S, Stevens LM, Stein DS: **Reconstitution of Torso signaling in cultured cells**
suggests a role for both Trunk and Torso-like in receptor activation. *Development* 2017,
144(4):677-686.

80.   Guraya SS: **Recent advances in the morphology, cytochemistry, and function of**
Balbiani's vitelline body in animal oocytes. *Int Rev Cytol* 1979, **59**:249-321.

81.   Kloc M, Bilinski S, Etkin LD: **The Balbiani body and germ cell determinants: 150 years**
later. *Curr Top Dev Biol* 2004, **59**:1-36.

82.   Hurd TR, Herrmann B, Sauerwald J, Sanny J, Grosch M, Lehmann R: **Long Oskar Controls**
Mitochondrial Inheritance in Drosophila melanogaster. *Dev Cell* 2016, **39**(5):560-571.

83.   Cox RT, Spradling AC: **Milton controls the early acquisition of mitochondria by**
Drosophila oocytes. *Development* 2006, **133**(17):3371-3377.

84.   Patil VS, Anand A, Chakrabarti A, Kai T: **The Tudor domain protein Tapas, a homolog of**
the vertebrate Tdrd7, functions in the piRNA pathway to regulate retrotransposons in
germline of Drosophila melanogaster. *BMC Biol* 2014, **12**:61.

85.   Gao M, Thomson TC, Creed TM, Tu S, Loganathan SN, Jackson CA, McCluskey P, Lin Y,
Collier SE, Weng Z *et al*: **Glycolytic enzymes localize to ribonucleoprotein granules in**
Drosophila germ cells, bind Tudor and protect from transposable
elements. *EMBO reports* 2015, **16**(3):379-386.

86.   Teng H, Sui X, Zhou C, Shen C, Yang Y, Zhang P, Guo X, Huo R: **Fatty acid degradation**
plays an essential role in proliferation of mouse female primordial germ cells via the
p53-dependent cell cycle regulation. *Cell Cycle* 2016, **15**(3):425-431.

87.   Pao SS, Paulsen IT, Saier MH: **Major facilitator superfamily**. *Microbiol Mol Biol R* 1998,
62(1):1-+.

88.   Yamada Y, Davis KD, Coffman CR: **Programmed cell death of primordial germ cells in**
Drosophila is regulated by p53 and the Outsiders monocarboxylate
transporter. *Development* 2008, **135**(2):207-216.

89.   Jaspers MHJ, Pflanz R, Riedel D, Kawelke S, Feussner I, Schuh R: **The fatty acyl-CoA**
reductase Waterproof mediates airway clearance in Drosophila. *Developmental*
Biology 2014, **385**(1):23-31.

90.   Hildebrandt A, Pflanz R, Behr M, Tarp T, Riedel D, Schuh R: **Bark beetle controls**
epithelial morphogenesis by septate junction maturation in Drosophila. *Dev Biol* 2015,
400(2):237-247.

91.   Tanaka T, Kato Y, Matsuda K, Hanyu-Nakamura K, Nakamura A: **Drosophila Mon2**
1017   **couples Oskar-induced endocytosis with actin remodeling for cortical anchorage of the**
1018   **germ plasm**. *Development* 2011, **138**(12):2523-2532.
1019   92.   Lerit DA, Gavis ER: **Transport of germ plasm on astral microtubules directs germ cell**
1020   **development in Drosophila**. *Curr Biol* 2011, **21**(6):439-448.
1021   93.   Ferree PM, McDonald K, Fasulo B, Sullivan W: **The origin of centrosomes in**
1022   **parthenogenetic hymenopteran insects**. *Curr Biol* 2006, **16**(8):801-807.
1023   94.   van Oostende Triplet C, Jaramillo Garcia M, Haji Bik H, Beaudet D, Piekny A: **Anillin**
1024   **interacts with microtubules and is part of the astral pathway that defines cortical**
1025   **domains**. *J Cell Sci* 2014, **127**(Pt 17):3699-3710.
1026   95.   Tse YC, Piekny A, Glotzer M: **Anillin promotes astral microtubule-directed cortical**
1027   **myosin polarization**. *Mol Biol Cell* 2011, **22**(17):3165-3175.
1028   96.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL: **TopHat2: accurate**
1029   **alignment of transcriptomes in the presence of insertions, deletions and gene fusions**.
1030   *Genome Biol* 2013, **14**(4):R36.
1031   97.   Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL,
1032   Pachter L: **Differential gene and transcript expression analysis of RNA-seq experiments**
1033   **with TopHat and Cufflinks**. *Nat Protoc* 2012, **7**(3):562-578.
1034   98.   Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL: **StringTie**
1035   **enables improved reconstruction of a transcriptome from RNA-seq reads**. *Nat*
1036   *Biotechnol* 2015, **33**(3):290-295.
1037   99.   Brent AE, Schweitzer R, Tabin CJ: **A somitic compartment of tendon progenitors**. *Cell*
1038   2003, **113**(2):235-248.
1039

1040

Figure Legends

Fig. 1. The behavior and fate of the oosome. Green marks the germ plasm (Nv-bark in situ hybridization), white (DAPI) marks the nuclei. A-D: the green shows the oosome shapes and localizations in different nuclear divisions during pre-blastoderm stage. E: the oosome flattens into the bud (green) as the nuclei reach the embryonic cortex. F: several pole cells immediately form from the bud during early syncytial blastoderm stage. G: the pole cells divide several times to make more and smaller cells by the time of cellularization. H: the pole cells are internalized during gastrulation and begin to migrate to the where the gonads will form. Anterior is at left, and dorsal is up in all figures.

1050

1051   **Fig. 2.** Genes expressed anteriorly in the pre-blastoderm stage. All embryos are aligned and

1052   grouped into three columns (Pre-blastoderm, Blastoderm and Gastrulation) according to their

1053   embryogenesis stages, with posterior side to the right and dorsal side on the top.

1054

1055   **Fig. 3.** Transcripts localized to the oosome that are subsequently maintained in pole cells. All

1056   embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and

1057   Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal

1058   side on the top.

1059

1060   **Fig. 4.** mRNAs localized in the oosome (Pre-blastoderm) but are excluded from the pole cells.

1061   All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and

1062   Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal

1063   side on the top.

1064

1065   **Fig. 5.** mRNAs strongly enriched in the posterior region of the embryos, but not enriched in the

1066   oosome. All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm

1067   and Gastrulation) according to their embryogenesis stages, with posterior side to the right and

1068   dorsal side on the top.

1069

Fig. 6. Functional analysis of a subset of germ plasm localized factors by eRNAi. A1-A4: eGFP

dsRNA injected embryos of increasing as the negative control. B1-B4: Nv-bark dsRNA injected

embryos, C1-C4: Nv-anillin dsRNA injected embryos, D1-D4: Nv-rrm dsRNA injected embryos,

E1-E4: Nv-coronin dsRNA injected embryos, F1-F4: Nv-innexin1 dsRNA injected embryos. All

the embryos are positioned with posterior side to the right and dorsal side on the top. The red in

B1-B4 are marked by Nv-nos, and the rest are marked by Nv-bark. DAPI (white) marks the

nuclei.

1077

Additional Files

Additional File 1. (.xlsx) Significantly enriched transcripts from both experiments. The

transcripts listed in "Compilation of Transcripts significant in the main analyses" were generated

using the sequencing data from the second experiment by NCBI annotation 102 with various

normalization parameters. The "Transcripts from the second experiment only found in OGS 2.0"

were the transcripts from the second experiment only found using the annotation OGS 2.0, but

not found using NCBI annotation 102. The "Transcripts from the first experiment" were the

transcripts only found in the first time sequencing data, but not in the second time sequencing

data.

Additional File 2. (.docx) Command jobs. The command jobs for analyses using the NCBI

annotation 102 and the annotation OGS 2.0 with various normalization parameters.

Additional File 3. (.xlsx) Original lists of the transcripts generated by Cuffdiff. This file

includes the original lists of the transcripts generated by Cuffdiff using the NCBI annotation 102

and the annotation OGS 2.0 with various normalization parameters.

Additional File 4. (.xlsx) Primer list. All primers used in this manuscript are listed. This list

includes the primers used for cloning the genes and making templates for the probes and

dsRNAs.

1095

1096

1097

1098

1099

1100

1101

	Pre-blastoderm	Blastoderm	Gastrulation
LOC103 315431	A1	A2	A3
LOC100 124035	B1	B2	B3
LOC1001 21321	C1	C2	C3
Nv-exdl2	D1	D2	D3
LOC100 680515	E1	E2	E3
Naswi2EG0 06979	F1	F2	F3
LOC1001 19882	G1	G2	G3
LOC1079 82211	H1	H2	H3
LOC100 313502	I1	I2	I3
LOC100 120674	J1	J2	J3
LOC100 115729	K1	K2	K3
Nv-mex3	L1	L2	L3
Nv-ucth	M1	M2	M3