1

# Transcriptomic and functional analysis of the oosome, a unique form of germplasm in the wasp *Nasonia vitripennis*

Honghu Quan and Jeremy Lynch

University of Illinois at Chicago, Chicago IL 60607 USA

## Abstract

The oosome is the germline determinant in the wasp *Nasonia vitripennis* and is homologous to the polar granules of *Drosophila*. Despite a common evolutionary origin and developmental role, the oosome is morphologically quite distinct from polar granules. It is a solid sphere that migrates within the cytoplasm before budding out and forming pole cells.  To gain an understanding of both the molecular basis of the novel form of the oosome, and the conserved essential features of germ plasm, we quantified and compared transcript levels between embryo fragments that contained the oosome, and those that did not. The identity of the localized transcripts indicated that *Nasonia* uses different molecules to carry out conserved germ plasm functions. In addition, functional testing of a sample of localized transcripts revealed potentially novel mechanisms of ribonucleoprotein assembly and cellularization in the wasp.

## Introduction

Germline establishment is a crucial event for sexually reproducing organisms. Germline cells are special in that they are able to generate all of the cell fates of the soma and to regenerate themselves. There are two major strategies to specify the germline among animals: zygotic induction and maternal provision. In zygotic induction, inductive signals from surrounding tissues drive the establishment of germline fate, usually relatively late in embryogenesis, after the transition from maternal to zygotic control (1). In contrast, in the maternal provision mode, the germ cells are specified by determinants called germ plasm that are synthesized and localized during oogenesis and are the first group cells formed during embryogenesis. Germline specification in this mode occurs very early in development, usually prior to the activation of the zygotic genome (1, 2). Classical experiments have shown that germ plasm is both necessary and sufficient to establish the germline fate (3-5).

It is likely that the maternal provision mode of germplasm evolved multiple times among the animals, and this reflected in the molecular basis of germplasm determinants (6-27). In vertebrates, germplasm (where it exists) is dependent on the maternal localization of *bucky ball* (28). In insects, the gene products of *oskar* are both necessary and sufficient to induce germ plasm, and thus, primordial germ cells (PGCs) (29-32). Downstream of these nucleators is a suite

of highly conserved germline-associated molecules (i.e., Vasa, Nanos, Tudor, etc.) that are

recruited to the germplasm where they presumably carry out functions ancestral for the animal

germline (30, 33-35).

There are several conserved properties of PGCs that must be conferred to naive cells by

germplasm.  One of these is a period of transcriptional quiescence germ cells undergo after being

specified. This feature makes the composition of a germplasm even more important since it

means that mRNAs and proteins critical for at least the early germ cell functions need to be

provided in the germplasm itself. Germ cells also usually become highly migratory later in

development as they seek to colonize the developing gonad. They are often highly enriched for

mitochondria and have specific metabolic needs (36). Since they carry the genome that will be

passed to future generations, germ cells have enhanced mechanisms to prevent DNA damage and

to reduce the activity of transposable elements (37, 38). Finally, they must have the capability to

induce pluripotency to their genome, as germline cells will be the source for all cell fates in the

eventual progeny.

Like all developmental processes, there is likely to be variation in the details of these conserved

features of germline determination, whether due to selective or neutral forces.  Pressures that

could impact the composition of the germplasm could be differential activity of transposable

elements in the genome, or a novel path for migration of the PGCs to the gonad, for example.

Novelties in embryogenesis may also drive germ plasm composition.  For example, in the

parthogenetic paedomorphic embryos of the midge *Miastor,* somatic nuclei undergo significant

chromosomal diminution in the early cleavages while the cell that takes up germplasm maintains

the full complement of chromosomes, indicating that the germplasm contains a component that

prevents the loss of chromosomes during the early cleavages (39).

Holometabolous insects are an ideal system with which to study how germplasm evolves, given

that it is ancestral in this clade (35), the unparalleled levels of diversity, and the strong baseline

understanding germplasm function derived from *Drosophila melanogaster*. Here we focus on the

parasitic wasp *Nasonia vitripennis* as a model to compare to the fruit fly. Like *Drosophila*,

*Nasonia* depends on Oskar, Vasa, and Tudor to assemble germ plasm (35, 40). However, in

contrast to the collection of small granules stably associated with the posterior pole that make up

the *Drosophila* germplasm, the *Nasonia* germplasm forms a very large, dense organelle called

the oosome (Figure 1). This highly divergent morphology strongly indicates that the composition

of the *Nasonia* oosome may diverge significantly from the polar granules of *Drosophila*.

**Fig. 1. The behavior and fate of the oosome.** Green marks the germplasm (*Nv-bark in situ* hybridization), white (DAPI) marks the nuclei.
A-D: the green shows the oosome shapes and localizations in different nuclear divisions during pre-blastoderm stage. E: the oosome
     flattens into the bud (green) as the nuclei reach the embryonic cortex. F: several pole cells immediately form from the bud during early
     syncytial blastoderm stage. G: the pole cells divide several times to make more and smaller cells by the time of cellularization. H: the pole
85   cells are internalized during gastrulation and begin to migrate to the where the gonads will form. Anterior is at left, and dorsal is up in all
     figures.

The behaviors of the oosome and the PGCs in *Nasonia* further imply a divergent composition of

the oosome. In freshly laid eggs, the oosome is tightly bound to the ventral-posterior cortex of

the embryo (Fig. 1 A). When the zygotic nucleus forms and moves into the interior of the

embryo, the oosome detaches from the cortex and coalesces into a dense, extremely large

structure in the same central column as the nuclei. It migrates anteriorly, sometimes being found

near to 50% egg length, before migrating back to the posterior pole (Fig. 1 B, C). As the

cleavage nuclei migrate toward the cortex, the oosome flattens into a crescent on the posterior

pole of the embryo while a large bud protrudes from the pole (Fig. 1 D). Typically, two or three

nuclei become associated with the bud and the oosome material. The bud pinches off, and the

nuclei rapidly individuate into pole cells (Fig. 1 E).  This is distinct from pole cell formation in

*Drosophila,* where each pole cell forms individually upon association with a certain critical

amount of pole plasm (41). After dividing a few times, the pole cells remain stable at the

posterior pole of the egg until gastrulation, when they migrate through the posterior epithelium

and coalesce into to two masses, presumably where the primordial gonads are developing (Fig. 1

F, G). This migration is distinct from *Drosophila* pole cells which remain external on the

posterior pole as the germ band undergoes massive extension well after gastrulation and migrate

internally through the gut wall (41).

From germ plasm assembly to the germ plasm behavior to PGC formation to PGC migration,

*Nasonia* and *Drosophila* share some fundamental aspects of germline establishment, but they

also have their own diverged features that fit in their own embryogenesis program. This raises

the question of which genes are the core components for the maternal provision mode and which

genes contribute to their own distinct features in germline development. To address these

questions, we compared the mRNA content of anterior and posterior halves of the pre-

blastoderm stage *Nasonia* embryos in an effort to identify the components specifically localized

to the oosome. We found only a few of mRNAs conserved in both fly polar granules and the

*Nasonia* oosome, such as *oskar (osk)*, *nanos (nos)*, and *ovo* (see Supplemental File 1). The rest

are all novel components germ plasm that either lack *Drosophila* homologs or have homologs in

*Drosophila* that do not play any roles in germline development. We performed functional studies

for a set of localized transcripts, all of which showed roles in the unique features of *Nasonia*

germline development, demonstrating the value of our approach to identify the molecular

sources of novelty among various insect lineages.

## Results

**Results**

**RNA-seq analyses of the anterior and posterior poles of the wasp *Nasonia* early embryos**

To identify the maternal transcripts in the oosome, we isolated the total RNA separately from

anterior and posterior poles of the pre-blastoderm stage *Nasonia vitripennis* embryos. Six

samples that comprised three sets of each pole were sequenced and subjected to differential

expression analysis, which are described in more detail in the Materials and Methods.

Our first attempt yielded encouraging, but mixed results. Several novel posteriorly enriched

factors were identified, and we did recover some known posterior factors (*Nv-osk* and *Nv-nos*).

However, other known localized factors such as *Nv-cad* and *Nv-dpp* were not found to be

statistically significant. In addition, one of the posterior samples was of poor quality, and could

not be used in the differential expression analyses, reducing the power of our approach. Since

we wanted results as comprehensive as possible, we repeated the experiment using a few

adjustments (see Materials and Methods) and obtained much more robust results. While this
second experiment was being prepared, we moved forward with expression and functional
analysis of novel candidates identified in the first. Since a handful of transcripts with confirmed
oosome localization were found only in the first experiment, the remainder of the results will
include these transcripts along with those identified in the second experiment.

All known maternally localized molecules (*Nv-osk*, *nos*, *dpp*, *cad*, *otd1*, and *gt*) were found with
high significance, giving confidence that our experimental design and analysis approaches were
appropriate for our goal of discovering the mRNAs localized to the oosome. Overall, we found
92 transcripts with apparent significant enrichment at the posterior pole. These ranged in levels
of enrichment from 1.4 to 55 times higher in the posterior fragments compared to anterior
fragments. Our analyses also uncovered anteriorly enriched mRNAs, of which there were 89,
with a range of fold enrichment from 1.4 to 10 times higher at the anterior.

**Novel transcripts localized in the anterior half of the *Nasonia* early embryos**
While the anterior factors are not the focus of this manuscript, some interesting observations
were made in examining a small sample of candidate mRNAs. Most transcripts are localized in
small domains at the anterior cortex and seem to extend toward the posterior in variable tendrils,
rather than being uniform or graded caps of anterior localization (Fig. 2 A1-K1). A notable
exception is the transcript of *Nasonia* homolog of *mex-3,* an RNA binding protein known for
controlling translation of orthologs of the posterior patterning factor *caudal* in the nematode *C.*
*elegans* and the beetle *Tribolium* (24, 42). *Nv-mex3* mRNA is localized in a broad domain
extending far toward the posterior of the embryo (Fig. 2 L1). *Nv-mex3* expression is highly
dynamic and variable in both the blastoderm and post-gastrular stages (Fig. 2 L2-L3, additional
images not shown).
Transient localization is the most common feature of the anteriorly localized transcripts.  Most
are ubiquitous or absent by the time the early syncytial blastoderm forms (Fig. 2 A2-H2, J1-K3,
M1-M3), except for *LOC100313502* which persists into the blastoderm stage where it forms a
anterior cap (Fig. 2 I1-I3). Among the anteriorly enriched transcripts, many have predicted
functions that may be relevant to egg activation (e.g. four encoding ion channels), and anterior-
posterior polarity and patterning (e.g. transcription and translation factors, see Supplemental file

1). There are also a large number of transcripts with no clear homologs outside of *Nasonia* see
(Supplemental File 1). One of these (LOC100119982, Fig. 2 G1-G3) is a member of a family
ankyrin domain containing molecules  that are specific to Chalcid wasps and appear to have
obtained a broad diversity of expression and potential function during *Nasonia* development
(manuscript in review).
170
Finally, a handful of molecules localized at both poles were detected.  These include the known
factors  *Nv-otd1*  (43) and *Nv-TollC* (JAL in preparation) *,* along with  *Nv-endoglucanase* (Fig. 3
E)*, Nv-insulin-like growth factor (Nv-igf,* Fig. 3 C)*, Nv-ucth* (a ubiquitin carboxy terminus
hydrolase, Fig. 2 M)) .  The fact that these were found despite lower apparent fold differences
between the two embryonic halves, further gave confidence that our analysis was robust enough
to detect even subtle germ plasm localization of the vast majority of mRNAs.
177
**General description of the novel transcripts localized in the posterior pole of the *Nasonia***
**early embryos**
From the two analyses, we identified 92 candidate transcripts that were statistically enriched at
the posterior half of the *Nasonia* embryos. We then isolated PCR products for 54 of these genes
and made probes to determine their expression patterns during early embryogenesis. We were
able to confirm 47 transcripts that are expressed posteriorly during pre-blastoderm stage. Of the
remaining transcripts, one of them was not successfully cloned, and the rest were successfully
cloned but did not have any specific, localized expression. Interpretation of these potential false
positives (as well as some false negatives) will be discussed later.  Among the 28 transcripts with
expression in the posterior half of the embryo, we grouped them into three categories based on
their expression patterns: A) genes expressed in oosome and with expression maintained in the
PGCs (Fig. 3), B) mRNAs localized in the oosome but then degraded in the PGCs (Fig. 4), C)
mRNAs strongly enriched in the posterior region of the embryo, but not incorporated in the
oosome (Fig. 5).
192
193

| | Pre-blastoderm | Blastoderm | Gastrulation |
|---|---|---|---|
| *LOC103 315431* | A1 | A2 | A3 |
| *LOC100 124035* | B1 | B2 | B3 |
| *LOC100 121321* | C1 | C2 | C3 |
| *Nv-exd12* | D1 | D2 | D3 |
| *LOC100 680515* | E1 | E2 | E3 |
| *Nasvi2EG 006979* | F1 | F2 | F3 |
| *LOC100 119882* | G1 | G2 | G3 |
| *LOC107 982211* | H1 | H2 | H3 |
| *LOC100 313502* | I1 | I2 | I3 |
| *LOC100 120674* | J1 | J2 | J3 |
| *LOC100 115729* | K1 | K2 | K3 |
| *Nv-mex3* | L1 | L2 | L3 |
| *Nv-ucth* | M1 | M2 | M3 |

**Fig. 2. Genes expressed anteriorly in the pre-blastoderm stage.** All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal side on the top.

**Highly germline-associated transcripts**

We identified 11 transcripts that are localized to the oosome and maintained in the PGCs at pole cell formation and beyond. We consider these to be the strongest candidates for having important roles in the specification and function of the PGCs but cannot exclude *a priori* that they have other (or no) important functions. Six transcripts are maintained throughout the stages of germline development followed in this manuscript: oosome, pole cells, and migrating germ cells. They include *Nasonia* homologs of the fly genes *bark beetle* (*Nv-bark*) (Fig. 1), *spt5* (*Nv-spt5*), *tejas* (*Nv-tdrd7*), *insulin-like growth factor* (*Nv-igf*), and two transcripts without fly homologs (*Nasvi2EG001470* and *Nv-endoglucanase*) (Fig. 3 A1-E3).

Among these, *Nv-bark* is the best germline marker, bearing strong and consistent germline association throughout early development, including expression in the late embryonic gonads (Fig. 1). In this respect, it is better than our previously favored marker, *Nv-nos*, which is downregulated significantly toward the end of PGC migration (40). Bark is a large transmembrane protein and is not expressed in the *Drosophila* germline. Its only known role is in stabilizing tricellular junction in epithelial cells during embryogenesis (44, 45). It is not clear how this function is relevant to germline function, and indicates a novel recruitment of this factor in the wasp.

*Nv-spt5* is significantly enriched in the oosome with low levels of ubiquitous expression in the rest of the embryos in pre-blastoderm stage (Fig. 3 A1). In the blastoderm and migrating germ cell stages expression is reduced but still enriched in the pole cells while the ubiquitous expression in the embryo persists (Fig. 3 A2-A3). Spt5 homologs are involved in regulating RNA polymerase progression during transcription (46), which might indicate that *Nv-Spt5* involves repressing the transcription in the germ cells.

*Nv-tdrd7* is present at appreciable levels throughout the bulk cytoplasm and is also strongly localized in the oosome (Fig. 3 B1). This pattern is well reflected in the quantification of mRNA

levels in the two halves of the embryo, which show significant numbers of reads coming from the anterior half of the embryo. At the blastoderm stage, *Nv-tdrd7* is moderately enriched in the pole cells and is zygotically expressed in a ventral-posterior patch (Fig. 3 B2), which was detected in our earlier analysis of dorsal-ventral patterning (47). After gastrulation, *Nv-tdrd7* is strongly upregulated in a group of cells that are near to where the germ cells migrated, but it is not clear if they are germ cells (Fig. 3 B3).

*Nv-igf* is initially expressed in a bipolar pattern (similar to *Nv-otd1*(43)), before becoming specific to the pole cells during the blastoderm stage and the migrating germ cells after gastrulation (Fig. 3 C1-C3).

*Nasvi2EG001470* encodes a short peptide of 80 amino acids and was not included in the most recent annotation of the *Nasonia* genome at NCBI, but was present in OGS 2.0 (48). A very similar sequence is annotated in the close relative *Trichomalopsis* (JAL personal observation), indicating that it is a *bona fide* transcript that is either novel, or very rapidly evolving. *Nasvi2EG001470* is strongly expressed in the oosome and pole cells, while levels markedly decrease in migrating germ cells (Fig. 3 D1-D3).

Besides the expression in the oosome and the pole cells, *Nv-endoglucanase* is initially expressed at both poles during pre-blastoderm stage and early blastoderm stage (Fig. 3 E1-E2). Later in blastoderm stage, the expression is down-regulated at the anterior pole and become specific to the pole cells (Fig. 3 E3). Proteins of this type are found extensively in Hemimetabola, Hymenoptera, and Coleoptera, and appear to have been lost at the origin of Lepidoptera and Diptera clades. It is intriguing that what roles this protein plays during *Nasonia* embryogenesis.

Five transcripts localized to the oosome enter and are maintained in the pole cells but are then downregulated in the migrating germ cells (Fig. 3 F1-J3). This set includes homologs of *Drosophila anillin* (*scraps*), *qin, and innexin1 (ogre)*, (*Nv-anillin*, *Nv-qin*, and *Nv-inx1*, respectively). Anillin is an actin binding protein that localizes to the contractile ring during cytokinesis (49). In *Drosophila* Anillin protein is localized in the cleavage furrows when forming the PGCs (50), but its mRNA is ubiquitous. The early oosome localization of *Nv-anillin* mRNA

suggests that the protein might also play roles in oosome outside of a potential conserved role in
pole cell formation.

*Nv-qin* encodes a protein containing tudor domains along with an E3 ubiquitin ligase domain. *qin*
is important in processing germline piRNAs, repressing retroelements assembling the nuage, and
proper completion of oogenesis in the fly (51-53). While *qin* has an important late role in
germline cells, it is only weakly and diffusely expressed during embryogenesis in *Drosophila*
(54). Its mammalian homolog *RNF17* is required for production of specific particles in the
germline nuage and for sperm development, but not for early germline specification (55).

*Nv-innexin1* encodes a putative gap junction protein whose fly homolog is most well-known for
its role in proper development and function of the nervous system (56). Other unexpected roles
for innexin proteins have been described and proposed in insects (57), but at the moment the
potential functional significance of the germline localization in *Nasonia* is unclear.

Two transcripts localized to the oosome and preserved in the pole cells have not clear
*Drosophila* homologs (Fig. 3 I1-J3). One of these (*Nv-C16orf87*) encodes a homolog of the
human protein *C16orf87* and is expressed in the posterior region as well as specifically in the
oosome in pre-blastoderm stage (Fig. 3 I1-I3). The protein of *Nv-C16orf87* belongs to the
uncharacterized protein family UPF0547, which contains the zinc-ribbon motif, and functions of
this protein and it homologs are not known. It appears that this gene has been lost specifically in
the Brachyceran fly lineage as it is found in beetles, moths, and some mosquitos, but not
*Drosophila* (JAL personal observation).

Finally, an ankyrin domain encoding transcript is strongly localized to the oosome and is taken
up into pole cells (Fig. 3 J1-J2). It later has a complex and dynamic pattern in the blastoderm
stages  (Fig. 3 J2-J3, additional images not shown). This transcript is a member of the newly
described CLANK (Chalcid Lineage-specific ANKyrin-domain gene) family, of which there are
nearly 200 in the *Nasonia* genome (Lynch and Pers, manuscript in review). To differentiate from
the others, we name it *Nasonia vitiripennis Oosome CLANK* (*Nv-OoCLANK*).

**Fig. 3. Transcripts localized to the oosome that are subsequently maintained in pole cells.**
All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal side on the top.

**Transcripts enriched in the oosome but excluded from pole cells**

A set of 12 genes is expressed in the oosome but not transported to the pole cells. We predicted

that these would have germline roles primarily in the oosome itself, or in the early stages of pole

cell formation. We also expect many transcripts in this set will have roles outside of germline

production, such as in embryonic patterning (as already known for *Nv-dpp* and *Nv-cad*) (58, 59).

Examples of what we consider potential embryonic patterning factors include: a CLIP protease

encoding message (*Nv-mp1*) related to fly Melanization Protease and Easter (Fig. 4 A1-A3), *Nv-*

*kayak* (Fig. 4 B1-B3) encoding at transcription factor downstream of JNK signaling(60), and *Nv-*

*elbow* (Fig. 4 C1-C3) encoding a single zinc-finger transcription factor (61). *Nv-kayak* is later

expressed in a dorsal domain, indicating a conserved role in extraembryonic patterning.

Interestingly, *Drosophila elbow* interacts with *orthodenticle* in specifying the ocelli and in

photoreceptor cell fate determination (62), and an intriguing possibility is that these proteins

work together in posterior patterning, with *Nv-elbow* possibly playing an important role in

differentiating the posterior targets of *otd* from the anterior ones.

Several oosome resident transcripts have suggestive functional annotations. For example, the

*Nasonia coronin* gene (*Nv-coronin,* (Fig. 4 D1-D3)) encodes a protein whose homologs are

known to bind and modulate actin, provide links between the actin and microtubule

cytoskeletons, and regulate endo- and exocytosis in several developmental contexts (63, 64). A

germline role for the *Drosophila* Coronin ortholog has not been observed. *Nasonia milton* (*Nv-*

*milton*) is another exciting transcript (Fig. 4 E1-E3). *Drosophila* Milton is an adaptor protein that

allows mitochondria to be loaded onto, and transported by, microtubule motors (65).

An oosome resident mRNA encodes a protein whose fly ortholog is uncharacterized, but whose

function may be relevant to oosome function. This is the *Nasonia* homolog of *CG4552* (*Nv-*

*CG4552*), which encodes a protein with a TBC25 domain (Fig. 4 F1-F3). Proteins with this

domain interact with Rabs to regulate membrane trafficking and dynamics. Such activities have

been shown to be crucial for Oskar function in the fly (66), and *Nv-CG4552* may play a

supporting role in regulating membrane dynamics in the wasp. Another suggestive localized

factor does not have clear orthologs outside of the hymenoptera, but it does have two predicted

RNA Recognition Motifs, therefore we name it *Nv-rrm* (Fig. 4 G1-G3). RRM domains bind

RNA and are components of proteins that regulate RNA localization and translation. This novel

lineage specific protein could therefore be involved in the localization of specific RNAs in the

oosome, or the regulation of translation of specific RNAs within it.

Many of the oosome localized transcripts do not have annotations that lead to simple hypotheses

about their roles in specifying the germline. One of these is *Nv-waterproof*, which encodes a

fatty acyl-CoA reductase. *Drosophila waterproof* produces the hydrophobic molecules that coat

the trachael tubes during *Drosophila* embryogenesis and is essential for gas filling of the trachea

(67). The protein's novel role in *Nasonia* germline is not clear and worth to investigate, as is its

early maternal expression in the oosome and later zygotic expression as the dorsal strip and in

the extraembryonic tissue. (Fig. 4 H1-H3). *Nv-CG42269* encodes a predicted organic ion

transporter protein whose *Drosophila* homolog (*CG42269*) has no described function (Fig. 4 I1-

I3).

Three oosome localized transcripts have no clear homologs in *Drosophila* or in other model

organisms. *LOC100123551* has a sterile alpha motif (SAM) domain, which might indicate

protein-protein or -RNA interactions (Fig. 4 J1-J3). *LOC103315681* contains weak similarity to

the N-terminal domain of Folded-gastrulation proteins (but is not a folded gastrulation ortholog)

(Fig. 4 K1-K3), while *Nasvi2EG022821* has no discernible conserved domains (Fig. 4 L1-L3).

These functions of these factors will be the object of future investigation.

**Fig. 4. mRNAs localized in the oosome (Pre-blastoderm) but are excluded from the pole cells.** All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal side on the top.

**Transcripts enriched in the posterior pole but not specifically the oosome**

There are five transcripts that are significantly enriched in the posterior region of the embryo, but do not show significant association with the oosome (Fig. 5). The early embryonic expression for these transcripts appears as a cap or broad posterior to anterior gradient of mRNA. The significance of such transcripts to oosome assembly or to germ cell formation is not clear. Two of this class are known developmental transcription factors: orthologs of Zerknuellt (68), and Mothers against dpp (69) (*Nv-zen* and *Nv-mad2, respectively* (there are two mad paralogs in *Nasonia*). Two transcripts are predicted to encode catalytic enzymes: a choline kinase homologous to the *Drosophila CG2201* (*Nv-CG2201*), and a homolog of the ADP ribosylation factor-like 4 protein (*Nv-ADPrf4*). Finally, *LOC100680197* and *LOC100677932* have no identifiable homologs outside of hymenoptera. *LOC100680197* encodes a protein with MYND-type zinc-fingers and a p27-like domain, while *LOC100677932* has no clear functional domains and do not have identifiable conserved domains.

**Fig. 5. mRNAs strongly enriched in the posterior region of the embryos, but not enriched in the oosome.** All embryos are aligned and grouped into three columns (Pre-blastoderm, Blastoderm and Gastrulation) according to their embryogenesis stages, with posterior side to the right and dorsal side on the top.

**Functional analysis of a subset of germplasm localized factors**

While localization of an mRNA to the oosome and pole cells may strongly suggest a function related to PGC specification, demonstration of any such function is required. We chose a sampling of five promising molecules for in depth functional analysis (*Nv-bark*, *Nv-anillin*, *Nv-rrm*, *Nv-coronin*, and *Nv-innexin*). We initially tried to apply our parental RNAi (pRNAi) approach, but quickly found that this was not the ideal approach. Most dsRNAs caused partial sterility, with most of the obtainable eggs being apparently normal escapers. To circumvent this problem, we developed a protocol for embryonic injection of dsRNA followed by fixation and *in situ* hybridization. As a negative control, we injected dsRNA against eGFP. While some damage was visible after injection, embryonic development was largely normal through the blastoderm stage in most of embryos (Fig. 6 A1-A4). We considered the disruptions of developmental events to be potential effects of a knockdown when they were only specific to the knockdown and were never observed in the negative control.

**Fig. 6. Functional analysis of a subset of germplasm localized factors by eRNAi.** A1-A4: eGFP dsRNA injected embryos of increasing as the negative control. B1-B4: *Nv-bark* dsRNA injected embryos, C1-C4: *Nv-anillin* dsRNA injected embryos, D1-D4: *Nv-rrm* dsRNA injected embryos, E1-E4: *Nv-coronin* dsRNA injected embryos, F1-F4: *Nv-innexin1* dsRNA injected embryos. All the embryos are positioned with posterior side to the right and dorsal side on the top. The red in B1-B4 are marked by *Nv-nos*, and the rest are marked by *Nv-bark*. DAPI (white) marks the nuclei.

One transcript that particularly captured our attention was *Nv-bark*, as it was the transcript most strongly and consistently associated with PGC specification over embryonic development. However, its potential function in the germline is not clear. Since it encodes a transmembrane protein involved in epithelial junctions in *Drosophila*, we speculated that it might have a role in mediating adhesion or migration of the pole cells once they were formed. Surprisingly, the phenotypes produced by knocking down this transcript showed a much earlier requirement for this transcript. In early embryos (before migration of nuclei to periphery), the oosome has lost its integrity as a single unit (Fig. 6 B1). Instead there are scattered particles of what appears to be oosome-like material, concentrated around the lateral cortex of the embryo. Later, at the time the pole cells would normally form, no budding is observed, and some loose

aggregates of germplasm like material remains attached to the cortex (Fig. 6 B2-B3). In some
cases, germplasm surrounds nuclei in a way similar to what is seen in pole cells (Fig. 6 B3, B4).
However, these nuclei remain part of the embryonic syncytium.

Like *bark beetle*, the mRNA of *Drosophila anillin* is not localizd to the polar granules (70).
Anillin protein, however, accumulates at the base of pole cells when they are budding in
*Drosophia* (50). Since Anillin is a major component of the contractile ring during mitotic
cytokinesis (49) and is enriched at the bud furrow during pole cell formation in *Drosophila* (50),
we predicted that the enrichment of *Nv-anillin* mRNA in the oosome and pole cells would be
related to *Nasonia's* unique way of forming a single large pole bud instead of several small ones,
as occurs in *Drosophila*. Surprisingly, the phenotype of *Nv-anillin* is indistinguishable from that
of *Nv-bark*:  The oosome does not form properly, and germplasm material remains bound to the
posterior cortex of the embryo, leading to the association of the syncytial nuclei at random
posterior locations (Fig. 6 C1-C4).  The same set of phenotypes is observed when we knock
down *Nv-rrm* with eRNAi (Fig. 6 D1-D4). Thus, three genes with very different predicted
functions all result in the same phenotype.

It is important to note at this point that the common phenotype of the above three knockdowns is
not the same as the complete loss of germ plasm activity. Such phenotypes are seen for *Nv-osk*,
*Nv-vas*, and *Nv-aubergine*. When these genes are knocked down, posterior mRNAs such as *Nv-*
*nos* take on a uniformly graded posterior cap and no enriched accumulation of germ plasm
markers is ever observed at the posterior (35, 40).  This indicates that these transcripts are
involved in the specific form of the oosome, but may not be essential for the production of
germline-like cells.

Finally, despite the fact that the oosome does not form or migrate through the embryo as in
wildtype, a bud like protrusion similar to the one that initiates pole cell formation is regularly
observed (Fig. 6 C3, Fig. 1 E), and this bud is not associated with the bulk of the germplasm like
material. This indicates that neither the oosome, nor its remnants induces the bud, and that bud
formation may be autonomous to the embryo. This has some precedent in *Drosophila*, where the
autonomous ability of the fly embryo to produce pole-cell like structures at both poles of the

embryo is revealed when Arf guanine exchange factor Steppke is reduced (71). However, in the

fly, normal global repression of pole-cell formation is overcome by germ plasm components

(primarily *germ-cell-less*), while in *Nasonia,* at least the initial budding appears to be germplasm

independent.

We tested the functions of two other transcripts with eRNAi: *Nv-coronin* and *Nv-innexin1*.

Knockdown of both genes left the oosome intact, and able to migrate through the embryonic

cytoplasm normally (Fig. 6 E1, F1).  However, in both cases, pole cell formation fails, indicating

that these genes have downstream functions that are specific to pole cell formation (Fig. 6 E2-E4,

F2-F4).  *Nv-coronin*'s functional annotation is consistent with a role in cellularization, as it is

predicted to interact with both the microtubule and actin cytoskeletons, both of which are crucial

for mitosis and cell formation.  While Innexin1 is most well known as a crucial component of

gap junctions and a role in the nervous system (56), a role for at least one innexin in cellurization

has been demonstrated in *Tribolium* (57), suggesting that the full potential of these proteins in

regulating cellular processes has not been fully explored.

**Discussion**

**RNA-seq analyses**

Our results have uncovered an unexpectedly large amount novelty in the mRNA content of the

germline determinant of the wasp *Nasonia vitripennis*.  This was achieved using RNAseq

followed by statistical detection of differential expression of mRNAs between the anterior and

posterior poles of the early embryo. The statistical predictions were then extended by *in situ*

hybridization and RNAi of candidate genes.

Our goal was to identify all transcripts specifically localized to the oosome, and it is worth

contemplating how close we came to achieve this goal. We used three different approaches that

varied in the strategy for mapping, quantifying transcripts and assessing differential expression.

These analyses agreed on the vast majority of the genes with putative significantly different

enrichment at the two poles, and most of the disagreement was at the margin of differential

expression that would be likely biologically significant. In addition, all analyses found genes

previously identified to be localized anteriorly or posteriorly.  We even detected all known genes

with both strong oosome localization and strong localization at the anterior pole. We thought these could be missed, because if the anteriorly localized mRNA population approached the same levels as that in the oosome, the enrichment might be obscured.

For these reasons, we think that we have uncovered the vast majority of mRNAs localized to the *Nasonia* oosome. Of course, no approach can guarantee comprehensiveness, and some molecules may have been missed for several reasons. For example, while the *Nasonia* genome is well annotated, it is possible that a very few genes are not represented in the genome assembly or the transcriptome annotations, and thus would not have been assessed. Indeed, we did identify slightly different sets of genes when using NCBI annotation 102 versus the OGS 2.0 annotation (48), which were created using different approaches to identify and predict genes. Most of the differences are at the margins of significance and low levels of differential expression. However, there are a handful of confirmed localized transcripts detected using annotation 102 that were missed in analyses with OGS 2.0. In addition, at least one transcript was not annotated in annotation 2.1, but was found in annotation 2.0 and was confirmed to be localized to the oosome (*Nasvi2EG001470*, Fig 3D). This is likely a false negative annotation in annotation 102, as a very similar sequence is found in the transcriptome of the very closely related wasp *Trichomalopsis sarcophagae* (72).

Another issue that can cause false negatives is the large number of *Nasonia* transcripts that overlap, which can lead to the concatenation of transcripts. If a localized transcript is fused to a ubiquitous, highly expressed transcript, the signal of localization can be lost. This problem is largely solved by preventing novel junctions when mapping with tophat2. However, this also seem to change the calculation of significance for some transcripts, so performing the analysis with and without novel junctions gives more complete results.

Finally, there are some unknown artifacts which may cause significantly enriched genes to be missed. A prime example of this is our discovery of *Nv-coronin*. A preliminary attempt at this experiment resulted in generally poor sequencing results, and a completely unusable replicate. In general, this analysis gave predictably worse results, where several known localized genes were not found to be significant. However, all of the molecules whose functions were analyzed in this

manuscript were found to be statistically significantly enriched at the posterior, including *Nv-*
*coronin*, and we proceeded to clone them to test their localization and function. In the subsequent
analyses based on high quality sequencing results (used as the basis for this manuscript), *Nv-*
*coronin* was excluded by cuffdiff for statistical testing for unknown reasons, despite showing
similar posterior enrichment that had previously been deemed statistically significant. Thus,
there is a potential for false negatives if this artifact affects several genes. Very few transcripts
in the "not tested" category show a similarly strong posterior enrichment, so we believe that this
effect is also small.

**Comparison of the oosome of *Nasonia* to the polar granules of *Drosophila***
Germ plasm in insect embryos must perform several different functions. First, it must be able to
concentrate and arrange the set of proteins and RNAs (e.g. Tud, Vas, *nos* mRNA, Piwi/Aub) that
are associated with germline fate in the posterior of the egg. Second, it must ensure its own
incorporation into the primordial germ cells as they form by interacting with the specialized
cytoskeletal structures that mediate pole cell formation. Germ plasm must also include molecules
involved specialized features of primordial germline cells. Such functions include the
concentration and selection of mitochondria, repression of transcription, repression of
transposons, and guidance of the germ cell migration to the gonad primordia.

On the other hand, we had already observed that the oosome's morphology and interaction with
the cytoskeleton (during its migration and formation of the pole cells) was quite distinct from
*Drosophila*, as is the way the pole cells migrate into the interior of the embryo. Therefore, we
expected to find a mixture of conservation and novelty when we examined the mRNA content of
the oosome. Indeed, this is what was found, but with a surprisingly strong bias toward novelty.

In terms of conservation, our analyses found *Nv-nos, Nv-osk* (already known factors found in
both oosome and polar granules) as well as *Nv-ovo*. We also found *Nv-aub* whose ortholog is
localized as protein, but not mRNA, in *Drosophila*. While there are a large number of polar
granule localized mRNAs that are not found in the oosome, we will only discuss a few that are
significant.

mRNA for *polar granule component (pgc)* encodes a small peptide (73), and is strongly localized to the posterior pole (74). Polar-granule-component protein has a crucial role in the global repression of transcription that occurs in pole cells upon their formation, through an interaction with the transcription elongation factor TEF-b (73). This repression is a widely conserved feature of primordial germ cells across animals, which makes it somewhat surprising that *pgc* appears to be a novelty in the *Drosophila* lineage (74).

Fascinatingly, TEF-b may be a unifying factor underlying, as it interacts with PIE-1 in the worm *C. elegans* to repress transcription (75). In line with this, we have found a highly conserved transcription elongation factor, *Nv-spt5* (Fig. 3 A1-A3), localized strongly to the oosome and pole cells. In human cells, Spt5 acts as an inhibitor or transcriptional elongation, until its C-terminal domain is phosphorylated in a TEF-b dependent manner (76). If an interaction between *Nasonia* Spt5 and TEF-b does have a role in regulating the cessation of transcription in *Nasonia* pole cells, it would be strong evidence for TEF-b being a core conserved component of the germline fate, whose interaction partners and regulators are extremely labile across lineages.

Another crucial *Drosophila* germ cell factor that is not present in the *Nasonia* oosome is *germ cell less (gcl)*. The Germ-cell-less protein itself is very highly conserved at the sequence level in *Nasonia,* but the mRNA showed no enrichment in our RNAseq experiments, and we independently confirmed by *in situ* hybridization that it is expressed uniformly throughout the early embryo (not shown). Gcl is important for the proper production of pole cells, apparently by regulating the orientation centrosome separation at the posterior pole, which is required for efficient pole cell formation and uptake polar granules by the pole cells (77). At the molecular level, Gcl seems to act by downregulating torso signaling, to allow the proper conditions for pole cells to form. The lack of Gcl function in the germline of *Nasonia* is consistent with the lack of Torso signaling at the termini in the wasp (78), making the need for Gcl redundant.  At the moment, it is not clear whether the use of Gcl in pole cell formation is a recent novelty in *Drosophila*, or whether it was present ancestrally, but lost in the Hymenopteran lineage.

A number of transcripts found in the *Nasonia* oosome are good candidates for generating essential PGC features. For example, a high concentration of mitochondria is a strongly

conserved feature of germ plasm and primordial germ cells across animals (79, 80). In

*Drosophila*, the long Oskar isoform plays an important role in concentrating mitochondria in the

pole plasm (81). But, since the long form of Oskar appears to be a novelty of *Drosophila* and its

close relatives, other molecules should be expected to perform this role in other species.

Suggestively, mRNA encoding a Milton ortholog was found strongly localized to the *Nasonia*

oosome. Milton acts an adaptor that loads mitochondria onto microtubule motors for transport

and localization within and between cells in *Drosophila* (82), and we propose that *Nasonia*

Milton plays a role in enriching mitochondria around the oosome and in the pole cells in the

wasp, and perhaps other insect species that lack the specialization of long Oskar isoform.

Another critical function for germ cells is the control of transposable elements, which is often

dependent on one or more Tudor domain containing protein. mRNAs for two Tudor domain

proteins are present in the oosome, including *Nv-qin* and *Nv-tdrd7*. Neither of them is enriched in

the polar granules or pole cells in *Drosophila*, but both have crucial roles during oogenesis to

reduce the activity of transposable elements (51, 52, 83). The presence of these additional Tudor

domain proteins may indicate that either there is an increased activity of transposable elements in

*Nasonia* that requires an earlier response, or perhaps other mechanisms are employed in

*Drosophila* to combat transposon activity in the early PGCs. Further sampling of germ plasm of

other insects should help to resolve these questions.

Germ cells are known to have a distinct metabolic profile from somatic cells, and this difference

is related to their pluripotent stem cell-like properties, and to the requirements of their migratory

properties (84, 85). Potentially related to this we have found that a *Nasonia* insulin-like growth

factor I mRNA is localized to the oosome. Interestingly, this mRNA encodes a short protein that

shows much stronger similarity to insulin proteins of vertebrates, than it does to any of the

insulin-like molecules of *Drosophila* (DILPs) (JAL personal observation). In addition, an mRNA

encoding a putative organic cation transporter (*Nv-CG42269)* containing a Major Facilitator

Superfamily (MFS) domain is strongly localized to the oosome. Such molecules are crucial for

regulating cellular metabolism and signaling at multiple levels, by controlling the trafficking of

many small organic molecules (including sugars) within and between cells (86, 87). Lipid

metabolism is also uniquely regulated in germ cells, and the identification of the *Nasonia*

homolog of the Acyl-CoA reductase *waterproof* may reflect this (88). Finally, we surprisingly

found a transcript encoding an protein similar to endoglucanases found in several insect lineages

(but absent from Diptera). Enzymes of this type are broadly defined by their ability to break

down polymers glucose. The substrate and potential role for this enzyme in the germline is not

yet known.

In addition to providing insight into the conserved functions of germ cell components, we also

found several molecules that do not have clear homology outside of the Hymenoptera, or in some

cases outside of *Nasonia* and its closest relatives. This includes a novel RNA recognition domain

containing protein whose function we analyzed in depth (*Nv-rrm* discussed in the following

section).  We also found that an mRNA encoding an ankyrin domain protein (*Nv-OoCLANK*

(LOC100679945)) that belongs to a family of proteins that underwent a massive amplification

within chalcid wasp lineage, and which appears to have entered the ancestral chalcid wasp by

horizontal transfer from an endosymbiont (Pers and Lynch, in review). Finally, a handful of

transcripts have no identifiable domains or homologs. The functional relevance of these

molecules will be an area of intense interest in the future.

**Unexpected functions of novel oosome components**

Early in our analysis we chose a handful of transcripts for functional analysis.  These were

chosen based on a combination of criteria that included high enrichment in the oosome, novelty,

and the potential to give phenotypes that we could characterize with the current set of functional

tools available to us in *Nasonia*. Three of these (*Nv-bark*, *Nv-anillin* and *Nv-rrm*) gave an

unexpected phenotype, where oosome-like material was not coalesced into the typical spherical

oosome structure, but rather was scattered in clumps attached to the plasma membrane near the

posterior pole. Eventually, these clumps of germ plasm-like material come into contact with

syncytial nuclei when they migrate to the cortex.  However, this remnant material is unable to

induce the pole cell fate.  It is important to reiterate that this phenotype is quite distinct from that

seen for genes that have a core role in oosome assembly (*Nv-osk*, *Nv-tud*, *Nv-vas* and *Nv-aub*

(not shown).  In these cases, there is no hint of the oosome, and mRNAs normally localized in

the oosome are distributed in homogenous caps at the posterior pole*,* rather than as discrete

clumps of material (35).

These phenotypes indicate these genes are involved in the coalescence of the oosome into single

entity within the central column of cytoplasm in the embryo, and/or maintenance of oosome

integrity.   The molecular bases of such functions are not completely clear at the moment for any

of these three genes. *Nasonia* Bark is a putative transmembrane protein and would be predicted

to be targeted to the membrane.  One hypothesis could be that *Nasonia* Bark targeted to the

membrane at the posterior pole of the embryo, releases and/or repels oosome material from

adhering to the plasma membrane, forcing the oosome to remain in the bulk cytoplasm where it

concentrated into a large sphere and moved around by strong cytoplasmic flows that occur

during the earliest cleavages. This model would also imply that interaction with the cortex

prevents oosome material from coalescing, leading to the scattered clumps we observe in *Nv-*

*bark* knockdowns.   Mechanistically, this could be related to the ability of Bark to induce

endocytosis (89)*,* a process that is associated with proper anchoring of the germ plasm and the

recruitment of specialized actin binding proteins to the posterior pole in *Drosophila* (90).

An alternative hypothesis is that *Nasonia* Bark produced in the oosome is not secreted but is

instead incorporated as an important structural component of the large, solid form of the oosome.

This is consistent with the structure of the protein, which contains main protein-protein

interaction domains of different types (44, 45). Testing these hypotheses will require in depth

analysis of the subcellular localization of *Nasonia* Bark during early embryogenesis, and

proteomic analysis of binding partners of *Nasonia* Bark.

The role of *Nv-anillin* in maintaining the stability of the oosome was also surprising and the

molecular basis of the phenotype will require further investigation. Anillin orthologs are well

known as actin binding proteins involved in assembling the contractile ring required to separate

cells in cytokinesis (49).  Anillin also plays a crucial and novel role in the specialized cytokinesis

of the *Drosophila* pole cells (50). While these known functions might have indicated that

oosome localization of *Nv-anillin* was related to an important role in the specialized polar bud

formed in *Nasonia*, RNAi showed that this protein has an earlier role in oosome

assembly/maintenance (Fig. 6 C1). Similar to *Nasonia* Bark, one possible function of *Nasonia*

Anillin is as a structural component of the oosome, which may or may not be related to its ability

to bind actin and associated proteins. Alternatively, *Nv-anillin* may act to release and/or repel the

oosome from the cortex, as proposed above for *Nv-bark*.

Anillin homologs have known functions which are directly related to the processes of release of

germ plasm from the embryonic plasma membrane. In *Drosophila*, the germ plasm is tightly

bound to the plasma membrane until the nuclei reach the posterior pole of the embryo. The

centrosomes associated with these nuclei mediate detachment of the pole plasm from the cortex

through interactions of the astral microtubules emanating from the centrosomes (91). This is in

contrast to *Nasonia,* where the oosome detaches from the cortex at about the same time as the

zygotic nucleus begins its first division at the anterior pole. While nuclei are lacking at this time,

numerous centrosomes are present, as they are provided maternally in a process characteristic of

many Hymenopteran embryos (92). *Nasonia* Anillin could be relevant to a model where astral

microtubules emanating from maternally provided centrosomes detach oosome material from the

cortex, because Anillin homologs have been shown to mediate interactions between the actin

cytoskeleton and cortical and subcortical microtubule arrays in multiple model systems (93, 94).

Again, in depth examination of the subcellular localization and interactions of *Nasonia* Anillin

will be required to completely understand its role in maintaining the oosome.

Based on the presence of only RNA recognition motifs in the protein, we predict that *Nv-rrm*

will have one of two likely roles. One possibility is that it is involved in translational regulation

of key regulators of oosome structure, presumably including *Nasonia* Anillin and *Nasonia* Bark.

Alternatively, *Nv-rrm* may be important in binding RNA and protein in order to maintain the

structural integrity of the oosome. In addition, neither of these possibilities is mutually exclusive.

The knockdowns of *Nv-coronin* and *Nv-innexin* had specific effects only on the formation of

pole cells, while the oosome appeared to remain intact. Coronin is an actin binding protein

associated with the formation of highly concentrated networks of F-actin (64). It seems likely

that *Nasonia* Coronin has an important role in organizing an actin cytoskeleton arrangement

specialized for the formation of the large polar bud that initiates pole cell formation.

The potential role of *Nv-innexin1* is somewhat more mysterious. Innexins are typically known as

components of the gap junctions that are found in some tightly integrated epithelial tissues. Such

junctions would not be expected of the motile pole cells. Interestingly, Innexin-7 (a paralogous

protein with a similar structure to *Nasonia* Innexin1) in the beetle *Tribolium* has a novel role in

cellularization of the syncytial blastoderm. Such a function for *Nasonia* Innexin1 would explain

the failure in pole cell formation we see after RNAi.

**Conclusion**

This work has revealed numerous unexpected mRNAs that are localized to the germ plasm of the

wasp *Nasonia.* The results have given insights on the potentially ancestral mechanisms used by

germ plasm to accomplish conserved required functions, such as a possible ancestral role for

Milton orthologs to bring mitochondria to the germplasm, a function replaced by the long Oskar

isoform in *Drosophila.* On the other hand, our results have identified numerous components that

are likely to be specific to *Nasonia* and its relatives in the parasitic wasp lineage (e.g., *Nv-*

*OoCLANK, Nv-rrm*, and the use of Nv-Bark and Nv-Anillin in assembling the oosome). Deeper

analyses of the functions of these molecules in *Nasonia*, and broader sampling of germ plasm in

other holometabolous insects will be required to determine the patterns of evolutionary change in

the germ plasm. Such analyses will be important because the germ plasm is a uniquely powerful

organelle that can rapidly drive naive nuclei into a highly specialized, yet functionally totipotent

state. Understanding how and why such a fundamental substance changes and is even lost in the

course of evolution will provide foundational insights into the mechanisms of cell fate

determination and the interaction of subcellular organelles and their cellular milieu.

**Materials and Methods**

**Sample preparation**

In order to identify the components of the maternally deposited mRNAs in the oosome located in

the posterior half of the parasitoid wasp *Nasonia vitripennis* embryos, we collected and bisected

the pre-blastoderm stage embryos (0-2 hours after egg lay at 25 °C) to detect the differential

expression levels between the anterior and posterior halves of the embryos.

The embryos were aligned with the anterior pole to the right on the ice-prechilled and heptane
glue-coated slide. Then a thin layer of halocarbon oil 700 (Sigma) was applied to cover the
embryos. The slide was transferred on the dry ice-prechilled "guillotine" and then was anchored
by tightening the screws on each end. After the halocarbon oil 700 was solidified on dry ice, put
the guillotine on the dry ice-prechilled stabilizer and transferred it under the dissection
microscope. The embryos were positioned to match the slot in the guillotine where the dry ice-
prechilled razor blade will be inserted into. After the embryos were cut, the anterior and posterior
halves of the embryos were immediately collected and transferred into the two 1.5mL non-stick
RNase-Free microfuge tubes (Ambion) with the dry ice-prechilled probe, separately. Three
biological replicates were created.

Total RNA was isolated from these six samples for library preparation. In the  library preparation
upon which this manuscript is based, we used around one microgram of total RNA from each
sample, in which 100ng of total RNA was from the *Nasonia* other 900ug was from a distantly
related parasitic wasp (*Melittobia digitata* (Pers and Lynch, in revision). Libraries were prepared
using the NEBNext Ultra Directional RNA Library Prep Kit for Illumina (NEB #E7420) in
conjunction with NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB
#E7490). Libraries were validated and quantified before being pooled and sequenced on an
Illumina HiSeq 2000 sequencer with a 100 bp paired-end protocol. Sequence files are available
in the NCBI SRA database under accession SRP156232.

**RNA sequencing data analysis**
The quality of the sequencing data was determined using FastQC software. The sequences were
processed by Cufflinks package for differential expression detection, using multiple variations on
the default parameters (job files in Supplemental File 2).  Briefly, raw reads were aligned to
annotation 102 of assembly 2.1 of the *N. vitripennis* genome
(https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Nasonia_vitripennis/102/) using
TopHat2(95). These results were either used directly in cuffdiff (96), or were further processed
using stringtie (97) to generate new transcriptome predictions and quantification. Various
normalization parameters were used, and each permutation gave slightly different results. In
addition, we mapped the reads using assembly 1.0 and annotation 2.0 (OGS 2.0) (48). All of the

commands jobs in this process of analysis are included in the Supplemental File 2. The cuffdiff

results of all permutations of the analysis are provided in Supplemental File 3. The computing

work was done by the High Performance Computing Cluster located at University of Illinois at

Chicago.

Probes and dsRNAs for the chosen genes were generated by the protocol described in (40).

Primers for generating these templates are provided in Supplemental File 4. Alkaline

phosphatase *in situ* hybridization was performed by the protocol described in (98).

**Embryonic RNAi**

In order to perform embryonic RNA interference (eRNAi) on the early *Nasonia* embryos to

study the germline candidate genes' functions, we created this workflow.

Around 30 pre-blastoderm stage embryos (0-1 hour after egg lay at 25 °C) were collected and

quickly aligned vertically on the heptane glue-coated 18mm×18mm coverslip. This coverslip

was transferred and anchored on the ice-prechilled slide by applying a thin layer of water. The

slide was then put in an air tight petri dish with proper amount of desiccant (Drierite with

indicator, 8 mesh, ACROS Organics) pre-chilled at 4 °C. The embryos were dehydrated in the

desiccant at 4 °C for 45 minutes. After dehydration, the embryos were covered with a layer of

halocarbon oil 700 and were ready for microinjection.

The dsRNAs were dissolved in Nuclease-Free Water (Ambion) at the concentration of 1mg/mL

and loaded into the Femtotips II Microinjection Capillary (Cat. No. 930000043, Eppendorf). The

constant pressure was set at 500hpa and the injection pressure was set at 250hpa. The pressures

needed to be adjusted accordingly to avoid bursting the embryos or sucking cytoplasm into the

capillaries. The process of injection was performed at room temperature and needed to be done

as soon as possible. After injection, the slide was transferred into a paper towel-moisturized petri

dish pre-warmed at 28 °C to incubate the injected embryos for specific developmental stages.

The embryogenesis of these embryos was stopped at pre-blastoderm stage (before the budding),

beginning of blastoderm stage (during budding), and later in blastoderm stage (pole cells formed

and/or after pole cell divisions). To stop the development, the coverslip was put into the heptane

to wash off the halocarbon oil 700 for three minutes, and then transferred into the 37%

formaldehyde-saturated heptane for 2-5 hours fixation in the dark with the embryos facing up.

After fixation, the coverslip was carefully taken out of the fixative, and flipped upside down to

gently press the embryos on a double-sided tape that was taped on a petri dish, so that all the

embryos can be anchored on the tape for dissection. Add about 15mL PBS with 1% Tween, use

the needle (BD PrecisionGlide Needle, 30G × 1) to carefully remove the eggshells from the

embryos. The dissected embryos were then transferred by pipette into the 1.5mL non-stick

RNase-Free microfuge tubes. The embryos were immediately dehydrated by 100% Methanol and

stored at -20 °C.

Before performing fluorescent *in situ* hybridization (FISH) on those eRNAi knocked out

embryos, they need to be rehydrated with a series of Methanol/PBT washes (75%, 50%, 25%).

The protocol for FISH was adapted from (40). A detailed protocol is available on request.

**Acknowledgments:**

We thank Daniel Pers for performing the combined *Nasonia/ Melittobia* RNAseq libraries, and

the UIC RRC Sequencing core for arranging the sequencing of our samples. This work was

supported by startup funds from UIC, and NIH grants 1R03HD087476 and 1R03HD078578.

**Competing Interests:**

The authors declare no competing interests.

# References

1.        Extavour CG, Akam M. Mechanisms of germ cell specification across the metazoans:

epigenesis and preformation. Development. 2003;130(24):5869-84. Epub 2003/11/05. doi:

10.1242/dev.00804. PubMed PMID: 14597570.

2. Extavour CG. Evolution of the bilaterian germ line: lineage origin and modulation of

specification mechanisms. Integr Comp Biol. 2007;47(5):770-85. Epub 2007/11/01. doi:

10.1093/icb/icm027. PubMed PMID: 21669758.

3. Hegner RW. Experiments with Chrysomelid Beetles. III. The Effects of Killing Parts of the

Eggs of Leptinotarsa decemlineata. Biological Bulletin 1911;20:237-51.

4. Illmensee K, and A. P. Mahowald. Transplantation of posterior polar plasm in Drosophila.

Induction of germ cells at the anterior pole of the egg. Proceedings of the National Academy of

Sciences. 1974;71(4):1016-20.

5. Illmensee K, and A. P. Mahowald. The autonomous function of germ plasm in a somatic

region of the Drosophila egg. Experimental cell research 1976;97(1):127-40.

6. Xia QY, Zhou, Z. Y., Lu, C., Cheng, D. J., Dai, F. Y., et al. A draft sequence for the genome

of the domesticated silkworm (Bombyx mori). Science. 2004;306:1937-40.

7. Nagy L, Riddiford, L., Kiguchi, K. . Morphogenesis in the Early Embryo of the

Lepidopteran Bombyx-Mori. Developmental Biology. 1994;165:137-51.

8. Juhn J, Marinotti, O., Calvo, E., James, A. A. Gene structure and expression of nanos (nos)

and oskar (osk) orthologues of the vector mosquito, Culex quinquefasciatus. Insect Molecular

Biology. 2008;17:545-52.

9. Suzuki N, Shimizu, S., Ando, H. . Early Embryology of the Alderfly, Sialis-Mitsuhashii

Okamoto (Megaloptera, Sialidae). International Journal of Insect

Morphology & Embryology. 1981;10:409-18.

10. Werren JH, Richards, S., Desjardins, C. A., Niehuis, O., Gadau, J. Functional and

Evolutionary Insights from the Genomes of Three Parasitoid Nasonia Species. Science.327:343-8.

11. Berg GJ, Gassner, G. Fine structure of the blastoderm embryo of the pink bollworm,

Pectinophora Gossypiella (Saunders) (lepidoptera: Gelechiidae). International Journal of Insect

Morphology and Embryology 1978;7:81–105.

12. Miura T, Braendle, C., Shingleton,  A., Sisk, G., Kambhampati, S., Stern, D. L. A

comparison of parthenogenetic and sexual embryogenesis of the pea aphid Acyrthosiphon

pisum (Hemiptera : Aphidoidea). Journal of Experimental Zoology Part B-Molecular and

Developmental Evolution 2003;295B:59-81. doi: 10.1002/jez.b.00003.

13.    Bull AL. Stages of Living Embryos in the Jewel Wasp Mormoniella-

(Nasonia)-Vitripennis-(Walker)(Hymenoptera, Pteromalidae). Journal of Insect Morphology &

Embryology. 1982;11:1-23.

14.    Klag J. Differentiation of primordial germ cells in the embryonic development of

Thermobia domestica. Pack. (Thysanura): an ultrastructural study. Journal of Embryology and

Experimental Morphology. 1977;38:93-114.

15.    Chang C-c, Dearden P, Akam M. Germ Line Development in the Grasshopper

Schistocerca gregaria: vasa As a Marker. Developmental Biology. 2002;252(1):100-18. doi:

10.1006/dbio.2002.0840.

16.    Consortium THGS. Insights into social insects from the genome of the honeybee Apis

mellifera. Nature. 2006;443(7114):931-49. doi: 10.1038/nature05260.

17.    Dearden PK. Germ cell development in the Honeybee (Apis mellifera); vasa and nanos

expression. BMC Dev Biol. 2006;6:6. Epub 2006/03/01. doi: 10.1186/1471-213X-6-6. PubMed

PMID: 16503992; PMCID: PMC1388196.

18.    Dearden PK, Wilson MJ, Sablan L, Osborne PW, Havler M, McNaughton E, Kimura K,

Milshina NV, Hasselmann M, Gempe T, Schioett M, Brown SJ, Elsik CG, Holland PW, Kadowaki T,

Beye M. Patterns of conservation and change in honey bee developmental genes. Genome Res.

2006;16(11):1376-84. Epub 2006/10/27. doi: 10.1101/gr.5108606. PubMed PMID: 17065607;

PMCID: PMC1626639.

19.    Juhn J, James AA. oskar gene expression in the vector mosquitoes, Anopheles gambiae

and Aedes aegypti. Insect Mol Biol. 2006;15(3):363-72. Epub 2006/06/08. doi: 10.1111/j.1365-

2583.2006.00655.x. PubMed PMID: 16756555.

20.    Nakao H, Hatakeyama M, Lee JM, Shimoda M, Kanda T. Expression pattern of Bombyx

vasa-like (BmVLG) protein and its implications in germ cell development. Dev Genes Evol.

2006;216(2):94-9. Epub 2005/11/02. doi: 10.1007/s00427-005-0033-8. PubMed PMID:

16261344.

21.    Schroder R. vasa mRNA accumulates at the posterior pole during blastoderm formation

in the flour beetle Tribolium castaneum. Dev Genes Evol. 2006;216(5):277-83. Epub 2006/02/17.

doi: 10.1007/s00427-005-0054-3. PubMed PMID: 16482438.

22.     Khila A, Abouheif E. Reproductive constraint is a developmental mechanism that

maintains social harmony in advanced ant societies. Proc Natl Acad Sci U S A.

2008;105(46):17884-9. Epub 2008/11/14. doi: 10.1073/pnas.0807351105. PubMed PMID:

19004767; PMCID: PMC2584687.

23.     Mito T, Nakamura T, Sarashina I, Chang CC, Ogawa S, Ohuchi H, Noji S. Dynamic

expression patterns of vasa during embryogenesis in the cricket Gryllus bimaculatus. Dev Genes

Evol. 2008;218(7):381-7. Epub 2008/06/11. doi: 10.1007/s00427-008-0226-z. PubMed PMID:

18542998.

24.     Tribolium Genome Sequencing C, Richards S, Gibbs RA, Weinstock GM, Brown SJ, Denell

R, Beeman RW, Gibbs R, Beeman RW, Brown SJ, Bucher G, Friedrich M, Grimmelikhuijzen CJ,

Klingler M, Lorenzen M, Richards S, Roth S, Schroder R, Tautz D, Zdobnov EM, Muzny D, Gibbs

RA, Weinstock GM, Attaway T, Bell S, Buhay CJ, Chandrabose MN, Chavez D, Clerk-Blankenburg

KP, Cree A, Dao M, Davis C, Chacko J, Dinh H, Dugan-Rocha S, Fowler G, Garner TT, Garnes J,

Gnirke A, Hawes A, Hernandez J, Hines S, Holder M, Hume J, Jhangiani SN, Joshi V, Khan ZM,

Jackson L, Kovar C, Kowis A, Lee S, Lewis LR, Margolis J, Morgan M, Nazareth LV, Nguyen N,

Okwuonu G, Parker D, Richards S, Ruiz SJ, Santibanez J, Savard J, Scherer SE, Schneider B,

Sodergren E, Tautz D, Vattahil S, Villasana D, White CS, Wright R, Park Y, Beeman RW, Lord J,

Oppert B, Lorenzen M, Brown S, Wang L, Savard J, Tautz D, Richards S, Weinstock G, Gibbs RA,

Liu Y, Worley K, Weinstock G, Elsik CG, Reese JT, Elhaik E, Landan G, Graur D, Arensburger P,

Atkinson P, Beeman RW, Beidler J, Brown SJ, Demuth JP, Drury DW, Du YZ, Fujiwara H, Lorenzen

M, Maselli V, Osanai M, Park Y, Robertson HM, Tu Z, Wang JJ, Wang S, Richards S, Song H,

Zhang L, Sodergren E, Werner D, Stanke M, Morgenstern B, Solovyev V, Kosarev P, Brown G,

Chen HC, Ermolaeva O, Hlavina W, Kapustin Y, Kiryutin B, Kitts P, Maglott D, Pruitt K, Sapojnikov

V, Souvorov A, Mackey AJ, Waterhouse RM, Wyder S, Zdobnov EM, Zdobnov EM, Wyder S,

Kriventseva EV, Kadowaki T, Bork P, Aranda M, Bao R, Beermann A, Berns N, Bolognesi R,

Bonneton F, Bopp D, Brown SJ, Bucher G, Butts T, Chaumot A, Denell RE, Ferrier DE, Friedrich M,

Gordon CM, Jindra M, Klingler M, Lan Q, Lattorff HM, Laudet V, von Levetsow C, Liu Z, Lutz R,

Lynch JA, da Fonseca RN, Posnien N, Reuter R, Roth S, Savard J, Schinko JB, Schmitt C,

Schoppmeier M, Schroder R, Shippy TD, Simonnet F, Marques-Souza H, Tautz D, Tomoyasu Y,

Trauner J, Van der Zee M, Vervoort M, Wittkopp N, Wimmer EA, Yang X, Jones AK, Sattelle DB,

Ebert PR, Nelson D, Scott JG, Beeman RW, Muthukrishnan S, Kramer KJ, Arakane Y, Beeman RW,

Zhu Q, Hogenkamp D, Dixit R, Oppert B, Jiang H, Zou Z, Marshall J, Elpidina E, Vinokurov K,

Oppert C, Zou Z, Evans J, Lu Z, Zhao P, Sumathipala N, Altincicek B, Vilcinskas A, Williams M,

Hultmark D, Hetru C, Jiang H, Grimmelikhuijzen CJ, Hauser F, Cazzamali G, Williamson M, Park Y,

Li B, Tanaka Y, Predel R, Neupert S, Schachtner J, Verleyen P, Raible F, Bork P, Friedrich M,

Walden KK, Robertson HM, Angeli S, Foret S, Bucher G, Schuetz S, Maleszka R, Wimmer EA,

Beeman RW, Lorenzen M, Tomoyasu Y, Miller SC, Grossmann D, Bucher G. The genome of the

model beetle and pest Tribolium castaneum. Nature. 2008;452(7190):949-55. Epub 2008/03/26.

doi: 10.1038/nature06784. PubMed PMID: 18362917.

25.    Consortium TIAG. Genome Sequence of the Pea Aphid Acyrthosiphon pisum. PLoS Biol.

2010. doi: 10.1371/

10.1371/journal.pbio.1000313.g001.

26.    Khila A, Abouheif E. Evaluating the role of reproductive constraints in ant social

evolution. Philos Trans R Soc Lond B Biol Sci. 2010;365(1540):617-30. Epub 2010/01/20. doi:

10.1098/rstb.2009.0257. PubMed PMID: 20083637; PMCID: PMC2817144.

27.    Shigenobu S, Bickel RD, Brisson JA, Butts T, Chang CC, Christiaens O, Davis GK, Duncan EJ,

Ferrier DE, Iga M, Janssen R, Lin GW, Lu HL, McGregor AP, Miura T, Smagghe G, Smith JM, van

der Zee M, Velarde RA, Wilson MJ, Dearden PK, Stern DL. Comprehensive survey of

developmental genes in the pea aphid, Acyrthosiphon pisum: frequent lineage-specific

duplications and losses of developmental genes. Insect Mol Biol. 2010;19 Suppl 2:47-62. Epub

2010/05/28. doi: 10.1111/j.1365-2583.2009.00944.x. PubMed PMID: 20482639.

28.    Bontems F, Stein A, Marlow F, Lyautey J, Gupta T, Mullins MC, Dosch R. Bucky ball

organizes germ plasm assembly in zebrafish. Curr Biol. 2009;19(5):414-22. Epub 2009/03/03.

doi: 10.1016/j.cub.2009.01.038. PubMed PMID: 19249209.

29.    Lehmann R, Nussleinvolhard C. Abdominal Segmentation, Pole Cell-Formation, and

Embryonic Polarity Require the Localized Activity of Oskar, a Maternal Gene in Drosophila. Cell.

1986;47(1):141-52. doi: Doi 10.1016/0092-8674(86)90375-2. PubMed PMID:

WOS:A1986E471700017.

30.    Ephrussi ALKD, L. K.; Lehmann, R. oskar organizes the germ plasm and directs

localization of the posterior determinant nanos. Cell. 1991;66(1):14.

31.    Ephrussi A, Lehmann R. Induction of Germ-Cell Formation by Oskar. Nature.

1992;358(6385):387-92. doi: Doi 10.1038/358387a0. PubMed PMID: WOS:A1992JF85300046.

32.    Rongo CG, E. R.; Lehmann, R. Localization of oskar RNA regulates oskar translation and

requires Oskar protein. Development. 1995;121(9):2737-46.

33.    Breitwieser W, Markussen, F. H., Horstmann, H., Ephrussi, A. Oskar protein interaction

with Vasa represents an essential step in polar granule assembly. Genes & Development.

1996;10:2179-88.

34.    Boswell RE, Mahowald AP. Tudor, a Gene Required for Assembly of the Germ Plasm in

Drosophila-Melanogaster. Cell. 1985;43(1):97-104. doi: Doi 10.1016/0092-8674(85)90015-7.

PubMed PMID: WOS:A1985AVE3900012.

35.    Lynch JA, Ozuak O, Khila A, Abouheif E, Desplan C, Roth S. The phylogenetic origin of

oskar coincided with the origin of maternally provisioned germ plasm and pole cells at the base

of the Holometabola. PLoS genetics. 2011;7(4):e1002029. Epub 2011/05/10. doi:

10.1371/journal.pgen.1002029. PubMed PMID: 21552321; PMCID: 3084197.

36.    Jaglarz MK, Howard, K. R. The active migration of Drosophila primordial germ cells.

Development. 1995;121:3495-503.

37.    Wylie A, Jones AE, D'Brot A, Lu WJ, Kurtz P, Moran JV, Rakheja D, Chen KS, Hammer RE,

Comerford SA, Amatruda JF, Abrams JM. p53 genes function to restrain mobile elements. Genes

Dev. 2016;30(1):64-77. doi: 10.1101/gad.266098.115. PubMed PMID: 26701264; PMCID:

PMC4701979.

38.    Tiwari B, Kurtz P, Jones AE, Wylie A, Amatruda JF, Boggupalli DP, Gonsalvez GB, Abrams

JM. Retrotransposons Mimic Germ Plasm Determinants to Promote Transgenerational

Inheritance. Curr Biol. 2017;27(19):3010-6 e3. doi: 10.1016/j.cub.2017.08.036. PubMed PMID:

28966088; PMCID: PMC5639916.

39.    Bantock CR. Experiments on chromosome elimination in the gall midge, Mayetiola

destructor. Journal of Embryology and Experimental Morphology. 1970;24:257-89.

40.    Lynch JA, Desplan C. Novel modes of localization and function of nanos in the wasp

Nasonia. Development. 2010;137(22):3813-21. Epub 2010/10/12. doi: 10.1242/dev.054213.

PubMed PMID: 20929949; PMCID: 3049278.

41.    Santos AC, Lehmann R. Germ cell specification and migration in Drosophila and beyond.

Curr Biol. 2004;14(14):R578-89. Epub 2004/07/23. doi: 10.1016/j.cub.2004.07.018. PubMed

PMID: 15268881.

42.    Draper BWM, C. C.; Bowerman, B.; Hardin, J.; Priess, J. R. . MEX-3 Is a KH Domain Protein

That Regulates Blastomere Identity in Early C. elegans Embryos. Cell. 1996;87:205-16.

43.    Lynch JA, Brent AE, Leaf DS, Pultz MA, Desplan C. Localized maternal orthodenticle

patterns anterior and posterior in the long germ wasp Nasonia. Nature. 2006;439(7077):728-32.

Epub 2006/02/10. doi: 10.1038/nature04445. PubMed PMID: 16467838.

44.    Byri S, Misra T, Syed ZA, Batz T, Shah J, Boril L, Glashauser J, Aegerter-Wilmsen T, Matzat

T, Moussian B, Uv A, Luschnig S. The Triple-Repeat Protein Anakonda Controls Epithelial

Tricellular Junction Formation in Drosophila. Dev Cell. 2015;33(5):535-48. Epub 2015/05/20. doi:

10.1016/j.devcel.2015.03.023. PubMed PMID: 25982676.

45.    Hildebrandt A, Pflanz R, Behr M, Tarp T, Riedel D, Schuh R. Bark beetle controls

epithelial morphogenesis by septate junction maturation in Drosophila. Dev Biol.

2015;400(2):237-47. Epub 2015/02/24. doi: 10.1016/j.ydbio.2015.02.008. PubMed PMID:

25704509.

46.    Jennings BH, Shah S, Yamaguchi Y, Seki M, Phillips RG, Handa H, Ish-Horowicz D. Locus-

specific requirements for Spt5 in transcriptional activation and repression in Drosophila. Curr

Biol. 2004;14(18):1680-4. Epub 2004/09/24. doi: 10.1016/j.cub.2004.08.066. PubMed PMID:

15380072.

47.    Pers D, Buchta T, Ozuak O, Wolff S, Pietsch JM, Memon MB, Roth S, Lynch JA. Global

analysis of dorsoventral patterning in the wasp Nasonia reveals extensive incorporation of

novelty in a regulatory network. BMC Biol. 2016;14:63. Epub 2016/08/03. doi: 10.1186/s12915-

016-0285-y. PubMed PMID: 27480122; PMCID: PMC4968023.

48.    Rago A, Gilbert DG, Choi JH, Sackton TB, Wang X, Kelkar YD, Werren JH, Colbourne JK.

OGS2: genome re-annotation of the jewel wasp Nasonia vitripennis. BMC Genomics.

2016;17:678. doi: 10.1186/s12864-016-2886-9. PubMed PMID: 27561358; PMCID:

PMC5000498.

49.     Field CM, Alberts BM. Anillin, a Contractile Ring Protein That Cycles from the Nucleus to

the Cell Cortex. J Cell Biol. 1995;131(1):165-78. doi: DOI 10.1083/jcb.131.1.165. PubMed PMID:

WOS:A1995RX83700014.

50.     Cinalli RM, Lehmann R. A spindle-independent cleavage pathway controls germ cell

formation in Drosophila. Nature cell biology. 2013;15(7):839-45. Epub 2013/06/04. doi:

10.1038/ncb2761. PubMed PMID: 23728423; PMCID: 3818562.

51.     Anand A, Kai T. The tudor domain protein kumo is required to assemble the nuage and

to generate germline piRNAs in Drosophila. EMBO J. 2012;31(4):870-82. Epub 2011/12/14. doi:

10.1038/emboj.2011.449. PubMed PMID: 22157814; PMCID: PMC3280549.

52.     Zhang Z, Koppetsch BS, Wang J, Tipping C, Weng Z, Theurkauf WE, Zamore PD. Antisense

piRNA amplification, but not piRNA production or nuage assembly, requires the Tudor-domain

protein Qin. EMBO J. 2014;33(6):536-9. Epub 2014/03/22. doi: 10.1002/embj.201384895.

PubMed PMID: 24652836; PMCID: PMC3989648.

53.     Anand A, Kai T. Response to Zhang et al. EMBO J. 2014;33(6):540-1. Epub 2014/03/22.

doi: 10.1002/embj.201387548. PubMed PMID: 24652837; PMCID: PMC3989649.

54.     Lecuyer E, Yoshida H, Parthasarathy N, Alm C, Babak T, Cerovina T, Hughes TR,

Tomancak P, Krause HM. Global analysis of mRNA localization reveals a prominent role in

organizing cellular architecture and function. Cell. 2007;131(1):174-87. doi:

10.1016/j.cell.2007.08.003. PubMed PMID: 17923096.

55.     Pan J, Goodheart M, Chuma S, Nakatsuji N, Page DC, Wang PJ. RNF17, a component of

the mammalian germ cell nuage, is essential for spermiogenesis. Development.

2005;132(18):4029-39. Epub 2005/08/12. doi: 10.1242/dev.02003. PubMed PMID: 16093322;

PMCID: PMC1994335.

56.     Watanabe T, Kankel, D. R. Molecular Cloning and Analysis olf( 1 )ogre, a Locus of

Drosophila melanogaster With Prominent Effects on the Postembryonic Development of the

Central Nervous System. Genetics. 1990;126:1033-44.

57.    van der Zee M, Benton MA, Vazquez-Faci T, Lamers GE, Jacobs CG, Rabouille C.

Innexin7a forms junctions that stabilize the basal membrane during cellularization of the

blastoderm in Tribolium castaneum. Development. 2015;142(12):2173-83. doi:

10.1242/dev.097113. PubMed PMID: 26015545.

58.    Olesnicky EC, Brent AE, Tonnes L, Walker M, Pultz MA, Leaf D, Desplan C. A caudal

mRNA gradient controls posterior development in the wasp Nasonia. Development.

2006;133(20):3973-82. Epub 2006/09/15. doi: 10.1242/dev.02576. PubMed PMID: 16971471.

59.    Ozuak O, Buchta T, Roth S, Lynch JA. Dorsoventral polarity of the Nasonia embryo

primarily relies on a BMP gradient formed without input from Toll. Curr Biol. 2014;24(20):2393-

8. Epub 2014/10/14. doi: 10.1016/j.cub.2014.08.035. PubMed PMID: 25308075.

60.    Riesgo-Escovar JR, Hafen E. Common and distinct roles of DFos and DJun during

Drosophila development. Science. 1997;278(5338):669-72. PubMed PMID: 9381174.

61.    Dorfman R, Glazer L, Weihe U, Wernet MF, Shilo BZ. Elbow and Noc define a family of

zinc finger proteins controlling morphogenesis of specific tracheal branches. Development.

2002;129(15):3585-96. PubMed PMID: 12117809.

62.    Wernet MF, Meier KM, Baumann-Klausener F, Dorfman R, Weihe U, Labhart T, Desplan

C. Genetic dissection of photoreceptor subtype specification by the Drosophila melanogaster

zinc finger proteins elbow and no ocelli. PLoS genetics. 2014;10(3):e1004210. Epub 2014/03/15.

doi: 10.1371/journal.pgen.1004210. PubMed PMID: 24625735; PMCID: PMC3953069.

63.    Rybakin V, Clemen CS. Coronin proteins as multifunctional regulators of the

cytoskeleton and membrane trafficking. Bioessays. 2005;27(6):625-32. doi: 10.1002/bies.20235.

PubMed PMID: 15892111.

64.    Gandhi M, Goode BL. Coronin: the double-edged sword of actin dynamics. Subcell

Biochem. 2008;48:72-87. Epub 2008/10/18. doi: 10.1007/978-0-387-09595-0_7. PubMed PMID:

18925372.

65.    Glater EE, Megeath LJ, Stowers RS, Schwarz TL. Axonal transport of mitochondria

requires milton to recruit kinesin heavy chain and is light chain independent. J Cell Biol.

2006;173(4):545-57. Epub 2006/05/24. doi: 10.1083/jcb.200601067. PubMed PMID: 16717129;

PMCID: PMC2063864.

66.    Tanaka T, Nakamura A. The endocytic pathway acts downstream of Oskar in Drosophila

germ plasm assembly. Development. 2008;135(6):1107-17. Epub 2008/02/15. doi:

10.1242/dev.017293. PubMed PMID: 18272590.

67.    Jaspers MH, Pflanz R, Riedel D, Kawelke S, Feussner I, Schuh R. The fatty acyl-CoA

reductase Waterproof mediates airway clearance in Drosophila. Dev Biol. 2014;385(1):23-31.

Epub 2013/11/05. doi: 10.1016/j.ydbio.2013.10.022. PubMed PMID: 24183938.

68.    Buchta T, Ozuak O, Stappert D, Roth S, Lynch JA. Patterning the dorsal-ventral axis of the

wasp Nasonia vitripennis. Dev Biol. 2013;381(1):189-202. doi: 10.1016/j.ydbio.2013.05.026.

PubMed PMID: 23735637.

69.    Ozuak O, Buchta T, Roth S, Lynch JA. Ancient and diverged TGF-beta signaling

components in Nasonia vitripennis. Dev Genes Evol. 2014;224(4-6):223-33. doi:

10.1007/s00427-014-0481-0. PubMed PMID: 25304164; PMCID: PMC4218986.

70.    Tomancak P, Berman BP, Beaton A, Weiszmann R, Kwan E, Hartenstein V, Celniker SE,

Rubin GM. Global analysis of patterns of gene expression during Drosophila embryogenesis.

Genome Biol. 2007;8(7):R145. doi: 10.1186/gb-2007-8-7-r145. PubMed PMID: 17645804;

PMCID: PMC2323238.

71.    Lee DM, Wilk R, Hu J, Krause HM, Harris TJ. Germ Cell Segregation from the Drosophila

Soma Is Controlled by an Inhibitory Threshold Set by the Arf-GEF Steppke. Genetics.

2015;200(3):863-72. Epub 2015/05/15. doi: 10.1534/genetics.115.176867. PubMed PMID:

25971667; PMCID: PMC4512548.

72.    Martinson EO, Mrinalini, Kelkar YD, Chang CH, Werren JH. The Evolution of Venom by

Co-option of Single-Copy Genes. Curr Biol. 2017;27(13):2007-13 e8. doi:

10.1016/j.cub.2017.05.032. PubMed PMID: 28648823; PMCID: PMC5719492.

73.    Hanyu-Nakamura K, Sonobe-Nojima H, Tanigawa A, Lasko P, Nakamura A. Drosophila

Pgc protein inhibits P-TEFb recruitment to chromatin in primordial germ cells. Nature.

2008;451(7179):730-3. Epub 2008/01/18. doi: 10.1038/nature06498. PubMed PMID: 18200011;

PMCID: PMC2719856.

74.    Nakamura A, Amikura R, Mukai M, Kobayashi S, Lasko PF. Requirement for a noncoding

RNA in Drosophila polar granules for germ cell establishment. Science. 1996;274(5295):2075-9.

Epub 1996/12/20. PubMed PMID: 8953037.

75.    Zhang F, Barboric M, Blackwell TK, Peterlin BM. A model of repression: CTD analogs and

PIE-1 inhibit transcriptional elongation by P-TEFb. Genes Dev. 2003;17(6):748-58. Epub

2003/03/26. doi: 10.1101/gad.1068203. PubMed PMID: 12651893; PMCID: PMC196018.

76.    Yamada T, Yamaguchi Y, Inukai N, Okamoto S, Mura T, Handa H. P-TEFb-mediated

phosphorylation of hSpt5 C-terminal repeats is critical for processive transcription elongation.

Mol Cell. 2006;21(2):227-37. doi: 10.1016/j.molcel.2005.11.024. PubMed PMID:

WOS:000234943000011.

77.    Lerit DA, Shebelut CW, Lawlor KJ, Rusan NM, Gavis ER, Schedl P, Deshpande G. Germ

Cell-less Promotes Centrosome Segregation to Induce Germ Cell Formation. Cell Rep.

2017;18(4):831-9. Epub 2017/01/26. doi: 10.1016/j.celrep.2016.12.074. PubMed PMID:

28122234; PMCID: PMC5327791.

78.    Amarnath S, Stevens LM, Stein DS. Reconstitution of Torso signaling in cultured cells

suggests a role for both Trunk and Torso-like in receptor activation. Development.

2017;144(4):677-86. doi: 10.1242/dev.146076. PubMed PMID: WOS:000394381100014.

79.    Guraya SS. Recent advances in the morphology, cytochemistry, and function of

Balbiani's vitelline body in animal oocytes. Int Rev Cytol. 1979;59:249-321. Epub 1979/01/01.

PubMed PMID: 385545.

80.    Kloc M, Bilinski S, Etkin LD. The Balbiani body and germ cell determinants: 150 years

later. Curr Top Dev Biol. 2004;59:1-36. Epub 2004/02/21. doi: 10.1016/S0070-2153(04)59001-4.

PubMed PMID: 14975245.

81.    Hurd TR, Herrmann B, Sauerwald J, Sanny J, Grosch M, Lehmann R. Long Oskar Controls

Mitochondrial Inheritance in Drosophila melanogaster. Dev Cell. 2016;39(5):560-71. Epub

2016/12/07. doi: 10.1016/j.devcel.2016.11.004. PubMed PMID: 27923120; PMCID:

PMC5147492.

82.    Cox RT, Spradling AC. Milton controls the early acquisition of mitochondria by

Drosophila oocytes. Development. 2006;133(17):3371-7. Epub 2006/08/05. doi:

10.1242/dev.02514. PubMed PMID: 16887820.

83.    Patil VS, Anand A, Chakrabarti A, Kai T. The Tudor domain protein Tapas, a homolog of

the vertebrate Tdrd7, functions in the piRNA pathway to regulate retrotransposons in germline

of Drosophila melanogaster. BMC Biol. 2014;12:61. doi: 10.1186/s12915-014-0061-9. PubMed

PMID: 25287931; PMCID: PMC4210518.

84.    Gao M, Thomson TC, Creed TM, Tu S, Loganathan SN, Jackson CA, McCluskey P, Lin Y,

Collier SE, Weng Z, Lasko P, Ohi MD, Arkov AL. Glycolytic enzymes localize to ribonucleoprotein

granules in <em>Drosophila</em> germ cells, bind Tudor and protect from transposable

elements. EMBO reports. 2015;16(3):379-86. doi: 10.15252/embr.201439694.

85.    Teng H, Sui X, Zhou C, Shen C, Yang Y, Zhang P, Guo X, Huo R. Fatty acid degradation

plays an essential role in proliferation of mouse female primordial germ cells via the p53-

dependent cell cycle regulation. Cell Cycle. 2016;15(3):425-31. doi:

10.1080/15384101.2015.1127473.

86.    Pao SS, Paulsen IT, Saier MH. Major facilitator superfamily. Microbiol Mol Biol R.

1998;62(1):1-+. PubMed PMID: WOS:000072489500001.

87.    Yamada Y, Davis KD, Coffman CR. Programmed cell death of primordial germ cells in

<em>Drosophila</em> is regulated by p53 and the Outsiders monocarboxylate transporter.

Development. 2008;135(2):207-16. doi: 10.1242/dev.010389.

88.    Jaspers MHJ, Pflanz R, Riedel D, Kawelke S, Feussner I, Schuh R. The fatty acyl-CoA

reductase Waterproof mediates airway clearance in Drosophila. Developmental Biology.

2014;385(1):23-31. doi: 10.1016/j.ydbio.2013.10.022. PubMed PMID: WOS:000328796100004.

89.    Hildebrandt A, Pflanz R, Behr M, Tarp T, Riedel D, Schuh R. Bark beetle controls

epithelial morphogenesis by septate junction maturation in Drosophila. Dev Biol.

2015;400(2):237-47. doi: 10.1016/j.ydbio.2015.02.008. PubMed PMID: WOS:000352466000007.

90.    Tanaka T, Kato Y, Matsuda K, Hanyu-Nakamura K, Nakamura A. Drosophila Mon2

couples Oskar-induced endocytosis with actin remodeling for cortical anchorage of the germ

plasm. Development. 2011;138(12):2523-32. Epub 2011/05/26. doi: 10.1242/dev.062208.

PubMed PMID: 21610029.

91.    Lerit DA, Gavis ER. Transport of germ plasm on astral microtubules directs germ cell

development in Drosophila. Curr Biol. 2011;21(6):439-48. doi: 10.1016/j.cub.2011.01.073.

PubMed PMID: 21376599; PMCID: PMC3062663.

92.    Ferree PM, McDonald K, Fasulo B, Sullivan W. The origin of centrosomes in

parthenogenetic hymenopteran insects. Curr Biol. 2006;16(8):801-7. doi:

10.1016/j.cub.2006.03.066. PubMed PMID: 16631588.

93.    van Oostende Triplet C, Jaramillo Garcia M, Haji Bik H, Beaudet D, Piekny A. Anillin

interacts with microtubules and is part of the astral pathway that defines cortical domains. J

Cell Sci. 2014;127(Pt 17):3699-710. doi: 10.1242/jcs.147504. PubMed PMID: 24994938.

94.    Tse YC, Piekny A, Glotzer M. Anillin promotes astral microtubule-directed cortical

myosin polarization. Mol Biol Cell. 2011;22(17):3165-75. doi: 10.1091/mbc.E11-05-0399.

PubMed PMID: 21737681; PMCID: PMC3164463.

95.    Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate

alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome

Biol. 2013;14(4):R36. doi: 10.1186/gb-2013-14-4-r36. PubMed PMID: 23618408; PMCID:

PMC4053844.

96.    Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL,

Pachter L. Differential gene and transcript expression analysis of RNA-seq experiments with

TopHat and Cufflinks. Nat Protoc. 2012;7(3):562-78. doi: 10.1038/nprot.2012.016. PubMed

PMID: 22383036; PMCID: PMC3334321.

97.    Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. StringTie

enables improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol.

2015;33(3):290-5. doi: 10.1038/nbt.3122. PubMed PMID: 25690850; PMCID: PMC4643835.

98.    Brent AE, Schweitzer R, Tabin CJ. A somitic compartment of tendon progenitors. Cell.

2003;113(2):235-48. Epub 2003/04/23. PubMed PMID: 12705871.

1133