Classification: Plant biology

Mechanical properties of the stigmatic cell wall mediate

pollen tube path in Arabidopsis

Lucie Riglet[1][§], Frédérique Rozier[1], Chie Kodera[1][£], Isabelle Fobis-Loisy[1] and Thierry Gaude[1][*]

[1]Laboratoire de Reproduction et Développement des Plantes, Université de Lyon, ENS de
Lyon, UCBL, INRA, CNRS, 46 Allée d'Italie, 69364 Lyon Cedex 07, France

* For correspondence : thierry.gaude@ens-lyon.fr

Present addresses :
[§]Sainsbury Laboratory, University of Cambridge, Bateman Street, Cambridge CB2 1LR, UK.
[£]Institut Jean-Pierre Bourgin, Institut National de la Recherche Agronomique (INRA),
AgroParisTech, CNRS, Université Paris-Saclay, 78000 Versailles, France.

Keywords: Pollen tube guidance, stigma papilla, KATANIN, mechanical anisotropy

ABSTRACT

Successful fertilization in angiosperms depends on the proper trajectory of pollen tubes through the pistil tissues to reach the ovules. Pollen tubes first grow within the cell wall of the papilla cells, applying pressure to the cell. Mechanical forces are known to play a major role in plant cell shape by controlling the orientation of cortical microtubules (CMTs), which in turn mediate deposition of cellulose microfibrils (CMFs). Here, by combining cell imaging and genetic approaches, we show that isotropic reorientation of CMTs and CMFs in aged and *katanin1-5* (*ktn1-5*) papilla cells is accompanied by a tendency of pollen tubes to coil around the papillae. Furthermore, we uncover that aged and *ktn1-5* papilla cells have a softer cell wall and provide less resistance to pollen tube growth. Our results reveal an unexpected role for KTN1 in pollen tube guidance by ensuring mechanical anisotropy of the papilla cell wall.

INTRODUCTION

Following deposition of dehydrated pollen grains on the receptive surface of the female organ, the stigma, pollen rehydrates, germinates and produces a pollen tube that carries the male gametes toward the ovules where the double fertilization takes place. This long itinerary through the different tissues of the pistil is finely controlled, avoiding misrouting of the pollen tube and hence assuring proper delivery of the sperm cells to the female gametes. In *Arabidopsis thaliana*, pollen tubes grow within the cell wall of papillae of the stigmatic epidermis, and then through the transmitting tissue of the style and ovary (Lennon and Lord, 2000). The transmitting tissue has an essential function in pollen tube guidance, providing chemical attractants and nutrients (Crawford and Yanofsky, 2008; Higashiyama and Hamamura, 2008). In contrast to these accumulating data showing the existence of factors mediating pollen tube growth in the pistil, whether guidance cues exist at the very early stage of pollen tube emergence and growth in the papilla cell wall remains largely unknown. The cell wall constitutes a stiff substrate and hence a mechanical barrier to pollen tube progression. There are numerous examples in animal cells demonstrating that mechanical properties of the cellular environment, and in particular rigidity, mediate cell signalling, proliferation, differentiation and migration (Discher et al., 2005; Ermis et al., 2018; Fu et al., 2010). In plant cells, cell wall rigidity depends mainly on its major component, cellulose, which is synthesized by plasma membrane-localized cellulose synthase complexes (CSCs) moving along cortical microtubule (CMT) tracks (Paredez et al., 2006). While penetrating the cell wall, the pollen tube exerts a pressure onto the stigmatic cell (Sanati Nezhad and Geitmann, 2013). Such physical forces are known to reorganise the cortical microtubules (CMTs), which by directing CSCs to the plasma membrane, reinforce wall stiffness by novel cellulose microfibril (CMF) synthesis (Paredez et al., 2006; Sampathkumar et al., 2014). Hence, there is an intricate interconnection between CMT organisation, CMF deposition and cell wall rigidity (Xiao et al.,

2016; Xiao and Anderson, 2016). A major regulatory element of CMT dynamics is the
KATANIN (KTN1) microtubule-severing enzyme, which allows CMT reorientation following
mechanical stimulation (Louveaux et al., 2016; Sampathkumar et al., 2014; Uyttewaal et al.,
2012). Here we investigated whether the CMT network of papilla cells might contribute to
pollen tube growth and guidance in stigmatic cells by combining cell imaging techniques,
genetic tools and atomic force microscopy. We show that isotropic reorientation of CMTs
occurs in aged Col-0 and *ktn1-5* papilla cells, which is accompanied by a change in the growth
direction of pollen tubes that tend to make coils on papillae. We demonstrate that both papilla
types exhibit softer cell walls that correlates with a faster growth rate of pollen tubes. Besides,
we show that CMT reorganisation is associated with isotropic rearrangement of CMFs. To
determine whether the coiled phenotype of pollen tubes could be a general feature of papillae
affected in cytoskeleton or cell wall composition, we tested a series of cell wall mutants
including the *xxt1 xxt2* double mutant that displays cytoskeleton orientation defects and
reduced stiffness of the cell wall. Remarkably, none used as female induces coiled growth of
wild-type pollen tubes. Altogether, our results indicate that cytoskeleton dynamics and
mechanical properties of the cell wall that both depend on KTN1 activity have a major role in
guiding early pollen tube growth in stigma papillae.

RESULTS

CMT dynamic pattern and pollen tube growth during stigma development. To assess the
functional role of stigmatic CMTs in pollen – papilla cell interaction, we first analysed their
organisation in papillae at stages 12 to 15 of stigma development as described (Smyth et al.,
1990) (Fig. 1*A* and *B*). We generated a transgenic line expressing the CMT marker MAP65.1-
citrine under the control of the stigma specific promoter SLR1 (Fobis-Loisy et al., 2007). Before
(stage 12) and at anthesis (stage 13), the CMTs were aligned perpendicularly to the
longitudinal axis of papilla cells and were highly anisotropic (median value of 0.40) (Fig. 1*C*
and *D*). At stage 14, when anthers extend above the stigma, the CMT pattern became less
organised, with a higher variability in anisotropy values. Finally, at stage 15 when the stigma
extends above anthers, CMT anisotropy had a median value of 0.09 indicative of an isotropic
orientation of CMTs (Fig. 1*C* and *D*). These findings reveal that the papilla CMT cytoskeleton
is dynamic during development, with a change of CMT array orientation from anisotropy to
isotropy. We then wondered whether this change in CMT organisation could be correlated with
pollen tube growth. To this end, we self-pollinated Col-0 papillae from stages 12 to 15 and
examined pollen tube growth one hour after pollination by scanning electron microscopy (SEM)
(Fig. 2*A*). At stage 12 and 13, we found most (~60%) pollen tubes to grow straight in the

papillae, whereas about 30% and 10% of tubes made half-turn or one turn around stigmatic

cells, respectively (Fig. 2*B*). At later stages of development, the tendency to coil around the

papillae dramatically increased, with more than 35% of pollen tubes at stage 14 and more than

55% at stage 15 making one or more than one turn around papillae. These results suggest

that CMT organisation in the papilla impacts the direction of growth of pollen tubes and that

loss of CMT anisotropy is associated with coiled growth.

Impaired CMT dynamics of papillae affects pollen tube growth direction. To confirm the

relation between stigma CMTs and pollen behaviour, we examined pollen tube growth on

stigmas of the *katanin1-5* mutant, which is known to exhibit reduced CMT array anisotropy in

root cells (Bichet et al., 2001; Burk et al., 2001; Burk and Ye, 2002). Because the CMT

organisation in *ktn1-5* papillae is unknown, we crossed *ktn1-5* with the MAP65-1-citrine marker

line and found that CMT arrays were more isotropic in *ktn1-5* papillae when compared with

those of the WT (Fig. 3*A* and *C*). Using SEM, we then analysed Col-0 pollen behaviour on

ktn1-5 stigmatic cells at stage 13 (Fig. 3*B*). We observed that Col-0 pollen tubes acquired a

strong tendency to coil around *ktn1-5* papillae, with above 60% of tubes making one or more

than one turn around papillae, sometimes making up to 6 turns, before reaching the base of

the cell (Fig. 3*D* and Fig. S1*A*). In some rare cases, pollen tubes even grew upward in the

ktn1-5 mutant and appeared blocked at the tip of the papilla (Fig. S1*B*). To test the direct

impact of stigma CMTs on pollen tube growth direction, we examined whether the

destabilization of CMTs in Col-0 papillae could affect pollen tube growth. To this end, we

treated stigmas by local application of the depolymerizing microtubule drug oryzalin in lanolin

pasted around the style. After 4 hours of drug treatment, no more CMT labelling was detected

in papillae, while CMTs were clearly visible in mock-treated (DMSO) stigmas (Fig. 4*A*). Stigmas

were then pollinated with Col-0 pollen and one hour later, pollen tubes were found turning on

drug-treated but not on control papillae (Fig. 4*B* and *C*). However, the number of coils was

significantly lower than on *ktn1-5* papillae. Indeed, 25% of the pollen tubes made at least 2

coils in *ktn1-5* papillae whereas this percentage represented only 4% on the oryzalin treated

stigmas (Fig. 3*D* and 4*C*). Altogether, these results confirm that stigmatic CMTs contribute to

the directional growth of pollen tubes in papilla cells.

Mechanical properties of the cell wall are disturbed in *ktn1-5* papilla cells. Because the

main role of CMTs in plant cells is to guide the trajectory of CSCs, thereby impacting the

mechanical anisotropy of the cell wall, we analysed the cell walls of Col-0 and *ktn1-5* papillae

following pollination. First, using Transmission Electron Microscopy (TEM), we found Col-0

pollen tubes to penetrate the cuticle and to grow between the two layers of the papilla cell wall,

as previously described (Kandasamy et al., 1994), in both Col-0 and *ktn1-5* papilla cells (Fig.

5*A*). We did not detect any significant difference in the ultrastructure of cell walls (Fig. S2). Interestingly, as the pollen tube progresses through the papilla cell wall, it generates a bump (external deformation) and an invagination (internal deformation) in the cell wall. As such deformations could reflect differences in wall properties, we quantified the external (extD) and internal (intD) deformation following pollination of Col-0 and *ktn1-5* stigmas. To visualize more clearly this deformation, we pollinated stigmas expressing the plasma membrane protein LTI6B fused to GFP (LTI6B-GFP) with pollen whose tube was labelled with the red fluorescent protein RFP driven by the ACT11 promoter (Rotman et al., 2005) (Fig. 5*B*). We found that Col-0 pollen tubes grew with almost equal extD and intD values in Col-0 cell wall. However, the ratio between extD and intD was about 3 when Col-0 pollen tubes grew in *ktn1-5* papilla cells (Fig. 5*B-D*). These quantitative data were consistent with the observation that pollen tubes appeared more prominent on *ktn1-5* stigmatic cells using SEM (Fig. 3*B* and Fig. S1). Assuming that stigmatic cells are pressurized by their turgor pressure, this may reflect the presence of softer walls in *ktn1-5* papillae. To test this hypothesis, we assessed the stiffness of Col-0 and *ktn1-5* papilla cell walls using Atomic Force Microscopy (AFM) with a 200 nm indentation. We found that cell wall stiffness in *ktn1*-5 papilla cells was about 30% lower than that in stage 13 WT cells (Fig. 5*E* and *F*). To confirm this result, we investigated the stiffness of the papilla cell wall on WT stigmas at stage 15, where increased coiled pollen tubes were detected (Fig. 2*A* and *B*). We found the cell wall to be softer than that of papillae at stage 13 but stiffer than that of the *ktn1-5* (Fig. 5*E* and *F*). We then reasoned that the presence of softer walls should also affect the pollen tube growth rate, stiffer walls reducing growth. Thus, measuring the growth rate would provide an indication of the resistance force encountered by the tube while growing in the papilla cell wall. We thus monitored the growth rate of Col-0 pollen tubes in Col-0 and *ktn1-5* papillae. We found that pollen tubes grew faster (~ x 1.8) within *ktn1-5* papillae (Fig. 5*G*). Similarly, we found that pollen tube growth on stage 15 papillae was faster (~ x 1.6) than on stage 13 papillae (Fig. 5*H*). These results suggest that the coiled phenotype is associated with a higher pollen tube growth rate and hence, that papilla cell walls of *ktn1-5* and stage-15 papillae exhibit less resistance to tube penetration. Altogether, these data suggest that KATANIN-dependent mechanical properties of papilla cell wall play a role in pollen tube guidance.

Defects in papilla cell-wall composition are not sufficient to affect Col-0 pollen tube behaviour. The *katanin1/fra2* mutant was initially described as a mutant impaired in cell wall biosynthesis and CMT array organisation (Burk et al., 2001). This prompted us to investigate whether other mutants affected in cell wall biogenesis might exhibit the coiled pollen tube phenotype. Indeed, in the simplest scenario, the presence of softer walls, whatever the cause, may be sufficient to induce extra coiling due to faster pollen tube growth. We selected mutants

impaired in the cellulose synthase complex (*kor1.1*, *prc1* and *any1*), hemicellulose

biosynthesis (*xxt1 xxt2*, *xyl1.4*) and pectin content (*qua2.1*) (Table S1). Strikingly, none of the

6 cell wall mutants displayed the coiled pollen tube phenotype (Fig. S3). This suggests that

the relation between CMT organisation, cell wall stiffness and pollen tube trajectory is stricter

than anticipated. Because CMTs guide cellulose deposition, they also control the directional

elongation of plant cells (Baskin, 2005). Indeed, *katanin1/fra2* cells are wider and shorter than

wild type (16, 18), and we may predict that the shape of papilla cells in the mutant might be

similarly altered. The contribution of stigmatic CMTs to pollen tube growth may thus be

mediated by papilla cell shape only; for instance, wider cells in *ktn1-5* would promote pollen

tube coiling. To check that possibility, we measured the length and width of papilla cells in Col-

0 (at stage 13 and 15), *ktn1-5, xxt1 xxt2* and *any1* (Fig. S4). We did not find any correlation

between papilla length and the coiled phenotype. Indeed, coiled phenotype was observed in

stage-15 WT papillae that were longer than those at stage-13, as well as in *ktn1-5* papillae that

had a length similar to stage-13 WT papillae. The correlation between papilla width and coiled

phenotype was also not clear-cut. As anticipated, *ktn1-5* mutant exhibited wider papillae than

Col-0. Stage-15 WT papillae were also significantly larger than stage-13 WT papillae, where

coils were observed. However, *xxt1 xxt2* and *any1* papillae were also wider than the WT but

did not display the coiled phenotype. Altogether, this suggests that papilla morphology is not

sufficient to explain the coiled phenotype. Because CMT disorganisation in papillae affects

both wall stiffness and mechanical anisotropy, we next investigated the relative contribution of

these two parameters in pollen tube growth. We focused our analysis on the *xxt1 xxt2* double

mutant. Indeed, this mutant was reported to display CMT orientation defects and reduced

stiffness of the cell wall in hypocotyl cells, when compared with the WT (Xiao et al., 2016). As

the pollen tube phenotype on *xxt1 xxt2* stigmas was similar to the WT, we wondered whether

the cell wall stiffness of stigmatic cells was actually affected in these mutant papillae. Using

AFM, we found the papilla cell wall of the mutant to be about 30% softer than that of Col-0

papillae, i.e. very similar to that of *ktn1-5* (Fig. 5*F*). However, despite this similarity, the pollen

tube growth rate in *xxt1 xxt2* stigmas was identical to Col-0 stigmas at stage 13 (Fig. 5*H*). In

addition, contrary to *ktn1-5*, pollen tubes were not prominent while growing in *xxt1 xxt2* papilla

cell walls (Fig. S3*G*). These results suggested that although *ktn1-5* and *xxt1 xxt2* papillae had

similar cell wall stiffness as inferred from AFM measurements, they exhibited different

mechanical constraints to pollen tube growth. Because of the close relationship between CMT

organisation, CMF deposition and cell wall rigidity (Xiao et al., 2016; Xiao and Anderson,

2016), we suspected that CMT and CMF organisation might be different between the two

mutants and be the possible causal agent of stiffness differences and the related tube coiled

phenotype observed in *ktn1-5*. To check this possibility, and to gain insight into the

predominant cellulose pattern in the stigmatic cell walls, we stained CMFs by using Direct Red

23. We found cellulose fibres to be highly aligned and slanted to the longitudinal axis of the
papilla in stage 13 Col-0 stigmas, whereas *ktn1-5* and stage 15 Col-0 papillae both displayed
a clearly disordered CMF pattern (Fig. 6a) with CMFs forming thicker and more spaced
bundles (Fig. 6b). By contrast, CMFs in *xxt1 xxt2* papillae organized in a dense and well
oriented pattern of fibres, mostly perpendicular to the papilla long axis, which resembled Col-
0 stage 13 CMF organisation (Fig. 6a and b). These results suggest that KTN1 loss-of-function
alters mechanical properties of the papilla cell wall in a complex manner, lowering both its
stiffness to pressure applied perpendicularly to its surface (e.g., by the AFM indenter) and its
resistance to pollen tube penetration. This latter modification appears as the main cause of the
pollen tube coiled phenotype. By contrast, loss of XXT1 and XXT2 functions, by perturbing cell
wall stiffness to indenter pressure only, does not induce change in pollen tube growth.
Altogether, our data suggest that the mechanical anisotropy of papilla cell walls plays a key
role in pollen tube trajectory.

DISCUSSION

How the pollen germinates a tube and how pollen tube growth is regulated have been the
object of many investigations (Mizuta and Higashiyama, 2018; Palanivelu and Tsukamoto,
2012). The use of in vitro pollen germination as well as semi-in vivo fertilization assays,
together with the analysis of mutants defective in pollen or ovule functions, have rapidly
expanded our knowledge of the mechanisms that sustain pollen tube growth, its guidance
toward the ovule and the final delivery of male gametes within the embryo sac (Cameron and
Geitmann, 2018; Dresselhaus and Franklin-Tong, 2013; Higashiyama and Yang, 2017). At the
cellular level, cytoskeleton has been extensively studied during pollen tube elongation (Fu,
2015), highlighting the critical role played by the actin microfilaments in pollen-tube tip growth
through delivery of materials for the biosynthesis of the plasma membrane and cell wall. By
contrast, CMTs seem to have a lower importance in this process, drugs affecting CMT
polymerisation having no significant effects on pollen-tube growth rate, yet altering the capacity
of pollen tubes to change their growth direction (Gossot and Geitmann, 2007).
Rearrangements of actin microfilaments (Iwano et al., 2007) and destabilization of CMTs
(Samuel et al., 2011) have been described in stigmas following compatible pollination in
Brassica species. However, CMT pattern and dynamics during papilla development have
never been reported. Our data show that during the course of stigma maturation, which is
associated with papilla cell elongation (Fig. S4*A* and *B*), CMT bundles progressively move
from perpendicular (anisotropic) to the elongation axis at stage 12 to disorganised (isotropic)
at stage 15. This correlates with the known CMT dynamics during elongation in plant cells,
where CMT arrays are highly anisotropic in young cells and become more isotropic as cells
differentiate (Baskin, 2005; Landrein and Hamant, 2013). The progressive randomisation of
CMT orientation we observed in papilla cells is accompanied by an increased coiled growth of
pollen tubes in papillae. Similarly, when CMTs are destabilized by the microtubule
depolymerizing drug oryzalin, coiled pollen tubes are more frequently observed compared with
untreated control stigmas. These results reveal a link between the stigmatic CMT cytoskeleton
organisation and the trajectory that pollen tube takes while growing in the papilla cell wall. The
fact that the most striking effect on pollen tube growth was found on *ktn1-5* mutant suggests
that the coiled phenotype depends not only on the CMT organisation but implicates other
factors. Among various phenotypic alterations described in loss-of-function mutants for *KTN1*,
are the impaired cell mechanical properties and cell elongation, defects in CMT organisation,
cell wall composition and CMF orientation (Burk et al., 2001; Burk and Ye, 2002; Ryden et al.,
2003). Interestingly, we found that the protuberance of pollen tubes at the surface of *ktn1-5*
papillae was associated with a faster growth rate and a lower rigidity of the cell wall compared
with Col-0. This places mechanics of the cell wall as a likely component involved in the coiled

phenotype. Surprisingly, of the six cell wall mutants we analysed, including *xxt1 xxt2* and *prc1*
known to exhibit both abnormal CMT organisation and softer cell walls, none induced the coiled
phenotype displayed by *ktn1-5* papillae. More remarkably, despite the similar stiffness of *xxt1*
xxt2 and *ktn1-5* papilla cell walls measured in our AFM experiments, pollen tubes behaved
differently on these two stigmas. This indicates that components other than cell wall stiffness
are involved in the pollen tube coiled phenotype and that mechanical properties of the cell wall
are not identical in the two mutants. Indeed, though sharing a similar alteration of cell wall
stiffness, the global plant morphology of these mutants is clearly distinct. The *ktn1-5* mutant
was described to have a severe reduction in cell length and an increase in cell width in all
organs (Burk et al., 2001). This cellular phenotype was attributed to the distorted deposition of
CMFs correlated with the isotropic orientation of CMTs, whereas in WT cells, CMFs like CMTs
are oriented perpendicularly to the elongation axis (Burk and Ye, 2002). Similarly, we found
that orientation of CMFs is also altered in *ktn1-5* papillae (Fig. 6) and this was associated with
larger papilla cells (Fig. S4*C* and *D*). Hence, we may assume that mechanical anisotropy,
described as the cell wall anisotropy made by the orientation of the rigid CMFs (Sassi et al.,
2014), is impaired in *ktn1-5* papillae. Contrary to the *ktn1-5* mutant, papillae of the *xxt1 xxt2*
double mutant display an anisotropic CMF pattern with parallel fibres approximatively oriented
transversally to the papilla axis. These data are in agreement with those reported for etiolated
hypocotyl cells, where CMFs are largely parallel to one another, straighter than in WT and
oriented almost transversely to the long axis of the cell (Xiao et al., 2016). In addition, we found
the papilla cell shape of *xxt1 xxt2* stigmas to be similar to Col-0, though with slightly wider
papillae (Fig. S3*G* and Fig. S4*E* and *F*). However, *xxt1 xxt2* papillae are less wide than *ktn1-5*
papillae, which is consistent with an anisotropic growth of the papillae in the *xxt1 xxt2* mutant
compared with *ktn1-5*. Apart from the similar stiffness of papilla cell walls as deduced from
AFM measurements, the main differences between *ktn1-5* and *xxt1 xxt2* cell walls are likely to
be the cell wall composition, orientation of CMFs and possible changes in molecular
connections between cell wall components and plasma membrane and/or cytoskeleton
proteins. Indeed, a recent proteomic study revealed that loss of KATANIN function is
associated with the decrease in abundance of several cytoskeleton proteins, such as profilin
1, actin-depolymerizing factor 3 and actin 7 (Takáč et al., 2017), whereas targeted quantitative
RT-PCR unveiled that several Microtubule-associated protein (MAP) and wall signal receptor
genes are downregulated in *xxt1 xxt2* (Xiao et al., 2016). In this latter study, KTN1 expression
level was shown to be unchanged compared with Col-0. Altogether, our study suggests that
KTN1, by maintaining the papilla mechanical anisotropy, has a key function in mediating early
pollen tube guidance on stigma papillae.
The coiled phenotype was not only observed in *ktn1-5* but also in the WT Col-0 papillae at
stage 15. Remarkably, several lines of evidence reveal that papilla cells from these two

genotypes share common features, such as CMT and CMF increased isotropy, decreased stiffness of the cell wall and less resistance of the wall to pollen tube growth. Isotropic orientation of the cytoskeleton at stage 15 is likely to relate to cell elongation, which is known to be accompanied by cytoskeleton reorganisation (Crowell et al., 2011; Zhang et al., 2014). At the organ level, it has been suggested that the mechanical anisotropy of the wall restrains organ emergence (Sassi et al., 2014). The authors propose that for the same wall stiffness, a cell wall with isotropic properties would lead to larger outgrowth than a wall with anisotropic properties. Our data are consistent with this hypothesis, albeit at the subcellular scale, since large protuberance of papilla wall following pollen tube growth is observed in *ktn1-5* papilla cells, exhibiting walls with isotropic properties.

It remains unclear how mechanical anisotropy guides pollen tube growth. We can hypothesise that as pollen tube grows inside the wall, it encounters recently deposited CMFs on the inner side of the wall (facing the cytoplasm) and older CMFs on the outer side. It is likely that these layers have different mechanical properties related to CMF orientation (Baskin, 2005). Pollen tubes may grow helically by default, as is the case for climbing plants around a cylindrical substrate but the presence of a mechanically reinforced inner wall may slow down and bias the trajectory of the pollen tube. It is worth noting that growth rate is slowed down when pollen tubes pass through a microgap of a microfluiding device, the tubes adapting their invasive force to the mechanical constraints (Nezhad et al., 2013). Based on measurements of pollen tube growth rates, our data show that the tube tip, while progressing in the papilla wall, senses the mechanical features of its environment and reacts accordingly. Hence, it reveals some unanticipated internal and hidden properties of the cell wall. Interestingly, recent work showed that axons were capable of adapting their growth rate and direction according to mechanical constraints, growing straighter on rigid substrate, underlining that mechanical signals are important regulators of pathfinding (Koser et al., 2016).

In the last decades, many studies pointed out that chemical but also mechanical components must be considered to be implicated in pollen tube growth direction (Cameron and Geitmann, 2018). Implication of chemical cues for pollen tube guidance in the stigma remains largely unknown, although the blue copper protein plantacyanin, when overexpressed in Arabidopsis stigmas, was reported to be a possible guidance factor through an as yet undiscovered mechanism (Dong et al., 2005). However, no defect in pollen tube growth directionality was detected in a knock-down plantacyanin mutant, questioning the actual role of this protein as a chemoattractant in Arabidopsis. In our study, we add mechanics as a key player in early pollen tube guidance in the papilla cell. Our results suggest that this role is mediated by a specific CMT/CMF organisation and mechanical anisotropy of the papilla cell, which both are dependent on KTN1. Importantly, these specific mechanical properties of the stigmatic cells prevent emerging pollen tubes to grow upward on papillae and straighten pollen

tube direction, helping the tube to find its correct path to the stylar transmitting tract. We uncovered a yet unexpected role in pollen tube guidance for KTN1, which was until now mostly known to be involved in plant development and stress-response regulation. In addition, our study also clearly unveils that the mechanical properties of one single cell (e.g., the stigmatic papilla) impact the behaviour of its neighbouring cell (e.g., the pollen tube).

MATERIALS AND METHODS

Plant Materials and Growth Conditions. *Arabidopsis thaliana*, ecotype Columbia (Col-0), *Arabidopsis* transgenic plants generated in this study and *Arabidopsis* mutants were grown in soil under long-day conditions (16 hours of light / 8 hours of dark, 21°C / 19°C) with a relative humidity around 60%. *ktn1-5* (SAIL_343_D12), *xxt1xxt2*, *prc1.1, qua2.1*, *xyl1.4*, *kor1.1* and *any1* mutant lines were described previously (Cavalier et al., 2008; Fagard et al., 2000; Fujita et al., 2013; Lin et al., 2013; Mouille et al., 2007; Nicol et al., 1998; Sampedro et al., 2010; Shoji et al., 2004). All mutants were in Col-0 background except *kor1.1* which was in WS.

Plasmid construction. We used the Gateway® technology (Life Technologies, USA) and two sets of Gateway®-compatible binary T-DNA destination vectors (Hellens et al., 2000; Karimi et al., 2002) for expression of transgenes in *A. thaliana*. The DNA fragment containing the *Brassica oleracea SLR1* promoter was inserted into the pDONP4-P1R vector. The 165 bp-*LTI6B* fragment was introduced into the pDONR207 vector. *MAP65* gene spanning the coding region from start to stop codons was introduced into the pDONR221 vector. CDS from citrine or GFP were cloned into the pDONP2R-P3 vector. Final constructs, pSLR1::MAP65-citrine and pSLR1::LTI6B-GFP were obtained by a three-fragment recombination system (Life Technologies) using the pK7m34GW and the pB7m34GW destination vectors, respectively. We generated a pAct11::RFP construct by amplifying the promoter of the *A. thaliana Actin 11* gene and cloning it into the pGreenII gateway vector in front of the RFP coding sequence.

Generation of transgenic lines and crossing. Transgenic lines were generated by *Agrobacterium tumefaciens*-mediated transformation of *A. thaliana* Col-0 as described (Logemann et al., 2006). Unique insertion lines, homozygous for the transgene were selected. We introduced the pSLR1::LTI6B-GFP or pSLR1::MAP65-citrine construct in *ktn1-5* background by crossing and further selecting the progeny on antibiotic containing medium.

Microscopy
Confocal microscopy. Flowers at stages 12 to 15 (Smyth et al., 1990) collected from fluorescent lines were emasculated and stigmas were observed under a Zeiss LSM800

366 microscope (AxioObserver Z1) using a 40x Plan-Apochromat objective (numerical aperture
367 1.3, oil immersion). Citrine was excited at 515 nm and fluorescence detected between 530 and
368 560nm. GFP was excited at 488 nm and fluorescent detected between 500 and 550 nm. RFP
369 was excited at 561 nm and fluorescent detected between 600 and 650 nm.

370 ***Live imaging.*** Flowers from stages 12 to 15 were emasculated and pollinated on plants with
371 mature pollen from the pACT11::RFP line. Immediately after pollination, stigmas were mounted
372 between two coverslips. To maintain a constant humidity without adding liquid directly on the
373 stigma surface, we use a wet piece of tissue in contact with the base of the stigma. Pollinated
374 stigmas were observed under a Zeiss microscope (AxioObserver Z1) equipped with a spinning
375 disk module (CSU-W1-T3, Yokogawa) using a 40x Plan-Apochromat objective (numerical
376 aperture 1.1, water immersion). Serial confocal images were acquired in the entire volume of
377 the stigma every 1 µm and every minute. Images were processed with Image J software and
378 pollen tube lengths were measured.

379 ***Atomic Force Microscopy.*** Pistils at stage 13 were placed straight in a 2% agar MS medium
380 and 0.8% low-melting agarose was added up to the base of papilla cells. AFM indentation
381 experiments were carried out with a Catalyst Bioscope (Bruker Nano Surface, Santa Barbara,
382 CA, USA) that was mounted on an optical microscope (MacroFluo, Leica, Germany) equipped
383 with a x10 objective. All quantitative measurements were performed using standard pyramidal
384 tips (RFESP-190 (Bruker)). The tip radius given by the manufacturer was 8-12 nm. The spring
385 constant of the cantilever was measured using the thermal tune method and was 35 N/m. The
386 deflection sensitivity of the cantilever was calibrated against a sapphire wafer. All experiments
387 were made in ambient air at room temperature. Matrix of 10x10 measurements (step 500 nm)
388 was obtained for each papilla, with a 1µN force. The Young's Modulus was estimated using
389 the Nanoscope Analysis (Bruker) software, using the Sneddon model with a < 200nm
390 indentation.

392 ***Environmental Scanning Electron Microscopy (SEM).*** Flowers from stages 12 to 15 were
393 emasculated and pollinated on plants with mature WT pollen. One hour after pollination, pistils
394 were cut in the middle of the ovary, deposited on a SEM platform and observed under Hirox
395 SEM SH-3000 at -20°C, with an accelerating voltage of 15kV. Images were processed with
396 ImageJ software and pollen tube direction was quantified by counting the number of turns
397 made by the tube, only on papillae that received one unique pollen grain.

398 ***Transmission Electron Microscopy.*** Stage 13 flowers were emasculated and pollinated on
399 plants with mature WT pollen. One hour after pollination, pistils were immersed in fixative
400 solution containing 2.5% glutaraldehyde and 2.5% paraformaldehyde in 0.1 M phosphate
401 buffer (pH 7.2) and after 4 rounds of 30 min vacuum, they were incubated in fixative for 12

hours at room temperature. Pistils were then washed in phosphate buffer and further fixed in
1% osmium tetroxide in 0.1 M phosphate buffer (pH 7.2) for 1.5 hours at room temperature.
After rinsing in phosphate buffer and distilled water, samples were dehydrated through an
ethanol series, impregnated in increasing concentrations of SPURR resin over a period of 3
days before being polymerized at 70°C for 18 h, sectioned (65 nm sections) and imaged at 80
kV using an FEI TEM tecnaiSpirit with 4 k x 4 k eagle CCD.

Anisotropy estimation. Flowers from MAP65 lines were emasculated and stigmas were
observed under confocal microscope. Images were processed with ImageJ software and
quantitative analyses of the average orientation and anisotropy of CMTs were performed using
FibrilTool, an ImageJ plug-in (Boudaoud et al., 2014). Anisotropy values range from 0 to 1; 0
indicates pure isotropy, and 1 pure anisotropy.

Cellulose staining. A droplet of Adigor™ was applied at 2.5% (v/v in water) on the stigma
during 4 hours to facilitate subsequent chemical treatments. As described for hypocotyls
(Landrein et al., 2013), pistils were incubated in 12.5% glacial acetic acid for 1 hour and
dehydrated in 100% and then 50% ethanol for 20 min each. Pistils were then washed in water
during 20 min and stored in 1M KOH during 2 days. Pistils were then stained using 0.02%
(w/v) Direct Red 23 dye (Sigma-Aldrich) during 4 hours and washed with distilled water. Pistils
were observed with a mRFP filter (excited at 561 nm) under a Zeiss LSM 880 confocal
microscope using a 63x Plan-Apochromat objective (numerical aperture 1.4, oil immersion).
Stigmas were imaged by taking a z-stack of 0.2 μm sections in the papilla cells. Image analysis
was performed by doing a 3D-projection and measuring the fluorescence intensity
longitudinally along the papilla axis using ImageJ software.

Membrane deformation estimation. Flowers from LTI6B lines were emasculated and
pollinated with mature pollen from the PACT11::RFP line. 20 minutes after pollination, stigmas
were observed under confocal microscope. Serial confocal images every 1 μm encompassing
the entire volume of the stigma were recorded and processed with ImageJ software. Plasma
membrane deformation was estimated by choosing the slide from the stack that corresponded
to the focus plan of the contact site with the RFP-labelled pollen tube. On the Bright field image
corresponding to the selected slide, a line was drawn connecting the two ridges of the
invagination of the papilla. Two perpendicular lines, one toward the exterior (ExtD) of the
papilla to the maximum point of deformation, and the other toward the interior (IntD) on the
GFP image were measured, respectively.

Oryzalin treatment. To avoid contact of pollen grains with liquid, we performed local
applications of oryzalin (Chemical service, Supelco) in lanolin pasted (Sassi et al., 2014),

around the style, just under the stigmatic cells, at 833 µg/mL (DMSO), for 4 hours at 21°C.

Oryzalin-treated pistils were pollinated with mature WT pollen and 1 hour after pollination

observed under SEM.

Statistical analysis. Graph and statistics were obtained with R software or Excel. Statistical

tests performed are specified in figure legends.

ACKNOWLEDGEMENTS

We thank O. Hamant for critical reading of the manuscript and fruitful discussion, A. Boudaoud,

S. Bovio and V. Battu for advice on AFM experiments, and the Sice and MechanoDevo team

members for discussion, P. Bolland, A. Lacroix and J. Berger for plant care and the PLATIM

imaging facility of the SFR Biosciences Gerland-Lyon Sud. We thank the Bordeaux Imaging

Centre especially L. Brocard and B. Batailler for TEM microscopy. We also thank the

BioMeca® society and Pascale Milani for the AFM measurements. L.R. was funded by a

fellowship from the French Ministry of Higher Education and Research. The work was

supported by Grant ANR-14-CE11-0021.

AUTHOR CONTRIBUTIONS

L.R. performed all the experiments and the image analysis, except TEM and AFM, which were

subcontracted. F.R. and C.K. contributed to cell imaging setup. L.R., T.G., and I.F.L., designed

the study and analysed the data. L.R. and T.G. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

Abasolo W, Eder M, Yamauchi K, Obel N, Reinecke A, Neumetzler L, Dunlop JWC, Mouille G, Pauly M, Höfte H, Burgert I. 2009. Pectin May Hinder the Unfolding of Xyloglucan Chains during Cell Deformation: Implications of the Mechanical Performance of Arabidopsis Hypocotyls with Pectin Alterations. *Molecular Plant* **2**:990–999. doi:10.1093/mp/ssp065

Baskin TI. 2005. Anisotropic Expansion of the Plant Cell Wall. *Annual Review of Cell and Developmental Biology* **21**:203–222. doi:10.1146/annurev.cellbio.20.082503.103053

Bichet A, Desnos T, Turner S, Grandjean O, Höfte H. 2001. BOTERO1 is required for normal orientation of cortical microtubules and anisotropic cell expansion in Arabidopsis. *The Plant Journal* **25**:137–148. doi:10.1111/j.1365-313X.2001.00946.x

Boudaoud A, Burian A, Borowska-Wykręt D, Uyttewaal M, Wrzalik R, Kwiatkowska D, Hamant O. 2014. FibrilTool, an ImageJ plug-in to quantify fibrillar structures in raw microscopy images. *Nature Protocols* **9**:457–463. doi:10.1038/nprot.2014.024

Burk DH, Liu B, Zhong R, Morrison WH, Ye Z-H. 2001. A Katanin-like Protein Regulates Normal Cell Wall Biosynthesis and Cell Elongation. *Plant Cell* **13**:807–828.

Burk DH, Ye Z-H. 2002. Alteration of Oriented Deposition of Cellulose Microfibrils by Mutation of a Katanin-Like Microtubule-Severing Protein. *Plant Cell* **14**:2145–2160. doi:10.1105/tpc.003947

Cameron C, Geitmann A. 2018. Cell mechanics of pollen tube growth. *Current Opinion in Genetics & Development*, Developmental mechanisms, patterning and evolution **51**:11–17. doi:10.1016/j.gde.2018.03.008

Cavalier DM, Lerouxel O, Neumetzler L, Yamauchi K, Reinecke A, Freshour G, Zabotina OA, Hahn MG, Burgert I, Pauly M, Raikhel NV, Keegstra K. 2008. Disrupting Two Arabidopsis thaliana Xylosyltransferase Genes Results in Plants Deficient in Xyloglucan, a Major Primary Cell Wall Component. *The Plant Cell* **20**:1519–1537. doi:10.1105/tpc.108.059873

Crawford BCW, Yanofsky MF. 2008. The Formation and Function of the Female Reproductive Tract in Flowering Plants. *Current Biology* **18**:R972–R978. doi:10.1016/j.cub.2008.08.010

Crowell EF, Timpano H, Desprez T, Franssen-Verheijen T, Emons A-M, Höfte H, Vernhettes S. 2011. Differential Regulation of Cellulose Orientation at the Inner and Outer Face of

Epidermal Cells in the Arabidopsis Hypocotyl. *The Plant Cell* **23**:2592–2605.
doi:10.1105/tpc.111.087338

Discher DE, Janmey P, Wang Y. 2005. Tissue Cells Feel and Respond to the Stiffness of Their
Substrate. *Science* **310**:1139–1143. doi:10.1126/science.1116995

Dong J, Kim ST, Lord EM. 2005. Plantacyanin Plays a Role in Reproduction in Arabidopsis.
Plant Physiology **138**:778–789. doi:10.1104/pp.105.063388

Dresselhaus T, Franklin-Tong N. 2013. Male–Female Crosstalk during Pollen Germination,
Tube Growth and Guidance, and Double Fertilization. *Molecular Plant* **6**:1018–1036.
doi:10.1093/mp/sst061

Ermis M, Antmen E, Hasirci V. 2018. Micro and Nanofabrication methods to control cell-
substrate interactions and cell behavior: A review from the tissue engineering
perspective. *Bioactive Materials* **3**:355–369. doi:10.1016/j.bioactmat.2018.05.005

Fagard M, Desnos T, Desprez T, Goubet F, Refregier G, Mouille G, McCann M, Rayon C,
Vernhettes S, Höfte H. 2000. PROCUSTE1 Encodes a Cellulose Synthase Required
for Normal Cell Elongation Specifically in Roots and Dark-Grown Hypocotyls of
Arabidopsis. *Plant Cell* **12**:2409–2423.

Fobis-Loisy I, Chambrier P, Gaude T. 2007. Genetic transformation of Arabidopsis lyrata:
specific expression of the green fluorescent protein (GFP) in pistil tissues. *Plant Cell*
Rep **26**:745–753. doi:10.1007/s00299-006-0282-7

Fu J, Wang Y-K, Yang MT, Desai RA, Yu X, Liu Z, Chen CS. 2010. Mechanical regulation of
cell function with geometrically modulated elastomeric substrates. *Nat Methods* **7**:733–
736. doi:10.1038/nmeth.1487

Fu Y. 2015. The cytoskeleton in the pollen tube. *Current Opinion in Plant Biology*, Cell biology
28:111–119. doi:10.1016/j.pbi.2015.10.004

Fujita M, Himmelspach R, Ward J, Whittington A, Hasenbein N, Liu C, Truong TT, Galway ME,
Mansfield SD, Hocart CH, Wasteneys GO. 2013. The anisotropy1 D604N mutation in
the Arabidopsis cellulose synthase1 catalytic domain reduces cell wall crystallinity and
the velocity of cellulose synthase complexes. *Plant Physiol* **162**:74–85.
doi:10.1104/pp.112.211565

Gossot O, Geitmann A. 2007. Pollen tube growth: coping with mechanical obstacles involves
the cytoskeleton. *Planta* **226**:405–416. doi:10.1007/s00425-007-0491-5

Hellens RP, Edwards EA, Leyland NR, Bean S, Mullineaux PM. 2000. pGreen: a versatile and
flexible binary Ti vector for Agrobacterium-mediated plant transformation. *Plant*
Molecular Biology **42**:819–832. doi:10.1023/A:1006496308160

Higashiyama T, Hamamura Y. 2008. Gametophytic pollen tube guidance. *Sex Plant Reprod*
21:17–26. doi:10.1007/s00497-007-0064-6

Higashiyama T, Yang W. 2017. Gametophytic Pollen Tube Guidance: Attractant Peptides,
Gametic Controls, and Receptors. *Plant Physiol* **173**:112–121.
doi:10.1104/pp.16.01571

His I, Driouich A. 2001. Altered pectin composition in primary cell walls of korrigan, a dwarf
mutant of Arabidopsis defcient in a membrane-bound endo-1,4-b-glucanase 11.

Iwano M, Shiba H, Matoba K, Miwa T, Funato M, Entani T, Nakayama P, Shimosato H,
Takaoka A, Isogai A, Takayama S. 2007. Actin Dynamics in Papilla Cells of Brassica
rapa during Self- and Cross-Pollination. *Plant Physiol* **144**:72–81.
doi:10.1104/pp.106.095273

Kandasamy MK, Nasrallah JB, Nasrallah ME. 1994. Pollen-pistil interactions and
developmental regulation of pollen tube growth in Arabidopsis. *Development*
120:3405–3418.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY™ vectors for Agrobacterium-mediated plant
transformation. *Trends in Plant Science* **7**:193–195. doi:10.1016/S1360-
1385(02)02251-3

Koser DE, Thompson AJ, Foster SK, Dwivedy A, Pillai EK, Sheridan GK, Svoboda H, Viana
M, da F. Costa L, Guck J, Holt CE, Franze K. 2016. Mechanosensing is critical for axon
growth in the developing brain. *Nat Neurosci* **19**:1592–1598. doi:10.1038/nn.4394

Landrein B, Hamant O. 2013. How mechanical stress controls microtubule behavior and
morphogenesis in plants: history, experiments and revisited theories. *The Plant Journal*
75:324–338. doi:10.1111/tpj.12188

Lei L, Zhang T, Strasser R, Lee CM, Gonneau M, Mach L, Vernhettes S, Kim SH, J. Cosgrove
D, Li S, Gu Y. 2014. The jiaoyao1 Mutant Is an Allele of korrigan1 That Abolishes
Endoglucanase Activity and Affects the Organization of Both Cellulose Microfibrils and
Microtubules in Arabidopsis. *Plant Cell* **26**:2601–2616. doi:10.1105/tpc.114.126193

Lennon KA, Lord EM. 2000. In vivo pollen tube cell of Arabidopsis thaliana I. Tube cell
cytoplasm and wall. *Protoplasma* **214**:45–56. doi:10.1007/BF02524261

Lin D, Cao L, Zhou Z, Zhu L, Ehrhardt D, Yang Z, Fu Y. 2013. Rho GTPase Signaling Activates
Microtubule Severing to Promote Microtubule Ordering in Arabidopsis. *Current Biology*
23:290–297. doi:10.1016/j.cub.2013.01.022

Logemann E, Birkenbihl RP, Ülker B, Somssich IE. 2006. An improved method for preparing
Agrobacterium cells that simplifies the Arabidopsis transformation protocol. *Plant*
Methods **2**:16. doi:10.1186/1746-4811-2-16

Louveaux M, Rochette S, Beauzamy L, Boudaoud A, Hamant O. 2016. The impact of
mechanical compression on cortical microtubules in Arabidopsis: a quantitative
pipeline. *Plant J* **88**:328–342. doi:10.1111/tpj.13290

MacKinnon IM, Šturcová A, Sugimoto-Shirasu K, His I, McCann MC, Jarvis MC. 2006. Cell-
wall structure and anisotropy in procuste, a cellulose synthase mutant of Arabidopsis
thaliana. *Planta* **224**:438–448. doi:10.1007/s00425-005-0208-6

Mizuta Y, Higashiyama T. 2018. Chemical signaling for pollen tube guidance at a glance.
Journal of Cell Science **131**:jcs208447. doi:10.1242/jcs.208447

Mouille G, Ralet M-C, Cavelier C, Eland C, Effroy D, Hématy K, McCartney L, Truong HN,
Gaudon V, Thibault J-F, Marchant A, Höfte H. 2007. Homogalacturonan synthesis in
Arabidopsis thaliana requires a Golgi-localized protein with a putative
methyltransferase domain. *The Plant Journal* **50**:605–614. doi:10.1111/j.1365-
313X.2007.03086.x

Nezhad AS, Naghavi M, Packirisamy M, Bhat R, Geitmann A. 2013. Quantification of cellular
penetrative forces using lab-on-a-chip technology and finite element modeling. *PNAS*
110:8093–8098. doi:10.1073/pnas.1221677110

Nicol F, His I, Jauneau A, Vernhettes S, Canut H, Höfte H. 1998. A plasma membrane-bound
putative endo-1,4-beta-D-glucanase is required for normal wall assembly and cell
elongation in Arabidopsis. *EMBO J* **17**:5563–5576. doi:10.1093/emboj/17.19.5563

Palanivelu R, Tsukamoto T. 2012. Pathfinding in angiosperm reproduction: pollen tube
guidance by pistils ensures successful double fertilization. *Wiley Interdisciplinary*
Reviews: Developmental Biology **1**:96–113. doi:10.1002/wdev.6

Panteris E, Adamakis I-DS, Daras G, Rigas S. 2014. Cortical microtubule patterning in roots
of Arabidopsis thaliana primary cell wall mutants reveals the bidirectional interplay with
cell expansion. *Plant Signal Behav* **9**. doi:10.4161/psb.28737

Paredez AR, Somerville CR, Ehrhardt DW. 2006. Visualization of Cellulose Synthase
Demonstrates Functional Association with Microtubules. *Science* **312**:1491–1495.
doi:10.1126/science.1126551

Rotman N, Durbarry A, Wardle A, Yang WC, Chaboud A, Faure J-E, Berger F, Twell D. 2005. A Novel Class of MYB Factors Controls Sperm-Cell Formation in Plants. *Current Biology* **15**:244–248. doi:10.1016/j.cub.2005.01.013

Ryden P, Sugimoto-Shirasu K, Smith AC, Findlay K, Reiter W-D, McCann MC. 2003. Tensile Properties of Arabidopsis Cell Walls Depend on Both a Xyloglucan Cross-Linked Microfibrillar Network and Rhamnogalacturonan II-Borate Complexes. *Plant Physiol* **132**:1033–1040. doi:10.1104/pp.103.021873

Sampathkumar A, Krupinski P, Wightman R, Milani P, Berquand A, Boudaoud A, Hamant O, Jönsson H, Meyerowitz EM. 2014. Subcellular and supracellular mechanical stress prescribes cytoskeleton behavior in Arabidopsis cotyledon pavement cells. *eLife Sciences* **3**:e01967. doi:10.7554/eLife.01967

Sampedro J, Pardo B, Gianzo C, Guitián E, Revilla G, Zarra I. 2010. Lack of α-Xylosidase Activity in Arabidopsis Alters Xyloglucan Composition and Results in Growth Defects. *Plant Physiol* **154**:1105–1115. doi:10.1104/pp.110.163212

Samuel MA, Tang W, Jamshed M, Northey J, Patel D, Smith D, Siu KWM, Muench DG, Wang Z-Y, Goring DR. 2011. Proteomic Analysis of Brassica Stigmatic Proteins Following the Self-incompatibility Reaction Reveals a Role for Microtubule Dynamics During Pollen Responses. *Mol Cell Proteomics* **10**. doi:10.1074/mcp.M111.011338

Sanati Nezhad A, Geitmann A. 2013. The cellular mechanics of an invasive lifestyle. *Journal of Experimental Botany* **64**:4709–4728. doi:10.1093/jxb/ert254

Sassi M, Ali O, Boudon F, Cloarec G, Abad U, Cellier C, Chen X, Gilles B, Milani P, Friml J, Vernoux T, Godin C, Hamant O, Traas J. 2014. An Auxin-Mediated Shift toward Growth Isotropy Promotes Organ Formation at the Shoot Meristem in Arabidopsis. *Current Biology* **24**:2335–2342. doi:10.1016/j.cub.2014.08.036

Shigeyama T, Watanabe A, Tokuchi K, Toh S, Sakurai N, Shibuya N, Kawakami N. 2016. α-Xylosidase plays essential roles in xyloglucan remodelling, maintenance of cell wall integrity, and seed germination in Arabidopsis thaliana. *J Exp Bot* **67**:5615–5629. doi:10.1093/jxb/erw321

Shoji T, Narita NN, Hayashi K, Asada J, Hamada T, Sonobe S, Nakajima K, Hashimoto T. 2004. Plant-Specific Microtubule-Associated Protein SPIRAL2 Is Required for Anisotropic Growth in Arabidopsis. *Plant Physiology* **136**:3933–3944. doi:10.1104/pp.104.051748

Smyth DR, Bowman JL, Meyerowitz EM. 1990. Early flower development in Arabidopsis. *The Plant Cell* **2**:755–767. doi:10.1105/tpc.2.8.755

Takáč T, Šamajová O, Pechan T, Luptovčiak I, Šamaj J. 2017. Feedback Microtubule Control
and Microtubule-Actin Cross-talk in *Arabidopsis* Revealed by Integrative Proteomic and
Cell Biology Analysis of *KATANIN 1* Mutants. *Molecular & Cellular Proteomics*
16:1591–1609. doi:10.1074/mcp.M117.068015

Uyttewaal M, Burian A, Alim K, Landrein B, Borowska-Wykręt D, Dedieu A, Peaucelle A,
Ludynia M, Traas J, Boudaoud A, Kwiatkowska D, Hamant O. 2012. Mechanical stress
acts via katanin to amplify differences in growth rate between adjacent cells in
Arabidopsis. *Cell* **149**:439–451. doi:10.1016/j.cell.2012.02.048

Verger S, Long Y, Boudaoud A, Hamant O. 2018. A tension-adhesion feedback loop in plant
epidermis. *eLife*. doi:10.7554/eLife.34460

Xiao C, Anderson CT. 2016. Interconnections between cell wall polymers, wall mechanics, and
cortical microtubules: Teasing out causes and consequences. *Plant Signal Behav* **11**.
doi:10.1080/15592324.2016.1215396

Xiao C, Zhang T, Zheng Y, Cosgrove DJ, Anderson CT. 2016. Xyloglucan Deficiency Disrupts
Microtubule Stability and Cellulose Biosynthesis in Arabidopsis, Altering Cell Growth
and Morphogenesis. *Plant Physiol* **170**:234–249. doi:10.1104/pp.15.01395

Zhang T, Mahgsoudy-Louyeh S, Tittmann B, Cosgrove DJ. 2014. Visualization of the
nanoscale pattern of recently-deposited cellulose microfibrils and matrix materials in
never-dried primary walls of the onion epidermis. *Cellulose* **21**:853–862.
doi:10.1007/s10570-013-9996-1

FIGURES

Fig. 1. CMT organisation during papilla cell development. (*A*) Flower development of *A.*
thaliana from developmental stages 12 to 15. Scale bar, 500 µm. (*B*) Upper view of the stigma
during development by SEM. Scale bar, 50 µm. (*C*) Confocal images of papilla cells expressing
MAP65-citrine at each stage of development. Scale bar, 5 µm. (*D*) Quantitative analysis of
CMT array anisotropy of papilla cells from stages 12 to 15. The red dots correspond to the
mean values of the three replicates. Statistical differences were calculated using a Shapiro-
Wilk test to evaluate the normality and then a Wilcoxon test, ***P < 0.01. N > 4 stigmas, n >
60 papilla cells for each stage.

Fig. 2. Pollen tube growth behaviour on papillae during development. (*A*) SEM images of Col-0 papillae pollinated with Col-0 pollen, one hour post pollination, from stages 12 to 15; pollen and tubes were artificially colorised. Scale bar, 5 µm. (*B*) Quantification of the number of turns (T) made by the pollen tube on papillae from stages 12 to 15. Data are expressed as mean +/- s.e.m. A Chi-Square test for independence (at 6 degree of freedom) was used to compare all stages and demonstrated that the number of turns was significantly different between stages (***$P < 0.01$). N corresponds to the number of stigmas analysed in this study and n, the number of papillae.

Fig. 3. Effect of CMT organisation on pollen tube path. (*A*) Confocal images of papilla cells expressing MAP65.1-citrine in Col-0 and *ktn1-5* at stage 13. Scale bars, 5 µm. (*B*) SEM images of Col-0 and *ktn1-5* papillae pollinated with Col-0 pollen grains; pollen and tubes were artificially colorised. Scale bar, 10 µm. (*C*) CMT anisotropy of Col-0 and *ktn1-5* papilla cells at stage 13. N(Col-0) = 10 stigmas, n(Col-0) = 106 papillae, N(*ktn1-5*) = 11stigmas, n(*ktn1-5*) = 114 papillae. Statistical differences were calculated using a Shapiro-Wilk test to evaluate the normality and then a Wilcoxon test with ***P < 0.01. (*D*) Quantification of the number of turns (T) made by Col-0 pollen tubes on *ktn1-5* and Col-0 papillae. Data are expressed as mean +/- s.e.m. Statistical difference was found between pollen tube path within *ktn1-5* and Col-0 papillae and was calculated using an adjusted Chi-Square test for homogeneity (2 degrees of freedom), ***P < 0.01. N(Col-0) = 27 stigmas, n(Col-0) = 251 papillae, N(*ktn1-5*) = 23 stigmas, n(*ktn1-5*) = 327 papillae.

Fig. 4. Local oryzalin application on Col-0 stigmas promotes MT destabilization and induces Col-0 pollen tube coils. (*A*) Col-0 papilla cells expressing MAP65.1-citrine after 4 hours of DMSO (top) or oryzalin (bottom) local treatment. (*B*) SEM images of DMSO- (top) or oryzalin-treated (bottom) Col-0 stigmas pollinated with Col-0 pollen grains; pollen and tubes were artificially colorised. (*A* and *B*) Scale bars, 5 µm. (*C*) Quantification of the number of turns (T) made by Col-0 pollen tubes on drug-treated and control papillae. Data are expressed as mean +/- s.e.m. Statistical difference was found between pollen tube path within DMSO (control) and oryzalin-treated papillae and was calculated using an adjusted Chi-Square test for homogeneity (2 degrees of freedom), ***$P < 0.01$. N(DMSO) = 12 stigmas, n(DMSO) = 117 papillae, N(oryzalin) = 16 stigmas, n(oryzalin) = 149 papillae.

703
Fig. 5. Mechanical properties of papilla cells. (*A*) Location of a Col-0 pollen tube in the cell-
wall of Col-0 and *ktn1-5* papillae by TEM. pt = pollen tube, C = stigma cuticle, sCW = stigma
cell wall, sp = stigma papilla. Scale bar, 1 μm. (*B*) Confocal images of Col-0 and *ktn1-5* papillae
expressing the plasma membrane marker LTI6B-GFP pollinated with an RFP-expressing
pollen. Scale bar, 5 μm. (*C*) Diagram showing the procedure used for evaluating the external
(extD) and internal (intD) deformations made by Col-0 pollen tubes. (*D*) External and internal
deformations caused by Col-0 pollen tube growth in Col-0 and *ktn1-5* papillae. (*E*) Drawing of
the AFM experimental setup. Dissected pistils were inserted in agarose medium and fixed with
low melting agarose for measurements. (*F*) Young's modulus values of the papilla cell wall for
Col-0 at stage 13 (N = 4 stigmas, n = 8 papillae), *ktn1-5* (N = 5 stigmas, n = 9 papillae)*, xxt1*

xxt2 (N = 4 stigmas, n = 11 papillae) and Col-0 at stage15 (N = 4 stigmas, n = 10 papillae. (*G*)

Mean of travel distances made by Col-0 pollen tubes in Col-0 and *ktn1-5* papillae. (*H*) Mean of

travel distances made by Col-0 pollen tubes in papillae of Col-0 at stage 13, Col-0 at stage 15,

xxt1 xxt2 at stage 13.

D-E and *G-H*, Statistical differences were calculated using a Shapiro-Wilk test to evaluate the

normality and then a T-test. *** P < 0.01, n.s. = non significant. For *H*, we found a significant

difference (***P < 0.01) between Col-0 at stage 13 and 15, but non significant (n.s.) between

Col-0 at stage 13 and *xxt1 xxt2* at stage 13.

Fig. 6. Cellulose microfibril organisation in Col-0 stage 13, *ktn1-5*, Col-0 stage 15 and

***xxt1 xxt2* papillae.** (*A*) 3D-projections of z-stack images of Col-0 stage 13, *ktn1-5*, Col-0 stage

15 and *xxt1 xxt2* papilla cells stained with Direct Red 23 dye. Scale bars, 10 μm. (*B*) Plot

profiles of the fluorescence intensity (AFU, Arbitrary Fluorescence Units) along the yellow

lines. Note the similarities of profiles between Col-0 stage 13 and *xxt1 xxt2* on one hand, and

between *ktn1-5* and Col-0 on the other hand.

SUPPLEMENTAL INFORMATION

Fig. S1. Col-0 pollen tube behaviour on Col-0 and *ktn1-5* papillae at stage 13. (*A* and *B*)

Top views of Col-0 (*A*) and *ktn1-5* (*B*) stigmas pollinated with Col-0 pollen grains. Scale bars,

20 µm. (*C* and *D*) Magnification of pollinated papilla cells. (*C*) Col-0 pollen tube grows mainly

straight to the direction of the ovules on Col-0 papillae whereas (*D*) it coils around and can

even grow upward on *ktn1-5* papilla cells (*). Scale bars, 5 µm.

Fig. S2. Ultrastructure of Col-0 and *ktn1-5* papilla cells. (*A*) TEM images of non-pollinated
Col-0 and *ktn1-5* papillae. Scale bar, 5 μm. (*B*) Both Col-0 and *ktn1-5* papillae show a similar
two-layered cell wall (*) delimited by a more electron dense thin layer (arrows) corresponding
to the cuticle. Scale bar, 1 μm.

Fig. S3. **Quantification of the number of coils made by Col-0 pollen tubes on papillae**
from cell-wall mutants at stage 13. (*A*) *xxt1 xxt2* x Col-0. N(Col-0) = 14 stigmas, n(Col-0) =
116 papillae, N(*xxt1 xxt2*) = 12 stigmas, n(*xxt1 xxt2*) = 116 papillae. (*B*) *prc1* x Col-0. N(Col-0)
= 14 stigmas, n(Col-0) = 115 papillae, N(*prc1*) = 12 stigmas, n(*prc1*) = 100 papillae. (*C*) *qua2.1*
x Col-0. N(Col-0) = 22 stigmas, n(Col-0) = 160 papillae, N(*qua2.1*) = 18 stigmas, n(*qua2.1*) =
162 papillae. (*D*) *any1* x Col-0. N(Col-0) = 14 stigmas, n(Col-0) = 91 papillae, N(*any1*) = 14

stigmas, n(*any1*) = 95 papillae. (*E*) *kor1.1* x Col-0. N(Col-0) = 14 stigmas, n(Col-0) = 141

papillae, N(*kor1.1*) = 11 stigmas, n(*kor1.1*) = 91 papillae. (*F*) *xyl1.4* x Col-0. N(Col-0) = 18

stigmas, n(Col-0) = 136 papillae, N(*xyl1.41*) = 16 stigmas, n(*xyl1.4*) = 108 papillae. Data are

expressed as mean +/- s.e.m. No statistical difference was found between pollen tube path

within the cell wall mutants and Col-0 papillae based on an adjusted Chi-Square test for

homogeneity (2 degrees of freedom). (*G*) SEM images of Col-0 and *xxt1 xxt2* stigmas

pollinated with Col-0 pollen grains. Scale bar, 10 μm.

Fig. S4. Size of the papilla cells of Col-0 and cell wall mutants. (*A*) Papilla cell length of Col-0 at stage 13 and stage 15. (*B*) Papilla cell width of Col-0 at stage 13 and stage 15. (*C*) Papilla cell length of Col-0 and *ktn1-5* at stage 13. (*D*) Papilla cell width of Col-0 and *ktn1-5* at stage 13. (*E*) Papilla cell length of Col-0, *xxt1 xxt2* and *any1* at stage 13. (*F*) Papilla cell width of Col-0, *xxt1 xxt2* and *any1* at stage 13. Statistical differences were calculated using a T-test. * P < 0.05, *** P < 0.01, n.s. = non significant. N=4 stigmas and n=120 papillae for each genotypes.

777   **Table S1. Cell-wall mutants tested**

Cell wall component	AGI/Name	Function*	Mutant	References
Cellulose	AT5G49720 *KOR1*	Endoglucanase 25	*kor1*	(His and Driouich, 2001; Lei et al., 2014; Nicol et al., 1998)
Cellulose	AT5G64740 *PRC1*	Cellulose synthaseA catalytic subunit 6	*prc1*	(Fagard et al., 2000; MacKinnon et al., 2006; Panteris et al., 2014; Xiao et al., 2016)
Cellulose	AT4G32410 *ANY1*	Cellulose synthaseA catalytic subunit 1	*any1*	(Fujita et al., 2013)
Hemicellulose	AT3G62720 *XXT1* AT4G02500 *XXT2*	Xyloglucan 6-xylosyltransferase 1 Xyloglucan 6-xylosyltransferase 2	*xxt1 xxt2*	(Cavalier et al., 2008; Xiao et al., 2016)
Hemicellulose	AT1G68560 *XYL1*	Alpha-xylosidase 1	*xyl1.4*	(Shigeyama et al., 2016)
Pectin	AT1G78240 *QUA2*	Probable Pectin methyltransferase	*qua2.1*	(Abasolo et al., 2009; Mouille et al., 2007; Verger et al., 2018)

778   *From https://www.uniprot.org

780