Title:

GenoDup Pipeline: a tool to detect genome duplication using the dS-based method

Authors: Yafei Mao[1*], Noriyuki Satoh[1]

Affiliations:

[1]Marine Genomics Unit, Okinawa Institute of Science and Technology Graduate

University, Onna, Okinawa 904-0495, Japan

*Correspondence to: Yafei Mao: yafei.mao@oist.jp

Yafei Mao: yafei.mao@oist.jp

Noriyuki Satoh: norisky@oist.jp

ABSTRACT

Understanding whole genome duplication (WGD), or polyploidy, is fundamental to

investigating the origin and diversification of organisms in evolutionary biology. The

wealth of genomic data generated by next generation sequencing (NGS) has resulted

in an urgent need for robust and accurate tools to detect WGD. Here, we present a

useful and user-friendly pipeline called GenoDup for inferring WGD using the dS-

based method. We have successfully applied GenoDup to identify WGD in empirical

data from both plants and animals. The GenoDup Pipeline provides a reliable and

useful tool to infer WGD from NGS data.

Keywords: Whole-genome duplication (WGD), Polyploidy, dS-based, Age

distribution, Software, GenoDup, Next generation sequencing (NGS)

INTRODUCTION

Whole (large-scale)-genome duplication (WGD), or polyploidy, has been regarded as

an evolutionary landmark in the origins and diversification of animals, plants, and

other evolutionary lineages (Van De Peer, et al. 2017). Previous studies have shown

that WGD plays an important role in enhancing speciation and reducing risks of

extinction. Moreover, evolutionary novelty can be generated by duplicated genes via

subfunctionalization, neofunctionalization, and dosage effects under WGD (Glasauer

and Neuhauss 2014; Van De Peer, et al. 2017). Identification of WGD is the first step

to understanding the impacts of WGD and the fates of duplicated genes. In addition,

WGD is now known to be a common event in plants, since the availability of genomic

data generated by next generation sequencing (NGS). However, recent studies also

suggest that WGD is a common evolutionary force in animals (Li, et al. 2018; Van De

Peer, et al. 2017). Hence, an easy-to-use pipeline is urgently needed to infer WGD

using NGS data.

There are three main approaches to infer WGD with NGS data (Tiley, et al. 2016).

First, identification of synteny blocks is the most straightforward method to detect

WGD, but it requires high-quality genome assembly, and sadly, many genomes have

not yet reached that assembly quality. Second, phylogenetic analysis of gene families

can unravel WGD when organisms have undergone extensive gene loss or genome

shuffling, but the uncertainties of gene tree reconstruction is a serious limitation and

heavy computation is required. Finally, analysis of rates of synonymous substitutions

per synonymous site (dS) of duplicated genes (the dS-based method or age

distribution method) is the most common and widely used approach to infer WGD.

The dS-based method is often a fragmented step-wise process (Vanneste, et al. 2014).

Multiple software packages are required to build gene pairs, align sequences, and

calculate dS values. Usually, there are two approaches to build gene pairs using the

dS-based method. First, gene pairs can be generated using gene family cluster

information as paralogous gene pairs. Second, gene pairs can be generated by synteny

information as anchor gene pairs. Consistent results from both approaches yield a

credible conclusion. DupPipe is a web-based method to infer WGD using the dS-

based method, but users need to wait for a long time to get results (Barker, et al.

2010). In addition, DupPipe only calculates dS values based on paralogous gene pairs.

Here, we develop an open-source script called GenoDup Pipeline, which integrates all

processes with only one command and it calculates dS values based on paralogous

gene pairs or/and anchor gene pairs.

IMPLEMENTATION

GenoDup Pipeline architecture

GenoDup Pipeline is written in Python with all alignment of sequences, building gene

pairs, and dS value calculations. BioPython must be installed and three more

executable dependencies are needed: MAFFT (Katoh, et al. 2002), Translatorx

(Abascal, et al. 2010), and Codeml package in PAML (Yang 2007). Nuclear protein

coding sequences and corresponded protein sequences are mandatory input files to

run GenoDup Pipeline. In addition, gene family cluster information or anchor gene

pairs information is another mandatory input file for paralogous gene pairs and anchor

gene pairs approach respectively. Once input files have been inputted appropriately, 3

subroutines in GenoDup run as follows (Fig.1).

(1). Alignment of gene pairs: GenoDup Pipeline can automatically align gene pairs

using MAFFT and Translatorx. Before performing gene pair alignments, GenoDup

first filters ambiguous sequences that contain 'N', and removes sequences that nuclear

protein coding sequences cannot be translated to match corresponding protein

sequences. Then, MAFFT is used to perform alignments of protein sequences with

parameters: localpair and maxiterate: 1000; and Translatorx are used to align nuclear

protein coding sequences based on alignments of protein sequences.

(2). Building gene pairs: there are two ways to build gene pairs in GenoDup Pipeline.

The first requires a gene family cluster information file and a number that the

maximum number of genes in a gene family should be analyzed, as input. Then,

GenoDup Pipeline builds $N(N-1)/2$ paralogous gene pairs within a gene family (N is

the number of genes in a gene family). We recommend OrthoMCL to generate gene

family cluster information (Li, et al. 2003). Second, it requires an anchor gene pair

information file. The anchor gene pair information file contains paralogs located on

same synteny blocks and we recommend MCScanX (Wang, et al. 2012) or i-ADHoRe

(Proost, et al. 2011) to generate anchor gene pair information.

(3). dS value calculations: based on alignments of protein coding sequences,

GenoDup can automatically build control files, required by Codeml, for each gene

pair. Codeml package in PAML is used to calculate dS values with parameters: noisy

= 9, verbose = 1, runmode = -2, seqtype = 1, CodonFreq = 2, model = 0, NSsites = 0,

icode = 0, fix_kappa = 0, kappa = 1, fix_omega = 0, and omega = 0.5.

All of the 3 subroutines above can automatically run in the GenoDup Pipeline, which

finally generates two directories and one text file as output. One directory called

aln_genefam contains all nuclear protein coding sequence alignments in Fasta format.

The other directory, called PAML_result contains all results generated by Codeml,

and a text file called dS_value.txt contains all dS values of gene pairs. We also

provided an R script (plot_GenoDup.r) to plot dS distributions.

Empirical data validation

To evaluate the performance of the GenoDup Pipeline, we applied it to two empirical

data: one is a model plant (*Arabidopsis thaliana*) and the other is a model animal

(*Oncorhynchus mykiss*). *Arabidopsis thaliana* has undergone two independent WGD

(alpha and beta WGD) and *Oncorhynchus mykiss* has experienced four independent

WGD (Ss4R, Ts3R, and Two-rounds WGD) (Berthelot, et al. 2014; Vanneste, et al.

2014).

We downloaded the nuclear protein coding sequences, protein sequences, and genome

annotation files of *Arabidopsis thaliana* from Ensembl Plants

(http://plants.ensembl.org/Arabidopsis_thaliana/Info/Index). We generated gene

family clusters with OrthoMCL and clustered 48,307 genes of *Arabidopsis thaliana*

into 5,962 gene families. We set the maximum number of genes in a gene family

analyzed in GenoDup Pipeline as 15 and generated 68,231 paralogous gene pairs. The

entire analysis ran in 8 .8 hours with 4 cores. The result of the analysis showed an

obvious peak (dS value range: 0.5~1) in dS distributions of paralogous gene pairs. On

the other hand, we generated anchor gene pair information with MCScanX. We used

MCScanX to generate 99,309 anchor gene pairs and the entire analysis ran in 76.8

hours with 4 cores. The result of the analysis showed the same peak (dS value range:

0.5~1) in the dS distribution of anchor gene pairs as the result based on gene family

cluster information.

We downloaded the nuclear protein coding sequences, protein sequences, and genome

annotation files of *Oncorhynchus mykiss* from GENEOSCOPE

(http://www.genoscope.cns.fr/trout/data/). We generated gene family cluster by

OrthoMCL and clustered 46,585 genes of *Oncorhynchus mykiss* into 6,562 gene

families. We set the maximum number of genes in a gene family analyzed in

GenoDup Pipeline as 15 and generated 42,888 paralogous gene pairs. The entire

analysis ran in 7.63 hours with 4 cores. The result of the analysis showed two obvious

peaks (dS value ranges: 0.1~0.5 and 1.2~2) in the dS distribution of paralogous gene

pairs. On the other hand, we generated anchor gene pair information identified by

MCScanX. We used MCScanX to generate 26,880 anchor gene pairs and the entire

analysis ran in 17.65 hours with 4 cores. The result of the analysis showed the same

peak (dS value range: 0.1~0.5) in the dS distribution of anchor gene pairs as the result

based on gene family cluster information.

Our GenoDup Pipeline represented consistent results as previous studies on

Arabidopsis thaliana and *Oncorhynchus mykiss*, respectively (Berthelot, et al. 2014;

Vanneste, et al. 2014), indicating that GenoDup Pipeline is a reliable tool to infer

WGD using the dS-based method.

DISCUSSION

The rapid development of NGS technologies has enabled generation of massive

amounts of data, allowing us to better understand the evolutionary history of all

organisms. WGD is suggested to have occurred in diverse organismal groups (Li, et al.

2018; Van De Peer, et al. 2017); thus, a reliable, efficient tool to detect WGD with

NGS data is greatly needed. We developed a reliable and easy-to-use tool called

GenoDup Pipeline to infer WGD using the dS-based method. GenoDup Pipeline is

written by Python and can be run with one command. It is easy to use for researchers

who have a little background in bioinformatics.

In GenoDup Pipeline, it is faster to run to take gene family cluster information as

input rather than to take anchor gene pair information as input. Because GenoDup

Pipeline aligns sequences for each gene family rather than for anchor gene pairs

individually, GenoDup Pipeline spends much time on alignments of anchor gene pairs

individually when taking anchor gene pair information as input.

Our empirical validation shows that the analysis of *Arabidopsis thaliana* generated

with GenoDup Pipeline presented an obvious signal for one WGD event (alpha WGD)

but the signal of the second WGD event (beta WGD) was lost. This result is

consistent with previous studies because the dS methods cannot infer WGD when

organisms have undergone extensive gene loss or genome shuffling, especially for

plants (Rabier, et al. 2014; Tiley, et al. 2016). Additionally the *Oncorhynchus mykiss*

analysis with GenoDup showed two obvious two WGD signals (Ss4R and Ts3R) in

the dS distribution of paralogous gene pairs because of high conservation in the

Oncorhynchus mykiss genome (Rabier, et al. 2014). Yet, The Ts3R signal was lost in

the distribution of anchor gene pairs because there were few anchor gene pairs in the

analysis. Moreover, the both analysis on *Oncorhynchus mykiss* do not present the

two-round WGD because it is too ancient to infer with the dS-based method.

Importantly, the dS-based method has been debated to generate artificial signals,

because of dS saturation when the dS value >1 (Vanneste, et al. 2012). Thus,

GenoDup Pipeline is suitable to infer WGD when the dS value <1, and other evidence

from phylogenetic analysis and/or synteny blocks analysis is needed to draw a

conclusive result.

In all, with more and increasingly large sequencing projects (1KP, 10KP-EBP, 1KITE,

and i5K) developing (Li, et al. 2018; Van De Peer, et al. 2017), we present a reliable

and user-friendly tool to infer WGD using the dS-based method.

ACKNOWLEDGEMENTS

This study was supported by OIST and was funded by JSPS grant (No. 17J00557 to

YFM). We thank Dr. Steven D. Aird for editing the manuscript.

Availability and requirements

Project name: GenoDup Pipeline

Source code: https://github.com/MaoYafei/GenoDup-Pipeline

Documentation: https://github.com/MaoYafei/GenoDup-

Pipeline/blob/master/README.md

Operating system: Linux (Red Hat, CentOS, and Ubuntu)

Programming language: Bash and Python

Other requirements: MAFFT, Translatorx, Codeml package in PAML

Funding

This study was supported by OIST and was funded by JSPS grant (No. 17J00557 to

YFM).

Competing Interests

The authors declare there are no competing interests.

Author Contributions:

Yafei Mao conceived and designed the experiments, performed the experiments,

analyzed the data, wrote the paper, and reviewed drafts of the paper.

Noriyuki Satoh supervised the project, wrote the paper, and reviewed drafts of the

paper.

Data Availability

The following information was supplied regarding data availability:

Github: https://github.com/MaoYafei/GenoDup-Pipeline

REFERENCES

Abascal, F., R. Zardoya, and M. J. Telford. 2010. TranslatorX: multiple alignment of

nucleotide sequences guided by amino acid translations. *Nucleic Acids Research* 38:

W7-13.

Barker, Michael S., et al. 2010. EvoPipes. net: bioinformatic tools for ecological and

evolutionary genomics. *Evolutionary Bioinformatics* 6:EBO. S5861.

Berthelot, Camille, et al. 2014. The rainbow trout genome provides novel insights into

evolution after whole-genome duplication in vertebrates. *Nature communications*

5:3657.

Glasauer, S. M. K., and S. C. F. Neuhauss. 2014. Whole-genome duplication in

teleost fishes and its evolutionary consequences. *Molecular Genetics and Genomics*

289(6):1045-1060.

Katoh, K., et al. 2002. MAFFT: a novel method for rapid multiple sequence alignment

based on fast Fourier transform. *Nucleic Acids Research* 30(14):3059-66.

Li, L., C. J. Stoeckert, Jr., and D. S. Roos. 2003. OrthoMCL: identification of

ortholog groups for eukaryotic genomes. *Genome Research* 13(9):2178-89.

Li, Zheng, et al. 2018. Multiple large-scale gene and genome duplications during the

evolution of hexapods. *Proceedings of the National Academy of Sciences*:201710791.

Proost, Sebastian, et al. 2011. i-ADHoRe 3.0—fast and sensitive detection of

genomic homology in extremely large data sets. *Nucleic acids research* 40(2):e11-

e11.

Rabier, C. E., T. Ta, and C. Ane. 2014. Detecting and Locating Whole Genome

Duplications on a Phylogeny: A Probabilistic Approach. *Molecular Biology and*

Evolution 31(3):750-762.

Tiley, G. P., C. Ane, and J. G. Burleigh. 2016. Evaluating and Characterizing Ancient

Whole-Genome Duplications in Plants with Gene Count Data. *Genome Biology and*

Evolution 8(4):1023-1037.

Van De Peer, Y., E. Mizrachi, and K. Marchal. 2017. The evolutionary significance of

polyploidy. *Nature Reviews Genetics* 18(7):411-424.

Vanneste, K., et al. 2014. Analysis of 41 plant genomes supports a wave of successful

genome duplications in association with the Cretaceous-Paleogene boundary. *Genome*

Research 24(8):1334-1347.

Vanneste, Kevin, Yves Van de Peer, and Steven Maere. 2012. Inference of genome

duplications from age distributions revisited. *Molecular biology and evolution*

30(1):177-190.

Wang, Y. P., et al. 2012. MCScanX: a toolkit for detection and evolutionary analysis

of gene synteny and collinearity. *Nucleic Acids Research* 40(7).

Yang, Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular*

biology and evolution 24(8):1586-91.

Table 1. Statistics of empirical data validation in GenoDup Pipeline

	Arabidopsis thaliana		*Oncorhynchus mykiss*	
	Paralogous gene pairs	Anchor gene pairs	Paralogous gene pairs	Anchor gene pairs
The number of gene pairs	68,231	99,309	42,888	26,880
Running Time (h)	8.88	76.8	7.63	17.65

Fig 1. Workflow in GenoDup Pipeline. Orange oval boxes represent input files.

Blue boxes represent three subroutines in the GenoDup Pipeline. Red boxes represent

output files generated by GenoDup Pipeline.

Fig 2. dS distributions of paralogous gene pairs and anchor gene pairs in

***Arabidopsis thaliana*.** The peak (dS value range: 0.5~1) marked with a red box

represents a signal as alpha WGD in *Arabidopsis thaliana*.

Fig 3. dS distributions of paralogous gene pairs and anchor gene pairs in

Oncorhynchus mykiss. The peaks (dS value ranges: 0.1~0.5 and 1.2~2) marked with

red boxes represent signals of the Ss4R and Ts3R WGDs of *Oncorhynchus mykiss*,

respectively, in the dS distributions of anchor gene pairs. The peak (dS value range:

0.1~0.5) marked with a red box represents the signal attributable to Ss4R WGD of

Oncorhynchus mykiss in the dS distributions of anchor gene pairs.