Different neural networks for conceptual retrieval in sighted and blind reveal the experiential bases of knowledge.

Authors:

Roberto Bottini[(1,2)*], Stefania Ferraro[(3)], Anna Nigri[(3)], Valeria Cuccarini[(3)], Maria Grazia Bruzzone[(3)], Olivier Collignon[(1,4)*]

Affiliations:

[1] Center for Mind/Brain Studies (CIMeC), University of Trento, Trento, Italy

[2] International School for Advanced Studies (SISSA), Trieste, Italy

[3] Neuroradiology Department, Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy.

[4] Institute of research in Psychology (IPSY) & Institute of Neuroscience (IoNS) - University of Louvain (UCL), Louvain, Belgium

*Correspondence to:

Roberto Bottini
CIMeC
Via delle regole 101, Mattarello, Italy
Email: bottini.r@gmail.com

Olivier Collignon
Institute of Psychology (IPSY) & Institute of Neuroscience (IoNS)- University of Louvain (UcL)
Email: olivier.collignon@uclouvain.be

Abstract

We investigated the experiential bases of knowledge by asking whether people that perceive the world in a different way also show a different neurobiology of concepts. We characterized the brain activity of early-blind and sighted individuals during a conceptual retrieval task in which participants rated the similarity between color and action concepts. Between-categories analysis showed that, whereas multimodal concepts (action) activated a similar fronto-temporal network in the sighted and blind, color knowledge activated partially different brain regions in the two groups, with the posterior portion of the right IPS being significantly more active in the sighted compared to the blind. Interestingly, regions that were similarly activated in sighted and blind during conceptual processing (lpMTG for action, and precuneus for color), showed, however, an increased task-dependent connectivity with occipital regions in the blind. Finally, within-category adaptation analysis showed that word-pairs referring to perceptually similar color or actions led to repetition-suppression in occipital visual areas in the sighted only, whereas adaptation was observed in language-related temporal regions in the blind. Our results show that visual deprivation changes the neural bases of conceptual retrieval, which is partially grounded in sensorimotor experience.

Keywords: Concepts; Color; Action; Blindness; Neuroplasticity; fMRI; Adaptation

Significance statement: (max 120 words)

Do people with different sensory experience conceive the world differently? We tested whether conceptual knowledge builds on sensory experience by looking at the neurobiology of concepts in early blind individuals. We show that cortical regions involved in the processing of multimodal concepts (actions) are mostly similar in blind and sighted, whereas strictly visual concepts (colors) activated partially different brain topographies. Moreover, brain regions classically involved in conceptual retrieval showed different connectivity profiles in the blind, working in concert with re-organized "visual" areas. Finally, we further demonstrate that perceptual distance between concepts is represented in the visual cortex of the sighted, but not the blind. Blindness changes how the brain implements conceptual knowledge, which is partially grounded in sensorimotor experience.

Introduction

As we think, we navigate and retrieve conceptual knowledge, but the nature of this knowledge is highly debated. One of the major sources of disagreement is whether thinking is more similar to the analog replay of our experience [1,2], or the symbolic computations of a Turing machine [3,4]. On one side, empiricist approaches suggest that concepts are based on sensorimotor representations: Thinking is simulating (or re-instantiating) our own sensorimotor experience of the world [5,6]. In support of this view, neuroimaging studies have shown that people activate sensorimotor areas when retrieving conceptual knowledge [7–9]. For instance, words referring to objects with specific colors or shapes (e.g., tomato, ball), or directly to color and shape properties (e.g., red, squared), may activate occipito-temporal areas such as V4 and LOC, respectively [10–13]. Similarly, words referring to action verbs or manipulable objects activate neural networks involved in action execution and observation, such as somatosensory areas, premotor cortices and temporal-occipital areas [10,14].

Yet, empiricist theories of concepts has been repeatedly challenged by more rationalistic approaches suggesting that knowledge is retrieved and represented via abstract symbols and propositional representations that are modality-invariant [15,16]. In this view, conceptual representations are particular configurations of interconnected symbols that are neither tactile, nor motor, nor visual. Thus, conceptual retrieval does not require the on-line activation of the sensorimotor system [16]. In this view, putative sensorimotor activations elicited during conceptual processing in several studies, could instead be interpreted as the activity of symbolic (experience-invariant) neural computations that may occur in the proximity of sensorimotor cortices [15]. One of the strongest line of support for this view comes from the cognitive neuroscience of blindness [15,17]. The reasoning is straightforward: If conceptual processing is largely

grounded into experience, people who experience the world in a different way should also show a different neurobiology of concepts, at least in part [18]. Congenital blindness is an ideal model to test this hypothesis, and the available data seem to support the symbolic view. In a seminal study [19], when sighted and blind were asked to retrieve information about geometric figures (e.g., is a pyramid curved?) or highly visual entities (e.g., is the dusk usually dark?), people in both groups activated the same fronto-temporal semantic retrieval system, but no visual (or other) areas were more active in sighted than blind. Similarly, in other experiments, knowledge about objects' shape [20], or small and manipulable objects [21] activated the lateral occipital complex; thinking about big non-manipulable objects activated the parahipoccampal place area [22]; and processing action verbs (compared to nouns) activated the left posterior middle temporal gyrus [23] in both sighted and blind. These results seem to suggest that blindness leaves the neurobiology of conceptual retrieval largely unchanged, and that experience plays a minor role in shaping mental representations[15].

However, previous studies involving congenitally blind people have some important limitations. Most of these studies investigated highly multimodal conceptual categories, such as shape, actions and tools, which can be perceived across different senses. In these cases, blind can re-map visual features to other senses minimizing the impact of visual deprivation [24], especially in brain regions where different sensory information seem to converge. For instance, the anterior ventral occipito-temporal cortex (aVOTC), that showed stunning functional resilience to visual deprivation [22,25], seems to support the integration of visual, auditory and tactile information in multimodal representations [8,9,26–28] that are crucial for categorical knowledge [29] (e.g., distinguish faces from houses, or animals from tools). At this level of the ventral-occipital hierarchy, different categories (e.g., object, animals, tools,

actions) are represented in a format that is relatively independent from visual appearance [30–32], thus more likely to be resilient to visual deprivation. In contrast, posterior occipital regions of the brain organize information based on a visual code that is largely independent of categorical membership [30,33]. Therefore, to the extent that posterior occipital areas encoding visual features are also recruited during conceptual retrieval [10,34,35], differences between sighted and blind should emerge in these regions, as predicted by empiricist theories.

In our study, we investigated the impact of blindness on how the brain implements conceptual knowledge, in two ways. First, we compared a highly multimodal category such as action, with a highly visual-unimodal category such as color. Since visual experience of actions can be remapped to experience in other senses, we predicted that the neural network involved in retrieving action-categories in the sighted will be preserved in the blind [23]. On the contrary, since color knowledge cannot be experienced with nonvisual senses, becoming an abstract property of objects for the blind [36], we predicted to see different brain areas involved in its representation, as a function of visual deprivation. Second, we investigated the neural activity related to within-category perceptual similarity (e.g. similar action Vs different actions). We predicted that perceptual similarity would be represented in posterior occipital areas in the sighted, but not in the blind. These results would (i) be in line with previous results showing that these areas encode visual features in the sighted [30,33], (ii) confirm their involvement in conceptual retrieval [10,34,35] and (iii) show, for the first time to our knowledge, that visual deprivation changes the neuro-biological basis of conceptual processing by precluding visual representations.

Results

Participants (18 Early Blind and 18 Sighted controls; see table S1 for demographic details) listened to pairs of action (e.g. kick - jump) or color words (e.g., red - yellow) in the MRI, and rated each pair in terms of similarity on a scale from 1 to 5 (1= very different; 5= very similar).

Univariate analysis: Sighted and blind activate a similar brain network for multimodal action concepts but not for unimodal (visual) color concepts.

Behavioral analysis: Reaction times analysis using a mixed ANOVA, with Category (action, color) as within-subject factor and Group (sighted, blind) as between-subjects factor, showed no difference between categories ($F(1,33)=2.37$, $p>0.05$, $\eta^2= 0.07$), between groups ($F(1,33)=0.074$, $p>0.05$, $\eta^2=0.002$) and no Category by Group interaction ($F(1,33)=0.69$, $p>0.05$, $\eta^2=0.02$).

fMRI analysis: The contrast Action > Color did not reveal any significant difference between groups, suggesting a comparable neural activity, across sighted and blind, during the retrieval of action concepts (see fig. S1 for details). Indeed, a conjunction analysis between groups showed a common significant activation in the left posterior middle temporal gyrus (lpMTG; Peak= -54, -61, 5; Fig 1A).

--- Figure 1 about here ---

On the other hand, analysis for the contrast Color > Action revealed a cluster in the right parietal cortex, in and around the right intraparietal sulcus (rIPS), showing higher activity for color concepts in sighted compared to blind (peak= 33, -43, 35; Fig. 1B).

A conjunction analysis between groups, for the contrast Color > Action, did not reveal any common activation between sighted and blind after correction for

multiple comparisons at the whole brain level, in line with the hypothesis that color-related conceptual processing engages partially different neural systems as a function of visual deprivation (see fig. S1 for details). However, displaying the conjunction results at a more lenient threshold (p<.001 uncorrected; Fig 1C), we could notice a unique common activity for color concepts in the right precuneus (peak= 6, -55, 26). Accordingly, analysis within groups showed a significant precuneus activity in the blind (peak= 6, -52, 20, p= .04) and a marginally significant activity in the sighted (peak= 0, -61, 29, p= .06), with no significant difference between groups (Table S1; Fig. S1).

Psychophysiological interactions: The precuneus and lpMTG display similar category selectivity across sighted and blind but show different connectivity profiles. We relied on Psychophysiological Interaction (PPI) analysis [37,38] to test whether regions that showed similar categorical preference across groups (lpMTG for action > color, and the precuneus for color > action) also maintain a similar connectivity profile in both groups. In particular, we tested the hypothesis that posterior occipital areas in the blind could be recruited during conceptual retrieval, and connect with conceptual hubs [39] such as the lpMTG and the Precuneus, as a consequence of neural reorganization [40,41] (See also Fig. S2 showing higher activity in the occipital cortex of EB versus SC when processing both conceptual domains compared to rest). With this aim, we selected two ROIs in the left and right middle occipital gyrus (MOG) that (i) are recruited in early blind during high-level conceptual tasks such as language processing and math [42,43]; (ii) show increased long range connectivity, in early blind, with extra-occipital regions (e.g., frontal, parietal and ventral temporal cortices) during resting state [42–45] and task-based (PPI) analysis [19].

PPI with seed in the lpMTG revealed an increase of action-selective functional connectivity in both occipital ROIs of blind people compared to their sighted counterpart (lMOG: t= 3.59, p= 0.02; rMOG: t=3.46, p=0.026; Fig 1D & 1E). Similarly, PPI with seed in the precuneus revealed an increase of color-selective functional connectivity in the occipital cortices of blind compared to sighted participants (lMOG: t= 3.12, p= 0.054; rMOG: t=4.09, p=0.007; Fig 1F and 1G). Albeit showing a similar activity profile in sighted and blind during conceptual processing, the lpMTG and the precuneus showed a different connectivity profile as a function of early visual deprivation. Such increase in task-based connectivity suggests that occipital areas in early blind can be flexibly recruited during conceptual processing in interaction with conceptual hubs.

On the other hand, PPI with seed on the right IPS, for the contrast Color>Action, did not reveal any significant color-specific connectivity profile neither in SC nor for the contrast SC>EB.

Adaptation analyses: Within-category similarity is encoded in occipital areas in the sighted but not in the blind. Classic univariate and functional connectivity analyses revealed a multifaceted impact of visual deprivation on conceptual retrieval. Yet, they did not reveal any preferential activity in the visual cortex of sighted more than blind, as would be predicted by an empiricist approach. With adaptation analyses we tested whether this difference emerges when we look for areas that encode perceptual similarity. The rationale was that the direct contrast between pairs with high versus low perceptual differences will induce a release from adaptation [46–48] therefore probing regions that are specifically sensitive to the *perceptual distance* between concepts. Our prediction was that that the finer grained encoding of

perceptual distances would be represented in posterior occipital areas in the sighted, but not in the blind.

Behavioral analysis: Similarity ratings were highly correlated between sighted and blind, both for action (r= .99) and color words (r= .93; Fig. 2A). In order to perform the adaptation analysis we divided the trials in similar pairs (e.g. red - orange) and different pairs (e.g. red - blue), based on each participant' subjective ratings. Rating distributions for each subject and category (color, action) were divided in 5 intervals with a similar number of items (see Method session for details). Stimulus-pairs in the first two intervals were labeled as different (low similarity ratings), the 3rd interval contained medium pairs, and the 4th-5th intervals similar pairs (high similarity ratings; Fig 2B). Overall, the average number of "different" trials was slightly larger than the "similar" ones (126 vs 115; F=8.41, p=0.007, η^2=0.20; Fig. 2C). However, there was no similarity by group interaction (F=0.18, p=0.67, η^2=0.004), indicating that this unbalance (that reflected personal judgments of similarity) was the same across SC and EB (fig. 2C). An analysis of reaction times showed that Medium pairs (not analyzed in fMRI) had on average longer latencies than Similar and Different ones (Main Effect of Similarity: F=21.07, p<0.001, η^2=0.38). This was expected since pairs that are neither similar nor different would require longer and more difficult judgments. Crucially, there was no difference in reaction times between different (Mean=1.80 sec, SD=0.39) and similar pairs (Mean=1.79 sec, SD=0.37; F=0.09, p=0.76, η^2=0.003), and no interaction between Similarity and Group (F=0.04, p=0.84, η^2=0.001; Fig 2D).

--- Figure 2 about here ---

fMRI analysis: to find brain areas that showed adaptation based on conceptual similarity, we looked at the contrast Different Pairs > Similar Pairs, with Medium pairs as a regressor of no interest. Action and color pairs were considered together since, at the whole brain level, we did not find a significant higher-order interaction between Similarity (different, similar) and Category (see the method section for analysis details). In the sighted, similar concepts led to repetition suppression in several occipital areas (See Fig. 3A), with a significant cluster in and around the left lingual gyrus (peak coordinates: -24, -70, -7). In the blind, instead, adaptation emerged in language-related areas with significant clusters along the middle and superior temporal gyrus, bilaterally (Peak coordinates RH: 57, -28, 8; LH: -60, -10, -7) and in the right precentral gyrus (Peak coordinates: 27, -25, 56; Fig. 3B). Importantly, no adaptation in posterior occipital areas was observed in the blind.

A comparison between groups showed greater adaptation in occipital cortices for sighted compared to blind (Fig. 3C), with peaks in the left superior occipital gyrus (-24, -91, 26), the left lingual gyrus (-24, -70, -7) and the right middle occipital gyrus (27, -85, 11). The contrast Blind > Sighted showed increased adaptation in posterior lateral temporal cortices (PLTC) bilaterally (Fig. 3D). Planned ROI analysis in PLTC, a region that consistently show repetition suppression for semantic similarity [49–51], revealed a significantly greater adaptation for similar concepts in blind more than sighted (Conceptual Similarity by Group interaction; lPLTC= -45 -31 20, t=3.23, P= 0.035; rPLTC= 45 -28 11, t=3.41, P= 0.024).

--- Figure 3 about here ---

Finally, we performed planned ROI analysis in the color-sensitive region at the posterior banks of the collateral sulcus (PCoS) corresponding to the V4-complex

[10,52] and the motion sensitive region V5 [53]. In area PCoS-V4 we found greater adaptation both for color and action in sighted compared to blind (Conceptual Similarity by Group interaction; peak= -24 -73 -10, t=4.26, P= 0.004; Fig 3 E-F). In contrast, the analysis in V5 showed that repetition suppression was specific for action concepts in the sighted and no adaptation was observed in the blind (Conceptual Similarity by Group by Category interaction; Peak: -51 -76 8, t=3.29, P= 0.037; Fig 3F).

Discussion

Embodied approaches to conceptual knowledge suggest that concepts are grounded in our sensory and motor experience of the world [2,7]. A straightforward hypothesis emerging from these theories is that people that perceive the world in a different way should also have different conceptual representations [18]. Congenital blindness offers an ideal test-bed for this hypothesis, which is crucial to validate empiricist theories of concepts: If seeing the color red adds nothing to the neural representation of the concept of red, which can be constructed equally by learning its properties through a verbal code, then the idea that we rely on distributed analogical simulations to construct and retrieve knowledge becomes necessarily weak [15].

In our study we tested this hypothesis by characterizing the brain activity of sighted and early blind individuals while they rated the similarity of action and color concepts in fMRI. Whereas the experience of action is highly multimodal, and accessible to blind people from the remaining senses, the sensory experience of color is purely visual, and inaccessible to the blind. This epistemological difference was reflected in the differential brain activations observed in our blind and sighted groups. Sighted and blind activated a similar fronto-temporal network to retrieve action knowledge, with the highest common activity located in the left posterior MTG,

and no brain area showing between-groups differences (see fig S1 for details). In contrast, color knowledge activated different brain regions in the two groups, with the posterior portion of the right IPS being significantly more active in the sighted compared to the blind. The IPS is known to be involved in the perception of color [52,54,55] as well as other visual features [56–58], and its anatomical position make it a good candidate to work at the interface between perceptual and conceptual representations [54]. In particular, it seems to be crucial for retrieving perceptually-based categorical knowledge [54,55]. Indeed, the peak of color specific activity that we found (peak coordinates: 33, -43, 35) is very close to the color-specific rIPS area found by Cheadle & Zeki (ref 54; peak coordinates: 30, -39, 45). The lack of visual input in blind people prevents the formation of perceptually-driven color representations, which may limit the contribution of the IPS during the retrieval of color knowledge. This is not the case for action representation, for which perceptual knowledge can be compensated by other senses (e.g., touch, audition). In sum, this result shows that, in the case of visual-unimodal categories for which direct experience cannot be compensated by nonvisual perception, conceptual retrieval is instantiated in blind and sighted through partially different neural networks.

However, in keeping with previous findings [19,20,23], we also found commonalities across the two groups. Both sighted and blind similarly engaged the lpMTG during action processing and the Precuneus during color processing. Interestingly, though, psychophysiological interactions (PPI) showed that early visual deprivation changes the connectivity profile of these regions, increasing their functional coupling with occipital regions in the blind. This result suggests that the occipital cortex in early blind is re-organized to extend its integration into conceptual selective networks, highlighting further how visual deprivation impact on the neurobiology of conceptual knowledge. Previous studies have shown that occipital

areas in the early blind are recruited for high-level conceptual tasks such as language processing, semantic retrieval and math [42,43,59–61], and that they increase long-range connectivity with frontal and parietal cortices during rest [42–45] and inferior temporal cortices during semantic judgments [19]. However, here we showed that early blind seem to rely on enhanced connectivity between occipital cortices and temporo-parietal conceptual hubs during conceptual processing. Albeit our data remain correlational, they suggest that EBs' occipital regions are not activated independently of other "classic" regions involved in conceptual retrieval, but they instead work in concert.

Classic univariate and PPI analysis showed that early visual deprivation modifies the neural network supporting the retrieval of color knowledge and the connectivity pattern of temporal and parietal areas involved in conceptual processing. Yet, these analyses failed to show any concept-related activity in visual areas, in sighted more than blind, as an empiricist account would predict. This is not surprising, since both action and color are highly-visual categories for sighted, and should both engage visual simulations.[1] In order to reveal perceptual-specific mechanisms of conceptual retrieval we needed to investigate finer-grained patterns of representations based on perceptual similarity. We did so with our adaptation analysis, probing regions that are specifically sensitive to the *perceptual distance* between concepts of a same class. Our results demonstrated that, within each conceptual domain (action and color), concepts that are judged by individual subject as perceptually different induced a release from adaptation in several posterior

---

[1] It is worth noticing that previous studies have found activations in the anterior fusiform gyrus (anterior V4) for the univariate contrast color > action [12,13,83]. However, all these experiments involved judgments about the color of objects, and words were presented visually. On the other hand, studies that used simple color words (e.g., red, blue, green) and auditory stimuli, failed to find color-related activity in the anterior fusiform [84–86], in line with our results. Indeed, color-specificity in anterior V4 may be specific for object-color or refer, more generally, to multimodal superficial properties of objects [10].

occipital regions (e.g., Lingual Gyrus, V5, PCoS-V4, middle and superior occipital gyri) in the sighted but not in the blind. These posterior occipital regions, in the sighted are known to encode visual features and to be sensitive to visual similarity independently of categorical membership [30,33,62,63]. Our data corroborate the hypothesis that these regions are also involved in conceptual retrieval [10,34] supporting representations related to visual experience that are not accessible to the blind.

The results of the adaptation analysis are crucial to support the hypothesis that conceptual retrieval consist on a partial replay of our perceptual experience [2,64], suggesting that visual features of objects and events, encoded in posterior occipital cortices and retrieved during conceptual processing, are not part of the conceptual representations of people that have never seen. Interestingly, our PPI analysis (see also Fig. S2) show that during conceptual processing the occipital cortices of blind are actively involved in conceptual retrieval too, paralleling the activation patterns of classic conceptual hubs such as the precuneus or the pMTG. This abnormal occipital activation in blind compared to sighted has been reported other times in the context of conceptual processing [19,43,59], and sometimes used as an argument against empiricist accounts: Not only visual cortices do not participate in conceptual retrieval in the sighted, but, if anything, they do it in the blind only [15,19]. In our study, we demonstrated that this finding is not at odds with empiricist arguments, showing that occipital cortices of both sighted and blind are involved in conceptual retrieval, but with different functions and probably at different levels of representation: Occipital cortex support sensorimotor simulations of visual features during conceptual retrieval in the sighted, whereas occipital cortices are re-organized to engage in more general processes related to conceptual retrieval in the blind (albeit these processes need to be better specified in future studies).

Another important result revealed by the adaptation analysis is that early blind participants showed greater repetition suppression in the posterior lateral-temporal cortices (PLTC), compared to sighted. Many studies before us have found that conceptually similar (e.g., dog - wolf) or semantically associated (e.g., dog - leash) words can lead to repetition suppression in PLTC [49–51]. Although it is still unclear what level of conceptual knowledge is represented in that region [49], there is some agreement that the PLTC stores auditory representations of words, that are connected to distributed semantic representations in the brain [49,65–67]. In this framework, the PLTC may work at the interface between wordforms and semantic knowledge [65,66], and a greater activity in the blind can index a larger use of verbal knowledge in this population [68,69] to compensate for the absence of visual information.

From a broader theoretical point of view, the results of this study are in line with a hierarchical model of conceptual representations based on progressive levels of abstraction [8–10,28,70]. At the top of the hierarchy, multimodal representations may co-exist with purely symbolic ones organized in a linguistic/propositional code [16]. It is possible that a large number of conceptual processes can take place involving only the higher-levels of representation, supported by conceptual hubs such as pMTG and precuneus [7,10,39], as well as category-sensitive regions in anterior VOTC [7,32] or language regions in temporal cortices [71,72]. Moreover, multimodal and abstract representations may be ideal to interact with lexical entries in the context of compositional [70], highly automatic [73] or shallow [74] semantic representations that are continuously required by natural language processing [70]. This level of representation can account for the obvious fact that congenitally blind people can think about colors and their perceptual properties [75,76] although they cannot simulate vision.

On the other hand, modality-specific simulation in sensory areas (e.g. visual, auditory, somatosensory), as the one highlighted by our adaptation analysis, may become central in deeper and more deliberate stages of conceptual processing, providing situated, specific and sometimes imaginistic representations. In this view, modality-specific simulations may be crucial to support the subjective experience of knowing providing the phenomenological qualia of conscious thinking [9,77]. Indeed, early blind people lack visual qualia, so that color and other visual features are necessarily represented as abstract entities or remapped on different senses [36]. In our study, we were able to show the neural basis of this phenomenological difference between the mental representations of sighted and early blind people.

Materials and methods

Participants

Thirty-six participants took part to this experiment: 18 early blinds (EB; 8F) and 18 sighted controls (SC; 8F). Participants were matched pairwise for gender, age, and years of education (Table S1).

All the blind participants lost sight at birth or before 3 years of age and all of them reported not having visual memories (Table S2). All participants were blindfolded during the task. The ethical committee of the Besta Neurological Institute approved this study (protocol fMRI_BP_001) and participants gave their informed consent before participation.

Stimuli

We selected six Italian color words (rosso/*red*, giallo/*yellow*, arancio/*orange*, verde/*green*, azzurro/*blue*, viola/*purple*), and six Italian action words (pugno/*punch*, graffio/*scratch*, schiaffo/*slap*, calcio/*kick*, salto/*jump*, corsa/*run*). Words were all highly

familiar nouns and were matched across categories (color, action), by number of letters (Color: mean= 5.83, sd= 0.98); Action: mean= 6, sd= 1.23), frequency (Zipf scale; Color: mean= 4.02, sd= 0.61; Action: mean= 4.18, sd= 0.4), and orthographic neighbors (Coltheart's N; Color: mean= 14, sd= 9.12; Action: mean= 15.33, sd= 12.42).

Auditory files were made using a voice synthesizer (talk to me), with a female voice, and edited into separated audio files with the same auditory properties (44100 Hz, 32 bit, mono, 78 dB of intensity). The original duration of each audio file (range 356 – 464 ms) was extended or compressed to 400 ms using the PSOLA (Pitch Synchronous Overlap and Add) algorithm and the sound-editing software Praat [78]. All the resulting audio files were highly intelligible.

Procedure

We designed a fast event-related fMRI paradigm during which participants listened to pairs of color and action words. In each trial the two words were played one after the other with a stimulus onset asynchrony (SOA) of 2000 ms.

The inter-trial interval ranged between 4000 and 16000 ms. Participants were asked to judge the similarity of the two colors or the two actions from 1 to 5 (1: very different, 5: very similar). Responses were collected via an ergonomic hand-shaped response box with five keys (Resonance Technology Inc.). All participants used their right hand to provide responses (thumb = very different, pinky = very similar). Participants were told that they had about 4 seconds to provide a response after the onset of the second word of the pair and they were encouraged to use all the scale (1 to 5). Furthermore, the instruction was to judge the similarity of colors and actions based on their perceptual properties (avoiding reference to emotion, valence, or other

non perceptual characteristics). Blind participants were told to judge color pairs on the basis of their knowledge about the perceptual similarity between colors.

Color and action words were presented in all possible within-category combination (15 color pairs, 15 action pairs). Each pair was presented twice in each run, in the two possible orders (e.g., red-yellow, yellow-red). Thus, there were 60 trials in each run and the experiment consisted in 5 runs of 7 minutes. Stimuli were pseudorandomized using optseq2 to optimize the sequence of presentation of the different conditions. Three different optimized lists of trials were used across runs. List order was counterbalanced across subjects.

One early blind was excluded from the analyses because the subject answered to less than 70% of the trials throughout the experiment due to sleepiness. One run of one sighted subject was excluded from the analysis because of a technical error during the acquisition, and two other runs (one in a sighted subject, one in a blind subject) were excluded since the subject answered to less than 70% of the trials in that specific run.

Conceptual similarity ratings

In order to perform the adaptation analysis, we divided the trials in similar pairs (e.g. red - orange) and different pairs (e.g. red - blue). We did so based on the participants' subjective ratings. For each participant we took the average rating for each of the 15 word-pairs in the action and color categories. Then we automatically divided the 15 pairs in 5 intervals (4 quantiles) of nearly equal size. This subdivision was performed using the function *quantile*, in R [79], that divides a probability distribution into contiguous intervals of equal probabilities (i.e., 20%). The pairs in the first two intervals were the different pairs (low ratings of similarity), the pairs in the 3rd interval were the medium pairs, and the pairs in the 4th and 5th intervals were the

similar pairs (See fig 2B). However, in some cases, ratings distributions were slightly unbalanced, due to the tendency of some subjects to find more "very different" pairs than "very similar" pairs. In these cases (8 subjects for action ratings [3 EB]; 4 subjects for Color Ratings [1 EB]), the automatic split in 5 equal interval was not possible. Thus, we set the boundary between the 2nd and 3rd interval at the ratings average (for that given subject), and set to the minimum (1 or 2, depending on the cases) the number of items in the 3rd interval (not analyzed), in order to balance as much as possible the number of pairs in the Different and Similar groups. This procedure made so that in these special cases (as well as in all the others), the rating values of different pairs were always below the mean, and the values of similar pairs was always above the mean. Fig. S3, S4, S5, S6, in the supplementary information, show subject-specific rating distributions.

MRI data acquisition

Brain images were acquired at the Neurological Institute Carlo Besta in Milano on a 3-Tesla scanner with a 32-channel head coil (Achieva TX; Philips Healthcare, Best, the Netherlands) and gradient echo planar imaging (EPI) sequences.

In the event-related experiment, we acquired 35 slices (voxel size 3 X 3 X 3.5) with no gap. The data in-plane matrix size were 64 X 64, field of view (FOV) 220mm X 220mm, time to repetition (TR)= 2 s, flip angle 90 degrees and time to echo (TE)= 30 ms. In all, 1210 whole-brain images were collected during the experimental sequence. The first 4 images of each run were excluded from the analysis for steady-state magnetization. Each participant performed 5 runs, with 242 volumes per run.

Anatomical data was acquired using a T1-weighted 3D-TFE sequence with the following parameters: 1 X 1 X 1 mm voxel size, 240 X 256 matrix size, 2.300 ms TR, 2.91 ms ET, 900 ms TI, 256 FoV, 160 slices.

MRI data analysis

We analyzed the fMRI data using SPM12 (www. fil.ion.ucl.ac.uk/spm/software/spm12/) and Matlab R2014b (The MathWorks, Inc.). **Preprocessing.** Preprocessing included slice timing correction of the functional time series [80], realignment of functional time series, coregistration of functional and anatomical data, spatial normalization to an echoplanar imaging template conforming to the Montreal Neurological Institute (MNI) space, and spatial smoothing [Gaussian kernel, 6 mm full-width at half-maximum (FWHM)]. Serial autocorrelation, assuming a first-order autoregressive model, was estimated using the pooled active voxels with a restricted maximum likelihood procedure, and the estimates were used to whiten the data and design matrices.

Data analysis. Following preprocessing steps, the analysis of fMRI data, based on a mixed-effects model, was conducted in two serial steps accounting, respectively, for fixed and random effects. In all the analysis the regressors for the conditions of interest consisted of a event-related boxcar function convolved with the canonical hemodynamic response function according to a variable epoch model [81]. Movement parameters derived from realignment of the functional volumes (translations in x, y, and z directions and rotations around x, y, and z axes), and a constant vector, were also included as covariates of no interest. We used a high-pass filter with a discrete cosine basis function and a cutoff period of 128 s to remove artifactual low-frequency trends.

Univariate analysis: For each subject, changes in regional brain responses were estimated through a general linear model including 2 regressors corresponding to the two categories Action and Color. The onset of each event was set at the beginning of the first word of the pair, the offset was determined by the subject response, thus included reaction time [81]. Linear contrasts tested for action-specific [Action > Color] and color-specific [Color > Action] BOLD activity.

These linear contrasts generated statistical parametric maps [SPM(T)]. The resulting contrast images were then further spatially smoothed (Gaussian kernel 5mm FWHM) and entered in a second-level analysis, corresponding to a random-effects model, accounting for inter-subject variance. One-sample t-tests were run on each group separately. Two-sample t-tests were then performed to compare these effects between groups (Blind vs Sighted) and to perform conjunction analyses to observe if the two groups presented similar activated networks for the two contrasts of interests.

Connectivity analysis: Psychophysiological interaction (PPI) analyses were computed to identify brain regions showing a significant change in the functional connectivity with seed regions (the right precuneus, the left pMTG and the rIPS) that showed a significant activation ($p<.001$, uncorrected) in the [(EB Conj. SC) X (Color > Action)] contrast, the [(EB Conj. SC) X (Action > Color)] contrast, and the [(SC > EB) X (Action > Color)] contrast respectively. In each individual, time series of activity (principal eigenvariate) were extracted from a 8 mm sphere centered on the nearest local maxima to the identified peaks in the second-level analysis (Note that centering the sphere on the peak itself does not change the ROI Analysis results, see Supplementary Information). New linear models were generated at the individual level, using three regressors. One regressor represented the psychological condition of interest (action or color trial). The second regressor was the physiological activity

extracted in the reference area. The third regressor represented the interaction of interest between the first (psychological) and the second (physiological) regressor. The design matrix also included movement parameters and a constant vector as regressors of no interest. A significant PPI indicated a change in the regression coefficients between any reported brain area and the seed region, related to the experimental conditions (Color>Action or Action>Color). Next, the individual summary statistic images obtained at the first-level (fixed-effects) analysis were spatially smoothed (5 mm FWHM Gaussian kernel) and entered in a second-level (random-effects) analysis using a two-sample t test contrasting the two groups.

Adaptation analysis: For each subject, the general linear model included 6 regressors corresponding to the 3 levels of similarity (different, medium, similar) in each condition (color, action). Color and Action pairs in the medium condition were modeled as regressors of no interest.

At the first level of analysis, linear contrasts tested for Repetition Suppression [Different > Similar] collapsing across categories (Action, Color). The same contrasts were then repeated within each category [Color Different > Color Similar; Action Different > Action Similar]. Finally, we tested for the Similarity by Category interactions, testing whether the adaptation was stronger in one category compared to the other (e.g., [Color Different > Color Similar] > [Action Different > Action Similar]).

These linear contrasts generated statistical parametric maps [SPM(T)]. The resulting contrast images were then further spatially smoothed (Gaussian kernel 5mm FWHM) and entered in a second-level analysis (RFX), corresponding to a random-effects model, accounting for inter-subject variance. One-sample t-tests were run on each group separately. Two-sample t-tests were then performed to compare these effects between groups (Blind vs Sighted).

Statistical inferences. At the whole brain level, statistical inference was made at a corrected cluster level of P < 0.05 FWE (with a standard voxel-level threshold of P < 0.001 uncorrected) and a minimum cluster-size of 50 voxels. ROI analysis based on Small Volume Correction were thresholded at p<0.05 FEW at the voxel level.

ROI definition and analysis. Occipital ROI for the PPI analyses were defined as following. Two peak-coordinates were taken from previous studies [42,43] showing the involvement of EB occipital areas in high-level functions such as language (left MOG [-36, -90, -1]) and mathematics (right MOG [33, -82, 9]). These areas also showed increased long range connectivity (in early blind) with frontal and parietal areas during rest [42–45].

The V4 and V5 ROI were drawn from the literature, considering both perceptual localizers, as well as evidence from semantic/conceptual task. We selected 3 peak coordinates for area V5. The first [-47, -78, -2] from a highly-cited study contrasting the perception of visual motion vs static images [53]. The second [-44, -74, 2] from a study [82] showing V5 sensitivity to motion sentences (e.g., "The wild horse crossed the barren field"). The third from a research on the on-line meta-analysis tool Neurosynth (http://neurosynth.org/) for the topic "action". In Neurosynth, the area in the occipital cortex with the highest action-related activation was indeed V5 (peak coordinates: -50, -72, 2). To avoid ROI proliferation, we averaged these 3 peak-coordinates in order to obtain a single peak (average peak: -47, -75, 1).

As for V4, we selected the color-sensitive occipital ROI considering perceptual localizers, as well as evidence of color-specific activity from semantic/conceptual task. Fernandino et al. [10] reported a color sensitive area in the left posterior collateral sulcus (ColS; at the intersection between the Lingual and the Fusyform gyrus; MNI peak coordinates: -16, -71, -12) associated with color-related

words. This peak is close to the posterior-V4 localization done by Beauchamps and colleagues (peak coordinates: -22, -82, -16) in a MRI version of the Farnsworth–Munsell 100-Hue Test [52]. A search in neurosynth with the keyword "color" also highlighted a left posterior color-sensitive region along the ColS with peak coordinates [-24, -90, -10]. We averaged these 3 peaks to find the center of our region of interest (average peak: -21, -81, -13).

The posterior lateral-temporal cortex ROI (PLTC) was taken from 3 studies showing semantic repetition suppression in that area. Bedny and colleagues [49] observed increased neural adaptation in PLTC (peak coordinates: 57, -36, 21) for repeated words (fan - fan), when the words were presented in a similar context (summer – fan; ceiling - fan), compared to when different context triggered different meanings (e.g., admirer – fan; ceiling – fan). This result conceptually replicated previous studies [50,51] showing semantic adaptation in the bilateral PLTC for related (e.g., dog - cat) vs unrelated (e.g., dog - apple) word pairs (peak coordinates: -42, -27, 9 and -51, -22, 8). These 3 peaks were averaged to find the center of our region of interest in both hemispheres (average peak: ±50, -28, 13).

All ROI analyses were performed using Small Volume Correction using spheres with a 10mm radius. Within the ROI, results were considered significant at a threshold of p<0.05, FEW-corrected. Here, and throughout the paper, brain coordinates are reported in MNI space.

Acknowledgement

This work was supported by a European Research Council starting grant (MADVIS grant #337573) attributed to OC. OC is a research associate at the Fond National de Recherche Scientifique de Belgique (FRS-FNRS). We wish to extend our gratitude to the Michela Picchetti, Mattia Verri and Alberto Redolfi for the technical

support during fMRI acquisition. We are also extremely thankful to our blind participants and the Unione Ciechi e Ipovedenti in Trento, Milano, Savona, Trieste, and the Blind Institute of Milano.

Author contributions:

R.B. and O.C. designed the research; R.B., S.F., A.N. and V.C. performed the research; R.B. analyzed the data in interaction with O.C.; R.B. and O.C. drafted the paper; all authors revised and edit the draft, and agreed on the final version of the manuscript.

References

1. Kosslyn, S. M., Ganis, G. & Thompson, W. L. Neural foundations of imagery. *Nat. Rev. Neurosci.* **2,** 635–642 (2001).

2. Barsalou, L. W. Perceptual Symbol Systems. *Behav. Brain Sci.* **30322,** 577– 660 (1999).

3. Fodor, J. *The language of thought.* (Harvard University Press, 1975).

4. Pylyshyn, Z. W. *Computation and cognition.* (MIT Press, 1984).

5. Lakoff, G. & Johnson, M. *Philosophy in the flesh.* (Basic Books, 1999).

6. Barsalou, L. W. Simulation, situated conceptualization, and prediction. *Philos. Trans. R. Soc. B Biol. Sci.* **364,** 1281–1289 (2009).

7. Binder, J. R. & Desai, R. H. The neurobiology of semantic memory. *Trends Cogn. Sci.* **15,** 527–536 (2011).

8. Barsalou, L. W. On Staying Grounded and Avoiding Quixotic Dead Ends. *Psychon. Bull. Rev.* **23,** 1122–1142 (2016).

9. Binder, J. R. *et al.* Toward a brain-based componential semantic representation. *Cogn. Neuropsychol.* **33,** 130–174 (2016).

10. Fernandino, L. *et al.* Concept representation reflects multimodal abstraction: A framework for embodied semantics. *Cereb. Cortex* 1–17 (2015). doi:10.1093/cercor/bhv020

11. Hsu, N. S., Frankland, S. M. & Thompson-Schill, S. L. Chromaticity of color perception and object color knowledge. *Neuropsychologia* **50,** 327–333 (2012).

12. Hsu, N. S., Kraemer, D. J. M., Oliver, R. T., Schlichting, M. L. & Thompson-schill, S. L. Color , Context , and Cognitive Style : Variations in Color Knowledge Retrieval as a Function of Task and Subject Variables. 2544–2557 (2011).

13. Simmons, W. K. *et al.* A common neural substrate for perceiving and knowing about color. *Neuropsychologia* **45,** 2802–2810 (2007).

14. Willems, R. M., Hagoort, P. & Casasanto, D. Body-specific representations of action verbs: neural evidence from right- and left-handers. *Psychol. Sci.* **21,** 67–74 (2010).

15. Bedny, M. & Saxe, R. Insights into the origins of knowledge from the cognitive neuroscience of blindness. *Cogn. Neuropsychol.* **29,** 56–84 (2012).

16. Mahon, B. Z. & Caramazza, A. A critical look at the embodied cognition hypothesis and a new proposal for grounding conceptual content. *J. Physiol. Paris* **102,** 59–70 (2008).

17. Ricciardi, E., Bonino, D., Pellegrini, S. & Pietrini, P. Mind the blind brain to understand the sighted one! Is there a supramodal cortical functional architecture? *Neurosci. Biobehav. Rev.* **41,** 64–77 (2014).

18. Casasanto, D. Different Bodies, Different Minds: The Body Specificity of Language and Thought. *Curr. Dir. Psychol. Sci.* **20,** 378–383 (2011).

19. Noppeney, U., Friston, K. J. & Price, C. J. Effects of visual deprivation on the

organization of the semantic system. *Brain* **126,** 1620–1627 (2003).

20. Peelen, M. V, He, C., Han, Z., Caramazza, A. & Bi, Y. Nonvisual and visual object shape representations in occipitotemporal cortex: evidence from congenitally blind and sighted adults. *J. Neurosci.* **34,** 163–70 (2014).

21. Peelen, M. V. *et al.* Tool selectivity in left occipitotemporal cortex develops without vision. *J. Cogn. Neurosci.* **25,** 1225–1234 (2013).

22. He, C. *et al.* Selectivity for large nonmanipulable objects in scene-selective visual cortex does not require visual experience. *Neuroimage* **79,** 1–9 (2013).

23. Bedny, M., Caramazza, a, Pascual-Leone, a & Saxe, R. Typical neural representations of action verbs develop without vision. *Cereb. Cortex* **22,** 286–93 (2012).

24. Dormal, G. *et al.* Functional Preference for Object Sounds and Voices in the Brain of Early Blind and Sighted Individuals. *J. Cogn. Neurosci.* **30,** 86–106 (2017).

25. van den Hurk, J., Van Baelen, M. & Op de Beeck, H. P. Development of visual category selectivity in ventral visual cortex does not require visual experience. *Proc. Natl. Acad. Sci.* 201612862 (2017). doi:10.1073/pnas.1612862114

26. Kassuba, T. *et al.* The left fusiform gyrus hosts trisensory representations of manipulable objects. *Neuroimage* **56,** 1566–1577 (2011).

27. Fernandino, L., Humphries, C. J., Conant, L. L., Seidenberg, M. S. & Binder, J. R. Heteromodal Cortical Areas Encode Sensory-Motor Features of Word Meaning. *J. Neurosci.* **36,** 9763–9769 (2016).

28. Martin, A. GRAPES-Grounding representations in action, perception, and emotion systems: How object properties and categories are represented in the human brain. *Psychon. Bull. Rev.* 979–990 (2015). doi:10.3758/s13423-015-0842-3

29. Grill-Spector, K. & Weiner, K. S. The functional architecture of the ventral temporal cortex and its role in categorization. *Nat. Rev. Neurosci.* **15,** 536–548 (2014).

30. Bracci, S. & Op de Beeck, H. Dissociations and associations between shape and category representations in the two visual pathways. *J. Neurosci.* **36,** 432–444 (2016).

31. Proklova, D., Kaiser, D. & Peelen, M. V. Disentangling Representations of Object Shape and Object Category in Human Visual Cortex: The Animate–Inanimate Distinction. *J. Cogn. Neurosci.* **28,** 680–692 (2016).

32. Peelen, M. V. & Downing, P. E. Category selectivity in human visual cortex: Beyond visual object recognition. *Neuropsychologia* **105,** 177–183 (2017).

33. Connolly, A. C. *et al.* The representation of biological classes in the human brain. *J. Neurosci.* **32,** 2608–18 (2012).

34. Borghesani, V. *et al.* Word meaning in the ventral visual path: a perceptual to conceptual gradient of semantic coding. *Neuroimage* **143,** 128–140 (2016).

35. Mitchell, T. M. *et al.* Predicting Human Brain Activity Associated with the Meanings of Nouns. *Science (80-.).* **320,** 1191–1195 (2008).

36. Connolly, A. C., Gleitman, L. R. & Thompson-Schill, S. L. Effect of congenital blindness on the semantic representation of some everyday concepts. *Proc. Natl. Acad. Sci. U. S. A.* **104,** 8241–6 (2007).

37. Friston, K. J. *et al.* Psychophysiological and modulatory interactions in neuroimaging. *Neuroimage* **6,** 218–229 (1997).

38. O'Reilly, J. X., Woolrich, M. W., Behrens, T. E. J., Smith, S. M. & Johansen-berg, H. Tools of the trade: Psychophysiological interactions and functional connectivity. *Soc. Cogn. Affect. Neurosci.* **7,** 604–609 (2012).

39. Binder, J. R., Desai, R. H., Graves, W. W. & Conant, L. L. Where is the

semantic system? A critical review and meta-analysis of 120 functional

neuroimaging studies. *Cereb. Cortex* **19,** 2767–2796 (2009).

40. Amedi, A., Raz, N., Pianka, P., Malach, R. & Zohary, E. Early 'visual' cortex

    activation correlates with superior verbal memory performance in the blind.

    Nat. Neurosci. **6,** 758–66 (2003).

41. Burton, H. Visual cortex activity in early and late blind people. *J. Neurosci.* **23,**

    4005–4011 (2003).

42. Kanjlia, S., Lane, C., Feigenson, L. & Bedny, M. Absence of visual experience

    modifies the neural basis of numerical thinking. *Proc. Natl. Acad. Sci.* **113,**

    201524982 (2016).

43. Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E. & Saxe, R.

    Language processing in the occipital cortex of congenitally blind adults. *Proc.*

    Natl. Acad. Sci. **108,** 4429–4434 (2011).

44. Liu, Y. *et al.* Whole brain functional connectivity in the early blind. *Brain* **130,**

    2085–2096 (2007).

45. Collignon, O. *et al.* Impact of blindness onset on the functional organization

    and the connectivity of the occipital cortex. *Brain* **136,** 2769–2783 (2013).

46. Barron, H. C., Garvert, M. M. & Behrens, T. E. J. Repetition suppression: a

    means to index neural representations using BOLD? *Philos. Trans. R. Soc. B*

    Biol. Sci. **371,** 20150355 (2016).

47. Wheatley, T., Weisberg, J., Beauchamp, M. S. & Martin, A. Automatic priming

    of semantically related words reduces activity in the fusiform gyrus. *J. Cogn.*

    Neurosci. **17,** 1871–85 (2005).

48. Grill-Spector, K., Henson, R. & Martin, A. Repetition and the brain: Neural

    models of stimulus-specific effects. *Trends Cogn. Sci.* **10,** 14–23 (2006).

49. Bedny, M., McGill, M. & Thompson-Schill, S. L. Semantic adaptation and

competition during word comprehension. *Cereb. Cortex* **18,** 2574–2585

(2008).

50.   Wible, C. G. *et al.* Connectivity among semantic associates: An fMRI study of

semantic priming. *Brain Lang.* **97,** 294–305 (2006).

51.   Kotz, S. A., Cappa, S. F., Cramon, D. Y. Von & Friederici, A. D. Modulation of

the Lexical – Semantic Network by Auditory Semantic Priming : An Event-

Related Functional MRI Study. *Neuroimage* **17,** 1761–1772 (2002).

52.   Beauchamp, M. S., Haxby, J. V, Jennings, J. E. & DeYoe, E. A. An fMRI

version of the farnsworth-munsell 100-hue test reveals multiple color-selective

areas in human ventral occipitotemporal cortex. *Cereb. Cortex* **9,** 257–263

(1999).

53.   Dumoulin, S. O. *et al.* A new anatomical landmark for reliable identification of

human area V5/MT: a quantitative analysis of sulcal patterning. *Cereb. Cortex*

10, 454–463 (2000).

54.   Cheadle, S. W. & Zeki, S. The role of parietal cortex in the formation of color

and motion based concepts. *Front. Hum. Neurosci.* **8,** 535 (2014).

55.   Zeki, S. & Stutters, J. Functional specialization and generalization for grouping

of stimuli based on colour and motion. *Neuroimage* **73,** 156–166 (2013).

56.   Xu, Y. The Role of the Superior Intraparietal Sulcus in Supporting Visual Short-

Term Memory for Multifeature Objects. *J. Neurosci.* **27,** 11676–11686 (2007).

57.   Swisher, J. D., Halko, M. A., Merabet, L. B., McMains, S. A. & Somers, D. C.

Visual Topography of Human Intraparietal Sulcus. *J. Neurosci.* **27,** 5326–5337

(2007).

58.   Grill-Spector, K. The neural basis of object perception. *Curr. Opin. Neurobiol.*

13, 159–166 (2003).

59.   Burton, H., Diamond, J. B. & McDermott, K. B. Dissociating cortical regions

activated by semantic and phonological tasks: a FMRI study in blind and sighted people. *J. Neurophysiol.* **90,** 1965–82 (2003).

60. Van Ackeren, M. J., Barbero, F., Mattioni, S., Bottini, R. & Collignon, O. Neuronal populations in the occipital cortex of the blind synchronize to the temporal dynamics of speech. *Elife* 1–45 (2018). doi:10.7554/eLife.31640

61. Crollen, V. *et al.* Recruitment of occipital cortex by arithmetic processing follows computational bias in early blind. *bioRxiv Prepr.* (2018).

62. Kriegeskorte, N. *et al.* Matching Categorical Object Representations in Inferior Temporal Cortex of Man and Monkey. *Neuron* **60,** 1126–1141 (2008).

63. Naselaris, T., Prenger, R. J., Kay, K. N., Oliver, M. & Gallant, J. L. Bayesian Reconstruction of Natural Images from Human Brain Activity. *Neuron* **63,** 902–915 (2009).

64. Harnad, S. The symbol grounding problem. *Phys. D* **42,** 335–346 (1990).

65. Hickok, G. & Poeppel, D. The cortical organization of speech processing. *Nat. Rev. Neurosci.* **8,** 393–402 (2007).

66. Gow, D. W. The cortical organization of lexical knowledge: A dual lexicon model of spoken language processing. *Brain Lang.* **121,** 273–288 (2012).

67. Mirman, D. *et al.* Neural organization of spoken language revealed by lesion-symptom mapping. *Nat. Commun.* **6,** 1–9 (2015).

68. Cattaneo, Z. *et al.* Imagery and spatial processes in blindness and visual impairment. *Neurosci. Biobehav. Rev.* **32,** 1346–1360 (2008).

69. Crollen, V. *et al.* Visual experience influences the interactions between fingers and numbers. *Cognition* **133,** 91–96 (2014).

70. Binder, J. R. In defense of abstract conceptual representations. *Psychon. Bull. Rev.* **23,** 1096–1108 (2016).

71. Anderson, A. J., Bruni, E., Lopopolo, A., Poesio, M. & Baroni, M. Reading

visually embodied meaning from the brain: Visually grounded computational models decode visual-object mental imagery induced by written text. *Neuroimage* **120,** 309–322 (2015).

72. Anderson, A. J. *et al.* Predicting neural activity patterns associated with sentences using a neurobiologically motivated model of semantic representation. *Cereb. Cortex* **27,** 4379–4395 (2017).

73. Bottini, R., Bucur, M. & Crepaldi, D. The nature of semantic priming by subliminal spatial words. Embodied or disembodied? *J. Exp. Psychol. Gen.* **145,** 1160–1176 (2016).

74. Ostarek, M. & Huettig, F. A Task-Dependent Causal Role for Low-Level Visual Processes in Spoken Word Comprehension. *J. Exp. Psychol. Learn. Mem. Cogn.* 1–32 (2017). doi:http://dx.doi.org/10.1037/xlm0000375

75. Barilari, M., de Heering, A., Crollen, V., Collignon, O. & Bottini, R. Is Red Heavier Than Yellow Even for Blind? *Iperception.* **9,** 204166951875912 (2018).

76. Marmor, G. Age at onset of blindness and the development of the semantics of color names. *J. Exp. Child Psychol.* **278,** 344–345 (1978).

77. Koch, C., Massimini, M., Boly, M. & Tononi, G. Neural correlates of consciousness: progress and problems. *Nat. Rev. Neurosci.* **17,** 307–321 (2016).

78. Boersma, P. & Weenink, D. Praat: doing phonetics by computer. (2018). at <http://www.praat.org/>

79. R Core Team. R: A language and environment for statistical computing. (2013).

80. Sladky, R. *et al.* Slice-timing effects and their correction in functional MRI. *Neuroimage* **58,** 588–594 (2011).

81. Grinband, J., Wager, T. D., Lindquist, M., Ferrera, V. P. & Hirsch, J. Detection of time-varying signals in event-related fMRI designs. *Neuroimage* **43,** 509–520 (2008).

82. Saygin, A. P., McCullough, S., Alac, M. & Emmorey, K. Modulation of BOLD response in motion-sensitive lateral temporal cortex by real and fictive motion sentences. *J. Cogn. Neurosci.* **22,** 2480–90 (2010).

83. Martin, A., Haxby, J. & Lalonde, F. Discrete Cortical Regions Associated with Knowledge of Color and Knowledge of Action. *Science (80-.).* **270,** 102–105 (1995).

84. Noppeney, U. & Price, C. J. Retrieval of visual, auditory, and abstract semantics. *Neuroimage* **15,** 917–926 (2002).

85. Pulvermüller, F. & Hauk, O. Category-specific conceptual processing of color and form in left fronto-temporal cortex. *Cereb. Cortex* **16,** 1193–201 (2006).

86. Noppeney, U. & Price, C. J. Functional imaging of the semantic system: Retrieval of sensory-experienced and verbally learned knowledge. *Brain Lang.* **84,** 120–133 (2001).

Figures

Figure 1

Figure 1. Classic univariate analysis and psychophysiological interactions. Regional BOLD responses are depicted over multiplanar slices and renders of the MNI-ICBM152 template. (A) Suprathreshold cluster (P<0.05 FEW corrected) showing common activity in the lpMTG for the contrast Action > Color in both sighted and early blind (conj., conjunction analysis); (B) Suprathreshold cluster (P<0.05 FEW corrected) showing greater activity in the rIPS, in sighted compared to early blind, for the contrast Color > Action; (C) Suprathreshold cluster (P<0.001 uncorrected; for illustrative purposes only) showing common activity in the precuneus for the contrast Color > Action in both sighted and early blind (conj., conjunction analysis); (D) Suprathreshold clusters (P<0.005 uncorrected; for illustrative purposes only) showing greater connectivity in the occipital areas of early blind people with the lpMTG (PPI for the contrast Action > Color); (E) Barplot (for illustrative purposes only) showing beta weights derived from PPI analysis, with seed in lpMTG, in sighted (gray) and blind (white), in the right and left middle occipital gyrus (MOG) for the contrast Action > Color (arbitrary unit ± SEM); (F) Suprathreshold clusters (P<0.005 uncorrected; for

illustrative purposes only) showing greater connectivity in the occipital areas of early blind people with the precuneus (PPI for the contrast Color > Action); (G) Barplot (for illustrative purposes only) showing beta weights derived from PPI analysis, with seed in precuneus, in sighted (gray) and blind (white), in the right and left middle occipital gyrus (MOG) for the contrast Color > Action (arbitrary unit ± SEM);

Figure 2

Figure 2. Adaptation, behavioral analysis. (A) Similarity judgments were highly correlated across groups both for actions and color; (B) Conceptual schema of the division of word pairs in "different" and "similar" based on subjective similarity ratings; (C) Barplot depicting the average number of items in the "different", "medium" and "similar" categories. The number of items in the "different" and "similar" categories is very similar across groups (number of trials ± SEM); (D) Barplot depicting the average reaction time in the "different", "medium", and "similar", the average RTs of the "different" and "similar" categories is very similar across groups (seconds ± SEM).

Figure 3

Figure 3. Adaptation, fMRI results. Regional BOLD responses are depicted over renders of the MNI-ICBM152 template. (A) Suprathreshold clusters showing neural adaptation for similar word pairs in the occipital cortices of sighted participants; (B) Suprathreshold clusters showing neural adaptation for similar word pairs in the temporal and somatosensory-motor cortices of early blind participants; (C) Suprathreshold clusters showing neural adaptation for similar word pairs in sighted compared to blind, and (D) blind compared to sighted; (E) Suprathreshold cluster showing neural adaptation for similar color-word pairs in the sighted, with regional activity localized in the left PCoS/V4; (F) Suprathreshold clusters showing neural adaptation for similar action-word pairs in the sighted, with regional activity spread in different occipital areas including the left PCoS/V4 and the left V5. Cluster threshold at $P<0.005$ uncorrected, for illustration only.

Supplementary Information for

Different neural networks for conceptual retrieval in sighted and blind reveal

the experiential bases of knowledge.

Roberto Bottini, Stefania Ferraro, Anna Nigri, Valeria Cuccarini, Maria Grazia

Bruzzone , Olivier Collignon

Email: bottini.r@gmail.com

This PDF file includes:

> Figs. S1 to S6
> Tables S1 to S5
> Supplementary text

Table S1. Demographic information of sighted and blind participants.

Sight Status	Age	Gender	Age of formal education	Sight Status	Age	Gender	Age of formal education
EB01	32	M	18	SC01	37	M	18
EB02	52	F	18	SC02	46	F	18
EB03	30	M	13	SC03	34	M	18
EB04	29	F	13	SC04	26	F	16
EB05	36	F	18	SC05	33	F	18
EB06	27	F	18	SC06	27	F	18
EB07	39	M	18	SC07	40	M	18
EB08	48	M	8	SC08	51	M	8
EB09	42	M	13	SC09	41	M	18
EB10	36	M	18	SC10	40	M	18
EB11	49	M	18	SC11	43	M	18
EB12	36	F	18	SC12	39	F	18
EB13	45	M	18	SC13	42	M	18
EB14	25	F	13	SC14	25	F	16
EB15	29	F	16	SC15	30	F	18
EB16	30	M	13	SC16	26	M	13
EB17	28	F	18	SC17	31	F	18
EB18	45	M	8	SC18	39	M	13
M	*36,56*		*15,39*		*36,11*		*16,67*
SD	*8,48*		*3,48*		*7,52*		*2,72*

Note. M = male; F = female; EB=early blind; SC=sighted control; M=mean; SD=standard deviation.

Table S2. Early blind participants, additional information

Subject	Handedness	Age onset blindness	Cause of blindness
EB01	32	0	Optic nerve Hypoplasia
EB02	52	1	Retinoblastoma
EB03	30	0	Congenital retinal dystrophy
EB04	29	2	Retinoblastoma
EB05	36	0	Congenital Microphtalmia
EB06	27	0	Congenital Microphtalmia
EB07	39	0	Retrolental fibroplasia
EB08	48	0	Optic nerve atrophy
EB09	42	0	Congenital retinal dystrophy
EB10	36	0	Retrolental fibroplasia
EB11	49	0	Retrolental fibroplasia
EB12	36	0	Retrolental fibroplasia
EB13	45	0	Retrolental fibroplasia
EB14	25	0	Leber's congenital amaurosis
EB15	29	3	Retinitis pigmentosa
EB16	30	0	Retrolental fibroplasia
EB17	28	0	Agenesis of the optic nerve
EB18	45	0	Glaucoma

—

Table S3. Regional responses for the comparison between action and color trials across sighted and early blind.

Area	k	Xmm	Ymm	Zmm	Z	P_{FWE}
Sighted, action > color						
L middle temporal gyrus	1134	-57	-61	8	5.80	<.001
L inferior frontal gyrus	S.C.	-48	29	-1	5.31	
L middle temporal gyrus	S.C.	-60	-43	29	5.15	
R middle temporal gyrus	474	60	-10	-7	5.14	<.001
	S.C.	57	2	-13	4.96	
R superior temporal gyrus	S.C.	51	-28	-1	4.74	
Sighted, color > action						
R orbital gyrus	65	27	35	-13	4.98	.009*
L orbital gyrus	71	-30	35	-13	4.90	.012*
precuneus	91	0	-61	29	4.11	.06 ‡
Blind, action > color						
L middle temporal gyrus	315	-54	-61	5	4.69	<.001
	S.C.	-42	-67	17	4.46	
	S.C.	-45	-52	14	3.49	
R calcarine/post. cingulate	325	9	-67	8	4.63	<.001
	S.C.	0	-76	11	4.07	
L calcarine	S.C.	-9	-82	11	4.07	
R middle temporal gyrus	203	57	-55	5	4.40	.003
	S.C.	42	-55	8	3.94	
	S.C.	63	-43	8	3.63	
R inferior frontal gyrus	256	42	20	5	4.21	<.001
R superior temporal pole	S.C.	48	11	-19	3.97	
R inferior frontal gyrus	S.C.	51	17	-1	3.80	
Blind, color > action						
R precuneus	109	6	-52	20	4.72	.036
L medial frontal gyrus	174	-9	59	-7	4.20	.008
	S.C.	6	53	-10	3.85	
Sighted ∩ Bind, action > color						
L middle temporal gyrus	155	-54	-61	5	4.69	.01
Sighted > Bind, color > action						
R intraparietal sulcus	146	33	-43	35	4.33	.013

Significance corrections are reported at the cluster level, unless otherwise specified; cluster size threshold = 50. FWE, family-wise error; L, left; R, right; S.C., same cluster.
*Brain activations significant after FWE correction at voxel-level over the whole brain.   ‡ Indicates marginally significant clusters ($P_{fwe} < 0.1$)

Figure S1. Regional BOLD responses are depicted over multiplanar slices and renders of the MNI-ICBM152 template. Suprathreshold cluster (P<0.05 FEW corrected) showing brain activity for the contrasts (A) Action > Color in SC; (B) Action > Color in EB; (C) Action > Color & SC > EB, with no significant cluster; (D) Action > Color & EB > SC, with no significant cluster; (E) Color > Action in SC; (F) Color > Action in EB; (G) Color > Action & SC > EB; (H)) Color > Action & EB > SC.

Table S4. Regional responses for psychophysiological interactions with seeds in lpMTG and Precuneus, across sighted and early blind.

Area	k	Xmm	Ymm	Zmm	Z	P_{FWE}
Seed in lpMTG						
Sighted						
L inferior parietal cortex	174	-45	-40	41	4.12	.006
	S.C.	-36	-52	44	3.94	
	S.C.	-30	-61	38	3.94	
Blind						
L middle occipital gyrus	75	-30	-91	-7	3.36	<.015^
Blind > Sighted						
L middle occipital gyrus	56	-36	-85	-7	3.28	.02^
R middle occipital gyrus	33	27	-82	14	3.17	.026^
Seed in rPrecuneus						
Blind						
L middle frontal gyrus	86	-42	50	5	4.76	.023*
L middle occipital gyrus	234	-24	-76	26	4.55	.001
L superior parietal lobe	S.C.	-30	-61	41	3.78	
L superior parietal lobe	S.C.	-30	-91	14	3.31	
R middle occipital gyrus	537	39	-85	20	3.78	<.001
	S.C.	33	-88	2	3.69	
R fusiform gyrus	S.C.	24	-52	-16	3.75	
Blind > Sighted						
L middle occipital gyrus	159	-27	-79	23	4.44	.007
	S.C.	-30	-91	17	3.63	
	S.C.	-42	-82	20	3.20	
R middle occipital gyrus	170	39	-85	20	3.95	<.001
	S.C.	39	-85	5	3.59	
R superior occipital gyrus	S.C.	24	-88	23	3.45	

Significance corrections are reported at the cluster level, unless otherwise specified; cluster size threshold = 50. FWE, family-wise error; L, left; R, right; S.C., same cluster.
*Brain activations significant after FWE voxel-level correction over the whole brain. ^Brain activation significant after FWE voxel correction over a small spherical volume (10-mm radius) at peak coordinates for right/left MOG.

Table S5. Regional responses for adaptation analysis (repetition suppression), across sighted and early blind.

Area	k	Xmm	Ymm	Zmm	Z	P$_{FWE}$
Sighted, different > similar						
L lingual gyrus	180	-24	-70	-7	4.33	.013
	S.C.	-15	-70	-10	4.32	
Blind, different > similar						
L middle temporal gyrus	119	-60	-10	-7	4.20	.049
	S.C.	-63	-22	2	3.90	
L superior temporal gyrus	S.C.	-51	-13	-1	3.39	
R superior temporal gyrus	323	57	-28	8	4.23	.001
	S.C.	48	-37	20	3.92	
	S.C.	54	-22	2	3.92	
L Postcentral gyrus	105	-48	-13	38	4.12	.07‡
R Precentral gyrus	121	27	-25	56	3.71	.02
	S.C.	36	-22	62	3.70	
	S.C.	36	-16	44	3.45	
Sighted > Blind, diff. > similar						
L superior occipital gyrus	141	-24	-91	26	4.35	.03
	S.C.	-9	-97	20	3.77	
L middle occipital gyrus	S.C.	-15	-100	14	3.75	
R middle occipital gyrus	250	27	-85	11	3.97	.003
	S.C.	36	-79	2	3.67	
R superior occipital gyrus	S.C.	27	-79	23	3.79	
L lingual gyrus	165	-24	-70	-7	3.91	.017
L middle occipital gyrus	S.C.	-39	-73	2	3.48	
L middle occipital gyrus	S.C.	-27	-94	2	3.25	
Blind > Sighted, diff. > similar						
L superior temporal gyrus		-45	-31	20	2.99	.035^
R superior temporal gyrus		45	-28	11	3.13	.024^

Significance corrections are reported at the cluster level, unless otherwise specified; cluster size threshold = 50. FWE, family-wise error; L, left; R, right; S.C., same cluster. ^Brain activation significant after FWE voxel correction over a small spherical volume (10-mm radius) at peak coordinates for right/left PLTC. ‡ Indicates marginally significant clusters (P$_{fwe}$<0.1)

Figure S2. Whole brain results for the contrast Stimulation > Rest in blind

more than sighted. FWE corrected at voxel level (p< .05).

Figure S3. Action pairs divided by Similarity ratings in Sighted. In blue dissimilar, green medium and red similar pairs. Dashed line is the average rating for that given subject. Note that dissimilar pairs are on average below average and similar pairs above average.

Figure S4. Action pairs divided by Similarity ratings in Blind. In blue dissimilar, green medium and red similar pairs. Dashed line is the average rating for that given subject. Note that dissimilar pairs are on average below average and similar pairs above average.

Figure S5. Color pairs divided by Similarity ratings in Sighted. In blue dissimilar, green medium and red similar pairs. Dashed line is the average rating for that given subject. Note that dissimilar pairs are on average below average and similar pairs above average.

Figure S6. Color pairs divided by Similarity ratings in Blind. In blue dissimilar, green medium and red similar pairs. Dashed line is the average rating for that given subject. Note that dissimilar pairs are on average below average and similar pairs above average.

Additional ROI analysis for PPI   51

For the original PPI analysis, in each individual, time series of activity (principal eigenvariate) were extracted from a 8 mm sphere centered on the nearest local maxima to the identified peaks in the second-level analysis (seed region). However, centering the sphere on the peak itself does not change the ROI Analysis results:

PPI with seed in the lpMTG revealed an increase of action-selective functional connectivity in both occipital ROIs of blind people compared to their sighted counterpart (lMOG: t= 3.33, p= 0.034; rMOG: t=3.21, p=0.043). Similarly, PPI with seed in the precuneus revealed an increase of color-selective functional connectivity in the occipital cortices of blind compared to sighted participants (lMOG: t= 3.44, p= 0.028; rMOG: t=3.81, p=0.012).