Different neural networks for conceptual retrieval in sighted and blind

Roberto Bottini[1,2]*, Stefania Ferraro[3], Anna Nigri[3], Valeria Cuccarini[3], Maria

Grazia Bruzzone[3], Olivier Collignon[1,4]*

[1] Center for Mind/Brain Studies (CIMeC), University of Trento, Trento, Italy (38123)

[2] International School for Advanced Studies (SISSA), Trieste, Italy (34136)

[3] Neuroradiology Department, Fondazione IRCCS Istituto Neurologico Carlo Besta,
Milan, Italy (20133)

[4] Institute of research in Psychology (IPSY) & Institute of Neuroscience (IoNS) -
University of Louvain (UCL), Louvain, Belgium (1348)

*Correspondence to:
Roberto Bottini; University of Trento (CIMeC)
Via delle regole 101, 38123, Mattarello, Italy
Email: bottini.r@gmail.com
15
16 Olivier Collignon; University of Louvain (UcL)
17 10, Place du Cardinal Mercier; 1348 Louvain-La-Neuve \ Belgium
18 Email: olivier.collignon@uclouvain.be
19

 # Pages: 37

 # Words: Abstract (193); Introduction (642); Discussion (1469).

Acknowledgement: This work was supported by a European Research Council
starting grant (MADVIS grant #337573) attributed to OC. OC is a research associate
at the Fond National de Recherche Scientifique de Belgique (FRS-FNRS). We wish to
extend our gratitude to the Michela Picchetti, Mattia Verri and Alberto Redolfi for the
technical support during fMRI acquisition. We are also extremely thankful to our blind
participants and the Unione Ciechi e Ipovedenti in Trento, Milano, Savona, Trieste,
and the Blind Institute of Milano. We also thank Yanchao Bi and Xiaoying Wang for
sharing brain maps from their previously published data.
30

31 **Author contributions:** R.B. and O.C. designed the research; R.B., S.F., A.N. and V.C.
32 performed the research; R.B. analyzed the data in interaction with O.C.; R.B. and O.C.
33 drafted the paper; all authors revised and edit the draft, and agreed on the final version
34 of the manuscript.
35

Abstract

We investigated the experiential bases of knowledge by asking whether people that

perceive the world in a different way also show a different neurobiology of concepts.

We characterized the brain activity of early-blind and sighted individuals during a

conceptual retrieval task in which participants rated the perceptual similarity between

color and action concepts evoked by spoken words. Adaptation analysis showed that

word-pairs referring to perceptually similar colors (e.g., red-orange) or actions (e.g.,

run-jump) led to repetition-suppression in occipital visual regions in the sighted,

regions that are known to encode visual features of objects and events, independently

of their category. Early blind showed instead adaptation for similar concepts in

language-related regions, but not in occipital cortices. Further analysis contrasting the

two categories (color and action), independently of item similarity, activated category-

sensitive regions in the pMTG (for actions) and the precuneus (for color) in both sighted

and blind. These two regions, however, showed a different connectivity profile as a

function of visual deprivation, increasing task-dependent connectivity with reorganized

occipital regions in the early blind. Overall, our results show that visual deprivation

changes the neural bases of conceptual retrieval, which is partially grounded in

sensorimotor experience.

Keywords: Concepts; Color; Action; Blindness; Neuroplasticity; fMRI; Adaptation

Significance Statement:

Do people with different sensory experience conceive the world differently? We

tested whether conceptual knowledge builds on sensory experience by looking at the

neurobiology of concepts in early blind individuals. Participants in fMRI heard pairs of

words referring to colors (e.g., green-blue) or actions (e.g., jump-run) and rated their

perceptual similarity. Perceptual similarity of colors and actions was represented in

occipital visual regions in the sighted, but in language-related regions in the blind.

Occipital regions in the blind, albeit not encoding perceptual similarity, were however

recruited during conceptual retrieval, working in concert with classic semantic hubs

such as the Precuneus and the lpMTG. Overall, visual deprivation changes the neural

bases of conceptual processing, which is partially grounded in sensorimotor

experience.

Introduction

As we think, we navigate and retrieve conceptual knowledge, but the nature of

this knowledge is highly debated. One of the major sources of disagreement is whether

thinking is more similar to the analog replay of our experience (Barsalou, 1999;

Kosslyn, Ganis, & Thompson, 2001), or the symbolic computations of a Turing

machine (Fodor, 1975; Pylyshyn, 1984).

Congenital blindness offers an ideal model to test whether thinking is simulating

our own experience of the world (Barsalou, 2009; Lakoff & Johnson, 1999). If

conceptual processing is largely grounded into experience, blind people, who

experience the world in a different way, should also show a different neurobiology of

concepts, at least in part (Casasanto, 2011). Several studies, however, seem to

provide evidence against this idea (Bedny & Saxe, 2012). For instance, when sighted

and blind were asked to retrieve information about highly visual entities, knowledge

about small and manipulable objects (Peelen et al., 2013) activated the lateral

temporal-occipital complex; thinking about big non-manipulable objects activated the

parahipoccampal place area (He et al., 2013); and processing action verbs (compared

to nouns) activated the left posterior middle temporal gyrus (Bedny et al. 2012) in both

groups. These results seem to suggest that blindness leaves the neurobiology of

conceptual retrieval largely unchanged, and that experience plays a minor role in

shaping mental representations (Bedny & Saxe, 2012; Uta Noppeney et al., 2003).

Alternatively, it is possible that previous studies relied on paradigms designed

to investigate categorical knowledge, a level of processing that is relatively resilient to

visual deprivation (Van Baelen, and Op de Beeck 2017). Indeed, categorical

boundaries (e.g., between objects, tools or animals) are drawn based on several

dimensions that exceed visual appearance (e.g., tokens from the same category do

not always look alike; Bracci and Op de Beeck 2016; Proklova, Kaiser, and Peelen

2016), and depend on functional, emotional and linguistic constraints (Peelen &

Downing, 2017). Indeed, brain regions that show category specificity, seem to

represent different categories in a format that is at least partially independent from

visual appearance (Bracci & Op de Beeck, 2016), and thus more likely to be resilient

to visual deprivation (Bedny et al. 2012; van den Hurk, Van Baelen, and Op de Beeck

2017; Wang et al. 2015).

Differences between sighted and blind may be more likely to emerge when

participants process the perceptual similarity of retrieved concepts (Martin, Douglas,

Newsome, Man, & Barense, 2018), a process known to be encoded in more posterior

(visual) occipital regions in the sighted (Fernandino et al. 2015; Borghesani et al. 2016;

Mitchell et al. 2008). To test this hypothesis we asked sighted and early blind

participants to rate pairs of concepts based on their perceptual similarity and analyzed

the data using a repetition suppression framework (Barron, Garvert, and Behrens

2016; Wheatley et al. 2005; Grill-Spector, Henson, and Martin 2006). We reasoned

that this method will allow to tag selectively the representation of low-level perceptual

features during conceptual retrieval (Horner & Henson, 2011; Mohr, Linder, Linden,

Kaiser, & Sireteanu, 2009). We predicted that perceptually similar concepts (within a

given category) might elicit neural adaptation in the occipital cortex of sighted (who will

retrieve their visual similarity), but not blind people. On the other hand, a direct

comparison between categories (independently of visual/perceptual similarity) should

highlight category-specific responses in more anterior/multimodal areas of the brain,

with limited differences between sighted and blind. This should be especially true for

conceptual categories that can be perceptually experienced both by sighted and blind

individuals. Therefore, in our experiment, we choose stimuli exemplars coming from

actions and color categories. Including colors, that can be experienced through vision

only, allowed us to test whether their different epistemological status (concrete vs

abstract), in sighted and blind, would influence their representation even in multimodal

areas of the brain that usually show resilience to visual deprivation (Striem-Amit,

Wang, Bi, & Caramazza, 2018).

Materials and methods

Participants

Thirty-six participants took part to this experiment: 18 early blinds (EB; 8F) and

18 sighted controls (SC; 8F). Participants were matched pairwise for gender, age, and

years of education (Table S1).

All the blind participants lost sight at birth or before 3 years of age and all of

them reported not having visual memories (Table S2). All participants were blindfolded

during the task. The ethical committee of the Besta Neurological Institute approved

this study (protocol fMRI_BP_001) and participants gave their informed consent before

participation.

Stimuli

We selected six Italian color words (rosso/*red*, giallo/*yellow*, arancio/*orange*,

verde/*green*, azzurro/*blue*, viola/*purple*), and six Italian action words (pugno/*punch*,

graffio/*scratch*, schiaffo/*slap*, calcio/*kick*, salto/*jump*, corsa/*run*). Words were all highly

familiar nouns and were matched across categories (color, action), by number of letters

(Color: mean= 5.83, sd= 0.98); Action: mean= 6, sd= 1.23), frequency (Zipf scale;

Color: mean= 4.02, sd= 0.61; Action: mean= 4.18, sd= 0.4), and orthographic

neighbors (Coltheart's N; Color: mean= 14, sd= 9.12; Action: mean= 15.33, sd= 12.42).

Auditory files were made using a voice synthesizer (talk to me), with a female

voice, and edited into separated audio files with the same auditory properties (44100

Hz, 32 bit, mono, 78 dB of intensity). The original duration of each audio file (range

356 – 464 ms) was extended or compressed to 400 ms using the PSOLA (Pitch

Synchronous Overlap and Add) algorithm and the sound-editing software Praat

(Boersma & Weenink, 2018). All the resulting audio files were highly intelligible.

Experimental Design

We designed a fast event-related fMRI paradigm during which participants

listened to pairs of color and action words. In each trial the two words were played one

after the other with a stimulus onset asynchrony (SOA) of 2000 ms.

The inter-trial interval ranged between 4000 and 16000 ms. Participants were

asked to judge the similarity of the two colors or the two actions from 1 to 5 (1: very

different, 5: very similar). Responses were collected via an ergonomic hand-shaped

response box with five keys (Resonance Technology Inc.). All participants used their

right hand to provide responses (thumb = very different, pinky = very similar).

Participants were told that they had about 4 seconds to provide a response after the

onset of the second word of the pair and they were encouraged to use all the scale (1

to 5). Furthermore, the instruction was to judge the similarity of colors and actions

based on their perceptual properties (avoiding reference to emotion, valence, or other

non perceptual characteristics). Blind participants were told to judge color pairs on the

basis of their knowledge about the perceptual similarity between colors.

Color and action words were presented in all possible within-category

combinations (15 color pairs, 15 action pairs). Each pair was presented twice in each

run, in the two possible orders (e.g., red-yellow, yellow-red). Thus, there were 60 trials

in each run and the experiment consisted in 5 runs of 7 minutes. Stimuli were

pseudorandomized using optseq2 to optimize the sequence of presentation of the

different conditions. Three different optimized lists of trials were used across runs. List

order was counterbalanced across subjects.

One early blind was excluded from the analyses because the subject answered

to less than 70% of the trials throughout the experiment due to sleepiness. One run of

one sighted subject was excluded from the analysis because of a technical error during

the acquisition, and two other runs (one in a sighted subject, one in a blind subject)

were excluded since the subject answered to less than 70% of the trials in that specific

run.

Conceptual similarity ratings

In order to perform the adaptation analysis, we divided the trials in similar pairs

(e.g. red - orange) and different pairs (e.g. red - blue). We did so based on the

participants' subjective ratings. For each participant we took the average rating for

each of the 15 word-pairs in the action and color categories. Then we automatically

divided the 15 pairs in 5 intervals (4 quantiles) of nearly equal size. This subdivision

was performed using the function *quantile*, in R (R Core Team, 2013), that divides a

probability distribution into contiguous intervals of equal probabilities (i.e., 20%). The

pairs in the first two intervals were the different pairs (low ratings of similarity), the pairs

in the 3rd interval were the medium pairs, and the pairs in the 4th and 5th intervals were

the similar pairs (See fig 2B). However, in some cases, ratings distributions were

slightly unbalanced, due to the tendency of some subjects to find more "very different"

pairs than "very similar" pairs. In these cases (8 subjects for action ratings [3 EB]; 4

subjects for Color Ratings [1 EB]), the automatic split in 5 equal intervals was not

possible. Thus, we set the boundary between the 2nd and 3rd interval at the ratings

average (for that given subject), and set to the minimum (1 or 2, depending on the

cases) the number of items in the 3rd interval (not analyzed), in order to balance as

much as possible the number of pairs in the Different and Similar groups. This

procedure made so that in these special cases (as well as in all the others), the rating

values of different pairs were always below the mean, and the values of similar pairs

was always above the mean. Fig. S4, S5, S6, S7, in the supplementary information,

show subject-specific rating distributions.

MRI data acquisition

Brain images were acquired at the Neurological Institute Carlo Besta in Milano

on a 3-Tesla scanner with a 32-channel head coil (Achieva TX; Philips Healthcare,

Best, the Netherlands) and gradient echo planar imaging (EPI) sequences.

In the event-related experiment, we acquired 35 slices (voxel size 3 X 3 X 3.5)

with no gap. The data in-plane matrix size were 64 X 64, field of view (FOV) 220mm X

220mm, time to repetition (TR)= 2 s, flip angle 90 degrees and time to echo (TE)= 30

ms. In all, 1210 whole-brain images were collected during the experimental sequence.

The first 4 images of each run were excluded from the analysis for steady-state

magnetization. Each participant performed 5 runs, with 242 volumes per run.

Anatomical data was acquired using a T1-weighted 3D-TFE sequence with the

following parameters: 1 X 1 X 1 mm voxel size, 240 X 256 matrix size, 2.300 ms TR,

2.91 ms ET, 900 ms TI, 256 FoV, 160 slices.

MRI data analysis

We analyzed the fMRI data using SPM12 (www.

fil.ion.ucl.ac.uk/spm/software/spm12/) and Matlab R2014b (The MathWorks, Inc.).

Preprocessing. Preprocessing included slice timing correction of the functional time

series (Sladky et al., 2011), realignment of functional time series, coregistration of

functional and anatomical data, spatial normalization to an echoplanar imaging

template conforming to the Montreal Neurological Institute (MNI) space, and spatial

smoothing [Gaussian kernel, 6 mm full-width at half-maximum (FWHM)]. Serial

autocorrelation, assuming a first-order autoregressive model, was estimated using the

pooled active voxels with a restricted maximum likelihood procedure, and the

estimates were used to whiten the data and design matrices.

Data analysis. Following preprocessing steps, the analysis of fMRI data,

based on a mixed-effects model, was conducted in two serial steps accounting,

respectively, for fixed and random effects. In all the analysis the regressors for the

conditions of interest consisted of an event-related boxcar function convolved with the

canonical hemodynamic response function according to a variable epoch model

(Grinband, Wager, Lindquist, Ferrera, & Hirsch, 2008). Movement parameters derived

from realignment of the functional volumes (translations in x, y, and z directions and

rotations around x, y, and z axes), and a constant vector, were also included as

covariates of no interest. We used a high-pass filter with a discrete cosine basis

function and a cutoff period of 128 s to remove artifactual low-frequency trends.

Adaptation analysis: For each subject, the general linear model included 6

regressors corresponding to the 3 levels of similarity (different, medium, similar) in

each condition (color, action). Color and Action pairs in the medium condition were

modeled as regressors of no interest.

At the first level of analysis, linear contrasts tested for Repetition Suppression

[Different > Similar] collapsing across categories (Action, Color). The same contrasts

were then repeated within each category [Color Different > Color Similar; Action

Different > Action Similar]. Finally, we tested for the Similarity by Category interactions,

testing whether the adaptation was stronger in one category compared to the other

(e.g., [Color Different > Color Similar] > [Action Different > Action Similar]).

These linear contrasts generated statistical parametric maps [SPM(T)]. The

resulting contrast images were then further spatially smoothed (Gaussian kernel 5mm

FWHM) and entered in a second-level analysis (RFX), corresponding to a random-

effects model, accounting for inter-subject variance. One-sample t-tests were run on

each group separately. Two-sample t-tests were then performed to compare these

effects between groups (Blind vs Sighted).

Univariate analysis: For each subject, changes in regional brain responses

were estimated through a general linear model including 2 regressors corresponding

to the two categories Action and Color. The onset of each event was set at the

beginning of the first word of the pair, the offset was determined by the subject

response, thus included reaction time (Grinband et al., 2008). Linear contrasts tested

for action-specific [Action > Color] and color-specific [Color > Action] BOLD activity.

These linear contrasts generated statistical parametric maps [SPM(T)]. The

resulting contrast images were then further spatially smoothed (Gaussian kernel 5mm

FWHM) and entered in a second-level analysis, corresponding to a random-effects

model, accounting for inter-subject variance. One-sample t-tests were run on each

group separately. Two-sample t-tests were then performed to compare these effects

between groups (Blind vs Sighted) and to perform conjunction analyses to observe if

the two groups presented similar activated networks for the two contrasts of interests.

Connectivity analysis: Psychophysiological interaction (PPI) analyses were

computed to identify brain regions showing a significant change in the functional

connectivity with seed regions (the right precuneus, the left pMTG and the rIPS) that

showed a significant activation ($p<.001$, uncorrected) in the [(EB Conj. SC) X (Color >

Action)] contrast, the [(EB Conj. SC) X (Action > Color)] contrast, and the [(SC > EB)

X (Action > Color)] contrast respectively. In each individual, time series of activity

(principal eigenvariate) were extracted from a 8 mm sphere centered on the nearest

local maxima to the identified peaks in the second-level analysis (Note that centering

the sphere on the peak itself does not change the ROI Analysis results, see

Supplementary Information). New linear models were generated at the individual level,

using three regressors. One regressor represented the psychological condition of

interest (action or color trial). The second regressor was the physiological activity

extracted in the reference area. The third regressor represented the interaction of

interest between the first (psychological) and the second (physiological) regressor. The

design matrix also included movement parameters and a constant vector as regressors

of no interest. A significant PPI indicated a change in the regression coefficients

between any reported brain area and the seed region, related to the experimental

conditions (Color>Action or Action>Color). Next, the individual summary statistic

images obtained at the first-level (fixed-effects) analysis were spatially smoothed (5

mm FWHM Gaussian kernel) and entered in a second-level (random-effects) analysis

using a two-sample t test contrasting the two groups.

ROI definition. Occipital ROI for the PPI analyses were defined as following.

Two peak-coordinates were taken from previous studies (Bedny et al. 2011; Kanjlia et

al. 2016) showing the involvement of EB occipital areas in high-level functions such as

language (left MOG [-36, -90, -1]) and mathematics (right MOG [33, -82, 9]). These

areas also showed increased long range connectivity (in early blind) with frontal and

parietal areas during rest (Bedny et al. 2011; Kanjlia et al. 2016; Liu et al. 2007;

Collignon et al. 2013).

The V4 and V5 ROI were drawn from the literature, considering both perceptual

localizers, as well as evidence from semantic/conceptual task. We selected 3 peak

coordinates for area V5. The first [-47, -78, -2] from a highly-cited study contrasting the

perception of visual motion vs static images (Dumoulin et al., 2000). The second [-44,

-74, 2] from a study (Saygin, McCullough, Alac, & Emmorey, 2010) showing V5

sensitivity to motion sentences (e.g., "The wild horse crossed the barren field"). The

third from a research on the on-line meta-analysis tool Neurosynth

(http://neurosynth.org/) for the topic "action". In Neurosynth, the area in the occipital

cortex with the highest action-related activation was indeed V5 (peak coordinates: -50,

-72, 2). To avoid ROI proliferation, we averaged these 3 peak-coordinates in order to

obtain a single peak (average peak: -47, -75, 1).

As for V4, we selected the color-sensitive occipital ROI considering perceptual

localizers, as well as evidence of color-specific activity from semantic/conceptual task.

Fernandino et al. (Fernandino et al., 2015) reported a color sensitive area in the left

posterior collateral sulcus (ColS; at the intersection between the Lingual and the

Fusyform gyrus; MNI peak coordinates: -16, -71, -12) associated with color-related

words. This peak is close to the posterior-V4 localization done by Beauchamps and

colleagues (peak coordinates: -22, -82, -16) in a MRI version of the Farnsworth–

Munsell 100-Hue Test (Beauchamp, Haxby, Jennings, & DeYoe, 1999). A search in

neurosynth with the keyword "color" also highlighted a left posterior color-sensitive

region along the ColS with peak coordinates [-24, -90, -10]. We averaged these 3

peaks to find the center of our region of interest (average peak: -21, -81, -13).

The posterior lateral-temporal cortex ROI (PLTC) was taken from 3 studies

showing semantic repetition suppression in that area. Bedny and colleagues (Bedny,

McGill, and Thompson-Schill 2008) observed increased neural adaptation in PLTC

(peak coordinates: 57, -36, 21) for repeated words (fan - fan), when the words were

presented in a similar context (summer – fan; ceiling - fan), compared to when different

context triggered different meanings (e.g., admirer – fan; ceiling – fan). This result

conceptually replicated previous studies (Kotz, Cappa, Cramon, & Friederici, 2002;

Wible et al., 2006) showing semantic adaptation in the bilateral PLTC for related (e.g.,

dog - cat) vs unrelated (e.g., dog - apple) word pairs (peak coordinates: -42, -27, 9 and

-51, -22, 8). These 3 peaks were averaged to find the center of our region of interest

in both hemispheres (average peak: ±50, -28, 13).

Statistical analysis.

At the whole brain level, statistical inference was made at a corrected cluster

level of P < 0.05 FWE (with a standard voxel-level threshold of P < 0.001 uncorrected)

and a minimum cluster-size of 50 voxels. ROI analysis based on Small Volume

Correction were thresholded at p<0.05 FEW at the voxel level.

All ROI analyses were performed using Small Volume Correction using

spheres with a 10mm radius centered around the ROI peak coordinates (see previous

session). Within the ROI, results were considered significant at a threshold of p<0.05,

FEW-corrected. Here, and throughout the paper, brain coordinates are reported in MNI

space.

Behavioral data, analysis code and t-maps from the main contrasts will be

made available on-line (https://www.biorxiv.org/content/early/2018/08/23/384552).

Row fMRI images will be made available upon request, following agreements with our

ethical board committee.

Results

Within-category similarity is encoded in occipital areas in the sighted but not in

the blind. The rationale behind adaptation analyses was that the direct contrast

between pairs with high versus low perceptual differences will display neural

adaptation (Barron et al., 2016; Grill-Spector, Henson, & Martin, 2006; Wheatley,

Weisberg, Beauchamp, & Martin, 2005) therefore probing regions that are specifically

sensitive to the *perceptual distance* between concepts.

Behavioral analysis: Similarity ratings were highly correlated between sighted and

blind, both for action (r= .99) and color concepts (r= .93; Fig. 1A). In order to perform

the adaptation analysis we divided the trials in similar pairs (e.g. red - orange) and

different pairs (e.g. red - blue), based on each participant' subjective ratings. Rating

distributions for each subject and category (color, action) were divided in 5 intervals

with a similar number of items (see Method session for details). Stimulus-pairs in the

first two intervals were labeled as different (low similarity ratings), the 3^{rd} interval

contained medium pairs, and the 4^{th}-5^{th} intervals similar pairs (high similarity ratings;

Fig 1B). Overall, the average number of "different" trials was slightly larger than the

"similar" ones (126 vs 115; $F(1,33)=8.41$, $p=0.007$, $\eta^2=0.20$; Fig. 1C). However, there

was no similarity by group interaction ($F(1,33)=0.18$, $p=0.67$, $\eta^2=0.004$), indicating

that this unbalance (that reflected personal judgments of similarity) was the same

across SC and EB (fig. 1C). An analysis of reaction times showed that Medium pairs

(not analyzed in fMRI) had on average longer latencies than Similar and Different ones

(Main Effect of Similarity: $F(2,66)=21.07$, $p<0.001$, $\eta^2=0.38$). This was expected since

pairs that are neither similar nor different would require longer and more difficult

judgments. Crucially, there was no difference in reaction times between different

(Mean=1.80 sec, SD=0.39) and similar pairs (Mean=1.79 sec, SD=0.37; $F(1,33)=0.09$,

$p=0.76$, $\eta^2=0.003$), and no interaction between Similarity and Group ($F(1,33)=0.04$,

$p=0.84$, $\eta^2=0.001$; Fig 1D).

Figure 1. Adaptation, behavioral analysis. (A) Similarity judgments were highly correlated
across groups both for actions and color; (B) Conceptual schema of the division of word pairs
in "different" and "similar" based on subjective similarity ratings; (C) Barplot depicting the
average number of items in the "different", "medium" and "similar" categories. The number of
items in the "different" and "similar" categories is very similar across groups (number of trials ±
SEM); (D) Barplot depicting the average reaction time in the "different", "medium", and "similar"
categories. The average RTs of the "different" and "similar" categories is very similar across
groups (seconds ± SEM).
10

fMRI analysis: to find brain areas that showed adaptation based on conceptual

similarity, we looked at the contrast Different Pairs > Similar Pairs, with Medium pairs

as a regressor of no interest. Action and color pairs were considered together since,

at the whole brain level, we did not find a significant higher-order interaction between

Similarity (different, similar) and Category (see the method section for analysis details).

In the sighted, similar concepts led to repetition suppression in several occipital areas

(See Fig. 2A), with a significant cluster in and around the left lingual gyrus (peak

coordinates: -24, -70, -7). In the blind, instead, adaptation emerged in language-related

areas with significant clusters along the middle and superior temporal gyrus, bilaterally

(Peak coordinates RH: 57, -28, 8; LH: -60, -10, -7) and in the right precentral gyrus

(Peak coordinates: 27, -25, 56; Fig. 2B). Importantly, no adaptation in posterior

occipital areas was observed in the blind.

A comparison between groups showed greater adaptation in occipital cortices

for sighted compared to blind (Fig. 2C), with peaks in the left superior occipital gyrus

(-24, -91, 26), the left lingual gyrus (-24, -70, -7) and the right middle occipital gyrus

(27, -85, 11). The contrast Blind > Sighted showed increased adaptation in posterior

lateral temporal cortices (PLTC) bilaterally (Fig. 2D). Planned ROI analysis in PLTC, a

region that consistently show repetition suppression for semantic similarity (Bedny,

McGill, and Thompson-Schill 2008; Wible et al. 2006; Kotz et al. 2002), revealed a

significantly greater adaptation for similar concepts in blind more than sighted

(Conceptual Similarity by Group interaction; lPLTC= -45 -31 20, t(33)=3.23, P= 0.035;

rPLTC= 45 -28 11, t(33)=3.41, P= 0.024).

Figure 2. Adaptation, fMRI results. Regional BOLD responses are rendered over Conte-
69 average midthickness surfaces. (A) Suprathreshold clusters showing neural adaptation for

similar word pairs in the occipital cortices of sighted participants; (B) Suprathreshold clusters
showing neural adaptation for similar word pairs in the temporal and somatosensory-motor
cortices of early blind participants; (C) Suprathreshold clusters showing neural adaptation for
similar word pairs in sighted compared to blind, and (D) blind compared to sighted; (E)
Suprathreshold cluster showing neural adaptation for similar color-word pairs in the sighted,
with regional activity precisely localized in the left PCoS/V4 (see Fernandino et al. 2015 for a
very similar result with a multivariate design); (F) Suprathreshold clusters showing neural
adaptation for similar action-word pairs in the sighted, with regional activity spread in different
occipital areas including the left PCoS/V4 and the left V5. Cluster threshold at P<0.005
uncorrected, for illustration only.
11

Finally, we performed planned ROI analysis in the color-sensitive region at the

posterior banks of the collateral sulcus (PCoS) corresponding to the V4-complex

(Beauchamp et al., 1999; Fernandino et al., 2015) and the motion sensitive region V5

(Dumoulin et al., 2000). In area PCoS-V4 we found greater adaptation both for color

and action in sighted compared to blind (Conceptual Similarity by Group interaction;

peak= -24 -73 -10, $t(33)=4.26$, $P= 0.004$; Fig 2 E-F). In contrast, the analysis in V5

showed that repetition suppression was specific for action concepts in the sighted and

no adaptation was observed in the blind (Conceptual Similarity by Group by Category

interaction; Peak: -51 -76 8, $t(33)=3.29$, $P= 0.037$; Fig 2F).

Brain regions active in sighted and blind when contrasting action and color

concepts. Subsequently we ran classic univariate analysis, comparing items across

categories independently of their similarity, to find category-specific activations across

sighted and blind. In these analyses, the two words in each pair were considered as a

single trial.

Behavioral analysis: Reaction times analysis using a mixed ANOVA, with Category

(action, color) as within-subject factor and Group (sighted, blind) as between-subjects

factor, showed no difference between categories ($F(1,33)=2.37$, $p>0.05$, $\eta^2= 0.07$),

between groups ($F(1,33)=0.074$, $p>0.05$, $\eta^2=0.002$) and no Category by Group

interaction ($F(1,33)=0.69$, $p>0.05$, $\eta^2=0.02$).

fMRI analysis: The contrast Action > Color did not reveal any significant difference

between groups, suggesting a comparable categorical representation of action

concepts, across sighted and blind (see fig. S2 for details). Indeed, a conjunction

analysis between groups showed a common significant activation in the left posterior

middle temporal gyrus (lpMTG; Peak= -54, -61, 5; Fig 3A).

On the other hand, a conjunction analysis between groups for Color > Action

did not reveal any common activation between sighted and blind after correction for

multiple comparisons at the whole brain level. However, displaying the conjunction

results at a more lenient threshold (p<.001 uncorrected; Fig 3C), we could notice a

unique common activity for color concepts in the right precuneus (peak= 6, -55, 26).

Accordingly, analysis within groups showed a significant precuneus activity in the blind

(peak= 6, -52, 20, p= .04) and a marginally significant activity in the sighted (peak= 0,

-61, 29, p= .06), with no significant difference between groups (Table S1; Fig. S1).

Further analysis for the contrast Color > Action revealed a cluster in the right

parietal cortex, in and around the right intraparietal sulcus (rIPS), showing higher

activity for color concepts in sighted compared to blind (peak= 33, -43, 35; Fig. 3B).

Altogether these results show similar patterns of activity during conceptual

processing in sighted and blind, when categorical preferences are investigated

(independently of perceptual similarity). As in previous results (Bedny et al., 2012; Uta

Noppeney et al., 2003; van den Hurk et al., 2017) such common activities are found

outside the posterior occipital cortex, in areas that are considered to be highly

polymodal such as the precuneus and the lpMTG (Binder, Desai, Graves, & Conant,

2009). Interestingly, even at this level of comparison we could find an effect of visual

deprivation, concerning color knowledge, that seem to involve the right IPS more in

sighted than blind.

Figure 3. Contrasts between categories and psychophysiological interactions. Regional
BOLD responses are rendered over Conte-69 average midthickness surfaces. (A)
Suprathreshold cluster (P<0.05 FEW corrected) showing common activity in the lpMTG for the
contrast Action > Color in both sighted and early blind (conj., conjunction analysis); (B)
Suprathreshold cluster (P<0.001 uncorrected; for illustrative purposes only) showing common
activity in the precuneus for the contrast Color > Action in both sighted and early blind (conj.,
conjunction analysis); (C) Suprathreshold cluster (P<0.05 FEW corrected) showing greater
activity in the rIPS, in sighted compared to early blind, for the contrast Color > Action; (D)
Suprathreshold clusters (P<0.005 uncorrected; for illustrative purposes only) showing greater
connectivity in the occipital areas of early blind people with the lpMTG (PPI for the contrast
Action > Color); (E) Barplot (for illustrative purposes only) showing beta weights derived from
PPI analysis, with seed in lpMTG, in sighted (gray) and blind (white), in the right and left middle
occipital gyrus (MOG) for the contrast Action > Color (arbitrary unit ± SEM); (F) Suprathreshold
clusters (P<0.005 uncorrected; for illustrative purposes only) showing greater connectivity in
the occipital areas of early blind people with the precuneus (PPI for the contrast Color > Action);
(G) Barplot (for illustrative purposes only) showing beta weights derived from PPI analysis, with
seed in precuneus, in sighted (gray) and blind (white), in the right and left middle occipital gyrus
(MOG) for the contrast Color > Action (arbitrary unit ± SEM).

Psychophysiological interactions: The precuneus and lpMTG display similar

category selectivity across sighted and blind but show different connectivity

profiles. Finally, we relied on Psychophysiological Interaction (PPI) analysis (Friston

et al., 1997; O'Reilly, Woolrich, Behrens, Smith, & Johansen-berg, 2012) to test

whether regions that showed similar categorical preference across groups (lpMTG for

action > color, and the precuneus for color > action) also maintain a similar connectivity

profile in both groups. In particular, we tested the hypothesis that posterior occipital

areas in the blind could be recruited during conceptual retrieval, and connect with

conceptual hubs (Binder et al., 2009) such as the lpMTG and the Precuneus, as a

consequence of neural reorganization (Amedi, Raz, Pianka, Malach, & Zohary, 2003;

Burton, 2003). With this aim, we selected two ROIs in the left and right middle occipital

gyrus (MOG) that are recruited in early blind during high-level conceptual tasks such

as language processing and math (Kanjlia et al. 2016; Bedny et al. 2011); and show

increased long range connectivity, in early blind, with extra-occipital regions (e.g.,

frontal, parietal and ventral temporal cortices) during resting state (Bedny et al. 2011;

Kanjlia et al. 2016; Liu et al. 2007; Collignon et al. 2013) and task-based (PPI) analysis

(Noppeney, Friston, and Price 2003).

PPI with seed in the lpMTG revealed an increase of action-selective functional

connectivity in both occipital ROIs of blind people compared to their sighted

counterpart (lMOG: $t(33)= 3.59$, $p= 0.02$; rMOG: $t(33)=3.46$, $p=0.026$; Fig 1D & 1E).

Similarly, PPI with seed in the precuneus revealed an increase of color-selective

functional connectivity in the occipital cortex of blind compared to sighted participants

(lMOG: $t(33)= 3.12$, $p= 0.054$; rMOG: $t(33)=4.09$, $p=0.007$; Fig 1F and 1G). Albeit

showing a similar activity profile in sighted and blind during conceptual processing, the

lpMTG and the precuneus showed a different connectivity profile as a function of early

visual deprivation. Such increase in task-based connectivity suggests that occipital

areas in early blind, albeit not coding for perceptual similarity as in the sighted, are

however active during conceptual retrieval and can be flexibly recruited in interaction

with conceptual hubs such as the precuneus and the pMTG.

Discussion

Embodied approaches to conceptual knowledge suggest that concepts are grounded in our sensory and motor experience of the world (Barsalou, 1999; Binder & Desai, 2011). A straightforward hypothesis emerging from these theories is that people that perceive the world in a different way should also have different conceptual representations (Casasanto, 2011).

In our study we tested this hypothesis by characterizing the brain activity of sighted and early blind individuals while they rated the perceptual similarity of action and color concepts in fMRI. In particular, we investigated which brain regions encode the perceptual similarity of retrieved concepts using an adaptation paradigm. Results in the sighted group showed that word-pairs referring to similar colors or actions induced repetition suppression in several posterior occipital regions, including areas V3, V4 and V5. In striking contrast, early blind participants did not show repetition suppression in posterior occipital areas but instead showed a greater adaptation for concept similarity in language-related regions, disclosing a different neurobiology of concepts as a function of visual deprivation.

Posterior occipital regions are known to encode visual features in the sighted and to be sensitive to visual similarity independently of categorical membership (Bracci and Op de Beeck 2016; Connolly et al. 2012; Kriegeskorte et al. 2008; Naselaris et al. 2009). Our data corroborate the hypothesis that these regions are also involved in conceptual retrieval (Borghesani et al., 2016; Fernandino et al., 2015) supporting information related to the visual appearance of objects and events, that is not available to blind people. This result is crucial to support the hypothesis that conceptual retrieval consists, in part, on replay of our perceptual experience (Barsalou, 1999; Harnad, 1990).

Since posterior occipital regions represent visual features also during

conceptual retrieval (Borghesani et al., 2016; Fernandino et al., 2015; Mitchell et al.,

2008), it is therefore expected, as observed in our study, that sighted and blind should

show different activity in these regions due to the lack of visual qualia in the blind

population. Our results are in line with a previous study showing that relatively anterior

regions such as the pMTG and part of the VOTC show a similar functional and

connectivity fingerprint in sighted and blind (Wang et al., 2015), whereas this similarity

decreases strikingly in more posterior occipital regions (i.e., approximately behind the

conventional line that separate the temporal from the occipital lobe; Wang et al. 2015).

Actually, the posterior occipital regions showing higher conceptual adaptation in

sighted compared to blind in our study tightly overlap with regions showing the lowest

functional and connectivity similarity between sighted and blind in Wang et al. study

(Wang et al. 2015; Fig. S2).

Reduced adaptation in the occipital cortex in our blind participant (compared to

sighted) coincides with stronger adaptation in the posterior lateral-temporal cortices

(PLTC). Several studies have found that conceptually similar (e.g., dog - wolf) or

semantically associated (e.g., dog - leash) words can lead to repetition suppression in

PLTC (Wible et al. 2006; Kotz et al. 2002; Bedny, McGill, and Thompson-Schill 2008).

Although it is still unclear what level of conceptual knowledge is represented in that

region (Bedny, McGill, and Thompson-Schill 2008), there is some agreement that the

PLTC stores auditory representations of words, that are connected to distributed

semantic representations in the brain (Hickok and Poeppel 2007; Gow 2012; Bedny,

McGill, and Thompson-Schill 2008; Mirman et al. 2015). In this framework, the PLTC

may work at the interface between wordforms and semantic knowledge (Gow, 2012;

Hickok & Poeppel, 2007), and a greater activity in the blind can index a larger use of

verbal knowledge in this population (Cattaneo et al., 2008; Crollen et al., 2014).

Directly contrasting different categories (Action Vs Color), independently of

within-category similarity, showed instead some commonalities in brain activity across

the two groups (Bedny et al. 2012; Noppeney, Friston, and Price 2003; Peelen et al.

2014). Both sighted and blind engaged the lpMTG during action processing and the

Precuneus during color processing. These regions are located outside of the visual

cortex, in polymodal areas, that typically display category-specificity in a format that is

at least partially independent from perceptual appearance (Bracci & Op de Beeck,

2016; Peelen & Downing, 2017) and largely resilient to visual deprivation (Bedny et al.,

2012; Uta Noppeney et al., 2003; Wang et al., 2015).

Interestingly, though, psychophysiological interactions (PPI) showed that early

visual deprivation changes the connectivity profile of these regions, increasing their

functional coupling with occipital regions in the blind. These results suggest that the

occipital cortex in early blind is re-organized to extend its integration into conceptual

selective networks, highlighting further how visual deprivation impact on the

neurobiology of conceptual knowledge. Previous studies have shown that occipital

areas in the early blind are recruited for high-level conceptual tasks such as language

processing, semantic retrieval and math (Bedny et al. 2011; Kanjlia et al. 2016; Burton,

Diamond, and McDermott 2003; Van Ackeren et al. 2018; Crollen et al. 2018; See also

Fig. S3 showing general higher activity in the blind occipital cortex during the auditory

presentation of words), and that they increase long-range connectivity with frontal and

parietal cortices during rest (Bedny et al. 2011; Kanjlia et al. 2016; Liu et al. 2007;

Collignon et al. 2013) and inferior temporal cortices during semantic judgments

(Noppeney, Friston, and Price 2003). Notably, graph-theoretic metrics of regional

cortical thickness covariance found that language and visual regions showed a pattern

of merging into shared modules in the blind but not in sighted (Hasson, Andric, Atilgan,

& Collignon, 2016). Extending those previous studies, we show here that early blinds

seem to rely on enhanced connectivity between occipital cortices and temporo-parietal

conceptual hubs during conceptual processing. Albeit our data remain correlational,

they suggest that EBs' occipital regions are not activated independently of other

"classic" regions involved in conceptual retrieval, but they instead work in concert.

These results suggest that occipital cortices are involved in conceptual retrieval

in both sighted and blind, but with different functions and probably at different levels of

representation. Occipital areas may support sensorimotor simulations of visual

features during conceptual retrieval in the sighted, showing adaptation for concepts

that refer to visually similar objects or events; on the other hand, in the blind, occipital

cortices do not encode perceptual similarity, but may be re-organized to engage in

more general processes related to conceptual retrieval (albeit these processes need

to be better specified; see for instance, Bedny 2017; Van Ackeren et al. 2018).

Outside the posterior occipital cortex, we found that the posterior portion of the

right IPS showed a stronger preference for color trials in the sighted compared to the

blind. The IPS is known to be involved in the perception of color (Beauchamp et al.,

1999; Cheadle & Zeki, 2014; Zeki & Stutters, 2013) as well as other visual features

(Grill-Spector, 2003; Swisher, Halko, Merabet, McMains, & Somers, 2007; Xu, 2007),

and its anatomical position make it a good candidate to work at the interface between

perceptual and conceptual representations (Cheadle & Zeki, 2014). In particular, it

seems to be crucial for retrieving perceptually-based categorical knowledge (Cheadle

& Zeki, 2014; Zeki & Stutters, 2013). Indeed, the peak of color-specific activity that we

found (peak coordinates: 33, -43, 35) is very close to the color-specific rIPS area found

by Cheadle & Zeki (2014; peak coordinates: 30, -39, 45). The lack of visual input in

blind people prevents the formation of perceptually-driven color representations, which

may limit the contribution of the IPS during the retrieval of color knowledge. This is not

the case for action representation, for which perceptual knowledge can be more easily

compensated by other senses (e.g., touch, audition) in the blind. The distinct impact of

blindness on the broader categorical responses to action and colors again highlight

how visual experience may selectively shape the neurobiology of conceptual

representations, even in putatively multimodal areas of the brain.

From a broader theoretical point of view, the results of this study are in line with a

hierarchical model of conceptual representations based on progressive levels of

abstraction (Barsalou, 2016; Binder, 2016; Binder et al., 2016; Fernandino et al., 2015;

A. Martin, 2015). At the top of the hierarchy, multimodal representations may co-exist

with purely symbolic ones organized in a linguistic/propositional code (Mahon &

Caramazza, 2008). This level of representation can account for the obvious fact that

congenitally blind people can think about colors and their perceptual properties

(Barilari, de Heering, Crollen, Collignon, & Bottini, 2018; Marmor, 1978) although they

cannot simulate vision. On the other hand, modality-specific simulation in sensory

areas (e.g. visual, auditory, somatosensory), as the one highlighted by our adaptation

analysis, may become central in deeper and more deliberate stages of conceptual

processing, providing situated, specific and sometimes imaginistic representations,

and eventually supporting the subjective experience of knowing providing the

phenomenological qualia of conscious thinking (Binder et al., 2016; Koch, Massimini,

Boly, & Tononi, 2016).

References

Amedi, A., Raz, N., Pianka, P., Malach, R., & Zohary, E. (2003). Early "visual" cortex activation correlates with superior verbal memory performance in the blind. *Nature Neuroscience, 6*(7), 758–66. http://doi.org/10.1038/nn1072

Barilari, M., de Heering, A., Crollen, V., Collignon, O., & Bottini, R. (2018). Is Red Heavier Than Yellow Even for Blind? *I-Perception, 9*(1), 204166951875912. http://doi.org/10.1177/2041669518759123

Barron, H. C., Garvert, M. M., & Behrens, T. E. J. (2016). Repetition suppression: a means to index neural representations using BOLD? *Philosophical Transactions of the Royal Society B: Biological Sciences, 371*(1705), 20150355. http://doi.org/10.1098/rstb.2015.0355

Barsalou, L. W. (1999). Perceptual Symbol Systems. *Behavioral and Brain Science, 30322*(September 2001), 577–660.

Barsalou, L. W. (2009). Simulation, situated conceptualization, and prediction. *Philosophical Transactions of the Royal Society B: Biological Sciences, 364*(1521), 1281–1289. http://doi.org/10.1098/rstb.2008.0319

Barsalou, L. W. (2016). On Staying Grounded and Avoiding Quixotic Dead Ends. *Psychonomic Bulletin & Review, 23*(4), 1122–1142. http://doi.org/10.3758/s13423-016-1028-3

Beauchamp, M. S., Haxby, J. V, Jennings, J. E., & DeYoe, E. A. (1999). An fMRI version of the farnsworth-munsell 100-hue test reveals multiple color-selective areas in human ventral occipitotemporal cortex. *Cerebral Cortex, 9*(3), 257–263. http://doi.org/10.1093/cercor/9.3.257

Bedny, M. (2017). Evidence from Blindness for a Cognitively Pluripotent Cortex. *Trends in Cognitive Sciences, 21*(9), 637–648. http://doi.org/10.1016/j.tics.2017.06.003

Bedny, M., Caramazza, a, Pascual-Leone, a, & Saxe, R. (2012). Typical neural representations of action verbs develop without vision. *Cerebral Cortex (New York, N.Y. : 1991), 22*(2), 286–93. http://doi.org/10.1093/cercor/bhr081

Bedny, M., McGill, M., & Thompson-Schill, S. L. (2008). Semantic adaptation and competition during word comprehension. *Cerebral Cortex, 18*(11), 2574–2585. http://doi.org/10.1093/cercor/bhn018

Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E., & Saxe, R. (2011). Language processing in the occipital cortex of congenitally blind adults. *Proceedings of the National Academy of Sciences, 108*(11), 4429–4434. http://doi.org/10.1073/pnas.1014818108

Bedny, M., & Saxe, R. (2012). Insights into the origins of knowledge from the cognitive neuroscience of blindness. *Cognitive Neuropsychology, 29*(1–2), 56–84. http://doi.org/10.1080/02643294.2012.713342

Binder, J. R. (2016). In defense of abstract conceptual representations. *Psychonomic Bulletin & Review, 23*(4), 1096–1108. http://doi.org/10.3758/s13423-015-0909-1

Binder, J. R., Conant, L. L., Humphries, C. J., Fernandino, L., Simons, S. B., Aguilar, M., & Desai, R. H. (2016). Toward a brain-based componential semantic representation. *Cognitive Neuropsychology, 33*(3–4), 130–174. http://doi.org/10.1080/02643294.2016.1147426

Binder, J. R., & Desai, R. H. (2011). The neurobiology of semantic memory. *Trends in Cognitive Sciences, 15*(11), 527–536. http://doi.org/10.1016/j.tics.2011.10.001

Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where is the semantic system? A critical review and meta-analysis of 120 functional neuroimaging studies. *Cerebral Cortex, 19*(December), 2767–2796. http://doi.org/10.1093/cercor/bhp055

Boersma, P., & Weenink, D. (2018). Praat: doing phonetics by computer. Retrieved

from http://www.praat.org/

Borghesani, V., Pedregosa, F., Buiatti, M., Amadon, A., Eger, E., & Piazza, M.

(2016). Word meaning in the ventral visual path: a perceptual to conceptual

gradient of semantic coding. *NeuroImage, 143*, 128–140.

http://doi.org/10.1016/j.neuroimage.2016.08.068

Bracci, S., & Op de Beeck, H. (2016). Dissociations and associations between shape

and category representations in the two visual pathways. *Journal of*

Neuroscience, 36(2), 432–444. http://doi.org/10.1523/JNEUROSCI.2314-

15.2016

Burton, H. (2003). Visual cortex activity in early and late blind people. *The Journal of*

Neuroscience : The Official Journal of the Society for Neuroscience, 23(10),

4005–4011.

Burton, H., Diamond, J. B., & McDermott, K. B. (2003). Dissociating cortical regions

activated by semantic and phonological tasks: a FMRI study in blind and sighted

people. *Journal of Neurophysiology, 90*(3), 1965–82.

http://doi.org/10.1152/jn.00279.2003

Casasanto, D. (2011). Different Bodies, Different Minds: The Body Specificity of

Language and Thought. *Current Directions in Psychological Science, 20*(6),

378–383. http://doi.org/10.1177/0963721411422058

Cattaneo, Z., Vecchi, T., Cornoldi, C., Mammarella, I., Bonino, D., Ricciardi, E., &

Pietrini, P. (2008). Imagery and spatial processes in blindness and visual

impairment. *Neuroscience and Biobehavioral Reviews, 32*(8), 1346–1360.

http://doi.org/10.1016/j.neubiorev.2008.05.002

Cheadle, S. W., & Zeki, S. (2014). The role of parietal cortex in the formation of color

and motion based concepts. *Frontiers in Human Neuroscience, 8*(July), 535.

http://doi.org/10.3389/fnhum.2014.00535

Collignon, O., Dormal, G., Albouy, G., Vandewalle, G., Voss, P., Phillips, C., & Lepore, F. (2013). Impact of blindness onset on the functional organization and the connectivity of the occipital cortex. *Brain*, *136*(9), 2769–2783. http://doi.org/10.1093/brain/awt176

Connolly, A. C., Guntupalli, J. S., Gors, J., Hanke, M., Halchenko, Y. O., Wu, Y.-C., … Haxby, J. V. (2012). The representation of biological classes in the human brain. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *32*(8), 2608–18. http://doi.org/10.1523/JNEUROSCI.5547-11.2012

Crollen, V., Lazzouni, L., Bellemare, A., Rezk, M., Lepore, F., Noel, M.-P., … Collignon, O. (2018). Recruitment of occipital cortex by arithmetic processing follows computational bias in early blind. *BioRxiv Preprint*.

Crollen, V., NoëL, M. P., Seron, X., Mahau, P., Lepore, F., & Collignon, O. (2014). Visual experience influences the interactions between fingers and numbers. *Cognition*, *133*(1), 91–96. http://doi.org/10.1016/j.cognition.2014.06.002

Dumoulin, S. O., Bittar, R. G., Kabani, N. J., Baker, C. L., Le Goualher, G., Bruce Pike, G., & Evans, A. C. (2000). A new anatomical landmark for reliable identification of human area V5/MT: a quantitative analysis of sulcal patterning. *Cerebral Cortex*, *10*(5), 454–463. http://doi.org/10.1093/cercor/10.5.454

Fernandino, L., Binder, J. R., Desai, R. H., Pendl, S. L., Humphries, C. J., Gross, W. L., … Seidenberg, M. S. (2015). Concept representation reflects multimodal abstraction: A framework for embodied semantics. *Cerebral Cortex*, 1–17. http://doi.org/10.1093/cercor/bhv020

Fodor, J. (1975). *The language of thought*. Cambridge: Harvard University Press.

Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., & Dolan, R. J. (1997).

Psychophysiological and modulatory interactions in neuroimaging. *Neuroimage*, *6*(3), 218–229. http://doi.org/10.1006/nimg.1997.0291

Gow, D. W. (2012). The cortical organization of lexical knowledge: A dual lexicon model of spoken language processing. *Brain and Language*, *121*(3), 273–288. http://doi.org/10.1016/j.bandl.2012.03.005

Grill-Spector, K. (2003). The neural basis of object perception. *Current Opinion in Neurobiology*, *13*(2), 159–166. http://doi.org/10.1016/S0959-4388(03)00040-0

Grill-Spector, K., Henson, R., & Martin, A. (2006). Repetition and the brain: Neural models of stimulus-specific effects. *Trends in Cognitive Sciences*, *10*(1), 14–23. http://doi.org/10.1016/j.tics.2005.11.006

Grinband, J., Wager, T. D., Lindquist, M., Ferrera, V. P., & Hirsch, J. (2008). Detection of time-varying signals in event-related fMRI designs. *NeuroImage*, *43*(3), 509–520. http://doi.org/10.1016/j.neuroimage.2008.07.065

Harnad, S. (1990). The symbol grounding problem. *Physica D*, *42*, 335–346. http://doi.org/10.1080/0952813X.2014.940139

Hasson, U., Andric, M., Atilgan, H., & Collignon, O. (2016). Congenital blindness is associated with large-scale reorganization of anatomical networks. *NeuroImage*, *128*, 362–372. http://doi.org/10.1016/j.neuroimage.2015.12.048

He, C., Peelen, M. V., Han, Z., Lin, N., Caramazza, A., & Bi, Y. (2013). Selectivity for large nonmanipulable objects in scene-selective visual cortex does not require visual experience. *NeuroImage*, *79*, 1–9. http://doi.org/10.1016/j.neuroimage.2013.04.051

Hickok, G., & Poeppel, D. (2007). The cortical organization of speech processing. *Nature Reviews Neuroscience*, *8*(May), 393–402. http://doi.org/10.1038/nrn2113

Horner, A. J., & Henson, R. N. (2011). Repetition suppression in occipitotemporal cortex despite negligible visual similarity: Evidence for postperceptual

processing? *Human Brain Mapping*, *32*(10), 1519–1534.

http://doi.org/10.1002/hbm.21124

Hsu, N. S., Kraemer, D. J. M., Oliver, R. T., Schlichting, M. L., & Thompson-Schill, S.

L. (2011). Color, context, and cognitive style: Variations in color knowledge

retrieval as a function of task and subject variables. *Journal of Cognitive*

Neuroscience, *23*(9), 2544–2557. http://doi.org/10.1162/jocn.2011.21619

Kanjlia, S., Lane, C., Feigenson, L., & Bedny, M. (2016). Absence of visual

experience modifies the neural basis of numerical thinking. *Proceedings of the*

National Academy of Sciences, *113*(40), 201524982.

http://doi.org/10.1073/pnas.1524982113

Koch, C., Massimini, M., Boly, M., & Tononi, G. (2016). Neural correlates of

consciousness: progress and problems. *Nature Reviews Neuroscience*, *17*(5),

307–321. http://doi.org/10.1038/nrn.2016.22

Kosslyn, S. M., Ganis, G., & Thompson, W. L. (2001). Neural foundations of imagery.

Nature Reviews Neuroscience, *2*(9), 635–642. http://doi.org/10.1038/35090055

Kotz, S. A., Cappa, S. F., Cramon, D. Y. Von, & Friederici, A. D. (2002). Modulation

of the Lexical – Semantic Network by Auditory Semantic Priming : An Event-

Related Functional MRI Study. *NeuroImage*, *17*, 1761–1772.

http://doi.org/10.1006/nimg.2002.1316

Kriegeskorte, N., Mur, M., Ruff, D. A., Kiani, R., Bodurka, J., Esteky, H., …

Bandettini, P. A. (2008). Matching Categorical Object Representations in Inferior

Temporal Cortex of Man and Monkey. *Neuron*, *60*(6), 1126–1141.

http://doi.org/10.1016/j.neuron.2008.10.043

Lakoff, G., & Johnson, M. (1999). *Philosophy in the flesh*. New York: Basic Books.

Liu, Y., Yu, C., Liang, M., Li, J., Tian, L., Zhou, Y., … Jiang, T. (2007). Whole brain

functional connectivity in the early blind. *Brain*, *130*(8), 2085–2096.

http://doi.org/10.1093/brain/awm121

Mahon, B. Z., & Caramazza, A. (2008). A critical look at the embodied cognition hypothesis and a new proposal for grounding conceptual content. *Journal of Physiology Paris*, *102*(1–3), 59–70. http://doi.org/10.1016/j.jphysparis.2008.03.004

Marmor, G. (1978). Age at onset of blindness and the development of the semantics of color names. *Journal of Experimental Child Psychology*, *278*, 344–345. Retrieved from http://www.sciencedirect.com/science/article/pii/0022096578900826

Martin, A. (2015). GRAPES-Grounding representations in action, perception, and emotion systems: How object properties and categories are represented in the human brain. *Psychonomic Bulletin & Review*, 979–990. http://doi.org/10.3758/s13423-015-0842-3

Martin, A., Haxby, J., & Lalonde, F. (1995). Discrete Cortical Regions Associated with Knowledge of Color and Knowledge of Action. *Science*, *270*(5233), 102–105. Retrieved from http://www.sciencemag.org/content/270/5233/102.short

Martin, C. B., Douglas, D., Newsome, R. N., Man, L. L. Y., & Barense, M. D. (2018). Integrative and distinctive coding of visual and conceptual object features in the ventral visual stream. *ELife*, *7*, 1–29. http://doi.org/10.7554/eLife.31873

Mirman, D., Chen, Q., Zhang, Y., Wang, Z., Faseyitan, O. K., Coslett, H. B., & Schwartz, M. F. (2015). Neural organization of spoken language revealed by lesion-symptom mapping. *Nature Communications*, *6*, 1–9. http://doi.org/10.1038/ncomms7762

Mitchell, T. M., Shinkareva, S. V., Carlson, A., Chang, K.-M., Malave, V. L., Mason, R. A., & Just, M. A. (2008). Predicting Human Brain Activity Associated with the Meanings of Nouns. *Science*, *320*(5880), 1191–1195.

http://doi.org/10.1126/science.1152876

Mohr, H. M., Linder, N. S., Linden, D. E. J., Kaiser, J., & Sireteanu, R. (2009).
Orientation-specific adaptation to mentally generated lines in human visual
cortex. *NeuroImage*, *47*(1), 384–391.
http://doi.org/10.1016/j.neuroimage.2009.03.045

Naselaris, T., Prenger, R. J., Kay, K. N., Oliver, M., & Gallant, J. L. (2009). Bayesian
Reconstruction of Natural Images from Human Brain Activity. *Neuron*, *63*(6),
902–915. http://doi.org/10.1016/j.neuron.2009.09.006

Noppeney, U., Friston, K. J., & Price, C. J. (2003). Effects of visual deprivation on the
organization of the semantic system. *Brain : A Journal of Neurology*, *126*(Pt 7),
1620–1627. http://doi.org/10.1093/brain/awg152

Noppeney, U., & Price, C. J. (2001). Functional imaging of the semantic system:
Retrieval of sensory-experienced and verbally learned knowledge. *Brain and
Language*, *84*, 120–133. http://doi.org/10.1016/S0093-934X(02)00525-4

Noppeney, U., & Price, C. J. (2002). Retrieval of visual, auditory, and abstract
semantics. *NeuroImage*, *15*, 917–926. http://doi.org/10.1006/nimg.2001.1016

O'Reilly, J. X., Woolrich, M. W., Behrens, T. E. J., Smith, S. M., & Johansen-berg, H.
(2012). Tools of the trade: Psychophysiological interactions and functional
connectivity. *Social Cognitive and Affective Neuroscience*, *7*(5), 604–609.
http://doi.org/10.1093/scan/nss055

Peelen, M. V., Bracci, S., Lu, X., Chenxi, H., Caramazza, A., & Bi, Y. (2013). Tool
selectivity in left occipitotemporal cortex develops without vision. *Journal of
Cognitive Neuroscience*, *25*(8), 1225–1234. http://doi.org/10.1162/jocn

Peelen, M. V., & Downing, P. E. (2017). Category selectivity in human visual cortex:
Beyond visual object recognition. *Neuropsychologia*, *105*(April), 177–183.
http://doi.org/10.1016/j.neuropsychologia.2017.03.033

Peelen, M. V, He, C., Han, Z., Caramazza, A., & Bi, Y. (2014). Nonvisual and visual object shape representations in occipitotemporal cortex: evidence from congenitally blind and sighted adults. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *34*(1), 163–70. http://doi.org/10.1523/JNEUROSCI.1114-13.2014

Proklova, D., Kaiser, D., & Peelen, M. V. (2016). Disentangling Representations of Object Shape and Object Category in Human Visual Cortex: The Animate– Inanimate Distinction. *Journal of Cognitive Neuroscience*, *28*(5), 680–692. http://doi.org/10.1162/jocn

Pulvermüller, F., & Hauk, O. (2006). Category-specific conceptual processing of color and form in left fronto-temporal cortex. *Cerebral Cortex (New York, N.Y. : 1991)*, *16*(8), 1193–201. http://doi.org/10.1093/cercor/bhj060

Pylyshyn, Z. W. (1984). *Computation and cognition*. Cambridge: MIT Press.

R Core Team. (2013). R: A language and environment for statistical computing.

Saygin, A. P., McCullough, S., Alac, M., & Emmorey, K. (2010). Modulation of BOLD response in motion-sensitive lateral temporal cortex by real and fictive motion sentences. *Journal of Cognitive Neuroscience*, *22*(11), 2480–90. http://doi.org/10.1162/jocn.2009.21388

Simmons, W. K., Ramjee, V., Beauchamp, M. S., McRae, K., Martin, A., & Barsalou, L. W. (2007). A common neural substrate for perceiving and knowing about color. *Neuropsychologia*, *45*(12), 2802–2810. http://doi.org/10.1016/j.neuropsychologia.2007.05.002

Sladky, R., Friston, K. J., Tröstl, J., Cunnington, R., Moser, E., & Windischberger, C. (2011). Slice-timing effects and their correction in functional MRI. *NeuroImage*, *58*(2), 588–594. http://doi.org/10.1016/j.neuroimage.2011.06.078

Striem-Amit, E., Wang, X., Bi, Y., & Caramazza, A. (2018). How do blind people

represent rainbows? Disentangling components of conceptual representations. *BioRxiv*, 287318. http://doi.org/10.1101/287318

Swisher, J. D., Halko, M. A., Merabet, L. B., McMains, S. A., & Somers, D. C. (2007). Visual Topography of Human Intraparietal Sulcus. *Journal of Neuroscience*, *27*(20), 5326–5337. http://doi.org/10.1523/JNEUROSCI.0991-07.2007

Van Ackeren, M. J., Barbero, F., Mattioni, S., Bottini, R., & Collignon, O. (2018). Neuronal populations in the occipital cortex of the blind synchronize to the temporal dynamics of speech. *ELife*, 1–45. http://doi.org/10.7554/eLife.31640

van den Hurk, J., Van Baelen, M., & Op de Beeck, H. P. (2017). Development of visual category selectivity in ventral visual cortex does not require visual experience. *Proceedings of the National Academy of Sciences*, 201612862. http://doi.org/10.1073/pnas.1612862114

Wang, X., Peelen, M. V., Han, Z., He, C., Caramazza, A., & Bi, Y. (2015). How Visual Is the Visual Cortex? Comparing Connectional and Functional Fingerprints between Congenitally Blind and Sighted Individuals. *Journal of Neuroscience*, *35*(36), 12545–12559. http://doi.org/10.1523/JNEUROSCI.3914-14.2015

Wheatley, T., Weisberg, J., Beauchamp, M. S., & Martin, A. (2005). Automatic priming of semantically related words reduces activity in the fusiform gyrus. *Journal of Cognitive Neuroscience*, *17*(12), 1871–85. http://doi.org/10.1162/089892905775008689

Wible, C. G., Han, S. D., Spencer, M. H., Kubicki, M., Niznikiewicz, M. H., Jolesz, F. A., … Nestor, P. (2006). Connectivity among semantic associates: An fMRI study of semantic priming. *Brain and Language*, *97*(3), 294–305. http://doi.org/10.1016/j.bandl.2005.11.006

Xu, Y. (2007). The Role of the Superior Intraparietal Sulcus in Supporting Visual Short-Term Memory for Multifeature Objects. *Journal of Neuroscience*, *27*(43),

1   11676–11686. http://doi.org/10.1523/JNEUROSCI.3545-07.2007

2 Zeki, S., & Stutters, J. (2013). Functional specialization and generalization for

3   grouping of stimuli based on colour and motion. *NeuroImage, 73*, 156–166.

4   http://doi.org/10.1016/j.neuroimage.2013.02.001