Motor Learning Without Moving: Proprioceptive and Predictive Hand Localization After Passive Visuoproprioceptive Discrepancy Training

Ahmed M. Mostafa, Bernard Marius 't Hart, Denise Y.P. Henriques

CVR / Kinesiology, York University, Toronto, Ontario, Canada

Corresponding author:
Bernard Marius 't Hart
thartbm@gmail.com

Abstract

An accurate estimate of limb position is necessary for movement planning. Where we localize our unseen hand after a reach depends on felt hand position, or proprioception, but this is usually neglected in favour of predicted sensory consequences based on efference copies of motor commands. Both sources of information should contribute, so here we set out to further investigate how much of hand localization depends on proprioception and how much on predicted sensory consequences. We use a passive training paradigm with rotated visual feedback that eliminates the possibility to update predicted sensory consequences, but still recalibrates proprioception. Localizing an unseen hand after self-generated movements uses both efference-based predictions as well as afferent proprioceptive signals, but after robot-generated movements only proprioception is available. We find direction changes in open-loop reaches as well as shifts in hand localization, both after robot- and self-generated hand movements. Both motor and proprioceptive changes are only slightly smaller as those after training with self-generated movements, confirming that proprioception plays a large role in estimating limb position and in planning movements.

Introduction

Sensory information is central to how we control all our movements. Our brain is even thought to use *predicted* sensory consequences derived from efferent copies of motor commands for motor control (1). When training with rotated visual feedback of the hand, we update these predictions (2). Additionally, such training leads to a recalibration of our sense of felt hand position - "proprioception" - to be more aligned with the distorted visual feedback (3). Both of these changes have been measured by asking people to localize where their unseen hand is – before and after training (4–8). While our lab has previously found that proprioception accounts for a large portion of the change in hand localization (9), it is far from clear how much each process contributes or how to tease them apart. Here we use passive training, that removes the need to update predicted sensory consequences, in an attempt to isolate the contribution of proprioception to hand localization.

The predicted sensory consequences of movements may play several important roles in motor control. Predicted sensory consequences allow us to correct our movements before sensory error signals are available, they can be used to select movements that best achieve our goals and they may inform us on the location of our limbs. Hence measuring predicted sensory consequences is valuable in movement research. In visuomotor rotation adaptation tasks, the actual sensory outcome is systematically different from the expected outcome, so that participants update the predictions on the outcome of the trained movements. In previous experiments, people were asked to make a movement and then indicate the location of, or "localize," their unseen, right hand, before and after training with rotated visual feedback (5,7). Since there was no visual information available to the participants, the predicted sensory

consequences of the movement should be used in localizing the unseen hand. Both studies found a significant shift in hand localization, providing evidence that predicted sensory consequences are indeed *40* updated as a result of visuomotor rotation adaptation.

However, our lab has shown that our sense of where we feel our hand to be, proprioception, is also reliably recalibrated after visuomotor rotation adaptation (3,10–18). This has also been shown by other labs (19) and a comparable proprioceptive change is induced with force-field adaptation (20). As proprioception also informs us on the location of our limbs, we have on occasion used a task that is very *45* similar to hand localization to investigate this (3,6,21,22). Although proprioceptive recalibration has been largely ignored as an explanation for changes in hand localization, we and others have shown that it accounts for a substantial part of the changes in localization, along with updates in predicted sensory consequences (9,19). Nevertheless, it is far from clear how much proprioception and prediction each contribute to hand localization.   *50*

Here we intend to further examine the contribution of proprioception to hand localization. To do this, we use passive training where a robot arm moves the participants' arm out so that the cursor always directly goes to the target (19,23,24). This means there is no efference copy available and no visuomotor error-signal, both of which are required to update predicted sensory consequences. However, there is a discrepancy between vision and proprioception that drives proprioceptive recalibration. Thus, changes in *55* hand localization after this type of training should be due to proprioceptive recalibration only. We use the same experimental protocal as before (9), so that we can compare localization shifts between the two different types of training, and can better assess the contributions of predicted sensory consequences and proprioception to hand localization.

Methods   *60*

We set out to test the relative contributions of proprioception and efference-based prediction to hand localization. We use visual training with robot-generated hand movements to prevent updates of predicted sensory consequences, but still elicit proprioceptive recalibration.

Participants

Twenty-five right-handed participants were recruited for this study. One participant was excluded for not *65* following task instructions, and three were excluded for low performance on a task that ensures attention during the passive training. All analyses presented here pertain to the remaining twenty-one participants (mean age: 20.1±2.3, 13 females and 8 males), but the data of the three low-performing participants is included in the online dataset [URL?]. All had normal or corrected-to-normal vision, and provided prior, written informed consent in accordance with the ethical guidelines set by the York Human subjects *70* Review Subcommittee and received credit toward an undergraduate psychology course. Participants were screened verbally, and all reported being right handed and not having any history of visual, neurological, and/or motor dysfunction.

Setup

Participants sat in a height-adjustable chair to ensure that they could easily see and reach all targets *75* presented on a reflective screen (see Fig 1). During all tasks, they held the vertical handle on a two-joint robot manipulandum (Interactive Motion Technologies Inc., Cambridge, MA) with their right hand so that their thumb rested on top of the handle. A monitor (Samsung 510 N, refresh rate 60 Hz) was mounted 11 cm above the reflective screen, such that images displayed on the monitor appeared to lie in the horizontal plane where the right hand was moving. The reflective screen was mounted horizontally 18 cm *80* above the robot manipulandum. A touch screen was mounted 13 cm underneath the reflective surface, so that subjects could indicate the location of the unseen right-hand locations (specifically the unseen thumb) with their visible left hand, which was lit up with a small spot light (only in localization tasks). The

Figure 1: Setup and tasks. a) Participants moved their unseen right hand with visual feedback on hand position provided through a mirror (middle surface) half-way between their hand and the monitor (top surface). A touchscreen located just above the hand was used to collect responses for the localization tasks (bottom surface). **b)** Training task. The target, shown as a yellow disc, is located 10 cm away from the home position at 45°. In the rotated training tasks, the cursor (shown here as a green circle) represents the hand position rotated 30° relative to the home position. **c)** No-cursor reach task. Targets are located 10 cm away from the home position at 15°, 25°, 35°, 45°, 55°, 65°, and 75°, shown by the yellow circles here (only one was shown on each trial). While reaching to one of these targets, no visual feedback on hand position is provided. **d)** Localization task. The participants' unseen, right hands moved out, and subsequently participants indicated the direction of the hand movement by indicating a location on an arc using a touch screen with their visible left index finger.

85   room lights were dimmed and the participants' view of their right hand was blocked by the reflective screen, as well as a dark cloth draped between the touch screen and participants' right shoulder.

Procedure

The first part of the experiment used training with a cursor aligned with the hand and the second part had training with a cursor rotated around the start position (Table 1). During the training with rotated feedback, the cursor was gradually rotated 30° clockwise. This introduces a discrepancy between the actual, or felt,

90   hand position and the visual feedback, that should evoke proprioceptive recalibration. However, the movements are robot generated, so that there are no predicted sensory consequences based on the outgoing motor command. Hence the prediction errors that are thought to lead to motor learning are absent as well. After both types of training, open-loop reaches as well as two kinds of hand localization tasks were done to test the effect of training on proprioceptive and predictive hand estimates.

aligned	rotated		
№ trials in session		Exposure training	Classic training
50	90	training	training
-	21	no-cursor reaches	no-cursor reaches
-	60	training	training
25	25	active delayed	active delayed
21	21	no-cursor reaches	no-cursor reaches
10	60	training	training
25	25	passive delayed	passive delayed
21	21	no-cursor reaches	no-cursor reaches
10	60	training	training
25	25	active delayed	(active online)
21	21	no-cursor reaches	no-cursor reaches
10	60	training	training
25	25	passive delayed	(passive online)
21	21	no-cursor reaches	no-cursor reaches

Table 1. Task order.   *95*

Exposure training

In what we call 'exposure training' the participants did not move their hand toward the target, but the robot did. In this task (Table 1, white rows), the right hand was represented by a cursor (green disk, 1 cm in diameter, Fig 1b) located directly above participant's thumb. The robot moved the participant's unseen right hand (and the cursor) along a direct path toward a visual yellow target disk and back to the starting *100* position (1 cm in diameter, Fig 1b). The home position was located approximately 20 cm in front of subjects and the visual target located 10 cm from the home position at 45° (Fig 1b). In order to make sure participants were paying attention to the cursor, the cursor was switched of for 2 screen refreshes (~33.3ms) on 50% of the trials at a random distance between 4 and 9 cm from the home position and participants were asked to report this using a button press with the left hand. Performance on this task *105* was used to screen participants.

During the first half of the experiment, the cursor and hand path were aligned during exposure training. In the second part of the experiment, the "rotated" session, the same training target at 45° was used, and the cursor kept moving straight to this target. However, the robot-generated hand path gradually rotated 30° CCW (Fig 1b) with respect to the visible target and the cursor in increments of *110* 0.75°/trial, so that the full rotation was reached after 45 trials. This mimics error-free responses to a gradual visuomotor rotation of 30° CW. The initial training consisted of 50 trials in the aligned part and 90 in the rotated part. In between open-loop reach tasks and localization tasks (Table 1, all gray rows) extra training tasks were done, each of which consisted of 10 trials in the aligned part of the experiment and 60 trials in the rotated part. *115*

No-cursor reaching

The trials in no-cursor reaching (Table 1, light gray rows) serve as a classical measure of motor adaptation. On each of these trials participants were asked to reach with their unseen right hand to one of 7 visual targets, without any visual feedback on hand position. The targets were 10 cm from the home position, located radially at: 15°, 25°, 35°, 45°, 55°, 65°, and 75° (Fig 1c). A trial started with the robot *120*

handle at the home position and, after 500 ms, the home position disappeared and the target appeared, cuing the participants to reach for the target. Once the participants thought they had reached the target they held their position for 250 ms, and the target and the home position disappeared, cuing participants to move back to the home position along a straight, constrained path to begin the next trial. If participants tried to move outside of the path, a resistance force, a stiffness of 2 N/(mm/s) and a viscous damping of 5 N/(mm/s), was generated perpendicular to the path. In every iteration of the no-cursor reach task, each target was reached three times, for a total of 21 reaches in pseudo-random order. The no-cursor reaching task was performed four times in the aligned part of the experiment and five times in the rotated part of the experiment.

Localization

In this task (Fig 1d; Table 1, dark gray rows) participants indicated where they thought their unseen right hand was after a movement. First, an arc appeared, spanning from 0° to 90° and located 10 cm away from the home position and the participants' unseen, right hand moved out from the home position in a direction somewhere within the arc. The hand was stopped by the robot at 10 cm from the home position and then, to prevent the online proprioceptive signals from overriding the predictive signals (5,9), the hand was moved back to the home position using the same kind of constrained path as used for the return movements in the no-cursor task. Participants indicated with the index finger of their visible, left hand on the touch screen mounted directly above the robot handle where they thought their trained hand had crossed the arc.

Crucially, there were two variations of this task. First, in the 'active' localization task participants generated the movement themselves, as they could freely move their unseen right hand from the home position to any point on the arc. Second, there was a 'passive' localization task where the robot moved the participants' hand out and back, to the same locations the participants moved to in the preceding 'active' localization task in a shuffled order (hence, active localization is done first). In active localization, participants have access to both proprioceptive information as well as an efference-based prediction of sensory consequences, but in passive localization, only proprioception should be available. The active and passive localization task each consisted of 25 trials, and each of the tasks was done a total of four times; twice after aligned and twice after rotated training.

Classic training

The paradigm described above is an exact replica of a paradigm we used earlier (9) with two exceptions. First, we used exposure training here, instead of the standard reach training with volitional movements, which we will call 'classic' training. Second, all localization is delayed until the right hand has returned to the home position in this study (Table 1), so that instead of both delayed and online localization we have two repetitions of each delayed localization task. With this paradigm we can compare changes in localization and no-cursor reaches change after exposure training with changes in the same measures after classic visuomotor adaptation training.

Analyses

Prior to any analyses, both the localization responses and the no-cursor reach data were visually inspected and trials where the participants did not follow task instruction were removed (e.g. several movements back and forth, or a touch-screen response on the home position, instead of on the white arc).

Localization

Localization responses were taken as the (signed) angular difference between vectors through the home position and the actual hand position as well as the location indicated on the touch screen. Prior to

analyses, idiosyncratic differences in performing this task were countered. Before conversion to degrees angle, a circle with a 10 cm radius was fit to the touch screen responses of each participant and the offset of this circle's centre was subtracted from all response coordinates, so that all responses fell close to the arc. Then, a smoothed spline was fit to every participant's response errors in each of the four localization *170* tasks (aligned vs. rotated and active vs. passive) and these were used to obtain localization errors at the same locations used for the no-cursor reaches (15°, 25°, 35°, 45°, 55°, 65° and 75°), but only if that location fell within the range of the data (i.e. we only interpolate). This way localization responses could be compared across participants despite the freely chosen reach directions. At the 15° degree location 7/21 participants didn't have an estimate in one or more of the four localization tasks (in the "classic" data *175* it was also 7/21). While that data is shown in the figures, we did not use it for analysis.

First we test if localization responses shifted following rotated exposure training compared to aligned. We then test if the shift in localization responses is different for active and passive localization, and we run analyses comparing localization after exposure training with localization after "classic" training. Finally, we explore the generalization of localization responses and if they are different between *180* the groups doing classic and exposure training.

Reach aftereffects

To assess any reach adaptation that may have occurred after exposure training we analyzed reach endpoints errors in no-cursor trials. Reach endpoint errors were the (signed) angular difference between a vector from the home position to reach endpoint and a vector from the home position to the target. We *185* obtained reach aftereffects by subtracting reach endpoint errors after aligned training from those after rotated training. No-cursor endpoint errors were analyzed to see if participants adapted the direction of their reaching movements after rotated exposure training. We also tested if any such change decayed, i.e. if it was the same immediately after exposure training, or when a localization task was done in between exposure training and no-cursor reaches. Furthermore, we test if the generalization of reach *190* aftereffects is different between exposure and classic training and if there is any generalization of reach aftereffects.

Pre-processing and analyses were done in R 3.4.3 (25) using lme4, lmerTest and various other packages. Most analyses used linear mixed effects models, since there is some missing localization data. These are "converted" to more readable ANOVA-like output, using a Satterthwaite approximation (26). Highly similar *195* results are obtained with a Chi-square approximation. Data, scripts and a notebook with analyses (27) are available on the Open Science Framework (https://osf.io/zfdth/) and GitHub (https://github.com/thartbm/exposureconsequences).

In short, this experiment allowed us to test how mere exposure to a visual-proprioceptive discrepancy changes both reach aftereffects and hand localization responses, and compare them with those obtained *200* after more regular, "classic" training.

Results

In this study we intend to further elucidate the relative contributions of (updated) predicted sensory consequences and (recalibrated) proprioception to hand localization. We can parcel out these contributions by measuring hand localization after both robot-generated and self-generated movements. *205* Finally we compare the data from the current experiment with those obtained in an earlier study that used an identical paradigm, but trained with self-generated movements, or "classic" training.

Figure 2: Hand localization. The shifts of the angles of touchscreen responses in all variations of the localization task, using spline-interpolated estimates for hand angles matching the reach targets in the no-cursor reach block. **a)** Localization shifts after expourure training. Dark blue: active localization shifts, Light blue: passive localization shifts. **b)** Localization shifts after classic training. Dark red: active localization shifts, Orange: passive localization shifts. The dashed line segments illustrate that the 15° data is not used for statistical analyses (see Methods). **c)** Generalization curves of active localization shifts after exposure training (blue) and classic training (red). Shaded areas: 95% confidence intervals for the peak of the generalization curve (red and blue lines through shaded area indicate 50% points). Downward black arrow: visual trained target. Upward black arrow: hand location during training.

Localization

First we test our hypothesis that exposure training does not lead to changes in predicted sensory consequences. Since the difference between active and passive localization stems only from the presence or absence of predicted sensory consequences, there should be no difference between the two if predicted sensory consequences are not changed by exposure training. At first glance, it seems there might be a difference between active and passive localization shifts in exposure training (see Fig 2a), although it is smaller than in classic training (Fig 2b).

First, to test if rotated exposure training induces changes in hand localization, we fit an LME to the localization errors throughout the workspace using *session* (aligned or rotated), *movement type* (active and passive) and *hand angle* (25°, 35°, 45°, 55°, 65° and 75°), and all interactions as fixed effects and *participant* as random effect. There was an effect of *session* ($F_{(1,450.5)}=155.8$; $p<0.001$), showing that exposure training leads to changes in hand localization. There was also an effect of *hand angle* ($F_{(5,450.6)}=6.54$; $p<.001$) and an interaction between *hand angle* and *session* ($F_{(5,450.3)}=8.25$; $p<.001$), but no other effects (all $p>.60$), which we'll explore below. Since localization responses shifted, for further tests we use the difference between hand localization after rotated training and after aligned training (as plotted in Fig 2).

If this shift in localization after exposure training partly reflects predicted sensory consequences, then shifts in active localization, that rely on both (recalibrated) proprioception and (updated) predictions should be different from shifts in passive localization that only rely on (recalibrated) proprioception. We fit an LME to the change in localization using *movement type* (active or passive localization) and *hand angle*, as well as their interaction as fixed effects and *participant* as random effect. There was no effect of *movement type* ($F_{(1,211.8)}=0.07$; $p=0.79$). There was an effect of *hand angle* ($F_{(5,212.2)}=10.8$; $p<0.001$), but no interaction between *hand angle* and *movement type* ($F_{(5,211.8)}=1.23$, $p=.29$). The lack of an effect of *movement type* suggests that predicted sensory consequences did not contribute to localization in this paradigm.

In order to compare hand localization shifts after exposure training with those after classic training (9), we fit an LME to localization shift using *training type* (exposure vs. classic), *movement type* (active vs. passive) and *hand angle* and all interactions as fixed effects and *participant* as random effect. There was a main effect of *movement type* (F(1,422.0)=6.22; p=.013) and of *hand angle* (F(5,422.7)=8.19; p<.001), as well as an interaction between *training type* and *hand angle* (F(5,422.7)=4.54; p<.001) and between *training type* and *movement type* (F(1,422.0)=4.48, p=.035), but there was no main effect of *training type* (F(1,39.1)=0.92, p=.34) and no other effects (all p>.14). These results also suggests that the magnitude of the shifts in localization are comparable between classic and exposure training, but that the pattern of generalization is different.

To address our main question, we will look at the interaction between *training type* (exposure vs. classic) and *movement type* (active vs. passive) we found above. Since there is no difference between active and passive localization shifts after exposure training alone, the interaction between *training type* and *movement type* should be caused by an effect of *movement type* on the localization shifts after classic training, as we found previously (9). This means that shifts in hand localization after exposure training indeed relies on recalibrated proprioception alone, while after classic training, there also is a contribution of predicted sensory consequences to active localization.

For completeness, we explore the potentially different generalization patterns of localization shifts after classic and exposure training (Fig 2c). The LME indicates no difference in overall amplitude of localization shifts between the groups, so the interaction between *training type* and *hand angle* might stem from a generalization that does not peak at the trained location after exposure training. Using the active localization shifts only (which are larger, and arguably more similar to reach aftereffects), we bootstrap a 95% confidence interval for the peak localization shift across participants in each group. Here we include the data at 15° where it is available. After classic training, the peak localization shift was at 48.7° (95% confidence: 31.6° - 62.8°; red area in Fig 2c), and after exposure training the peak localization shift was at 62.1° (95% confidence: 50.0° - 75.6°; blue are in Fig 2c). This means that peak localization after classic training is lower than after exposure training, but not vice versa. Also note that the confidence interval for the peak localization shift after classic training includes the trained target (45°), but not after exposure training (5). In short, the LME for localization shifts indicates a different generalization curve after exposure and classic training, which could be partially explained by a different position of the peak localization shift after exposure or classic training.

Reach aftereffects

Apart from proprioception and prediction, we want to see if rotated exposure training has any effect on open-loop reaches and if these are robust. We measure whether participants adapted their reach directions by assessing their reach errors in no-cursor reach trials after aligned and rotated exposure training. In Fig 3, the changes in no-cursor endpoint errors, or reach aftereffects, appear to be well over 5°. First, to test if exposure training affects open-loop reach direction, we fit a linear mixed effects model (LME) to reach endpoint error using *session* (aligned; all blocks, or rotated; only the first block immediately after training) and *target* (15°, 25°, 35°, 45°, 55°, 65° and 75°), as well as their interactions as fixed effects and *participant* as random effect. There is an effect of *session* (F(1,260)=93.81, p<.001), that is: exposure training leads to substantial reach aftereffects. There was no effect of *target* (F(6,260)=1.07, p=.37) and no interaction (F(6,260)=1.00, p=.42). Since there is an effect of *session*, we now take the differences in reach endpoint errors between the rotated and aligned session for every participant and target as reach aftereffects, and use those for all further analyses.

To see if reach aftereffects decayed during the localization tasks, we compared reach aftereffects in the initial no-cursor block, that immediately followed training, with those in the later blocks that followed a localization task. We fit an LME to the reach aftereffects with *iteration* (initial vs. later no-cursor blocks) and *target* (as above) as well as their interaction as fixed effects and *participant* as random effect. There is no effect of *iteration* (F(1,260)=2.72, p=0.10). There was an effect of *target* (F(6,260)=6.29, p<.001) but no interaction (F(6,260)=0.58, p=0.74). Hence, reach aftereffects were not appreciably different right after

Figure 3: Reach aftereffects. Changes of the angle of reach endpoints in the no-cursor tasks. **a)** Reach aftereffects across the experiment. Light blue: first no-cursor task in the rotated session (immediately following training), Dark blue, dashed line: he other four repetitions of the task (with localization in between training and no-cursor tasks). **b)** Reach aftereffects after classic and exposure training. Blue: exposure training, Red: classic training. **c)** Generalization curves of reach aftereffects after exposure training (blue) and classic training (red). Shaded areas: 95% confidence intervals for the peak of the generalization curve (red and blue lines through shaded area indicate 50% points). Downward black arrow: visual trained target. Upward black arrow: hand location during training.

training and after the localization tasks. In other words, there was likely no noticeable decay of reach aftereffects during the localization tasks, so that we can collapse the data across iterations.

285 Now we compare the reach aftereffects after classic training with those after exposure training (Fig 3b). It appears as if there is little overall difference in magnitude, but there might be a shift of the generalization curve. We fit an LME to reach aftereffects with *training type* (classic vs. exposure), *target* (as above) as well as their interaction as fixed effects and *participant* as random effect. There is no main effect of *training* (F(1,40)=0.11, p=.74) indicating approximately equal magnitude of reach aftereffects after the two training types. There is an effect of *target* (F(6,240)=8.36, p<.001), indicating that reach

290 aftereffects exhibit some form of a generalization curve. There is also an interaction between *training type* and *target* (F(6,240)=2.27, p=.038), indicating these generalization curves are different after the two training types.

We will explore these potentially different generalization curves here. In Fig 3c we can observe that reach aftereffects after exposure training seem not to peak at the trained target direction of 45° but a

295 more forward direction. We test this by taking the 95% confidence interval of the centre of a normal curve fit to this data, bootstrapped across participants, and find that the median peak of the generalization curve of reach aftereffects after exposure training is at 66.3°, with a 95% confidence interval ranging from 49.3° to 78.9° this would indeed suggest that the reach aftereffects after exposure training do not generalize around the trained direction of 45°, but at a more counter clockwise location. For classic training,

300 generalization of reach aftereffects peaks at 53.2°, with a 95% confidence interval spanning 42.0° to 66.4°. So for classic training the 95% confidence interval for peak reach aftereffects does include the trained target. These confidence intervals also indicates that generalization of reach aftereffects does not peak at different target position after exposure and classic training. However, we can also observe that the full curve was not sampled after exposure training, so that curve fitting is not optimal. This means that

305 – given our data – the interaction between *target* and *training type* found in the LME above can't be explained by a shifted generalization curve.

In summary, our main hypotheses are confirmed; exposure training leads to shifts in hand localization that are not different for active or passive localizatoin, while movement type does have an effect on localization shift after classic training. Exposure training also causes robust reach aftereffects that are of comparable size to those found with classic training. There is some evidence that the generalization of *310* both localization shifts and reach aftereffects are different after the two training types, and it appears this can partially be explained by a different peak of the generalization curves, but our data and analyses are not definitive.

Discussion

The position of limbs is important for planning and evaluating movements, and can be estimated through *315* predicted sensory consequences, as well as visual and proprioceptive feedback. As in a previous study (9) here we quantify the contributions of predicted sensory consequences and proprioceptive recalibration to where we localize our hand after training with altered visual feedback of the hand. In classical adaptation paradigms, both predictions are updated and proprioception is recalibrated. Predictions are updated when they don't match actual sensory consequences, and proprioception is recalibrated when it *320* doesn't match visual feedback. In this study we use "exposure" training, where the participants do not have volitional control of their movements. By design. this should eliminate efference copies and prevent updating predicted consequences of movements, but since the proprioceptive and visual feedback is the same, exposure training still allows proprioceptive recalibration. Before and after training, participants localize their hand, both after "active," self-generated movements that allow using predicted sensory *325* consequences, and after "passive," robot-generated movements that only allow using proprioception. We calculate the training-induced shift in both types of localization given the same actual hand position. After classic training we previously reported larger shifts in active localization as compared to passive (9). As we expected, after exposure training there are substantial shifts in localization, but no difference between active and passive localization, indicating that predictions are not updated after exposure training. *330* Furthermore, we find that exposure training evokes substantial and robust reach aftereffects, indicating that recalibrated proprioception is used to plan movements.

      Our lab previously investigated proprioceptive recalibration and reach aftereffects following visuomotor adaptation with classic training and matched exposure training. There we also found that proprioceptive recalibration is of similar magnitude in both training paradigms, but unlike here, reach *335* aftereffects are usually much larger with classic training (13,22–24,28). And while proprioceptive recalibration and reach aftereffects do proportionally increase with gradual increases in rotation size for classical training, they do not for exposure training (24). The similar magnitude of proprioceptive recalibration and reach aftereffects following exposure training, but not classical training, suggest that this sensory recalibration is partly driving this modest change in movements. The effect of exposure training *340* on movements is also demonstrated by savings and interference from exposure training to subsequent classic training (29) and transfer of exposure training effects from one hand to the other (30). In the current study, we further demonstrate that exposure training affects movements and proprioception, but also measure its potential effect on predictive estimates.

      Results similar to what we find here were reported in a study by Cameron and colleagues (19), *345* using gain modulation of visual feedback of single-joint hand movements around the elbow. Their within-subjects experiment included both training with volitional movements as well as with passive movements and also tested perception of movements that were either passive or active. They too found a robust change in passive perception of hand movement (using a different measure), and these changes did not differ between the two types of training. Similarly, they found shifts in what we might call "active *350* localization," although the task is different, after both training types. Like here, these shifts are larger after classic training as compared to exposure training. They also found that passive exposure leads to reach aftereffects, although these were smaller than those produced following "classical" training with altered visual gain. Both our findings, and those of Cameron et al. (19) indicate that updating predicted sensory

355   consequences requires volitionally controlled movements that lead to prediction errors, while proprioception recalibrates equally in both types of training, and that recalibrated proprioception affects open-loop reaches. Our combined results suggest that updates in predicted sensory consequences only provide a partial explanation for motor learning.

          In our classical training group, we not only see shifts for passive localization but even larger shifts
360   for active localization which is consistent with a change in both proprioception and an update in predictions. In our exposure training, we did not find a consistent difference between active and passive localization, and none at the trained direction. Assuming that predictions were not updated in exposure training, a maximum likelihood estimate (MLE) or "optimal integration" (31) would predict that active localization should shift less than passive localization after exposure training. But of course this is not the
365   case in our findings (although it is the case for Cameron et al., 2012). This suggests that perhaps these two signals are not optimally integrated which is consistent with our comparisons of the variance between passive and active localization. In 't Hart and Henriques (9), we tested the prediction derived from MLE that hand localization with two signals – proprioception and prediction in active localization – should be more reliable, i.e. have lower variance, than hand localization with only one signal – proprioception only in
370   passive localization. However, we found no difference in variance between active and passive localization, and recently replicated this in a much larger dataset (32). Taken together, this suggests these different sources of information about unseen hand location are not optimally integrated. While localizing the unseen hand is less precise than locating (pointing to) a remembered visual target or a seen and felt hand location, we find that these bimodal estimates are rarely integrated optimally (33–35), although
375   others have (36). A more recent study (37) has also shed doubt on whether "optimal" or "Bayesian" integration is used for locating the hand with two afferent signals. Analogously, here we again can't find evidence that afferent and efferent information combine as a maximum likelihood estimate.

          It seems clear that the cerebellum plays a role in motor learning as it appears to compute predicted sensory consequences, i.e. it implements a forward model. (38–40) People with cerebellar
380   damage do worse on motor learning tasks (41–44), and show decreased shifts in hand localization tasks following motor learning (5,7). This highlights that the cerebellum, and likely predicted sensory consequences, are important for motor learning, but does not explain the remaining shifts in hand localization. We previously found that proprioceptive recalibration is intact in people with mild cerebellar ataxia and that it is similar following exposure and classical training with a gradually introduced cursor
385   rotation (28). The remaining changes in hand localization found in cerebellar patients can be attributed to recalibrated proprioception which should be intact (18). Analogously, here we show that in a paradigm that stops updates of predictions of sensory consequences, as supposedly in people with cerebellar damage, we still see substantial shifts in localization. Again, the remaining localization shifts can be explained if, along with predictions, we use afferent recalibrated proprioceptive estimates to localize our
390   hand.

Generalization

          We do find some evidence that, after exposure training, the generalization curves for localization shifts are not centred on the visual location of the trained target; they don't peak at 45° but at ~62°. In contrast, after classic training the peak of the generalization curve does peak close to the training target.
395   It is possible that proprioceptive recalibration is not anchored to the visual goal of the training task as it is not a requirement to feel your hand at any specific point; rather the visual cursor has to be brought to a visual target to end a trial. Although we can't substantiate this here, the generalization curves of the reach aftereffects seem to mimic the generalization curves of localization shifts, suggesting a relationship between changes in state estimates and changes in movements. This needs to be tested further, but if
400   these effects are true, they may provide insight into how state estimates are used to produce movements, and also may lead to a way method to disentangle the influence of recalibrated proprioception and more traditional updated internal models on motor changes, such as reach aftereffects. Either way, even though the experiment was not designed to investigate this, the shifted generalization curves of changes in

localization after classic and exposure training suggest they are generated by different mechanisms. This is in line with our earlier findings that proprioception generalizes differently from reach adaptation (11,45).   *405*

Conclusion

To sum up, after a training paradigm designed to prevent updating of predicted sensory consequences but allow recalibration of proprioception, we find substantial changes in where people localize their hand. This means that recalibrated proprioceptive estimates can explain shifts in localization. Since our participants also changed the direction of open loop reaches, recalibrated proprioception seems to guide   *410* motor planning. Finally, we have some evidence that after exposure training, the shift in hand localization does not generalize around the trained target location, confirming that different mechanisms underly proprioceptive recalibration and motor adaptation.

Acknowledgements

This work was supported by a Canadian Network for Research and Innovation in Machining Technology   *415* NSERC Operating grant (DYPH) and the German Research Foundation (DFG) under grant no. HA 6861/2-1 (BMtH). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. We thank Shanaathanan Modchalingam for assistance in collecting the data.

References   *420*

1. Duhamel J-R, Colby CL, Goldberg ME. The Updating of the Representation of Visual Space in Parietal Cortex by Intended Eye Movements. Science (80-). 1992;255(5040):90–2.

2. Wolpert DM, Ghahramani Z, Jordan MI. An internal model for sensorimotor integration. Science [Internet]. 1995 Sep 29 [cited 2018 Feb 9];269(5232):1880–2. Available from: http://www.ncbi.nlm.nih.gov/pubmed/7569931

3. Cressman EK, Henriques DYP. Sensory Recalibration of Hand Position Following Visuomotor Adaptation. J Neurophysiol [Internet]. 2009;102(6):3505–18. Available from: http://jn.physiology.org/cgi/doi/10.1152/jn.00514.2009

4. Clayton HA, Cressman EK, Henriques DYP. Proprioceptive sensitivity in Ehlers-Danlos syndrome patients. Exp Brain Res. 2013;

5. Izawa J, Criscimagna-Hemminger SE, Shadmehr R. Cerebellar Contributions to Reach Adaptation and Learning Sensory Consequences of Action. J Neurosci [Internet]. 2012;32(12):4230–9. Available from: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.6353-11.2012

6. Jones SAH, Fiehler K, Henriques DYP. A task-dependent effect of memory and hand-target on proprioceptive localization. Neuropsychologia [Internet]. 2012;50(7):1462–70. Available from: http://dx.doi.org/10.1016/j.neuropsychologia.2012.02.031

7. Synofzik M, Lindner A, Thier P. The Cerebellum Updates Predictions about the Visual Consequences of One's Behavior. Curr Biol [Internet]. 2008 Jun 3;18(11):814–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18514520

8. Yavari F, Mahdavi S, Towhidkhah F, Ahmadi-Pajouh M-A, Ekhtiari H, Darainy M. Cerebellum as a forward but not inverse model in visuomotor adaptation task: a tDCS-based and modeling study. Exp Brain Res [Internet]. 2016 Apr 26 [cited 2018 Jul 29];234(4):997–1012. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26706039

9. 't Hart BM, Henriques DYP. Separating predicted and perceived sensory consequences of motor learning. PLoS One. 2016;11(9).

10. Mostafa AA, Salomonczyk D, Cressman EK, Henriques DYP. Intermanual transfer and proprioceptive recalibration following training with translated visual feedback of the hand. Exp Brain Res. 2014;232(6):1639–51.

11. Mostafa AA, Kamran-Disfani R, Bahari-Kashani G, Cressman EK, Henriques DYP. Generalization of reach adaptation and proprioceptive recalibration at different distances in the workspace. Exp Brain Res [Internet]. 2015 Mar 6 [cited 2018 Aug 1];233(3):817–27. Available from: http://link.springer.com/10.1007/s00221-014-4157-9

12. Maksimovic S, Cressman EK. Long-term retention of proprioceptive recalibration. Neuropsychologia [Internet]. 2018 Jun [cited 2018 Aug 1];114:65–76. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29654883

13. Henriques DYP, Cressman EK. Visuomotor adaptation and proprioceptive recalibration. J Mot Behav. 2012;44(6):435–44.

14. Cressman EK, Salomonczyk D, Henriques DYP. Visuomotor adaptation and proprioceptive recalibration in older adults. Exp Brain Res [Internet]. 2010 Sep 18 [cited 2018 Aug 1];205(4):533–44. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20717800

15. Barkley V, Salomonczyk D, Cressman EK, Henriques DYP. Reach adaptation and proprioceptive recalibration following terminal visual feedback of the hand. Front Hum Neurosci [Internet]. 2014;8(September):1–11. Available from: http://journal.frontiersin.org/article/10.3389/fnhum.2014.00705/abstract

16. Salomonczyk D, Cressman EK, Henriques DYP. Proprioceptive recalibration following prolonged training and increasing distortions in visuomotor adaptation. Neuropsychologia [Internet]. 2011;49(11):3053–62. Available from: http://dx.doi.org/10.1016/j.neuropsychologia.2011.07.006

17. Nourouzpour N, Salomonczyk D, Cressman EK, Henriques DYP. Retention of proprioceptive recalibration following visuomotor adaptation. Exp Brain Res. 2014;233(3):1019–29.

18. Henriques DYP, Filippopulos F, Straube A, Eggert T. The cerebellum is not necessary for visually driven recalibration of hand proprioception. Neuropsychologia [Internet]. 2014 [cited 2018 Jul 29];64:195–204. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25278133

19. Cameron BD, Franks IM, Inglis JT, Chua R. The adaptability of self-action perception and movement control when the limb is passively versus actively moved. Conscious Cogn [Internet]. 2012;21(1):4–17. Available from: http://dx.doi.org/10.1016/j.concog.2010.11.006

20. Ostry DJ, Gribble PL. Sensory Plasticity in Human Motor Learning. Trends Neurosci [Internet]. 2016 Feb [cited 2018 Aug 1];39(2):114–23. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26774345

21. Clayton HA, Cressman EK, Henriques DYP. The effect of visuomotor adaptation on proprioceptive localization: The contributions of perceptual and motor changes. Exp Brain Res. 2014;232(7):2073–86.

22. Ruttle JE, 't Hart BM, Henriques DYP. The fast contribution of visual-proprioceptive discrepancy to reach aftereffects and proprioceptive recalibration. PLoS One [Internet]. 2018 [cited 2018 Jul 29];13(7):e0200621. Available from: http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0200621

23. Cressman EK, Henriques DYP. Reach Adaptation and Proprioceptive Recalibration Following Exposure to Misaligned Sensory Input. J Neurophysiol [Internet]. 2010;103(4):1888–95. Available from: http://jn.physiology.org/cgi/doi/10.1152/jn.01002.2009

24. Salomonczyk D, Cressman EK, Henriques DYP. The role of the cross-sensory error signal in visuomotor adaptation. Exp Brain Res. 2013;228(3):313–25.

25. Team RC. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing.; 2017. Available from: https://www.r-project.org/

26. Luke SG. Evaluating significance in linear mixed-effects models in R. 2017;(September 2016):1494–502.

27. Mostafa AA, 't Hart BM, Henriques DY. Motor Learning Without Moving. 2017 [cited 2018 Jul 30]; Available from: https://osf.io/zfdth/

28. Henriques DYP, Filippopulos F, Straube A, Eggert T. The cerebellum is not necessary for visually driven recalibration of hand proprioception. Neuropsychologia [Internet]. 2014;64:195–204. Available from: http://dx.doi.org/10.1016/j.neuropsychologia.2014.09.029

29. Sakamoto T, Kondo T. Visuomotor learning by passive motor experience. Front Hum Neurosci [Internet]. 2015 May 15 [cited 2018 Jul 30];9:279. Available from: http://www.frontiersin.org/Human_Neuroscience/10.3389/fnhum.2015.00279/abstract

30. Bao S, Lei Y, Wang J. Experiencing a reaching task passively with one arm while adapting to a visuomotor rotation with the other can lead to substantial transfer of motor learning across the arms. Neurosci Lett [Internet]. 2017 Jan 18 [cited 2018 Jul 29];638:109–13. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27988346

31. Ernst MO, Banks MS. Humans integrate visual and haptic information in a statistically optimal fashion. Nature. 2002;

32. 't Hart BM, Ayala M, Henriques DYP. Testing maximum likelihood estimates of hand position based on proprioception and prediction [Internet]. 2018 [cited 2018 Aug 1]. Available from: https://osf.io/wv3db/

33. Fiehler K, Rösler F, Henriques DYP. Interaction between gaze and visual and proprioceptive position judgements. Exp Brain Res [Internet]. 2010 Jun 29 [cited 2018 Jul 29];203(3):485–98. Available from: http://link.springer.com/10.1007/s00221-010-2251-1

34. Byrne PA, Henriques DYP. When more is less: Increasing allocentric visual information can switch visual-proprioceptive combination from an optimal to sub-optimal process. Neuropsychologia [Internet]. 2013;51(1):26–37. Available from: http://dx.doi.org/10.1016/j.neuropsychologia.2012.10.008

35. Jones SAH, Henriques DYP. Memory for proprioceptive and multisensory targets is partially coded relative to gaze. Neuropsychologia. 2010;

36. Beers RJ van. When Feeling Is More Important Than Seeing\rin Sensorimotor Adaptation. 2002;12(02):834–7.

37. Mikula L, Gaveau V, Pisella L, Khan AZ, Blohm G. Learned rather than online relative weighting of visual-

proprioceptive sensory cues. J Neurophysiol [Internet]. 2018 May 1 [cited 2018 Jul 29];119(5):1981–92. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29465322

38. Bastian AJ. Learning to predict the future: the cerebellum adapts feedforward movement control. Curr Opin Neurobiol [Internet]. 2006 Dec [cited 2018 Aug 1];16(6):645–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17071073

39. Shadmehr R, Krakauer JW. A computational neuroanatomy for motor control. Exp Brain Res. 2008;185(3):359–81.

40. Wolpert DM, Miall RC, Kawato M. Internal models in the cerebellum. Trends Cogn Sci. 1998;2(9):338–47.

41. Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT. Throwing while looking through prisms. II. Specificity and storage of multiple gaze-throw calibrations. Brain [Internet]. 1996 Aug [cited 2018 Aug 1];119 (Pt 4):1199–211. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8813283

42. Maschke M, Gomez CM, Ebner TJ, Konczak J. Hereditary Cerebellar Ataxia Progressively Impairs Force Adaptation During Goal-Directed Arm Movements. J Neurophysiol [Internet]. 2004 Jan [cited 2018 Aug 1];91(1):230–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/13679403

43. Criscimagna-Hemminger SE, Bastian AJ, Shadmehr R. Size of Error Affects Cerebellar Contributions to Motor Learning. J Neurophysiol [Internet]. 2010 Apr [cited 2018 Aug 1];103(4):2275–84. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20164398

44. Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, et al. Cerebellar regions involved in adaptation to force field and visuomotor perturbation. J Neurophysiol [Internet]. 2012 Jan [cited 2018 Aug 1];107(1):134–47. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21975446

45. Mostafa AA, Kamran-Disfani R, Bahari-Kashani G, Cressman EK, Henriques DYP. Generalization of reach adaptation and proprioceptive recalibration at different distances in the workspace. Exp Brain Res. 2014;233(3):817–27.