Transcriptome analyses of tumor-adjacent somatic tissues reveal genes co-expressed with transposable elements

Nicky Chung[1,*], G.M. Jonaid[1,*], Sophia Quinton[1,*], Austin Ross[1,*], Corinne E. Sexton[1],

Adrian Alberto[2], Cody Clymer[2], Daphnie Churchill[2], Omar Navarro Leija[2] and Mira V.

Han[1,3,§]

[1] School of Life Sciences, University of Nevada, Las Vegas, NV 89154, USA

[2] Department of Computer Science, University of Nevada, Las Vegas, NV 89154, USA

[3] Nevada Institute of Personalized Medicine, Las Vegas, NV 89154, USA

[*] These authors contributed equally to this work

[§] Corresponding author

Email addresses:

      NC: chungn1@unlv.nevada.edu

      GMJ: jonaid@unlv.nevada.edu

      SQ: quints1@unlv.nevada.edu

      AR: rossa38@unlv.nevada.edu

      CS: csexton@unlv.nevada.edu

      AA: albera3@unlv.nevada.edu

      CC: clymer@unlv.nevada.edu

      DC: churchi2@unlv.nevada.edu

      ON: navar106@unlv.nevada.edu

      MVH: mira.han@unlv.edu

Abstract

Background

Despite the long-held assumption that transposons are normally only expressed in the germ-line, recent evidence shows that transcripts of transposable element (TE) sequences are frequently found in the somatic cells. However, the extent of variation in TE transcript levels across different tissues and different individuals are unknown, and the co-expression between TEs and host gene mRNAs have not been examined.

Results

Here we report the variation in TE derived transcript levels across tissues and between individuals observed in the non-tumorous tissues collected for The Cancer Genome Atlas. We found core TE co-expression modules consisting mainly of transposons, showing correlated expression across broad classes of TEs. Despite this co-expression within tissues, there are individual TE loci that exhibit tissue-specific expression patterns, when compared across tissues. The core TE modules were negatively correlated with other gene modules that consisted of immune response genes in interferon signaling. KRAB Zinc Finger Proteins (KZFPs) were over-represented gene members of the TE modules, showing positive correlation across multiple tissues. But we did not find overlap between TE-KZFP pairs that are co-expressed and TE-KZFP pairs that are bound in published ChIP-seq studies.

Conclusions

We find unexpected variation in TE derived transcripts, within and across non-tumorous tissues. We describe a broad view of the RNA state for non-tumorous tissues exhibiting higher level of TE transcripts. Tissues with higher level of TE transcripts have a broad range of TEs co-expressed, with high expression of a large number of KZFPs, and lower RNA levels of immune genes.

Keywords

Transposon, TE, L1HS, RNA-seq, co-expression, mitochondria, KRAB zinc finger

Background

Although transposable elements (TEs) have been studied for a long time, their

ubiquitous and highly tissue-specific expression patterns are starting to be appreciated

only recently. The fact that TEs compose close to 40% of the human genome is

frequently emphasized, but the fact that there is observable amount of TE derived

transcripts in human RNA-seq data has mostly been ignored or regarded as a nuisance

without any functional relevance [1]. LINE elements have long been thought to be

expressed only in the germline cells [2–4]. But, both full-length and partial transcripts of

LINEs are frequently found in the somatic cells [4–6] with large variation in expression

levels across tissue types, and among different individuals [7]. The level of TE

expression is especially pronounced in cancer cells [8–11], and cell lines [12], but are

also observed in neurogenesis [13] and normal somatic tissue. Faulkner et al. in 2009,

was the first study to provide a global picture of the significant contribution of

retrotransposons to human transcriptome in multiple tissue types [14]. This report

showed that 6-30% of transcripts had transcription start sites located within

transposons, and these transposons were expressed in a tissue-specific manner and

influenced the transcription of nearby genes. The results were extended by Djebali et al.

in 2012 showing again the tissue-specificity of transposon expression, and that most of

these transcripts are found in the nuclear part of the cell [15]. In addition to the tissue-

specific expression of TEs, important regulatory roles for TEs are emerging (reviewed in

[16]). Observations include contribution to transcription start sites [14], source of

transcription factor binding sites [17], source of long non-coding RNAs [18], active

transcription during early development [19], and even critical function similar to long

non-coding RNAs that guide chromatin-remodeling complexes to specific loci in the

genome [20].

Although there are many reports of TE expression in the somatic cells, there is still a

large gap in our understanding of how TE expression is repressed and de-repressed in

human somatic cells. Based on what we have learned so far, TE expression is regulated

through multiple layers, consisting of transcription factors, epigenetic modification,

PIWI-interacting RNAs (piRNAs), RNA interference (RNAi), and posttranscriptional host

factors. Recently, two different approaches of genome-wide screening have identified

proteins that regulate different aspects of the activities of LINE elements. CRISPR–Cas9

screen was used to identify proteins that restrict LINE activity [21]. The protein MORC2

and the human silencing hub (HUSH) complex was shown to selectively bind

evolutionarily young, full-length LINEs located within euchromatic environments, and

promote deposition of histone H3 Lys9 trimethylation (H3K9me3) for transcriptional

silencing [21]. And through proteomics approaches, two studies have recently identified

the localization of ORF1 and ORF2 proteins and its interacting partners [22], and the

timing of the entrance of the ORF2 protein complex into the nucleus [23]. But, no study

has yet examined the correlation in transcript levels of host mRNA and transposon RNA.

Recently, high-throughput RNA-seq data of various types of cancer samples and their

normal counterparts have become available in The Cancer Genome Atlas (TCGA) [24–

26]. By focusing on the non-tumorous tissue samples from TCGA, we can access

thousands of natural experiments across various types of tissues that show variation in

TE transcript levels, and obtain a global picture of TE expression and regulation in

humans. An important strength of the TCGA dataset is the large number of samples

collected for each tissue type and the high depth of the RNA-seq experiment, with a

median of about 150M reads per sample, which is several times larger than a usual

RNA-seq library. The variation in TE transcript levels observed in multiple samples

within each tissue, allowed us to analyze the co-expression patterns between host genes

and TEs for the first time. We hypothesized that genes that regulate the transcription

level of TEs would show correlation in expression levels with the TE transcripts. Since

the samples are collected from fresh-frozen tissues, TE transcript levels are observed *in*

vivo, complementing the studies that focus on retrotransposition assays or transposon

expression in human cell lines.

We first summarize the survey of TE expression variation found in the RNA-seq data

from 697 samples of cancer-adjacent non-tumorous tissue. We confirm the earlier

findings that TE expression varies across tissue types. Transcript levels of individual TE

loci are highly tissue specific and within each family only a few individual loci are highly

expressed, contributing to the bulk of the transposon transcripts at the family level. We

also find large variation in total TE transcript level across individual samples within

each tissue type.

Although, transposons have strong tissue-specific patterns at the locus level, we also

found that the majority of TEs show global co-expression at the family level across

samples. By analyzing the co-expression between these TEs and individual genes, we

found co-expression modules of TEs and genes replicated across tissues.

Results

TE derived transcripts are quantified across 16 tissues and 697 samples of tumor adjacent controls.

We re-aligned and quantified TE derived transcripts from the RNA sequencing data of

697 samples across 16 tissues collected as non-tumorous controls for the TCGA project

(supplementary Table 1). The library sizes for these samples range from 50M reads at

the minimum, to up to 390M reads, with a median at about 149M reads (75M pairs).

Although all tissues included in this study were sequenced using the HiSeq 2000

platform, esophagus and stomach samples were sequenced separately at British

Columbia Genome Sciences Centre (BCGSC), with higher sequencing depth on average

(median 227M reads). The proportion of reads that do not map to annotated genes were

different between the later samples sequenced at University of North Carolina at Chapel

Hill (UNC), and the earlier BCGSC sequenced samples, with BCGSC samples having more

reads (median 177M) not mapping to annotated genes and discarded, while UNC

samples had less reads discarded (median 97M), possibly due to the difference in poly-A

enrichment protocol (MultiMACS mRNA isolation kit *vs.* TruSeq RNA Library Prep Kit

[27]) among many other differences, including read length. Because of these differences,

when comparing across tissue samples, we had to consider esophagus and stomach

tissues separately, and they could not be compared against the rest of the tissues.

Despite the differences in the overall sequencing depth and overall proportion of reads

mapping to genes, we found that the DESeq2 normalization method normalizes the

reads effectively and the correlation due to library size disappears after normalization

within tissues (Supplementary Figure 1). Our co-expression analysis is done within each

tissue separately. We also replicate our results found in esophagus and stomach with

similar results found in at least one other tissue.

Although we find reads mapping to TEs in all the samples that we have examined, the

overall transcripts coming from TEs are still a relatively tiny proportion of the total

library. The total number of reads mapping to TEs ranged from 137K to 2.1M with a

median of 615K for UNC samples, and ranged from 282K to 3.3M with a median of 835K

for BCGSC samples. This count excludes some of the potential read-thru transcripts as

described in the next section. The total read counts across all TEs amount to about 1.2%

(UNC) or 1.7% (BCGSC) of the total reads mapping to known genes in non-tumorous

tissue samples.

TE reads originating from pre-mRNAs or retained introns are corrected by
comparing the read depths of the flanking introns.

There have been previous reports of transposon reads coming from pre-mRNA or

retained introns in the mature RNA of genes that contain TE sequences in their introns

[28]. The extent of this problem can be partially estimated by comparing the read

depths of the transposon to the read depths of the flanking introns. If the reads mapping

to TEs are part of the pre-mRNA or retained introns, we should see continuous mapping

of reads that span the introns flanking the TE of interest, and observe reads that map

across the intron-TE boundaries. We can also partially correct for this problem by

utilizing the read depths in the flanking introns to proportionally reduce the number of

total reads mapped to TEs. The approach is described below.

$$R_{IL} = \frac{count_{IL}}{len_{IL} - read\_len}$$

$$count'_{TE} = \begin{cases} count_{TE} - count_{TE} \times \dfrac{R_{IL} + R_{IR}}{2R_{TE}} & , if\ \dfrac{R_{IL} + R_{IR}}{2R_{TE}} < 1 \\ 0 & , otherwise \end{cases}$$

TE : focal TE
IL : intron left to TE. IR : intron right to TE.
R_{IL} : read depth of the intron left to the TE
$count_{IL}$: read counts mapped to the intron left to the TE (includes multi-mapped
reads)
len_{IL} : length of the left intron
$read\_len$: length of the sequencing read
$count_{TE}'$: count of reads mapped to TE after the correction.

We modified the software TEtranscripts [29], following this approach, to discount the

TE read counts based on the read depths of the surrounding introns. By looking for

large differences after correcting by flanking read depth, we identified TEs that are most

frequently transcribed as part of the introns (Table 1). We also found cases where the

method corrected for erroneous TE quantifications due to TEs embedded within long

non-coding RNAs (lncRNAs). For example, an AluSx1 element on chromosome Y at

position 21153222 (AluSx1_dup59209) had very high transcript levels with an average

read count of 18863 in thyroid and head and neck tissue, but the Alu element is

embedded in a lncRNA gene called *TTTY14*. The reads mapping here are counted as

AluSx1 transcripts based on the UCSC TE annotation, but in the alignment, we see that

there are reads spanning the boundaries of AluSx1_dup59209, and almost all the reads

mapped in the region are uniquely mapped reads. It looks to be a case of an *Alu*

domestication, where an *Alu* insertion or a secondary duplication of an original *Alu*

insertion became part of a testis specific RNA gene [30]. Three examples of AluSx1, L2a,

and L1MA7, where the read counts for the transposons are reduced to zero are

visualized in Supplementary Figure 2. AluSx1_dup59209(chrY:21153222-21153521) is

embedded within an exon of gene *TTY14*. L2a_dup21781(chr2:113980079-113981081)

is embedded in an intron of *PAX8*. L1MA7_dup4297 (chr8:134015602-134015763) is

embedded in an intron of gene *TG*. In all three cases, read counts for the focal TEs were

reduced to zero after the correction described above, and the reads mapping to these

TEs did not contribute to the overall TE family count. If one is interested in transposable

element transcript level that is not part of a longer RNA molecule, it is important to take

into account the read depths of the flanking introns or exons, especially the latest non-

coding RNA gene annotations, when quantifying repeat element transcripts in the

genome.

Relying on uniquely mapped reads for repeat quantification results in
quantification biased for mappable elements.

Due to the difficulty of mapping reads to repeat elements, one of the approaches taken

is to count only the reads that map to a unique position in the genome. But this

approach has repeatedly been shown to produce results that are worse than

expectation-maximization [31,32], and can lead to serious biases. If we only count

uniquely mapped reads in our analysis, not only did we throw away from 10.7% to up to

45% (median 14.2%) of the total TE transcripts, we threw away data in a biased

manner, such that we ended up "quantifying mappability" instead of "quantifying

transcripts". This problem is especially pronounced when quantifying the young and

active L1HS element. To assess the effect of alignment on quantification, we tried two

different alignments, one based on the STAR aligner with up to 200 multi-mapped

positions, and the other based on Bowtie1 with only a single best alignment position,

discarding all reads that do not have a unique best mapping. Figure 1 shows two cases

that illustrate the limitations of either of the approaches. Figure 1 a. shows an example

of a full length L1HS locus on chromosome X: 75453754-75459553 with high

mappability (48base mappability shown at the top of the panel) due to many

accumulated mutations in its sequence. The top panel shows the bowtie1 alignment,

allowing only uniquely mapping reads with a single best hit, and the bottom panel

shows the STAR alignment with multi-mapping up to 200 mappings per read. In the

STAR alignment, we can see erroneously split read alignments at the 3' end that result

in reads mapping across greater than 10K distances, that shows a limitation of a splicing

oriented alignment software. The transcription for this element does not start at the 5'

end of the full length, but there is clear and unambiguous transcription starting from

about 1500 bases in, that are congruent between both alignments. In Figure 1 b. it

shows another full length L1HS locus on chromosome X: 11953208- 11959433, this

time a young element with very low mappability. Comparing the top and bottom panels,

we can see that with the unique mapping we are ignoring all the reads that are perfectly

mapping to this locus, but also map to multiple other locations. There is a huge pile-up

at the 5' end of the full length. If we look at the reads mapping to the 5' end of this locus,

their NH tags show numbers ranging from 2 to 4, meaning that they are mapping to two

to four alternative locations in the genome. Considering that L1HS loci containing the 5'

ends are more likely to be full length elements, these reads are more likely to be coming

from one of the few full length L1HS loci in the genome, but, we end up ignoring these

reads if we are only counting uniquely mapping reads. On the other hand, with multi-

mapping, we end up quantifying with large uncertainty on whether the reads piled up in

this region are really transcribed from this particular locus. This is evident by the small

regions of extremely high pile-ups that reflect fragments that are found in the genome

with high frequency. Although, we should point out that we don't count all the reads

aligning here at face value, since the expectation maximization algorithm will down

weigh the counts of reads by the number of places it maps to.

Comparing the mappability of the two examples, we can see that the uniquely mapping

approach preferentially counts reads coming from older TE loci with higher

mappability. This can also be shown by correlating the locus level read counts for each

L1HS element against the length of uniquely mappable positions in each TE locus

(Figure 2 a.). For the unique mapping approach, we see there is significant correlation

between the locus level quantification and the total length of uniquely mappable

positions within that locus (*p-value* = 5.523e-07 for sum of read counts across all

samples, and *p-value* = 1.072e-13 for maximum read count among all samples). The

multi mapping approach with Expectation Maximization does not show that bias for

uniquely mappable regions (*p-value*= 0.60 for sum and *p-value* = 0.08 for max) (Figure 2

b.).

Consequently, there is limited correlation in the locus level quantification of L1HS

between the uniquely mapped reads and the multi-mapped reads (Figure 2 c.). This

shows the difficulty of quantifying young active elements, such as L1HS using genome-

wide RNA-seq data. Due to these limitations, analysis on L1HS in this study has been

done at the family level. The family level quantification of L1HS still shows variability

based on the read mapping approach (Supplementary Figure 3 e.), but it shows stronger

correlation than the locus level quantification. We still wanted to utilize the abundance

of RNA-seq data available for studying L1HS transcription, and glean information on

L1HS from these data. Based on the observation that 3' ends of L1HS are frequently

represented in fragmented L1HS loci, while the 5' ends are more frequent in full length

L1HS loci, we decided to use the read counts of the 5' end of the element as a measure

that better represents the transcript level of full-length L1HS transcripts in the sample.

All the following analysis on L1HS expression are based on the reads mapped to L1HS

sequences in the genome that align with the first 300 bases of the 5' end of the L1HS

consensus sequence and allowing for multi-mapping.

On the other hand, we found that quantification of older elements showed very strong

correlation between the two approaches, unique mapping and multi-mapping, reflecting

the higher mappability of older elements in the genome (Supplementary Figure 3). For

the locus level co-expression analysis with the Zinc Finger Proteins, we limited our

analysis to older elements that are 100% uniquely mappable across its sequences with a

48-base read length. All of our main results are qualitatively replicated in the data with

uniquely mapped reads aligned with bowtie, except for the results regarding the 5' end

of L1HS expression.

TE expression shows tissue-specific expression patterns at the locus level
among somatic tissues

There have been multiple reports of tissue specific expression of TEs in the human

genome, starting from Faulkner et al. in 2009 [14] to Philippe et al. 2017 [12] more

recently. We also found highly distinct tissue specificity in TE transcripts in the TCGA

data, such that that we could cluster each sample into their broader tissue groupings,

based on locus level TE expression patterns alone without relying on any genes at all.

Figure 3 shows the clustering of tissues for family level and locus level quantification of

LTRs, DNA transposons, SINEs and LINEs. We used normalized mutual information

between the different clustering results and the ground truth (the true tissue group) to

evaluate the quality of clustering. Normalized mutual information was compared for

clustering results based on gene expression, family level TE expression, locus level TE

expression and random assignments. We found that the locus-level TE expression was

as predictive of tissue groupings as the genes (Figure 3.i, Supplementary Table 2). LTRs,

DNA transposons, LINEs and SINEs gave similar clustering accuracy as the genes. The

TE family expression levels did not have enough information to cluster the tissues

correctly. We note here, that these are loci selected by the rank of variance in $\log_2$

normalized read count across all samples regardless of tissue type, and we haven't done

any differential expression analysis to identify the markers that are the most

informative for accurate tissue classification. Thus, the classification performance we

observe here is not the optimal performance that we could get if we were to decide on

the markers based on a trained classifier. When we excluded TEs that are within 1K,

10K, and 100K of the start and ends of the genes, the accuracy declined, so part of the

tissue specificity is due to co-location with tissue specific genes. But, even when relying

on TEs 100K away from any known genes, we saw that tissue specific information was

largely retained. On the other hand, when we focused on younger elements, HERVs

within LTRs and young L1s within LINEs, there was a large reduction in information

content, especially for young L1s. Clustering based on locus level expression for L1HS,

L1PA2 and L1PA3 was not any better than clustering based on family level expression

of all LINEs. We suspect this is due to the lower locus level mappability and large

uncertainty in locus level expression quantification for young L1 elements. Figure 4

shows fifteen representative TE loci that show tissue specific expression. These loci are

chosen from TEs that are 100K away from the start or end of any gene.

The granular levels of locus specific TE expression contained tissue-specific

information, but, the overall transcript level of TE classes did not show significant

variation across tissues (Figure 5 a.). The higher expression levels for TEs seen for

esophagus and stomach is confounded with the differences in sequencing protocol

described above, so they are not directly comparable to the rest of the tissues. When

focusing on 300 bases in the 5' end of L1HS, it showed some variation across tissues

with higher levels in the head and neck tissues and lower levels in the liver, consistent

with the previous observation in adult human tissues [6] and in human cell lines [12],

albeit with large within tissue variance (Figure 5 b.). Although, we cannot directly

compare L1HS expression in esophagus and stomach to the rest of the other tissues, we

can tell that there is clear transcription of the 5' end of L1HS in esophagus and stomach.

Figure 5 b. shows the normalized read counts in log2 scale, with a median of more than

500 reads mapping to 300 bases at the 5' end of L1HS for esophagus and stomach

(median library size 227M reads). There have been observations of full length L1HS

expressed in the adult esophagus and stomach tissue, at about 80% and 150% relative

to the levels in HeLa cells [6], and active L1 retrotransposition in premalignant

precursor legions of esophageal adenocarcinoma [33]

Co-expression analysis of intergenic TEs identifies core TE modules and
correlated Zinc Finger Proteins.

Co-expression network analysis is an appropriate approach to examine the co-

expression across different TE families and host genes together. In order to identify the

common gene/TE modules that are correlated across different tissues, we did a

consensus network analysis across tissues using the weighted gene co-expression

network analysis in the WGCNA package [34]. For the TE family transcripts in this

analysis, we only included intergenic TEs, *i.e.* we only counted reads mapping to TEs

that are 1Kb away from any start and end of known genes. We identified 61 modules

across 11 group of tissues, combining certain tissue types together as a broader group

(colon and rectum, esophagus and stomach, kidneys, lungs). Among the 20531 genes

and 992 TE families that were quantified in the 697 samples, 18670 genes and 923 TE

families had enough expression level and variation to be included in the network

analysis. Among those 19593 genes and TEs, 9599 genes and 658 TEs were clustered

into a module of co-expression, while 9336 did not belong to any defined module. The

list of modules, correlations between the modules, and topological adjacency matrix

that defines the modules are visualized for the breast tissue in Supplementary Figure 4.

Visualization for other tissues were similar, as we looked for consensus modules across

all tissues. There were only a few modules that contained TE transcripts: only seven

modules contained more than ten TE families within the module. Supplementary Table

3 shows the distribution of TE families in these seven modules. We considered modules

M8, M21, M38 and M45 as core TE modules, as their membership mainly consisted of

TE families as the majority (marked by * in Supplementary Figure 4).

The correlation between closely related TE subfamilies is expected because reads from

transposons that map to sequences that are indistinguishable between subfamilies are

assigned to multiple subfamilies with proportional weight by TEtranscripts using an

Expectation-Maximization algorithm. Closely related families such as L1HS and L1PAs

also share common regulatory elements at the 5' end. But, we find that the correlated

TE families in a TE module span different classes of TEs, and are replicated even when

counting uniquely mapped reads only. Considering there is no sequence similarity

between the SINEs, DNA transposons, LTRs and the LINEs, the correlation among these

diverse class of transposons is probably due to a common regulation, or dys-regulation,

that is de-repressing these transposons at the same time. There have been reports of

such co-expression of ERVs and LINEs in cancerous tissues [35,36], possibly through

concordant hypomethylation [37].

There was one class of host genes that were frequently found as members of the TE co-expression modules, and they were the KRAB Zinc Finger Proteins (KZFPs). Table 2 shows the list of KZFPs that were identified as TE module members.

Expression of immune genes are negatively correlated with intergenic TE expression.

Once we identified modules consisting mostly of transposon families, we also examined whether any co-expression modules were negatively correlated with TE modules. We found two modules, M33 and M35, that showed consistent negative correlation across tissues. The genes included in these modules were genes involved in innate immune system, interferon signaling, immunoproteasome, etc (Figure 6). Figure 6 shows the enriched annotation terms detected for both modules through the Reactome database, the top 30 genes with highest module membership for the two modules, and the correlation plot between TEs in the TE modules M8, M21, M38 and M45, and genes in module M33 and M35 in the tissues breast (Figure 6d) and esophagus/stomach (Figure 6e). Only genes and TEs that show greater than 0.6 Pearson correlation with the representative profile of the module in all tissues have been included in the correlation plot. We observe high correlation within groups and contrasting negative correlation between groups.

Co-expression analysis including intronic TEs reveals negative correlation between intronic TE expression and mitochondrial gene expression.

When we include intronic TE transcripts in the overall TE expression levels, the co-expression analysis led to a different picture from the analysis of intergenic TEs. When intronic TEs are included, a single module, N1, emerges as the dominant TE module, containing 612 out of 848 TE families (72%) that was assigned a module membership. In fact, N1 consists of 72% of all TE families but only 2% of all genes.

Table 3 shows genes that are significantly correlated with module N1 in multiple

tissues. A pattern immediately noticeable is that there are many pseudogenes, intronic

transcripts, antisense RNAs, and long non-coding RNAs on the list. It looks like with

intronic TEs, we are detecting a cell state that is dysregulated in splicing or mRNA

quality control, and as a result, we are seeing a global elevation of pervasive

transcription that is generally non-functional. Multiple protein coding genes on the list

are involved in mRNA splicing regulation, such as *NCRNA00201*, an isoform of HNRNPU

which showed strong correlation with the intronic TE module in seven different tissues,

as well as *CCNL2*, *LUC7* and *LUC7L3*, perhaps as a response to the dysregulated splicing.

Another interesting gene in the list is *NKTR*, hinting at the presence of immune cells in

the tissue samples with high intronic TE expression. This is in contrast with the negative

correlation we observe with immune genes and intergenic TE expression.

The module that was negatively correlated with the intronic TE module (N1) included

co-expression clusters consisting of mitochondrial proteins and ribosomal proteins

(N4). N4 was the only module that was consistently negatively correlated with N1 with

less than -0.7 correlation coefficient across all tissues. Figure 7 shows the correlation

plots between TEs in N1, and genes in the mitochondrial gene module N4, for breast and

esophagus. Enriched annotation terms for the genes found in the Reactome database

are centered around translation and mitochondria. One intriguing possibility may be

that the failed splicing and mRNA control is leading to a suppression of translation that

in turn leads to reduced RNA levels of mitochondrial genes and ribosome.

We again saw an enrichment of KZFPs as members of the intronic TE module N1, and

another module N10, that was positively correlated with N1 (Supplementary Table 4).

The list of KZFPs had some overlap with the KZFPs co-expressed with intergenic TE

modules, but there were some differences as well. We combined the 22 KZFPs in

module N1 and N10 and examined whether there were any common transcription

factor binding for these genes found in the ENCODE ChIP-seq data with the EnrichR

database [38]. The region near these KZFPs were enriched with binding of *GABPA,* a

regulator of nuclear encoded mitochondrial genes, in multiple cell lines (Supplementary

Figure 5). This was interesting, given the negative correlation observed with intronic

transposons and nuclear encoded mitochondrial gene expression described above.

Genes co-expressed with L1HS include genes regulating major signaling
pathways, chromatin, and stress response.

Given the interest in the active element L1HS, and the uncertainty in L1HS

quantification, we decided to limit the quantification to the 5' region of L1HS, and

examine the host genes that are specifically correlated to the expression of 5' region of

L1HS without regard to the co-expression modules. In order to control for the

correlation with other TEs, especially intronic TEs, we included the representative

profile of N1 as a covariate into our linear model. One concern with co-expression

analysis is positional overlap. There were 14 genes that overlapped with the L1HS loci

we were counting the reads from. Only 1 of the 14 genes, *RAB3GAP2*, showed significant

correlation with L1HS 5', and was removed from the final list. 56 genes were identified

as negatively correlated, and 77 genes were identified as positively correlated with

L1HS 5' in at least two tissues (Figure 8, Supplementary Table 5). Notable genes include

RASA1, RASA2, *RRAS, EGFR* and *MAPK1*, in the Ras-MAPK pathway, *ECSIT, TAB3* and

TRAF6, regulators of the NF-κB pathway, *RNASEH2C*, a known L1HS repressor [39],

TET2, known to bind to and demethylate young L1s [40], *THAP7*, a histone tail binding

transcription repressor [41], and *DDI2*, a protease that cleaves and activates

NFE2L1/NRF1 [42]. Multiple genes in the respiratory electron transport pathway,

ECSIT, NDUFA1, NDUFA8, NDUFB10, NDUFB8, SURF1, UQCR11, UQCRB, were negatively correlated with L1HS 5', even after controlling for the covariation with intronic TEs, N1. Whole list of genes are reported in Supplementary Table 5.

We checked whether the list of our negatively correlated genes were overlapping with the genes identified through CRISPR–Cas9 screen [21]. Of the 56 negatively correlated genes, three genes, *RNASEH2C*, *HAUS7*, *RNF166* were also on the list of 253 secondary screen hits. There was no overlap among the 77 positively correlated genes.

We also checked whether there were transcription factors known to bind to L1HS sequence [43] in our list. Of the 77 positively correlated genes, four genes, *YY1*, *REST*, *ELF1*, *ZBTB33* were identified to bind to L1HS [43]. There was no overlap among the 56 negatively correlated genes. To check if the same transcription factors are regulating the correlated genes and L1HS, we also checked what kind of TF binding is observed in the upstream of our correlated genes. There were a few enrichment of ENCODE transcription factor binding upstream of our list of correlated genes (Supplementary Figure 6), but except for YY1, the enriched TFs did not overlap with the list of Sun et al. [43]

TE module expression is correlated with radiation exposure in thyroid tissue.
We examined whether any of the clinical variables were associated with the TE module expression or the L1HS expression levels. We tested the variables age, days to death, pathological stage, T staging, N staging, M staging, gender, radiation and race for each tissue type. No variable was found to be associated with L1HS 5' expression. Radiation therapy was the only clinical variable associated with module N1 (intronic TE module) expression in the non-tumorous tissue of thyroid (p-val = 0.00894, Supplementary Figure 7).

Co-expressed TEs and KRAB-ZFPs show limited overlap with ChIP-seq binding

Based on the positive correlation observed among KZFPs and TE modules, and existing literature on the role of KZFPs for TE repression, we decided to examine the correlated expression of all pairs of 979 TE families and 366 KZFPs. The most striking pattern observed was that KZFPs and TEs show overwhelmingly positive correlation and little negative correlation. Chromosome 19, where the majority of the KZFPs are clustered, is also the chromosome with the highest density of transposable elements. This unique structure of chromosome 19 may lead to TEs embedded in KZFP genes erroneously identified as co-expressed. We avoid the confounding effect of positional overlap between TEs and KZFPs by only counting reads mapping to TEs that are in the intergenic region 1Kb away from any genes. There may be residual correlation due to shared genomic environment of a larger scale, such as the chromatin state. But, that doesn't explain all the positive correlation, because, when we look at the locus level correlation, we find that the individual TE loci correlated with the ZNFs are scattered across all chromosomes, and not necessarily enriched on chromosome 19.

The co-expression between KZFPs and TEs were observed across almost all TE families, as 794 TE families had at least one co-expressed ZFP in at least one tissue. Certain ZFPs, such as *ZNF621*, *ZNF780B*, *ZNF84*, *ZNF33A*, and *ZNF662*, showed correlation with a wide range of TE families in multiple tissues. TE-KZFP pairs, HERVK14-int:*ZNF814*, MER57A-int:*ZNF621*, and MSTB-int:*ZNF41* were the most frequent pair-wise co-expression observed between TE families and KZFPs, found positively correlated in six different tissues. The ZFPs that were negatively correlated with TEs were *ZNF511* and *ZNF32*, but, they are not classified as KZFPs as they do not have a KRAB domain.

We looked at the family level co-expression between TE families and KZFPs and tested

the overlap against the KZFP bound TE family enrichment reported in the ChIP-exo

study (GSE78099 [44]). We found that there is a statistically significant association

between co-expression and binding (*p-value* < 2.2e-16). But, the number of overlapping

pairs were very small. Figure 9 shows the overlap between co-expression and binding

enrichment. We only mark the co-expression found in at least two tissues, and we have

omitted the TE-KZFP combinations that have neither co-expression nor binding

enrichment from the figure. The total combinations tested that overlap between the two

datasets is 200,889 (221 KZFP x 909 TE families). 4138 pairwise co-expression was

observed in at least two tissues. Of those, only 119 was enriched for binding in the ChIP-

exo study [44].

To check how the co-expression is observed at individual TE loci, we took the TE family-

KZFP pairs that show correlated expression, and further tested co-expression between

individual TE loci of the correlated TE family against the KZFP of interest. With

correlations at the locus level, we were able to examine the locus level co-expression

and compare it directly to the binding peaks reported in Imbeault et al [44]. Of the 6258

co-expressed TE loci where the KZFP had been assayed with ChIP-exo, there were only

4 that were bound by the same KZFP. We do not have a good explanation for why there

is a lack of overlap between co-expression and binding at the locus level, when there

was at least some amount of overlap at the family level. It looks like the co-expression

we observe is a result of indirect interactions, and not necessarily direct binding.

We also observed that at the locus level, there was not a lot of overlap between the TEs

that are bound by KZFPs in [44], and the TEs that are expressed in the TCGA non-

tumourous tissues, regardless of the co-expression relationship with KZFPs. Here

"expressed" means there is at least one sample in our data with more than five reads

mapped to the TE locus, and "binding" means that there is a peak detected in the

GSE78099 ChIP-seq data overlapping with the TE locus with a +-250bp buffer. Figure 10

shows the overall breakdown of the 4.5 million transposons annotated in hg19 UCSC

Repeatmasker track. Statistically, there is more overlap than expected (*p-value* < 2.2e-

16) between binding and expression, but, the overall proportion of TEs that are both

expressed in at least one sample and bound by at least one ZFP are a tiny proportion

(2.6%) of all TEs in the genome.

One interesting pattern did emerge when we examined the overlap with epigenetic

marks of candidate cis-Regulatory Elements defined in the ENCODE data [45]. We were

more likely to see an enhancer-like mark (DNase + H3K27ac) for TE loci that are

expressed compared to non-expressed TEs, and we were more likely to see a promoter-

like mark (DNase + H3K4me3) for ZFP bound TE loci compared to TEs with no binding

(Figure 10). The 2.6% of TE loci that are expressed in at least one sample and bound by

at least one ZFP showed the highest proportion of both promoter-like marks and

enhancer-like marks. When we divide the TE loci into gene regions (genes including

introns and +- 1K flanking region) and intergenic regions (1K away from start and end

of genes), the overall pattern remained the same, except that TE loci were twice as likely

to be expressed if they are close to genes compared to intergenic regions, and the TE

loci were twice as likely to be overlapping with the promoter-like marks

(Supplementary Figure 8). The enhancer-like marks showed no difference between

gene regions and intergenic regions, and the CTCF marks increased in the intergenic

regions.

Discussion

Limitations to the quantification and correction

Quantifying transposon transcripts is a difficult problem, due to their ambiguity in short read mapping because of repeated content in the reference genome. Current state of the art methods rely on Expectation-Maximization to account for the uncertainty in multi-mapped reads [29]. Focusing only on uniquely mapped reads doesn't really solve this problem, and will lead to biased quantification, favoring older elements with higher mappability. Scott et al. have demonstrated that by relying on unique mutations found within individual L1HS loci, and by including sequences of non-reference polymorphic L1HS loci, it is possible to identify the source of the L1HS activity with substantial success [46]. But, in our study we did not attempt to identify the individual loci of L1HS transcription, and instead focused on the totality of reads mapping to regions of annotated L1HS that align to the 5' end of L1HS consensus sequence.

Another complication in TE transcript quantification is that TEs are frequently embedded within introns that are transcribed before they are processed, or sometimes fail to be spliced out, or embedded within exons or non-coding RNAs that are expressed in different conditions [28]. To account for this source of error, we introduced a method to correct for TE reads coming from retained introns or pre-mRNA. Although we observed large corrections for specific transposable elements embedded within introns, the correction is not complete. We can tell this from the observation that the co-expression profiles of intronic TEs are different from the co-expression profiles of intergenic TEs away from the genes. The genes co-expressed with intronic TEs include pseudogenes, intronic transcripts, anti-sense transcripts and genes with functions in splicing. A more accurate approach would be to correct for the read counts from retained introns before the EM algorithm based on the read depth of uniquely mapped

reads, and then run EM based on the corrected counts. But, estimating the read depth of

the repeat region using uniquely mapped reads is a difficult problem. The effective

length of the uniquely mapped region is difficult to estimate, because again mappability

varies from locus to locus for any TE, depending on the unique mutations it has

accumulated. So, for this study, we decided to use the easier approach to run EM first,

and probabilistically assign the TE reads, and then correct based on the expected read

depth across the length of the TE locus. An important future study would be to study the

mappability of individual TE loci carefully, including the known polymorphic sites, and

to design a software for TE quantification that can take into account the mappability of

each locus in its EM algorithm, as well as correct for the retained introns while

considering the effective length of the uniquely mappable region within the TE.

Despite these limits, the main results of co-expression analysis were not affected by the

quantification. Most of the results in the paper were replicated when quantification was

done on uniquely mapped reads only. The only results that changed between the multi-

mapped approach vs. the unique mapping approach were the genes correlated with the

L1HS 5' expression level. For those, we decided to report on results from the multi-

mapped reads rather than the unique reads, because of the bias of the uniquely mapped

reads we described above.

Stress, immune response and TE expression

Initially, when we started the project, our goal was to identify candidate genes involved

in transposon control, based on the co-expression analysis. But, once the analysis was

done, the results were pointing to what induces TE expression, rather than what

suppresses TE expression. Among the genes known to function in transposon control,

RNaseH2C (Figure 8), *HAUS7*, and *RNF166* [21] showed negative correlation with L1HS.

But several well-known genes with functions in transposon control, *e.g. MORC2, SIRT6,*

KAP1, SAMHD1, MOV10, ZAP, C12orf35 (human ortholog of *RESF1*), etc. are missing in

our list of significantly correlated transcripts. Instead, the major theme that emerged

from our results is signal transduction, immune response, and stress response as seen

in the correlation between L1HS and *DDI2*, Ras-MAPK and NF-κB pathway. In humans,

various stresses have been shown to induce LINE1 transcription or activation including

chemical compounds [47–49], radiation [50,51], oxidative stress [52] and aging [53].

Most of these studies have observed L1 activity *in vitro*, by exposing cultured cells to

stress factors and assaying the retrotransposition activity.

The negative correlation we find between TE expression and immune gene activity has

been reported before in gastrointestinal cancer samples. Jung et al. have shown that the

L1 retrotransposition rate is inversely correlated with expression of immunologic

response genes [54]. Here, we extend those results and show that the negative

correlation between TE expression and immune response is a pattern found in non-

tumorous samples as well, across different tissues and different classes of TEs. This

relationship is confusing, since it is opposite of the positive correlation we find between

L1HS and NF-κB pathway genes (Figure 8), and opposite of the pattern observed in

several cancer studies, where DNA hypomethylation and expression of endogeneous

retrovirus activates interferon signaling [55–57]. Immune active environment

surrounding these tumor adjacent cells plus nucleic acids in the extra-cellular

environment coming from cancer nearby may be putting the tumor adjacent cells in an

antiviral state. It is known that interferon signaling induces proteins that act against

viruses. *ZAP* is one example that degrades viral RNA as well as RNA of LINEs and Alus

[58], although *ZAP* does not show correlated expression with the TE modules in our

data. We hypothesize that such cell states may reduce transposon transcripts with higher sensitivity through RNA degradation and chromatin remodeling.

The tissue samples in this study are not representative of "normal" cells, as they are collected as controls from tissue adjacent to cancer cells. Although they are not undergoing the molecular changes associated with malignant transformation, they could be under the influence of nearby environment, with changes in pH levels, inflammation, and infiltration of immune cells. The inclusion criteria for TCGA does not allow patients with any prior systemic chemotherapy or any other neoadjuvant therapy, but it does allow local radiation, and we observe that past local radiation is associated with higher TE expression levels in adjacent cells in thyroid tissues. Given the characteristics of the samples, the variation in TE expression levels or the co-expression pattern we observe in this study may be due to cancer-associated stress. Future studies will be needed to confirm whether the results are replicated in true normal tissue.

TEs and KRAB-ZFPs

ChIP-Seq studies on KRAB-ZFPs have identified extensive binding between this family of proteins and transposable elements [44,59], implying a role for suppressing TE expression. KRAB domain is a well-known repressor domain and together with the co-factor KAP1 (TRIM28), the KZFP-KAP1 complex has been shown to silence both exogenous retroviruses and endogenous retroelements during embryonic development [60,61]. Based on this observation, and the pattern of co-evolution of retroviral LTRs and the C2H2-Zinc Finger gene family, it has been hypothesized that the KRAB-ZFPs function in transposable element suppression [62]. But except for a few KRAB-ZFPs, most members do not have a characterized function. In an alternative hypothesis, instead of its original role in silencing, it was proposed that KRAB-ZFPs may also have a

role in controlling domesticated transposable elements that contribute to the host transcription regulation network [63]. In our co-expression analysis, we found overwhelming positive correlation between KZFPs and TEs across all classes of TEs. This positive correlation was observed whether we are counting multi-mapped reads or uniquely mapped reads, and whether we are counting TEs close to genes, or TEs in the intergenic regions. Despite this robust positive correlation, we found that the co-expressed relationship showed limited correspondence with published ChIP-seq binding results. There was statistically meaningful but very small number of overlap at the family level, and almost no overlap at the locus level. The co-expression we observe seems to be largely an indirect relationship, and not a result of direct binding. There have been observations of a unique chromatin state that is shared between ZFP clusters and repeat classes by the Roadmap Epigenomics project [64]. This chromatin state, termed ZNF/Rpts, is characterized by H3K36me3 marks co-occuring with H3K9me3 marks and high DNA methylation. It is possible that local chromatin environment that is co-regulated at a larger scale is responsible for the correlation at the RNA level.

Conclusions

TE derived transcripts in the non-tumourous tissues show large variation across tissues, and across individuals. Co-expression network analysis within tissues revealed general co-expression of TEs across all classes. It also found strong co-expression between TEs and KRAB-Zinc Finger Proteins that are replicated in multiple tissues, but not congruent with direct binding of TE-ZFP relationships assayed through ChIP-seq. We also found negative correlation between intronic TEs and mitochondrial genes, and between intergenic TEs and immune response genes, replicated in multiple tissues.

1   Methods

2   RNA-Seq and gene expression quantification in the non-tumorous tissues.

We used the gene level quantification provided by The Cancer Genome Atlas (TCGA) for

the gene expressions [24–26]. We collected gene level quantifications for 697 samples

from TCGA. We focused on cancer types that had at least 10 control samples of RNA-seq

data, collected from non-tumorous tissue adjacent to the cancer tissue. As a result, 16

different tissue types were included in our analysis: BLCA (Bladder urothelial

carcinoma) , BRCA (Breast carcinoma), COAD (Colon adenocarcinoma), ESCA

(Esophageal adenocarcinoma), HNSC (Head and neck squamous cell carcinoma), KICH

(kidney chromophobe), KIRC (kidney renal clear cell carcinoma), KIRP (Kidney renal

papillary cell carcinoma), LIHC (Liver hepatocellular carcinoma), LUAD (Lung

adenocarcinoma), LUSC (Lung squamous cell carcinoma), PRAD (Prostate

adenocarcinoma), READ (Rectum adenocarcinoma), STAD (Stomach adenocarcinoma),

THCA (Thyroid carcinoma) and UCEC (Uterine Corpus Endometrial Carcinoma).

Number of samples for each tissue is described in Supplementary Table 1. Although we

will use the acronym for the cancer type to describe these tissues, we emphasize again

that all our samples come from the non-tumorous tissues collected from the same organ

of the same patient with the cancer. The cancer tissue samples were not included in our

analysis.

Methods for sequencing and data processing of RNA using the RNA-seq protocol for all

tissues except esophagus and stomach have been previously described for TCGA in [24–

26]. Briefly, RNA was extracted, prepared into poly(A) enriched Illumina TruSeq mRNA

libraries, sequenced by Illumina HiSeq2000 (resulting in paired 48-nt reads), and

subjected to quality control. Sequencing for esophagus and stomach was done

differently from other tissues and have been described in [27]. Briefly, poly A+ mRNA

was purified using MultiMACS mRNA isolation kit on MultiMACS 96 separator, and

double stranded cDNA was synthesized using the Superscript Double-Stranded cDNA

synthesis kit. Following the library preparation protocol described in [27], the final DNA

was sequenced on Illumina HiSeq2000 with paired end 75-nt reads. RNA reads were

aligned to the hg19 genome assembly using Mapsplice [65]. Gene expression was

quantified for the transcript models corresponding to the TCGA GAF2.1 using RSEM

[31]. We used the raw_count values in the .rsem.genes.results files, rounded to an

integer, as the gene level quantification.

Quantifying TE derived transcripts at the locus and family level

We collected RNA-seq level 1 binary alignment files (.bam files) for 697 samples

(Supplementary Table 1) from TCGA. The bam files were then converted to fastq and

realigned to the hg19 reference genome using STAR and Bowtie1. With the STAR

alignment, we allowed up to 200 mappings for every read (--outFilterMultimapNmax

200 --winAnchorMultimapNmax 200). With the Bowtie1 alignment, we only allowed

the single best alignment for each read, and if there were multiple best alignments, the

read was discarded from the final alignment (-m 1 -S -y -v 3 -X 1000 --max). We used a

modified version of the software TEtranscripts [29] for quantifying the reads mapping

to annotated transposons. TEtranscripts is a software that can quantify both gene and

TE transcript levels from RNAseq experiments. It takes into account the ambiguously

mapped TE-associated reads by proportionally assigning read counts to the

corresponding TE families using an Expectation-Maximization algorithm. We

implemented two modification to the original TEtranscripts software. 1) We modified it

to report read counts for each individual TE locus in the reference genome in addition to

the family level counts. 2) We developed a function to discount the read counts by

removing read counts that correspond to transcripts containing TE sequences that

originate from pre-mRNA or retained introns in the mature RNA [28]. Downstream

analyses were done using the discounted quantification based on multi-mapped reads

and the uniquely mapped quantification for both the STAR alignment and the Bowtie1

alignment, to assess the impact of uncertainty in multi-mapped reads.

The retrotransposon annotations used were generated from the RepeatMasker tables,

obtained from the UCSC genome database and provided by TEtranscripts. For

quantifying reads mapping to the TE flanking introns we generated gtf files containing

1) the TE flanking intron positions, 2) the intergenic TE positions, and 3) the exonic TE

positions (TEs that fall within an exon, including non-coding RNA genes). In case of

intronic TEs, we use the algorithm described above to discount the transcripts from pre-

mRNA or retained introns. In case of intergenic TEs, we count all EM estimated reads

mapped to TEs without any discount. In case of exonic TEs, we ignore those counts

altogether, and the exonic TEs do not contribute to the locus count nor the family level

count.

Normalization and transformation of read counts

After quantifying the reads mapping to annotated genes and TEs, both the gene level

counts, and the TE counts were normalized between samples across all tissue types

with DEseq2. We used the default "median ratio method" for normalization in DESeq2

[66]. Briefly, the scaling factor for each sample is calculated as median of the ratio, for

each gene, of its read count to its geometric mean across all samples. The assumption of

the median ratio method is that most genes are not consistently differentially expressed

between tissues. If there is systematic difference in ratio between samples, the median

ratio will capture the size relationship. But, this assumption may be violated when we

are comparing large number of tissues types at the same time, since a large proportion

of the genes may be differentially expressed in at least one tissue type, or one of the

tissues may be extremely biased in their number of differentially expressed genes. In

order to achieve more robust normalization, we used a two-step normalization method

called the differentially expressed genes elimination strategy (DEGES) [67]. We

performed preliminary normalization using the "median ratio method", filtered out

potential differentially expressed genes in the data, found a subset of robust non-

differentially expressed genes, and used the subset to perform the second round of

"median ratio normalization". The resulting pairwise MA plot between tissues after

normalization showed better normalization compared to the regular one-step

normalization. The size factors for each sample obtained from the two-step

normalization on gene counts were then used to normalize the TE quantifications of the

same sample. The normalized counts were log2 transformed using the variance

stabilizing transformation function in DESeq2 [66,68] for downstream analysis.

Clustering of samples by expression pattern.

We cluster the samples using the "average" method (=UPGMA) in the *hclust* function of

R, and visualize the clusters with the ComplexHeatmap package [69]. The top 150 genes

or TEs, with the largest variances on the log2 transformed read counts were used for

clustering. We did not select genes by any measure of differential expression across

tissues. These genes were simply the genes showing the largest variance in read count

across all 697 samples, regardless of tissue type. We exclude genes and TEs on X and Y

chromosomes. Based on the log2 read count of the top 150 TEs, a dissimilarity matrix is

calculated and used for the clustering and visualization. The average method of *hclust*

computes all pairwise dissimilarities between the members of the two clusters and

considers the average as the distance between the two clusters. Hierarchical clustering

starts with each sample assigned to its own cluster and then proceeds iteratively, at

each stage joining the two most similar clusters, continuing until there is just a single

cluster. For the locus level TE expression, we filtered out all loci that had less than 5

read counts for every sample. To compare the clustering of samples based on gene

expression, TE expression and random assignment, we used the normalized Mutual

Information (NMI) measure [70]. The hierarchical clusters were cut off at k = 16, the

number of different tissue types. Because the resulting clusters were not accurate

enough to distinguish between similar tissues, we used a broader tissue grouping to

compare with the clusters. The tissues were grouped to 10 broader types based on

preliminary clustering: bladder/endometrium (BLCA, UCEC), breast (BRCA), liver

(LIHC), colon/rectum (COAD, READ), esophagus/stomach (ESCA, STAD), head and neck

— the squamous epithelium in the mucosal surfaces inside the mouth, nose, and throat

(HNSC), kidney (KICH, KIRC, KIRP), lung (LUAD, LUSC), prostate (PRAD), and thyroid

(THCA). The broader tissue type of each sample was used as the ground truth. Each

resulting cluster was then assigned a group label based on the majority tissue type.

Normalized mutual information was calculated by comparing the labels from the

clustering to the true class labels. Random assignment clusters were generated by

permuting the tissue types with and without replacement 100 times, and the mean NMI

was reported.

Co-expression network analysis with TEs and host genes

Weighted correlation network analysis was done with the WGCNA package [34]. We

start with the signed pair-wise correlation matrix across the expression levels

(normalized $\log_2$ read counts) of all genes and TE families. We calculate the adjacency

matrix by raising the correlation matrix to the power of 14, power parameter selected

using the scale free topology measure, effectively suppressing the low correlations due

to noise. Topological overlap based distance matrix (TOM) is calculated using the

network topology resulting from the adjacency matrix. This procedure was repeated for

each tissue, and a consensus TOM was calculated across all tissues. We used hierarchical

clustering on this consensus topological overlap matrix to identify clusters (modules)

that are shared across tissues. A representative gene expression profile of the module is

defined by the first principal component of the expression levels of all members in each

module. The representative profile is compared between each module to identify

positive and negative correlation between modules.

Correlated expression between genes and L1HS 5'

We blasted all the L1HS instances annotated in repeatmasker against the L1HS

consensus sequence and identified the regions aligning to the 300 bases of the 5' end of

the consensus sequence. We counted all the reads mapping to the list of L1HS 5' ends

and normalized them with the same size factor described above. We used log2

transformed value of this normalized read count as the variable representing L1HS

transcript level. Correlation between gene and L1HS 5' transcripts were tested in each

tissue groups separately, in bladder, breast, liver, colon/rectum, stomach/esophagus,

head and neck, kidney, lung, prostate and thyroid. We tested 20532 genes for each

tissue group using a linear model with log2 L1HS 5' expression as the dependent

variable, and log2 gene expression as the independent variable. For a gene to be

included in our test, it had to be present in at least eight individual patients. We also

required that the gene be expressed with a minimum RPM of 2 in 75% of the samples to

be included in the dataset. In addition to the radiation therapy for thyroid tissue, we

considered effective library size (sum of all normalized counts) and the batch ID

provided by the TCGA project as additional covariates. Since there was significant co-

expression across all TE classes especially for the intronic TEs, we included the

expression profile of the intronic TE module N1 identified during the co-expression

network analysis as a covariate in our linear model. The linear model we used is

described below.

$$\log_2 L1HS \ \sim \log_2 gene \ + \ \log_2 effective\_library\_size \ + \ batch \ + \ radiation \\ + \log_2 N1 \, profile$$

We tested all combination of linear models that can be created by including or excluding

these variables. Second-order Akaike Information Criterion (AIC$_c$) was used to select the

best linear model. We used the coefficient and p-value from the best model to calculate

the q-values. Genes with q-value < 0.0001 in at least two tissues were identified as

correlated genes.

Correlation between TE and KRAB-ZFPs

To understand the positive correlation between TEs and KRAB-ZFPs, we looked at the

correlation between each KZFPs and TEs at the family level and at the individual TE

locus level in different tissue types. We tested the correlation for 366 KRAB Zinc Finger

Proteins that were identified in Imbeault et al. [44] and also found in our gene

expression data. Because the search space of pairwise combinations of KZFP and

individual TE loci was too large, we examined the relationship in a step-wise approach.

In the first step, we tested the correlation between all pairwise combinations of 366

KZFPs and 979 TE subfamilies using the TE quantification at the family level in each

tissue type. Then, in the second step, once the significantly correlated KZFP and TE

family was identified, we focused on those pairs. We tested the correlation between the

expression of the significant KZFP and the expression of each individual locus of the

significant TE family in the tissue where the initial co-expression was found to identify

individual TE loci that are co-expressed with the KZFP.

Overlap between co-expression and binding was examined at the family level and at the

locus level. At the family level, we downloaded the family enrichment results from

Imbeault et al. [44] and identified pairs of TE families and KZFP that had an enrichment score greater than 1. We compared those families enriched with binding of specific KZFPs to our co-expression results, to check if the TE families were co-expressed with the same KZFPs. At the locus level, we compared the co-expressed TE loci with the binding peaks reported in the dataset GSE78099. We took +- 250 bp around the boundary of peaks and found overlap with TE annotations from Repeatmasker. We checked if the TE locus overlapping with ChIP-seq peaks were found to be co-expressed with any KZFPs.

List of Abbreviations

TE: Transposable Element

KZFP: KRAB Zinc Finger Protein

TCGA: The Cancer Genome Atlas

Declaration

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable

Availability of data and material

The datasets generated and/or analysed during the current study are available in the github repository, *https://github.com/HanLabUNLV/TEcoex*. The modified version of the TEtranscripts software [29] and the required gtf files can be found at *https://github.com/HanLabUNLV/tetoolkit*.

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by the National Institutes of Health [R15GM116108, P20GM121325 to M.V.H.], and by the National Science Foundation [1750532 to M.V.H].

Authors' contributions

NC performed the co-expression analysis. GMJ performed TEtranscripts quantification. SQ performed the REC score analysis. NC, AR analyzed the KZFP-TE locus pairwise co-expression and KZFP motif search. CS re-ran the pipeline with the Bowtie and STAR alignment program. AA, CC, DC, ON assisted with the analyses. MVH designed the experiments, modified the TEtranscripts software, analyzed and interpreted the data. MVH wrote the manuscript with the help of all other authors. All authors read and approved the final manuscript.

Acknowledgements

We thank the TCGA and GTEx project teams for making the data available.

References

1. Slotkin RK. The case for not masking away repetitive DNA. Mob DNA. 2018;9:15.

2. Branciforte D, Martin SL. Developmental and cell type specificity of LINE-1 expression in mouse testis: implications for transposition. Mol Cell Biol. 1994;14:2584–92.

3. Trelogan SA, Martin SL. Tightly regulated, developmentally specific expression of the first open reading frame from LINE-1 during mouse embryogenesis. Proc Natl Acad Sci. 1995;92:1520–4.

4. Ergün S, Buschmann C, Heukeshoven J, Dammann K, Schnieders F, Lauke H, et al. Cell Type-specific Expression of LINE-1 Open Reading Frames 1 and 2 in Fetal and Adult Human Tissues. J Biol Chem. 2004;279:27753–63.

5. Kubo S, Seleme M del C, Soifer HS, Perez JLG, Moran JV, Kazazian HH, et al. L1 retrotransposition in nondividing and primary human somatic cells. Proc Natl Acad Sci. 2006;103:8036–41.

6. Belancio VP, Roy-Engel AM, Pochampally RR, Deininger P. Somatic expression of
LINE-1 elements in human tissues. Nucleic Acids Res. 2010;38:3909–22.
7. Rangwala SH, Zhang L, Kazazian HH. Many LINE1 elements contribute to the
transcriptome of human somatic cells. Genome Biol. 2009;10:R100.
8. Skowronski J, Singer MF. Expression of a cytoplasmic LINE-1 transcript is regulated in
a human teratocarcinoma cell line. Proc Natl Acad Sci U S A. 1985;82:6050–4.
9. Bratthauer GL, Cardiff RD, Fanning TG. Expression of LINE-1 retrotransposons in
human breast cancer. Cancer. 1994;73:2333–6.
10. Rodić N, Sharma R, Sharma R, Zampella J, Dai L, Taylor MS, et al. Long Interspersed
Element-1 Protein Expression Is a Hallmark of Many Human Cancers. Am J Pathol.
2014;184:1280–6.
11. Bratthauer GL, Fanning TG. Active LINE-1 retrotransposons in human testicular
cancer. Oncogene. 1992;7:507–10.
12. Philippe C, Vargas-Landin DB, Doucet AJ, Essen D van, Vera-Otarola J, Kuciak M, et al.
Activation of individual L1 retrotransposon instances is restricted to cell-type
dependent permissive loci. eLife. 2016;5:e13926.
13. Muotri AR, Chu VT, Marchetto MCN, Deng W, Moran JV, Gage FH. Somatic mosaicism
in neuronal precursor cells mediated by L1 retrotransposition. Nature. 2005;435:903.
14. Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, et al. The regulated
retrotransposon transcriptome of mammalian cells. Nat Genet. 2009;41:563–71.
15. Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, et al. Landscape of
transcription in human cells. Nature. 2012;489:101–8.
16. Chuong EB, Elde NC, Feschotte C. Regulatory activities of transposable elements:
from conflicts to benefits. Nat Rev Genet. 2016;18:71.
17. Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, et al. Widespread contribution of
transposable elements to the innovation of gene regulatory networks. Genome Res.
2014;24:1963–76.
18. Kapusta A, Kronenberg Z, Lynch VJ, Zhuo X, Ramsay L, Bourque G, et al.
Transposable Elements Are Major Contributors to the Origin, Diversification, and
Regulation of Vertebrate Long Noncoding RNAs. PLOS Genet. 2013;9:e1003470.
19. Jachowicz JW, Bing X, Pontabry J, Bošković A, Rando OJ, Torres-Padilla M-E. LINE-1
activation after fertilization regulates global chromatin accessibility in the early mouse
embryo. Nat Genet. 2017;49:1502.
20. Percharde M, Lin C-J, Yin Y, Guan J, Peixoto GA, Bulut-Karslioglu A, et al. A LINE1-
Nucleolin Partnership Regulates Early Development and ESC Identity. Cell.
2018;174:391-405.e19.
21. Liu N, Lee CH, Swigut T, Grow E, Gu B, Bassik MC, et al. Selective silencing of
euchromatic L1s revealed by genome-wide screens for L1 regulators. Nature.
2017;553:228.
22. Taylor MS, Altukhov I, Molloy KR, Mita P, Jiang H, Adney EM, et al. Dissection of
affinity captured LINE-1 macromolecular complexes. eLife. 2018;7:e30094.
23. Mita P, Wudzinska A, Sun X, Andrade J, Nayak S, Kahler DJ, et al. LINE-1 protein
localization and functional dynamics during the cell cycle. eLife. 2018;7:e30058.
24. The Cancer Genome Atlas Network. Comprehensive molecular characterization of
human colon and rectal cancer. Nature. 2012;487:330.
25. The Cancer Genome Atlas Network. Comprehensive molecular portraits of human
breast tumours. Nature. 2012;490:61.
26. The Cancer Genome Atlas Research Network. Comprehensive genomic
characterization of squamous cell lung cancers. Nature. 2012;489:519.

27. The Cancer Genome Atlas Research Network, Bass AJ, Thorsson V, Shmulevich I, Reynolds SM, Miller M, et al. Comprehensive molecular characterization of gastric adenocarcinoma. Nature. 2014;513:202.

28. Deininger P, Morales ME, White TB, Baddoo M, Hedges DJ, Servant G, et al. A comprehensive approach to expression of L1 loci. Nucleic Acids Res. 2017;45:e31–e31.

29. Jin Y, Tam OH, Paniagua E, Hammell M. TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. Bioinforma Oxf Engl. 2015;31:3593–9.

30. Britten RJ. Mobile elements inserted in the distant past have taken on important functions. Gene. 1997;205:177–82.

31. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics. 2011;12:323.

32. Yang WR, Ardeljan D, Pacyna CN, Payer LM, Burns KH. SQuIRE reveals locus-specific regulation of interspersed repeat expression. Nucleic Acids Res. 2019;47:e27–e27.

33. Doucet-O'Hare TT, Rodić N, Sharma R, Darbari I, Abril G, Choi JA, et al. LINE-1 expression and retrotransposition in Barrett's esophagus and esophageal carcinoma. Proc Natl Acad Sci. 2015;112:E4894.

34. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC Bioinformatics. 2008;9:559.

35. Desai N, Sajed D, Arora KS, Solovyov A, Rajurkar M, Bledsoe JR, et al. Diverse repetitive element RNA expression defines epigenetic and immunologic features of colon cancer. JCI Insight. 2017;2:e91078.

36. Solovyov A, Vabret N, Arora KS, Snyder A, Funt SA, Bajorin DF, et al. Global Cancer Transcriptome Quantifies Repeat Element Polarization between Immunotherapy Responsive and T Cell Suppressive Classes. Cell Rep. 2018;23:512–21.

37. Menendez L, Benigno BB, McDonald JF. L1 and HERV-W retrotransposons are hypomethylated in human ovarian carcinomas. Mol Cancer. 2004;3:12.

38. Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res. 2016;44:W90–7.

39. Choi J, Hwang S-Y, Ahn K. Interplay between RNASEH2 and MOV10 controls LINE-1 retrotransposition. Nucleic Acids Res. 2018;46:1912–26.

40. de la Rica L, Deniz Ö, Cheng KCL, Todd CD, Cruz C, Houseley J, et al. TET-dependent regulation of retrotransposable elements in mouse embryonic stem cells. Genome Biol. 2016;17:234.

41. Macfarlan T, Kutney S, Altman B, Montross R, Yu J, Chakravarti D. Human THAP7 Is a Chromatin-associated, Histone Tail-binding Protein That Represses Transcription via Recruitment of HDAC3 and Nuclear Hormone Receptor Corepressor. J Biol Chem. 2005;280:7346–58.

42. Koizumi S, Irie T, Hirayama S, Sakurai Y, Yashiroda H, Naguro I, et al. The aspartyl protease DDI2 activates Nrf1 to compensate for proteasome dysfunction. Dikic I, editor. eLife. 2016;5:e18357.

43. Sun X, Wang X, Tang Z, Grivainis M, Kahler D, Yun C, et al. Transcription factor profiling reveals molecular choreography and key regulators of human retrotransposon expression. Proc Natl Acad Sci. 2018;115:E5526.

44. Imbeault M, Helleboid P-Y, Trono D. KRAB zinc-finger proteins contribute to the evolution of gene regulatory networks. Nature. 2017;543:550–4.

45. The ENCODE Project Consortium, Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis
CA, et al. An integrated encyclopedia of DNA elements in the human genome. Nature.
2012;489:57.
46. Scott EC, Gardner EJ, Masood A, Chuang NT, Vertino PM, Devine SE. A hot L1
retrotransposon evades somatic repression and initiates human colorectal cancer.
Genome Res. 2016;26:745–55.
47. Okudaira N, Iijima K, Koyama T, Minemoto Y, Kano S, Mimori A, et al. Induction of
long interspersed nucleotide element-1 (L1) retrotransposition by 6-formylindolo[3,2-
b]carbazole (FICZ), a tryptophan photoproduct. Proc Natl Acad Sci. 2010;107:18487–
18492.
48. Stribinskis V, Ramos KS. Activation of Human Long Interspersed Nuclear Element 1
Retrotransposition by Benzo(a)pyrene, an Ubiquitous Environmental Carcinogen.
Cancer Res. 2006;66:2616–2620.
49. Terasaki N, Goodier JL, Cheung LE, Wang YJ, Kajikawa M, Kazazian HH Jr, et al. In
Vitro Screening for Compounds That Enhance Human L1 Mobilization. PLOS ONE.
2013;8:e74629.
50. Banaz-Yaşar F, Gedik N, Karahan S, Diaz-Carballo D, Bongartz BM, Ergün S. LINE-1
Retrotransposition Events Regulate Gene Expression After X-Ray Irradiation. DNA Cell
Biol. 2012;31:1458–67.
51. Farkash EA, Kao GD, Horman SR, Prak ETL. Gamma radiation increases
endonuclease-dependent L1 retrotransposition in a cultured cell assay. Nucleic Acids
Res. 2006;34:1196–204.
52. Giorgi G, Marcantonio P, Del Re B. LINE-1 retrotransposition in human
neuroblastoma cells is affected by oxidative stress. Cell Tissue Res. 2011;346:383–91.
53. Van Meter M, Kashyap M, Rezazadeh S, Geneva AJ, Morello TD, Seluanov A, et al.
SIRT6 represses LINE1 retrotransposons by ribosylating KAP1 but this repression fails
with stress and age. Nat Commun. 2014;5:5011.
54. Jung H, Choi JK, Lee EA. Immune signatures correlate with L1 retrotransposition in
gastrointestinal cancers. Genome Res [Internet]. 2018; Available from:
http://genome.cshlp.org/content/early/2018/07/03/gr.231837.117.abstract
55. Chiappinelli KB, Strissel PL, Desrichard A, Li H, Henke C, Akman B, et al. Inhibiting
DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including
Endogenous Retroviruses. Cell. 2015;162:974–86.
56. Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, et al. DNA-
Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by
Endogenous Transcripts. Cell. 2015;162:961–73.
57. Haffner MC, Taheri D, Luidy-Imada E, Palsgrove DN, Eich M-L, Netto GJ, et al.
Hypomethylation, endogenous retrovirus expression, and interferon signaling in
testicular germ cell tumors. Proc Natl Acad Sci. 2018;115:E8580.
58. Moldovan JB, Moran JV. The Zinc-Finger Antiviral Protein ZAP Inhibits LINE and Alu
Retrotransposition. PLOS Genet. 2015;11:e1005121.
59. Najafabadi HS, Mnaimneh S, Schmitges FW, Garton M, Lam KN, Yang A, et al. C2H2
zinc finger proteins greatly expand the human regulatory lexicon. Nat Biotechnol.
2015;33:555–62.
60. Wolf D, Goff SP. Embryonic stem cells use ZFP809 to silence retroviral DNAs. Nature.
2009;458:1201–4.
61. Jacobs FMJ, Greenberg D, Nguyen N, Haeussler M, Ewing AD, Katzman S, et al. An
evolutionary arms race between KRAB zinc-finger genes *ZNF91/93* and SVA/L1
retrotransposons. Nature. 2014;516:242–5.

62. Rowe HM, Trono D. Dynamic control of endogenous retroviruses during development. Virology. 2011;411:273–87.

63. Trono D. Transposable Elements, Polydactyl Proteins, and the Genesis of Human-Specific Transcription Networks. Cold Spring Harb Symp Quant Biol. 2015;80:281–8.

64. Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, et al. Integrative analysis of 111 reference human epigenomes. Nature. 2015;518:317.

65. Wang K, Singh D, Zeng Z, Coleman SJ, Huang Y, Savich GL, et al. MapSplice: Accurate mapping of RNA-seq reads for splice junction discovery. Nucleic Acids Res. 2010;38:e178–e178.

66. Anders S, Huber W. Differential expression analysis for sequence count data. Genome Biol. 2010;11:R106.

67. Kadota K, Nishiyama T, Shimizu K. A normalization strategy for comparing tag count data. Algorithms Mol Biol. 2012;7:5.

68. Huber W, von HA, Sueltmann H, Poustka A, Vingron M. Parameter estimation for the calibration and variance stabilization of microarray data. Stat Appl Genet Mol Biol. 2003;2:Article3.

69. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics. 2016;32:2847–9.

70. Cover TM, Thomas JA. Elements of Information Theory. New York: Wiley & Sons; 1991.

71. Caracausi M, Piovesan A, Antonaros F, Strippoli P, Vitale L, Pelleri MC. Systematic identification of human housekeeping genes possibly useful as references in gene expression studies. Mol Med Rep. 2017;16:2397–410.

72. Eisenberg E, Levanon EY. Human housekeeping genes, revisited. Trends Genet TIG. 2013;29:569–74.

Figures

Figure 1. Comparison of read alignment on full length L1HS with multi-mapping and unique mapping.

Reads mapped to two different full length L1HS elements from a stomach tissue sample

(A4GY) visualized through IGV. a. L1HS_dup967, a 5799nt length element on

chromosome X: 75453754-75459553. b. a 6225nt length element L1HS_dup924 on

chromosome X: 11953208- 11959433. Chromosomal locations, 48 base mappability

calculated with GEM, Bowtie1 alignment allowing only uniquely mapped reads with

single best hit, STAR alignment allowing multi-mapped reads up to 200 mapping for

each read, gene annotation and Repeatmasker TE annotation are shown from top to

bottom. Red lines mark the boundary of the L1HS elements with 5' and 3' noted.

Figure 2. Relationship between read counts for each L1HS locus, and the mappability of the locus.

Log_2 transformed total read counts mapped to each L1HS locus in the hg19 genome,

summed across all samples in our dataset are plotted against the total uniquely

mappable positions (number of positions with mappability score = 1 based on 48bp

mappability calculated with GEM) for each locus. L1HS loci with zero read counts are

marked at -1 instead of –infinity. a. read counts from Bowtie1 alignment with uniquely

mapped reads only b. read counts from STAR alignment allowing multi-mapped reads

up to 200. c. comparison of read counts for each L1HS locus between the uniquely

mapped reads and multi-mapped reads.

Figure 3. Tissue clustering based on transposable elements

Heatmap showing the tissue clustering results based on the top 150 TEs with the largest variance of each class. With the color bars at the top of the heatmap, the upper color labels show tissue types and the lower color labels show broader tissue groupings. a-d. clustering based on family level quantification. e-h. clustering based on locus level quantification. i. Clustering quality measured by Mutual Information for clustering results based on family level TE quantification, locus level quantification for TEs 100Kb away from genes, locus level quantification for TEs 1Kb away from genes, and locus level TE quantification without filtering for gene proximity.

Figure 4. representative transposable element loci that show tissue-specific expression

15 TE loci with tissue-specific expression was identified as representative examples among intergenic TEs that are 100Kb away from start and end of genes. Heatmap color reflects the z-score of normalized $\log_2$ read counts across samples.

Figure 5. Variation in TE expression across tissues and individuals

a. Normalized and log-transformed sum of all read counts mapping to the TEs of each class, LINE, DNA, SINE and LTR are shown as violin plots. Mean read count of a set of housekeeping genes (Supplementary Table 6) are plotted as a reference. Horizontal line across the violin plot represents the median value across all samples of that tissue type. b. Normalized and log-transformed sum of all read counts mapping to 300 bp at the 5' end of L1HS are plotted as a violin plot. Same set of housekeeping genes used in a. are plotted as a reference.

Figure 6. co-expression modules with immune genes are negatively correlated with TE modules.

a. enriched annotations for genes belonging to module M33 identified in the Reactome database. b. enriched annotations for genes belonging to module M35 identified in the Reactome database. c. top 30 genes with highest module membership (high correlation with representative profile of the module) for each module. d-e. correlation plot showing high within group correlation and negative between group correlation between the TEs in the TE modules and the immune genes in modules M33 and M35. d. data from breast, and e. data from esophagus. Color label on top of the correlation plot show different classes of TEs, and genes that are annotated with the GO term "immune system process".

Figure 7. co-expression modules with mitochondrial genes and ribosome genes are negatively correlated with intronic TEs.

a. enriched annotations for genes belonging to module N4 identified in the Reactome database. b-c. correlation plot showing high within group correlation and negative between group correlation between the TEs in the intronic TE module N1, and the mitochondrial and ribosomal genes in modules N4. b. shows data from breast, and c. shows data from esophagus. Color label on top of the correlation plot show different classes of TEs, and genes that are annotated with the GO term "mitochondrion" and "ribosome".

Figure 8. Genes that show positive and negative correlation with the transcript level of L1HS 5' end.

Gene that show significant positive and negative correlation with L1HS 5' in multiple tissues. Esophagus and stomach are combined as one tissue group.

Figure 9. Overlap between co-expressed KZFP-TE family pairs and TE families enriched for KZFP binding.

a. KZFP-TE families co-expressed in at least two tissues are marked with pink, TE

families enriched for KZFP binding are marked with yellow, and TE families that are

both bound by KZFP and co-expressed with same KZFP are marked with green. KZFPs

that show overlap of binding and co-expression for multiple TE families are labeled

along the vertical axis. b-c. categorization of co-expression and KZFP binding for all

200,889 KZFP-TE family pair-wise combinations (221 KZFP x 909 TE families), that

have both expression and ChIP-exo data. b. counts co-expression significant in at least

one tissue. c. counts co-expression significant in at least two tissues.

Figure 10. Overlap between expression and KZFP binding for TE loci

a. categorization of all 4,496,028 TE loci annotated in hg19 RepeatMasker by expression

and KZFP binding. b. Overlap of expression and KZFP binding with ENCODE Candidate

Regulatory Element marks. c. Proportion of each category of TEs that are marked with

ENCODE Candidate Regulatory Element marks.

Tables

Table 1. Transposon loci that show large difference after correcting for pre-mRNA/retained introns.

a. Transposon loci embedded within introns or exons of genes that frequently result in the largest correction in each sample. Locus id, genomic location, surrounding gene and structure the TE is embedded in, and the maximum number of reads removed in a sample.

locus	chr	start	end	surrounding gene	TE embedded in	# of samples	Max correction
MIRc_dup47590	8	22021288	22021431	*SFTPC*	Intron 4, Exon 5	105	428021
AluY_dup80589	12	69747275	69747567	*LYZ*	Exon 4	62	313317
MIRb_dup137684	10	81315669	81315913	*SFTPA2*	Exon 5	106	266566
MIRb_dup137689	10	81374907	81375150	*SFTPA1*	Exon 5	106	230394
MIR3_dup57107	10	81316603	81316678	*SFTPA2*	Exon 5	103	90241
AluSz6_dup3320	1	207102295	207102608	*PIGR*	Exon 11	124	59130
MIRc_dup74805	12	50351953	50352157	*AQP2*	Exon 4	109	57581
LTR39_dup404	6	160102172	160102969	*SOD2*	Exon 4, Intron 7	119	34545
AluSx1_dup59209	Y	21153222	21153521	*TTTY14*	Exon 1	305	25867
AluJb_dup119100	17	16344881	16345132	*C17orf76-AS1*	Intron 4, Exon 5	253	21915

Table 2. KZFP gene members in core TE modules

KZFP genes that are members of the core TE modules, and module M3 that is correlated

with core TE modules.

core TE modules	KZFP	chromosome	KZFPs in correlated module M3	chromosome
	HKR1	19	KZFP169	9
	KZFP226	19	KZFP202	11
	KZFP682	19	KZFP266	19
	KZFP789	7	KZFP300	5
M8	KZFP814	19	KZFP320	19
	KZFP404	19	KZFP431	19
	KZFP418	19	KZFP439	19
	KZFP589	3	KZFP44	19
M21	KZFP75A	19	KZFP587	19
M38	KZFP117	7	KZFP662	3
	KZFP334	20	KZFP7	8
	KZFP493	19	KZFP700	19
	KZFP506	19	KZFP708	19
	KZFP721	4	KZFP714	19
M45	KZFP737	19	KZFP732	4
			KZFP83	19
			KZFP841	19

Table 3. Gene members in the intronic TE module N1.

Genes that are members of the intronic TE module N1, the number of tissues they are

assigned to module N1 in, synonyms to gene names, full gene names.

Gene Symbol	tissues	Synonyms	Full gene name
NCRNA00201	7	HNRNPU	heterogeneous nuclear ribonucleoprotein U
AHSA2	6	AHSA2P	activator of HSP90 ATPase homolog 2, pseudogene
CCNL2	6	CCNL2	cyclin L2
CG030	5	N4BP2L2-IT2	N4BPL2 intronic transcript 2
FAM13AOS	5	FAM13A-AS1	FAM13A antisense RNA 1
MDM4	5	MDM4	MDM4 regulator of p53
NKTR	5	NKTR	natural killer cell triggering receptor
SLC25A27	5	UCP4	solute carrier family 25 member 27
ANKRD36	4	ANKRD36	ankyrin repeat domain 36
LOC100190986	4	LOC100190986	uncharacterized LOC100190986
LOC440944	4	THUMPD3-AS1, SETD5-AS1	THUMPD3 antisense RNA 1
LOC91316	4	GUSBP11	GUSB pseudogene 11
LUC7L	4	LUC7L	LUC7 like
LUC7L3	4	LUC7L3	LUC7 like 3 pre-mRNA splicing factor
NCRNA00105	4	ASMTL-AS1	ASMTL antisense RNA 1
OGT	4	OGT	O-linked N-acetylglucosamine (GlcNAc) transferase
SEC31B	4	SEC31B	SEC31 homolog B, COPII coat complex component
KZFP789	4	KZFP789	zinc finger protein 789

Figure 1

Figure 2

read counts summed across samples for each L1HS locus

a. uniquely mapped reads

coef: 0.001752
p-val: 5.523e-07

b. multi-mapped reads

coef: 0.0001353
p-val: 0.6038

c.

Figure 3

i. Tissue clustering Mutual Information

Figure 4

HERVIP10F-int_dup69:HERVIP10F-int:ERV1:LTR:-   Chr3:5733265-5733969

L4_dup13748:L4:RTE:LINE:-   Chr14:25735703-25735768

ERV3-16A3_I-int_dup4366:ERV3-16A3_I-int:ERVL:LTR:-   Chr11:119797856-119801574

L1PA11_dup956:L1PA11:L1:LINE:+   Chr4:27721476-27721737

HERVE-int_dup79:HERVE-int:ERV1:LTR:-   Chr7:6990536-6998376

AluSz_dup79518:AluSz:Alu:SINE:-   Chr16:76016076-76016361

L1PA3_dup778:L1PA3:L1:LINE:-   Chr2:13612138-13613422

SST1_dup58:SST1:centr:Satellite:-   Chr9:42251223-42252481

MLT1D_dup442:MLT1D:ERVLMaLR:LTR:+   Chr1:67997610-67998059

L1PA3_dup778:L1PA3:L1:LINE:-   Chr2:13612138-13613422

LTR7B_dup240:LTR7B:ERV1:LTR:+   Chr4:80617583-80618104

LTR12C_dup844:LTR12C:ERV1:LTR:+   Chr4:187811087-187812497

REP522_dup155:REP522:telo:Satellite:-   Chr16:33806736-33807068

LTR30_dup97:LTR30:ERV1:LTR:-   Chr10:54784934-54785648

L1MA3_dup4399:L1MA3:L1:LINE:+   Chr8:31240505-31244899

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

a.

b.

	ZNF bound	ZNF unbound	total
co-expressed	540	25326	25866
not-coexpressed	1870	173153	175023
total	2410	198479	200889

c.

	ZNF bound	ZNF unbound	total
co-expressed	119	4019	4138
not-coexpressed	2291	194460	196751
total	2410	198479	200889

Figure 10

a.

	bound	unbound	total
expressed	117154	585717	702871
nonexpressed	362642	3430515	3793157
total	479796	4016232	4496028

b.

ENCODE	RNA-seq	ZFP ChIP	overlap cnt	total cnt
		bound	16370	117154
	expressed	unboud	33505	585717
		bound	28032	362642
DNase+H3K4me3	nonexpressed	unboud	91819	3430515
		bound	31205	117154
	expressed	unboud	139049	585717
		bound	79953	362642
DNase+H3K27ac	nonexpressed	unboud	572925	3430515
		bound	2137	117154
	expressed	unboud	6924	585717
		bound	6179	362642
DNase+CTCFonly	nonexpressed	unboud	39254	3430515

c.

proportion of TEs overlapping with ENCODE marks

■ H3K4me3   ■ H3K27ac   ■ CTCFonly   ■ no mark