***In vivo* characterization of candidate genes for heart rate variability identifies**
**culprits for sinoatrial pauses and arrests**

Benedikt von der Heyde[1,2], Anastasia Emmanouilidou[1,2], Tiffany Klingström[1,2],
Eugenia Mazzaferro[1,2], Silvia Vicenzi[1,2], Sitaf Jumaa[1,2], Olga Dethlefsen[3,4], Harold
Snieder[5], Eco de Geus[6], Erik Ingelsson[2,7,8,9], Amin Allalou[2,10], Hannah L. Brooke[11],
Marcel den Hoed[#1,2]

[1]Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala,
Sweden.
[2]Science for Life Laboratory, Uppsala University, Uppsala, Sweden.
[3]Science for Life Laboratory, Stockholm University, Stockholm, Sweden.
[4]National Bioinformatics Infrastructure Sweden, Stockholm University, Stockholm,
Sweden.
[5]Department of Epidemiology, University of Groningen, University Medical Center
Groningen, Groningen, the Netherlands.
[6]Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam,
the Netherlands.
[7]Department of Medical Sciences, Molecular Epidemiology, Uppsala University,
Uppsala, Sweden.
[8]Department of Medicine, Division of Cardiovascular Medicine, Stanford University
School of Medicine, Stanford, CA, USA.
[9]Stanford Cardiovascular Institute, Stanford University, Stanford, CA 94305.
[10]Department of Information Technology, Visual Information and Interaction,
Uppsala University, Uppsala, Sweden.
[11]Department of Public Health and Caring Sciences, Uppsala University, Uppsala,
Sweden.

[#]Corresponding author: marcel.den_hoed@igp.uu.se

30
31 Running title: Causal genes for sinoatrial arrest in HRV loci
32
33 Keywords: heart rate variability, CRISPR-Cas9, zebrafish, sinoatrial arrest, GWAS

**ABSTRACT**

A meta-analysis of genome-wide association studies (GWAS) recently identified eight loci that are associated with heart rate variability (HRV) in data from 53,174 individuals. However, functional follow-up experiments - aiming to identify and characterize causal genes in these loci - have not yet been performed. We developed an image- and CRISPR-Cas9-based pipeline to systematically characterize candidate genes for HRV in live zebrafish embryos and larvae. Nine zebrafish orthologues of six human candidate genes were targeted simultaneously in fertilized eggs from fish that transgenically express GFP on smooth muscle cells (Tg(*acta2:GFP*)), to visualize the beating heart. An automated analysis of 30s recordings of 384 live zebrafish atria at 2 and 5 days post-fertilization helped identify genes that influence HRV (*kiaa1755* and *gngt1*); heart rate (*kiaa1755*); sinoatrial pauses and arrests (*syt10*, *hcn4* and *kiaa1755*); and early cardiac development (*gngt1, neo1a*). Hence, comprehensively characterizing candidate genes in GWAS-identified loci for HRV *in vivo* helped us identify previously unanticipated culprits for life-threatening cardiac arrhythmias.

## INTRODUCTION

Heart rate variability (HRV) reflects the inter-beat variation of heart rate. HRV is controlled by the sinoatrial node, which receives input from the autonomic nervous system. Autonomic imbalance, which has been associated with work stress and other modifiable or non-modifiable risk factors (Thayer et al. 2010), is reflected in lower HRV. Lower HRV has been associated with higher cardiac morbidity and mortality (de Bruyne et al. 1999), and with a higher risk of all-cause mortality (Dekker et al. 1997). HRV can be quantified non-invasively using the RR interval of a 12-lead ECG, making HRV a useful clinical marker for perturbations of the autonomic nervous system. However, the genetic basis of HRV remains largely elusive.

Recently, we and others identified the first loci that are robustly associated with HRV, using a meta-analysis of genome-wide association studies (GWAS) with data from 53,174 participants (Nolte et al. 2017). Eleven of the 17 associated single nucleotide polymorphisms (SNPs) overlapped with loci that we previously identified as being associated with resting heart rate (den Hoed et al. 2013). The heart rate associated loci in turn had on aggregate been associated with altered cardiac conduction and risk of sick sinus syndrome (den Hoed et al. 2013). *In silico* functional annotation of the five loci that are associated with both HRV (Nolte et al. 2017) and heart rate (den Hoed et al. 2013) previously resulted in the prioritization of six candidate genes that are anticipated to be causal (Nolte et al. 2017). Functional follow-up experiments - ideally *in vivo* - are required to conclude if these genes are indeed causal, and examine if they influence HRV, heart rate, and/or cardiac conduction.

Mouse models are commonly used in cardiac research, but mice show substantial differences in cardiac rate and electrophysiology compared with humans (Poon and

Brand 2013). Inherently, these differences complicate extrapolation of results from

mouse models to humans (Poon and Brand 2013). Additionally, rodents are not

suitable for high-throughput, *in vivo* screens of cardiac rhythm and rate. Such screens

are essential to systematically characterize positional candidate genes in the large

number of loci that have now been identified by GWAS for cardiac rhythm, rate

(Eppinga et al. 2016) and conduction (Arking et al. 2014). Hence, novel model

systems that facilitate systematic, *in vivo* characterization of a large number of

candidate genes are desirable.

In recent years, the zebrafish has become an important model system for genetic

and drug screens for human disease (Lieschke and Currie 2007; MacRae and Peterson

2015). Amongst the many advantages of zebrafish larvae as a model system are their

short generation time, large number of offspring produced regularly, relatively low

maintenance costs, rapid early development, and optical transparency. Zebrafish have

a fully functional, beating heart at ~24 hours post-fertilization, and the ECG of the

two-chambered zebrafish heart is comparable to that of humans (Liu et al. 2016).

Fluorescently labeled transgenes facilitate visualization of cell types and tissues of

interest, which can now be accomplished in high-throughput thanks to advances in

automated positioning of non-embedded, live zebrafish embryos (Pardo-Martin et al.

2013). In addition, the zebrafish has a well-annotated genome, with orthologues of at

least 71.4% of human genes (Howe et al. 2013). These genes can be targeted

efficiently and in a multiplexed manner (i.e. in multiple genes simultaneously) thanks

to recent advances in Clustered, Regulatory Interspaced, Short Palindromic Repeats

(CRISPR) and CRISPR-associated systems (Cas) (Varshney et al. 2016). All

characteristics combined make zebrafish embryos an attractive model system to

systematically characterize candidate genes for cardiac rhythm, rate and conduction.

The aim of this study was to objectively characterize the most promising candidate

genes in GWAS-identified loci for HRV and heart rate for a role in cardiac rhythm,

rate and function using a large-scale, image-based screen in zebrafish embryos and

larvae.

**RESULTS**

*Experimental pipeline*

Our experimental pipeline is outlined in Figure 1. Founder mutants ($F_0$ generation) were generated in which all nine zebrafish orthologues of six human candidate genes were targeted using a multiplexed CRISPR-Cas9 approach (Table 1, Supplemental Table S1) (Varshney et al. 2016). Generating mutants in a background with a transgenically expressed fluorescent label on smooth muscle cells (Tg(*acta2*:*gfp*)) allowed us to visualize the beating atrium with minimal background noise. $F_0$ mutants were in-crossed, and the atria of 384 $F_1$ embryos were recorded for 30s at 2 days post fertilization (dpf). In addition, we captured bright field images of the embryos to quantify body length, surface area and volume. Of the 384 imaged embryos, 326 were imaged again at 5dpf (Supplemental Fig. S1), to capture genetic effects on cardiac outcomes and body size at two key stages of early cardiac development (Hou et al. 2014).

Eight loci associated with HRV[1].
18 genes prioritized for functional follow-up[1].

Six human genes selected for follow-up,
represented by nine orthologues in zebrafish.

CRISPR-gRNA design,
synthesis and efficiency testing.

Raising of CRISPR-Cas9 induced mutant founder fish ($F_0$).
Crossing of $F_0$-Generation for experiments.

Acquistion of 30 s videos of the atrium of
384 embryos (2dpf) & larvae (5dpf).

Automatic quantification of 30 s videos
using a custom-written MatLab script.

Paired-end sequencing of CRISPR-targeted
sites using a MiSeq.

Splitting, merging, QC-ing and aligning
of sequencing data.

Transcript-specific variant calling and
annotation using Ensembl's VEP.

Genetic association analysis using hierarchical linear
models and multiple logistic regression analyses.

**Fig. 1**: Experimental pipeline describing the workflow from candidate gene selection to trait association. dpf = days post fertilization. VEP = Variant Effect Predictor.
1: Nolte et al. 2017

**Table 1: Selected candidate genes and their zebrafish orthologues in GWAS-identified heart rate variability associated loci**

| rsID | Chr | Human gene | ENSG | Zebrafish orthologue | ENSDARG |
|---|---|---|---|---|---|
| rs4262, rs180238 | 7 | GNG11 | ENSG00000127920 | gngt1 | ENSDARG00000035798 |
| rs7980799, rs1351682, rs1384598 | 12 | SYT10 | ENSG00000110975 | syt10 | ENSDARG00000045750 |
| rs4899412, rs2052015, rs2529471, rs36423 | 14 | RGS6 | ENSG00000182732 | rgs6 | ENSDARG00000015627 |
| rs2680344 | 15 | HCN4 | ENSG00000138622 | hcn4 | ENSDARG00000061685 |
| | | | | hcn4l | ENSDARG00000074419 |
| rs1812835 | 15 | NEO1 | ENSG00000067141 | neo1a | ENSDARG00000102855 |
| | | | | neo1b | ENSDARG00000075100 |
| rs6123471 | 20 | KIAA1755 | ENSG00000149633 | quo (kiaa1755_1) | ENSDARG00000073684 |
| | | | | si:dkey-65j6.2 (kiaa1755_2) | ENSDARG00000103586 |

rsID refers to the lead SNP(s) of the GWAS-identified loci (Nolte et. al, 2017). Multiple independent SNPs per locus can be present. For simplicity, *quo* is referred to as *kiaa1755_1*, and *si:dkey-65j6.2* as *kiaa1755_2*.

*Descriptive information*

At 2dpf, some embryos showed sinoatrial pauses (n=39, 10.3%, Supplemental Movie

S1) and arrests (n=36, 9.5%, Supplemental Movie S2); cardiac edema (n=15, 4.0%);

or uncontrolled atrial contractions (n=1, 0.2%, Supplemental Movie S3). At 5dpf, a

subset of larvae displayed sinoatrial pauses (n=9, 2.7%); sinoatrial arrests (n=3,

0.9%); cardiac edema (n=14, 4.2%); uncontrolled atrial contractions (n=9, 2.7%); or

an abnormal cardiac morphology and impaired cardiac contractility (n=6, 1.8%,

Supplemental Movie S4). In the 279 embryos without cardiac abnormalities that

survived quality control at 2dpf (Supplemental Fig. S1), HRV was 15.5±10 ms

(mean±SD) and heart rate 208±24 bpm. In the 291 larvae included in the analysis at

5dpf, HRV was 13.18±17.06 ms and heart rate 265±35 bpm (Table 2).

**Table 2: Descriptive information on HRV and heart rate at 2 and 5dpf**

| Age | Trait | Mean | SD | 1st | Median | 3rd |
|-----|-------|------|-----|-----|--------|-----|
|     | SDNN  | 19.52 | 12.60 | 9.84 | 17.39 | 26.55 |
|     | RMSSD | 11.60 | 8.93 | 6.43 | 8.73 | 13.12 |
| 2   | HRV   | 15.56 | 10.04 | 8.74 | 12.96 | 19.65 |
|     | HR    | 208 | 24 | 192 | 207 | 225 |
|     | SDNN  | 13.30 | 17.18 | 5.18 | 7.60 | 11.72 |
|     | RMSSD | 13.05 | 18.26 | 5.93 | 7.48 | 12.01 |
| 5   | HRV   | 13.18 | 17.06 | 5.65 | 7.72 | 12.06 |
|     | HR    | 265 | 35 | 244 | 261 | 286 |

Information based on data from 279 embryos at 2 days post-fertilization (dpf) and 291 larvae at 5 dpf. HRV was calculated by averaging SDNN and RMSSD. HRV, SDNN and RMSSD are shown in ms, whereas HR is shown in beats per min. 1st and 3rd refers to the respective quartile. HRV: heart rate variability; SDNN: standard deviation of the normal-to-normal interval; RMSSD: root mean square of successive heart beat interval differences; HR: heart rate.

After imaging at 5dpf, all larvae were sequenced at the nine CRISPR-Cas9

targeted and three putative off-target sites (2x250 bp paired end), and transcript-

specific dosages were calculated by weighting the number of mutated alleles by their

predicted impact on protein function based on Ensembl's variant effect predictor

(VEP). At the three putative off-target sites, no CRISPR-Cas9 induced mutations

were identified (Supplemental Table S2). A total of 164 unique mutations were

identified across the nine CRISPR-targeted sites, ranging from three unique mutations

in *hcn4l* to 37 in *kiaa1755_2* (Supplemental Table S3). Frameshift mutations were

most common (48.5%), followed by missense variants (24.5%), in-frame deletions

(14.7%), and synonymous variants (5.5%). Eighty-five, sixty-nine and nine variants

were predicted to have a high, moderate, and low impact on protein function,

respectively (Supplemental Table S3). Mutant allele frequencies ranged from 4.1%

for *hcn4l* to 92.0% for *neo1b* (Supplemental Table S4).

*Genetic effects on cardiac rhythm, rate and function*

We examined the effect of the induced mutations on HRV and heart rate after

excluding cases with cardiac abnormalities that would drive effects on HRV and heart

rate, i.e. sinoatrial pauses or arrests, uncontrolled atrial contractions, cardiac edema,

and abnormal cardiac morphology or contractility (primary analyses, Supplemental

Fig. S1). We also examined the impact of mutations on cardiac abnormalities for

events that affected at least five embryos (2dpf) or larvae (5dpf), and we examined the

effect of mutations on body size (secondary analyses, Supplemental Fig. S1).

Each additional mutated allele in *gngt1* was associated with higher odds of

abnormal cardiac morphology and impaired cardiac contractility at 2dpf

(Supplemental Table S5), as well as with higher odds of cardiac edema and a higher

HRV at 5dpf (Figure 2, Supplemental Table S6-S7). *Gngt1* mutants also had a smaller

dorsal body surface area at 5dpf.

Each additional mutated allele in the main transcript of *kiaa1755_1* was associated

with a higher HRV and a trend for a higher heart rate at 2dpf (Figure 2, Supplemental

Tables S6-S7). Mutations in *kiaa1755_1* were also associated with a larger body

volume at 5dpf (Supplemental Table S8). Embryos with mutations in *kiaa1755_2*

tended to be shorter and smaller at 2dpf (Supplemental Table S8), and had higher

odds of sinoatrial pauses and a higher heart rate at 5dpf (Figure 2, Supplemental

Tables S5-S7).

Each additional mutated allele in *hcn4* was associated with higher odds of

sinoatrial pauses and arrests at 2dpf (Supplemental Table S5). In larvae free from

severe cardiac abnormalities, each additional mutated allele in *hcn4* was associated

with a trend for higher heart rate at 5dpf (Figure 2, Supplemental Tables S6-S7),

without affecting body size (Supplemental Table S8). Mutations in *hcn4l* were

positively associated with lateral body surface area at 5dpf (Supplemental Table S8),

but not with cardiac rhythm, rate, or abnormalities at 2dpf or 5dpf.

Mutations in *neo1a* and *neo1b* did not affect cardiac rhythm, rate, or abnormalities

at 2dpf. However, embryos with mutations in *neo1a* tended to be shorter and had a

larger dorsal body surface area normalized for length. Embryos with mutations in

*neo1b* had a larger lateral body surface area at 2dpf. At 5dpf, mutations in *neo1a* were

associated with lower odds of abnormal cardiac morphology and impaired cardiac

contractility, uncontrolled atrial contractions (trend), and edema (Supplemental Table

S5), but not with body size.

Each additional mutated allele in *syt10* was associated with higher odds of a

sinoatrial pause at 2dpf (Supplemental Table S5), but did not affect HRV, heart rate

or body size (Supplemental Tables S6-S8).

**Fig. 2**: Effect of mutations in candidate genes on heart rhythm and rate at 2 (n=279) and 5dpf (n=291). Dots and whiskers show the effect size and its 95% confidence interval for each additional mutated allele, weighted by its predicted effect on protein function. Effects were adjusted for the number of mutated alleles in the other genes, as well as for heart rate or heart rate variability and time of day (fixed factor), with larvae nested in batches (random factor). Genes are ordered by chromosomal position in humans.

Five of the six human candidate genes taken forward for experimental follow-up

showed at least one relevant cardiac phenotype in zebrafish embryos or larvae (i.e.

*GNGT1*, *SYT10*, *HCN4*, *NEO1*, *KIAA1755*). Only a trend towards an effect - on

sinoatrial pauses at 2dpf - was observed for the *RGS6* orthologue, for which the

mutant allele frequency was only 4%. We next examined the druggability of the five

genes for which we showed most evidence of a causal role using the drug-gene

interaction (DGI) database. This only highlighted *HCN4* as being druggable (Cotto et

al. 2018). Indeed, ivabradine is already being used as an open channel blocker of

*HCN4* (Böhm et al. 2010). We next identified interaction partners of the proteins

encoded by the five genes using GeneMania (Warde-Farley et al. 2010) and STRING

(Szklarczyk et al. 2017). A total of 49 interacting partners were identified, 31 of

which were considered to be druggable or belong to a "favorable" protein family.

Seven of these are targeted by FDA-approved medication (Supplemental Table S9),

amongst which are several anti-hypertensive agents (e.g. Telmisartan, Iloprost and

Diazoxide), as well as a neuromuscular blocking agent (i.e. Botulinum toxin type A).

It would be worthwhile to examine if these agents can be repurposed to prevent

cardiac arrhythmias in individuals at risk, or whether cardiac arrhythmias represent

on-target, adverse side effects of these drugs.

**DISCUSSION**

Large-scale, *in vivo* follow-up studies of candidate genes in GWAS-identified loci remain sparse. Here we present an objective, image-based pipeline to systematically characterize candidate genes for cardiac rhythm, rate, and conduction-related disorders. Using a zebrafish model system, we identified orthologues of genes in HRV and heart rate associated loci that influence not only the complex traits for which they were identified (i.e. *kiaa1755*, *gngt1*), but also body size (*kiaa1755, hcn4, gngt1, neo1a*, *neo1b*), early cardiac development (*gngt1, neo1a, neo1b*), and/or the odds of sinoatrial pauses and arrests (*hcn4, syt10, kiaa1755*). These results add weight to the notion that GWAS-identified common variants for complex traits can flag genes for which rare, detrimental mutations cause severe, early-onset disorders (Arking et al. 2014; Flannick et al. 2016; Ganna et al. 2018). We show here that systematic characterization of candidate genes in GWAS-identified loci for complex traits using an image- and CRISPR-Cas9-based zebrafish model system can help identify clinically relevant causal genes, and aid confirmation of the role of such genes in whole-exome sequencing efforts in humans by reducing the multiple testing burden (Flannick et al. 2018).

The HRV increasing T-allele of rs4262 in the 5' untranslated region of *GNG11* is associated with a lower expression of *GNG11* in the tibial artery, atrial appendage and aorta (GTEx (Battle et al. 2017)). This is directionally consistent with 5dpf *gngt1* mutant zebrafish larvae - i.e. mutants for the zebrafish orthologue of the human paralogues *GNGT1* and *GNG11* (Lagman et al. 2012) - having a higher HRV. *GNG11* encodes the γ-subunit of a G-protein complex, which interacts with G-protein-coupled inwardly rectifying potassium channels (GIRKs) that transduce signals rapidly enough to induce changes in HRV (Mark and Herlitze 2000). The zebrafish heart

normally has a tube-like appearance at 28 hours post fertilization (hpf) (Bakkers

2011). Higher odds of abnormal cardiac morphology, contractility and edema with

each additional mutated *gngt1* allele at 5 but not 2dpf suggests that this gene affects

cardiac development between 2 and 5dpf.

Non-synonymous SNPs in *KIAA1755* have been identified in GWAS for HRV

(Nolte et al. 2017) and heart rate (den Hoed et al. 2013). In humans, the C-allele of

the HRV-associated rs6123471 in the 3' UTR of *KIAA1755* is associated with a lower

expression of *KIAA1755* in brain and aorta (GTEx (Battle et al. 2017)), a higher HRV

(Nolte et al. 2017), and a lower heart rate (den Hoed et al. 2013). Our results show

that mutations in *kiaa1755_1* predispose to a higher HRV and, in a sensitivity

analysis, a higher heart rate at 2dpf. Mutations in *kiaa1755_2* on the other hand result

in a higher heart rate and higher odds of sinoatrial pauses at 5dpf. Hence, a higher

HRV at 2dpf for each additional mutated allele in *kiaa1755_1* is directionally

consistent with results in humans, while the effects of mutations in *kiaa1755_1* and

*kiaa1755_2* on heart rate are not. Genetic effects on HRV and heart rate were

mutually adjusted in our study, which may have revealed the unbiased direction of

effect for mutations in *KIAA1755* on heart rate. In other words: the GWAS-identified

association of the *KIAA1755* locus - and possibly other loci - with heart rate may have

been driven by HRV. This would explain why the eleven HRV-associated loci that

showed evidence of an association with heart rate all did so in the expected (i.e.

opposite) direction from a phenotypic point of view (Nolte et al. 2017). *KIAA1755* is

a previously uncharacterized gene that shows a broad expression pattern, including

different brain regions and the left atrial appendage (GTEx (Battle et al. 2017)).

Future mechanistic studies are required to distill how *KIAA1755* influences heart

rhythm and rate.

The locus harboring *HCN4* has been flagged in GWAS for HRV (Nolte et al.

2017), heart rate (den Hoed et al. 2013) and atrial fibrillation (Ellinor et al. 2012).

*HCN4* is arguably the most well-characterized gene in the context of cardiac rhythm.

It belongs to the family of "funny" channels, aptly named for being activated upon

hyperpolarization. It is expressed in the sinoatrial node (Ludwig et al. 1999; Shi et al.

1999) and plays an important role in cardiac pace making (Stieber et al. 2003). The

heart-rate lowering agent ivabradine (Bucchi et al. 2013) is an open channel blocker

of *HCN4*, and reduces cardiovascular and all-cause mortality in heart failure patients

(Swedberg et al. 2010). Hcn4$^{-/-}$ mice die prenatally, and analyses of their hearts show

no arrhythmias and a lower heart rate compared with wildtype and heterozygotes

(Stieber et al. 2003). In line with this, humans that are heterozygous for an *HCN4* loss

of function mutation are typically characterized by bradycardia. We identified nine

zebrafish embryos that were compound heterozygous for mutations in *hcn4*, and one

embryo that was compound heterozygous for *hcn4l*, without a deviation from HWE

for either gene (Supplemental Table S4). Four of the nine compound heterozygous

larvae for *hcn4*, and six of the eight larvae that were heterozygous for mutations in

*hcn4* as well as *hcn4l* showed a sinoatrial arrest before or during the atrial recording at

2dpf. Zebrafish embryos can survive without a functional heart at this stage, thanks to

tissue oxygenation by diffusion (Burggren and Pinder 1991). This allowed us to

witness genetically driven cardiac arrests that would have been lethal in embryos of

other species. At 5dpf, larvae carrying mutations in *hcn4* that were free from severe

cardiac abnormalities tended to have a higher heart rate than larvae without CRISPR-

induced mutations. This unexpected result may reflect overcompensation by *hcn4l* or

cardiac compensation in response to unobserved sinoatrial arrests outside the 30s

recording.

In addition to the locus in *HCN4*, the GWAS for HRV identified an independent

locus in the adjacent *NEO1* (Nolte et al. 2017). Neogenin-1 is a transmembrane

receptor that is involved in cell migration, angiogenesis and neuronal survival

(reviewed by Wilson et al. (Wilson and Key 2007)). In line with our results,

morpholino-mediated knockdown of *neo1a* expression in zebrafish larvae was

previously shown to result in shorter, thicker larvae due to defects in somitogenesis,

with loss of neuronal differentiation due to malformation of the neural tube

(Mawdsley et al. 2004). *NEO1* has a pro-apoptotic function in humans (Wilson and

Key 2007), so the cardioprotective effect we observed in 5dpf mutants may result

from inefficient propagation of apoptotic signals by specific effectors. Inhibiting

NEO1 in early development may thus help protect cardiac integrity - perhaps

temporarily - but will likely have adverse effects due to inadequate apoptosis at later

stages of development.

*SYT10* belongs to the family of synaptogamins: transmembrane proteins that are

involved in regulation of membrane trafficking in neurons (Sudhof 2002), which is

important for neurotransmitter release (Chapman 2008). Variants in/near *SYT10* have

previously been associated with HRV (Nolte et al. 2017) as well as with heart rate

(den Hoed et al. 2013) and heart rate recovery (Verweij et al. 2018). We observed an

effect of mutations in *syt10* on the odds of sinoatrial pauses at 2dpf. Since *SYT10* is

mainly expressed in brain, tibial nerve and pituitary (GTEx (Battle et al. 2017)),

mutations in *syt10* may affect the odds of sinoatrial pauses through altered

neurotransmitter release.

Of the five genes for which we show causal effects, only *HCN4* is highlighted as

being druggable by DGIdb. The protein encoded by *HCN4* has already been targeted

using several compounds (Stieber et al. 2006). Although four of the five causal genes

are not highlighted as being druggable by current pharmaceutical standards, targeting

via antisense oligonucleotides or antibodies can still be explored. Expanding our

search revealed several druggable interaction partners of causal genes that are already

targeted by FDA-approved anti-hypertensive and neuromuscular blocking agents,

amongst others. For existing drugs that target interaction partners of the five causal

genes, it is worthwhile examining their effect on cardiac rhythm, rate and

development, since repurposing FDA-approved drugs would imply the quickest and

safest route to the clinic, while quantifying possibly unknown adverse side effects

would also be informative.

A few limitations of our study should be discussed. Firstly, acquiring 30s

recordings implies that false negatives for sinoatrial pauses or arrests are inevitable. In

fact, we observed 54 pauses or arrests at 2dpf (14.3%) and eight at 5dpf (2.5%) while

positioning the larva for imaging, without observing abnormalities during the

subsequent 30s recording. This suggests that at least 15.9% of all embryos without a

sinoatrial pause at 2dpf are false negatives, which negatively affects the statistical

power to find genetic effects. This limitation will have resulted in conservative effect

estimates. Secondly, CRISPR guide-RNAs with predicted off-targets free from

mismatches were avoided. However, two of the selected targets - i.e. for *hcn4* and

*kiaa1755_2* - have exonic off-targets with three mismatches (Supplemental Table S1).

Human orthologues of potential off-targets *dclk1* and *galnt10* have previously been

associated with heart-rate variability-related traits (Newton-Cheh et al. 2007) (*dclk1*),

as well as with carotid intima-media thickness (Dong et al. 2015) and body mass

index (Locke et al. 2015; Justice et al. 2017; Ng et al. 2017) (*galnt10*). Sequencing

larvae at the three predicted off-target regions did not show any CRISPR-Cas9

induced mutations (Supplemental Table S2). In line with predictions by Varshney *et*

*al.*, off-target effects are thus unlikely to have influenced our results (Varshney et al.

2015). Thirdly, we in-crossed mosaic founder mutants ($F_0$) and phenotypically

screened and sequenced the $F_1$ generation. For some genes, this yielded a small

number of larvae with 0 or 2 mutated alleles (i.e. *rgs6*, *hcn4*, and *hcn4l*), resulting in

a lower power to find true genetic effects for these genes. In spite of this limitation,

we were able to detect significant effects of mutations in all genes except *rgs6*, for

which we observed a trend for an effect on the odds of sinoatrial pauses at 2dpf.

Furthermore, screening the $F_0$ or $F_2$ generation instead has its own advantages and

disadvantages, and we argue that non-conclusive results for a small subset of

candidate genes in the primary screen is not problematic when aiming to

systematically characterize a large number of candidate genes. Finally, we recorded

the atrium only, to enable a higher frame rate, a higher resolution in time for HRV,

and a higher statistical power to detect small genetic effects on HRV. As a result, any

ventricular abnormalities were not registered, and uncontrolled atrial contractions

may thus reflect atrial fibrillation, premature atrial contractions, high atrial rate, or

atrial tachycardia. Additional studies with a larger frame and longer acquisition time

are required to replicate and clarify the effect of mutations in *neo1a* on uncontrolled

atrial contractions.

Our study benefits from a repeated measures design, which enabled us to capture

genetic effects at different stages of early development in zebrafish. Furthermore, the

throughput of the setup allowed us to examine the effect of multiple genes

simultaneously, in an unprecedented sample size for *in vivo* genetic screens. Our

results demonstrate that a large sample size is paramount to robustly detect genetic

effects on complex traits in zebrafish embryos and larvae, even for functional

knockout mutations. Identifying CRISPR-Cas9-induced mutations allele-specifically

helped us classify larvae as heterozygous and compound heterozygous, which in turn

helped pinpoint the true effect of mutations in these genes.

Our study is based on an objective, high-throughput imaging approach and aimed

to identify and characterize causal genes in GWAS-identified loci for HRV. In all

genes that show an effect on HRV (*kiaa1755* and *gngt1*), directions of effect were

directionally consistent with eQTL associations in humans (based on GTEx (Battle et

al. 2017)). This emphasizes the merit of our model system and the robustness of our

findings. Furthermore, our results extend beyond the complex outcomes for which

these genes were prioritized, and helped identify *KIAA1755*, *HCN4* and *SYT10* as

likely effectors of detrimental mutations that predispose to sinoatrial pauses and

arrests.

**METHODS**

*Candidate gene selection*

Candidate genes in GWAS-identified loci for HRV were identified as described in detail in Nolte et al. (2017). Of the 18 identified candidate genes, six were selected for experimental follow-up. This selection was based on overlap with findings from GWAS for heart rate (*KIAA1755, SYT10, HCN4, GNG11*) (den Hoed et al. 2013), as well as with results from eQTL analyses in sinoatrial node and brain (*RGS6*). Additional candidate genes from the same or nearby loci were also selected for experimental follow-up, i.e. *NEO1*, which resides next to *HCN4*. Zebrafish orthologues of the human genes were identified using Ensembl, as well as using a comprehensive synteny search using Genomicus (Louis et al. 2015) (Supplemental Table S1). Of the selected genes, *GNG11, SYT10* and *RGS6* have one orthologue in zebrafish, and *KIAA1755, HCN4* and *NEO1* each have two orthologues, resulting in a total of nine zebrafish orthologues for six human candidate genes (Table 1). For simplicity, the *KIAA1755* orthologues *quo* and *si:dkey-65j6.2* have been referred to as *kiaa1755_1* and *kiaa1755_2*.

*Mutagenesis*

All nine zebrafish genes were targeted together using a multiplexed CRISPR-Cas9 approach described recently (Varshney et al. 2016). Briefly, guide-RNAs (gRNAs) were selected using ChopChop (Labun et al. 2016) and CRISPRscan (Moreno-Mateos et al. 2015) (Supplemental Table S1), based on their predicted efficiency, a moderate to high GC-content, location in an early exon, and absence of predicted off-target effects without mismatches. Oligonucleotides were designed as described (Varshney

et al. 2016), consisting of a T7 or SP6 promotor sequence (for gRNAs starting with

'GG' or 'GA', respectively), a gene-specific gRNA-target sequence, and an overlap

sequence to a generic gRNA. The gene-specific oligonucleotides were annealed to a

generic 80 bp long oligonucleotide at 98°C for 2 mins, 50°C for 10 mins, and 72°C

for 10 mins. The products were checked for correct length on a 2% agarose gel. The

oligonucleotides were subsequently transcribed *in vitro* using the manufacturer's

instructions (TranscriptAid T7 high yield transcription kit / MEGAscript SP6

transcription kit, both ThermoFisher Scientific, Waltham, USA). The gRNAs were

purified, after which the integrity of the purified gRNAs was examined on a 2%

agarose gel. The zebrafish codon-optimized plasmid pT3TS-nls-zCas9-nls was used

as a template to produce Cas9 mRNA (Jao et al. 2013). The plasmid was linearized

with Xba1, and then purified using the Qiaprep Spin Miniprep kit (Qiagen, Hilden,

Germany). The DNA was transcribed using the mMESSAGE mMACHINE T3

Transcription Kit (ThermoFisher Scientific, Waltham, USA), followed by LiCl

precipitation. The quality of the RNA was confirmed on a 1% agarose gel.

*Husbandry & microinjections*

A zebrafish line with GFP-labelled α-smooth muscle cells Tg(*acta2*:GFP) (Whitesell

et al. 2014) was used to visualize the beating heart. To this end, eggs from an in-cross

of Tg(*acta2*:GFP) fish were co-injected with a mix of Cas9 mRNA (final

concentration 150 ng/µl) and all nine gRNAs (final concentration 25 ng/µl each) in a

total volume of 2nL at the single-cell stage. CRISPR-Cas9 injected embryos were

optically screened for the presence of Tg(*acta2*:GFP) at 2 days post fertilization (dpf),

using an automated fluorescence microscope (EVOS FL Cell imaging system,

ThermoFisher Scientific, Waltham, USA). Tg(*acta2*:GFP) carriers were retained and

raised to adulthood in systems with circulating, filtered and temperature controlled water (Aquaneering, Inc, San Diego, CA). All procedures and husbandry were conducted in accordance with Swedish and European regulations, and have been approved by the Uppsala University Ethical Committee for Animal Research (C142/13 and C14/16).

*Experimental procedure imaging*

The founder mutants ($F_0$ generation) were only used for reproduction. After crossing the founder mutants, $F_1$ embryos were used for experiments. Eggs were collected after $F_0$ fish were allowed to reproduce for 45 mins to minimize variation in developmental stage. Fertilized eggs were placed in an incubator at 28.5°C. At 1dpf, embryos were dechorionated using pronase (Roche Diagnostics, Mannheim, Germany).

At 2dpf, embryos were removed from the incubator and allowed to adapt to controlled room temperature (21.5 °C) for 20 mins. Individual embryos were exposed to 100 μg/ml Tricaine (MS-222, Sigma-Aldrich, Darmstadt, Germany) for 1 min before being aspirated, positioned in the field of view of a fluorescence microscope, and oriented dorsally using a Vertebrate Automated Screening Technology (VAST) BioImager (Union Biometrica Inc., Geel, Belgium). We subsequently acquired twelve whole-body images, one image every 30 degrees of rotation, using the camera of the VAST BioImager to quantify body length, dorsal and lateral surface area and volume, as well as the presence or absence of cardiac edema. The VAST BioImager then positioned and oriented the larva to visualize the beating atrium and triggered the upright Leica DM6000B fluorescence microscope to start imaging using a HCX APO L 40X/0.80 W objective and L5 ET, k filter system (Micromedic AB, Stockholm,

Sweden). Images of the beating atrium were acquired for 30s at a frame rate of 152

frames/s using a DFC365 FX high-speed CCD camera (Micromedic AB, Stockholm,

Sweden). After acquisition, the larvae were dispensed into a 96-well plate, rinsed

from tricaine, and placed back into the incubator. The procedure was repeated at 5dpf

(i.e. the larval stage), to allow genetic effects that influence HRV and heart rate

differently at different stages of development to be captured (Hou et al. 2014). After

imaging at 5dpf, the larvae were once again dispensed into 96-well plates, sacrificed,

and stored at -80°C for further processing.

*Quantification of cardiac traits and body size*

A custom-written MATLAB script was used to convert the images acquired by the

CCD camera into quantitative traits. To acquire the heart rate, each frame of the

sequence was correlated with a template frame. The repeating pattern yields a

periodic graph from the correlation values and by detecting the peaks in the graph we

can assess the heart rate. The template frame should represent one of the extreme

states in the cardiac cycle, i.e. end-systole or end-diastole. To detect these frames, we

examined the correlation between the first 100 frames. The combination of frames

that showed the lowest correlation corresponded to the heart being in opposite states.

One of these frames was chosen as the template (Niazi et al. 2009).

This numeric information was subsequently used to quantify: 1) heart rate as the

inverse of RR-interval; 2) the standard deviation of the normal-to-normal RR interval

(SDNN); and 3) the root mean square of successive heart beat interval differences

(RMSSD). Finally, a graph of pixel changes over time was generated across the 30s

recording to help annotate the script's performance. The files containing the inter-

beat-intervals were used to objectively quantify sinoatrial pauses (i.e. the atrium stops

beating for longer than 3x the median inter-beat-interval of the larva, Supplemental

Movie S1) and sinoatrial arrests (i.e. the atrium stops beating for longer than 2s,

Supplemental Movie S2) using a custom-written Stata script. The graphs of pixel

changes over time were also used to identify larvae with other abnormalities in

cardiac rhythm. Such abnormalities were annotated as: uncontrolled atrial

contractions (Supplemental Movie S3); abnormal cardiac morphology (i.e. a tube-like

atrium, Supplemental Movie S4); or impaired cardiac contractility (i.e. a vibrating

rather than a contracting atrium, Supplemental Movie S4). These phenotypes were

annotated independently by two investigators (B.v.d.H. and M.d.H.), resulting in an

initial concordance rate >90%. Discrepancies in annotation were discussed and re-

evaluated to reach consensus.

Bright-field images of the larvae were used to assess body length, dorsal and lateral

surface area, and volume. Images were automatically segmented and quantified using

a custom-written CellProfiler (Lamprecht et al. 2007) pipeline, followed by manual

annotation for segmentation quality. Larvae with suboptimal segmentation quality due

to the presence of an air-bubble, a bent body, an incomplete rotation within the

capillary, partial capturing of the larva, or an over-estimation of size were replaced by

images with a 180° difference in rotation, or excluded if that image was also sub-

optimally segmented. The larva was excluded from the analysis for body volume if

more than four of the 12 images had a bad segmentation. Imaging, image

quantification and image quality control were all performed blinded to the sequencing

results.

*Quality control of phenotype data*

A series of quality control steps was performed to ensure only high-quality data was

included in the genetic association analysis (Supplemental Fig. S1). First, graphs

indicating that one or more true beats were missed by the script were removed from

the analysis *a priori* (Supplemental Fig. S1). Second, larvae with any secondary

phenotype (i.e. uncontrolled atrial contractions, sinoatrial pause or arrest, edema,

abnormal morphology and/or reduced contractility) were excluded from the analyses

for HRV and heart rate (primary analysis) (Supplemental Fig. S1). Third, genetic

effects on these secondary traits were examined at 2dpf and 5dpf for traits with at

least five cases (secondary analysis). Furthermore, embryos and larvae that showed a

sinoatrial pause or arrest in between positioning and video acquisition, but not during

the recording, were excluded from the analysis for sinoatrial pauses and arrests, since

we cannot ascertain case status in the same rigorous manner for such samples, but

they are not appropriate controls either.

*Sample preparation for sequencing*

After imaging at 5dpf, larvae were sacrificed by prolonged exposure to tricaine, and

DNA was extracted by exposure to lysis buffer (10mM Tris-HCl pH8, 50mM KCl,

1mM EDTA, 0.3% Tween 20, 0.3% Igepal) and proteinase K (Roche Diagnostics,

Mannheim, Germany) for 2 h at 55°C, followed by 10 min at 95°C to deactivate the

proteinase K. Gene-specific primers (150bp-300bp) amplifying gRNA-targeted and

putative off-target regions in *dclk1b* and both *galnt10* orthologues (Supplemental

Table S1) were distilled from ChopChop (Labun et al. 2016) and Primer3 (Koressaar

and Remm 2007), and Illumina adaptor-sequences were added. The first PCR was

conducted by denaturation at 98°C for 30s; amplification for 35 cycles at 98°C for 10s, 62°C for 30s and 72°C for 30s; followed by a final extension at 72°C for 2 mins. Amplified PCR products were cleaned using magnetic beads (Mag-Bind PCR Clean-up Kit, Omega Bio-tek Inc. Norcross, GA). The purified products were used as a template for the second PCR, in which Illumina Nextera DNA library sequences were attached to allow multiplexed sequencing of all CRISPR-targeted sites across 384 larvae in a single lane. The second PCR amplification was performed by denaturation at 98°C for 30s; amplification for 25 cycles at 98°C for 10s, 66°C for 30s and 72°C for 30s; followed by a final extension at 72°C for 2 mins. Products were then purified using magnetic beads. All liquid handling was performed using a Hamilton Nimbus robot equipped with a 96-head (Hamilton robotics, Kista, Sweden). Samples were pooled and sequenced in a single lane on a MiSeq (300 bp paired-end, Illumina Inc., San Diego, CA) at the National Genomics Infrastructure, Sweden.

*Processing of sequencing data*

A custom-written bioinformatics pipeline was developed in collaboration with the National Bioinformatics Infrastructure Sweden to prepare .fastq files for analysis. First, a custom-written script was used to de-multiplex the .fastq files by gene and well. PEAR (Zhang et al. 2014) was then used to merge paired-end reads, followed by removal of low-quality reads using FastX (Pearson et al. 1997). The reads were then mapped to the wildtype zebrafish genome (Zv11) using STAR (Dobin et al. 2013). Next, we converted files from .sam to .bam format using samtools (Li et al. 2009), after which variants - mostly indels and SNVs - were called allele specifically using a custom-written variant calling algorithm in R (Danio rerio Identification of Variants by Haplotype - DIVaH). A summary of all unique sequences identified for the

orthologues and their respective alignment report string (Concise Idiosyncratic

Gapped Alignment Report (CIGAR)) is shown in Supplemental Table S2. All unique

variants (Supplemental Table S3) located within ±30bps of the CRISPR-targeted sites

that were identified across the two alleles were subsequently pooled, and used for

functional annotation using Ensembl's VEP (McLaren et al. 2016). Transcript-

specific dosage scores were then calculated by retaining the variant with the highest

predicted impact on protein function, and assigning it a score of 0 (no effect), 0.2

(modifier variant), 0.33 (low), 0.66 (moderate) or 1 (high). Transcript-specific dosage

scores were subsequently calculated by summing the allele-specific scores at each

embryo and target site. Since all transcripts within a target site were affected virtually

identically, we only used the main transcript of each target site for the genetic

association analysis.

Most larvae had successfully called sequences in all nine CRISPR-targeted sites

(n=288, 75%). Of the 96 remaining embryos, 83 larvae (22%) had a missing sequence

at one target site, and 11 larvae (3%) had two missing sequences. The remaining two

larvae (0.5%) had three and four missing sequences, respectively. These two larvae

have been excluded from the genetic association analysis. For the remaining larvae,

the mean dosage of the transcript was imputed for missing calls. At a gene-level, a

median of 380 larvae were successfully sequenced and called across all nine main

transcripts, and all but one gene had successfully called mutations in ≥375 larvae. For

*neo1b*, calling failed in 77 larvae (Supplemental Table S4). The imputed mean dosage

for the main transcript of *neo1b* was still included in the genetic association analysis,

since the mutant allele frequency in successfully called larvae was high (i.e. 0.798),

and using an imputed dosage was anticipated to influence the results less than to

discard the gene while it had been targeted. However, the results for *neo1b* should be

interpreted in light of its call rate. The mutant allele frequency was low for *hcn4*,

*hcn4l* and *rgs6*. For the remaining orthologues we had at least 10 wildtype and 10

compound heterozygous mutants. All three putative off-targets regions (*dclk1b*, both

*galnt10* orthologues, Supplemental Table S1) were classified as wildtype across all

larvae and had no CRISPR-Cas9 induced mutations within ±30bps of the cut-site

(Supplemental Table S2).

*Interrogating the druggability of the candidate genes*

All human orthologues of the zebrafish genes that we found an effect for were

interrogated in the drug-gene interaction database (DGIdb) v3.0.2 (Cotto et al. 2018).

The human interaction partners of the candidates (Supplemental Table S9) were

distilled from STRING v10.5 (Szklarczyk et al. 2017) and GeneMania (Warde-Farley

et al. 2010). The GeneMania search was limited to physical interactions and pathway

data.

*Statistical analysis*

The standard deviation of NN-intervals (SDNN) and the root mean square of

successive differences (RMSSD) were strongly correlated at 2dpf ($r^2$=0.77) and 5dpf

($r^2$=0.85), so a composite endpoint 'HRV' was calculated as the average of SDNN

and RMSSD. In larvae free from sinoatrial pauses and arrests, abnormal cardiac

morphology, impaired cardiac contractility, and edema HRV and heart rate at 2dpf

(n=279) or 5dpf (n=291) were inverse-normally transformed to ensure a normal

distribution, and to allow comparison of effect sizes across traits. Please see the

Supplemental methods for a detailed description of the cardiac phenotypes that were

used as exclusion criteria for the analysis for HRV and heart rate. Genetic effects on

HRV and heart rate were subsequently examined using hierarchical linear models at

2dpf and 5dpf separately (Stata's xtmixed), mutually adjusted for the other outcome

(i.e. heart rate and HRV) and time of imaging (fixed factors), and with larvae nested

in six batches (random factor with fixed slope). Embryos and larvae in which a

sinoatrial pause and/or arrest was observed in between positioning and recording were

excluded from the analysis (see Supplemental Table S7 for a sensitivity analysis). In

secondary analyses, genetic effects were examined for cardiac abnormalities that were

observed in ≥5 larvae at 2dpf and/or 5dpf, using a logistic regression analysis. Genetic

effects on body size were also examined, using hierarchical linear models on inverse-

normally transformed outcomes.

For dichotomous as well as continuous outcomes, genetic effects were examined

using an additive model, with dosage scores for all target sites as independent

exposures in the same model, i.e. mutually adjusted for effects of mutations in the

other orthologues. Missing genotypes were imputed at the mean in this analysis. For

each outcome, association analyses were examined for the main transcript of the

orthologue. We performed two independent primary and six independent secondary

tests at 2 and 5dpf – on different data – and considered $P$-values <0.05 to reflect

statistically significant effects. All statistical analyses were performed using Stata MP

14.2.

## DATA ACCESS

Raw data and scripts will be made publicly available after publication if desirable. Supplemental videos have been deposited at https://www.dropbox.com/s/3dboz7nq3emkz6z/Supp_videos.zip?dl=0.

## ACKNOWLEDGMENTS

The computations were performed on resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under Project SNIC b2015283. The authors would like to acknowledge support from Science for Life Laboratory, the National Genomics Infrastructure (NGI) and UPPMAX for providing assistance in massive parallel sequencing and computational infrastructure. Support from the National Bioinformatics Infrastructure Sweden (NBIS) is also gratefully acknowledged. MdH is a fellow of the Swedish Heart-Lung Foundation (20170872) and is supported by project grants from the Swedish Heart-Lung Foundation (20140543, 20170678), the Swedish Research Council (2015-03657), and NIH/NIDDK (R01DK106236, R01DK107786, U01DK105554).

**AUTHOR CONTRIBUTIONS**: B.v.d.H. and M.d.H. conceived the study design. B.v.d.H., A.E., T.K., S.V. and S.J. performed the experiments. B.v.d.H., A.E., E.M., O.D. and M.d.H. conceived the sequencing quality control and variant calling pipeline. B.v.d.H., A.A., H.L.B. and M.d.H. performed the image quantification and statistical analysis. B.v.d.H. and M.d.H. wrote the manuscript. All authors provided critical feedback to the manuscript.

634 **DISCLOSURE DECLARATION**

635 There are no competing interests.

**REFERENCES**

Arking DE Pulit SL Crotti L van der Harst P Munroe PB Koopmann TT Sotoodehnia N Rossin EJ Morley M Wang X et al. 2014. Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat Genet* **46**: 826-836.

Bakkers J. 2011. Zebrafish as a model to study cardiac development and human cardiac disease. *Cardiovasc Res* **91**: 279-288.

Battle A, Brown CD, Engelhardt BE, Montgomery SB. 2017. Genetic effects on gene expression across human tissues. *Nature* **550**: 204-213.

Böhm M, Swedberg K, Komajda M, Borer JS, Ford I, Dubost-Brama A, Lerebours G, Tavazzi L. 2010. Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *The Lancet* **376**: 886-894.

Bucchi A, Baruscotti M, Nardini M, Barbuti A, Micheloni S, Bolognesi M, DiFrancesco D. 2013. Identification of the molecular site of ivabradine binding to HCN4 channels. *PLoS One* **8**: e53132.

Burggren WW, Pinder AW. 1991. Ontogeny of cardiovascular and respiratory physiology in lower vertebrates. *Annu Rev Physiol* **53**: 107-135.

Chapman ER. 2008. How does synaptotagmin trigger neurotransmitter release? *Annu Rev Biochem* **77**: 615-641.

Cotto KC, Wagner AH, Feng YY, Kiwala S, Coffman AC, Spies G, Wollam A, Spies NC, Griffith OL, Griffith M. 2018. DGIdb 3.0: a redesign and expansion of the drug-gene interaction database. *Nucleic Acids Res* **46**: D1068-D1073.

de Bruyne MC, Kors JA, Hoes AW, Klootwijk P, Dekker JM, Hofman A, van Bemmel JH, Grobbee DE. 1999. Both decreased and increased heart rate variability on the standard 10-second electrocardiogram predict cardiac mortality in the elderly: the Rotterdam Study. *Am J Epidemiol* **150**: 1282-1288.

Dekker JM, Schouten EG, Klootwijk P, Pool J, Swenne CA, Kromhout D. 1997. Heart rate variability from short electrocardiographic recordings predicts mortality from all causes in middle-aged and elderly men - The zutphen study. *Am J Epidemiol* **145**: 899-908.

den Hoed M Eijgelsheim M Esko T Brundel BJ Peal DS Evans DM Nolte IM Segre AV Holm H Handsaker RE et al. 2013. Identification of heart rate-associated loci and their effects on cardiac conduction and rhythm disorders. *Nat Genet* **45**: 621-631.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15-21.

Dong C, Della-Morte D, Beecham A, Wang L, Cabral D, Blanton SH, Sacco RL, Rundek T. 2015. Genetic variants in LEKR1 and GALNT10 modulate sex-difference in carotid intima-media thickness: a genome-wide interaction study. *Atherosclerosis* **240**: 462-467.

Ellinor PT, Lunetta KL, Albert CM, Glazer NL, Ritchie MD, Smith AV, Arking DE, Muller-Nurasyid M, Krijthe BP, Lubitz SA et al. 2012. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat Genet* **44**: 670-675.

Eppinga RN, Hagemeijer Y, Burgess S, Hinds DA, Stefansson K, Gudbjartsson DF, van Veldhuisen DJ, Munroe PB, Verweij N, van der Harst P. 2016. Identification of genomic loci associated with resting heart rate and

shared genetic predictors with all-cause mortality. *Nat Genet* **48**: 1557-
1563.
Flannick J, Johansson S, Njolstad PR. 2016. Common and rare forms of diabetes
mellitus: towards a continuum of diabetes subtypes. *Nat Rev Endocrinol*
**12**: 394-406.
Flannick JA, Mercader JMM, Fuchsberger C, Udler MS, Mahajan A, Wessel J,
Altshuler D, Burtt NP, Scott L, Morris A et al. 2018. Genetic discovery and
translational decision support from exome sequencing of 20,791 type 2
diabetes cases and 24,440 controls from five ancestries. *bioRxiv*
doi:10.1101/371450.
Ganna A, Satterstrom FK, Zekavat SM, Das I, Kurki MI, Churchhouse C, Alfoldi J,
Martin AR, Havulinna AS, Byrnes A et al. 2018. Quantifying the Impact of
Rare and Ultra-rare Coding Variation across the Phenotypic Spectrum. *Am*
*J Hum Genet* **102**: 1204-1211.
Hou JH, Kralj JM, Douglass AD, Engert F, Cohen AE. 2014. Simultaneous mapping
of membrane voltage and calcium in zebrafish heart in vivo reveals
chamber-specific developmental transitions in ionic currents. *Front*
*Physiol* **5**: 344.
Howe K Clark MD Torroja CF Torrance J Berthelot C Muffato M Collins JE
Humphray S McLaren K Matthews L et al. 2013. The zebrafish reference
genome sequence and its relationship to the human genome. *Nature* **496**:
498-503.
Jao LE, Wente SR, Chen W. 2013. Efficient multiplex biallelic zebrafish genome
editing using a CRISPR nuclease system. *Proc Natl Acad Sci U S A* **110**:
13904-13909.
Justice AE Winkler TW Feitosa MF Graff M Fisher VA Young K Barata L Deng X
Czajkowski J Hadley D et al. 2017. Genome-wide meta-analysis of 241,258
adults accounting for smoking behaviour identifies novel loci for obesity
traits. *Nature communications* **8**: 14977.
Koressaar T, Remm M. 2007. Enhancements and modifications of primer design
program Primer3. *Bioinformatics* **23**: 1289-1291.
Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. 2016. CHOPCHOP v2: a
web tool for the next generation of CRISPR genome engineering. *Nucleic*
*Acids Res* **44**: W272-276.
Lagman D, Sundstrom G, Ocampo Daza D, Abalo XM, Larhammar D. 2012.
Expansion of transducin subunit gene families in early vertebrate
tetraploidizations. *Genomics* **100**: 203-211.
Lamprecht MR, Sabatini DM, Carpenter AE. 2007. CellProfiler: free, versatile
software for automated biological image analysis. *Biotechniques* **42**: 71-
75.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R. 2009. The Sequence Alignment/Map format and SAMtools.
*Bioinformatics* **25**: 2078-2079.
Lieschke GJ, Currie PD. 2007. Animal models of human disease: zebrafish swim
into view. *Nat Rev Genet* **8**: 353-367.
Liu CC, Li L, Lam YW, Siu CW, Cheng SH. 2016. Improvement of surface ECG
recording in adult zebrafish reveals that the value of this model exceeds
our expectation. *Sci Rep* **6**: 25073.

Locke AE Kahali B Berndt SI Justice AE Pers TH Day FR Powell C Vedantam S Buchkovich ML Yang J et al. 2015. Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**: 197-206.

Louis A, Nguyen NT, Muffato M, Roest Crollius H. 2015. Genomicus update 2015: KaryoView and MatrixView provide a genome-wide perspective to multispecies comparative genomics. *Nucleic Acids Res* **43**: D682-689.

Ludwig A, Zong X, Stieber J, Hullin R, Hofmann F, Biel M. 1999. Two pacemaker channels from human heart with profoundly different activation kinetics. *EMBO J* **18**: 2323-2329.

MacRae CA, Peterson RT. 2015. Zebrafish as tools for drug discovery. *Nat Rev Drug Discov* **14**: 721-731.

Mark MD, Herlitze S. 2000. G-protein mediated gating of inward-rectifier K+ channels. *Eur J Biochem* **267**: 5830-5836.

Mawdsley DJ, Cooper HM, Hogan BM, Cody SH, Lieschke GJ, Heath JK. 2004. The Netrin receptor Neogenin is required for neural tube formation and somitogenesis in zebrafish. *Dev Biol* **269**: 302-315.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, Flicek P, Cunningham F. 2016. The Ensembl Variant Effect Predictor. *Genome Biol* **17**: 122.

Moreno-Mateos MA, Vejnar CE, Beaudoin JD, Fernandez JP, Mis EK, Khokha MK, Giraldez AJ. 2015. CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. *Nat Methods* **12**: 982-988.

Newton-Cheh C, Guo CY, Wang TJ, O'Donnell C J, Levy D, Larson MG. 2007. Genome-wide association study of electrocardiographic and heart rate variability traits: the Framingham Heart Study. *BMC Med Genet* **8 Suppl 1**: S7.

Ng MCY Graff M Lu Y Justice AE Mudgal P Liu CT Young K Yanek LR Feitosa MF Wojczynski MK et al. 2017. Discovery and fine-mapping of adiposity loci using high density imputation of genome-wide association studies in individuals of African ancestry: African Ancestry Anthropometry Genetics Consortium. *PLoS Genet* **13**: e1006719.

Niazi MKK, Nilsson MF, Danielsson BR, Bengtsson E. 2009. Fully Automatic Heart Beat Rate Determination in Digital Video Recordings of Rat Embryos. doi:http://urn.kb.se/resolve?urn=urn:nbn:se:uu:diva-88052.

Nolte IM Munoz ML Tragante V Amare AT Jansen R Vaez A von der Heyde B Avery CL Bis JC Dierckx B et al. 2017. Genetic loci associated with heart rate variability and their effects on cardiac disease risk. *Nature communications* **8**: 15805.

Pardo-Martin C, Allalou A, Medina J, Eimon PM, Wahlby C, Fatih Yanik M. 2013. High-throughput hyperdimensional vertebrate phenotyping. *Nature communications* **4**: 1467.

Pearson WR, Wood T, Zhang Z, Miller W. 1997. Comparison of DNA sequences with protein sequences. *Genomics* **46**: 24-36.

Poon KL, Brand T. 2013. The zebrafish model system in cardiovascular research: A tiny fish with mighty prospects. *Glob Cardiol Sci Pract* **2013**: 9-28.

Shi W, Wymore R, Yu H, Wu J, Wymore RT, Pan Z, Robinson RB, Dixon JE, McKinnon D, Cohen IS. 1999. Distribution and prevalence of hyperpolarization-activated cation channel (HCN) mRNA expression in cardiac tissues. *Circ Res* **85**: e1-6.

Stieber J, Herrmann S, Feil S, Loster J, Feil R, Biel M, Hofmann F, Ludwig A. 2003.
The hyperpolarization-activated channel HCN4 is required for the
generation of pacemaker action potentials in the embryonic heart. *Proc*
*Natl Acad Sci U S A* **100**: 15235-15240.
Stieber J, Wieland K, Stockl G, Ludwig A, Hofmann F. 2006. Bradycardic and
proarrhythmic properties of sinus node inhibitors. *Mol Pharmacol* **69**:
1328-1337.
Sudhof TC. 2002. Synaptotagmins: why so many? *J Biol Chem* **277**: 7629-7632.
Swedberg K, Komajda M, Bohm M, Borer JS, Ford I, Dubost-Brama A, Lerebours
G, Tavazzi L. 2010. Ivabradine and outcomes in chronic heart failure
(SHIFT): a randomised placebo-controlled study. *Lancet* **376**: 875-885.
Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A,
Doncheva NT, Roth A, Bork P et al. 2017. The STRING database in 2017:
quality-controlled protein-protein association networks, made broadly
accessible. *Nucleic Acids Res* **45**: D362-d368.
Thayer JF, Yamamoto SS, Brosschot JF. 2010. The relationship of autonomic
imbalance, heart rate variability and cardiovascular disease risk factors.
*Int J Cardiol* **141**: 122-131.
Varshney GK, Carrington B, Pei W, Bishop K, Chen Z, Fan C, Xu L, Jones M, LaFave
MC, Ledin J et al. 2016. A high-throughput functional genomics workflow
based on CRISPR/Cas9-mediated targeted mutagenesis in zebrafish. *Nat*
*Protoc* **11**: 2357-2375.
Varshney GK, Pei W, LaFave MC, Idol J, Xu L, Gallardo V, Carrington B, Bishop K,
Jones M, Li M et al. 2015. High-throughput gene targeting and
phenotyping in zebrafish using CRISPR/Cas9. *Genome Res* **25**: 1030-1042.
Verweij N, van de Vegte YJ, van der Harst P. 2018. Genetic study links
components of the autonomous nervous system to heart-rate profile
during exercise. *Nature communications* **9**: 898.
Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz M,
Grouios C, Kazi F, Lopes CT et al. 2010. The GeneMANIA prediction server:
biological network integration for gene prioritization and predicting gene
function. *Nucleic Acids Res* **38**: W214-220.
Whitesell TR, Kennedy RM, Carter AD, Rollins EL, Georgijevic S, Santoro MM,
Childs SJ. 2014. An alpha-smooth muscle actin (acta2/alphasma)
zebrafish transgenic line marking vascular mural cells and visceral
smooth muscle cells. *PLoS One* **9**: e90590.
Wilson NH, Key B. 2007. Neogenin: one receptor, many functions. *Int J Biochem*
*Cell Biol* **39**: 874-878.
Zhang J, Kobert K, Flouri T, Stamatakis A. 2014. PEAR: a fast and accurate
Illumina Paired-End reAd mergeR. *Bioinformatics* **30**: 614-620.

384 embryos imaged at 2dpf

Secondary trait analysis:
Cardiac traits (left), body size analysis (right)

Sinoatrial pauses (n=39) & arrests (n=36),
uncontrolled atrial contraction (n=1),
edema (n=15), fish moved during
recording/missed beats (n=38)

Embryos excluded due to
suboptimal segmentation
and/or presence of edema:
length (n=46), dorsal (n=56),
lateral (n=78), volume (n=79)

Primary analysis:
279 embryos included in HRV and
heart rate analysis at 2dpf

384 embryos imaged at 2dpf

Larvae excluded due to
technical issues* (n=41), death (n=3)
or larva not detected by VAST (n=14)

326 larvae imaged at 5dpf

Secondary trait analysis:
Cardiac traits (left), body size analysis (right)

Sinoatrial pauses (n=9) & arrests (n=3),
uncontrolled atrial contraction (n=9),
abnormal morphology/reduced contractility (n=6)
edema (n=14), missed beats (n=1)

Larvae excluded due to
suboptimal segmentation
and/or presence of edema:
length (n=84), dorsal (n=93),
lateral (n=70), volume (n=128)

Primary analysis:
291 larvae included in HRV and
heart rate analysis at 5dpf

**Supplementary Figure S1**: Flow chart showing the number of included larvae in the analysis and reasons for exclusions. Embryos/larvae can show more than one phenotype. *Technical issues refers to not being able to image due to microscope complications. VAST: Vertebrate automated screening technology.

**Supplemental Table 1: Overview of zebrafish orthologues and designed gRNAs**

| Zebrafish orthologue | ENSDARG | Conservation Zebrafish - Human | Conservation Zebrafish - Spotted gar | Main transcript | Target sequence gRNA | Exon target (transcript name) | CRISPRscan rank | ChopChop rank | Activity test rank | Predicted off-targets | Nr of mismatches | Genomic location of potential off-target 1 | ENSDARG (gene-ID) of potential off-target 1, Feature | ENSG (gene-ID) of potential off-target 1 | Genomic location of potential off-target 2 | ENSDARG (gene-ID) of potential off-target 2, Feature | ENSG (gene-ID) of potential off-target 2 | Primer forward | Primer reverse | Amplicon size |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| gngt1 | ENSDARG00000035798 | 17 | 21 | ENSDART00000051950 | gGACCTGGATAAGGCGAAGA | 2/3 (gngt1-001); 1/2 (gngt1-002) | 62 | 4/13 (yellow) | moderate | 1 | 3 | chr25:17488676 | ENSDARG00000051959 (zgc:66449), intron (3/6), intron (4/4) | no human orthologue | | | | GTCCATTTTCCAGAGTGGAGAC | gCCCATAACACAATTGCTTACC | 207 |
| syt10 | ENSDARG00000045750 | 0 | 1 | ENSDART00000136049 | GGCTGATGCCGTCCTCCTGTG | 1/7 (syt10-001); 1/7 (syt10-201) | 36 | 6/129 (green) | low | 0 | | | | | | | | TAGGATATGGACTTGGACGCAT | TTACCTGGTATGTTGCTCTCCA | 227 |
| rgs6 | ENSDARG00000015627 | 9 | 20 | ENSDART00000135513 | GGCCTCTCAGGACCAAGG | 1/17 (rgs6-201); 2/18 (rgs6-001) | 62 | 1/107 (green) | low | 0 | | | | | | | | AAGGCTAATTCATTGTCTGGGA | GCATGTCAAATGTAAAGGCAAA | 169 |
| hcn4 | ENSDARG00000061685 | 6 | 3 | ENSDART00000136140 | GGAGAGCCCTCCTGGAGCCG | 1/8 (hcn4-001) | 59 | 11/280 (green) | high | 2 | 3/3 | chr21:45844184 | ENSDARG00000103825 (galnt10), exon (1/3) | ENSG00000164574 (GALNT10) | chrUn_KN150614v1:3424 | ENSDARG00000100453 (galnt10), exon (6/8) | ENSG00000164574 (GALNT10) | CCTTTCGTCCATCACCAGTC | CCGCTTGATAATACCTCTCGTC | 230 |
| hcn4l | ENSDARG00000074419 | 3 | 3 | ENSDART00000088249 | gGCCTGGATAGTGTTTAACG | 2/8 (hcn4l-201); 1/7 (hcn4l-001) | / | 2/191 (green) | low | 0 | | | | | | | | TTACTGGGACCTGATCATGCTT | AAGATGTAATCCACAGGGATGG | 235 |
| neo1a | ENSDARG00000102855 | 1 | 5 | ENSDART00000158160 | GGAGCCGTCGGATACACTAG | 1/4 (neo1a-201); 2/29 (neo1a-201); 2/30 (neo1a-202) | 48 | 2/55 (green) | very high | 0 | | | | | | | | TTCTGTGTCTCCACAGGATCAG | CCTCATCGGGTTTATTGTGTTT | 215 |
| neo1b | ENSDARG00000075100 | 2 | 6 | ENSDART00000115280 | GGACAGAGATGCTCGGCCTG | 3/29 (neo1b-201) | 58 | 18/354 (green) | very high | 0 | | | | | | | | CAGCTCTCTCTCCATTGGCTT | TATGGTGCAACGGTAGAGTCC | 206 |
| quo (kiaa1755_1) | ENSDARG00000073684 | 2 | 5 | ENSDART00000109679 | GGAGATCCGGTTGCCCTGTG | 2/22 (quo-001); 3/23 (quo-201) | 83 | 12/331 (green) | very high | 0 | | | | | | | | ATGACCGTAAGTCAAGGGAAAA | TGAATCTTGTCTAATGGCGTTG | 153 |
| si:dkey-65j6.2 (kiaa1755_2) | ENSDARG00000103586 | 3 | 5 | ENSDART00000158832 | GGACGGTTTGGCTCCAGCAG | 1/23 (si:dkey-65j6.2-201); 3/25 (si:dkey-65j6.2-001) | 59 | 14/303 (green) | very high | 2 | 3/3 | chr15:32969618 | ENSDARG00000104664 (dclk1b), exon (3/16), exon (4/17) | ENSG00000133083 (DCLK1) | chr22:28819880 | ENSDARG00000071083 (si:dkeyp-34c12.1), Intron (7/7), Exon (8/9) | no human orthologue | AGCCATTACACCCAAAGAAAA | GGGAATCATTTCGAAGTAGCTG | 152 |

Table showing zebrafish genes, main transcript used for the analysis and gRNAs; rsID refers to the lead SNP(s) of the GWAS-identified loci (Nolte et. al, 2017), multiple independent SNPs per locus can be present; For simplicity, quo is referred to as *kiaa1755_1*, and si:dkey-65j6.2 as *kiaa1755_2*; Conservation refers to the number of genes that are preserved in the same locus across species according to Genomicus; A small "g" in the gRNA target sequence indicates a manually introduced nu...

**Supplemental Table 2: Unique Cigars**

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S119M | 498 | 336 | 381 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 207M | 517 | 267 | 372 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S69M1D3S46M | 544 | 69 | 405 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG ATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 151M6D50M | 500 | 63 | 364 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG ATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S63M6D50M | 584 | 13 | 479 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGATGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 158M1S48M | 504 | 4 | 221 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S68M3D48M | 606 | 3 | 119 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGTAACCCAGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S72M4I3S44M | 526 | 2 | 143 |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATAATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 45M1S41M1S32M1S36M1D3S46M | 48 | 1 | - |
| *gngt1* | ATCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 1S44M1S41M1S69M1D3S46M | 10 | 1 | - |
| *gngt1* | CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA | 87M1S119M | 3 | 1 | - |
| *syt10* | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACAGAGGACGGCATCAGCCTGTGCCAAA GGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 227M | 1024 | 518 | 644 |
| *syt10* | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACAAGAGGACGGCATCAGCCTGTGCCAA AGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCA C | 126M1I101M | 735 | 46 | 462 |
| *syt10* | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGACGGCATCAGCCTGTGCCAAAGGGCTCTC CAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 122M8D97M | 606 | 45 | 397 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTACAGCTGATGCGTACGGGAATGGAGAGGACG GCATCAGCCTGTGCCAAAGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAAT GCGCGGATATATTTCCAC | 120M1S9I2M1S2M8I101M | 793 | 43 | 405 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACATGGACGGCATCAGCCTGTGCCAAAG GGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 127M1S1D98M | 368 | 42 | 246 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACGGCATCAGCCTGTGCCAAAGGGCTCTCCAAATAGTGACAG AGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 109M21D97M | 667 | 19 | 267 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGGACGGCATCAGCCTGTGCCAAAGGGCTCT CCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 122M7D98M | 963 | 16 | 431 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACGGCATCAGCGGCATCAGCCTGTGCC AAAGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTC CAC | 126M1S1M3I2S97M | 937 | 11 | 645 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACGGCATCAGCCTGTGCCAAAGGGCTCT CCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 124M7D96M | 502 | 11 | 300 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCCTGTCAGCCTGTGCCAAAGGGCTCTCCAA ATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 124M1S10D2S90M | 524 | 4 | 229 |
| syt10 | CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGAGGACGGCATCAGCCTGTGCCAAAGGGCT CTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC | 122M5D100M | 1247 | 3 | 670 |
| rgs6 | GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCACCTTGGTCC TGAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGA ATGAGTCCCAGAACAGG | 169M | 1431 | 728 | 813 |
| rgs6 | GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCACTTGGTCCT GAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAA TGAGTCCCAGAACAGG | 68M1D100M | 784 | 24 | 453 |
| rgs6 | GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCATTGGTCCT GAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAA TGAGTCCCAGAACAGG | 68M2D99M | 540 | 7 | 195 |
| rgs6 | GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCATAGTCCTG AGAGGCCTGAGCCATGCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACA GAAAAGGAATGAGTCCCAGAACAGG | 68M3D1M1S12M10I84M | 920 | 4 | 287 |
| rgs6 | GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGTGGTCCTGAGAGGCCTG AGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAATGAGTCCCAG AACAGG | 59M12D98M | 1424 | 1 | - |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTC GAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGCTCCAGGAG GGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTG CC | 230M | 1380 | 638 | 789 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGCCGCCACCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 142M1S1M4I1S85M | 725 | 48 | 343 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCATCCGCGCCGCCACCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 136M1S5M7I1S87M | 1314 | 16 | 225 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGCTCTCCTCCAGGAGCTCCAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 143M5I7M6I80M | 799 | 15 | 383 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 135M8D87M | 917 | 13 | 563 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCCCTCCAGGAGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 141M1D1S87M | 987 | 11 | 377 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCACCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 139M1S1M4D85M | 1302 | 10 | 587 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGGCCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 143M3I87M | 694 | 2 | 49 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCCATCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC | 98M11I43M2S1D86M | 346 | 2 | 36 |
| hcn4 | GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGTGGACGTGCC | 142M3D85M | 222 | 1 | - |
| hcn4l | AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG | 115M1S89M1S29M | 1076 | 697 | 671 |
| hcn4l | AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACCAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGAATGATGATCAGGTTGCCCACCATTAGAAG | 235M | 765 | 38 | 389 |
| hcn4l | AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG | 115M1S34M2D53M1S29M | 585 | 18 | 295 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| hcn4l | AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTG TACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCAGGC GGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG | 115M1S28M16D45M1S29M | 523 | 9 | 176 |
| hcn4l | AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTG TACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGT TTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCC ACCATTAGAAG | 115M1S33M1I56M1S29M | 1218 | 4 | 915 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACTAGCGGTGCGAGGAGCGCCAGTTC TCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGC TTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 157M1S57M | 941 | 151 | 691 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTGCGAGGAGCGCCAGTTCTCGCCAGTTCTCCTCAA CTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGAT GAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 40M16D10M10I91M1S57M | 957 | 95 | 545 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACGGTGCGAGGAGCGCCAGTTCTCCTC AACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAG ATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 50M4D103M1S57M | 899 | 82 | 475 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTC CACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGA CAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 40M16D101M1S57M | 696 | 72 | 413 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACGAGGAGCGCCAGTTCTCCTCAACTG CTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAG CGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 50M9D98M1S57M | 589 | 72 | 305 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTC AGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCG CAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 42M12D103M1S57M | 703 | 66 | 359 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTG CCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATG GCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 38M31D88M1S57M | 985 | 57 | 521 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACAGCGGTGCGAGGAGCGCCAGTTCTCC TCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTC AGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 49M3D105M1S57M | 960 | 20 | 653 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACGAGGAGCGCCAGTTCTCCTCAACTGCT CAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGC GCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 48M11D98M1S57M | 633 | 20 | 415 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATAGCGGTGCGAGGAGCGCCAGTTCTCCTC AACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAG ATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 46M5D106M1S57M | 1026 | 19 | 474 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATTCTCCTCAACTGCTCAGTCCACAGCGAGT CTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGG CCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 46M26D85M1S57M | 672 | 16 | 404 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACATCCATTCAGCGGTGCGA GGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTT TCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTC C | 11M9I39M5I2S105M1S57M | 688 | 14 | 336 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTC CTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCG ATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 34M28D95M1S57M | 807 | 11 | 252 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACAAAGCGAGGAGCGCCAGTTCTCCTCA ACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGA TGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 50M6D2S99M1S57M | 569 | 10 | 314 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTG CCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATG GCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 40M31D86M1S57M | 1179 | 9 | 312 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACCAGCGGTGCGAGGAGCGCCAGTTC TCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGC TTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 51M1S105M1S57M | 1569 | 6 | 1406 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACCAGCGCCAGTTCTCCTCAACTGCTC AGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCG CAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 51M1S12D93M1S57M | 1010 | 6 | 552 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACGGTGCGGAGCGGTGCGAGGAGCG CCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCA GCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 51M6I1S105M1S57M | 760 | 6 | 543 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGAGCGGTGCGAGGAGCGCCAGTTCTCCTCAA CTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGAT GAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 45M7D105M1S57M | 1451 | 5 | 663 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATAGTCGGATAGTTCTCCTCAACTGCTCAGT CCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAG ACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 48M2S1M11D3M4D1S87M1S57M | 754 | 5 | 361 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACTAGCGGTGCGAGGAGCGCCAGTTC TCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGC TTCAGACGAACGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 160M1S54M | 734 | 4 | 441 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCCTGGCGAGGAGCGCCAGTTCTCCTCAACTGCTC AGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCG CAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 43M7D2M1S5D99M1S57M | 376 | 2 | 40 |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATGTAGCGGTGCGAGGAGTGTAGCGGTGCG AGGAGCGCCAGTTCTCCTCAACTTCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCT TTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACT CC | 47M2S1M14I1S33M1S72M1S57M | 1242 | 1 | - |
| neo1a | CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACTAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCC ACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGAC AGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC | 31M17D109M1S57M | 698 | 1 | - |
| neo1b | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCGTTCGCAGCGGTGGGACTGAGATGCTGAGCT GTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTGTGGAGCTGGACAGCAGGGTCATT CAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 123M6D64M1S12M | 733 | 186 | 617 |
| neo1b | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCGTTCGCAGCGGTGGGACTGAGATGCTGAGCT GTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCTGTGGAGCTGGACAGCAGG GTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 128M1D77M | 632 | 84 | 483 |
| neo1b | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCGTTCGCAGCGGTGGGACTGAGATGCTGAGCT GTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCTGTGGAGCTGGACAGCAGG CAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 129M4I77M | 991 | 77 | 818 |
| neo1b | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCGTTCGCAGCGGTGGGACTGAGATGCTGAGCT GTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCTGGACAGCAGGGTCATT CAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 129M6D71M | 700 | 75 | 505 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 121M9D63M1S12M | 739 | 33 | 686 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 193M1S12M | 231 | 28 | 194 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 123M6D77M | 970 | 27 | 739 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCAGGGTCTGGGTCTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 129M8I3M4I74M | 915 | 24 | 914 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 206M | 564 | 21 | 525 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 125M4D77M | 689 | 18 | 469 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 126M7D60M1S12M | 503 | 11 | 266 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 125M17D51M1S12M | 960 | 10 | 710 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGGGACATTCAGGACATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 120M19D5M4S2M1S55M | 801 | 10 | 567 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 117M22D54M1S12M | 1170 | 3 | 685 |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCACTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC | 108M1S84M1S12M | 6 | 1 | - |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 129M2I77M | 5 | 1 | - |
| *neo1b* | TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCACTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC | 108M1S97M | 3 | 1 | - |
| *kiaa1755_1* | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S90M | 997 | 119 | 507 |
| *kiaa1755_1* | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S8M14D68M | 1117 | 99 | 569 |
| *kiaa1755_1* | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S15M7D68M | 1175 | 80 | 598 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACTTGTAGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S20M11I12M1S41M3I2S56M1S13M | 882 | 68 | 903 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S38M10D51M1S13M | 541 | 57 | 417 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S42M3I57M1S13M | 780 | 55 | 596 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S17M17D56M | 1077 | 38 | 490 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S18M1S1D70M | 876 | 30 | 459 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S99M1S13M | 775 | 30 | 588 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S42M1S1I1S55M1S13M | 822 | 22 | 547 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCACAAGTCATTCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S11M3D5M1S1M1S2M5D1S60M | 1190 | 21 | 408 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCATGACGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S17M1S1I1M1S70M | 1469 | 20 | 683 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S17M5D68M | 1365 | 18 | 561 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S18M1I72M | 1602 | 14 | 834 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATAATGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S15M81D1M1S34M1S13M | 1070 | 14 | 568 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S14M7D69M | 1063 | 14 | 794 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACTAACAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S4M1S1M15D1S68M | 876 | 12 | 386 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCAACCGGACTCAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S17M8I73M | 950 | 10 | 426 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCAACCGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGA | 6M1S32M1S40M3S56M1S13M | 618 | 9 | 283 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S41M1I58M1S13M | 581 | 6 | 250 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S42M1S1I1S69M | 145 | 6 | 23 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACTTGTAGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S20M11I12M1S41M3I2S70M | 136 | 6 | 89 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA | 6M1S32M1S41M4D54M1S13M | 1101 | 5 | 323 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S113M | 120 | 4 | 48 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S42M3I71M | 107 | 2 | 65 |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCGGCAAGTCATTCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S32M1S22M1S17M4I2M1S1M1S2M2S64M | 1184 | 1 | - |
| kiaa1755_1 | CGCAGAGTCTCCTGACCAACATAATGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 6M1S15M81D1M1S48M | 233 | 1 | - |
| kiaa1755_1 | CGCAGACTCTCCTGACCAACATATGACCCAACTTATCACAAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA | 153M | 25 | 1 | - |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S22M21D45M1S2M1S11M | 985 | 120 | 638 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGCTTGGCTCCAGCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S88M1S2M1S11M | 951 | 78 | 613 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S19M21D48M1S2M1S11M | 886 | 63 | 460 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S22M29D37M1S2M1S11M | 1070 | 62 | 464 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 82M3D67M | 772 | 62 | 333 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 78M15D59M | 856 | 59 | 414 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCATCAGAGAAACACAGAGAAATGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 81M1D2M1S3M1S1M11I3S59M | 829 | 55 | 457 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 81M16D55M | 935 | 35 | 587 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 76M12D64M | 837 | 33 | 470 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 152M | 564 | 32 | 351 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S22M14D52M1S2M1S11M | 964 | 26 | 370 |
| kiaa1755_2 | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S33M3D52M1S2M1S11M | 793 | 20 | 426 |

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGGACCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S25M1S8D1M1S52M1S2M1S11M | 787 | 19 | 595 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S26M21D41M1S2M1S11M | 1035 | 18 | 370 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S29M11D48M1S2M1S11M | 657 | 17 | 443 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S23M17D48M1S2M1S11M | 1152 | 15 | 331 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTAAAAATAAGGCTAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 77M8I4M5D66M | 584 | 11 | 214 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 71M21D60M | 1110 | 9 | 532 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGCACATCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 74M5S1M5D67M | 970 | 9 | 382 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 73M31D48M | 976 | 4 | 130 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGATGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 73M1S1M12D65M | 726 | 4 | 358 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTATGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAGAGGACAGAAG | 48M1S26M1S1M16D44M1S2M1S11M | 1411 | 3 | 177 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 68M21D63M | 913 | 3 | 550 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTATGGTCGACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG | 75M12D1S6M2D56M | 834 | 3 | 128 |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGTCGAACAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S30M1D2M1S1M1S52M1S2M1S11M | 562 | 1 | - |
| *kiaa1755_2* | TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG | 48M1S21M26D41M1S2M1S11M | 558 | 1 | - |
| *dclk1b* | ATGTGTGTGGATCAATTGAGCCTTACAAGATGCAGGAATATACTAAAAACGTCAACCCTAACTGGTCGGTCAACATGAGAACCACTGCCGGAGCAAAACCGCCCACCTCTCTGGCCAGTGCCAAGGGCGGGTCTATAGAAACCAAGGATAACAGGGATTTCATCCGACCCAAACTGGTGACCATAATCCGTAGCGGGGTAAAGCCTCGGAAAGCCATCCGAATCCTTCTCAACAA | 235M | 1638 | 377 | 1281 |
| *dclk1b* | ATGTGTGTGGATCAATTGAGCCTTACAAGATGCAGGAATATACTAAAAACGTCAACCCTAACTGGTCGGTCAACATGAGAACCACTGCCGGAGCAAAACCGCCCACCTCTCTGGCCAGTGCCAAGGGCGGGTCTATAGAAACCAAGGATAACAGGGATTTCATCCGACCCAAACTGGTGACCATAATCCGTAGTGGGGTAAAGCCTCGGAAAGCCATCCGAATCCTTCTCAACAA | 193M1S41M | 1078 | 102 | 877 |
| *galnt10chr21* | GCGTAACGTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCGATCCGCCTGGAACCATGCCTGAGGAGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACACTGCTAACTGCTAATA | 174M | 1246 | 198 | 994 |
| *galnt10chr21* | GCGTAACGTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCGATCCGCCTGGAACCATGCCTGAGGAGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACACTGCTAACTGCTAATA | 48M1S125M | 952 | 364 | 1044 |
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCGATCCGCCTGGAACCATGCCTGAGGAGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACACACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGT | 41M1S135M12D27M | 272 | 326 | 207 |

*Continued Supplemental Table 2*

| Orthologue | Sequence | Cigar | Mean Nr reads | Nr alleles | SD reads |
|---|---|---|---|---|---|
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT | 216M | 148 | 187 | 106 |
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGT | 41M1S135M14D25M | 157 | 136 | 94 |
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGT | 41M1S135M8D31M | 147 | 82 | 107 |
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT | 214M2N | 136 | 13 | 127 |
| *galnt10scaff* | GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGT | 41M1S135M10D29M | 4 | 2 | 1 |

All unique sequences identified for targeted orthologues and their respective alignment report string (Cigar). Mean number of reads, the number of alleles the cigar was found in and the standard deviation (SD) are reported. Cigar letters: M=match, D=deletion, I=insertion, S=substitution.

**Supplemental Table 3: Unique variants and affected transcripts**

| Orthologue | Chr | ENSDARG | Affected transcripts | Start (bp) | End (bp) | Mutation | Annotation | Base pair change | VEP impact | Nr affected alleles |
|---|---|---|---|---|---|---|---|---|---|---|
| *gngt1* | 19 | ENSDARG00000035798 | ENSDART00000051950 ENSDART00000139083 | 40856888 | 40856893 | AAGGCG/- | inframe_deletion | -6 | MODERATE | 75 |
| | | | | 40856893 | 40856895 | GAA/- | inframe_deletion | -3 | MODERATE | 3 |
| | | | | 40856894 | 40856894 | A/- | frameshift_variant | -1 | HIGH | 69 |
| | | | | 40856895 | 40856895 | A/T | missense_variant | 0 | MODERATE | 3 |
| | | | | 40856895 | 40856897 | AGA/TTT | missense_variant | 0 | MODERATE | 69 |
| | | | | 40856897 | 40856896 | -/TAAC | stop_gained,frameshift_variant | 4 | HIGH | 2 |
| | | | | 40856897 | 40856899 | ATG/CCA | missense_variant | 0 | MODERATE | 2 |
| *syt10* | 4 | ENSDARG00000045750 | ENSDART00000136049 | 10888933 | 10888953 | GACATGAATGTCCGCACAGAG/- | inframe_deletion | -21 | MODERATE | 19 |
| | | | | 10888944 | 10888944 | C/A | synonymous_variant | 0 | LOW | 42 |
| | | | | 10888945 | 10888944 | -/CAGCTGATG | inframe_insertion | 9 | MODERATE | 42 |
| | | | | 10888946 | 10888950 | GCACA/- | frameshift_variant | -5 | HIGH | 3 |
| | | | | 10888946 | 10888952 | GCACAGA/- | frameshift_variant | -7 | HIGH | 15 |
| | | | | 10888946 | 10888953 | GCACAGAG/- | frameshift_variant | -8 | HIGH | 40 |
| | | | | 10888947 | 10888947 | C/T | synonymous_variant | 0 | LOW | 42 |
| | | | | 10888948 | 10888948 | A/C | missense_variant | 0 | MODERATE | 4 |
| | | | | 10888948 | 10888954 | ACAGAGG/- | frameshift_variant | -7 | HIGH | 11 |
| | | | | 10888949 | 10888958 | CAGAGGACGG/- | frameshift_variant | -10 | HIGH | 4 |
| | | | | 10888950 | 10888949 | -/GGGAATGG | frameshift_variant | 8 | HIGH | 42 |
| | | | | 10888950 | 10888949 | -/A | frameshift_variant | 1 | HIGH | 44 |
| | | | | 10888950 | 10888950 | A/G | synonymous_variant | 0 | LOW | 11 |
| | | | | 10888951 | 10888951 | G/T | stop_gained | 0 | HIGH | 42 |
| | | | | 10888952 | 10888951 | -/CAT | stop_gained,protein_altering_variant | 3 | HIGH | 11 |
| | | | | 10888952 | 10888952 | A/- | frameshift_variant | -1 | HIGH | 42 |
| | | | | 10888952 | 10888953 | AG/CA | missense_variant | 0 | MODERATE | 11 |
| | | | | 10888959 | 10888960 | CA/TG | missense_variant | 0 | MODERATE | 4 |
| *rgs6* | 20 | ENSDARG00000015627 | ENSDART00000135513 ENSDART00000184779 | 28638811 | 28638822 | CCCACCGCACCT/- | inframe_deletion | -12 | MODERATE | 1 |
| | | | | 28638820 | 28638820 | C/- | frameshift_variant | -1 | HIGH | 24 |
| | | | | 28638820 | 28638821 | CC/- | frameshift_variant | -2 | HIGH | 7 |
| | | | | 28638820 | 28638822 | CCT/- | inframe_deletion | -3 | MODERATE | 4 |
| | | | | 28638824 | 28638824 | G/A | stop_gained | 0 | HIGH | 4 |
| | | | | 28638837 | 28638836 | -/CTGAGCCATG | frameshift_variant | 10 | HIGH | 4 |
| *hcn4* | 18 | ENSDARG00000061685 | ENSDART00000136140 ENSDART00000189186 | 1143637 | 1143644 | ACCCGCGG/- | frameshift_variant | -8 | HIGH | 13 |
| | | | | 1143638 | 1143638 | C/T | missense_variant | 0 | MODERATE | 16 |
| | | | | 1143641 | 1143641 | G/A | missense_variant | 0 | MODERATE | 9 |
| | | | | 1143643 | 1143643 | G/- | frameshift_variant | -1 | HIGH | 11 |
| | | | | 1143643 | 1143644 | GG/CA | missense_variant | 0 | MODERATE | 2 |
| | | | | 1143643 | 1143646 | GGCT/- | frameshift_variant | -4 | HIGH | 9 |
| | | | | 1143644 | 1143643 | -/CCGCCAC | frameshift_variant | 7 | HIGH | 16 |
| | | | | 1143644 | 1143644 | G/C | missense_variant | 0 | MODERATE | 75 |
| | | | | 1143644 | 1143646 | GCT/- | inframe_deletion | -3 | MODERATE | 1 |
| | | | | 1143645 | 1143644 | -/CTCTC | frameshift_variant | 5 | HIGH | 15 |
| | | | | 1143645 | 1143644 | -/GCC | protein_altering_variant | 3 | MODERATE | 2 |
| | | | | 1143645 | 1143645 | C/- | frameshift_variant | -1 | HIGH | 2 |
| | | | | 1143646 | 1143645 | -/GCCA | frameshift_variant | 4 | HIGH | 48 |
| | | | | 1143646 | 1143646 | T/C | synonymous_variant | 0 | LOW | 48 |
| | | | | 1143652 | 1143651 | -/AGCTCC | inframe_insertion | 6 | MODERATE | 15 |

| Orthologue | Chr | ENSDARG | Affected transcripts | Start (bp) | End (bp) | Mutation | Annotation | Base pair change | VEP impact | Nr affected alleles |
|---|---|---|---|---|---|---|---|---|---|---|
| *hcn4l* | 25 | ENSDARG00000074419 | ENSDART00000088249 ENSDART00000148940 | 29273086 | 29273101 | CCACGTTAAACACTAT/- | frameshift_variant | -16 | HIGH | 9 |
| | | | | 29273091 | 29273090 | -/T | frameshift_variant | 1 | HIGH | 4 |
| | | | | 29273092 | 29273093 | TA/- | frameshift_variant | -2 | HIGH | 18 |
| *neo1a* | 7 | ENSDARG00000102855 | ENSDART00000158160 ENSDART00000163261 ENSDART00000164768 ENSDART00000181629 | 53879228 | 53879244 | CACGGAGCCGTCGGATA/- | frameshift_variant | -17 | HIGH | 1 |
| | | | | 53879231 | 53879258 | GGAGCCGTCGGATACACTAGCGGTGCGA/- | frameshift_variant | -28 | HIGH | 11 |
| | | | | 53879235 | 53879265 | CCGTCGGATACACTAGCGGTGCGAGGAGCGC/- | frameshift_variant | -31 | HIGH | 57 |
| | | | | 53879237 | 53879252 | GTCGGATACACTAGCG/- | frameshift_variant | -16 | HIGH | 157 |
| | | | | 53879237 | 53879267 | GTCGGATACACTAGCGGTGCGAGGAGCGCCA/- | frameshift_variant | -31 | HIGH | 9 |
| | | | | 53879239 | 53879250 | CGGATACACTAG/- | inframe_deletion | -12 | MODERATE | 66 |
| | | | | 53879240 | 53879246 | GGATACA/- | frameshift_variant | -7 | HIGH | 2 |
| | | | | 53879242 | 53879248 | ATACACT/- | frameshift_variant | -7 | HIGH | 5 |
| | | | | 53879243 | 53879247 | TACAC/- | frameshift_variant | -5 | HIGH | 17 |
| | | | | 53879243 | 53879268 | TACACTAGCGGTGCGAGGAGCGCCAG/- | frameshift_variant | -26 | HIGH | 16 |
| | | | | 53879244 | 53879245 | AC/GT | missense_variant | 0 | MODERATE | 1 |
| | | | | 53879245 | 53879246 | CA/GT | missense_variant | 0 | MODERATE | 5 |
| | | | | 53879245 | 53879255 | CACTAGCGGTG/- | frameshift_variant | -11 | HIGH | 20 |
| | | | | 53879246 | 53879248 | ACT/- | inframe_deletion | -3 | MODERATE | 20 |
| | | | | 53879247 | 53879246 | -/GCGGTGCGAGGAGT | frameshift_variant | 14 | HIGH | 1 |
| | | | | 53879247 | 53879246 | -/TCCAT | frameshift_variant | 5 | HIGH | 14 |
| | | | | 53879247 | 53879247 | C/G | missense_variant | 0 | MODERATE | 1 |
| | | | | 53879247 | 53879248 | CT/TC | missense_variant | 0 | MODERATE | 14 |
| | | | | 53879247 | 53879250 | CTAG/- | frameshift_variant | -4 | HIGH | 82 |
| | | | | 53879247 | 53879252 | CTAGCG/- | inframe_deletion | -6 | MODERATE | 10 |
| | | | | 53879247 | 53879255 | CTAGCGGTG/- | inframe_deletion | -9 | MODERATE | 69 |
| | | | | 53879248 | 53879247 | -/GGTGCG | protein_altering_variant | 6 | MODERATE | 6 |
| | | | | 53879248 | 53879248 | T/G | missense_variant | 0 | MODERATE | 6 |
| | | | | 53879248 | 53879248 | T/C | missense_variant | 0 | MODERATE | 10 |
| | | | | 53879248 | 53879258 | TAGCGGTGCGA/- | frameshift_variant | -11 | HIGH | 5 |
| | | | | 53879249 | 53879249 | A/G | synonymous_variant | 0 | LOW | 2 |
| | | | | 53879249 | 53879260 | AGCGGTGCGAGG/- | inframe_deletion | -12 | MODERATE | 5 |
| | | | | 53879250 | 53879254 | GCGGT/- | frameshift_variant | -5 | HIGH | 2 |
| | | | | 53879253 | 53879254 | GT/AA | missense_variant | 0 | MODERATE | 10 |
| | | | | 53879262 | 53879265 | GCGC/- | frameshift_variant | -4 | HIGH | 5 |
| | | | | 53879263 | 53879262 | -/CGCCAGTTCT | frameshift_variant | 10 | HIGH | 89 |
| | | | | 53879266 | 53879266 | C/T | missense_variant | 0 | MODERATE | 5 |
| *neo1b* | 25 | ENSDARG00000075100 | ENSDART00000115280 | 2899698 | 2899698 | G/A | missense_variant | 0 | MODERATE | 2 |
| | | | | 2899707 | 2899728 | GAGATGCTCGGCCTGTGGAGCT/- | frameshift_variant | -22 | HIGH | 2 |
| | | | | 2899710 | 2899728 | ATGCTCGGCCTGTGGAGCT/- | frameshift_variant | -19 | HIGH | 7 |
| | | | | 2899711 | 2899719 | TGCTCGGCC/- | inframe_deletion | -9 | MODERATE | 19 |
| | | | | 2899713 | 2899718 | CTCGGC/- | inframe_deletion | -6 | MODERATE | 120 |
| | | | | 2899715 | 2899718 | CGGC/- | frameshift_variant | -4 | HIGH | 13 |
| | | | | 2899715 | 2899731 | CGGCCTGTGGAGCTGGA/- | frameshift_variant | -17 | HIGH | 6 |
| | | | | 2899716 | 2899722 | GGCCTGT/- | frameshift_variant | -7 | HIGH | 7 |
| | | | | 2899718 | 2899718 | C/- | frameshift_variant | -1 | HIGH | 58 |
| | | | | 2899719 | 2899718 | -/AGCAGGGT | frameshift_variant | 8 | HIGH | 15 |
| | | | | 2899719 | 2899718 | -/TCGG | frameshift_variant | 4 | HIGH | 50 |
| | | | | 2899719 | 2899718 | -/AG | frameshift_variant | 2 | HIGH | 1 |
| | | | | 2899719 | 2899724 | CTGTGG/- | inframe_deletion | -6 | MODERATE | 50 |
| | | | | 2899722 | 2899721 | -/GGTC | frameshift_variant | 4 | HIGH | 15 |
| | | | | 2899734 | 2899737 | GCAG/TTCA | missense_variant | 0 | MODERATE | 7 |
| | | | | 2899740 | 2899740 | T/A | missense_variant | 0 | MODERATE | 7 |

| Orthologue | Chr | ENSDARG | Affected transcripts | Start (bp) | End (bp) | Mutation | Annotation | Base pair change | VEP impact | Nr affected alleles |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | 2051654 | 2051734 | ATGACCCAACTTATCACAAGCATGGTCCAAGCAAGACAGCATCAC AAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAG/- | inframe_deletion | -81 | MODERATE | 13 |
| | | | | 2051699 | 2051699 | A/T | stop_gained | 0 | HIGH | 10 |
| | | | | 2051701 | 2051715 | GTCATTCCTCACAGG/- | inframe_deletion | -15 | MODERATE | 10 |
| | | | | 2051703 | 2051716 | CATTCCTCACAGGG/- | frameshift_variant | -14 | HIGH | 61 |
| | | | | 2051706 | 2051708 | TCC/- | inframe_deletion | -3 | MODERATE | 17 |
| | | | | 2051709 | 2051715 | TCACAGG/- | frameshift_variant | -7 | HIGH | 9 |
| | | | | 2051710 | 2051716 | CACAGGG/- | frameshift_variant | -7 | HIGH | 53 |
| | | | | 2051710 | 2051719 | CACAGGGCAA/- | frameshift_variant | -10 | HIGH | 51 |
| | | | | 2051712 | 2051711 | -/CCGG | frameshift_variant | 4 | HIGH | 1 |
| | | | | 2051712 | 2051711 | -/ACCGGACT | frameshift_variant | 8 | HIGH | 7 |
| | | | | 2051712 | 2051712 | C/T | missense_variant | 0 | MODERATE | 15 |
| | | | | 2051712 | 2051714 | CAG/ACC | missense_variant | 0 | MODERATE | 9 |
| | | | | 2051712 | 2051716 | CAGGG/- | frameshift_variant | -5 | HIGH | 14 |
| | | | ENSDART00000109679 | 2051712 | 2051728 | CAGGGCAACCGGATCTC/- | frameshift_variant | -17 | HIGH | 23 |
| | | | ENSDART00000153568 | 2051713 | 2051712 | -/G | frameshift_variant | 1 | HIGH | 15 |
| *kiaa1755_1* | 6 | ENSDARG00000073684 | ENSDART00000187502 | 2051713 | 2051712 | -/A | frameshift_variant | 1 | HIGH | 19 |
| | | | ENSDART00000187544 | 2051713 | 2051712 | -/TTG | stop_gained,inframe_insertion | 3 | HIGH | 70 |
| | | | ENSDART00000191165 | 2051713 | 2051713 | A/C | synonymous_variant | 0 | LOW | 22 |
| | | | | 2051713 | 2051714 | AG/TA | missense_variant | 0 | MODERATE | 70 |
| | | | | 2051713 | 2051716 | AGGG/- | frameshift_variant | -4 | HIGH | 4 |
| | | | | 2051714 | 2051713 | -/ACC | inframe_insertion | 3 | MODERATE | 51 |
| | | | | 2051714 | 2051714 | G/- | frameshift_variant | -1 | HIGH | 22 |
| | | | | 2051714 | 2051714 | G/A | missense_variant | 0 | MODERATE | 40 |
| | | | | 2051714 | 2051714 | G/C | missense_variant | 0 | MODERATE | 15 |
| | | | | 2051715 | 2051714 | -/C | frameshift_variant | 1 | HIGH | 22 |
| | | | | 2051715 | 2051715 | G/C | missense_variant | 0 | MODERATE | 22 |
| | | | | 2051716 | 2051716 | G/T | synonymous_variant | 0 | LOW | 18 |
| | | | | 2051716 | 2051716 | G/A | synonymous_variant | 0 | LOW | 10 |
| | | | | 2051719 | 2051720 | AC/TT | missense_variant | 0 | MODERATE | 1 |
| | | | | 2051719 | 2051723 | ACCGG/- | frameshift_variant | -5 | HIGH | 17 |
| | | | | 2051724 | 2051724 | A/T | missense_variant | 0 | MODERATE | 17 |
| | | | | 2051736 | 2051736 | C/A | missense_variant | 0 | MODERATE | 13 |

| Orthologue | Chr | ENSDARG | Affected transcripts | Start (bp) | End (bp) | Mutation | Annotation | Base pair change | VEP impact | Nr affected alleles |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | 43213133 | 43213153 | GGACGGTTTGGCTCCAGCAGT/- | inframe_deletion | -21 | MODERATE | 62 |
| | | | | 43213135 | 43213160 | ACGGTTTGGCTCCAGCAGTGGTCGGC/- | frameshift_variant | -26 | HIGH | 1 |
| | | | | 43213136 | 43213149 | CGGTTTGGCTCCAG/- | frameshift_variant | -14 | HIGH | 21 |
| | | | | 43213136 | 43213156 | CGGTTTGGCTCCAGCAGTGGT/- | inframe_deletion | -21 | MODERATE | 101 |
| | | | | 43213136 | 43213164 | CGGTTTGGCTCCAGCAGTGGTCGGCACAT/- | frameshift_variant | -29 | HIGH | 50 |
| | | | | 43213137 | 43213153 | GGTTTGGCTCCAGCAGT/- | frameshift_variant | -17 | HIGH | 15 |
| | | | | 43213138 | 43213138 | G/A | missense_variant | 0 | MODERATE | 4 |
| | | | | 43213138 | 43213168 | GTTTGGCTCCAGCAGTGGTCGGCACATCAGA/- | frameshift_variant | -31 | HIGH | 3 |
| | | | | 43213139 | 43213139 | T/G | synonymous_variant | 0 | LOW | 18 |
| | | | | 43213139 | 43213143 | TTTGG/CACAT | missense_variant | 0 | MODERATE | 9 |
| | | | | 43213140 | 43213140 | T/A | missense_variant | 0 | MODERATE | 3 |
| | | | | 43213140 | 43213147 | TTGGCTCC/- | frameshift_variant | -8 | HIGH | 18 |
| | | | | 43213140 | 43213151 | TTGGCTCCAGCA/- | inframe_deletion | -12 | MODERATE | 7 |
| | | | | 43213140 | 43213160 | TTGGCTCCAGCAGTGGTCGGC/- | inframe_deletion | -21 | MODERATE | 12 |
| | | | | 43213141 | 43213152 | TGGCTCCAGCAG/- | inframe_deletion | -12 | MODERATE | 32 |
| | | | ENSDART00000158832 | 43213142 | 43213141 | -/AAAAATAA | frameshift_variant | 8 | HIGH | 7 |
| | | | ENSDART00000164262 | 43213142 | 43213157 | GGCTCCAGCAGTGGTC/- | frameshift_variant | -16 | HIGH | 3 |
| *kiaa1755_2* | 23 | ENSDARG00000103586 | ENSDART00000186065 | 43213143 | 43213153 | GCTCCAGCAGT/- | frameshift_variant | -11 | HIGH | 13 |
| | | | ENSDART00000190315 | 43213143 | 43213157 | GCTCCAGCAGTGGTC/- | inframe_deletion | -15 | MODERATE | 57 |
| | | | ENSDART00000193300 | 43213144 | 43213144 | C/- | frameshift_variant | -1 | HIGH | 1 |
| | | | | 43213145 | 43213149 | TCCAG/- | frameshift_variant | -5 | HIGH | 9 |
| | | | | 43213146 | 43213146 | C/- | frameshift_variant | -1 | HIGH | 41 |
| | | | | 43213146 | 43213150 | CCAGC/- | frameshift_variant | -5 | HIGH | 7 |
| | | | | 43213146 | 43213161 | CCAGCAGTGGTCGGCA/- | frameshift_variant | -16 | HIGH | 34 |
| | | | | 43213147 | 43213147 | C/G | missense_variant | 0 | MODERATE | 1 |
| | | | | 43213147 | 43213149 | CAG/- | inframe_deletion | -3 | MODERATE | 69 |
| | | | | 43213149 | 43213149 | G/T | missense_variant | 0 | MODERATE | 41 |
| | | | | 43213149 | 43213149 | G/C | missense_variant | 0 | MODERATE | 18 |
| | | | | 43213149 | 43213149 | G/A | missense_variant | 0 | MODERATE | 1 |
| | | | | 43213152 | 43213152 | G/A | missense_variant | 0 | MODERATE | 3 |
| | | | | 43213153 | 43213153 | T/A | missense_variant | 0 | MODERATE | 41 |
| | | | | 43213155 | 43213154 | -/AAACACAGAGA | frameshift_variant | 11 | HIGH | 41 |
| | | | | 43213155 | 43213157 | GTC/AAT | missense_variant | 0 | MODERATE | 41 |
| | | | | 43213159 | 43213160 | GC/- | frameshift_variant | -2 | HIGH | 3 |

The predicted impact of unique CRISPR-Cas9 induced mutations on protein function was assessed using Ensembl's Variant Effect Predictor (VEP). An allele- and target-specific dosage score was subsequently calculated for each larva by weighting the mutation with the highest predicted impact by a factor 0.33, 0.66, or 1 for mutations with a low, moderate or high predicted impact, respectively. Modification of expression was only observed for transcripts of genes located near the targeted gene. Start and end coordinates are based on GRCz11. For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 4: Mutant allele frequencies**

| Gene | Nr mutated alleles | | | Missing | Sample call rate | Mutant allele freq | Expected nr mutated alleles | | | $P_{HWE}$ |
|---|---|---|---|---|---|---|---|---|---|---|
| | 0 | 1 | 2 | | | | 0 | 1 | 2 | |
| gngt1 | 234 | 132 | 12 | 4 | 0.990 | 0.206 | 238 | 124 | 16 | 2.70E-01 |
| syt10 | 193 | 131 | 54 | 4 | 0.990 | 0.316 | 177 | 163 | 38 | 2.10E-04 |
| rgs6 | 346 | 34 | 1 | 1 | 0.997 | 0.047 | 346 | 34 | 1 | 5.79E-01 |
| hcn4 | 268 | 100 | 9 | 5 | 0.987 | 0.156 | 268 | 100 | 9 | 4.69E-01 |
| hcn4l | 351 | 29 | 1 | 1 | 0.997 | 0.041 | 351 | 30 | 1 | - |
| neo1a | 20 | 113 | 241 | 8 | 0.979 | 0.795 | 16 | 122 | 237 | 1.16E-01 |
| neo1b | 15 | 19 | 271 | 77 | 0.798 | 0.920 | 2 | 45 | 258 | 2.12E-13 |
| kiaa1755_1 | 55 | 42 | 283 | 2 | 0.995 | 0.800 | 15 | 122 | 243 | 3.33E-32 |
| kiaa1755_2 | 24 | 60 | 295 | 3 | 0.992 | 0.858 | 8 | 93 | 279 | 5.65E-10 |

Number of larvae with 0, 1 and 2 mutated alleles, as well as the number of larvae with a missing call. *P*-value for a Hardy-Weinberg equilibrium (HWE) exact test, considering a ±30 bp window around the CRISPR targeted site as a single locus (*P*<2.9E-3 is significant after Bonferroni correction). Larvae with more than two missing genotypes were excluded from the statistical analysis (i.e. n_total=382). For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 5: Transcript-specific association analysis for dichotomous cardiac outcomes**

| | Sinoatrial pause | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | 2dpf (n=323, 39 cases) | | | | 5dpf (n=318, 9 cases) | | | |
| Gene | OR | LCI | UCI | P | OR | LCI | UCI | P |
| *gngt1* | 1.581 | 0.711 | 3.517 | 2.61E-01 | 0.123 | 0.009 | 1.649 | 1.13E-01 |
| *syt10* | 1.734 | 1.018 | 2.953 | 4.27E-02 | 0.732 | 0.241 | 2.221 | 5.81E-01 |
| *rgs6* | 2.209 | 0.862 | 5.666 | 9.90E-02 | 3.862 | 0.637 | 23.411 | 1.42E-01 |
| *hcn4* | 3.041 | 1.656 | 5.584 | 3.34E-04 | 1.209 | 0.359 | 4.072 | 7.59E-01 |
| *hcn4l* | 1.269 | 0.327 | 4.930 | 7.31E-01 | 3.112 | 0.328 | 29.534 | 3.23E-01 |
| *neo1a* | 0.936 | 0.465 | 1.881 | 8.52E-01 | 0.747 | 0.218 | 2.558 | 6.42E-01 |
| *neo1b* | 2.014 | 0.705 | 5.749 | 1.91E-01 | 0.379 | 0.081 | 1.785 | 2.20E-01 |
| *kiaa1755_1* | 0.749 | 0.426 | 1.318 | 3.16E-01 | 0.582 | 0.219 | 1.547 | 2.78E-01 |
| *kiaa1755_2* | 1.045 | 0.479 | 2.278 | 9.12E-01 | 8.170 | 1.077 | 61.988 | 4.22E-02 |
| time of day | 0.932 | 0.704 | 1.234 | 6.25E-01 | 1.079 | 0.583 | 1.998 | 8.09E-01 |
| intercept | 0.030 | 0.003 | 0.298 | 2.81E-03 | 0.017 | 0.000 | 0.830 | 4.00E-02 |

| | Sinoatrial arrest | | | |
|---|---|---|---|---|
| | 2dpf (n=321, 36 cases) | | | |
| Gene | OR | LCI | UCI | P |
| *gngt1* | 1.586 | 0.707 | 3.556 | 2.63E-01 |
| *syt10* | 1.639 | 0.952 | 2.821 | 7.47E-02 |
| *rgs6* | 1.988 | 0.737 | 5.357 | 1.75E-01 |
| *hcn4* | 2.906 | 1.556 | 5.428 | 8.19E-04 |
| *hcn4l* | 1.267 | 0.331 | 4.852 | 7.30E-01 |
| *neo1a* | 1.016 | 0.491 | 2.101 | 9.67E-01 |
| *neo1b* | 1.590 | 0.549 | 4.605 | 3.93E-01 |
| *kiaa1755_1* | 0.823 | 0.458 | 1.480 | 5.16E-01 |
| *kiaa1755_2* | 1.143 | 0.507 | 2.578 | 7.47E-01 |
| time of day | 0.928 | 0.697 | 1.236 | 6.10E-01 |
| intercept | 0.030 | 0.003 | 0.303 | 3.02E-03 |

| | Cardiac edema | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| | 2dpf (n=369, 15 cases) | | | | 5dpf (n=322, 14 cases) | | | |
| Gene | OR | LCI | UCI | P | OR | LCI | UCI | P |
| *gngt1* | 1.518 | 0.489 | 4.706 | 4.70E-01 | 7.780 | 1.755 | 34.480 | 6.92E-03 |
| *syt10* | 0.584 | 0.226 | 1.509 | 2.67E-01 | 0.901 | 0.291 | 2.787 | 8.57E-01 |
| *rgs6* | 0.702 | 0.086 | 5.739 | 7.42E-01 | 2.730 | 0.456 | 16.336 | 2.71E-01 |
| *hcn4* | 0.546 | 0.146 | 2.043 | 3.68E-01 | 0.380 | 0.065 | 2.213 | 2.82E-01 |
| *hcn4l* | 0.800 | 0.111 | 5.770 | 8.25E-01 | 0.039 | 0.000 | 63.816 | 3.90E-01 |
| *neo1a* | 0.844 | 0.309 | 2.307 | 7.41E-01 | 0.253 | 0.080 | 0.796 | 1.88E-02 |
| *neo1b* | 1.154 | 0.328 | 4.064 | 8.23E-01 | 0.217 | 0.046 | 1.028 | 5.42E-02 |
| *kiaa1755_1* | 1.137 | 0.505 | 2.563 | 7.56E-01 | 2.126 | 0.604 | 7.486 | 2.40E-01 |
| *kiaa1755_2* | 1.317 | 0.430 | 4.032 | 6.29E-01 | 2.011 | 0.414 | 9.761 | 3.86E-01 |
| time of day | 0.963 | 0.633 | 1.465 | 8.59E-01 | 1.675 | 0.930 | 3.016 | 8.57E-02 |
| intercept | 0.036 | 0.002 | 0.697 | 2.78E-02 | 0.017 | 0.001 | 0.513 | 1.90E-02 |

*Continued Supplemental Table 5*

| Gene | Uncontrolled atrial contractions 5dpf (n=326, 9 cases) | | | |
|---|---|---|---|---|
| | OR | LCI | UCI | P |
| gngt1 | 2.578 | 0.502 | 13.247 | 2.57E-01 |
| syt10 | 0.343 | 0.065 | 1.813 | 2.08E-01 |
| rgs6 | 1.313 | 0.138 | 12.517 | 8.13E-01 |
| hcn4 | 1.310 | 0.339 | 5.064 | 6.96E-01 |
| hcn4l | - | - | - | - |
| neo1a | 0.294 | 0.085 | 1.012 | 5.23E-02 |
| neo1b | 0.988 | 0.203 | 4.810 | 9.88E-01 |
| kiaa1755_1 | 0.644 | 0.230 | 1.802 | 4.02E-01 |
| kiaa1755_2 | 2.107 | 0.352 | 12.615 | 4.14E-01 |
| time of day | 0.648 | 0.335 | 1.251 | 1.96E-01 |
| intercept | 0.245 | 0.004 | 14.946 | 5.03E-01 |

| Gene | Abnormal morphology and impaired contractility 5dpf (n=326, 6 cases) | | | |
|---|---|---|---|---|
| | OR | LCI | UCI | P |
| gngt1 | 41.908 | 3.402 | 516.185 | 3.55E-03 |
| syt10 | 0.928 | 0.206 | 4.187 | 9.23E-01 |
| rgs6 | - | - | - | - |
| hcn4 | 0.348 | 0.021 | 5.885 | 4.64E-01 |
| hcn4l | - | - | - | - |
| neo1a | 0.15 | 0.026 | 0.851 | 3.21E-02 |
| neo1b | 0.145 | 0.015 | 1.379 | 9.29E-02 |
| kiaa1755_1 | 1.915 | 0.371 | 9.882 | 4.38E-01 |
| kiaa1755_2 | 0.628 | 0.099 | 3.98 | 6.21E-01 |
| time of day | 1.316 | 0.582 | 2.978 | 5.09E-01 |
| intercept | 0.076 | 0.001 | 5.739 | 2.43E-01 |

Associations of cardiac outcomes with the number of mutated alleles across the nine orthologues, weighted by their predicted effect on protein function. At 2 and 5 days post fertilization (dpf), associations were analyzed using logistic regression for outcomes that were observed in at least five larvae. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors. '-' indicates that parameters could not be estimated.  For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 6: Transcript-specific association analysis for HRV and heart rate**

| Age | n | Factor | Gene | Heart rate variability | | | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | | | Effect size | SE | LCI | UCI | P |
| 2 | 279 | fixed effects | gngt1 | -0.052 | 0.107 | -0.261 | 0.157 | 6.25E-01 |
| | | | syt10 | -0.070 | 0.077 | -0.221 | 0.081 | 3.62E-01 |
| | | | rgs6 | 0.000 | 0.175 | -0.343 | 0.344 | 9.98E-01 |
| | | | hcn4 | -0.004 | 0.101 | -0.202 | 0.194 | 9.70E-01 |
| | | | hcn4l | 0.165 | 0.178 | -0.184 | 0.514 | 3.55E-01 |
| | | | neo1a | -0.028 | 0.095 | -0.214 | 0.158 | 7.69E-01 |
| | | | neo1b | -0.065 | 0.113 | -0.287 | 0.158 | 5.68E-01 |
| | | | kiaa1755_1 | 0.152 | 0.076 | 0.003 | 0.302 | 4.61E-02 |
| | | | kiaa1755_2 | 0.074 | 0.097 | -0.115 | 0.264 | 4.42E-01 |
| | | | time of day | 0.503 | 0.045 | 0.414 | 0.592 | 1.77E-28 |
| | | | Heart rate (in SD) | -0.459 | 0.056 | -0.570 | -0.348 | 4.59E-16 |
| | | | intercept | -1.614 | 0.300 | -2.202 | -1.026 | 7.54E-08 |
| | | random effects | variation by batch | 0.340 | 0.113 | 0.177 | 0.652 | - |
| | | | residual | 0.765 | 0.033 | 0.704 | 0.832 | - |
| 5 | 291 | fixed effects | gngt1 | 0.278 | 0.108 | 0.067 | 0.489 | 9.81E-03 |
| | | | syt10 | 0.047 | 0.071 | -0.093 | 0.187 | 5.12E-01 |
| | | | rgs6 | -0.009 | 0.157 | -0.317 | 0.299 | 9.56E-01 |
| | | | hcn4 | -0.032 | 0.095 | -0.219 | 0.154 | 7.35E-01 |
| | | | hcn4l | 0.142 | 0.177 | -0.205 | 0.489 | 4.22E-01 |
| | | | neo1a | 0.017 | 0.096 | -0.171 | 0.204 | 8.62E-01 |
| | | | neo1b | 0.077 | 0.121 | -0.160 | 0.313 | 5.25E-01 |
| | | | kiaa1755_1 | 0.000 | 0.077 | -0.150 | 0.150 | 1.00E+00 |
| | | | kiaa1755_2 | 0.008 | 0.100 | -0.189 | 0.204 | 9.40E-01 |
| | | | time of day | 0.524 | 0.052 | 0.422 | 0.626 | 7.54E-24 |
| | | | Heart rate (in SD) | -0.645 | 0.054 | -0.751 | -0.540 | 2.86E-33 |
| | | | intercept | -1.893 | 0.383 | -2.643 | -1.142 | 7.67E-07 |
| | | random effects | variation by batch | 0.540 | 0.185 | 0.276 | 1.056 | - |
| | | | residual | 0.769 | 0.032 | 0.708 | 0.834 | - |

| Age | n | Factor | Gene | Heart rate | | | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | | | Effect size | SE | LCI | UCI | P |
| 2 | 279 | fixed effects | gngt1 | -0.061 | 0.102 | -0.261 | 0.139 | 5.48E-01 |
| | | | syt10 | -0.026 | 0.074 | -0.170 | 0.118 | 7.22E-01 |
| | | | rgs6 | -0.045 | 0.168 | -0.374 | 0.284 | 7.88E-01 |
| | | | hcn4 | 0.047 | 0.096 | -0.142 | 0.236 | 6.27E-01 |
| | | | hcn4l | 0.105 | 0.169 | -0.226 | 0.436 | 5.35E-01 |
| | | | neo1a | 0.055 | 0.090 | -0.122 | 0.233 | 5.39E-01 |
| | | | neo1b | -0.121 | 0.108 | -0.333 | 0.091 | 2.62E-01 |
| | | | kiaa1755_1 | 0.120 | 0.073 | -0.022 | 0.263 | 9.83E-02 |
| | | | kiaa1755_2 | 0.078 | 0.092 | -0.104 | 0.259 | 4.01E-01 |
| | | | time of day | 0.550 | 0.040 | 0.472 | 0.627 | 9.14E-44 |
| | | | HRV (in SD) | -0.408 | 0.051 | -0.508 | -0.308 | 1.23E-15 |
| | | | intercept | -1.770 | 0.258 | -2.276 | -1.264 | 7.14E-12 |
| | | random effects | variation by batch | 0.185 | 0.077 | 0.082 | 0.420 | - |
| | | | residual | 0.734 | 0.031 | 0.675 | 0.798 | - |
| 5 | 291 | fixed effects | gngt1 | 0.094 | 0.097 | -0.095 | 0.283 | 3.31E-01 |
| | | | syt10 | -0.033 | 0.063 | -0.157 | 0.091 | 6.02E-01 |
| | | | rgs6 | -0.087 | 0.139 | -0.360 | 0.186 | 5.32E-01 |
| | | | hcn4 | 0.150 | 0.084 | -0.014 | 0.315 | 7.37E-02 |
| | | | hcn4l | 0.204 | 0.157 | -0.104 | 0.512 | 1.94E-01 |
| | | | neo1a | 0.002 | 0.085 | -0.165 | 0.168 | 9.82E-01 |
| | | | neo1b | 0.006 | 0.107 | -0.204 | 0.216 | 9.57E-01 |
| | | | kiaa1755_1 | 0.031 | 0.068 | -0.102 | 0.164 | 6.50E-01 |
| | | | kiaa1755_2 | 0.197 | 0.088 | 0.024 | 0.369 | 2.59E-02 |
| | | | time of day | 0.564 | 0.043 | 0.480 | 0.648 | 1.97E-39 |
| | | | HRV (in SD) | -0.516 | 0.043 | -0.599 | -0.432 | 7.79E-34 |
| | | | intercept | -2.137 | 0.375 | -2.872 | -1.402 | 1.20E-08 |
| | | random effects | variation by batch | 0.619 | 0.204 | 0.324 | 1.181 | - |
| | | | residual | 0.682 | 0.029 | 0.629 | 0.740 | - |

Associations of heart rate variability (HRV) and heart rate (mutually adjusted) with the number of mutated alleles across the nine orthologues, weighted by their predicted effect on protein function. Associations were examined for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata), using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Associations were adjusted for time of day and the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 7: Transcript-specific association analysis for heart rate variability and heart rate in larvae without a sinoatrial pause before or after the recording**

| Age | n | Factor | Gene | Heart rate variability | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | Effect size | SE | LCI | UCI | P |
| 2 | 234 | fixed effects | gngt1 | -0.066 | 0.122 | -0.306 | 0.173 | 5.88E-01 |
| | | | syt10 | -0.058 | 0.084 | -0.223 | 0.107 | 4.91E-01 |
| | | | rgs6 | 0.036 | 0.201 | -0.358 | 0.429 | 8.60E-01 |
| | | | hcn4 | 0.029 | 0.111 | -0.189 | 0.247 | 7.95E-01 |
| | | | hcn4l | 0.171 | 0.206 | -0.232 | 0.575 | 4.05E-01 |
| | | | neo1a | -0.056 | 0.105 | -0.263 | 0.151 | 5.95E-01 |
| | | | neo1b | -0.094 | 0.125 | -0.338 | 0.150 | 4.50E-01 |
| | | | kiaa1755_1 | 0.180 | 0.082 | 0.019 | 0.340 | 2.81E-02 |
| | | | kiaa1755_2 | 0.093 | 0.102 | -0.107 | 0.292 | 3.64E-01 |
| | | | time of day | 0.555 | 0.050 | 0.456 | 0.654 | 4.10E-28 |
| | | | Heart rate (in SD) | -0.534 | 0.063 | -0.658 | -0.411 | 1.81E-17 |
| | | | intercept | -1.770 | 0.316 | -2.388 | -1.151 | 2.05E-08 |
| | | random effects | variation by batch | 0.344 | 0.118 | 0.176 | 0.674 | - |
| | | | residual | 0.769 | 0.036 | 0.702 | 0.843 | - |
| 5 | 283 | fixed effects | gngt1 | 0.264 | 0.106 | 0.056 | 0.472 | 1.27E-02 |
| | | | syt10 | -0.005 | 0.071 | -0.144 | 0.134 | 9.42E-01 |
| | | | rgs6 | -0.025 | 0.157 | -0.333 | 0.282 | 8.72E-01 |
| | | | hcn4 | -0.042 | 0.094 | -0.226 | 0.142 | 6.56E-01 |
| | | | hcn4l | 0.135 | 0.173 | -0.203 | 0.474 | 4.33E-01 |
| | | | neo1a | -0.002 | 0.094 | -0.185 | 0.182 | 9.86E-01 |
| | | | neo1b | 0.055 | 0.118 | -0.176 | 0.286 | 6.41E-01 |
| | | | kiaa1755_1 | 0.020 | 0.077 | -0.131 | 0.170 | 7.99E-01 |
| | | | kiaa1755_2 | 0.081 | 0.102 | -0.119 | 0.281 | 4.28E-01 |
| | | | time of day | 0.534 | 0.051 | 0.434 | 0.633 | 8.87E-26 |
| | | | Heart rate (in SD) | -0.623 | 0.053 | -0.727 | -0.519 | 5.00E-32 |
| | | | intercept | -1.978 | 0.372 | -2.707 | -1.250 | 1.01E-07 |
| | | random effects | variation by batch | 0.516 | 0.178 | 0.263 | 1.013 | - |
| | | | residual | 0.747 | 0.032 | 0.687 | 0.812 | - |

| Age | N | Factor | Gene | Heart rate | | | | |
|---|---|---|---|---|---|---|---|---|
| | | | | Effect size | SE | LCI | UCI | P |
| 2 | 234 | fixed effects | gngt1 | -0.008 | 0.112 | -0.228 | 0.212 | 9.43E-01 |
| | | | syt10 | -0.012 | 0.077 | -0.162 | 0.139 | 8.78E-01 |
| | | | rgs6 | -0.114 | 0.184 | -0.474 | 0.245 | 5.33E-01 |
| | | | hcn4 | 0.041 | 0.101 | -0.157 | 0.240 | 6.82E-01 |
| | | | hcn4l | 0.048 | 0.187 | -0.317 | 0.414 | 7.95E-01 |
| | | | neo1a | 0.050 | 0.095 | -0.137 | 0.236 | 6.01E-01 |
| | | | neo1b | -0.165 | 0.113 | -0.386 | 0.057 | 1.45E-01 |
| | | | kiaa1755_1 | 0.153 | 0.075 | 0.007 | 0.300 | 4.00E-02 |
| | | | kiaa1755_2 | 0.113 | 0.093 | -0.069 | 0.295 | 2.24E-01 |
| | | | time of day | 0.586 | 0.041 | 0.505 | 0.666 | 5.48E-46 |
| | | | HRV (in SD) | -0.434 | 0.051 | -0.535 | -0.333 | 3.28E-17 |
| | | | intercept | -1.890 | 0.256 | -2.392 | -1.388 | 1.60E-13 |
| | | random effects | variation by batch | 0.171 | 0.077 | 0.071 | 0.414 | - |
| | | | residual | 0.705 | 0.033 | 0.643 | 0.773 | - |
| 5 | 283 | fixed effects | gngt1 | 0.099 | 0.098 | -0.093 | 0.291 | 3.11E-01 |
| | | | syt10 | -0.056 | 0.065 | -0.183 | 0.072 | 3.92E-01 |
| | | | rgs6 | -0.085 | 0.144 | -0.366 | 0.197 | 5.55E-01 |
| | | | hcn4 | 0.157 | 0.085 | -0.010 | 0.325 | 6.54E-02 |
| | | | hcn4l | 0.198 | 0.158 | -0.111 | 0.508 | 2.09E-01 |
| | | | neo1a | 0.007 | 0.086 | -0.161 | 0.176 | 9.31E-01 |
| | | | neo1b | -0.002 | 0.108 | -0.213 | 0.210 | 9.86E-01 |
| | | | kiaa1755_1 | 0.037 | 0.070 | -0.100 | 0.175 | 5.94E-01 |
| | | | kiaa1755_2 | 0.208 | 0.093 | 0.027 | 0.390 | 2.47E-02 |
| | | | time of day | 0.574 | 0.044 | 0.488 | 0.659 | 2.21E-39 |
| | | | HRV (in SD) | -0.530 | 0.045 | -0.617 | -0.443 | 1.35E-32 |
| | | | intercept | -2.187 | 0.377 | -2.927 | -1.448 | 6.58E-09 |
| | | random effects | variation by batch | 0.617 | 0.204 | 0.323 | 1.178 | - |
| | | | residual | 0.684 | 0.029 | 0.629 | 0.743 | - |

Associations of heart rate variability (HRV) and heart rate (mutually adjusted) with the number of mutated alleles across the nine orthologues, weighted by their predicted effect on protein function. Associations were examined for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata), using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Larvae with a sinoatrial pause before or after the recording were excluded from the analysis in this sensitivity analysis. Analyses were adjusted for time of day and the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 8: Transcript-specific association analyses for body size**

### Body length

| Factor | Gene | 2dpf (n=332) | | | | | 5dpf (n=240) | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Effect size | SE | LCI | UCI | P | Effect size | SE | LCI | UCI | P |
| | *gngt1* | -0.053 | 0.114 | -0.278 | 0.171 | 6.41E-01 | 0.027 | 0.112 | -0.193 | 0.246 | 8.12E-01 |
| | *syt10* | 0.086 | 0.080 | -0.070 | 0.242 | 2.79E-01 | 0.040 | 0.073 | -0.103 | 0.183 | 5.85E-01 |
| | *rgs6* | 0.015 | 0.171 | -0.321 | 0.350 | 9.31E-01 | -0.271 | 0.157 | -0.577 | 0.036 | 8.38E-02 |
| | *hcn4* | -0.092 | 0.103 | -0.294 | 0.109 | 3.69E-01 | 0.050 | 0.096 | -0.138 | 0.238 | 6.02E-01 |
| | *hcn4l* | 0.159 | 0.204 | -0.241 | 0.559 | 4.37E-01 | -0.003 | 0.208 | -0.411 | 0.406 | 9.90E-01 |
| fixed effects | *neo1a* | -0.201 | 0.103 | -0.403 | 0.001 | 5.08E-02 | 0.061 | 0.094 | -0.123 | 0.246 | 5.16E-01 |
| | *neo1b* | -0.117 | 0.119 | -0.350 | 0.116 | 3.26E-01 | -0.079 | 0.118 | -0.311 | 0.152 | 5.02E-01 |
| | *kiaa1755_1* | 0.107 | 0.078 | -0.046 | 0.260 | 1.70E-01 | -0.005 | 0.075 | -0.152 | 0.143 | 9.50E-01 |
| | *kiaa1755_2* | -0.208 | 0.107 | -0.417 | 0.001 | 5.12E-02 | -0.071 | 0.107 | -0.281 | 0.139 | 5.07E-01 |
| | Time of day | 0.113 | 0.042 | 0.030 | 0.195 | 7.49E-03 | 0.081 | 0.048 | -0.013 | 0.175 | 9.26E-02 |
| | Intercept | 0.166 | 0.330 | -0.480 | 0.812 | 6.15E-01 | -0.235 | 0.429 | -1.077 | 0.607 | 5.84E-01 |
| random effects | Variation by batch | 0.407 | 0.131 | 0.217 | 0.764 | - | 0.711 | 0.231 | 0.376 | 1.345 | - |
| | Residual | 0.897 | 0.035 | 0.831 | 0.969 | - | 0.719 | 0.033 | 0.657 | 0.787 | - |

### Dorsal body surface area

| Factor | Gene | 2dpf (n=322) | | | | | 5dpf (n=233) | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Effect size | SE | LCI | UCI | P | Effect size | SE | LCI | UCI | P |
| | *gngt1* | 0.112 | 0.109 | -0.101 | 0.324 | 3.04E-01 | -0.328 | 0.136 | -0.595 | -0.060 | 1.63E-02 |
| | *syt10* | -0.057 | 0.076 | -0.206 | 0.091 | 4.48E-01 | -0.040 | 0.087 | -0.211 | 0.132 | 6.48E-01 |
| | *rgs6* | 0.038 | 0.163 | -0.282 | 0.358 | 8.16E-01 | -0.135 | 0.191 | -0.508 | 0.239 | 4.80E-01 |
| | *hcn4* | -0.106 | 0.098 | -0.298 | 0.086 | 2.80E-01 | -0.016 | 0.113 | -0.238 | 0.205 | 8.84E-01 |
| | *hcn4l* | -0.151 | 0.196 | -0.535 | 0.233 | 4.41E-01 | 0.359 | 0.243 | -0.118 | 0.835 | 1.41E-01 |
| fixed effects | *neo1a* | 0.218 | 0.099 | 0.025 | 0.412 | 2.66E-02 | 0.025 | 0.113 | -0.195 | 0.246 | 8.23E-01 |
| | *neo1b* | -0.100 | 0.115 | -0.325 | 0.126 | 3.86E-01 | -0.075 | 0.147 | -0.363 | 0.213 | 6.12E-01 |
| | *kiaa1755_1* | 0.016 | 0.075 | -0.131 | 0.163 | 8.32E-01 | 0.006 | 0.090 | -0.170 | 0.183 | 9.43E-01 |
| | *kiaa1755_2* | 0.060 | 0.102 | -0.139 | 0.259 | 5.55E-01 | -0.015 | 0.127 | -0.264 | 0.233 | 9.03E-01 |
| | Time of day | -0.046 | 0.040 | -0.125 | 0.032 | 2.49E-01 | -0.084 | 0.055 | -0.192 | 0.024 | 1.26E-01 |
| | Intercept | -0.152 | 0.338 | -0.815 | 0.510 | 6.52E-01 | 0.386 | 0.383 | -0.365 | 1.137 | 3.13E-01 |
| random effects | Variation by batch | 0.492 | 0.154 | 0.267 | 0.910 | - | 0.325 | 0.127 | 0.151 | 0.697 | - |
| | Residual | 0.842 | 0.034 | 0.779 | 0.911 | - | 0.852 | 0.040 | 0.777 | 0.934 | - |

### Lateral body surface area

| Factor | Gene | 2dpf (n=300) | | | | | 5dpf (n=256) | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Effect size | SE | LCI | UCI | P | Effect size | SE | LCI | UCI | P |
| | *gngt1* | -0.056 | 0.121 | -0.293 | 0.181 | 6.41E-01 | -0.063 | 0.124 | -0.306 | 0.179 | 6.09E-01 |
| | *syt10* | 0.042 | 0.085 | -0.124 | 0.208 | 6.21E-01 | 0.027 | 0.083 | -0.135 | 0.189 | 7.45E-01 |
| | *rgs6* | -0.128 | 0.183 | -0.487 | 0.231 | 4.84E-01 | -0.022 | 0.175 | -0.365 | 0.322 | 9.01E-01 |
| | *hcn4* | 0.151 | 0.108 | -0.061 | 0.363 | 1.64E-01 | -0.087 | 0.107 | -0.297 | 0.122 | 4.14E-01 |
| | *hcn4l* | -0.047 | 0.221 | -0.479 | 0.385 | 8.32E-01 | 0.442 | 0.216 | 0.019 | 0.864 | 4.04E-02 |
| fixed effects | *neo1a* | -0.036 | 0.110 | -0.252 | 0.180 | 7.42E-01 | -0.024 | 0.107 | -0.234 | 0.186 | 8.23E-01 |
| | *neo1b* | 0.257 | 0.130 | 0.002 | 0.512 | 4.80E-02 | -0.096 | 0.141 | -0.373 | 0.180 | 4.96E-01 |
| | *kiaa1755_1* | 0.027 | 0.085 | -0.140 | 0.194 | 7.52E-01 | 0.099 | 0.088 | -0.073 | 0.271 | 2.61E-01 |
| | *kiaa1755_2* | -0.224 | 0.117 | -0.454 | 0.005 | 5.53E-02 | 0.018 | 0.120 | -0.216 | 0.252 | 8.79E-01 |
| | Time of day | -0.049 | 0.045 | -0.138 | 0.039 | 2.72E-01 | -0.070 | 0.052 | -0.173 | 0.032 | 1.79E-01 |
| | Intercept | 0.061 | 0.335 | -0.595 | 0.718 | 8.55E-01 | 0.073 | 0.391 | -0.694 | 0.839 | 8.53E-01 |
| random effects | Variation by batch | 0.340 | 0.117 | 0.173 | 0.669 | - | 0.471 | 0.162 | 0.240 | 0.924 | - |
| | Residual | 0.914 | 0.038 | 0.843 | 0.991 | - | 0.829 | 0.037 | 0.760 | 0.905 | - |

### Body volume

| Factor | Gene | 2dpf (n=299) | | | | | 5dpf (n=198) | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Effect size | SE | LCI | UCI | P | Effect size | SE | LCI | UCI | P |
| | *gngt1* | -0.051 | 0.111 | -0.270 | 0.167 | 6.45E-01 | -0.071 | 0.132 | -0.330 | 0.187 | 5.88E-01 |
| | *syt10* | 0.008 | 0.079 | -0.147 | 0.163 | 9.20E-01 | 0.046 | 0.082 | -0.114 | 0.207 | 5.73E-01 |
| | *rgs6* | -0.250 | 0.181 | -0.606 | 0.106 | 1.68E-01 | -0.202 | 0.173 | -0.542 | 0.137 | 2.43E-01 |
| | *hcn4* | 0.123 | 0.099 | -0.070 | 0.317 | 2.12E-01 | -0.044 | 0.104 | -0.249 | 0.160 | 6.71E-01 |
| | *hcn4l* | -0.066 | 0.194 | -0.446 | 0.314 | 7.35E-01 | 0.319 | 0.246 | -0.163 | 0.800 | 1.94E-01 |
| fixed effects | *neo1a* | 0.105 | 0.103 | -0.097 | 0.307 | 3.07E-01 | 0.043 | 0.105 | -0.162 | 0.248 | 6.82E-01 |
| | *neo1b* | 0.175 | 0.121 | -0.063 | 0.413 | 1.49E-01 | -0.061 | 0.146 | -0.347 | 0.225 | 6.78E-01 |
| | *kiaa1755_1* | -0.035 | 0.079 | -0.190 | 0.120 | 6.57E-01 | 0.192 | 0.083 | 0.028 | 0.355 | 2.14E-02 |
| | *kiaa1755_2* | -0.019 | 0.106 | -0.226 | 0.188 | 8.58E-01 | -0.036 | 0.124 | -0.279 | 0.207 | 7.72E-01 |
| | Time of day | -0.056 | 0.042 | -0.139 | 0.027 | 1.85E-01 | -0.053 | 0.053 | -0.158 | 0.051 | 3.18E-01 |
| | Intercept | -0.184 | 0.345 | -0.860 | 0.492 | 5.94E-01 | -0.296 | 0.441 | -1.160 | 0.569 | 5.03E-01 |
| random effects | Variation by batch | 0.496 | 0.155 | 0.268 | 0.916 | - | 0.598 | 0.199 | 0.311 | 1.149 | - |
| | Residual | 0.841 | 0.035 | 0.776 | 0.912 | - | 0.733 | 0.037 | 0.663 | 0.809 | - |

Associations were analyzed for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata) using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). For simplicity, *kiaa1755_1* refers to *quo*, and *kiaa1755_2* to *si:dkey-65j6.2*.

**Supplemental Table 9: Druggability of interacting partners**

| Candidate | Interaction partner | DGIdb categories | Drug available? (y/n) | Drug name (FDA approved) | Drug category |
|---|---|---|---|---|---|
| | GNAI1 | Tumor supressor | yes | | |
| | GNAI3 | | | | |
| | GNAL | | | | |
| | GNAS | Druggable genome, Drug resistance, Clinically actionable | yes | Fluorouracil | Anti-neoplastic agents |
| | GNAT1 | | | | |
| | GNB1 | | | | |
| | GNB2 | Transporter, Ion channel | | | |
| | GNB3 | | | | |
| | GNB4 | | | | |
| | GNB5 | Transporter, Ion channel | | | |
| GNGT1 | OR10W1 | Druggable genome, G-protein coupled receptor | | | |
| | OR1L6 | Druggable genome, G-protein coupled receptor | | | |
| | OR2AJ1 | Druggable genome, G-protein coupled receptor | | | |
| | OR4C5 | Druggable genome, G-protein coupled receptor | | | |
| | OR4F21 | Druggable genome, G-protein coupled receptor | | | |
| | OR4K15 | Druggable genome, G-protein coupled receptor | | | |
| | OR51H1 | Druggable genome, G-protein coupled receptor | | | |
| | PDC | Phospolipase | | | |
| | PDCL | | | | |
| | PLEKHB1 | | | | |
| | RHO | Druggable genome, G-protein coupled receptor | yes | Halothane | Anesthetics |
| | | | | Telmisartan, Irbesartan, Valsartan, Eprosartan, Losartan | Anti-hypertensive agents |
| | | | | Atorvastatin | Anti-cholesterimic agents |
| | AGTR1 | Druggable genome, G-protein coupled receptor, Phospolipase | yes | Cyclosporine | Immunosuppresive agents |
| | | | | Indomethacin, Dexamethasone | Anti-inflammatory agents |
| | | | | Levodopa | Anti-parkinson agents |
| | GNAI1 | Tumor supressor | yes | | |
| | GNAI3 | | | | |
| | GNAT1 | | | | |
| | GNAT2 | G-protein coupled receptor | | | |
| GNG11 | GNAT3 | | | | |
| | GNB1 | | | | |
| | GNB2 | Transporter, Ion channel | | | |
| | GNB3 | | | | |
| | GNB4 | | | | |
| | NMB | Druggable genome, Hormone activity | | | |
| | PDC | Phospolipase | | | |
| | PTGIR | Druggable genome, G-protein coupled receptor | yes | Iloprost, Treprostinil, Selexipag, Epoprostenol | Anti-hypertensive agents |
| | | | | Dinoprost thromethamine, Misoprostol | Abortifacient agents |
| | SMURF1 | | | | |
| | BOP1 | Tumor supressor | | | |
| | SNAP23 | Druggable genome | | | |
| | SNAP25 | Druggable genome | yes | Diazoxide | Anti-hypertensive agents |
| SYT10 | | | | Botulinum toxin type A | Neuromuscular blocking agents |
| | SNAP29 | Druggable genome | | | |
| | SNAP47 | | | | |
| | SYT6 | | | | |
| | HCN1 | Druggable genome, Transporter, Ion channel | yes | Ivabradine | Channel blocker |
| HCN4 | HCN2 | Druggable genome, Transporter, Ion channel | yes | | |
| | HCN3 | Druggable genome, Transporter, Ion channel | yes | Ivabradine | Channel blocker |
| | PEX5L | | | | |
| | CDH15 | | | | |
| | CDON | | | | |
| | HFE2 | Kinase, Serine-threonine kinase, Cell surface | | | |
| | NTN1 | Druggable genome | | | |
| | NTN3 | Druggable genome | | | |
| NEO1 | NTN4 | Druggable genome | | | |
| | PITPNA | Transporter | | | |
| | PTK2 | Druggable genome, Kinase, Tyrosine kinase, Lipid kinase | yes | | |
| | RGMA | | | | |
| | RGMB | | | | |
| | TTN | Druggable genome, Kinase, Serine-threonine kinase, Tyrosine kinase | | | |
| | UNC5B | | | | |
| KIAA1755 | - | | | | |

Human candidate genes were examined for their interaction partners using STRING and Genemania, and their classification according to the Drug gene interaction database (DGIdb). Proteins that are already targeted by a drug are indicated by a "y". A selection of FDA-approved drugs was distilled from DGIdb, including their categorization.