Translating GWAS-identified loci for cardiac rhythm and rate using an *in vivo* image- and CRISPR-Cas9-based approach

Benedikt von der Heyde[1,2], Anastasia Emmanouilidou[1,2], Tiffany Klingström[1,2], Eugenia Mazzaferro[1,2], Silvia Vicenzi[1,2], Sitaf Jumaa[1,2], Olga Dethlefsen[3,4], Harold Snieder[5], Eco de Geus[6], Erik Ingelsson[2,7,8,9], Amin Allalou[2,10], Hannah L. Brooke[11], Marcel den Hoed[#1,2]

[1]The Beijer laboratory and Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden.
[2]Science for Life Laboratory, Uppsala University, Uppsala, Sweden.
[3]Science for Life Laboratory, Stockholm University, Stockholm, Sweden.
[4]National Bioinformatics Infrastructure Sweden, Stockholm University, Stockholm, Sweden.
[5]Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands.
[6]Department of Biological Psychology, Vrije Universiteit Amsterdam, Amsterdam, the Netherlands.
[7]Department of Medical Sciences, Molecular Epidemiology, Uppsala University, Uppsala, Sweden.
[8]Department of Medicine, Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA, USA.
[9]Stanford Cardiovascular Institute, Stanford University, Stanford, CA 94305.
[10]Department of Information Technology, Division of Visual Information and Interaction, Uppsala University, Uppsala, Sweden.
[11]Department of Public Health and Caring Sciences, Uppsala University, Uppsala, Sweden.

Corresponding author: Marcel den Hoed

Abstract

A meta-analysis of genome-wide association studies (GWAS) identified eight loci

that are associated with heart rate variability (HRV) in data from 53,174 individuals.

However, functional follow-up experiments - aiming to identify and characterize

causal genes in these loci - have not yet been performed. We developed an image- and

CRISPR-Cas9-based pipeline to systematically characterize candidate genes for HRV

in live zebrafish embryos and larvae. Nine zebrafish orthologues of six human

candidate genes were targeted simultaneously in fertilized eggs from fish that

transgenically express GFP on smooth muscle cells (Tg(*acta2:GFP*)), to visualize the

beating heart using a fluorescence microscope. An automated analysis of repeated 30s

recordings of 381 live zebrafish atria at 2 and 5 days post-fertilization highlighted

genes that influence HRV (*hcn4* and *si:dkey-65j6.2*); heart rate (*rgs6 and hcn4*) and

the risk of sinoatrial pauses and arrests (*hcn4*). Hence, our screen confirmed the role

of established genes for heart rate and rhythm (*rgs6* and *hcn4*), and highlighted a

novel gene implicated in HRV (*si:dkey-65j6.2*).

Introduction

Heart rate variability (HRV) reflects the inter-beat variation of heart rate. HRV is controlled by the sinoatrial node, which receives input from the autonomic nervous system. Autonomic imbalance has been associated with work stress and other modifiable or non-modifiable risk factors[1], and is reflected in lower HRV. Lower HRV has been associated with higher cardiac morbidity and mortality[2], and with a higher risk of all-cause mortality[3]. HRV can be quantified non-invasively using the RR interval of a 12-lead ECG, making HRV a useful clinical marker for perturbations of the autonomic nervous system. However, the genetic basis of HRV remains largely elusive.

Recently, we and others identified the first loci that are robustly associated with HRV, using a meta-analysis of genome-wide association studies (GWAS) with data from 53,174 participants[4]. Eleven of the 17 associated single nucleotide polymorphisms (SNPs) overlapped with loci that are also associated with resting heart rate[5]. The heart rate associated loci in turn had on aggregate been associated with altered cardiac conduction and risk of sick sinus syndrome[5]. *In silico* functional annotation of the five loci that are associated with both HRV[6] and heart rate[5] previously resulted in the prioritization of six candidate genes that are anticipated to be causal[6]. Functional follow-up experiments - ideally *in vivo* - are required to conclude if these genes are indeed causal, and examine if they influence HRV, heart rate, and/or cardiac conduction.

Mouse models are commonly used in cardiac research, but mice show substantial differences in cardiac rate and electrophysiology when compared with humans[7]. Inherently, these differences complicate extrapolation of results from mouse models to humans[7]. Additionally, rodents are not suitable for high-throughput, *in vivo*

characterization of cardiac rhythm and rate. Such screens are essential to

systematically characterize positional candidate genes in the large number of loci that

have now been identified by GWAS for cardiac rhythm[4], rate[8] and conduction[9].

Hence, novel model systems that facilitate systematic, *in vivo* characterization of a

large number of candidate genes are desirable.

In recent years, the zebrafish has become an important model system for genetic

and drug screens for human disease[10,11]. Despite morphological differences to the

human heart, zebrafish have a functional, beating heart at ~24 hours post-fertilization,

and the ECG of the two-chambered zebrafish heart is similar to that of humans[12].

Conditions affecting cardiac electrophysiology have previously been successfully

modeled in zebrafish. For example *nkx2.5* has been demonstrated to be necessary for

establishing HRV[13], mutations in *kcnh2* were used to model long QT syndrome[14],

trpm7 was shown to influence heart rate and risk for sinoatrial pauses[15], and *hcn4*

knockdown modeled sick sinus syndrome in zebrafish analogous to the human

condition[16]. Fluorescently labeled transgenes facilitate visualization of cell types and

tissues of interest, which can now be accomplished in high-throughput thanks to

advances in automated positioning of non-embedded, live zebrafish embryos[17]. The

automated positioning system has already been used to identify and characterize small

molecules for myelination[18] and thyroid disease[19]. In addition, the zebrafish has a

well-annotated genome, with orthologues of at least 71.4% of human genes[20]. These

genes can be targeted efficiently and in a multiplexed manner thanks to recent

advances in Clustered, Regulatory Interspaced, Short Palindromic Repeats (CRISPR)

and CRISPR-associated systems (Cas)[21]. All characteristics combined make zebrafish

embryos and larvae an attractive model system to systematically characterize

candidate genes for cardiac rhythm, rate and conduction.

The aim of this study was to objectively characterize the most promising candidate

genes in GWAS-identified loci for HRV as well as heart rate for a role in cardiac

rhythm, rate and function using a large-scale, image-based screen in zebrafish

embryos and larvae.

Results

Descriptive results

At 2dpf, some embryos showed sinoatrial pauses (n=39, 10.3%, Supplementary Movie 1) and arrests (n=36, 9.5%, Supplementary Movie 2); cardiac edema (n=16, 4.2%); or uncontrolled atrial contractions (n=1, 0.2%, Supplementary Movie 3). Three larvae died between 2 and 5dpf (Supplementary Figure 1). At 5dpf, fewer larvae displayed sinoatrial pauses (n=9, 2.7%, one of which had a sinoatrial pause at 2dpf); sinoatrial arrests (n=3, 0.9%, none of which had a sinoatrial arrest at 2dpf); and cardiac edema (n=15, 4.5%, nine of which had an edema at 2dpf); while more larvae showed uncontrolled atrial contractions (n=9, 2.7%, none of which had uncontrolled atrial contraction at 2dpf); or an abnormal cardiac morphology and impaired cardiac contractility (n=6, 1.8%, Supplementary Movie 4).

After imaging at 5dpf, all larvae were sequenced at the nine CRISPR-Cas9 targeted sites (Figure 1, Table 1), as well as at three *in silico* predicted off-target sites (Supplementary Table 1). Transcript-specific dosages were calculated by weighting the number of mutated alleles by their predicted impact on protein function based on Ensembl's variant effect predictor (VEP). No CRISPR-Cas9 induced mutations were identified at any of the predicted off-target sites (Supplementary Table 2). A total of 169 unique mutations were identified across the nine CRISPR-targeted sites, ranging from three unique mutations in *hcn4l* to 34 in *si:dkey-65j6.2* (i.e. *KIAA1755*) (Supplementary Table 3). Frameshift mutations were most common (47.9%), followed by missense variants (25.4%), in-frame deletions (14.2%), and synonymous variants (5.9%). Eighty-seven, seventy-two and ten variants were predicted to have a high, moderate, and low impact on protein function, respectively (Supplementary

Table 3). Mutant allele frequencies ranged from 4.1% for *hcn4l* to 93.4% for *neo1b*

(Supplementary Table 4).

Genetic effects on cardiac rhythm and rate

Each additional mutated allele in *rgs6* was associated with a lower heart rate at 2dpf,

independently of HRV (Figure 2, Supplementary Table 5). Larvae with CRISPR-

induced mutations in *rgs6* were shorter at 2 and 5 dpf, and had a smaller dorsal area at

5dpf (Supplementary Figure 2, Supplementary Table 6). These results were mostly

driven by heterozygous larvae, since only one larva carried two mutated *rgs6* alleles

(Supplementary Table 4).

At 2dpf, *hcn4* influenced heart rate independently of HRV when larvae with two

functionally knocked alleles were compared with wildtypes (Supplementary Table 7),

and may affect HRV when adjusting for heart rate (Supplementary Table 5).

Furthermore, *hcn4*-affected embryos had higher odds of a sinoatrial pause or arrest at

2dpf (Table 2, Supplementary Table 8). At 5dpf, *hcn4* affected heart rate as well as

HRV, with effect sizes attenuated but not abolished when adjusting for the other trait

(Supplementary Table 5,7). The odds of a sinoatrial arrest was similar at 5dpf

compared with 2dpf, but did not reach significance at 5dpf (Supplementary Table 8).

Each additional mutated allele in the main transcript of *si:dkey-65j6.2* (i.e.

KIAA1755) tended to be associated with a higher HRV at 2dpf as well as at 5dpf

(Figure 2, Supplementary Table 5). Additionally, *si:dkey-65j6.2*-affected larvae

tended to be smaller at 2 and 5dpf (Supplementary Figure 2, Supplementary Tables 6,

9). Larvae with mutations in *quo* (i.e. the other *KIAA1755* orthologue) were larger at 2

and 5dpf (Supplementary Figure 2, Supplementary Tables 6, 9).

Larvae with two vs. zero functionally knocked *syt10* alleles tended to have a higher

odds of sinoatrial pauses at 2 and 5dpf (Supplementary Tables 8, 10).

Larvae with mutations in *neo1a* tended to be larger at 2dpf (Supplementary Figure

2, Supplementary Tables 6, 9).

Discussion

Large-scale, *in vivo* follow-up studies of candidate genes in GWAS-identified loci remain sparse. Here we present an objective, image-based pipeline to systematically characterize candidate genes for cardiac rhythm, rate, and conduction-related disorders. Using a zebrafish model system, we confirmed the roles of genes previously implicated in heart rate and rhythm (*rgs6* and *hcn4*), provided a novel gene influencing cardiac rhythm (*si:dkey-65j6.2*, i.e. *KIAA1755*), and observed an effect of previously unanticipated genes on early growth and development (*rgs6*, *quo*, *si:dkey-65j6.2*, *neo1a*). Additionally, we confirmed that knockdown of *hcn4* results in sinoatrial pauses[16]. These results add weight to the notion that GWAS-identified common variants for complex traits can flag genes for which rare, detrimental mutations cause severe, early-onset disorders[9,22,23]. We show here that systematic characterization of candidate genes in GWAS-identified loci for complex traits using an image- and CRISPR-Cas9-based zebrafish model system can help identify clinically relevant causal genes, and aid confirmation of the role of such genes in whole-exome sequencing efforts in humans by reducing the multiple testing burden[24].

Regulator of G protein signaling 6 (RGS6) plays a role in the parasympathetic regulation of heart rate[28] and is a negative regulator of muscarinic signaling, thus decreasing HRV to prevent bradycardia. Common variants near *RGS6* have been identified in GWAS for resting heart rate[8,25,26], heart rate recovery after exercise[27] and HRV[4]. An eQTL analysis using GTEx data showed that the minor T-allele in rs4899412 that is associated with lower HRV is also associated with a higher expression of *RGS6* in esophagus and tibial nerve. This is directionally consistent with humans with loss-of-function variants in *RGS6* showing higher heart rate variability[29]. Also in line with this, mice deficient in *Rgs6* (*Rgs6*[-/-]) were previously

characterized by bradycardia, higher HRV and susceptibility to atrioventricular block[30]. In our study, zebrafish embryos with CRISPR-induced mutations in *rgs6* had a lower heart rate at 2dpf, independently of HRV. Hence, our results for heart rate are in line with studies in murine models. However, we did not detect an effect of mutations in *rgs6* on HRV in our study, possibly due to the low mutant allele frequency of *rgs6*.

The locus harboring *HCN4* has been identified in GWAS for HRV[6], heart rate[5] and atrial fibrillation[31]. *HCN4* is arguably the most well characterized gene in the context of cardiac rhythm. It belongs to the family of "funny" channels, aptly named for being activated upon hyperpolarization and non-selective permeability by Na^+ and K^+. It is expressed in the sinoatrial node[32,33] and plays an important role in cardiac pace making[34]. The heart rate lowering agent ivabradine[35] is an open channel blocker of *HCN4*, and reduces cardiovascular and all-cause mortality in heart failure patients[36]. $Hcn4^{-/-}$ mice die prenatally, and analyses of their hearts show no arrhythmias and a lower heart rate compared with wildtype and heterozygotes[34]. In line with this, humans that are heterozygous for an *HCN4* loss of function mutation are typically characterized by bradycardia. Comparably, knockdown of *hcn4* in zebrafish using morpholinos resulted in bradycardia and sinoatrial pauses[16]. We identified nine zebrafish embryos that were compound heterozygous for mutations that result in a functional knock of *hcn4*, and one embryo that was compound heterozygous for such mutations in *hcn4l*. Mutations in both orthologues did not deviate from HWE (Supplementary Table 4). Four of the nine compound heterozygous larvae for *hcn4*, and four of the eight larvae that were heterozygous for mutations in *hcn4* as well as *hcn4l* showed a sinoatrial arrest before the atrial recording was initiated at 2dpf. Zebrafish embryos can survive without a functional heart at this stage, thanks to

adequate tissue oxygenation by diffusion[37]. This allowed us to witness genetically

driven cardiac arrests that would have been lethal in embryos of other species.

However, the direction of effect on heart rate that we observed is unexpected. It most

likely reflects overcompensation by *hcn4l*, or another gene that has been induced by

hcn4 knockout[38]. We cannot rule out that some of the CRISPR-Cas9 induced

mutations represent a gain-of-function either. Our gRNA binds in the first exon of

hcn4, and cuts in the vicinity of the ion-transport domain, as annotated by protein

families (PFAM)[39]. A gain-of-function variant was identified in humans that

increased *HCN4*'s susceptibility to cAMP[40]. In line with this, overexpression of *Hcn4*

in mice resulted in lower HRV[41]. Hence, we may have induced an *hcn4* gain-of-

function mutation in a subset of the larvae.

Non-synonymous SNPs in *KIAA1755* have been identified in GWAS for HRV[6]

and heart rate[5]. In our analysis, *si:dkey-65j6.2*-affected larvae have a higher HRV at 2

and 5dpf, largely independently of heart rate. In humans, the C-allele of the HRV-

associated rs6123471 in the 3' UTR of *KIAA1755* is associated with a lower

expression of *KIAA1755* in brain and aorta (GTEx[42]), a higher HRV[6], and a lower

heart rate[5]. Hence, a higher HRV at 2 and 5dpf for each additional mutated allele in

si:dkey-65j6.2 is directionally consistent with results in humans. Genetic effects on

HRV and heart rate were mutually adjusted in our sensitivity analysis, which may

have revealed the unbiased direction of effect for mutations in *KIAA1755* on HRV. In

other words: the GWAS-identified association of the *KIAA1755* locus - and possibly

other loci - with heart rate may have been driven by HRV. This would explain why

the eleven HRV-associated loci that showed evidence of an association with heart rate

all did so in the expected (i.e. opposite) direction from a phenotypic point of view[6].

KIAA1755 is a previously uncharacterized gene that shows a broad expression pattern,

including different brain regions and the left atrial appendage (GTEx[42]). Future

mechanistic studies are required to distill how *KIAA1755* influences heart rhythm.

SYT10 belongs to the family of synaptogamins: transmembrane proteins that are

involved in regulation of membrane trafficking in neurons[43], which is important for

neurotransmitter release[44]. Variants in or near *SYT10* have previously been associated

with HRV[6] as well as with heart rate [5] and heart rate recovery[27]. Since *SYT10* is

mainly expressed in brain, tibial nerve and pituitary (GTEx[42]). The trend towards

higher odds of sinoatrial pauses in larvae with CRISPR-induced mutations in *syt10*

may be mediated by altered neurotransmitter release.

In addition to the additive analysis, we were able to compare embryos and larvae

with two functionally knocked out alleles with wildtypes for six out of nine orthologs.

These analyses largely confirmed the association of the additive analysis, but also

revealed novel associations. Screening the F_0 or F_2 generation instead would have had

their own advantages and disadvantages. Wu and colleagues described a powerful

approach to efficiently disrupt and analyze F_0 larvae[45]. However, this approach

requires sham injected controls, and is not compatible with multiplexing of many

gRNAs. Hence, we argue that non-conclusive results for a subset of candidate genes

in the screen is not problematic when systematically characterizing a large number of

candidate genes.

The large differences in mutant allele frequency across target genes can be

attributed to several factors. First, the mosaic founders were in-crossed six times,

while random mating was allowed. Hence, we did not ascertain which fish generated

the offspring in each mating. Secondly, while all gRNAs were pre-tested, mutated

alleles that didn't affect the germline were not included in our screen. Thirdly, larvae

that carry embryonic lethal variants were not included in the screen.

Four limitations of our study should be discussed. Firstly, acquiring 30s recordings

implies that false negatives for sinoatrial pauses or arrests are inevitable. We decided

to exclude the 54 embryos and eight larvae with a sinoatrial pause or arrest during

positioning for imaging from the analysis, because the case status of these samples

cannot be confirmed objectively; yet they are not appropriate controls either. This

limitation will have resulted in conservative effect estimates for sinoatrial pauses and

arrests. Secondly, CRISPR guide-RNAs with predicted off-target effects free from

mismatches were avoided. However, two of the selected targets - i.e. for *hcn4* and

si:dkey-65j6.2 - had predicted exonic off-targets with three mismatches at the time we

designed the gRNAs (Supplementary Table 1). Human orthologues of potential off-

target genes *dclk1* and *galnt10* have previously been associated with heart rate

variability-related traits[46] (*dclk1*), as well as with carotid intima-media thickness[47] and

body mass index[48-50] (*galnt10*). However, sequencing larvae at the three predicted off-

target regions did not show any CRISPR-Cas9 induced mutations (Supplementary

Table 2). In line with predictions by Varshney *et al.*, off-target effects are thus

unlikely to have influenced our results[51]. We cannot exclude the possibility of there

being other, non-predicted off-target effects though. Thirdly, we in-crossed mosaic

founder mutants (F_0) and phenotypically screened and sequenced the F_1 generation.

For some genes, this yielded a small number of larvae with 0 or 2 mutated alleles (i.e.

for *rgs6*, *hcn4*, and *hcn4l*), resulting in a lower statistical power to find true genetic

effects - if present - for these genes. In spite of this limitation, we were able to detect

significant effects of mutations in *rgs6* and *hcn4*. Finally, we recorded the atrium

only, to enable a higher frame rate, a higher resolution in time for HRV, and a higher

statistical power to detect small genetic effects on HRV. As a result, any ventricular

abnormalities were not registered, and uncontrolled atrial contractions may thus

reflect atrial fibrillation, premature atrial contractions, high atrial rate, or atrial

tachycardia.

Strengths of our study include its repeated measures design, which enabled us to

capture genetic effects at different stages of early development in zebrafish.

Furthermore, the throughput of the setup allowed us to examine the effect of multiple

genes simultaneously, in an unprecedented sample size for *in vivo* genetic screens.

Our results demonstrate that a large sample size is paramount to robustly detect

genetic effects on complex traits in zebrafish embryos and larvae when screening the

F_1 generation, even for functional knockout mutations. Identifying CRISPR-Cas9-

induced mutations allele-specifically using a custom-written algorithm helped us

classify larvae as heterozygous and compound heterozygous, which in turn helped

pinpoint the true effect of mutations in these genes. Also, our study is based on a

high-throughput imaging approach with objective, automated quantification, as

compared with more manual annotation of previous studies[5]. Finally, for all genes that

showed an effect on HRV, directions of effect were directionally consistent with

eQTL associations in humans. This further emphasizes the strength of our model

system and the robustness of our findings.

In conclusion, our large-scale imaging approach highlights that zebrafish embryos

and larvae can be used for rapid and comprehensive follow-up of GWAS-prioritized

candidate genes. This will likely increase our understanding of the underlying biology

of heart rate and rhythm, and potentially yield novel drug targets.

Methods

Candidate gene selection

Candidate genes in GWAS-identified loci for HRV were identified as described in detail in Nolte et al.[4]. Of the 18 identified candidate genes, six were selected for experimental follow-up. This selection was based on overlap with findings from GWAS for heart rate (*KIAA1755, SYT10, HCN4, GNG11*)[5], as well as with results from eQTL analyses in sinoatrial node and brain (*RGS6*). Additional candidate genes from the same or nearby loci were also selected for experimental follow-up, i.e. *NEO1*, which resides next to *HCN4*. Zebrafish orthologues of the human genes were identified using Ensembl, as well as using a comprehensive synteny search using Genomicus[52] (Supplementary Table 1). Of the selected genes, *GNG11, SYT10* and *RGS6* have one orthologue in zebrafish, and *KIAA1755, HCN4* and *NEO1* each have two orthologues, resulting in a total of nine zebrafish orthologues for six human candidate genes (Table 1).

Mutagenesis

All nine zebrafish genes were targeted together using a multiplexed CRISPR-Cas9 approach described recently[21]. Briefly, guide-RNAs (gRNAs) were selected using ChopChop[53] and CRISPRscan[54] (Supplementary Table 1), based on their predicted efficiency, a moderate to high GC-content, location in an early exon, and absence of predicted off-target effects without mismatches. Oligonucleotides were designed as described[21], consisting of a T7 or SP6 promotor sequence (for gRNAs starting with 'GG' or 'GA', respectively), a gene-specific gRNA-target sequence, and an overlap sequence to a generic gRNA. The gene-specific oligonucleotides were annealed to a

generic 80 bp long oligonucleotide at 98°C for 2 mins, 50°C for 10 mins, and 72°C

for 10 mins. The products were checked for correct length on a 2% agarose gel. The

oligonucleotides were subsequently transcribed *in vitro* using the manufacturer's

instructions (TranscriptAid T7 high yield transcription kit / MEGAscript SP6

transcription kit, both ThermoFisher Scientific, Waltham, USA). The gRNAs were

purified, after which the integrity of the purified gRNAs was examined on a 2%

agarose gel. The zebrafish codon-optimized plasmid pT3TS-nls-zCas9-nls was used

as a template to produce Cas9 mRNA[55]. The plasmid was linearized with Xba1, and

then purified using the Qiaprep Spin Miniprep kit (Qiagen, Hilden, Germany). The

DNA was transcribed using the mMESSAGE mMACHINE T3 Transcription Kit

(ThermoFisher Scientific, Waltham, USA), followed by LiCl precipitation. The

quality of the RNA was confirmed on a 1% agarose gel.

Husbandry & microinjections

A zebrafish line with GFP-labelled α-smooth muscle cells Tg(*acta2*:GFP)[56] was used

to visualize the beating heart. To this end, eggs from an in-cross of Tg(*acta2*:GFP)

fish were co-injected with a mix of Cas9 mRNA (final concentration 150 ng/μl) and

all nine gRNAs (final concentration 25 ng/μl each) in a total volume of 2nL at the

single-cell stage. CRISPR-Cas9 injected embryos were optically screened for the

presence of Tg(*acta2*:GFP) at 2 days post fertilization (dpf), using an automated

fluorescence microscope (EVOS FL Cell imaging system, ThermoFisher Scientific,

Waltham, USA). Tg(*acta2*:GFP) carriers were retained and raised to adulthood in

systems with circulating, filtered and temperature controlled water (Aquaneering, Inc,

San Diego, CA). All procedures and husbandry were conducted in accordance with

Swedish and European regulations, and have been approved by the Uppsala

University Ethical Committee for Animal Research (C142/13 and C14/16).

Experimental procedure imaging

The mosaic founder mutants (F_0 generation) were only used for reproduction. After

crossing the founder mutants, F_1 embryos were used for experiments. To reach the

experimental sample size, founder mutants were incrossed six times. At all occasions,

mating was entirely random. Eggs were collected after F_0 fish were allowed to

reproduce for 45 mins to minimize variation in developmental stage. Fertilized eggs

were placed in an incubator at 28.5°C. At 1dpf, embryos were dechorionated using

pronase (Roche Diagnostics, Mannheim, Germany).

At 2dpf, embryos were removed from the incubator and allowed to adapt to controlled

room temperature (21.5 °C) for 20 mins. Individual embryos were exposed to 100

μg/ml Tricaine (MS-222, Sigma-Aldrich, Darmstadt, Germany) for 1 min before

being aspirated, positioned in the field of view of a fluorescence microscope, and

oriented dorsally using a Vertebrate Automated Screening Technology (VAST)

BioImager (Union Biometrica Inc., Geel, Belgium). We subsequently acquired twelve

whole-body images, one image every 30 degrees of rotation, using the camera of the

VAST BioImager to quantify body length, dorsal and lateral surface area and volume,

as well as the presence or absence of cardiac edema. The VAST BioImager then

positioned and oriented the larva to visualize the beating atrium and triggered the

upright Leica DM6000B fluorescence microscope to start imaging using a HCX APO

L 40X/0.80 W objective and L5 ET, k filter system (Micromedic AB, Stockholm,

Sweden). Images of the beating atrium were acquired for 30s at a frame rate of 152

frames/s using a DFC365 FX high-speed CCD camera (Micromedic AB, Stockholm,

Sweden). After acquisition, the larvae were dispensed into a 96-well plate, rinsed

from tricaine, and placed back into the incubator. The procedure was repeated at 5dpf

(i.e. the larval stage), to allow genetic effects that influence HRV and heart rate

differently at different stages of development to be captured[57]. After imaging at 5dpf,

the larvae were once again dispensed into 96-well plates, sacrificed, and stored at -

80°C for further processing.

Quantification of cardiac traits and body size

A custom-written MATLAB script was used to convert the images acquired by the

CCD camera into quantitative traits. To acquire the heart rate, each frame of the

sequence was correlated with a template frame. The repeating pattern yields a

periodic graph from the correlation values and by detecting the peaks in the graph we

can assess the heart rate. The template frame should represent one of the extreme

states in the cardiac cycle, i.e. end-systole or end-diastole. To detect these frames, we

examined the correlation between the first 100 frames. The combination of frames

that showed the lowest correlation corresponded to the heart being in opposite states.

One of these frames was chosen as the template [58].

This numeric information was subsequently used to quantify: 1) heart rate as the

inverse of RR-interval; 2) the standard deviation of the normal-to-normal RR interval

(SDNN); and 3) the root mean square of successive heart beat interval differences

(RMSSD). Finally, a graph of pixel changes over time was generated across the 30s

recording to help annotate the script's performance. The files containing the inter-

beat-intervals were used to objectively quantify sinoatrial pauses (i.e. the atrium stops

beating for longer than 3x the median inter-beat-interval of the larva, Supplementary Movie 1) and sinoatrial arrests (i.e. the atrium stops beating for longer than 2s, Supplementary Movie 2) using a custom-written Stata script. The graphs of pixel changes over time were also used to identify larvae with other abnormalities in cardiac rhythm. Such abnormalities were annotated as: uncontrolled atrial contractions (Supplementary Movie 3); abnormal cardiac morphology (i.e. a tube-like atrium, Supplementary Movie 4); or impaired cardiac contractility (i.e. a vibrating rather than a contracting atrium, Supplementary Movie 4). These phenotypes were annotated independently by two investigators (B.v.d.H. and M.d.H.), resulting in an initial concordance rate >90%. Discrepancies in annotation were discussed and re-evaluated to reach consensus.

Bright-field images of the larvae were used to assess body length, dorsal and lateral surface area, and volume. Images were automatically segmented and quantified using a custom-written CellProfiler[59] pipeline, followed by manual annotation for segmentation quality. Larvae with suboptimal segmentation quality due to the presence of an air-bubble, a bent body, an incomplete rotation within the capillary, partial capturing of the larva, or an over-estimation of size were replaced by images with a 180° difference in rotation, or excluded if that image was also sub-optimally segmented. The larva was excluded from the analysis for body volume if more than four of the 12 images had a bad segmentation. Imaging, image quantification and image quality control were all performed blinded to the sequencing results.

Quality control of phenotype data

A series of quality control steps was performed to ensure only high-quality data was included in the genetic association analysis (Supplementary Fig. 1). First, graphs indicating that one or more true beats were missed by the script were removed from the analysis *a priori* (Supplementary Fig. 1). Second, embryos and larvae showing uncontrolled atrial contractions, sinoatrial pauses or arrests, edemas, abnormal morphology and/or reduced contractility, and embryos/larvae annotated to have a sinoatrial pause while positioning before video acquisition were excluded from the analyses for HRV and heart rate (Supplementary Fig. 1). Third, genetic effects on sinoatrial pauses and arrests were examined at 2dpf and 5dpf. For the remainder of the cardiac abnormalities, too few cases were observed to justify a separate association analysis. Furthermore, embryos and larvae that showed a sinoatrial pause or arrest in between positioning and video acquisition, but not during the recording, were excluded from the analysis for sinoatrial pauses and arrests, since we cannot ascertain case status in the same rigorous manner for such samples, but they are not appropriate controls either.

Sample preparation for sequencing

After imaging at 5dpf, larvae were sacrificed and DNA was extracted by exposure to lysis buffer (10mM Tris-HCl pH8, 50mM KCl, 1mM EDTA, 0.3% Tween 20, 0.3% Igepal) and proteinase K (Roche Diagnostics, Mannheim, Germany) for 2 h at 55°C, followed by 10 min at 95°C to deactivate the proteinase K. Gene-specific primers (150bp-300bp) amplifying gRNA-targeted and putative off-target regions in *dclk1b* and both *galnt10* orthologues (Supplementary Table 1) were distilled from

ChopChop[53] and Primer3[60], and Illumina adaptor-sequences were added.

Additionally, we included 96 uninjected Tg(*acta2*:*GFP*) larvae for sequencing across

all gene-specific regions to adjust for natural occurring polymorphisms. The first PCR

was conducted by denaturation at 98°C for 30s; amplification for 35 cycles at 98°C

for 10s, 62°C for 30s and 72°C for 30s; followed by a final extension at 72°C for 2

mins. Amplified PCR products were cleaned using magnetic beads (Mag-Bind PCR

Clean-up Kit, Omega Bio-tek Inc. Norcross, GA). The purified products were used as

a template for the second PCR, in which Illumina Nextera DNA library sequences

were attached to allow multiplexed sequencing of all CRISPR-targeted sites across

384 larvae in a single lane. The second PCR amplification was performed by

denaturation at 98°C for 30s; amplification for 25 cycles at 98°C for 10s, 66°C for 30s

and 72°C for 30s; followed by a final extension at 72°C for 2 mins. Products were

then purified using magnetic beads. All liquid handling was performed using a

Hamilton Nimbus robot equipped with a 96-head (Hamilton robotics, Kista, Sweden).

Samples were pooled and sequenced in a single lane on a MiSeq (300 bp paired-end,

Illumina Inc., San Diego, CA) at the National Genomics Infrastructure, Sweden.

Processing of sequencing data

A custom-written bioinformatics pipeline was developed in collaboration with the

National Bioinformatics Infrastructure Sweden to prepare .fastq files for analysis.

First, a custom-written script was used to de-multiplex the .fastq files by gene and

well. PEAR[61] was then used to merge paired-end reads, followed by removal of low-

quality reads using FastX[62]. The reads were then mapped to the wildtype zebrafish

genome (Zv11) using STAR[63]. Next, we converted files from .sam to .bam format

using samtools[64], after which variants - mostly indels and SNVs - were called allele

specifically using a custom-written variant calling algorithm in R (Danio rerio

Identification of Variants by Haplotype - DIVaH). A summary of all unique

sequences identified for the orthologues and their respective alignment report string

(Concise Idiosyncratic Gapped Alignment Report (CIGAR)) is shown in

Supplementary Table 2. All unique variants (Supplementary Table 3) located within

±30bps of the CRISPR-targeted sites that were identified across the two alleles were

subsequently pooled, and used for functional annotation using Ensembl's VEP [65].

Naturally occuring variants based on sequencing of uninjected Tg*(acta2:gfp)* larvae

were excluded. Noteworthy, all variants per transcript get annotated individually,

rather than on a per transcript basis. In absence of a continuous score provided by

variant prediction algorithms, we attributed a score of 0.33, 0.66 and 1, and that pilot

experiments using *hcn4* and heart rate suggested that this was a more powerful

approach than not weighting.

Transcript-specific dosage scores were then calculated by retaining the variant with

the highest predicted impact on protein function, and assigning it a score of 0 (no

effect), 0.2 (modifier variant), 0.33 (low), 0.66 (moderate) or 1 (high). Transcript-

specific dosage scores were subsequently calculated by summing the allele-specific

scores at each embryo and target site. Since all transcripts within a target site were

affected virtually identically, we only used the main transcript of each target site for

the genetic association analysis.

Most larvae had successfully called sequences in all nine CRISPR-targeted sites.

Larvae with more than three missing sequences were excluded from the genetic

association analysis. For the remaining larvae, the mean dosage of the transcript was

imputed for missing calls. For *neo1b*, calling failed in 78 larvae (Supplementary Table

4). The imputed mean dosage for the main transcript of *neo1b* was still included in the

genetic association analysis, since the mutant allele frequency in successfully called

larvae was high (i.e. 0.934), and using an imputed dosage was anticipated to influence

the results less than to discard the gene while it had been targeted. However, the

results for *neo1b* should be interpreted in light of its call rate. The mutant allele

frequency was low for *hcn4*, *hcn4l* and *rgs6*. All three putative off-targets regions

(*dclk1b*, both *galnt10* orthologues, Supplementary Table 1) were classified as

wildtype across all larvae and had no CRISPR-Cas9 induced mutations within ±30bps

of the cut-site (Supplementary Table 2).

Statistical analysis

The standard deviation of NN-intervals (SDNN) and the root mean square of

successive differences (RMSSD) were strongly correlated at 2dpf (r^2=0.78) and 5dpf

(r^2=0.83), so a composite endpoint 'HRV' was calculated as the average of SDNN

and RMSSD. In larvae free from sinoatrial pauses and arrests, abnormal cardiac

morphology, impaired cardiac contractility, and edema HRV and heart rate at 2dpf

(n=279) or 5dpf (n=293) were inverse-normally transformed to ensure a normal

distribution, and to allow comparison of effect sizes across traits. Genetic effects on

HRV and heart rate were subsequently examined using hierarchical linear models at

2dpf and 5dpf separately (Stata's xtmixed), mutually adjusted for the time of imaging

(fixed factors), and with larvae nested in six batches (random factor with fixed slope).

Embryos and larvae in which a sinoatrial pause and/or arrest was observed in between

positioning and recording were excluded from the analysis. In a sensitivity analysis,

we mutually adjusted for the other outcome (i.e. heart rate and HRV).

In further analyses, genetic effects were examined for sinoatrial pause and arrest at

2dpf, as well as sinoatrial pause at 5dpf using a logistic regression analysis. Genetic

effects on body size were also examined, using hierarchical linear models on inverse-

normally transformed outcomes.

For dichotomous as well as continuous outcomes, genetic effects were examined

using an additive model, with dosage scores for all target sites as independent

exposures in the same model, i.e. mutually adjusted for effects of mutations in the

other orthologues. Additionally, larvae with two functionally knocked alleles were

compared with larvae free from CRISPR-induced mutations for six out of nine

orthologues. Missing genotypes were imputed at the mean in this analysis. For each

outcome, association analyses were examined for the main transcript of the

orthologue. *P*-values <0.05 were considered to reflect statistically significant effects.

All statistical analyses were performed using Stata MP 14.2.

Acknowledgments

The computations were performed on resources provided by SNIC through Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under Project SNIC b2015283. The authors would like to acknowledge support from Science for Life Laboratory, the National Genomics Infrastructure (NGI) and UPPMAX for providing assistance in massive parallel sequencing and computational infrastructure. Support from the National Bioinformatics Infrastructure Sweden (NBIS) is also gratefully acknowledged. Constructive discussions with the Genome Engineering Zebrafish (GEZ) facility are also acknowledged.

Sources of funding

MdH is a fellow of the Swedish Heart-Lung Foundation (20170872) and is supported by project grants from the Swedish Heart-Lung Foundation (20140543, 20170678), the Swedish Research Council (2015-03657), and NIH/NIDDK (R01DK106236, R01DK107786, U01DK105554).

Disclosures

None.

REFERENCES

Thayer, J. F., Yamamoto, S. S. & Brosschot, J. F. The relationship of autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *Int. J. Cardiol.* **141**, 122-131, (2010).

de Bruyne, M. C. *et al.* Both decreased and increased heart rate variability on the standard 10-second electrocardiogram predict cardiac mortality in the elderly: the Rotterdam Study. *Am. J. Epidemiol.* **150**, 1282-1288, (1999).

Dekker, J. M. *et al.* Heart rate variability from short electrocardiographic recordings predicts mortality from all causes in middle-aged and elderly men - The zutphen study. *Am. J. Epidemiol.* **145**, 899-908, (1997).

Nolte, I. M. *et al.* Genetic loci associated with heart rate variability and their effects on cardiac disease risk. *Nature communications* **8**, 15805, (2017).

den Hoed, M. *et al.* Identification of heart rate-associated loci and their effects on cardiac conduction and rhythm disorders. *Nat. Genet.* **45**, 621-631, (2013).

Nolte, I. M. *et al.* Genetic loci associated with heart rate variability and their effects on cardiac disease risk. *Nature communications* **8**, 15805, (2017).

Poon, K. L. & Brand, T. The zebrafish model system in cardiovascular research: A tiny fish with mighty prospects. *Glob Cardiol Sci Pract* **2013**, 9-28, (2013).

Eppinga, R. N. *et al.* Identification of genomic loci associated with resting heart rate and shared genetic predictors with all-cause mortality. *Nat. Genet.* **48**, 1557-1563, (2016).

Arking, D. E. *et al.* Genetic association study of QT interval highlights role for calcium signaling pathways in myocardial repolarization. *Nat. Genet.* **46**, 826-836, (2014).

Lieschke, G. J. & Currie, P. D. Animal models of human disease: zebrafish swim into view. *Nature reviews. Genetics* **8**, 353-367, (2007).

MacRae, C. A. & Peterson, R. T. Zebrafish as tools for drug discovery. *Nature reviews. Drug discovery* **14**, 721-731, (2015).

Liu, C. C., Li, L., Lam, Y. W., Siu, C. W. & Cheng, S. H. Improvement of surface ECG recording in adult zebrafish reveals that the value of this model exceeds our expectation. *Sci. Rep.* **6**, 25073, (2016).

Harrington, J. K., Sorabella, R., Tercek, A., Isler, J. R. & Targoff, K. L. Nkx2.5 is essential to establish normal heart rate variability in the zebrafish embryo. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **313**, R265-r271, (2017).

Arnaout, R. *et al.* Zebrafish model for human long QT syndrome. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11316-11321, (2007).

Sah, R. *et al.* Ion channel-kinase TRPM7 is required for maintaining cardiac automaticity. *Proc. Natl. Acad. Sci. U. S. A.* **110**, E3037-3046, (2013).

Jou, C. J. *et al.* A Functional Assay for Sick Sinus Syndrome Genetic Variants. *Cell. Physiol. Biochem.* **42**, 2021-2029, (2017).

Pardo-Martin, C. *et al.* High-throughput hyperdimensional vertebrate phenotyping. *Nature communications* **4**, 1467, (2013).

Early, J. J. *et al.* An automated high-resolution in vivo screen in zebrafish to identify chemical regulators of myelination. *eLife* **7**, (2018).

Jarque, S. *et al.* An automated screening method for detecting compounds with goitrogenic activity using transgenic zebrafish embryos. *PLoS One* **13**, e0203087, (2018).

20   Howe, K. *et al.* The zebrafish reference genome sequence and its relationship
to the human genome. *Nature* **496**, 498-503, (2013).
21   Varshney, G. K. *et al.* A high-throughput functional genomics workflow based
on CRISPR/Cas9-mediated targeted mutagenesis in zebrafish. *Nat. Protoc.* **11**,
2357-2375, (2016).
22   Flannick, J., Johansson, S. & Njolstad, P. R. Common and rare forms of
diabetes mellitus: towards a continuum of diabetes subtypes. *Nat. Rev.*
Endocrinol. **12**, 394-406, (2016).
23   Ganna, A. *et al.* Quantifying the Impact of Rare and Ultra-rare Coding
Variation across the Phenotypic Spectrum. *Am. J. Hum. Genet.* **102**, 1204-
1211, (2018).
24   Flannick, J. A. *et al.* Genetic discovery and translational decision support from
exome sequencing of 20,791 type 2 diabetes cases and 24,440 controls from
five ancestries. *bioRxiv*, (2018).
25   Ramirez, J. *et al.* Thirty loci identified for heart rate response to exercise and
recovery implicate autonomic nervous system. *Nature communications* **9**,
1947, (2018).
26   Kerr, K. F. *et al.* Genome-wide association study of heart rate and its
variability in Hispanic/Latino cohorts. *Heart Rhythm* **14**, 1675-1684, (2017).
27   Verweij, N., van de Vegte, Y. J. & van der Harst, P. Genetic study links
components of the autonomous nervous system to heart-rate profile during
exercise. *Nature communications* **9**, 898, (2018).
28   Yang, J. *et al.* RGS6, a modulator of parasympathetic activation in heart. *Circ.*
Res. **107**, 1345-1349, (2010).
29   Posokhova, E. *et al.* Essential role of the m2R-RGS6-IKACh pathway in
controlling intrinsic heart rate variability. *PLoS One* **8**, e76973, (2013).
30   Wydeven, N., Posokhova, E., Xia, Z., Martemyanov, K. A. & Wickman, K.
RGS6, but not RGS4, is the dominant regulator of G protein signaling (RGS)
modulator of the parasympathetic regulation of mouse heart rate. *J. Biol.*
Chem. **289**, 2440-2449, (2014).
31   Ellinor, P. T. *et al.* Meta-analysis identifies six new susceptibility loci for
atrial fibrillation. *Nat. Genet.* **44**, 670-675, (2012).
32   Shi, W. *et al.* Distribution and prevalence of hyperpolarization-activated
cation channel (HCN) mRNA expression in cardiac tissues. *Circ. Res.* **85**, e1-
6, (1999).
33   Ludwig, A. *et al.* Two pacemaker channels from human heart with profoundly
different activation kinetics. *EMBO J.* **18**, 2323-2329, (1999).
34   Stieber, J. *et al.* The hyperpolarization-activated channel HCN4 is required for
the generation of pacemaker action potentials in the embryonic heart. *Proc.*
Natl. Acad. Sci. U. S. A. **100**, 15235-15240, (2003).
35   Bucchi, A. *et al.* Identification of the molecular site of ivabradine binding to
HCN4 channels. *PLoS One* **8**, e53132, (2013).
36   Swedberg, K. *et al.* Ivabradine and outcomes in chronic heart failure (SHIFT):
a randomised placebo-controlled study. *Lancet* **376**, 875-885, (2010).
37   Burggren, W. W. & Pinder, A. W. Ontogeny of cardiovascular and respiratory
physiology in lower vertebrates. *Annu. Rev. Physiol.* **53**, 107-135, (1991).
38   Rossi, A. *et al.* Genetic compensation induced by deleterious mutations but
not gene knockdowns. *Nature* **524**, 230-233, (2015).
39   El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic Acids*
Res. **47**, D427-d432, (2019).

| 650 | 40 | Baruscotti, M. *et al.* A gain-of-function mutation in the cardiac pacemaker HCN4 channel increasing cAMP sensitivity is associated with familial Inappropriate Sinus Tachycardia. *Eur. Heart J.* **38**, 280-288, (2017). |

650 40 Baruscotti, M. *et al.* A gain-of-function mutation in the cardiac pacemaker HCN4 channel increasing cAMP sensitivity is associated with familial Inappropriate Sinus Tachycardia. *Eur. Heart J.* **38**, 280-288, (2017).

651
652

41 Kozasa, Y. *et al.* HCN4 pacemaker channels attenuate the parasympathetic response and stabilize the spontaneous firing of the sinoatrial node. *J. Physiol.* **596**, 809-825, (2018).

42 Battle, A., Brown, C. D., Engelhardt, B. E. & Montgomery, S. B. Genetic effects on gene expression across human tissues. *Nature* **550**, 204-213, (2017).

43 Sudhof, T. C. Synaptotagmins: why so many? *J. Biol. Chem.* **277**, 7629-7632, (2002).

44 Chapman, E. R. How does synaptotagmin trigger neurotransmitter release? *Annu. Rev. Biochem.* **77**, 615-641, (2008).

45 Wu, R. S. *et al.* A Rapid Method for Directed Gene Knockout for Screening in G0 Zebrafish. *Dev. Cell* **46**, 112-125.e114, (2018).

46 Newton-Cheh, C. *et al.* Genome-wide association study of electrocardiographic and heart rate variability traits: the Framingham Heart Study. *BMC Med. Genet.* **8 Suppl 1**, S7, (2007).

47 Dong, C. *et al.* Genetic variants in LEKR1 and GALNT10 modulate sex-difference in carotid intima-media thickness: a genome-wide interaction study. *Atherosclerosis* **240**, 462-467, (2015).

48 Locke, A. E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**, 197-206, (2015).

49 Justice, A. E. *et al.* Genome-wide meta-analysis of 241,258 adults accounting for smoking behaviour identifies novel loci for obesity traits. *Nature communications* **8**, 14977, (2017).

50 Ng, M. C. Y. *et al.* Discovery and fine-mapping of adiposity loci using high density imputation of genome-wide association studies in individuals of African ancestry: African Ancestry Anthropometry Genetics Consortium. *PLoS genetics* **13**, e1006719, (2017).

51 Varshney, G. K. *et al.* High-throughput gene targeting and phenotyping in zebrafish using CRISPR/Cas9. *Genome Res.* **25**, 1030-1042, (2015).

52 Louis, A., Nguyen, N. T., Muffato, M. & Roest Crollius, H. Genomicus update 2015: KaryoView and MatrixView provide a genome-wide perspective to multispecies comparative genomics. *Nucleic Acids Res.* **43**, D682-689, (2015).

53 Labun, K., Montague, T. G., Gagnon, J. A., Thyme, S. B. & Valen, E. CHOPCHOP v2: a web tool for the next generation of CRISPR genome engineering. *Nucleic Acids Res.* **44**, W272-276, (2016).

54 Moreno-Mateos, M. A. *et al.* CRISPRscan: designing highly efficient sgRNAs for CRISPR-Cas9 targeting in vivo. *Nature methods* **12**, 982-988, (2015).

55 Jao, L. E., Wente, S. R. & Chen, W. Efficient multiplex biallelic zebrafish genome editing using a CRISPR nuclease system. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 13904-13909, (2013).

56 Whitesell, T. R. *et al.* An alpha-smooth muscle actin (acta2/alphasma) zebrafish transgenic line marking vascular mural cells and visceral smooth muscle cells. *PLoS One* **9**, e90590, (2014).

57 Hou, J. H., Kralj, J. M., Douglass, A. D., Engert, F. & Cohen, A. E. Simultaneous mapping of membrane voltage and calcium in zebrafish heart in vivo reveals chamber-specific developmental transitions in ionic currents. *Front. Physiol.* **5**, 344, (2014).

58   Niazi, M. K. K., Nilsson, M. F., Danielsson, B. R. & Bengtsson, E. Fully
Automatic Heart Beat Rate Determination in Digital Video Recordings of Rat
Embryos. (2009).

59   Lamprecht, M. R., Sabatini, D. M. & Carpenter, A. E. CellProfiler: free,
versatile software for automated biological image analysis. *Biotechniques* **42**,
71-75, (2007).

60   Koressaar, T. & Remm, M. Enhancements and modifications of primer design
program Primer3. *Bioinformatics* **23**, 1289-1291, (2007).

61   Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate
Illumina Paired-End reAd mergeR. *Bioinformatics* **30**, 614-620, (2014).

62   Pearson, W. R., Wood, T., Zhang, Z. & Miller, W. Comparison of DNA
sequences with protein sequences. *Genomics* **46**, 24-36, (1997).

63   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
29, 15-21, (2013).

64   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**, 2078-2079, (2009).

65   McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**,
122, (2016).

718
719

Eight loci associated with HRV[1].
18 genes prioritized for functional follow-up[1].

Six human genes selected for follow-up,
represented by nine orthologues in zebrafish.

CRISPR-gRNA design,
synthesis and efficiency testing.

Raising of CRISPR-Cas9 induced mutant founder fish (F_0).
Crossing of F_0-Generation for experiments.

Acquistion of 30 s videos of the atrium of
381 embryos (2dpf) & larvae (5dpf).

Automatic quantification of 30 s videos
using a custom-written MatLab script.

Paired-end sequencing of CRISPR-targeted
sites using a MiSeq.

Splitting, merging, QC-ing and aligning
of sequencing data.

Transcript-specific variant calling and
annotation using Ensembl's VEP.

Genetic association analysis using hierarchical linear
models and multiple logistic regression analyses.

Figure 1: Experimental pipeline describing the workflow from candidate gene selection to trait association. dpf = days post fertilization. VEP = Variant Effect Predictor.
1: Nolte et al. 2017

Figure 2: Effect of mutations in candidate genes on heart rhythm and rate at 2 (n=234) and 5dpf (n=285). Dots and whiskers show the effect size and its 95% confidence interval for each additional mutated allele (solid line), weighted by its predicted effect on protein function. Dotted bars indicate the results of comparing larvae with two functionally knocked alleles with wildtype larvae. Effects were adjusted for the number of mutated alleles in the other genes, as well as time of day (fixed factor), with larvae nested in batches (random factor).

Table 1: Selected candidate genes and their zebrafish orthologues in GWAS-identified heart rate variability associated loci

rsID	Chr	Human gene	ENSG	Zebrafish orthologue	ENSDARG
rs4262, rs180238	7	*GNG11*	ENSG00000127920	*gngt1*	ENSDARG00000035798
rs7980799, rs1351682, rs1384598	12	*SYT10*	ENSG00000110975	*syt10*	ENSDARG00000045750
rs4899412, rs2052015, rs2529471, rs36423	14	*RGS6*	ENSG00000182732	*rgs6*	ENSDARG00000015627
rs2680344	15	*HCN4*	ENSG00000138622	*hcn4*	ENSDARG00000061685
				hcn4l	ENSDARG00000074419
rs1812835	15	*NEO1*	ENSG00000067141	*neo1a*	ENSDARG00000102855
				neo1b	ENSDARG00000075100
rs6123471	20	*KIAA1755*	ENSG00000149633	*quo*	ENSDARG00000073684
				si:dkey-65j6.2	ENSDARG00000103586

rsID refers to the lead SNP(s) of the GWAS-identified loci (Nolte et. al, 2017). Multiple independent SNPs per locus can be present.

Table 2: Effects of mutations in *hcn4* on sinoatrial pauses and arrests at 2dpf

Age	Gene	Outcome	n total	n cases	OR	LCI	UCI	P
2	*hcn4*	SA arrest	248	36	2.644	1.402	4.986	2.66E-03
		SA pause	297	39	2.749	1.461	5.171	1.71E-03

Associations of sinoatrial pauses and arrests with the number of mutated alleles in *hcn4*, weighted by their predicted effect on protein function. Associations were analyzed using logistic regression, and were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors.

381 embryos imaged at 2dpf

Secondary trait analysis:
Cardiac traits (left), body size analysis (right)

Sinoatrial pauses (n=39) & arrests (n=36),
uncontrolled atrial contraction (n=1),
edema (n=15), fish moved during
recording/missed beats (n=38),
sinoatrial event before recording (n=54)

Embryos excluded due to
suboptimal segmentation
and/or presence of edema:
length (n=46), dorsal (n=56),
lateral (n=78), volume (n=79)

Primary analysis:
234 embryos included in HRV and
heart rate analysis at 2dpf

381 embryos imaged at 2dpf

Larvae excluded due to
technical issues* (n=41), death (n=3)
or larva not detected by VAST (n=11)

326 larvae imaged at 5dpf

Secondary trait analysis:
Cardiac traits (left), body size analysis (right)

Sinoatrial pauses (n=9) & arrests (n=3),
uncontrolled atrial contraction (n=9),
abnormal morphology/reduced contractility (n=6)
edema (n=14), missed beats (n=1),
sinoatrial event before recording (n=8)

Larvae excluded due to
suboptimal segmentation
and/or presence of edema:
length (n=84), dorsal (n=93),
lateral (n=70), volume (n=128)

Primary analysis:
285 larvae included in HRV and
heart rate analysis at 5dpf

Supplementary Figure 1: Flow chart showing the number of included larvae in the analysis and reasons for exclusions. Embryos/larvae can show more than one phenotype. *Technical issues refers to not being able to image due to microscope complications. VAST: Vertebrate automated screening technology.

Supplementary Figure 2: Effect of mutations in candidate genes on length, dorsal and lateral area, and volume at 2 (upper part) and 5dpf (lower part). Dots and whiskers show the effect size and its 95% confidence interval for each additional mutated allele (solid line), weighted by its predicted effect on protein function. Dotted bars indicate the results of comparing larvae with two functionally knocked alleles with wildtype larvae. Effects were adjusted for the number of mutated alleles in the other genes, as well as time of day (fixed factor), with larvae nested in batches (random factor).

Supplementary Table 1: Overview of zebrafish orthologues and designed gRNAs

Zebrafish orthologue	ENSDARG	Conservation Zebrafish - Human	Conservation Zebrafish - Spotted gar	Main transcript	Target sequence gRNA	Exon target (transcript name)	CRISPR-scan rank	ChopChop rank	Activity test rank	Predicted off-targets	Nr of mismatches	Genomic location of potential off-target 1	ENSDARG (gene-ID) of potential off-target 1, Feature	ENSG (gene-ID) of potential off-target 1	Genomic location of potential off-target 2	ENSDARG (gene-ID) of potential off-target 2, Feature	ENSG (gene-ID) of potential off-target 2	Primer forward	Primer reverse	Amplicon size
gngt1	ENSDARG00000035798	17	21	ENSDART00000051950	gGACCTGGATAAGGCGAAGA	2/3 (gngt1-001); 1/2 (gngt1-002)	62	4/13 (yellow)	moderate	1	3	chr25:17488676	ENSDARG00000051959 (*zgc:66449*), intron (3/6), intron (4/4)	no human orthologue				GTCCATTTTCCAGAGTGGAGAC	gCCCATAACACAATTGCTTACC	207
syt10	ENSDARG00000045750	0	1	ENSDART00000136049	GGCTGATGCCGTCCTCTGTG	1/7 (syt10-001); 1/7 (syt10-201)	36	6/129 (green)	low	0								TAGGATATGGACTTGGACGCAT	TTACCTGGTATGTTGCTCTCCA	227
rgs6	ENSDARG00000015627	9	20	ENSDART00000135513	GGCCTCTCAGGACCAAGG	1/17 (rgs6-201); 2/18 (rgs6-001)	62	1/107 (green)	low	0								AAGGCTAATTCATTGTCTGGGA	GCATGTCAAATGTAAAGGCAAA	169
hcn4	ENSDARG00000061685	6	3	ENSDART00000136140	GGAGAGCCCTCCTGGAGCCG	1/8 (hcn4-001)	59	11/280 (green)	high	2	3/3	chr21:45844184	ENSDARG00000103825 (*galnt10*), exon (1/3)	ENSG00000164574 (GALNT10)	chrUn_KN150614v1:3424	ENSDARG00000100453 (*galnt10*), exon (6/8)	ENSG00000164574 (GALNT10)	CCTTTCGTCCATCACCAGTC	CCGCTTGATAATACCTCTCGTC	230
hcn4l	ENSDARG00000074419	3	3	ENSDART00000088249	gGCCTGGATAGTGTTTAACG	2/8 (hcn4l-201); 1/7 (hcn4l-001)	/	2/191 (green)	low	0								TTACTGGGACCTGATCATGCTT	AAGATGTAATCCACAGGGATGG	235
neo1a	ENSDARG00000102855	1	5	ENSDART00000158160	GGAGCCGTCGGATACACTAG	1/4 (neo1a-001); 2/29 (neo1a-201); 2/30 (neo1a-202)	48	2/55 (green)	very high	0								TTCTGTGTCTCCACAGGATCAG	CCTCATCGGGTTTATTGTGTTT	215
neo1b	ENSDARG00000075100	2	6	ENSDART00000115280	GGACAGAGATGCTCGGCCTG	3/29 (neo1b-201)	58	18/354 (green)	very high	0								CAGCTCTCTCTCTCATTGGCTT	TATGGTGCAACGGTAGAGTCC	206
quo	ENSDARG00000073684	2	5	ENSDART00000109679	GGAGATCCGGTTGCCCTGTG	2/22 (quo-001); 3/23 (quo-201)	83	12/331 (green)	very high	0								ATGACCGTAAGTCAAGGGAAAA	TGAATCTTGTCTAATGGCGTTG	153
si:dkey-65j6.2	ENSDARG00000103586	3	5	ENSDART00000158832	GGACGGTTTGGCTCCAGCAG	1/23 (si:dkey-65j6.2-201); 3/25 (si:dkey-65j6.2-001)	59	14/303 (green)	very high	2	3/3	chr15:32969618	ENSDARG00000104664 (*dclk1b*), exon (3/16), exon (4/17)	ENSG00000133083 (DCLK1)	chr22:28819880	ENSDARG00000071083 (*si:dkeyp-34c12.1*), Intron (7/7), Exon (8/9)	no human orthologue	AGCCATTACACCCAAAGAAAAA	GGGAATCATTTCGAAGTAGCTG	152

Table showing zebrafish genes, main transcript used for the analysis and gRNAs; rsID refers to the lead SNP(s) of the GWAS-identified loci (Nolte et. al, 2017), multiple independent SNPs per locus can be present; For simplicity, *quo* is referred to as *kiaa1755_1*, and *si:dkey-65j6.2* as *kiaa1755_2*; Conservation refers to the number of genes that are preserved in the same locus across species according to Genomicus; A small "g" in the gRNA target sequence indicates a manually introduced nucleotide change to guanine; Exon target lists the targeted transcripts and their respective exons; ChopChop rank indicates the rank of the gRNA chosen by ChopChop and its rating (green being a "good"-predicted target, yellow an average target, details can be found on the ChopChop homepage; Activity test rank quantification: Low: 8 of 8 larvae showed wildtype sequence and fewer than 4 of 8 also contained an indel sequence, Moderate: 8 of 8 larvae showed the wildtype sequence and >4 of 8 also contained an indel sequence, High: 8 of 8 larvae showed wildtype as well as indel sequences, Very high: Fewer than 4 of 8 larvae showed wildtype sequence and all larvae showed indels; ENSDARG of potential off-target refers to the ENSEMBL gene-ID of the potential off-target and where the potential target is located (Feature), zgc:66449 and si:dkeyp-34c12.1 have no human orthologue, whereas galnt10 and dckl1b may exert an effect on heart rate and/or rhythm.

Supplementary Table 2: Unique Cigars

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S119M	498	336	381
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	207M	517	267	372
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S69M1D3S46M	544	69	405
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG ATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	151M6D50M	500	63	364
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG ATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S63M6D50M	584	13	479
gngt1	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATATGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGATGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	158M1S48M	504	4	221
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S68M3D48M	606	3	119
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGTAACCCAGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S72M4I3S44M	526	2	143
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATAATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	45M1S41M1S32M1S36M1D3S46M	48	1	-
	ATCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAACATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGTTTTGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	1S44M1S41M1S69M1D3S46M	10	1	-
	CTCATCGTAAGCAGCAGTAAGTTTGCTTATTTCCAGTATATTGAATATCAAATATAATCCATTCATTCAACACAACAT GTAAATGATGTGATTGTTTGATTTCATGCAGAAAATGCCGATCATAGATGTAGAAAATATGACTGACCTGGATAAG GCGAAGATGGAAGTAACCCAGCTCAAAACTGAGGTGAAGCTAGAAAGGGCGAA	87M1S119M	3	1	-
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACAGAGGACGGCATCAGCCTGTGCCCAAA GGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	227M	1024	518	644
syt10	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACAAGAGGACGGCATCAGCCTGTGCCAA AGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCA C	126M1I101M	735	46	462
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGACGGCATCAGCCTGTGCCCAAAGGGCTCTC CAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	122M8D97M	606	45	397

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTACAGCTGATGCGTACGGGAATGGAGAGGACG GCATCAGCCTGTGCCAAAGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAAT GCGCGGATATATTTCCAC	120M1S9I2M1S2M8I101M	793	43	405
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACATGGACGGCATCAGCCTGTGCCAAAG GGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	127M1S1D98M	368	42	246
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACGGCATCAGCCTGTGCCAAAGGGCTCTCCAAATAGTGACAG AGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	109M21D97M	667	19	267
syt10	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGGACGGCATCAGCCTGTGCCAAAGGGCTCT CCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	122M7D98M	963	16	431
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACGGCATCAGCGGCATCAGCCTGTGCC AAAGGGCTCTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTC CAC	126M1S1M3I2S97M	937	11	645
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCACGGCATCAGCCTGTGCCAAAGGGCTCT CCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	124M7D96M	502	11	300
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGCCTGTCAGCCTGTGCCAAAGGGCTCTCCAA ATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	124M1S10D2S90M	524	4	229
	CGGTCTCTCTCCATTCCCGTACGCATCAGCTGCACTCAGCAGCTCACATGGTGCCCCAATGAACTCTCGCCCTCACG CATCGTCCAACATACAATGGATTAACAGGAAAGACATGAATGTCCGAGGACGGCATCAGCCTGTGCCAAAGGGCT CTCCAAATAGTGACAGAGCTGTGCCTCGCCGGGCACGTCGACCGGGAGAAATGCGCGGATATATTTCCAC	122M5D100M	1247	3	670
	GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCACAGAAAGGA TGAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGA ATGAGTCCCAGAACAGG	169M	1431	728	813
	GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCACTTGGTCCT GAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAA TGAGTCCCAGAACAGG	68M1D100M	784	24	453
rgs6	GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCATTGGTCCT GAGAGGCCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAA TGAGTCCCAGAACAGG	68M2D99M	540	7	195
	GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGCCCACCGCATAGTCCTG AGAGGCCTGAGCCATGCTGAGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACA GAAAAGGAATGAGTCCCAGAACAGG	68M3D1M1S12M10I84M	920	4	287
	GTGACAGTACCTTCCTGTATGCAATCATGTTGGGTGAGTCCTCATCTGGGTCGGCTATGTGGTCCTGAGAGGCCTG AGCCATGCTTGATTGCTGGGTCTCTCTTTCTGTGTCTGTGGGGAACGGGCACAACAGAAAAGGAATGAGTCCCAG AACAGG	59M12D98M	1424	1	-
hcn4	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTC GAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGCTCCAGGAG GGCTCTCCTGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTG CC	230M	1380	638	789

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
hcn4	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGCCGCCACCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	142M1S1M4I1S85M	725	48	343
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCATCCGCGCCGCCACCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	136M1S5M7I1S87M	1314	16	225
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGCTCTCCTCCAGGAGCTCCAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	143M5I7M6I80M	799	15	383
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	135M8D87M	917	13	563
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	141M1D1S87M	987	11	377
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCACCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	139M1S1M4D85M	1302	10	587
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGGGCCCTCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	143M3I87M	694	2	49
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCCATCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	98M11I43M2S1D86M	346	2	36
	GTCCGGCAGTATCTGCTCGATGCCTTCCAGCTTAATAAAGCCAGCCTGCTGATCCGGGATGCCCTGATGCTGGCTCGAACCGGCGGCTTCCTCTCCGCCTCCCCCGCCGCTCTGCGCACCGGGCTCCGCGCCGCCACCCGCGCCAGGAGGGCTCTCCTCGCTAGGAGTGACCTCTCCGTCCGTGATCAGACGCCTTTCCTCGGCTACATCCGAGTCGTGGACGTGCC	142M3D85M	222	1	-
hcn4l	AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG	115M1S89M1S29M	1076	697	671
	AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACCAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGAATGATGATCAGGTTGCCCACCATTAGAAG	235M	765	38	389
	AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG	115M1S34M2D53M1S29M	585	18	295

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
hcn4l	AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG	115M1S28M16D45M1S29M	523	9	176
	AGGAGAGGAAATCCACGGCAAACCAGCTGCGCAAGTACCTGACCTTGATCTGCTGAGGGTCCAGGATGATTTCTGTACTGTCCTCTTTCACAATCCCCGTGCGGAAGTTCAGCACGAGGTCGACCAGGAAGAAGGTGTCGGAGACCACGTTTAAACACTATCCAGGCGGGTGTGTGTTCGTCTTTGAAGAAGGTGATGCCCACGGGGATGATGATCAGGTTGCCCACCATTAGAAG	115M1S33M1I56M1S29M	1218	4	915
neo1a	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACTAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	157M1S57M	941	151	691
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTGCGAGGAGCGCCAGTTCTCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	40M16D10M10I91M1S57M	957	95	545
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	50M4D103M1S57M	899	82	475
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	40M16D101M1S57M	696	72	413
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	50M9D98M1S57M	589	72	305
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	42M12D103M1S57M	703	66	359
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	38M31D88M1S57M	985	57	521
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	49M3D105M1S57M	960	20	653
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	48M11D98M1S57M	633	20	415
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCGCGTCGGATAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	46M5D106M1S57M	1026	19	474
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	46M26D85M1S57M	672	16	404
	CAGTGAGGAACTTCAGCCCATTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACATCCATTCAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	11M9I39M5I2S105M1S57M	688	14	336
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	34M28D95M1S57M	807	11	252

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
neo1a	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACAAAGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	50M6D2S99M1S57M	569	10	314
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	40M31D86M1S57M	1179	9	312
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACCAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	51M1S105M1S57M	1569	6	1406
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACCAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	51M1S12D93M1S57M	1010	6	552
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACGGTGCGGAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	51M6I1S105M1S57M	760	6	543
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	45M7D105M1S57M	1451	5	663
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATAGTCGGATAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	48M2S1M11D3M4D1S87M1S57M	754	5	361
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCGGATACACTAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGACGAACGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	160M1S54M	734	4	441
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGTCCTGGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	43M7D2M1S5D99M1S57M	376	2	40
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACGGAGCCGGATGTAGCGGTGCGGTGCGAGGAGTGTAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTTCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	47M2S1M14I1S33M1S72M1S57M	1242	1	-
	CAGTGAGGAACTTCAGCCCATTCTGGTTCTCCACTAGCGGTGCGAGGAGCGCCAGTTCTCCTCAACTGCTCAGTCCACAGCGAGTCTCCTGCCAAGATCGAGTGGAAAAAAGATGGCTCTTTCCTCAGCCTGGCTTCAGATGAGCGCAGACAGGTCCTGGCCGATGGCTCTCTGCTCATCAGCTCAGTGGTGCACTCC	31M17D109M1S57M	698	1	-
neo1b	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	123M6D64M1S12M	1295	155	1734
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	128M1D77M	1334	100	1800
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	129M6D71M	1207	87	1570

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCTCGGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	129M4I77M	1430	78	1952
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	123M6D77M	1342	33	2035
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCAGGGTCTGGGTCTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	129M8I3M4I74M	1573	32	2534
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	121M9D63M1S12M	1017	27	1531
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	206M	679	24	1019
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	125M4D77M	1884	22	2143
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	193M1S12M	347	16	543
neo1b	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	126M7D60M1S12M	345	14	675
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	125M17D51M1S12M	1504	9	1182
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGGGACATTCAGGACATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	120M19D5M4S2M1S55M	1278	9	1869
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCACAGGCTGACGCC	117M22D54M1S12M	1197	2	
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCATTCAGATGCTCATGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	126M3S1M1S1M2S1M2S1M2S66M	8	2	1
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	122M14D70M	4	2	1
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATGCTCGGCAGCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	129M2I77M	49	1	
	TGGCAGGTCTTCCGCGGTTCCTCAGTCAGCCGGAGGGTGTTTCTGTTCGCAGCGGTGGGACTGAGATGCTGAGCTGTGAGGTGACGGAGGAGCTGGTCCCGTTCACCCGCTGGGACAGAGATCTGTGGAGCTGGACAGCAGGGTCATTCAGCTGCCCAGCGGAGCTCTGGTCATCAGCAACGCCTCGCAGGCTGACGCC	122M7D77M	9	1	

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S90M	997	119	507
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S8M14D68M	1117	99	569
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S15M7D68M	1175	80	598
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACTTGTAGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S20M11I12M1S41M3I2S56M1S13M	882	68	903
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S38M10D51M1S13M	541	57	417
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S42M3I57M1S13M	780	55	596
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S17M17D56M	1077	38	490
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S18M1S1D70M	876	30	459
quo	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S99M1S13M	775	30	588
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S42M1S1I1S55M1S13M	822	22	547
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCACAAGTCATTCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S11M3D5M1S1M1S2M5D1S60M	1190	21	408
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCATGACGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S17M1S1I1M1S70M	1469	20	683
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S17M5D68M	1365	18	561
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACAAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S18M1I72M	1602	14	834
	CGCAGAGTCTCCTGACCAACATAATGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S15M81D1M1S34M1S13M	1070	14	568
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S14M7D69M	1063	14	794
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACTAACAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S4M1S1M15D1S68M	876	12	386
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCAACCGGACTCAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S17M8I73M	950	10	426

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCAACCGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S40M3S56M1S13M	618	9	283
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S41M1I58M1S13M	581	6	250
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S42M1S1I1S69M	145	6	23
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACTTGTAGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S20M11I12M1S41M3I2S70M	136	6	89
quo	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGATATCCCTGTCTGAA	6M1S32M1S41M4D54M1S13M	1101	5	323
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S113M	120	4	48
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAACCGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S42M3I71M	107	2	65
	CGCAGAGTCTCCTGACCAACATATGACCCAACTTATCACCAGCATGGTCCAAGCAAGACAGCCTCACAAGTCATTCCTCACCGGCAAGTCATTCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S32M1S22M1S17M4I2M1S1M1S2M2S64M	1184	1	-
	CGCAGAGTCTCCTGACCAACATAATGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	6M1S15M81D1M1S48M	233	1	-
	CGCAGACTCTCCTGACCAACATATGACCCAACTTATCACAAGCATGGTCCAAGCAAGACAGCATCACAAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAGACTGTACTGGGAACTAAGGCCTTACCAGATAAAAGACATCCCTGTCTGAA	153M	25	1	-
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S22M21D45M1S2M1S11M	985	120	638
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S88M1S2M1S11M	951	78	613
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S19M21D48M1S2M1S11M	886	63	460
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S22M29D37M1S2M1S11M	1070	62	464
si:dkey-65j6.2	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	82M3D67M	772	62	333
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	78M15D59M	856	59	414
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCATCAGAGAAACACAGAGAAATGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	81M1D2M1S3M1S1M11I3S59M	829	55	457
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	81M16D55M	935	35	587

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	76M12D64M	837	33	470
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	152M	564	32	351
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S22M14D52M1S2M1S11M	964	26	370
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGCTCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S33M3D52M1S2M1S11M	793	20	426
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGGACCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S25M1S8D1M1S52M1S2M1S11M	787	19	595
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S26M21D41M1S2M1S11M	1035	18	370
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTTTGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S29M11D48M1S2M1S11M	657	17	443
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S23M17D48M1S2M1S11M	1152	15	331
si:dkey-65j6.2	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTAAAAATAAGGCTAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	77M8I4M5D66M	584	11	214
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	71M21D60M	1110	9	532
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGCACATCCAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	74M5S1M5D67M	970	9	382
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	73M31D48M	976	4	130
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGATGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	73M1S1M12D65M	726	4	358
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACGGTATGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S26M1S1M16D44M1S2M1S11M	1411	3	177
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	68M21D63M	913	3	550
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTATGGTCGACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAACCAGGAGGACAGAAG	75M12D1S6M2D56M	834	3	128
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGCAAATCCAGAGCGCATACCGGGACGGTTTGGTCGAACAGTGGTCGGCACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S30M1D2M1S1M1S52M1S2M1S11M	562	1	-
	TCTGATCATCCTAATGAAGCAGAAACTGCTGACCCCTGCGGTGAGGAGAAAATCCAGAGCGCATACCGGGACATCAGAGAAATGCCACATTGTGATTCAGGGACAGTCAAATCAAGAGGACAGAAG	48M1S21M26D41M1S2M1S11M	558	1	-
dclk1b	ATGTGTGTGGATCAATTGAGCCTTACAAGATGCAGGAATATACTAAAAACGTCAACCCTAACTGGTCGGTCAACATGAGAACCACTGCCGGAGCAAAACCGCCCACCTCTCTGGCCAGTGCCAAGGGCGGGTCTATAGAAACCAAGGATAACAGGGATTTCATCCGACCCCAAACTGGTGACCATAATCCGTAGCGGGGTAAAGCCTCGGAAAGCCATCCGAATCCTTCTCAACAA	235M	1638	377	1281

Orthologue	Sequence	Cigar	Mean Nr reads	Nr alleles	SD reads
dclk1b	ATGTGTGTGGATCAATTGAGCCTTACAAGATGCAGGAATATACTAAAAACGTCAACCCTAACTGGTCGGTCAACAT GAGAACCACTGCCGGAGCAAAACCGCCCACCTCTCTGGCCAGTGCCAAGGGCGGGTCTATAGAAACCAAGGATAA CAGGGATTTCATCCGACCCAAACTGGTGACCATAATCCGTAGTGGGGTAAAGCCTCGGAAAGCCATCCGAATCCTT CTCAACAA	193M1S41M	1078	102	877
galnt10chr21	GCGTAACGTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCCGATCCGCCTGGAACCATG CCTGAGGAGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAG AGACACACACTGCTAACTGCTAATA	174M	1246	198	994
	GCGTAACGTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATG CCTGAGGAGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAG AGACACACACTGCTAACTGCTAATA	48M1S125M	952	364	1044
galnt10scaff	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGT	41M1S135M12D27M	272	326	207
	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT	216M	148	187	106
	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGT	41M1S135M14D25M	157	136	94
	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGTGT	41M1S135M8D31M	147	82	107
	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCCGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTCTGTGTGTGTGTGTGTGTGTGTGTGTGTGTGT	214M2N	136	13	127
	GTGGGCACTGGGCTGTGTGTGGAGAGCAAACACTTCTCCTCTGGGTCCCCGATCCGCCTGGAACCATGCCTGAGG AGCCGCGGAGAGCCCAGCTGGAGCCACGGGCAGGTGAGACACGCACCGCTAACAGCGCTAGCACTAGAGACACA CACTGCTAACTGCTAATATGGCTAACTCTGTGTGTGTGTGTGTGTGTGTGTGT	41M1S135M10D29M	4	2	1

All unique sequences identified for targeted orthologues and their respective alignment report string (Cigar). Mean number of reads, the number of alleles the cigar was found in and the standard deviation (SD) are reported. Cigar letters: M=match, D=deletion, I=insertion, S=substitution. The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 3: Unique variants and affected transcripts

Orthologue	Chr	ENSDARG	Affected transcripts	Start (bp)	End (bp)	Mutation	Annotation	Base pair change	VEP impact	Nr affected alleles
				40 856 888	40 856 893	AAGGCG/-	inframe deletion	-6	moderate	75
				40 856 893	40 856 895	GAA/-	inframe deletion	-3	moderate	3
gngt1	19	ENSDARG00000035798	ENSDART00000051950 ENSDART00000139083	40 856 894	40 856 894	A/-	frameshift variant	-1	high	69
				40 856 895	40 856 895	A/T	missense variant	0	moderate	3
				40 856 895	40 856 897	AGA/TTT	missense variant	0	moderate	69
				40 856 897	40 856 896	-/TAAC	stop gained,frameshift variant	4	high	2
				40 856 897	40 856 899	ATG/CCA	missense variant	0	moderate	2
				10 888 933	10 888 953	GACATGAATGTCCGCACAGAG/-	inframe deletion	-21	moderate	19
				10 888 944	10 888 944	C/A	synonymous variant	0	low	42
				10 888 945	10 888 944	-/CAGCTGATG	inframe insertion	9	moderate	42
				10 888 946	10 888 950	GCACA/-	frameshift variant	-5	high	3
				10 888 946	10 888 952	GCACAGA/-	frameshift variant	-7	high	15
				10 888 946	10 888 953	GCACAGAG/-	frameshift variant	-8	high	40
				10 888 947	10 888 947	C/T	synonymous variant	0	low	42
				10 888 948	10 888 948	A/C	missense variant	0	moderate	4
syt10	4	ENSDARG00000045750	ENSDART00000136049	10 888 948	10 888 954	ACAGAGG/-	frameshift variant	-7	high	11
				10 888 949	10 888 958	CAGAGGACGG/-	frameshift variant	-10	high	4
				10 888 950	10 888 949	-/GGGAATGG	frameshift variant	8	high	42
				10 888 950	10 888 949	-/A	frameshift variant	1	high	44
				10 888 950	10 888 950	A/G	synonymous variant	0	low	11
				10 888 951	10 888 951	G/T	stop gained	0	high	42
				10 888 952	10 888 951	-/CAT	stop gained,protein altering variant	3	high	11
				10 888 952	10 888 952	A/-	frameshift variant	-1	high	42
				10 888 952	10 888 953	AG/CA	missense variant	0	moderate	11
				10 888 959	10 888 960	CA/TG	missense variant	0	moderate	4
				28 638 811	28 638 822	CCCACCGCACCT/-	inframe deletion	-12	moderate	1
				28 638 820	28 638 820	C/-	frameshift variant	-1	high	24
rgs6	20	ENSDARG00000015627	ENSDART00000135513 ENSDART00000184779	28 638 820	28 638 821	CC/-	frameshift variant	-2	high	7
				28 638 820	28 638 822	CCT/-	inframe deletion	-3	moderate	4
				28 638 824	28 638 824	G/A	stop gained	0	high	4
				28 638 837	28 638 836	-/CTGAGCCATG	frameshift variant	10	high	4
				1 143 637	1 143 644	ACCCGCGG/-	frameshift variant	-8	high	13
				1 143 638	1 143 638	C/T	missense variant	0	moderate	16
				1 143 641	1 143 641	G/A	missense variant	0	moderate	9
				1 143 643	1 143 643	G/-	frameshift variant	-1	high	11
				1 143 643	1 143 644	GG/CA	missense variant	0	moderate	2
				1 143 643	1 143 646	GGCT/-	frameshift variant	-4	high	9
				1 143 644	1 143 643	-/CCGCCAC	frameshift variant	7	high	16
hcn4	18	ENSDARG00000061685	ENSDART00000136140 ENSDART00000189186	1 143 644	1 143 644	G/C	missense variant	0	moderate	75
				1 143 644	1 143 646	GCT/-	inframe deletion	-3	moderate	1
				1 143 645	1 143 644	-/CTCTC	frameshift variant	5	high	15
				1 143 645	1 143 644	-/GCC	protein altering variant	3	moderate	2
				1 143 645	1 143 645	C/-	frameshift variant	-1	high	2
				1 143 646	1 143 645	-/GCCA	frameshift variant	4	high	48
				1 143 646	1 143 646	T/C	synonymous variant	0	low	48
				1 143 652	1 143 651	-/AGCTCC	inframe insertion	6	moderate	15

Orthologue	Chr	ENSDARG	Affected transcripts	Start (bp)	End (bp)	Mutation	Annotation	Base pair change	VEP impact	Nr affected alleles
hcn4l	25	ENSDARG00000074419	ENSDART00000088249 ENSDART00000148940	29 273 086	29 273 101	CCACGTTAAACACTAT/-	frameshift variant	-16	high	9
				29 273 091	29 273 090	-/T	frameshift variant	1	high	4
				29 273 092	29 273 093	TA/-	frameshift variant	-2	high	18
neo1a	7	ENSDARG00000102855	ENSDART00000158160 ENSDART00000163261 ENSDART00000164768 ENSDART00000181629	53 879 228	53 879 244	CACGGAGCCGTCGGATA/-	frameshift variant	-17	high	1
				53 879 231	53 879 258	GGAGCCGTCGGATACACTAGCGGTGCGA/-	frameshift variant	-28	high	11
				53 879 235	53 879 265	CCGTCGGATACACTAGCGGTGCGAGGAGCGC/-	frameshift variant	-31	high	57
				53 879 237	53 879 252	GTCGGATACACTAGCG/-	frameshift variant	-16	high	157
				53 879 237	53 879 267	GTCGGATACACTAGCGGTGCGAGGAGCGCCA/-	frameshift variant	-31	high	9
				53 879 239	53 879 250	CGGATACACTAG/-	inframe deletion	-12	moderate	66
				53 879 240	53 879 246	GGATACA/-	frameshift variant	-7	high	2
				53 879 242	53 879 248	ATACACT/-	frameshift variant	-7	high	5
				53 879 243	53 879 247	TACAC/-	frameshift variant	-5	high	17
				53 879 243	53 879 268	TACACTAGCGGTGCGAGGAGCGCCAG/-	frameshift variant	-26	high	16
				53 879 244	53 879 245	AC/GT	missense variant	0	moderate	1
				53 879 245	53 879 246	CA/GT	missense variant	0	moderate	5
				53 879 245	53 879 255	CACTAGCGGTG/-	frameshift variant	-11	high	20
				53 879 246	53 879 248	ACT/-	inframe deletion	-3	moderate	20
				53 879 247	53 879 246	-/GCGGTGCGAGGAGT	frameshift variant	14	high	1
				53 879 247	53 879 246	-/TCCAT	frameshift variant	5	high	14
				53 879 247	53 879 247	C/G	missense variant	0	moderate	1
				53 879 247	53 879 248	CT/TC	missense variant	0	moderate	14
				53 879 247	53 879 250	CTAG/-	frameshift variant	-4	high	82
				53 879 247	53 879 252	CTAGCG/-	inframe deletion	-6	moderate	10
				53 879 247	53 879 255	CTAGCGGTG/-	inframe deletion	-9	moderate	69
				53 879 248	53 879 247	-/GGTGCG	protein altering variant	6	moderate	6
				53 879 248	53 879 248	T/G	missense variant	0	moderate	6
				53 879 248	53 879 248	T/C	missense variant	0	moderate	10
				53 879 248	53 879 258	TAGCGGTGCGA/-	frameshift variant	-11	high	5
				53 879 249	53 879 249	A/G	synonymous variant	0	low	2
				53 879 249	53 879 260	AGCGGTGCGAGG/-	inframe deletion	-12	moderate	5
				53 879 250	53 879 254	GCGGT/-	frameshift variant	-5	high	2
				53 879 253	53 879 254	GT/AA	missense variant	0	moderate	10
				53 879 262	53 879 265	GCGC/-	frameshift variant	-4	high	5
				53 879 263	53 879 262	-/CGCCAGTTCT	frameshift variant	10	high	89
				53 879 266	53 879 266	C/T	missense variant	0	moderate	5

Continued Supplementary Table 3

Orthologue	Chr	ENSDARG	Affected transcripts	Start (bp)	End (bp)	Mutation	Annotation	Base pair change	VEP impact	Nr affected alleles
				2 899 719	2 899 719	C/-	frameshift variant	-1	high	65
				2 899 722	2 899 721	-/GGTC	frameshift variant	4	high	18
				2 899 719	2 899 718	-/TCGG	frameshift variant	4	high	51
				2 899 710	2 899 728	ATGCTCGGCCTGTGGAGCT/-	frameshift variant	-19	high	7
				2 899 718	2 899 724	CCTGTGG/-	frameshift variant	-7	high	9
				2 899 716	2 899 719	GGCC/-	frameshift variant	-4	high	15
				2 899 708	2 899 729	AGATGCTCGGCCTGTGGAGCTG/-	frameshift variant	-22	high	1
				2 899 715	2 899 728	CGGCCTGTGGAGCT/-	frameshift variant	-14	high	2
				2 899 712	2 899 718	GCTCGGC/-	frameshift variant	-7	high	1
				2 899 719	2 899 718	-/AGCAGGGT	frameshift variant	8	high	18
neo1b	25	ENSDARG00000075100	ENSDART00000115280	2 899 719	2 899 718	-/AG	frameshift variant	2	high	1
				2 899 716	2 899 732	GGCCTGTGGAGCTGGAC/-	frameshift variant	-17	high	5
				2 899 719	2 899 724	CTGTGG/-	inframe deletion	-6	moderate	59
				2 899 713	2 899 721	CTCGGCCTG/-	inframe deletion	-9	moderate	18
				2 899 722	2 899 723	TG/AT	missense variant	0	moderate	2
				2 899 725	2 899 726	AG/CT	missense variant	0	moderate	2
				2 899 734	2 899 737	GCAG/TTCA	missense variant	0	moderate	7
				2 899 728	2 899 729	TG/AT	missense variant	0	moderate	2
				2 899 716	2 899 718	GGC/ATT	missense variant	0	moderate	2
				2 899 715	2 899 720	CGGCCT/-	inframe deletion	-6	moderate	108
				2 899 740	2 899 740	T/A	missense variant	0	moderate	7
				2 899 720	2 899 720	T/A	synonymous variant	0	low	2

Orthologue	Chr	ENSDARG	Affected transcripts	Start (bp)	End (bp)	Mutation	Annotation	Base pair change	VEP impact	Nr affected alleles
				2 051 654	2 051 734	ATGACCCAACTTATCACAAGCATGGTCCAAGCAAGACAGCATCAC AAGTCATTCCTCACAGGGCAACCGGATCTCCCTCAG/-	inframe deletion	-81	moderate	13
				2 051 699	2 051 699	A/T	stop gained	0	high	10
				2 051 701	2 051 715	GTCATTCCTCACAGG/-	inframe deletion	-15	moderate	10
				2 051 703	2 051 716	CATTCCTCACAGGG/-	frameshift variant	-14	high	61
				2 051 706	2 051 708	TCC/-	inframe deletion	-3	moderate	17
				2 051 709	2 051 715	TCACAGG/-	frameshift variant	-7	high	9
				2 051 710	2 051 716	CACAGGG/-	frameshift variant	-7	high	53
				2 051 710	2 051 719	CACAGGGCAA/-	frameshift variant	-10	high	51
				2 051 712	2 051 711	-/CCGG	frameshift variant	4	high	1
				2 051 712	2 051 711	-/ACCGGACT	frameshift variant	8	high	7
				2 051 712	2 051 712	C/T	missense variant	0	moderate	15
				2 051 712	2 051 714	CAG/ACC	missense variant	0	moderate	9
				2 051 712	2 051 716	CAGGG/-	frameshift variant	-5	high	14
			ENSDART00000109679	2 051 712	2 051 728	CAGGGCAACCGGATCTC/-	frameshift variant	-17	high	23
			ENSDART00000153568	2 051 713	2 051 712	-/G	frameshift variant	1	high	15
quo	6	ENSDARG00000073684	ENSDART00000187502	2 051 713	2 051 712	-/A	frameshift variant	1	high	19
			ENSDART00000187544	2 051 713	2 051 712	-/TTG	stop gained,inframe insertion	3	high	70
			ENSDART00000191165	2 051 713	2 051 713	A/C	synonymous variant	0	low	22
				2 051 713	2 051 714	AG/TA	missense variant	0	moderate	70
				2 051 713	2 051 716	AGGG/-	frameshift variant	-4	high	4
				2 051 714	2 051 713	-/ACC	inframe insertion	3	moderate	51
				2 051 714	2 051 714	G/-	frameshift variant	-1	high	22
				2 051 714	2 051 714	G/A	missense variant	0	moderate	40
				2 051 714	2 051 714	G/C	missense variant	0	moderate	15
				2 051 715	2 051 714	-/C	frameshift variant	1	high	22
				2 051 715	2 051 715	G/C	missense variant	0	moderate	22
				2 051 716	2 051 716	G/T	synonymous variant	0	low	18
				2 051 716	2 051 716	G/A	synonymous variant	0	low	10
				2 051 719	2 051 720	AC/TT	missense variant	0	moderate	1
				2 051 719	2 051 723	ACCGG/-	frameshift variant	-5	high	17
				2 051 724	2 051 724	A/T	missense variant	0	moderate	17
				2 051 736	2 051 736	C/A	missense variant	0	moderate	13

Orthologue	Chr	ENSDARG	Affected transcripts	Start (bp)	End (bp)	Mutation	Annotation	Base pair change	VEP impact	Nr affected alleles
				43 213 133	43 213 153	GGACGGTTTGGCTCCAGCAGT/-	inframe deletion	-21	moderate	62
				43 213 135	43 213 160	ACGGTTTGGCTCCAGCAGTGGTCGGC/-	frameshift variant	-26	high	1
				43 213 136	43 213 149	CGGTTTGGCTCCAG/-	frameshift variant	-14	high	21
				43 213 136	43 213 156	CGGTTTGGCTCCAGCAGTGGT/-	inframe deletion	-21	moderate	101
				43 213 136	43 213 164	CGGTTTGGCTCCAGCAGTGGTCGGCACAT/-	frameshift variant	-29	high	50
				43 213 137	43 213 153	GGTTTGGCTCCAGCAGT/-	frameshift variant	-17	high	15
				43 213 138	43 213 138	G/A	missense variant	0	moderate	4
				43 213 138	43 213 168	GTTTGGCTCCAGCAGTGGTCGGCACATCAGA/-	frameshift variant	-31	high	3
				43 213 139	43 213 139	T/G	synonymous variant	0	low	18
				43 213 139	43 213 143	TTTGG/CACAT	missense variant	0	moderate	9
				43 213 140	43 213 140	T/A	missense variant	0	moderate	3
				43 213 140	43 213 147	TTGGCTCC/-	frameshift variant	-8	high	18
				43 213 140	43 213 151	TTGGCTCCAGCA/-	inframe deletion	-12	moderate	7
				43 213 140	43 213 160	TTGGCTCCAGCAGTGGTCGGC/-	inframe deletion	-21	moderate	12
				43 213 141	43 213 152	TGGCTCCAGCAG/-	inframe deletion	-12	moderate	32
				43 213 142	43 213 141	-/AAAAATAA	frameshift variant	8	high	7
			ENSDART00000158832	43 213 142	43 213 157	GGCTCCAGCAGTGGTC/-	frameshift variant	-16	high	3
			ENSDART00000164262	43 213 143	43 213 153	GCTCCAGCAGT/-	frameshift variant	-11	high	13
si:dkey-65j6..2	23	ENSDARG00000103586	ENSDART00000186065	43 213 143	43 213 157	GCTCCAGCAGTGGTC/-	inframe deletion	-15	moderate	57
			ENSDART00000190315	43 213 144	43 213 144	C/-	frameshift variant	-1	high	1
			ENSDART00000193300	43 213 145	43 213 149	TCCAG/-	frameshift variant	-5	high	9
				43 213 146	43 213 146	C/-	frameshift variant	-1	high	41
				43 213 146	43 213 150	CCAGC/-	frameshift variant	-5	high	7
				43 213 146	43 213 161	CCAGCAGTGGTCGGCA/-	frameshift variant	-16	high	34
				43 213 147	43 213 147	C/G	missense variant	0	moderate	1
				43 213 147	43 213 149	CAG/-	inframe deletion	-3	moderate	69
				43 213 149	43 213 149	G/T	missense variant	0	moderate	41
				43 213 149	43 213 149	G/C	missense variant	0	moderate	18
				43 213 149	43 213 149	G/A	missense variant	0	moderate	1
				43 213 152	43 213 152	G/A	missense variant	0	moderate	3
				43 213 153	43 213 153	T/A	missense variant	0	moderate	41
				43 213 155	43 213 154	-/AAACACAGAGA	frameshift variant	11	high	41
				43 213 155	43 213 157	GTC/AAT	missense variant	0	moderate	41
				43 213 159	43 213 160	GC/-	frameshift variant	-2	high	3

The predicted impact of unique CRISPR-Cas9 induced mutations on protein function was assessed using Ensembl's Variant Effect Predictor (VEP). An allele- and target-specific dosage score was subsequently calculated for each larva by weighting the mutation with the highest predicted impact by a factor 0.33, 0.66, or 1 for mutations with a low, moderate or high predicted impact, respectively. Modification of expression was only observed for transcripts of genes located near the targeted gene. Start and end coordinates are based on GRCz11. The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 4: Mutant allele frequencies

Gene	Nr mutated alleles			Missing	Sample call rate	Mutant allele freq	Expected nr mutated alleles			P_{HWE}
	0	1	2				0	1	2	
gngt1	234	130	12	5	0.987	0.205	238	122	16	2.33E-01
syt10	192	130	54	5	0.987	0.316	176	163	38	9.82E-05
rgs6	344	34	1	2	0.995	0.047	344	34	1	8.69E-01
hcn4	266	100	9	6	0.984	0.157	266	99	9	9.12E-01
hcn4l	349	29	1	2	0.995	0.041	349	30	1	6.32E-01
neo1a	19	113	240	9	0.976	0.797	15	120	236	2.39E-01
neo1b	14	12	277	78	0.795	0.934	1	37	264	3.24E-32
quo	55	42	281	3	0.992	0.799	15	121	241	4.69E-37
si:dkey-65j6.2	22	60	295	4	0.990	0.862	7	90	280	1.34E-10

Number of larvae with 0, 1 and 2 mutated alleles, as well as the number of larvae with a missing call. *P*-value for a Hardy-Weinberg equilibrium (HWE) exact test, considering a ±30 bp window around the CRISPR targeted site as a single locus ($P<2.9E-3$ is significant after Bonferroni correction). Larvae with more than two missing genotypes were excluded from the statistical analysis (i.e. n_total=381). The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 5: Transcript-specific association analysis for heart rate variability and heart rate

Age	Outcome	n	Factor	Gene	Model 1					Model 2				
					Effect size	SE	LCI	UCI	P	Effect size	SE	LCI	UCI	P
2	Heart rate	234	fixed	gngt1	-0.080	0.131	-0.337	0.176	5.41E-01	-0.009	0.116	-0.236	0.217	9.37E-01
				syt10	-0.003	0.088	-0.176	0.170	9.75E-01	-0.006	0.078	-0.159	0.147	9.38E-01
				rgs6	-0.477	0.213	-0.895	-0.060	2.49E-02	-0.376	0.188	-0.745	-0.008	4.53E-02
				hcn4	-0.043	0.116	-0.271	0.184	7.08E-01	0.023	0.105	-0.182	0.227	8.29E-01
				hcn4l	0.096	0.182	-0.259	0.452	5.96E-01	0.148	0.162	-0.169	0.465	3.59E-01
				neo1a	0.110	0.118	-0.121	0.341	3.49E-01	0.138	0.107	-0.072	0.348	1.99E-01
				neo1b	0.006	0.120	-0.230	0.241	9.61E-01	0.062	0.107	-0.149	0.272	5.65E-01
				quo	-0.076	0.084	-0.240	0.088	3.64E-01	0.000	0.076	-0.149	0.149	1.00E+00
				si:dkey-65j6.2	0.059	0.108	-0.152	0.270	5.84E-01	0.167	0.097	-0.023	0.358	8.49E-02
				time of day	0.459	0.042	0.378	0.541	2.07E-28	0.579	0.040	0.500	0.658	1.55E-46
				HRV (in SD)	-	-	-	-	-	-0.424	0.052	-0.526	-0.321	6.06E-16
				intercept	-1.353	0.280	-1.902	-0.805	1.33E-06	-2.148	0.285	-2.706	-1.590	4.47E-14
			random	variation by batch	0.031	0.233	0.000	66000	-	0.098	0.097	0.014	0.680	-
				residual	0.810	0.038	0.738	0.889	-	0.712	0.034	0.649	0.781	-
2	HRV	234	fixed	gngt1	0.155	0.141	-0.120	0.431	2.69E-01	0.112	0.124	-0.131	0.354	3.66E-01
				syt10	-0.027	0.096	-0.214	0.161	7.78E-01	-0.032	0.084	-0.197	0.132	7.00E-01
				rgs6	0.270	0.230	-0.181	0.720	2.41E-01	0.031	0.204	-0.370	0.431	8.80E-01
				hcn4	0.246	0.129	-0.008	0.499	5.75E-02	0.209	0.114	-0.015	0.432	6.71E-02
				hcn4l	0.083	0.200	-0.309	0.476	6.78E-01	0.110	0.176	-0.236	0.455	5.34E-01
				neo1a	0.011	0.139	-0.261	0.283	9.37E-01	0.046	0.123	-0.195	0.288	7.06E-01
				neo1b	0.172	0.133	-0.088	0.432	1.94E-01	0.172	0.117	-0.057	0.400	1.41E-01
				quo	0.116	0.097	-0.074	0.307	2.32E-01	0.072	0.086	-0.097	0.241	4.03E-01
				si:dkey-65j6.2	0.188	0.121	-0.050	0.426	1.21E-01	0.221	0.107	0.012	0.431	3.87E-02
				time of day	0.296	0.047	0.204	0.388	2.80E-10	0.526	0.050	0.428	0.624	5.05E-26
				Heart rate (in SD)	-	-	-	-	-	-0.513	0.062	-0.636	-0.391	2.07E-16
				intercept	-1.718	0.381	-2.466	-0.970	6.68E-06	-2.362	0.353	-3.054	-1.670	2.20E-11
			random	variation by batch	0.310	0.128	0.138	0.697	-	0.306	0.119	0.143	0.657	-
				residual	0.869	0.041	0.792	0.952	-	0.763	0.036	0.696	0.837	-
5	Heart rate	285	fixed	gngt1	0.001	0.123	-0.240	0.242	9.96E-01	0.068	0.101	-0.130	0.266	5.01E-01
				syt10	-0.018	0.083	-0.181	0.145	8.31E-01	-0.049	0.068	-0.183	0.085	4.75E-01
				rgs6	-0.170	0.179	-0.520	0.181	3.43E-01	-0.062	0.147	-0.351	0.227	6.75E-01
				hcn4	0.338	0.106	0.130	0.545	1.44E-03	0.157	0.089	-0.016	0.331	7.58E-02
				hcn4l	-0.035	0.202	-0.430	0.360	8.63E-01	0.024	0.166	-0.301	0.350	8.83E-01
				neo1a	-0.139	0.115	-0.365	0.087	2.27E-01	-0.090	0.095	-0.277	0.097	3.47E-01
				neo1b	0.136	0.131	-0.121	0.392	3.00E-01	0.083	0.108	-0.128	0.294	4.42E-01
				quo	0.011	0.089	-0.165	0.186	9.04E-01	0.022	0.074	-0.123	0.168	7.62E-01
				si:dkey-65j6.2	-0.076	0.113	-0.297	0.146	5.02E-01	0.030	0.094	-0.153	0.214	7.48E-01
				time of day	0.410	0.051	0.310	0.509	6.80E-16	0.565	0.044	0.479	0.651	5.89E-38
				HRV (in SD)	-	-	-	-	-	-0.524	0.045	-0.613	-0.435	7.90E-31
				intercept	-1.243	0.421	-2.069	-0.417	3.17E-03	-1.858	0.406	-2.654	-1.063	4.70E-06
			random	variation by batch	0.509	0.181	0.253	1.023	-	0.615	0.206	0.319	1.186	-
				residual	0.845	0.036	0.778	0.918	-	0.694	0.029	0.639	0.754	-
5	HRV	285	fixed	gngt1	0.125	0.133	-0.136	0.386	3.47E-01	0.130	0.109	-0.084	0.344	2.35E-01
				syt10	-0.063	0.089	-0.237	0.111	4.76E-01	-0.066	0.074	-0.211	0.078	3.69E-01
				rgs6	0.231	0.192	-0.145	0.608	2.29E-01	0.100	0.159	-0.212	0.412	5.30E-01
				hcn4	-0.286	0.112	-0.506	-0.066	1.09E-02	-0.129	0.096	-0.317	0.059	1.78E-01
				hcn4l	0.056	0.211	-0.358	0.470	7.90E-01	0.080	0.179	-0.272	0.431	6.56E-01
				neo1a	0.117	0.118	-0.114	0.348	3.22E-01	0.035	0.103	-0.167	0.237	7.34E-01
				neo1b	-0.014	0.139	-0.286	0.257	9.17E-01	0.002	0.116	-0.226	0.231	9.83E-01
				quo	0.072	0.092	-0.107	0.251	4.31E-01	0.053	0.080	-0.103	0.209	5.06E-01
				si:dkey-65j6.2	0.243	0.119	0.010	0.476	4.08E-02	0.173	0.101	-0.024	0.370	8.59E-02
				time of day	0.241	0.050	0.143	0.339	1.51E-06	0.521	0.050	0.423	0.620	2.36E-25
				Heart rate (in SD)	-	-	-	-	-	-0.601	0.053	-0.705	-0.498	3.75E-30
				intercept	-1.293	0.366	-2.010	-0.575	4.14E-04	-1.981	0.386	-2.738	-1.223	2.97E-07
			random	variation by batch	0.145	0.092	0.042	0.500	-	0.477	0.170	0.237	0.959	-
				residual	0.915	0.039	0.842	0.994	-	0.751	0.032	0.692	0.816	-

Associations of heart rate variability (HRV) and heart rate with the number of mutated alleles across the nine orthologues, weighted by their predicted effect on protein function. Associations were examined for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata), using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Associations were adjusted for time of day and the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). Additionally, Model 2 includes mutual adjustment of heart rate and HRV. The genes quo and si:dkey-65j6.2 are orthologues of the human KIAA1755.

Supplementary Table 6: Transcript-specific association analyses for body size

Outcome	Factor	Gene		2dpf						5dpf				
			n	Effect size	SE	LCI	UCI	P	n	Effect size	SE	LCI	UCI	P
		gngt1		-0.120	0.115	-0.346	0.106	2.97E-01		0.043	0.115	-0.182	0.268	7.05E-01
		syt10		0.147	0.080	-0.009	0.303	6.54E-02		0.095	0.076	-0.053	0.243	2.09E-01
		rgs6		-0.444	0.179	-0.795	-0.094	1.31E-02		-0.467	0.181	-0.822	-0.113	9.65E-03
		hcn4		-0.131	0.101	-0.330	0.067	1.96E-01		0.080	0.094	-0.104	0.265	3.95E-01
		hcn4l		0.126	0.200	-0.265	0.518	5.28E-01		-0.075	0.192	-0.452	0.301	6.95E-01
	fixed	neo1a		-0.029	0.110	-0.244	0.185	7.89E-01		-0.136	0.108	-0.347	0.076	2.09E-01
Length		neo1b	332	0.075	0.122	-0.163	0.314	5.37E-01	242	0.045	0.127	-0.205	0.294	7.26E-01
		quo		-0.008	0.083	-0.171	0.156	9.27E-01		-0.026	0.082	-0.186	0.134	7.52E-01
		si:dkey-65j6.2		0.045	0.107	-0.165	0.254	6.77E-01		-0.087	0.105	-0.293	0.119	4.08E-01
		time of day		0.111	0.042	0.029	0.193	7.88E-03		0.091	0.047	-0.001	0.183	5.25E-02
		intercept		-0.490	0.371	-1.216	0.237	1.86E-01		-0.131	0.449	-1.011	0.750	7.71E-01
	random	variation by batch		0.395	0.129	0.208	0.748	-		0.689	0.227	0.361	1.315	-
		residual		0.897	0.035	0.830	0.968	-		0.703	0.032	0.643	0.770	-
		gngt1		0.025	0.110	-0.190	0.240	8.20E-01		-0.091	0.137	-0.360	0.178	5.08E-01
		syt10		0.052	0.076	-0.096	0.201	4.89E-01		0.048	0.090	-0.129	0.225	5.99E-01
		rgs6		-0.031	0.168	-0.361	0.299	8.55E-01		-0.462	0.223	-0.899	-0.024	3.86E-02
		hcn4		-0.022	0.096	-0.210	0.167	8.22E-01		-0.090	0.110	-0.305	0.126	4.14E-01
		hcn4l		0.066	0.188	-0.303	0.435	7.25E-01		0.097	0.222	-0.338	0.531	6.63E-01
	fixed	neo1a		0.208	0.105	0.002	0.414	4.76E-02		-0.032	0.125	-0.277	0.214	8.01E-01
Dorsal area		neo1b	322	0.003	0.117	-0.226	0.232	9.81E-01	234	-0.139	0.154	-0.441	0.163	3.66E-01
		quo		0.052	0.080	-0.105	0.210	5.16E-01		0.267	0.095	0.081	0.452	4.92E-03
		si:dkey-65j6.2		-0.184	0.103	-0.386	0.019	7.50E-02		-0.274	0.123	-0.515	-0.034	2.54E-02
		time of day		-0.052	0.040	-0.131	0.026	1.89E-01		-0.072	0.053	-0.176	0.032	1.74E-01
		intercept		-0.049	0.365	-0.765	0.668	8.94E-01		0.428	0.406	-0.369	1.225	2.92E-01
	random	variation by batch		0.437	0.138	0.236	0.810	-		0.300	0.129	0.130	0.696	-
		residual		0.842	0.033	0.779	0.910	-		0.824	0.039	0.751	0.903	-
		gngt1		0.052	0.124	-0.192	0.295	6.76E-01		-0.172	0.119	-0.405	0.060	1.47E-01
		syt10		0.086	0.087	-0.084	0.255	3.24E-01		-0.007	0.082	-0.167	0.153	9.30E-01
		rgs6		-0.063	0.182	-0.419	0.293	7.27E-01		-0.074	0.175	-0.417	0.269	6.72E-01
		hcn4		-0.137	0.106	-0.344	0.070	1.95E-01		-0.115	0.102	-0.314	0.084	2.57E-01
		hcn4l		0.053	0.208	-0.354	0.460	7.99E-01		0.195	0.200	-0.196	0.586	3.29E-01
	fixed	neo1a		0.235	0.112	0.015	0.454	3.61E-02		-0.034	0.122	-0.272	0.204	7.80E-01
Lateral area		neo1b	300	-0.051	0.127	-0.300	0.199	6.89E-01	258	-0.138	0.148	-0.428	0.151	3.48E-01
		quo		0.139	0.088	-0.033	0.312	1.13E-01		0.262	0.092	0.080	0.443	4.66E-03
		si:dkey-65j6.2		-0.248	0.111	-0.466	-0.030	2.57E-02		-0.402	0.117	-0.632	-0.173	5.79E-04
		time of day		-0.047	0.044	-0.134	0.040	2.91E-01		-0.074	0.050	-0.172	0.023	1.35E-01
		intercept		-0.001	0.355	-0.697	0.695	9.97E-01		0.595	0.415	-0.217	1.408	1.51E-01
	random	variation by batch		0.265	0.104	0.123	0.570	-		0.440	0.154	0.221	0.876	-
		residual		0.908	0.038	0.837	0.985	-		0.796	0.035	0.729	0.868	-
		gngt1		0.088	0.110	-0.129	0.304	4.27E-01		-0.187	0.129	-0.441	0.066	1.47E-01
		syt10		0.052	0.076	-0.097	0.202	4.93E-01		-0.023	0.083	-0.186	0.141	7.87E-01
		rgs6		0.093	0.174	-0.249	0.435	5.93E-01		-0.042	0.180	-0.394	0.310	8.13E-01
		hcn4		-0.014	0.096	-0.203	0.174	8.80E-01		0.017	0.102	-0.182	0.217	8.64E-01
		hcn4l		-0.033	0.185	-0.395	0.330	8.59E-01		0.092	0.229	-0.357	0.541	6.87E-01
	fixed	neo1a		0.314	0.103	0.113	0.515	2.23E-03		0.033	0.117	-0.196	0.262	7.76E-01
Volume		neo1b	299	-0.062	0.111	-0.279	0.154	5.73E-01	199	0.105	0.143	-0.176	0.386	4.64E-01
		quo		0.153	0.078	0.000	0.305	4.98E-02		0.150	0.091	-0.028	0.328	9.82E-02
		si:dkey-65j6.2		-0.239	0.100	-0.435	-0.042	1.74E-02		-0.398	0.116	-0.626	-0.170	6.23E-04
		time of day		-0.065	0.040	-0.144	0.015	1.10E-01		-0.075	0.052	-0.177	0.027	1.50E-01
		intercept		-0.153	0.355	-0.849	0.544	6.67E-01		0.198	0.448	-0.679	1.075	6.58E-01
	random	variation by batch		0.427	0.136	0.229	0.796	-		0.572	0.196	0.292	1.120	-
		residual		0.806	0.033	0.743	0.874	-		0.718	0.037	0.650	0.794	-

Associations were analyzed for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata) using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 7: Comparison of larvae with two vs. zero functionally knocked alleles for heart rate variability and heart rate

Gene	Age	n	Outcome	Model 1					Model 2				
				Effect size	SE	LCI	UCI	P	Effect size	SE	LCI	UCI	P
syt10	2	146	Heart rate	-0.115	0.199	-0.506	0.276	5.64E-01	-0.103	0.176	-0.449	0.243	5.59E-01
			HRV	-0.073	0.205	-0.475	0.329	7.22E-01	-0.128	0.182	-0.485	0.229	4.82E-01
	5	180	Heart rate	-0.184	0.189	-0.555	0.187	3.31E-01	-0.198	0.152	-0.496	0.101	1.95E-01
			HRV	-0.066	0.191	-0.440	0.308	7.30E-01	-0.133	0.157	-0.441	0.174	3.96E-01
hcn4	2	183	Heart rate	0.695	0.358	-0.006	1.396	5.20E-02	0.674	0.309	0.068	1.280	2.93E-02
			HRV	0.148	0.399	-0.634	0.930	7.11E-01	0.520	0.344	-0.153	1.194	1.30E-01
	5	212	Heart rate	1.208	0.321	0.579	1.836	1.66E-04	0.444	0.278	-0.100	0.988	1.10E-01
			HRV	-1.349	0.338	-2.010	-0.687	6.48E-05	-0.759	0.294	-1.335	-0.182	9.91E-03
neo1a	2	70	Heart rate	0.102	0.294	-0.473	0.677	7.28E-01	0.216	0.264	-0.302	0.734	4.15E-01
			HRV	0.029	0.343	-0.643	0.702	9.32E-01	0.094	0.317	-0.527	0.714	7.67E-01
	5	93	Heart rate	0.166	0.274	-0.372	0.704	5.45E-01	0.113	0.247	-0.371	0.597	6.47E-01
			HRV	0.089	0.286	-0.473	0.650	7.57E-01	0.140	0.249	-0.348	0.628	5.75E-01
neo1b	2	80	Heart rate	0.153	0.262	-0.360	0.667	5.58E-01	0.225	0.243	-0.251	0.701	3.55E-01
			HRV	0.292	0.308	-0.312	0.896	3.43E-01	0.332	0.290	-0.236	0.901	2.52E-01
	5	89	Heart rate	0.025	0.325	-0.612	0.661	9.40E-01	-0.073	0.276	-0.614	0.469	7.92E-01
			HRV	0.053	0.333	-0.600	0.706	8.74E-01	-0.092	0.289	-0.658	0.474	7.50E-01
quo	2	170	Heart rate	-0.104	0.178	-0.454	0.245	5.58E-01	0.006	0.167	-0.321	0.332	9.73E-01
			HRV	0.227	0.206	-0.176	0.631	2.70E-01	0.150	0.193	-0.228	0.528	4.36E-01
	5	214	Heart rate	-0.065	0.192	-0.441	0.310	7.34E-01	0.012	0.163	-0.308	0.332	9.40E-01
			HRV	0.240	0.193	-0.139	0.618	2.14E-01	0.203	0.177	-0.144	0.550	2.52E-01
si:dkey-65j6.2	2	60	Heart rate	-0.110	0.226	-0.552	0.332	6.27E-01	0.027	0.216	-0.397	0.452	8.99E-01
			HRV	0.291	0.298	-0.294	0.876	3.29E-01	0.224	0.281	-0.327	0.775	4.25E-01
	5	67	Heart rate	0.111	0.298	-0.473	0.695	7.08E-01	0.160	0.275	-0.378	0.698	5.60E-01
			HRV	0.364	0.350	-0.322	1.051	2.98E-01	0.395	0.328	-0.248	1.038	2.29E-01

Associations of heart rate variability (HRV) and heart rate by comparison of embryos with two functionally knocked alleles with wildtypes. Associations were examined for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata), using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. In model 1, heart rate and HRV are not mutually adjusted for each other. Associations were adjusted for time of day and the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). Model 2 includes additional mutual adjustment of heart rate and HRV. The genes quo and si:dkey-65j6.2 are orthologues of the human KIAA1755.

Supplementary Table 8: Transcript-specific association analysis for dichotomous cardiac outcomes

Age	Outcome	n controls	n cases	Gene	OR	LCI	UCI	P
				gngt1	0.951	0.387	2.336	9.12E-01
				syt10	1.447	0.792	2.642	2.30E-01
				rgs6	0.855	0.193	3.799	8.37E-01
				hcn4	2.644	1.402	4.986	2.66E-03
				neo1a	1.165	0.499	2.717	7.24E-01
				neo1b	1.694	0.434	6.604	4.48E-01
2	SA arrest	248	36	*quo*	0.764	0.404	1.442	4.06E-01
				si:dkey-65j6.2	0.750	0.327	1.720	4.97E-01
				time of day	1.117	0.829	1.506	4.66E-01
				intercept	0.008	0.000	0.414	1.64E-02
				Batch 4	5.656	0.523	61.178	1.54E-01
				Batch 2	11.656	1.342	101.224	2.60E-02
				Batch 3	5.798	0.665	50.571	1.12E-01
				Batch 1	7.540	0.770	73.835	8.27E-02
				gngt1	1.163	0.494	2.740	7.29E-01
				syt10	1.467	0.810	2.654	2.06E-01
				rgs6	0.788	0.175	3.553	7.57E-01
				hcn4	2.749	1.461	5.171	1.71E-03
				neo1a	1.176	0.507	2.731	7.06E-01
				neo1b	2.445	0.658	9.080	1.82E-01
				quo	0.746	0.401	1.389	3.56E-01
2	SA pause	297	39	*si:dkey-65j6.2*	0.755	0.339	1.685	4.93E-01
				time of day	1.147	0.856	1.539	3.59E-01
				intercept	0.004	0.000	0.185	4.77E-03
				Batch 2	14.723	1.714	126.443	1.42E-02
				Batch 3	5.507	0.633	47.917	1.22E-01
				Batch 4	4.948	0.457	53.573	1.88E-01
				Batch 1	7.380	0.755	72.106	8.57E-02
				Batch 5	0.890	0.042	18.778	9.40E-01
				gngt1	0.309	0.037	2.572	2.77E-01
				syt10	0.236	0.050	1.128	7.05E-02
				rgs6	1.049	0.068	16.223	9.73E-01
				hcn4	2.460	0.680	8.899	1.70E-01
				hcn4l	0.937	0.086	10.181	9.58E-01
				neo1a	0.897	0.190	4.232	8.91E-01
				neo1b	1.561	0.162	15.023	7.00E-01
5	SA pause	321	9	*quo*	0.672	0.221	2.043	4.84E-01
				si:dkey-65j6.2	3.533	0.421	29.662	2.45E-01
				time of day	1.066	0.514	2.209	8.64E-01
				intercept	0.016	0.000	7.788	1.90E-01
				Batch 1	0.240	0.011	5.447	3.70E-01
				Batch 3	0.214	0.015	3.049	2.55E-01
				Batch 5	0.213	0.005	8.956	4.18E-01
				Batch 2	0.703	0.073	6.750	7.60E-01

Associations of cardiac outcomes with the number of mutated alleles across the nine orthologues, weighted by their predicted effect on protein function. At 2 and 5 days post fertilization (dpf), associations were analyzed using logistic regression for outcomes where number of observed cases and genotype frequency allowed. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors. The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 9: Comparison of larvae with two vs. zero functionally knocked alleles for body size

Gene	Outcome	2dpf						5dpf					
		n	Effect size	SE	LCI	UCI	P	n	Effect size	SE	LCI	UCI	P
syt10	Length	208	0.207	0.174	-0.135	0.548	2.35E-01	153	0.174	0.182	-0.182	0.530	3.38E-01
	Dorsal area	200	0.022	0.174	-0.320	0.363	9.01E-01	145	0.101	0.204	-0.299	0.501	6.20E-01
	Lateral area	182	0.002	0.200	-0.390	0.393	9.93E-01	155	-0.122	0.193	-0.501	0.257	5.28E-01
	Volume	186	-0.028	0.173	-0.366	0.311	8.73E-01	128	-0.170	0.201	-0.564	0.224	3.98E-01
hcn4	Length	241	0.026	0.306	-0.573	0.625	9.33E-01	177	0.130	0.291	-0.439	0.700	6.54E-01
	Dorsal area	233	-0.020	0.309	-0.626	0.586	9.49E-01	171	0.440	0.319	-0.185	1.065	1.67E-01
	Lateral area	217	0.154	0.321	-0.475	0.783	6.31E-01	183	0.300	0.326	-0.339	0.940	3.57E-01
	Volume	216	0.264	0.303	-0.330	0.858	3.83E-01	141	0.474	0.280	-0.075	1.024	9.06E-02
neo1a	Length	107	-0.188	0.263	-0.703	0.328	4.75E-01	77	-0.317	0.317	-0.939	0.305	3.17E-01
	Dorsal area	104	0.707	0.246	0.224	1.189	4.08E-03	75	0.242	0.340	-0.424	0.907	4.76E-01
	Lateral area	97	-0.072	0.280	-0.621	0.476	7.96E-01	84	0.202	0.356	-0.496	0.901	5.70E-01
	Volume	94	0.516	0.266	-0.005	1.037	5.23E-02	69	-0.164	0.261	-0.676	0.347	5.29E-01
neo1b	Length	117	0.241	0.311	-0.369	0.851	4.38E-01	76	0.378	0.339	-0.286	1.042	2.64E-01
	Dorsal area	112	-0.056	0.296	-0.635	0.523	8.49E-01	72	-0.206	0.430	-1.050	0.637	6.32E-01
	Lateral area	106	0.343	0.336	-0.315	1.001	3.07E-01	74	0.209	0.394	-0.564	0.982	5.96E-01
	Volume	110	0.170	0.284	-0.386	0.726	5.49E-01	62	0.181	0.373	-0.550	0.913	6.27E-01
quo	Length	241	-0.115	0.182	-0.471	0.241	5.27E-01	180	0.078	0.171	-0.256	0.413	6.46E-01
	Dorsal area	235	0.238	0.178	-0.110	0.587	1.80E-01	175	0.545	0.210	0.133	0.957	9.52E-03
	Lateral area	223	0.244	0.198	-0.144	0.632	2.18E-01	188	0.394	0.212	-0.021	0.809	6.30E-02
	Volume	218	0.331	0.174	-0.010	0.671	5.70E-02	146	0.175	0.197	-0.211	0.560	3.75E-01
si:dkey-65j6.2	Length	80	0.221	0.325	-0.416	0.858	4.96E-01	53	-0.107	0.345	-0.782	0.569	7.57E-01
	Dorsal area	78	-0.110	0.266	-0.631	0.412	6.80E-01	51	-0.566	0.317	-1.188	0.056	7.45E-02
	Lateral area	74	-0.489	0.295	-1.067	0.089	9.72E-02	57	-0.622	0.389	-1.384	0.139	1.09E-01
	Volume	75	-0.300	0.262	-0.813	0.212	2.51E-01	44	-0.732	0.338	-1.395	-0.070	3.03E-02

Associations with body size by comparison of embryos with two functionally knocked alleles with wildtypes. Associations were analyzed for the main transcript of each targeted zebrafish orthologue using hierarchical linear models (xtmixed in Stata) using inverse-normally transformed outcomes, so effect sizes and SEs can be interpreted as z-score units. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors, with larvae nested in batches (random factor). The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.

Supplementary Table 10: Comparison of larvae with two vs. zero functionally knocked alleles for cardiac traits at 2dpf

Gene	Outcome	n controls	n cases	OR	LCI	UCI	P
syt10	SA arrest	141	36	3.480	0.789	15.347	9.95E-02
	SA pause	183	39	3.451	0.792	15.043	9.92E-02
hcn4	SA arrest	149	36	3.666	0.655	20.525	1.39E-01
	SA pause	150	39	3.667	0.633	21.249	1.47E-01
neo1a	SA arrest	72	36	1.200	0.148	9.717	8.65E-01
	SA pause	72	39	1.200	0.148	9.717	8.65E-01
quo	SA arrest	150	36	0.553	0.136	2.239	4.06E-01
	SA pause	190	39	0.513	0.132	1.993	3.35E-01
si:dkey-65j6.2	SA arrest	47	36	0.145	0.009	2.389	1.77E-01
	SA pause	47	39	0.145	0.009	2.389	1.77E-01

Associations of cardiac outcomes by comparison of embryos with two functionally knocked alleles and wildtypes. Associations were analyzed using logistic regression for outcomes where number of observed cases and genotype frequency allowed. Associations were adjusted for time of day and for the weighted number of mutated alleles in the other genes as fixed factors. The genes *quo* and *si:dkey-65j6.2* are orthologues of the human *KIAA1755*.