Efa6 regulates axon growth, branching and maintenance by eliminating off-track
microtubules at the cortex

Yue Qu*[,1], Ines Hahn*[,#,1], Meredith Lees[1], Jill Parkin[1], André Voelzmann[1], Karel Dorey[1], Alex
Rathbone[2], Claire Friel[2], Viki Allan[1], Pilar Okenve Ramos[3], Natalia Sanchez-Soriano[3],
Andreas Prokop[1,#]

1) The University of Manchester, Manchester Academic Health Science Centre, Faculty of
Biology, Medicine and Health, School of Biology, Manchester, UK

2) The University of Nottingham, School of Life Sciences, Faculty of Medicine & Health
Sciences, Nottingham, UK

3) Department of Cellular and Molecular Physiology, Institute of Translational Medicine,
University of Liverpool, Liverpool, United Kingdom

Running title: The role of Efa6 in axon maintenance

Key words: *Drosophila*, neurodegeneration, axons, actin, cytoskeleton, microtubules

*　　authors contributed equally
#　　authors for correspondence:
The University of Manchester
Faculty of Life Sciences
Oxford Road
Manchester M13 9PT
Tel: +44-(0)161-27-51556
Fax: +44-(0)161-27-51505
Andreas.Prokop@manchester.ac.uk
Ines.Hahn@manchester.ac.uk

Abstract

Axons are the enormously long, cable-like neuronal extensions that wire our nervous system.
The formation and plastic maintenance of these delicate structures requires parallel bundles
of microtubules (MTs), which form the structural backbones and highways for life-sustaining
transport in axons. In ageing and certain axonopathies, axonal MTs lose their bundled
appearances forming areas of disorganisation. We use *Drosophila* to study various
mechanisms that promote axonal MT bundle organisation and dynamics during development
and maintenance. Here we report important roles of the membrane-associated protein Efa6 in
axonal MT regulation relevant for axonal growth, branching and MT bundle maintenance. Efa6
acts as a typical cortical MT collapse factor by eliminating MTs that approach the axonal
plasma membrane. This action abolishes MTs that have left the axon bundle, thus providing a

quality control mechanism that prevents MT disorganisation during the formation and
maintenance of axons. Furthermore, we find this action to reduce axonal growth and branch
formation in cultured neurons and *in vivo*. Using structure-function analyses, we show that
cytoplasmic localisation of a small N-terminal peptide motif alone is sufficient to eliminate
entire MT networks in mouse fibroblasts, and loss of whole axons in *Drosophila* primary
neurons. This highly destructive activity becomes meaningful when restricted to the cell
plasma membrane via the plekstrin homology domain present in the Efa6 C-terminus. Our
data provide a paradigm for the fundamental and disease-relevant role of cortical collapse
factors in neurons.

Introduction

Axons are the cable-like neuronal extensions that wire the nervous system. They are typically
0.5-10 μm in diameter, but can be up to a meter long in humans (Debanne et al., 2011;
Prokop, 2013a). It is a fascinating challenge to understand how axons can extend over these
enormous distances and branch in orderly manners, but also how these delicate structures
can be maintained for a lifetime, i.e. many decades in humans. It is not surprising that we
gradually lose about half of our axons towards old age (Adalbert and Coleman, 2012; Medana
and Esiri, 2003) and that axon decay is a prominent neurodegenerative phenomenon (Fang
and Bonini, 2012; Wang et al., 2012).

The morphology and structural dynamics of axons essentially depends on continuous
parallel bundles of microtubules (MTs) running along their entire length to provide structural
support. During early stages of axon growth, amoeboid structures at axon tips, called growth
cones, navigate along pre-defined paths. They constantly elongate the axonal MT bundle
through a process where single MTs projecting into the periphery of growth cones get
stabilised and serve as a guide for elongation of the entire MT bundle (Dent et al., 2011;
Lowery and van Vactor, 2009; Prokop et al., 2013). If such events occur in parallel, the growth
cone can split and bifurcate into two branches (Acebes and Ferrus, 2000). Also interstitial
branching along the axon can occur. It depends on MT bundles in the shaft becoming
disorganised so that off-track MTs can extend into the periphery to seed new collateral
branches (Kalil and Dent, 2014; Lewis et al., 2013; Yu et al., 2008).

Axon maintenance depends on these MT bundles as well; they drive the dynamic
morphogenetic changes underlying neuronal plasticity, and also to form the tracks required for
life-sustaining cargo transport between cell bodies and synaptic endings (Prokop, 2013a). For
this, MTs have to maintain their proper bundled confirmation. MTs that become unbundled
and disorganised in axon swellings of ageing neurons are potential predictors of axon decay
(Adalbert and Coleman, 2012; Fiala et al., 2007). MT bundle maintenance is expected to
require constant renewal through polymerisation to prevent senescence (Voelzmann et al.,
2016a), but also other classes of MT regulatory proteins seem to play important roles. This is
strongly suggested by mouse models of neurodegeneration which display axonal MT
disorganisation and link to MT regulators; reported examples are spastin-associated spastic
paraplegia (SPG4; Online Mendelian Inheritance in Man®/OMIM reference #182601) or
spectraplakin-linked hereditary sensory and autonomic neuropathy (HSAN type VI; #614653)
(Bernier and Kothary, 1998; Dalpe et al., 1998; Fassier et al., 2013; Fiala et al., 2007; Tarrade
et al., 2006).

We study the mechanisms underlying the formation, maintenance and decay of axonal

MT bundles and use neurons of the fruit fly *Drosophila melanogaster* as an efficient model to
this end (Prokop et al., 2013). Our studies of over 40 cytoskeletal regulators performed in this
system led us to propose the model of "local axon homeostasis" (Prokop, 2016; Voelzmann et
al., 2016a). It proposes that the force-enriched environment of axons promotes MT
disorganisation, requiring therefore the action of MT regulating factors to tame MTs into
bundles. One example is the MT guidance mechanism of the spectraplakin Short stop (Shot)
(Alves-Silva et al., 2012). It binds simultaneously to cortical actin via its N-terminus and to the
tips of polymerising MTs through its C-terminus, thus guiding extending MTs in parallel to the
axonal cortex and laying them out into bundles; in the absence of Shot function, MTs leave
their bundled arrangements and go off-track (Alves-Silva et al., 2012), and Shot might
therefore serve as a paradigm for MT disorganisation observed in spectraplakin-linked HSAN
type VI (see above) (Voelzmann et al., 2017).

Here we show that Efa6 (Exchange factor for Arf6) acts as a complementary factor to
Shot in MT bundle maintenance. Like its *C. elegans* homologue (*Ce*Efa6) (O'Rourke et al.,
2010), it acts as a cortical collapse factor, in that it eliminates MTs at the plasma membrane of
fly neurons and mouse fibroblasts. Through this action, Efa6 maintains MT bundles (through
eliminating accidental off-track MTs) and negatively regulates axon growth and branch
formation (through preventing MT entry into filopodia). Since the morphological phenotypes
we describe resemble those of disease-relevant neuronal collapse factors in mammals, our
findings provide conceptual explanations for how cortical collapse factors contribute to the
growth and maintenance of neurons.

Methods

<u>Fly stocks</u>

Loss-of-function mutant stocks used in this study were the two deficiencies uncovering the
Efa6 locus *Df(3R)Exel6273* (94B2-94B11 or 3R:22,530,780..22,530,780) and *Df(3R)ED6091i*
(94B5-94C4 or 3R:22,587,681..22,587,681), *shot³* (the strongest available allele of *short stop*)
(Kolodziej et al., 1995; Sánchez-Soriano et al., 2009), and the null mutant alleles *Efa6^{KO#1}*,
Efa6^{GX6[w-]}, *Efa6^{GX6[w+]}* and *Arf51F^{GX16[w-]}* (all genomically engineered precise deletions) (Huang
et al., 2009). Gal4 driver lines used were the pan-neuronal lines *sca-Gal4* (strongest in
embryos) (Sánchez-Soriano et al., 2010a) and *elav-Gal4* (1st and 3rd chromosomal, both
expressing at all stages) (Luo et al., 1994), as well as the *ato-Gal4* line expressing in a subset
of neurons in the adult brain (Hassan et al., 2000; Voelzmann et al., 2016b) and *GMR31F10-*
Gal4 (Bloomington #49685) expressing in a subset of lamina neurons. Lines for targeted
gene expression were *UAS-Efa6^{RNAi}* (VDRC #42321), *UAS-Gal80^{ts}* (Zeidler et al., 2004), *UAS-*
Eb1-GFP (Alves-Silva et al., 2012), *UAS-α-tubulin84B* (Grieder et al., 2000) and *UAS-*
tdTomato (Zschätzsch et al., 2014). Efa6 expression was detected via the genomically
engineered *Efa6-GFP* allele, where a GFP was inserted after the last amino acid in exon 14
(Huang et al., 2009).

<u>*Drosophila* primary cell culture</u>

Drosophila primary neuron cultures were performed as published previously (Prokop et al.,
2012; Qu et al., 2017). In brief, stage 11 embryos were treated for 1 min with bleach to
remove the chorion, sterilized for ~30 s in 70% ethanol, washed in sterile Schneider's/FCS,

and eventually homogenized with micro-pestles in 1.5 centrifuge tubes containing 21 embryos
per 100 µl dispersion medium (Prokop et al., 2012) and left to incubated for 5 min at 37°C.
Cells are washed with Schneider's medium (Gibco), spun down for 4 mins at 650 g,
supernatant was removed and cells re-suspended in 90 µl of Schneider's medium containing
20% fetal calf serum (Gibco). 30 µl drops were placed on cover slips. Cells were allowed to
adhere for 90-120 min either directly on glass or on cover slips coated with a 5 µg/ml solution
of concanavalin A, and then grown as a hanging drop culture for hours or days at 26°C as
indicated.

To abolish maternal rescue of mutants, i.e. masking of the mutant phenotype caused by
deposition of normal gene product from the healthy gene copy of the heterozygous mothers in
the oocyte (Prokop, 2013b), we used a pre-culture strategy (Prokop et al., 2012; Sánchez-
Soriano et al., 2010b) where cells were kept for 5 days in a tube before they were plated on a
coverslip.

For the transfection of *Drosophila* primary neurons, a quantity of 70-75 embryos per 100 µl
dispersion medium was used. After the washing step and centrifugation, cells were re-
suspended in 100 µl transfection medium [final media containing 0.1-0.5 µg DNA and 2 µl
Lipofecatmine 2000 (L2000)]. To generate this media, dilutions of 0.1-0.5 µg DNA in 50 µl
Schneider's medium and 2 µl L2000 in 50 µl Schneider's medium were prepared, then mixed
together and incubated at room temperature for 5-30 mins, before being added to the cells in
centrifuge tubes where they were kept for 24 hrs at 26°C. Cells were then treated again with
dispersion medium, re-suspended in culture medium and plated out as described above.

For temporally controlled knock-down experiments we used flies carrying the driver construct
elav-Gal4, the knock-down construct *UAS-Efa6-RNAi*, and the temperature-sensitive Gal4
inhibitor *UAS-Gal80[ts]*, all in parallel. At the restrictive temperature of 19°C, Gal80[ts] blocks
Gal4-induced expression of *Efa6-RNAi*, and this repressive action is removed at the
permissive temperature of 27°C where Gal80[ts] is non-functional. Control neurons were from
flies carrying only the *Gal4/Gal80* (control 1 in Fig. 4K) or only the *Efa6-RNAi* transgene
(control 2).

<u>Fibroblast cell culture</u>

NIH/3T3 fibroblasts were grown in DMEM supplemented with 1% glutamine
(Invitrogen), 1% penicillin/streptomycin (Invitrogen) and 10% FCS in culture dishes (100 mm
with vents; Fisher Scientific UK Ltd) at 37°C in a humidified incubator at 5% CO_2. Cells were
split every 2-3 d, washed with pre-warmed PBS, incubated with 4 ml of Trypsin-EDTA (T-E) at
37°C for 5 min, then suspended in 7 ml of fresh culture medium and eventually diluted (1/3-
1/20 dilution) in a culture dish containing 10 ml culture media.

For transfection of NIH/3T3 cells, 2 ml cell solution (~10^5 cells per ml) were first
transferred to 6-well plates, and grown overnight to double cell density. 2 µg of DNA and 2 µl
Plus reagent (Invitrogen) were added to 1 ml serum-free media in a centrifuge tube, incubated
for 5 mins at RT, then 6 µl Lipofectamine (Invitrogen) were added, and incubated at RT for 25
mins. Cells in the 6-well plate were washed with serum-free medium and 25 mins later
DNA/Lipofectamine was mixed into the medium (1/1 dilution). Plates were incubated for 3 hrs
at 37°C, washed with 2 ml PBS, 400 µl trypsin were added for 5 mins (37°C), then 3 ml
complete medium; cells were suspended and added in 1 ml aliquots to 35 mm glass-bottom
dishes (MatTek) coated with fibronectin [300 µl of 5 µg/ml fibronectin (Sigma-Aldrich) placed
in the center of a MatTek dish for 1 hr at 37°C, then washed with PBS]; 1 ml of medium was

added and cells grown for 6 hrs or 24 hrs at 37°C in a CO_2 incubator. For live imaging, the
medium was replaced with 2 ml Ham's F-12 medium + 4% FCS.

Dissection of adult heads

To analyse the function of Efa6 in MT bundle integrity in medulla axons *in vivo*, flies were
aged at 29°C. Flies were maintain in groups of up to 20 flies of the same gender (Stefana et
al., 2017) and changed into new tubes every 3-4 days. Brain dissections were performed in
Dulbecco's PBS (Sigma, RNBF2227) after briefly sedating them on ice. Dissected brains with
their laminas and eyes attached were placed into a drop of Dulbecco's PBS on MatTek glass
bottom dishes (P35G1.5-14C), covered by coverslips and immediately imaged with a 3i
Marianas Spinning Disk Confocal Microscope.

To measure branching in *ato-Gal4 Drosophila* neurons, adult brains were dissected in
Dulbecco's PBS and fixed with 4% PFA for 15 min. Antibody staining and washes were
performed with PBS supplemented with 0.3% Triton X-100. Specimens were embedded in
Vectashield (VectorLabs).

Immunohistochemistry

Primary fly neurons and fibroblasts were fixed in 4% paraformaldehyde (PFA) in 0.05 M
phosphate buffer (PB; pH 7–7.2) for 30 min at room temperature (RT); for anti-Eb1 staining,
ice-cold +TIP fix (90% methanol, 3% formaldehyde, 5 mM sodium carbonate, pH 9; stored at -
80°C and added to the cells) (Rogers et al., 2002) was added for 10 mins. Adult brains were
dissected out of their head case in PBS and fixed with 4% PFA in PBS for 1 hr, followed by a
1hr wash in PBT.

Antibody staining and washes were performed with PBT. Staining reagents: anti-
tubulin (clone DM1A, mouse, 1:1000, Sigma; alternatively, clone YL1/2, rat, 1:500, Millipore
Bioscience Research Reagents); anti-DmEb1 (gift from H. Ohkura; rabbit, 1:2000) (Elliott et
al., 2005); anti-Elav (mouse, 1:1000, DHB); anti-GFP (goat, 1:500, Abcam); Cy3-conjugated
anti-HRP (goat, 1:100, Jackson ImmunoResearch); F-actin was stained with Phalloidin
conjugated with TRITC/Alexa647, FITC or Atto647N (1:100 or 1:500; Invitrogen and Sigma).
Specimens were embedded in ProLong Gold Antifade Mountant.

Microscopy and data analysis

Standard documentation was performed with AxioCam monochrome digital cameras (Carl
Zeiss Ltd.) mounted on BX50WI or BX51 Olympus compound fluorescent microscopes. For
the analysis of *Drosophila* primary neurons, we used two well established parameters (Alves-
Silva et al., 2012; Sánchez-Soriano et al., 2010a): axon length (from cell body to growth cone
tip; measured using the segmented line tool of ImageJ) and the degree of MT disorganisation
in axons which was either measured as binary score or ratio (percentage of neurons showing
obvious MT disorganisation in their axons) or as "MT disorganisation index" (MDI) (Qu et al.,
2017): the area of disorganisation was measured using the freehand selection in ImageJ; this
value was then divided by axon length (see above) multiplied by 0.5 μm (typical axon diameter,
thus approximating the expected area of the axon if it were not disorganised). For Eb1::GFP
comet counts, neurons were subdivided into axon shaft and growth cones (GC): the proximal
GC border was set where the axon widens up (broader GCs) or where filopodia density

increases significantly (narrow GCs). MT loss in fibroblasts was assessed on randomly
chosen images of successfully transfected, GFP-expressing fibroblasts, stained for tubulin
and actin. Images were derived from at least 2 independent experimental repeats performed
on different days, for each of which at least 3 independent culture wells were analysed by
taking a minimum of 20 images per well. Due to major differences in plasma membrane
versus cytoplasmic localisation of constructs, their expression strengths could not be
standardised. Assuming a comparable expression strength distribution, we therefore analyse
all transfected cells in the images and assigned them to three categories: MTs intact,
damaged or gone (Fig. 8). To avoid bias, image analyses were performed blindly, i.e. the
genotype or treatment of specimens was masked. To analyse ruffle formation in fibroblasts,
cells were stained with actin and classified (with or without ruffles).

To measure the degree of branching, we measured axonal projections of dorsal cluster
neurons in the medulla, which is part of the optic lobe in the adult brain (Hassan et al., 2000;
Voelzmann et al., 2016b). These neurons were labelled by expressing *UAS-myr-tdTomato* via
the *ato-Gal4* driver either alone (control), together with *UAS-Efa6^{RNAi}* or together with *UAS-
Efa6-FL-GFP*. We analysed them in young brains (2-5 d after eclosure of flies from their pupal
case) or old brains (15-18 d). Z-stacks of adult fly brains (optic lobe area) were taken with a
Leica DM6000 B microscope and extracted with Leica MM AF Premier software. They were
imaged from anterior and the number of branches was quantified manually. Branches were
defined as the protrusions from the DC neuron axons in the medulla. Branches in fly primary
neurons at 5DIV were also counted manually and defined as MT protrusions from main axon.

To measure MT disorganisation in the optic lobe of adult flies, *GMR31F10-Gal4*
(Bloomington #49685) was used to express *UAS-α-tubulin84B-GFP* (Grieder et al., 2000) in a
subset of lamina axons which projects within well-ordered medulla columns (Prokop and
Meinertzhagen, 2006). Flies were left to age for 26-27 days (about half their life expectancy)
and then their brains were dissected out. A section of the medulla columns comprising the 4
most proximal axonal terminals was used to quantify the number of swellings and regions with
disorganised MTs.

Time lapse imaging of cultured primary neurons (in Schneider's/FCS) and fibroblasts
(in Ham's F-12/FCS) was performed on a Delta Vision Core (Applied Precision) restoration
microscope using a [*100x/1.40 UPlan SAPO (Oil)*] objective and the Sedat Quad filter set
(*Chroma #89000*). Images were collected using a Coolsnap HQ2 (Photometrics) camera. The
temperature was set to 26°C for fly neurons and 37°C for fibroblasts. Time lapse movies were
constructed from images taken every 2 s for 2 mins. To analyse MT dynamics, Eb1::GFP
comets were tracked manually using the "manual tracking" plug-in of ImageJ.

For statistical analyses, Kruskal–Wallis one-way ANOVA with *post hoc* Dunn's test or
Mann–Whitney Rank Sum Tests (indicated as P_{MW}) were used to compare groups, and χ2
tests (indicated as P_{X2}) were used to compare percentages. All raw data of our analyses are
provided as supplementary Excel/Prism files.

<u>Molecular biology</u>

EGFP tags are based on pcDNA3-EGFP or pUAST-EGFP. All *Drosophila melanogaster* efa6
constructs are based on cDNA cloneIP15395 (Uniprot isoform C, intron removed).
Caenorhabditis elegans efa-6 (Y55D9A.1a) constructs are derived from pCZGY1125-efa-6-
pcr8 (kindly provided by Andrew Chisholm). *Homo sapiens* PSD(1) (ENST00000406432.5,
isoform 202) constructs were PCR-amplified from pLC32-hu-psd1-pcr8 vector (kindly provided

by Andrew Chisholm). *Homo sapiens* PSD2 (ENST00000274710.3, isoform 201, 771aa)
constructs were PCR-amplified from pLC33-hu-psd2-pcr8 vector (kindly provided by Andrew
Chisholm). *Homo sapiens* PSD3 was PCR-amplified from pLC34 hu-psd3-pcr8 vector (kindly
provided Andrew Chisholm). Note that the PSD3 cDNA clone is most closely related to
isoform 201 (ENST00000286485.12: 513aa) and therefore lacks the putative N-terminus
found in isoform 202 (ENST00000327040.12). However, the putative MTED core sequence is
encoded in the C-terminal PH domain (Fig.8C), not the potential N-terminus. *Homo sapiens*
PSD4 (ENST00000441564.7, isoform 205) was PCR-amplified from pLC35-hu-psd4-pcr8
vector (kindly provided by Andrew Chisholm).The CAAX motif is derived from human KRAS.
The *Dm*Efa6-NtermΔSxiP::EGFP (aa1-410) insert was synthesised by GeneArt Express
(ThermoFisher). All construct were cloned using standard (SOE) PCR/ligation based methods,
and constructs and inserts are detailed in Table T1. To generate transgenic fly lines,
P[acman]M-6-attB-UAS-1-3-4 constructs were integrated into *PBac{yellow[+]-attP-*
3B}VK00031 (Bloomington line #9748) via PhiC31 mediated recombination (outsourced to
Bestgene Inc.).

final vector	Source	insert
pcDNA3-EGFP	Addgene	*XhoI*-EGFP-*XbaI*
pUAST-AscI-PacI-EGFP	this study	*KpnI, AscI, PacI*-EGFP-*XbaI*
***pUAST-DmEfa6FL-EGFP* (aa1-1387)**	this study	*KpnI, AscI*-**kozak**-Efa6 (aa1-1387)-**GSGSGS-EGFP**-*PacI, XbaI*
***P[acman]M-6-attB-UAS-1-3-4-DmEfa6FL-EGFP* (aa1-1387)**	this study	*AscI*-**kozak**-*Dm*Efa6 (aa1-1387)-**GSGSGS-EGFP**-*PacI*
***pcDNA3.1-DmEfa6FL-EGFP* (aa1-1387)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6 (aa1-1387)-**GSGSGS-EGFP**-*PacI, XbaI*
***pUAST-DmEfa6ΔCterm-EGFP* (aa1-894)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6Δ**Cterm** (aa1-894)-**GSGSGS-EGFP**-*PacI, XbaI*
***P[acman]M-6-attB-UAS-1-3-4-DmEfa6ΔCterm-EGFP* (aa1-894)**	this study	*AscI*-**kozak**-*Dm*Efa6Δ**Cterm** (aa1-894)-**GSGSGS-EGFP**-*PacI*
***pcDNA3.1-DmEfa6ΔCterm-EGFP* (aa1-894)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6Δ**Cterm** (aa1-894)-**GSGSGS-EGFP**-*PacI, XbaI*
***pUAST-DmEfa6-Nterm-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP**-*PacI, XbaI*
***P[acman]M-6-attB-UAS-1-3-4-DmEfa6-Nterm-EGFP* (aa1-410)**	this study	*AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP**-*PacI*
***pcDNA3.1-DmEfa6-Nterm-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP**-*PacI, XbaI*
***pUAST-DmEfa6-Nterm-CAAX-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP-CAAX[KRAS]**-*PacI, XbaI*
***P[acman]M-6-attB-UAS-1-3-4-DmEfa6-Nterm-CAAX-EGFP* (aa1-410)**	this study	*AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP-CAAX[KRAS]**-*PacI*
***pcDNA3.1-DmEfa6-Nterm-CAAX-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS-EGFP-CAAX[KRAS]**-*PacI, XbaI*
***pUAST-DmEfa6-NtermΔSxiP-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-NtermΔ**SxiP** (aa1-410; **SQIP>AAAA; SRIP>AAAA**)-**GSGSGS-EGFP**-*PacI, XbaI*
***pcDNA3.1-DmEfa6-NtermΔSxiP-EGFP* (aa1-410)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-NtermΔ**SxiP** (aa1-410; **SQIP>AAAA; SRIP>AAAA**)-**GSGSGS-EGFP**-*PacI, XbaI*
***pUAST-DmEfa6-NtermΔMTED-EGFP* (aa1-300)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-NtermΔ**MTED** (aa1-300)-**GSGSGS-EGFP**-*PacI, XbaI*
***pcDNA3.1-DmEfa6-NtermΔMTED-EGFP* (aa1-300)**	this study	*KpnI, AscI*-**kozak**-*Dm*Efa6-NtermΔ**MTED** (aa1-300)-**GSGSGS-EGFP**-*PacI, XbaI*

pUAST-DmEfa6∆Nerm-EGFP (aa851-1387)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Dm*Efa6∆**Nerm** (aa851-1387)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-DmEfa6∆Nerm-EGFP (aa851-1387)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Dm*Efa6∆**Nerm** (aa851-1387)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pUAST-DmEfa6-MTED-EGFP (aa322-341)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Dm*Efa6-MTED (aa322-341)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-DmEfa6-MTED-EGFP (aa322-341)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Dm*Efa6-MTED (aa322-341)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-CeEfa6-FL-EGFP (aa1-816)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Ce*Efa6 (aa1-816)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-CeEfa6-Nterm-EGFP (aa1-152)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Ce*Efa6-Nterm (aa1-152)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-CeEfa6-MTED-EGFP (aa24-42)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Ce*Efa6-MTED (aa24-42)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD(1)-FL-EGFP (aa1-1024)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD(1) (aa1-1024)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD(1)-Nterm-EGFP (aa1-280)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD(1)-Nterm (aa1-280)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD(1)-MTED-EGFP (aa31-49)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD(1)-MTED (aa31-49)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD2-FL-EGFP (aa1-771)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD2 (aa1-771)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD3-EGFP (aa515-1047)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD3 (aa515-1047)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-HsPSD4-FL-EGFP (aa1-1027)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD4 (aa1-1027)-**GSGSGS**-<u>*NotI*</u>-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-co-HsPSD(1)-MTED-EGFP (aa31-49)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Hs*PSD(1)-MTED (aa31-49)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-co-CeEfa6-MTED-EGFP (aa24-42)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Ce*Efa6-MTED (aa24-42)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pcDNA3.1-co-DmEfa6-MTED-EGFP (aa322-341)	this study	<u>*KpnI, AscI*</u>-**kozak**-*Dm*Efa6-MTED (aa322-341)-**GSGSGS**-**EGFP**-<u>*PacI, XbaI*</u>
pCS107-DmEfa6-Nterm-EGFP	this study	<u>*NotI*</u>-**kozak**-*Dm*Efa6-Nterm (aa1-410)-**GSGSGS**-**EGFP**-<u>*StuI*</u>
RFP (Xenopus injection)	tba	tba
pFastBac-His6-MCAK-EGFP-StrepII	tba	His6-MCAK::EGFP-StrepII
pFastBac-His6-DmEfa6∆Cterm-EGFP-StrepII (aa1-894)	this study	His6-*Dm*Efa6∆**Cterm**::EGFP-StrepII (aa1-894)

Tab. T1. co=codon optimised; *Dm*=*Droshpila melanogaster*; *Ce*=*Caenorhabditis elegans*;
Hs=*Homo sapiens*

<u>*In silico* analyses</u>

To generate the **phylogenetic tree** of Efa6/PSD full length isoforms and N-terms of different
species (see Fig. S3), their amino acid sequences were aligned using Muscle or ClustalO
(Goujon et al., 2010; McWilliam et al., 2013; Sievers et al., 2011). ProtTest (Abascal et al.,
2005; Darriba et al., 2011) was used to determine amino acid frequencies in the protein
datasets and to identify the optimal amino acid substitution model to be used for the Bayesian
inference (VT+I+G+F). CUDA-Beagle-optimised MrBayes (Ronquist et al., 2012) was run
using the VT+I+G+F model [prset statefreqpr=fixed(empirical); lset rates=invgamma] using 5
chains (1 heated) and 9 parallel runs until the runs converged and standard deviation of split
frequencies were below 0.015 (0.06 for N-terms); PSRF+ was 1.000 and min ESS was >1300
for the TL, alpha and pinvar parameters. The *Drosophila melanogaster* Sec7-PH domain-
containing protein Steppke was used as outgroup in the full length tree. Archaeopteryx (Han
and Zmasek, 2009) was used to depict the MrBayes consensus tree showing branch lengths

(amino acid substitutions per site) and Bayesian posterior probabilities.

To **identify a potential MTED in PSD(1)**, previously identified Efa6 MTED motifs
(O'Rourke et al., 2010) of 18 orthologues were aligned to derive an amino acid logo (Suppl.
Tab. not available here). Further orthologues were identified and used to refine the logo.
Invariant sites and sites with restricted amino acid substitutions were determined (most
prominently MxG-stretch). Stretches containing the invariant MxG stretch were aligned among
vertebrate species to identify potential candidates. Berkley's Weblogo server (Crooks et al.,
2004) was used to generate amino acid sequence logos for each phylum using MTED
(ExxxMxGE/D) and MTED-like (MxGE/D) amino acid stretches.

In vitro analyses

Protein Expression and Purification: *Drosophila* Efa6-ΔCterm was cloned into a modified
pFastBac vector containing an N-terminal *His6* tag and C-terminal *eGFP* and *StrepII* tags.
Recombinant protein was expressed in Sf9 insect cells for 72 hours using a Baculovirus
system. The protein was purified via a two-step protocol. Cleared lysate was loaded onto a
1ml HisTrap column (GE Healthcare) in Ni-Affinity buffer [50 mM Tris pH 7.5, 300 mM NaCl,
10 % (v/v) glycerol, 1 mMM was shipped as lyophilized powder with a purity of 95.2%. Upon
arrival peptide was dissolved in ultrapure water and stored at -20°C.

MT binding assays: GMPCPP-stabilised, rhodamine-labeled MTs were adhered to
the surface of flow chambers as described earlier (Helenius et al., 2006). 20 nM Efa6-
ΔCterm::GFP (in BRB20 pH 6.9, 75mM KCl, 0.05% Tween20, 0.1 mg/ml BSA, 1% 2-
mercaptoethanol, 40mM glucose, 40 mg/ml glucose oxidase, 16 mg/ml catalase), 20 nM
MCAK::GFP (in the same buffer plus 1 mM ATP and 1 mM taxol) was introduced to the MT-
containing channel. Images were recorded using a Zeiss Observer.Z1 microscope equipped
with a Zeiss Laser TIRF 3 module, QuantEM 512SC EMCDD camera (Photometrics) and
100x objective (Zeiss, alphaPlanApo/1.46NA oil). Images of rhodamine-labeled MTs using a
lamp as the excitation source and GFP fluorescence using TIRF illumination via a 488 nm
laser were collected as described earlier (Patel et al., 2016). For both rhodamine and GFP
imaging an exposure time of 100 ms was used. The mean GFP intensity on individual MTs
was determined from the mean pixel intensity of lines drawn along the long-axis of individual
microtubules in Fiji (Schindelin et al., 2012). The rhodamine signal was used to locate the
position of MTs in the GFP images. Intensity from a region of background was subtracted.

MT depolymerisation assays: Single-cycled, rhodamine-labeled, GMPCPP-stabilised
MTs were adhered to the surface of flow chambers as described in (Helenius et al., 2006).
Images of a field of fluorescent microtubules (25% rhodamine-labeled tubulin) were recorded
using a Zeiss Observer.Z1 microscope, collecting 1 image every 5 s with an exposure time of
100 ms. Efa6-ΔCterm::GFP (14 nM), MCAK (40 nM) in solution (BRB20 pH 6.9, 75mM KCl,
1mM ATP, 0.05% Tween 20, 0.1 mg/ml BSA, 1% 2-mercaptoethanol, 40mM glucose, 40
mg/ml glucose oxidase, 16 mg/ml catalase) were added to the channel 1 min after acquisition
had commenced. De-polymerisation rates were determined from plots of the length of
individual microtubules versus time, obtained by thresholding and particle analysis of images
using Fiji (Schindelin et al., 2012).

***Xenopus* oocyte assays**: cytosol extracts from *Xenopus* oocytes were obtained as
described in (Allan and Vale, 1991). **MT depolymerisation** was assessed in a microscopic
flow chamber (Vale and Toyoshima, 1988) where *Xenopus* cytosol (1 µl cytosol diluted with 20
µl acetate buffer) was incubated for 20 min to allow MTs to polymerise. Then cytosol was

exchanged by flow through with Efa6-ΔCterm::GFP, MCAK or synthetic MTED peptide (all 20
nM in acetate buffer pH 7.4: 100 mM K-Acetate, 3 mM Mg-Acetate, 5 mM EGTA, 10 mM
HEPES), and MT length changes observed by recording 10 random fields via VE-DIC
microscopy (Allan, 1993; Allan and Vale, 1991). **MT polymerisation** was analysed in a
microscope flow cell containing 9 µl diluted *Xenopus* cytosol (see above) to which 1 µl acetate
buffer was added, either alone or containing 20nM MTED. After 10 min, 20 random fields were
recorded via VE-DIC microscopy for each condition and the numbers of MTs per field counted.

For the *in vivo* assay, *Xenopus* embryos were injected in one blastomere at the 4-cell stage
with 200 ng of mRNA encoding Efa6-Nterm::GFP or mCherry alone. The embryos were
imaged at stage 10.25 (Heasman, 2006) with a Leica fluorescent stereoscope.

Results

<u>Efa6 is a negative regulator of axonal growth and branching in cultured neurons and *in vivo*</u>

To evaluate the function of Efa6 in regulating axonal microtubules (MTs) we first determined
its expression in the nervous system. We used a genomically engineered fly line in which the
endogenous *Efa6* gene was GFP-tagged (*Efa6-GFP*) (Huang et al., 2009). These animals
express *Efa6-GFP* throughout the CNS at larval and adult stages, likely in most or even all
neurons (Fig. 1F-I). We cultured primary neurons from this fly line to analyse the subcellular
distribution of Efa6. In young neurons (6 hrs *in vitro;* 6 HIV) as well as mature neurons (5 days
in vitro; 5DIV), Efa6 was expressed throughout cell bodies and axons (Fig. 1A-E).

To determine the importance of Efa6 for axonal morphogenesis, we manipulated Efa6
expression levels and analysed axon growth and collateral branching. We used several fly
lines with decreased or abolished Efa6 expression: Efa6 knock-down (*Efa6-RNAi*),
overlapping deficiencies uncovering the entire *Efa6* gene locus (*Efa6^{Def}*), or different loss-of-
function mutant alleles generated by genomic engineering (*Efa6$^{KO\#1}$*, *Efa6$^{GX6[w-]}$*, *Efa6$^{GX6[w+]}$*). In
all these conditions, axon length at 6 HIV was increased over wild-type by at least 20% (Fig.
2A-D). We also generated transgenic fly lines for the targeted expression of Efa6-FL::GFP.
When expressing it in primary neurons these cells showed a 20% reduction in axon length
(2C,D). Together these results suggested Efa6 to exhibit a growth reducing role.

When assessing mature primary neurons at 5 days *in vitro* (DIV; Fig.2I), we found that
Efa6$^{KO\#1}$ homozygous mutant neurons showed almost two times more collateral branches
than wildtype neurons (Fig.2F,I). Conversely, branching was reduced by 21 % in neurons
expressing Efa6-FL::GFP (Fig.2G,I). These results are consistent with Efa6 acting as a
negative regulator of axon growth and branching.

To extend these studies to neurons *in vivo*, we measured a *myr-tdTomato* labelled subset of
axonal projections in the optic lobe of adult brains (Fig.3, see Methods) (Hassan et al., 2000;
Voelzmann et al., 2016b). We compared axon branching in young (2-5 d after eclosure of flies
from their pupal case) and old brains (15-18 d; Figs.3, S1) of wildtype, Efa6-RNAi and Efa6-
GFP overexpressing flies. We found that *Efa6* knock-down in dorsal cluster neurons caused a
significant increase in branch numbers both in young (29%) and in old brains (38%). In
contrast, over-expression of Efa6::GFP strongly decreased branch numbers by 33% in young
and 28% in old brains, respectively (Fig.3G).

In these experiments, Efa6-FL::GFP had an intriguing further effect: Only 57% of young brains
had any axon protrusion in the medulla region, compared to 88% in controls (Figs.3C,H, S1B).

Since axons are eventually present in old brains of Efa6-FL::GFP expressing flies (Fig.3F,H,
S1D), we concluded that this phenotype reflects delayed outgrowth and is likely the *in vivo*
equivalent to short axon growth observed upon Efa6 over-expression in primary neurons
(green bars in Fig.2D).

These data obtained from neurons in culture and in the adult brain, clearly demonstrate that
the presence of Efa6 in neurons is relevant for their morphology by acting as a negative
regulator of axonal growth and branching.

Efa6 helps maintain axonal MT bundle integrity in cultured neurons

Our analyses revealed a further phenotype: In neurons depleted of Efa6, we found regions
where axonal MTs were not arranged into bundles but were curled up and criss-crossing each
other, especially in mature neurons at 5 DIV (curved arrow in Fig. 2F).

We analysed various Efa6-depleted conditions at 6 HIV and found that the relative amount of
neurons showing regions of MT disorganisation was increased 1.5 to 2.5 fold when
normalised to wild-type controls (Fig. S2). Although many neurons showed some degree of
visible MT disorganisation already at early stages, the extent of MT disorganisation in
individual neurons tended to be fairly mild. To quantify the strength of this phenotype we used
a more refined readout by measuring the area of MT disorganisation per axon area (referred
to as 'MT disorganisation index', MDI) (Qu et al., 2017). MDI measurements in *Efa6* mutants
revealed a mild 1.3 fold increase in MT disorganisation in young neurons which gradually
worsened from 2.3 fold at 5 DIV to ~4 fold at 10 DIV (Fig.4A-F,I; all normalised to controls). To
test whether this gradual increase might be due to wild-type *Efa6* gene product deposited in
the *Efa6* mutant embryos by their heterozygous mothers (Prokop, 2013b), we used a pre-
culture strategy to remove potential maternal Efa6 contributions (see Methods) (Prokop et al.,
2012; Sánchez-Soriano et al., 2010b). After 5 days in pre-culture, we still found a low amount
of MT disorganisation in young neurons and a subsequent gradual increase to severe
phenotypes over the following days (Fig.4J).

This finding argues for a continued role of Efa6 in preventing MT disorganisation during
development as well as in mature neurons. To test whether Efa6 regulates MT organisation in
mature neurons, we used a temperature-based conditional knock-down technique (GAL80[ts];
see Methods) to induce *Efa6* knock-down after 3 DIV when neurons have long undergone
synaptic differentiation (Küppers-Munther et al., 2004; Prokop et al., 2012). Following this
delayed onset, cells were kept for 4 further days under *Efa6* knock-down conditions and fixed
at day 7. We found a significant increase in MT disorganisation in the experimental neurons,
much stronger than in parallel control neurons without knock-down (Fig. 4K). In contrast, at
3DIV (i.e. before onset of knock-down) there was no difference in MT disorganisation between
experimental and control neurons. This clearly indicates that MT disorganisation upon Efa6
deficiency occurs even in fully differentiated neurons and is not restricted to developmental
stages.

In contrast to increased MT disorganisation upon Efa6 loss of function, over-expression of
Efa6-FL::GFP for 5 days in wild-type neurons caused a reduction below the baseline levels
measured in control cells (cultured in parallel without the expression construct; Fig.4I). Our
various findings support therefore a role of Efa6 in preventing MT disorganisation in
developing and mature neurons.

    Efa6 helps maintain axonal MT bundle integrity *in vivo*

    We then assessed whether a role of Efa6 in MT bundle maintenance is relevant *in vivo*. We
    labelled MTs in a subset of lamina neurons in the adult optic lobe (Prokop and Meinertzhagen,
    2006) by expressing α-*tubulin84B-GFP* either alone (*GMR-tub* controls), or together with
    Efa6^{RNAi} to knock down *Efa6* specifically in these neurons (*GMR-tub-Efa6^{IR}*; see Methods for
    details).

    We analysed 26-27 day old flies and found that Efa6 knock-down caused a doubling in the
    occurrence of axonal swellings with disorganised axonal MTs: the average of total swellings
    per column section was increased from 0.3 in controls to 0.65 swellings upon Efa6 knock-
    down, and about a third of these contained disorganised MTs (*GMR-tub-Efa6^{IR}*: 0.23 per
    column section; *GMR-tub*: 0.13; Fig.5). These data suggested that our findings in cultured
    neurons are relevant *in vivo*.

    Taken together, we identified three different morphological features of axons that are
    regulated by Efa6: their growth, branching and MT bundle organisation. To study the
    underlying mechanisms we first performed subcellular analyses.

    Efa6 eliminates MTs in growth cones

    Based on the literature, Efa6 may cause these morphological effects through changes in the
    actin cytoskeleton, either as a potential activator of small Arf GTPases via its central Sec7
    domain (D'Souza-Schorey and Chavrier, 2006; Huang et al., 2009), or through its other C-
    terminal domains (see below) (Derrien et al., 2002; Franco et al., 1999; Macia et al., 2008).
    Alternatively, it could act directly on MTs, as was suggested by studies of *Ce*Efa6 in oocytes
    where it eliminates MTs at the plasma membrane and therefore acts as a cortical collapse
    factor (O'Rourke et al., 2010). We focussed on the latter possibility and studied potential
    effects of Efa6 on MT behaviours.

    For this, we distinguished between different subcellular compartments of neurons, known to
    display very different behaviours: in the dynamically shaped, amoeboid growth cones (GCs) at
    the tips of growing axons, MTs splay out towards the membrane in the GC periphery (Prokop
    et al., 2013). In contrast, in the cable-like axon shafts, most MTs extend within bundles and
    are kept away from the membrane (Prokop, 2013a). If Efa6 eliminates MT at the membrane,
    manipulations of its functions should therefore affect MTs in GCs and axon shafts differently.
    We therefore analysed these two sub-neuronal compartments separately, starting with GCs.

    In GCs of *Efa6* mutant neurons at 6 HIV, we found an increase in MT polymerisation events:
    the total number of Eb1 comets was increased compared to wild-type controls (Fig. 6I).
    Eb1::GFP comet velocity was unaffected (wildtype: 0.27µm/s ± 0.2SEM; *Efa6*: 0.30µm/s ±
    0.2SEM), but their lifetime was ~1.4 times longer in mutant GCs (Fig. 6M). Compared to
    control neurons, Eb1::GFP comets in *Efa6* mutant neurons appeared to vanish less often
    when reaching the GC periphery, where they could occasionally be observed to undergo
    persistent curved extensions along the periphery (Suppl. Movies not available here).

    We reasoned that, if there are more polymerising MTs in GCs of Efa6-depleted neurons, this
    should increase their chances of entering into filopodia. In agreement with this prediction, we
    found considerably more *Efa6* mutant than control neurons with at least one Eb1 comet or
    tubulin-stained MT in GC filopodia (Fig.6J, K). Also per GC, the average number of filopodia
    containing comets or MTs was increased, whilst the total number of filopodia per GC was
    comparable (~11 per neuron in wild-type and ~10 in *Efa6*). Consistent with this finding, the

Efa6 helps maintain axonal MT bundle integrity *in vivo*

We then assessed whether a role of Efa6 in MT bundle maintenance is relevant *in vivo*. We
labelled MTs in a subset of lamina neurons in the adult optic lobe (Prokop and Meinertzhagen,
2006) by expressing α-*tubulin84B-GFP* either alone (*GMR-tub* controls), or together with
Efa6^{RNAi} to knock down *Efa6* specifically in these neurons (*GMR-tub-Efa6^{IR}*; see Methods for
details).

We analysed 26-27 day old flies and found that Efa6 knock-down caused a doubling in the
occurrence of axonal swellings with disorganised axonal MTs: the average of total swellings
per column section was increased from 0.3 in controls to 0.65 swellings upon Efa6 knock-
down, and about a third of these contained disorganised MTs (*GMR-tub-Efa6^{IR}*: 0.23 per
column section; *GMR-tub*: 0.13; Fig.5). These data suggested that our findings in cultured
neurons are relevant *in vivo*.

Taken together, we identified three different morphological features of axons that are
regulated by Efa6: their growth, branching and MT bundle organisation. To study the
underlying mechanisms we first performed subcellular analyses.

Efa6 eliminates MTs in growth cones

Based on the literature, Efa6 may cause these morphological effects through changes in the
actin cytoskeleton, either as a potential activator of small Arf GTPases via its central Sec7
domain (D'Souza-Schorey and Chavrier, 2006; Huang et al., 2009), or through its other C-
terminal domains (see below) (Derrien et al., 2002; Franco et al., 1999; Macia et al., 2008).
Alternatively, it could act directly on MTs, as was suggested by studies of *Ce*Efa6 in oocytes
where it eliminates MTs at the plasma membrane and therefore acts as a cortical collapse
factor (O'Rourke et al., 2010). We focussed on the latter possibility and studied potential
effects of Efa6 on MT behaviours.

For this, we distinguished between different subcellular compartments of neurons, known to
display very different behaviours: in the dynamically shaped, amoeboid growth cones (GCs) at
the tips of growing axons, MTs splay out towards the membrane in the GC periphery (Prokop
et al., 2013). In contrast, in the cable-like axon shafts, most MTs extend within bundles and
are kept away from the membrane (Prokop, 2013a). If Efa6 eliminates MT at the membrane,
manipulations of its functions should therefore affect MTs in GCs and axon shafts differently.
We therefore analysed these two sub-neuronal compartments separately, starting with GCs.

In GCs of *Efa6* mutant neurons at 6 HIV, we found an increase in MT polymerisation events:
the total number of Eb1 comets was increased compared to wild-type controls (Fig. 6I).
Eb1::GFP comet velocity was unaffected (wildtype: 0.27µm/s ± 0.2SEM; *Efa6*: 0.30µm/s ±
0.2SEM), but their lifetime was ~1.4 times longer in mutant GCs (Fig. 6M). Compared to
control neurons, Eb1::GFP comets in *Efa6* mutant neurons appeared to vanish less often
when reaching the GC periphery, where they could occasionally be observed to undergo
persistent curved extensions along the periphery (Suppl. Movies not available here).

We reasoned that, if there are more polymerising MTs in GCs of Efa6-depleted neurons, this
should increase their chances of entering into filopodia. In agreement with this prediction, we
found considerably more *Efa6* mutant than control neurons with at least one Eb1 comet or
tubulin-stained MT in GC filopodia (Fig.6J, K). Also per GC, the average number of filopodia
containing comets or MTs was increased, whilst the total number of filopodia per GC was
comparable (~11 per neuron in wild-type and ~10 in *Efa6*). Consistent with this finding, the

lifetime of Eb1 comets persisting at the tip of filopodia was increased from ~2 s in wildtype to
~6 s in *Efa6* mutant neurons (Fig. 6M). Upon live-imaging we even observed highly unusual
cases where comets at the tip of filopodia moved backwards, seemingly pushed back by the
retracting filopodial tip (Suppl. Movies not available here).

Whilst Efa6 depletion clearly favoured MT polymerisation events at the cortex and in filopodia
of GCs, pan-neuronal expression of Efa6-FL::GFP in wildtype background had the opposite
effect. We found that transgenically expressed Efa6-FL::GFP localised to cell bodies, axons
and growth cones, as observed with the endogenous protein (Figs.1B,C). But increased levels
upon over-expression caused a reduction in the number of neurons containing Eb1 comets or
MTs in GC filopodia, and also the number of such filopodia per GC was reduced (green bars
in Fig.6).

Therefore, Efa6 negatively regulates the entry of MTs into GC filopodia likely by eliminating
the overall number of polymerising MTs and reducing their dwell time in filopodia, as would be
in agreement with a role of Efa6 in eliminating MTs at the cell membrane. As detailed in the
discussion, the increased extension of MTs into filopodia in *Efa6* mutant neurons matches the
observed morphological phenotypes: filopodial entry of MTs is considered an important
prerequisite for axon growth and can explain the observed increase in axon length (Prokop et
al., 2013); it could also contribute to increase in branch numbers by promoting GC splitting
events (Acebes and Ferrus, 2000).

<u>Efa6 eliminates MTs in axonal shaft filopodia</u>

We applied the same subcellular readouts to axon shafts where MTs are highly bundled and
expected to be kept away from the membrane. Accordingly, we found that Eb1 comet
numbers and lifetimes in axon shafts were not at all or only moderately affected by loss or
gain of Efa6 (Fig. 7A,B), a behaviour very different from that found in GCs (Fig.6). Other
parameters were comparable: like in GCs, Eb1 comet velocities were not changed in axon
shafts (Fig.7C) and also antero- versus retrograde movement was unaffected (~75% *versus*
~25%; Fig. 7D). Also similar to GCs, axon shafts displayed a strong increase in MTs entering
filopodia: the percentage of neurons containing at least one Eb1 comet or MT in shaft filopodia
was increased (Fig. 7E,F), as was the number of shaft filopodia per neuron that contained Eb1
comets or tubulin-stained MTs (~15-fold in *Efa6* mutant neurons, even when normalised to
axon length; Fig. 7G,H). In contrast, Efa6-FL::GFP caused a strong reduction of these
parameters relative to control neurons (Fig. 7E-H).

Taken together, also in axon shafts our data are consistent with a model in which Efa6
eliminates MTs at the cortex. It seems that Efa6 does not influence MTs which are kept in
axonal bundles, but negatively impacts on the few MTs which leave the axon bundle towards
the cell membrane. By eliminating only MTs that have escaped axonal bundles, Efa6 would
provide a quality control mechanism that prevents MT disorganisation. This is consistent with
the slow onset and gradual increase of MT disorganisation we observed upon Efa6 deficiency
(Fig.4I,J). A condition where off-track MTs are favoured is also expected to promote interstitial
branch formation (Kalil and Dent, 2014; Lewis et al., 2013) and could therefore contribute to
the branching phenotypes we observed (Figs.2 and 3).

<u>The N-terminus of Efa6 can eliminate whole axons in neurons</u>

We next established which region of the Efa6 protein carries MT eliminating capabilities. For

this, we first compared the domain structure and phylogenetic relationship of Efa6 proteins
from 30 species. These analyses revealed that all chordate proteins are rather distant from
invertebrates, and arthropods form a clear subgroup within the invertebrates (Fig. S3A). When
aligning the Efa6 proteins, the C-terminal halves appear relatively consistent, in most species
containing a putative pleckstrin homology (PH; potential membrane association) (Macia et al.,
2008), Sec7 (potential activation of ARF GTPases) and coiled-coil (CC) domain (Franco et al.,
1999) (Fig.8A, S4). In contrast, the N-termini are mainly unstructured and reveals enormous
length differences among species (see more details below); notably, phylogenetic relationship
analyses with the N-terminus alone reveal a very similar tree as the full length comparison
(Fig. S3A,B). We therefore generated two complementary constructs which reflect these two
very different parts of Efa6: *Efa6-ΔCterm-GFP* encoding the entire N-terminal half upstream of
the Sec7 domain, and *Efa6-ΔNterm-GFP* encoding all C-terminal domains from Sec7
onwards.

To functionally assess these Efa6 variants, we developed a transfection method for
Drosophila primary neurons (see Methods) and validated it by transfecting *Efa6-FL-GFP*.
When analysing these transfected primary neurons, we found that they had the same sub-
cellular localisation of Efa6-FL::GFP, reminiscent of neurons expressing the same protein
from transgenic constructs (Fig.1B,C). Axon length upon transfection was even further
decreased than in the transgenic constellation, likely due to higher copy numbers of the
expression construct (Figs. S5B, 2C,D).

We then transfected *Efa6-ΔCterm-GFP* or *Efa6-ΔNterm-GFP*. Both variants localised
throughout neurons (Fig.S5C,H), but only expression of the N-terminus (Efa6-ΔCterm::GFP)
caused phenotypes where axons were severely reduced or frequently even absent. Notably,
Efa6-ΔCterm::GFP induced considerably stronger phenotypes than observed with Efa6-
FL::GFP. To quantify potential axon loss, we labelled neurons with the pan-neuronal marker
Elav (Robinow and White, 1991) and found that only ~36% had an axon when expressing
Efa6-ΔCterm::GFP, as compared to ~57% with Efa6-FL::GFP, and ~74% with either GFP or
Efa6-ΔNterm::GFP (Fig.8B).

These results clearly indicated that the N-terminus is responsible for the axon growth-
inhibiting function of Efa6. The C-terminus which includes the Arf6 activating Sec7 domain
(Huang et al., 2009) appears dispensable for this. Accordingly, also functional deficiency of its
potential effector protein Arf6 in primary neurons (*Arf51F*[GX16[w-]] null mutant allele) had no
obvious effects on axon morphology (data not shown).

<u>Membrane association lowers the effects of N-terminal MT elimination in neurons</u>

As described above, the power of Efa6-ΔCterm::GFP in eliminating axons is considerably
greater than observed with EFA6-FL::GFP. This is likely due to the fact that Efa6-
ΔCterm::GFP lacks the membrane-associating PH domain, so that it can diffuse into the core
area of axons and eliminate MTs that are usually out of reach for the full length Efa6 protein.

To test this possibility, we generated two constructs: one encoding a shorter N-terminal
variant of Efa6 (*Efa6-Nterm-GFP*; Fig.8B), and another where a membrane-associating CAAX
domain was fused to this short N-terminus version (Hancock et al., 1991) (*Efa6-Nterm-GFP-
CAAX*; Fig.8B). Both variants localised throughout neurons (Fig.S5D,I). However, whereas
Efa6-Nterm::GFP showed the same strong axon-abolishing effects as observed with Efa6-
ΔCterm::GFP, the phenotypes induced by the Efa6-Nterm::GFP::CAAX variant were much

weaker and comparable to Efa6-FL::GFP (Fig.8B). Similarly, at 5 DIV, the transgenically
expressed full length and CAAX versions show levels in axonal side branch reductions (Fig.
2H,I).These findings suggest that membrane tethering of the Efa6 N-terminus lowers the
harmful effects it has in axons, with important implications for axon morphology.

<u>Membrane association lowers the effects of N-terminal MT elimination also in fibroblasts</u>

To complement these studies and achieve a higher subcellular resolution, we used NIH3T3
mouse fibroblasts as well-established heterologous cellular models for the study of MT
regulating *Drosophila* proteins (Alves-Silva et al., 2012).

24 hrs after transfection, fibroblasts with moderate Efa6-FL::GFP expression showed strong
GFP staining along the circumference and in areas of membrane folds, indicative of
membrane association (open arrow heads and curved arrow in Figs.S6B and S7B).
Membrane association was further suggested by a strong membrane ruffle phenotype (curved
arrows in Figs.S8B), which was similarly observed when expressing the human Efa6
homologues PSD(1) (Fig. S8E) and reported from other studies with human PSDs (Derrien et
al., 2002; Franco et al., 1999; Macia et al., 2008).

As already observed in neurons, Efa6 was able to eliminate MTs also in fibroblasts: in cells
expressing moderate levels of Efa6-FL::GFP, MTs show a strong tendency to vanish along
the cell fringes (Figs.8B, S9B), consistent with predominantly membrane-based MT
elimination. However, with increasing expression levels, Efa6-FL::GFP was also found in the
cytoplasm, suggesting that membrane-association might become saturated (Fig.S7C). In
these cases, we found a concentration-dependent MT depletion effect, with higher Efa6-FL
expression levels leading to more severe MT loss, and 16% of transfected cells lacking MTs
completely (Fig. S7D).

When transfecting *Efa6-ΔCterm-GFP* or *Efa6-Nterm-GFP* expression constructs, MT
depletion was more than doubled, likewise corroborating our findings in neurons (Fig. 8B). As
expected, the expressed protein failed to localise to membranes; even at low expression
levels, it was found exclusively in the cytoplasm and the nucleus (double chevrons in
Fig.S6C,D) - of which the nuclear localisation is a known phenomenon for GFP-tagged
proteins and not necessarily caused by the N-terminal sequence (Alves-Silva et al., 2012;
Seibel et al., 2007). We next used the N-terminal construct fused to the membrane-
associating CAAX domain and found it to be present at cell edges and absent from the
nucleus (as observed with Efa6-FL::GFP; Figs. S6B,I). Notably, the MT depletion phenotype
was much milder with Efa6-Nterm::GFP::CAAX than with Efa6-Nterm::GFP (Fig. 8B). This
again reproduced our findings in neurons and further supported the conclusion that
membrane localisation of the Efa6 N-terminus correlates with milder MT phenotypes.

<u>The Efa6 C-terminus regulates cortical actin in fibroblasts but does not cause MT elimination</u>

In further agreement with our findings in neurons, we found that the C-terminal construct does
not affect MTs in fibroblasts but has a conserved role in cortical actin regulation instead: Efa6-
ΔNterm::GFP localises to the plasma membrane in fibroblasts and was found in the cytoplasm
only upon increased expression levels (Fig.S6H). But even at high levels, it had no obvious
effects on MT networks, and cells were indistinguishable from GFP-transfected controls
(Fig.8B). Notably, we observed that Efa6-ΔNterm::GFP induced strong membrane ruffling, as
we had similarly observed with the full length construct of Efa6 and human PSD(1) (Fig.

S8B,D,E). This suggests conserved roles of the C-terminus in cortical actin regulation,
because also the human Efa6 homologues PSD(1) and PSD4 use their C-terminal PH and
CC domains to induce actin cytoskeletal re-organisation and membrane ruffle formation, as
observed in TRVb-1 and baby hamster kidney cells (Derrien et al., 2002; Franco et al., 1999;
Macia et al., 2008).

To sum up, our fibroblast studies confirm the expected membrane-tethered nature of the
CAAX and full length variants of Efa6, whereas the the ΔCterm or Nterm versions clearly
localise to the cytoplasm (and nucleus). Studies in both fibroblasts and neurons consistently
reveal that the membrane-tethered versions cause far milder MT-eliminating effects. Since we
found no MT-eliminating activity in the C-terminus, all our findings are in agreement with a
model where endogenous Efa6 displays MT-eliminating functionality in its N-terminus which is
restricted to the cortex via its C-terminus.

<u>The putative PDZ domain and functional SxIP motifs are not required for MT elimination</u>

We next aimed to identify the motifs in the N-terminus required for MT elimination. The N-
terminal half of the *Drosophila* Efa6 protein is mostly unstructured and comprises almost 900
amino acids (aa), which is unusually long compared to most other species (Figs. 8A,C and
S4A). Our CAAX-related experiments with the 410 aa long Efa6-Nterm::GFP variant had
already shown that most if not all MT-eliminating function localises to this stretch of the protein
which contains three different domains/motifs (Fig.8A,B): a putative PDZ domain (aa16-88),
two SxIP motifs surrounded by positive charges (aa 233-6 and 262-5), and a motif of 18aa
displaying 89% similarity with a motif in the N-terminus of *Ce*Efa6 (O'Rourke et al., 2010).

Of these, PDZ domains usually provide anchorage to transmembrane proteins (Ponting et al.,
1997), but might not do so in Efa6 because the N-terminus seems to fail to attach to
membranes in neurons and fibroblasts (Fig.S6C,D). Furthermore, detailed phylogenetic
analyses of the N-terminus, revealed that the PDZ-like domain seems to be a feature mainly
of insect versions of Efa6 (Fig.S4A). We therefore did not analyse it further.

SxIP motifs tend to mediate binding to Eb1 if surrounded by positive charges (Honnappa et
al., 2009), and such a function could potentially aid in capturing MTs for elimination. The SxIP
sites of Efa6 seem indeed to be functional, since we found that fibroblasts expressing Efa6-
Nterm::GFP displayed clear cases of MT tip tracking behaviour, but only in some very rare
cases where MTs had not yet been abolished by the construct (Suppl. Movies not available
here).

To test whether these SxIP motifs are relevant for MT-eliminating functions of Efa6, we
generated an *Efa6-Nterm^ΔSxIP::GFP* variant in which the two motifs were replaced by four
alanines, respectively. Live imaging of fibroblasts transfected with this construct never
revealed any tip tracking, but all other properties of this construct were comparable to Efa6-
Nterm::GFP: it showed the same cytoplasmic and nuclear localisation in fibroblasts and was
found in all compartments of primary *Drosophila* neurons; in both systems it had the same
drastic MT-eliminating effects (Figs.8B, S5E, S6E).

Therefore, Eb1 interaction does not appear to be a major requirement for Efa6-mediated MT
elimination. In agreement with this finding, SxIP motifs are also not widely conserved in other
species, and we could only identify them in a subset of insects (primarily flies) and molluscs,
and a derived SxLP motif in a number of vertebrate/mammalian species (Fig. S4A).

<u>A conserved N-terminal motif is crucial for MT-elimination</u>

We next analysed the conserved 18aa motif, because its worm homologue is known to display
MT-eliminating properties (O'Rourke et al., 2010). To assess potential roles of this putative
MT-eliminating domain (MTED) in the *Drosophila* Efa6 N-terminus, we generated two further
expression constructs (Fig. 8B): *Efa6-Nterm^{ΔMTED}-GFP* (truncated immediately before the
MTED-encoding sequence) and *Efa6-MTED-GFP* (encoding only the MTED).

Similar to Efa6-Nterm::GFP, both proteins localised throughout cell bodies and axons when
transfected into neurons (Fig.S5D,F,G), and to the cytoplasm and nucleus when expressed in
fibroblasts (Fig.S6D,F,G). However, the phenotypes they induced were very different from
each other - but consistent between neurons and fibroblasts (Fig.8B): Efa6-Nterm^{ΔMTED}::GFP
caused no obvious phenotypes (Figs.S5F, S9G), whereas MTED::GFP alone was sufficient to
cause severe loss of axons and entire MT networks in fibroblast, although these effects were
slightly milder than with Efa6-Nterm::GFP (Figs.S5D, 8B and S9D). Notably, transfection rates
with *Efa6-MTED-GFP* in fibroblasts were consistently very low and cell densities were
reduced compared to controls, suggesting that MTED induces cell lethality through an
unknown mechanism. Notwithstanding, our data clearly indicated MTED as a key motif
required for Efa6-induced MT elimination.

Surprisingly, our phylogenetic analyses revealed that also the MTED domain was not well
conserved across the animal kingdom; it was predicted primarily for nematodes, arthropods
and molluscs (Fig. S4A). To test, whether the presence of a MTED motif correlates with MT-
eliminating capabilities, we used a whole range of different constructs: full length versions of
*Ce*Efa6, *Drosophila* Efa6 and all four human PSDs (Fig. 8C), as well as Nterm versions of
*Ce*Efa6, fly Efa6 and human PSD(1) (Fig. 8D). Furthermore, we we deduced a MTED
consensus sequence from 39 *Efa6* genes (details in Fig.S4B, Suppl. Tab. not available here),
identified the most likely human MTED-like sequence [position 31-49aa of PSD(1); Fig. 8A]
and synthesised codon optimised versions of the human as well as the fly and worm MTEDs
(Fig. 8E).

When transfected into fibroblasts, we found a clear correlation: all three fly and all three worm
constructs had strong MT eliminating properties, whereas none of the 6 human constructs
(PSD1-4 full length, PSD(1)-Nterm, PSD(1)-MTED) showed any such effect (Fig.8C-E).
Therefore, the presence of a well conserved canonical MTED seems to be a good predictor
for MT eliminating capabilities of Efa6 proteins.

We next aimed to unravel the mechanism through which MTEDs act, and carried out a series
of experiments. However, experiments with purified Efa6-Nterm::GFP as well as synthetically
generated MTED were not able to reproduce MT-destabilising functions in two independent *in
vitro* assays. Firstly, we observed that Efa6-ΔCterm::GFP and the positive control protein
MCAK::GFP (a MT-destabilising kinesin) both bound to GMPCPP-stabilised MTs, but only
MCAK caused MT destabilisation (Fig.S10A,B). Secondly, also in *Xenopus* oocyte extracts
only MCAK caused destabilisation (Fig.S10C,D); this was surprising because mRNA
injections of *Efa6-Nterm-GFP* mRNA into *Xenopus* eggs generated cell division defects
(Fig.S1F). Therefore, MT defects seem to occur when the construct is expressed in *Xenopus*
oocytes *in vivo* (as is true for *Drosophila* neurons and mouse fibroblasts), but cannot be
reproduced when using purified proteins in the context of the same cytoplasm (or using
GMPCPP-stabilised MTs).

Taken together, Efa6-mediated MT elimination is driven solely or primarily by the MTED motif,
through a mechanism that remains elusive at present. Although this motif and its functions are

restricted to invertebrates, the morphological and sub-cellular phenotypes associated with
Efa6-mediated MT elimination in *Drosophila* neurons provide conceptual understanding that
can serve as a paradigm for mammalian cortical collapse factors and their contributions to
disease-relevant aspects of axon morphology and maintenance (e.g. Kif21A; see Discussion).

<u>Efa6 and Shot cooperate in MT bundle formation and maintenance</u>

As a fundamental conceptual model for axonal MTs, we recently introduced the hypothesis of
"local axon homeostasis". It proposes that the high-force environment inside axons causes
MTs to become disorganised by default, and that MT regulators are required to 'tame' them
into bundles (Prokop, 2016; Voelzmann et al., 2016a). In this model, the cortical MT-
elimination function of Efa6 could serve as a quality control mechanism, eliminating off-track
MTs that have accidentally left the bundle.

To test this possibility, we made use of spectraplakins, which are the most prominent neuronal
factors required for the bundled organisation of axonal MTs in mammals and fly (Bernier and
Kothary, 1998; Dalpe et al., 1998; Sánchez-Soriano et al., 2009; Voelzmann et al., 2017). In
Drosophila, spectraplakins are represented by the single *short stop* (*shot*) gene, and its
functional loss in *Drosophila* primary neurons was shown to cause a severe increase in axonal
off-track MTs (Alves-Silva et al., 2012). Accordingly, we found that the MDI value of *shot*[3]
mutant neurons was rather high (239% ± 14SEM normalised to wild-type controls) compared
to the *Efa6*[GX6[w-]] mutant phenotype at that same stage (155% ± 14SEM ; Fig. 4M). When
combining both mutations in *shot*[3] *Efa6*[GX6[w-]] double mutant neurons (416% ± 24SEM), MT
disorganisation was even further increased. This suggests complementary functions of both
proteins, consistent with our hypothesis that Efa6 can ameliorate *shot* mutant phenotypes by
eliminating a substantial fraction of the aberrant off-track MTs caused by loss of Shot; the fact
that Shot deficiency alone shows MT disorganisation, might indicate that the cortical
elimination function of Efa6 becomes saturated by the sheer amount of off-track MTs in these
neurons (Fig. 9; see Discussion).

In a complementary strategy, we expressed *Efa6-RNAi* in a *shot*[3/+] heterozygous mutant
background, and found that the MDI was clearly enhanced at both 6 HIV and 5 DIV when
compared to knock-down in wild-type background (Fig. 4L). This enhancement was also
observed *in vivo* in our optic lobe model where MTs are labelled with α-tubulin84B-GFP in a
subset of lamina neurons (Fig.4L). In our original experiment, we knocked down *Efa6*
specifically in these neurons and found an increase in axons with unbundled MTs (Fig.5B, E).
Here we combined *Efa6* knock down in those neurons with a heterozygous *shot* mutant
background and found a strong enhancement in axon swellings with MT disorganisation, as
compared to *Efa6* knock-down or *shot*[3/+] heterozygous condition alone (Fig.5). These data
suggest that Efa6 and Shot perform complementary functions that contribute to axonal MT
bundle maintenance also *in vivo*.

Discussion

<u>Efa6 is an essential component of the machinery that regulates MT behaviours underpinning</u>
<u>axon morphology</u>

Axons are the structures that wire the brain, fundamental to nervous system function. To

understand how axons are formed during development, maintained in a plastic state thereafter
but also pushed into degeneration by pathological conditions, we need more profound
knowledge of the dynamics of the MT bundles at their core, and how they are regulated by
MT-binding proteins (Voelzmann et al., 2016a). Here we show that the cortical collapse factor
Efa6 is required for these bundled MT arrangements. We propose that Efa6 acts as a quality
control factor which eliminates off-track MTs that extend away from the MT bundle towards
the axonal plasma membrane. We propose this function to impact on axon growth, branching
and axonal MT bundle maintenance.

Various complementary observations support our conclusions: (1) In neurons, loss of Efa6
causes an increase in Eb1 comet numbers specifically in GCs, and a rise in MTs entering
filopodia throughout axons. (2) Upon loss of Efa6, MTs have longer half-lives and gradually
arrange into winding trajectories instead of parallel ones. (3) Over-expression of Efa6-FL::GFP
fundamentally inverts the above phenotypes. (4) Speed and antero- *versus* retrograde
directionality of MT polymerisation events are unaffected in Efa6-deficient neurons, indicating
that Efa6 is not an instructive regulator of these events. (5) Structure-function analyses of
Efa6 in neurons and fibroblasts demonstrate that its MT-elimination function occurs
compartmentalised at the membrane.

Figure 10 demonstrates how all these findings can be integrated into a consistent model,
based on our current knowledge of MT behaviours in GCs (Dent and Gertler, 2003; Prokop et
al., 2013) and in axon shafts (Fig.9A) (Prokop, 2016; Voelzmann et al., 2016a). We propose
that Efa6-mediated cortical MT elimination exists in a well-balanced equilibrium, sufficient to
eliminate occasional off-track MTs (thus preventing MT disorganisation), but still permitting
intended morphogenetic changes, such as axon growth or branching. Both these
morphogenetic events are enhanced upon loss of Efa6 and reduced upon Efa6 over-
expression, further illustrating the balanced nature of the system. Such a model where active
mechanisms maintaining MTs in bundles protect them against cortical collapse would also
solve the long known conundrum of how axonal MTs extend hundreds of micrometres in the
immediate vicinity of the cell cortex in axons, whereas they tend to interact with the cortex in
non-neuronal cells by being either eliminated or captured (Fukata et al., 2002; Kaverina et al.,
1998).

<u>Roles of Efa6 during axonal growth, branching and MT bundle maintenance</u>

In the case of axon growth, MTs constantly polymerise towards the periphery of GCs and a
fraction of them overcomes elimination and enters filopodia; such events may also be actively
regulated by guidance mechanisms, such as actin-MT cross-linkage mediated by
spectraplakins or drebrin-Eb3 (Alves-Silva et al., 2012; Geraldo et al., 2008). MTs entering
filopodia have the chance of actively contributing to axon extension; this is consistent with the
widely accepted protrusion-engorgement-consolidation model of axon growth which proposes
that stabilised MTs in filopodia can seed axon elongation events (Aletta and Greene, 1988;
Goldberg and Burmeister, 1986; Prokop et al., 2013). Loss of Efa6 would favour this mode of
growth (and thus explain increased axon lengths observed in these neurons) in two ways:
firstly through allowing more MTs to enter filopodia and, secondly, by allowing them to dwell in
filopodia for longer thus enhancing the likelihood of their stabilisation (yellow arrows in
Fig.9C). This scenario could also explain why loss of *Efa6* in *C. elegans* improves axon re-
growth after injury and growth overshoot during development (Chen et al., 2015; Chen et al.,
2011). Furthermore, it provides a potential mechanism for enhanced branching of Efa6-
deficient neurons, since entry of multiple MTs in different filopodia at the same time may

cause simultaneous growth events in the same GC, thus leading to GC splitting as one mode
of branch formation (yellow arrows in Fig.9C) (Acebes and Ferrus, 2000).

Axon branching can also occur along axon shafts, although inhibitory mechanisms at the
shaft's cortex (including Efa6) as well as MT bundling mechanisms seem to prevent this from
happening randomly (Fig.9A). Accordingly, current models propose that collateral branching
requires the active generation (e.g. through MT severing) and then stabilisation of off-track
MTs that can enter shaft filopodia (Kalil and Dent, 2014; Lewis et al., 2013; Yu et al., 2008).
This scenario bears reminiscence with our observations in Efa6 deficient neurons: we find
greater numbers of MTs in shaft filopodia at 6 HIV, which then correlate with enhanced axonal
branch numbers in mature neurons - both in culture and *in vivo* in the adult brain (red arrow in
Fig.9C); this effect was inverted upon Efa6 over-expression. Therefore, also interstitial branch
formation may contribute to the observed *Efa6* mutant phenotypes. It remains to be seen
whether branching events require active regulation of Efa6, for example through signalling
events triggering sub-cellular re-localisation of Efa6 away from this site (Chen et al., 2015), or
the functional inactivation of Efa6. But the simplest mode would be that Efa6 is
homogeneously distributed at well-balanced concentrations, so that the sheer increase in off-
track MTs can outnumber Efa6-mediated elimination (Fig.9D).

This latter model view is supported by increased off-track MTs in *shot* mutant neurons: Efa6
can eliminate many but not all of these, as is indicated by strong MT disorganisation in single
shot mutant conditions which becomes enhanced when Efa6 is taken out in addition (*shot*
Efa6 double-mutant neurons). Such a scenario would also explain the shortened lifetime of
MT comets observed in *shot* mutant neurons (Alves-Silva et al., 2012), potentially caused by
the MT-eliminating functions of Efa6. In this scenario, it makes sense that loss of Shot causes
substantial MT disorganisation from early development onwards. In contrast, off-track MTs are
rare events in the presence of Shot so that *Efa6*-deficient phenotypes are initially modest; but
off-track MTs are given the chance to persist and accumulate over time, thus leading to
gradual increase in MT disorganisation (Fig.4I,J). We therefore propose that Efa6 plays a
continued role in MT quality control, from developmental stages onwards right through into the
late adult stage. Persistent quality control is required because MT polymerisation continues in
mature neurons as a mechanism to prevent senescence of axonal MT bundles (Qu et al.,
2017; Voelzmann et al., 2016a). Such a model of continued quality control is supported by our
finding that Efa6 is expressed at all these stages on the one hand (Fig. 1), and by the late
onset of MT disorganisation upon delayed knock-down in mature neurons on the other (Fig.
4K).

Clearly, the MT disorganisation phenotypes are far less pronounced *in vivo* than in culture, but
this is expected when considering that axons *in vivo* tend to be arranged into closely adhered
nerve tracts and surrounded by glial cells, i.e. have a much more rigid plasma membrane and
cortex, able to resist MTs pushing out to form swellings and areas of disorganisation.

<u>Evolutionary and mechanistic considerations of Efa6 function</u>

We found that the mammalian Efa6 proteins encoded by four *PSD* genes, do not contain a
canonical MTED and lack any MT collapse function. However, we propose that comparable
functional contributions are taken over by different types of cortical MT collapse factors. For
example, the kinesin-4 family member Kif21A was shown to act as a cortical collapse factor
and is expressed throughout neurons in mammals (van der Vaart et al., 2013). The R954W
mutation of Kif21A links to the eye movement disorder 'congenital fibrosis of extraocular

muscles' (OMIM reference #135700) (Tiab et al., 2004) and was shown to cause a shift of the
protein's localisation in cultured hippocampal neurons from axons towards GCs, expected to
cause a loss-of-function condition in the axon and gain-of-function in GCs (van der Vaart et
al., 2013). Increased protein levels in GCs correlate with reduced axon growth (as similarly
observed with Efa6 over-expression); decreased levels in proximal axons correlate with a
local increase in side branches (as similarly observed upon Efa6 loss-of-function). Therefore,
our findings with Efa6 can serve as a paradigm to guide research into the disease-relevant
roles of cortical collapse factors in higher organisms.

Our studies together with previous work on *Ce*Efa6 (O'Rourke et al., 2010) clearly indicate a
central role for the MTED in MT elimination, but the molecular mechanisms through which
they perform this function remain unfortunately elusive for now. In principle, MT elimination
can occur from their ends (like kinesin-8 or the kinesin-13 Kif2/MCAK) (Brouhard and Rice,
2014) or by destroying MTs along their shafts (like katanin or spastin) (Sharp and Ross,
2012). It can be achieved by depolymerisation, severing, buckling and breaking, or extrusion
of tubulins along their shafts. Since the MTED is only a very short peptide, it would not be
surprising if it acted indirectly through activating MT destabilisers or inhibiting MT
polymerisers, which is why we complemented our *in vitro* analyses with studies in *Xenopus*
oocyte extracts (Fig. S10C). But it could also block MT polymerisation processes by binding
tubulins directly. In any case, the small size of MTEDs might come in handy as experimental
tools to eliminate MTs, with potentially complementary properties to existing factors such as
kinesin-13s (Moore et al., 2005; Schimizzi et al., 2010), Stathmin (Marklund et al., 1996),
Spastin or Katanin (Eckert et al., 2012; Trotta et al., 2004).

In contrast, other regions of the Efa6 N-terminus are poorly conserved even among
invertebrates and seem not to contribute to MT elimination in prominent ways. For example,
SxIP sites of Efa6 mediate plus end tracking (Fig.M5), and MT plus end capture seemed an
attractive candidate mechanism to facilitate cortical MT collapse. However, we found the SxIP
sites to be dispensable for MT elimination (Fig.8B), and our *in silico* analyses of Efa6 proteins
failed to reveal any other Eb1-binding motifs, such as CAP-Gly or LxxPTPh sites (Akhmanova
and Steinmetz, 2008; Kumar et al., 2017). The scenario for Efa6 appears therefore
comparable to that of MCAK which is a prominent tip-tracker but does not require this property
for its MT depolymerising activity (Moore et al., 2005).

However, the C-terminal domains of Efa6 seem to display some degree of functional
conservation. So far, work on mammalian PSDs has revealed functions for C-terminal
domains in regulating ARF6, ARF1 or ARL14 during actin cytoskeletal reorganisation and
membrane ruffling, tumour formation and immune regulation (Derrien et al., 2002; Paul et al.,
2011; Pils et al., 2005). Our finding that PSD(1) and C-terminal Efa6 constructs cause similar
membrane ruffling phenotypes in fibroblasts (Figs.S6 and S8) suggests that some conserved
functions reside in this region and might contribute, together with N-terminally mediated MT
collapse, to the neuronal or non-neuronal defects that cause semi-lethality displayed by *Efa6*
mutant flies (data not shown).

<u>Main conclusions and future perspectives</u>

We propose Efa6 to act as a cortical collapse factor which is important for the regulation of
axonal MTs and, in turn, axon growth, maintenance and branching. Although this function is
evolutionarily not widely conserved, our findings provide a helpful paradigm for studies on
other classes of cortical collapse factors also in mammalian neurons. Further important

research avenues will be to identify the mechanism of Efa6-mediated MT elimination, not only to better understand how it can be regulated in axons but also to explore potential roles for MTEDs as molecular tools in cell biological research.

Acknowledgements

This work was made possible through funding by the BBSRC (BB/I002448/1, BB/P020151/1, BB/L000717/1, BB/M007553/1) to A.P., by parents as well as the Faculty of Life Sciences to Y.Q., by the Leverhulme Trust to I.H. (ECF-2017-247), by the German Research Council (DFG) to A.V. (VO 2071/1-1), and by the BBSRC to N.S.S. (BB/M007456/1) and KD (BB/J005983/1). The Manchester Bioimaging Facility microscopes used in this study were purchased with grants from the BBSRC, The Wellcome Trust and The University of Manchester Strategic Fund. The Fly Facility has been supported by funds from The University of Manchester and the Wellcome Trust (087742/Z/08/Z). We thank Tom Millard and Marvin Bentley for very helpful comments on the manuscript, Simon Lowell for advice on the phylogenetic analyses, Hiro Ohkura for kindly providing DmEb1 antibody and Andrew Chisholm for the C.elegans Efa6 and human PSD constructs. Stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study.

References

Abascal, F., Zardoya, R., and Posada, D. (2005). ProtTest: selection of best-fit models of protein evolution. Bioinformatics *21*, 2104-2105.

Acebes, A., and Ferrus, A. (2000). Cellular and molecular features of axon collaterals and dendrites. Trends Neurosci *23*, 557-565.

Adalbert, R., and Coleman, M.P. (2012). Axon pathology in age-related neurodegenerative disorders. Neuropathol Appl Neurobiol *39*, 90–108.

Akhmanova, A., and Steinmetz, M.O. (2008). Tracking the ends: a dynamic protein network controls the fate of microtubule tips. Nat Rev Mol Cell Biol *9*, 309-322.

Aletta, J.M., and Greene, L.A. (1988). Growth cone configuration and advance: a time-lapse study using video-enhanced differential interference contrast microscopy. J Neurosci *8*, 1425-1435.

Allan, V.J. (1993). Assay of membrane motility in interphase and metaphase Xenopus extracts. Methods Cell Biol *39*, 203-226.

Allan, V.J., and Vale, R.D. (1991). Cell cycle control of microtubule-based membrane transport and tubule formation in vitro. J Cell Biol *113*, 347-359.

Alves-Silva, J., Sánchez-Soriano, N., Beaven, R., Klein, M., Parkin, J., Millard, T., Bellen, H., Venken, K.J.T., Ballestrem, C., Kammerer, R.A.*, et al.* (2012). Spectraplakins promote microtubule-mediated axonal growth by functioning as structural microtubule-associated proteins and EB1-dependent +TIPs (Tip Interacting Proteins). J Neurosci *32*, 9143-9158.

Bernier, G., and Kothary, R. (1998). Prenatal onset of axonopathy in Dystonia musculorum mice. Dev Genet *22*, 160-168.

Bettencourt da Cruz, A., Schwarzel, M., Schulze, S., Niyyati, M., Heisenberg, M., and Kretzschmar, D. (2005). Disruption of the MAP1B-related protein FUTSCH leads to changes in the neuronal cytoskeleton, axonal transport defects, and progressive neurodegeneration in *Drosophila*. Mol Biol Cell *16*, 2433-2442.

Brouhard, G.J., and Rice, L.M. (2014). The contribution of aß-tubulin curvature to microtubule dynamics. J Cell Biol *207*, 323-334.

Chen, L., Chuang, M., Koorman, T., Boxem, M., Jin, Y., and Chisholm, A.D. (2015). Axon injury triggers EFA-6 mediated destabilization of axonal microtubules via TACC and doublecortin like kinase. eLife *4*, e08695.

Chen, L., Wang, Z., Ghosh-Roy, A., Hubert, T., Yan, D., O'Rourke, S., Bowerman, B., Wu, Z., Jin, Y., and Chisholm, A.D. (2011). Axon regeneration pathways identified by systematic genetic screening in *C. elegans*. Neuron *71*, 1043-1057.

Crooks, G.E., Hon, G., Chandonia, J.M., and Brenner, S.E. (2004). WebLogo: a sequence logo generator. Genome Res *14*, 1188-1190.

D'Souza-Schorey, C., and Chavrier, P. (2006). ARF proteins: roles in membrane traffic and beyond. Nature Rev Molec Cell Biol *7*, 347.

Dalpe, G., Leclerc, N., Vallee, A., Messer, A., Mathieu, M., De Repentigny, Y., and Kothary, R. (1998). Dystonin is essential for maintaining neuronal cytoskeleton organization. Mol Cell Neurosci *10*, 243-257.

Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2011). ProtTest 3: fast selection of best-fit models of protein evolution. Bioinformatics *27*, 1164-1165.

Debanne, D., Campanac, E., Bialowas, A., Carlier, E., and Alcaraz, G. (2011). Axon physiology. Physiol Rev *91*, 555-602.

Dent, E.W., and Gertler, F.B. (2003). Cytoskeletal dynamics and transport in growth cone motility and axon guidance. Neuron *40*, 209-227.

Dent, E.W., Gupton, S.L., and Gertler, F.B. (2011). The growth cone cytoskeleton in axon outgrowth and guidance. Cold Spring Harb Perspect Biol *3*, a001800.

Derrien, V., Couillault, C., Franco, M., Martineau, S., Montcourrier, P., Houlgatte, R., and Chavrier, P. (2002). A conserved C-terminal domain of EFA6-family ARF6-guanine nucleotide exchange factors induces lengthening of microvilli-like membrane protrusions. Journal of cell science *115*, 2867-2879.

Eckert, T., Le, D.T.-V., Link, S., Friedmann, L., and Woehlke, G. (2012). Spastin's microtubule-binding properties and comparison to katanin. PLoS One *7*, e50161.

Elliott, S.L., Cullen, C.F., Wrobel, N., Kernan, M.J., and Ohkura, H. (2005). EB1 is essential during *Drosophila* development and plays a crucial role in the integrity of chordotonal mechanosensory organs. Mol Biol Cell *16*, 891-901.

Fang, Y., and Bonini, N.M. (2012). Axon degeneration and regeneration: insights from *Drosophila* models of nerve injury. Annu Rev Cell Dev Biol *28*, 575-597.

Fassier, C., Tarrade, A., Peris, L., Courageot, S., Mailly, P., Dalard, C., Delga, S., Roblot, N., Lefevre, J., Job, D.*, et al.* (2013). Microtubule-targeting drugs rescue axonal swellings in cortical neurons from spastin knockout mice. Dis Model Mech *6*, 72-83.

Fiala, J.C., Feinberg, M., Peters, A., and Barbas, H. (2007). Mitochondrial degeneration in dystrophic neurites of senile plaques may lead to extracellular deposition of fine filaments. Brain Struct Funct *212*, 195-207.

Franco, M., Peters, P.J., Boretto, J., van Donselaar, E., Neri, A., D'Souza-Schorey, C., and Chavrier, P. (1999). EFA6, a sec7 domain-containing exchange factor for ARF6, coordinates membrane recycling and actin cytoskeleton organization. EMBO J *18*, 1480-1491.

Fukata, M., Watanabe, T., Noritake, J., Nakagawa, M., Yamaga, M., Kuroda, S., Matsuura, Y., Iwamatsu, A., Perez, F., and Kaibuchi, K. (2002). Rac1 and Cdc42 capture microtubules through IQGAP1 and CLIP-170. Cell *109*, 873–885.

Geraldo, S., Khanzada, U.K., Parsons, M., Chilton, J.K., and Gordon-Weeks, P.R. (2008). Targeting of the F-actin-binding protein drebrin by the microtubule plus-tip protein EB3 is required for neuritogenesis. Nat Cell Biol *10*, 1181-1189.

Goldberg, D.J., and Burmeister, D.W. (1986). Stages in axon formation: observations of growth of *Aplysia* axons in culture using video-enhanced contrast-differential interference contrast microscopy. J Cell Biol *103*, 1921-1931.

Goujon, M., McWilliam, H., Li, W., Valentin, F., Squizzato, S., Paern, J., and Lopez, R. (2010). A new bioinformatics analysis tools framework at EMBL-EBI. Nucleic Acids Res *38*, W695-699.

Grieder, N.C., de Cuevas, M., and Spradling, A.C. (2000). The fusome organizes the microtubule network during oocyte differentiation in *Drosophila*. Development *127*, 4253-4264.

Han, M.V., and Zmasek, C.M. (2009). phyloXML: XML for evolutionary biology and comparative genomics. BMC Bioinformatics *10*, 356.

992 Hancock, J.F., Cadwallader, K., Paterson, H., and Marshall, C.J. (1991). A CAAX or a CAAL motif and
993   a second signal are sufficient for plasma membrane targeting of ras proteins. EMBO J *10*, 4033-
994   4039.

995 Hassan, B.A., Bermingham, N.A., He, Y., Sun, Y., Jan, Y.N., Zoghbi, H.Y., and Bellen, H.J. (2000).
996   *atonal* regulates neurite arborization but does not act as a proneural gene in the *Drosophila* brain.
997   Neuron *25*, 549-561.

998 Heasman, J. (2006). Patterning the early Xenopus embryo. Development *133*, 1205-1217.

999 Helenius, J., Brouhard, G., Kalaidzidis, Y., Diez, S., and Howard, J. (2006). The depolymerizing kinesin
1000   MCAK uses lattice diffusion to rapidly target microtubule ends. Nature *441*, 115-119.

1001 Hinrichs, M.H., Jalal, A., Brenner, B., Mandelkow, E., Kumar, S., and Scholz, T. (2012). Tau protein
1002   diffuses along the microtubule lattice. J Biol Chem *287*, 38559-38568.

1003 Honnappa, S., Gouveia, S.M., Weisbrich, A., Damberger, F.F., Bhavesh, N.S., Jawhari, H., Grigoriev, I.,
1004   van Rijssel, F.J., Buey, R.M., Lawera, A.*, et al.* (2009). An EB1-binding motif acts as a microtubule
1005   tip localization signal. Cell *138*, 366-376.

1006 Huang, J., Zhou, W., Dong, W., Watson, A.M., and Hong, Y. (2009). From the Cover: Directed, efficient,
1007   and versatile modifications of the *Drosophila* genome by genomic engineering. Proc Natl Acad Sci U
1008   S A *106*, 8284-8289.

1009 Kalil, K., and Dent, E.W. (2014). Branch management: mechanisms of axon branching in the
1010   developing vertebrate CNS. Nat Rev Neurosci *15*, 7-18.

1011 Kaverina, I., Rottner, K., and Small, J.V. (1998). Targeting, capture, and stabilization of microtubules at
1012   early focal adhesions. J Cell Biol *142*, 181-190.

1013 Kolodziej, P.A., Jan, L.Y., and Jan, Y.N. (1995). Mutations that affect the length, fasciculation, or ventral
1014   orientation of specific sensory axons in the *Drosophila* embryo. Neuron *15*, 273-286.

1015 Krieg, M., Stühmer, J., Cueva, J.G., Fetter, R., Spilker, K., Cremers, D., Shen, K., Dunn, A.R., and
1016   Goodman, M.B. (2017). Genetic defects in β-spectrin and tau sensitize C. elegans axons to
1017   movement-induced damage via torque-tension coupling. Elife *6*, e20172.

1018 Kumar, A., Manatschal, C., Rai, A., Grigoriev, I., Degen, M.S., Jaussi, R., Kretzschmar, I., Prota, A.E.,
1019   Volkmer, R., Kammerer, R.A.*, et al.* (2017). Short Linear Sequence Motif LxxPTPh Targets Diverse
1020   Proteins to Growing Microtubule Ends. Structure *25*, 924-932 e924.

1021 Küppers-Munther, B., Letzkus, J., Lüer, K., Technau, G., Schmidt, H., and Prokop, A. (2004). A new
1022   culturing strategy optimises *Drosophila* primary cell cultures for structural and functional analyses.
1023   Dev Biol *269*, 459-478.

1024 Lewis, T.L., Jr., Courchet, J., and Polleux, F. (2013). Cell biology in neuroscience: Cellular and
1025   molecular mechanisms underlying axon formation, growth, and branching. J Cell Biol *202*, 837-848.

1026 Lowery, L.A., and van Vactor, D. (2009). The trip of the tip: understanding the growth cone machinery.
1027   Nat Rev Mol Cell Biol *10*, 332-343.

1028 Luo, L., Liao, Y.J., Jan, L.Y., and Jan, Y.N. (1994). Distinct morphogenetic functions of similar small
1029   GTPases: *Drosophila* Drac1 is involved in axonal outgrowth and myoblast fusion. Genes Dev *8*,
1030   1787-1802.

1031 Macia, E., Partisani, M., Favard, C., Mortier, E., Zimmermann, P., Carlier, M.F., Gounon, P., Luton, F.,
1032   and Franco, M. (2008). The pleckstrin homology domain of the Arf6-specific exchange factor EFA6
1033   localizes to the plasma membrane by interacting with phosphatidylinositol 4,5-bisphosphate and F-
1034   actin. J Biol Chem *283*, 19836-19844.

1035 Marklund, U., Larsson, N., Gradin, H.M., Brattsand, G., and Gullberg, M. (1996). Oncoprotein 18 is a
1036   phosphorylation-responsive regulator of microtubule dynamics. EMBO J *15*, 5290-5298.

1037 McWilliam, H., Li, W., Uludag, M., Squizzato, S., Park, Y.M., Buso, N., Cowley, A.P., and Lopez, R.
1038   (2013). Analysis Tool Web Services from the EMBL-EBI. Nucleic Acids Res *41*, W597-600.

1039 Medana, I.M., and Esiri, M.M. (2003). Axonal damage: a key predictor of outcome in human CNS
1040   diseases. Brain *126*, 515-530.

1041 Moore, A.T., Rankin, K.E., von Dassow, G., Peris, L., Wagenbach, M., Ovechkina, Y., Andrieux, A.,
1042   Job, D., and Wordeman, L. (2005). MCAK associates with the tips of polymerizing microtubules. J
1043   Cell Biol *169*, 391-397.

1044 O'Rourke, S.M., Christensen, S.N., and Bowerman, B. (2010). *Caenorhabditis elegans* EFA-6 limits
1045   microtubule growth at the cell cortex. Nat Cell Biol *12*, 1235-1241.

1046 Patel, J.T., Belsham, H.R., Rathbone, A.J., Wickstead, B., Gell, C., and Friel, C.T. (2016). The family-
1047   specific alpha4-helix of the kinesin-13, MCAK, is critical to microtubule end recognition. Open Biol *6*.

Paul, P., van den Hoorn, T., Jongsma, Marlieke L.M., Bakker, Mark J., Hengeveld, R., Janssen, L., Cresswell, P., Egan, David A., van Ham, M., ten Brinke, A.*, et al.* (2011). A Genome-wide Multidimensional RNAi Screen Reveals Pathways Controlling MHC Class II Antigen Presentation. Cell *145*, 268-283.

Pils, D., Horak, P., Gleiss, A., Sax, C., Fabjani, G., Moebus, V.J., Zielinski, C., Reinthaller, A., Zeillinger, R., and Krainer, M. (2005). Five genes from chromosomal band 8p22 are significantly down-regulated in ovarian carcinoma. Cancer *104*, 2417-2429.

Ponting, C.P., Phillips, C., Davies, K.E., and Blake, D.J. (1997). PDZ domains: targeting signalling molecules to sub-membranous sites. Bioessays *19*, 469-479.

Prokop, A. (2013a). The intricate relationship between microtubules and their associated motor proteins during axon growth and maintenance. Neur Dev *8*, 17.

Prokop, A. (2013b). A rough guide to *Drosophila* mating schemes. figshare, dx.doi.org/10.6084/m6089.figshare.106631.

Prokop, A. (2016). Fruit flies in biological research. Biological Sciences Review *28*, 10-14.

Prokop, A., Beaven, R., Qu, Y., and Sánchez-Soriano, N. (2013). Using fly genetics to dissect the cytoskeletal machinery of neurons during axonal growth and maintenance. J Cell Sci *126*, 2331-2341.

Prokop, A., Küppers-Munther, B., and Sánchez-Soriano, N. (2012). Using primary neuron cultures of *Drosophila* to analyse neuronal circuit formation and function. In The making and un-making of neuronal circuits in *Drosophila*, B.A. Hassan, ed. (New York, Humana Press), pp. 225-247.

Prokop, A., and Meinertzhagen, I.A. (2006). Development and structure of synaptic contacts in *Drosophila*. Semin Cell Dev Biol *17*, 20-30.

Qu, Y., Hahn, I., Webb, S.E.D., Pearce, S.P., and Prokop, A. (2017). Periodic actin structures in neuronal axons are required to maintain microtubules. Mol Biol Cell *28* 296-308.

Robinow, S., and White, K. (1991). Characterization and spatial distribution of the Elav protein during *Drosophila melanogaster* development. J Neurobiol *22*, 443-461.

Rogers, S.L., Rogers, G.C., Sharp, D.J., and Vale, R.D. (2002). *Drosophila* EB1 is important for proper assembly, dynamics, and positioning of the mitotic spindle. J Cell Biol *158*, 873-884.

Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Hohna, S., Larget, B., Liu, L., Suchard, M.A., and Huelsenbeck, J.P. (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol *61*, 539-542.

Sánchez-Soriano, N., Gonçalves-Pimentel, C., Beaven, R., Haessler, U., Ofner, L., Ballestrem, C., and Prokop, A. (2010a). *Drosophila* growth cones: a genetically tractable platform for the analysis of axonal growth dynamics. Dev Neurobiol *70*, 58-71.

Sánchez-Soriano, N., Gonçalves-Pimentel, C., Beaven, R., and Prokop, A. (2010b). Using *Drosophila* growth cones to dissect F-actin network regulation at the cellular level. Paper presented at: Actin 2009 (Bristol).

Sánchez-Soriano, N., Travis, M., Dajas-Bailador, F., Goncalves-Pimentel, C., Whitmarsh, A.J., and Prokop, A. (2009). Mouse ACF7 and *Drosophila* Short stop modulate filopodia formation and microtubule organisation during neuronal growth. Journal of cell science *122*, 2534-2542.

Schimizzi, G.V., Currie, J.D., and Rogers, S.L. (2010). Expression levels of a kinesin-13 microtubule depolymerase modulates the effectiveness of anti-microtubule agents. PLoS One *5*, e11381.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*, 676-682.

Seibel, N.M., Eljouni, J., Nalaskowski, M.M., and Hampe, W. (2007). Nuclear localization of enhanced green fluorescent protein homomultimers. Analytical Biochemistry *368*, 95-99.

Sharp, D.J., and Ross, J.L. (2012). Microtubule-severing enzymes at the cutting edge. Journal of cell science *125*, 2561-2569.

Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Soding, J.*, et al.* (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol *7*, 539.

Stefana, M.I., Driscoll, P.C., Obata, F., Pengelly, A.R., Newell, C.L., MacRae, J.I., and Gould, A.P. (2017). Developmental diet regulates *Drosophila* lifespan via lipid autotoxins. Nat Commun *8*, 1384.

Tarrade, A., Fassier, C., Courageot, S., Charvin, D., Vitte, J., Peris, L., Thorel, A., Mouisel, E., Fonknechten, N., Roblot, N.*, et al.* (2006). A mutation of spastin is responsible for swellings and

impairment of transport in a region of axon characterized by changes in microtubule composition.
Hum Mol Genet *15*, 3544-3558.

Tiab, L., Manzi, V.d.A., Borruat, F.-X., Munier, F., and Schorderet, D. (2004). Mutation analysis of
KIF21A in congenital fibrosis of the extraocular muscles (CFEOM) patients. Ophthalmic Genetics *25*,
241-246.

Trotta, N., Orso, G., Rossetto, M.G., Daga, A., and Broadie, K. (2004). The hereditary spastic
paraplegia gene, *spastin*, regulates microtubule stability to modulate synaptic structure and function.
Current Biology *14*, 1135-1147.

Vale, R.D., and Toyoshima, Y.Y. (1988). Rotation and translocation of microtubules in vitro induced by
dyneins from Tetrahymena cilia. Cell *52*, 459-469.

van der Vaart, B., van Riel, Wilhelmina E., Doodhi, H., Kevenaar, Josta T., Katrukha, Eugene A., Gumy,
L., Bouchet, Benjamin P., Grigoriev, I., Spangler, Samantha A., Yu, Ka L.*, et al.* (2013). CFEOM1-
associated kinesin KIF21A is a cortical microtubule growth inhibitor. Developmental Cell *27*, 145-
160.

Voelzmann, A., Hahn, I., Pearce, S., Sánchez-Soriano, N.P., and Prokop, A. (2016a). A conceptual
view at microtubule plus end dynamics in neuronal axons. Brain Res Bulletin *126*, 226-237.

Voelzmann, A., Liew, Y.-T., Qu, Y., Hahn, I., Melero, C., Sánchez-Soriano, N., and Prokop, A. (2017).
Drosophila Short stop as a paradigm for the role and regulation of spectraplakins. Sem Cell Dev Biol
69, 40-57.

Voelzmann, A., Okenve-Ramos, P., Qu, Y., Chojnowska-Monga, M., del Caño-Espinel, M., Prokop, A.,
and Sánchez-Soriano, N. (2016b). Tau and spectraplakins promote synapse formation and
maintenance through Jun kinase and neuronal trafficking. eLife *5*, e14694.

Wang, J.T., Medress, Z.A., and Barres, B.A. (2012). Axon degeneration: molecular mechanisms of a
self-destruction pathway. J Cell Biol *196*, 7-18.

Yu, W., Qiang, L., Solowska, J.M., Karabay, A., Korulu, S., and Baas, P.W. (2008). The microtubule-
severing proteins spastin and katanin participate differently in the formation of axonal branches. Mol
Biol Cell *19*, 1485-1498.

Zeidler, M.P., Tan, C., Bellaiche, Y., Cherry, S., Hader, S., Gayko, U., and Perrimon, N. (2004).
Temperature-sensitive control of protein activity by conditionally splicing inteins. Nat Biotech *22*,
871-876.

Zschätzsch, M., Oliva, C., Langen, M., De Geest, N., Ozel, M.N., Williamson, W.R., Lemon, W.C.,
Soldano, A., Munck, S., Hiesinger, P.R.*, et al.* (2014). Regulation of branching dynamics by axon-
intrinsic asymmetries in Tyrosine Kinase Receptor signaling. Elife *3*, e01699.

1137

Figures

1139

Fig. 1 Efa6 is expressed throughout neurons at all developmental stages

A-E) Images of primary *Drosophila* neurons at 6HIV or 5DIV (at indicated bottom right),

stained for tubulin (magenta) and GFP (green); control neurons are wild-type (wt) or express

elav-Gal4-driven nuclear GFP (*elav / nl-GFP*), whereas further neurons are either derived

from the endogenously tagged Efa6::GFP line or express Efa6-FL::GFP under the control of

sca-Gal4 (*sca / Efa6-FL::GFP*); asterisks indicate cell bodies and arrow the axon tips. F-I)

Late larval CNSs at about 4d of development from egg lay (L3; F,G) and adult CNSs at over

10d of development (H,I) derived from control wild-type animals (wt) or the Efa6::GFP line

(*Efa6::GFP*), stained for GFP and actin (Phalloidin, only larval preparations); OL, optic lobe;

Br, central brain; vNC, ventral nerve cord. Scale bar in A represent 15µm in A-C, 25µm in D

and E, 75µm in F and G, 130µm in H and I.

1151

Fig. 2. Efa6 regulates axonal length and branch numbers in primary *Drosophila* neurons.

Examples of primary *Drosophila* neurons at 6HIV (A-C) or at 5DIV (E-H), all stained for actin

(magenta) and tubulin (green); neurons are either wild-type controls (A, E), Efa6-deficient (B,
F), expressing Efa6-FL::GFP (C, G), or expressing Efa6-Nterm-CAAX::GFP (H); asterisks
indicate cell bodies, arrows point at axon tips, arrow heads at axon branches, the curved
arrow at an area of MT disorganisation; the scale bar in G represents 10µm in A-C and 20µm
in E-H. Quantification of axon lengths at 6HIV (D) and of axonal branch numbers (I); different
genotypes are colour-coded: grey, wild-type controls; blue, different *Efa6* loss-of-function
conditions; green, neurons over-expressing Efa6 variants; data represent fold-change relative
to wild-type controls (indicated as horizontal dashed "ctrl" line); they are shown as single data
points and a bar indicating mean ± SEM data; P values from Mann-Whitney tests are given
above each column, sample numbers at the bottom of each bar.

1164

Fig. 3. Efa6 regulates axon branching in the adult fly brain. A-F) Brains (medulla region of the
optic lobe in oblique view) of young (2-5 d after eclosure; top) and old flies (15-18 d; bottom)
driving *UAS-myr-tdTomato* via the *ato-Gal4* driver in dorsal cluster neurons (example neurons
are traced in magenta for axons and green for side branches); flies are either wild-type
controls (left) or display *ato-Gal4*-driven knock-down of Efa6 (middle) or over-expression of
Efa6-FL::GFP (right). G,H) Quantification of data for wild-type (wt; grey), Efa6 knock-down
(blue) and Efa6-FL::GFP over-expression (green), displaying the number of primary branches
per axon (G), all representing fold-change relative to wild-type controls (indicated as horizontal
dashed "ctrl" line); data are shown as bars indicating mean ± SEM in G accompanied by
single data point and binary score normalised to wt in H.; P values from Mann-Whitney or Chi^2
tests are respectively given above each column, sample numbers at the bottom of each bar
(in G number of medullas before and number of axons after slash). Scale bar in D represents
60µm in all figures.

1178

Fig. 4. Efa6 helps to maintain axonal MT bundles in *Drosophila* neurons. A-H) Images of
primary neurons at 6HIV (left), 5DIV (middle) and 10DIV (right), stained for tubulin (green) and
actin (magenta), derived from embryos that were either wildtype (wt, A-C), *Efa6* null mutant
(D-F), homozygous for *shot^3* (G) or *shot^3 Efa6$^{GX6[w-]}$* double-mutant (shot Efa6) (H); arrows
point at axon tip, arrow heads at areas of MT disorganisation, and asterisks indicate the cell
bodies; the scale bar in A represents 10µm for 6HIV neurons and 25µm for 5DIV and 10DIV
neurons. I-M) Quantitative analyses of MT disorganisation (measured as MT disorganisation
index, MDI) in different experimental contexts (as indicated above graphs); different
genotypes are colour-coded: grey, wild-type controls; blue, *Efa6* loss-of-function; light/dark
orange, *shot^3* in hetero-/homozygosis; green, neurons over-expressing Efa-FL::GFP; data are
shown as single data points and a bar indicating mean ± SEM, all representing fold-change
relative to wild-type controls (indicated as horizontal dashed "ctrl" line); P values from Mann-
Whitney tests in I, J) and Kruskal–Wallis one-way ANOVA with *post hoc* Dunn's test in K-M)
are given above each column, sample numbers at the bottom of each bar.

1193

Fig. 5 Efa6 is required for axonal MT bundle maintenance in adult fly brains. A-C) Medulla
region of adult brains at 26-27 days after eclosure, all carrying the *GMR31F10-Gal4* driver and
UAS-α-tubulin84B-GFP (*GMR-tub*) which together stain MTs in a subset of lamina neuron
axons that terminate in the medulla; the other specimens co-express *Efa6-RNAi* either alone
(*GMR-tub-Efa6^{IR}* in B) or combined with a *shot$^{3/+}$* heterozygous mutant background (*GMR-tub-
Efa6^{IR} shot$^{3/+}$* in C); white/black arrows indicate axonal swellings without/with MT

disorganisation; rectangles outlined by red dashed lines are shown as 2.5 fold magnified
insets where white arrow heads point at disorganised MTs; the scale bar in A represents
15µm in all images. D, E) Quantitative analyses of all axonal swelling (D) or swellings with MT
disorganisation (E); different genotypes are colour-coded as indicated; bars show mean ±
SEM, all representing fold-change relative to wild-type controls (indicated as horizontal
dashed line); P values from Kruskal–Wallis one-way tests are given above each column,
sample numbers at the bottom of each bar.

1207

Fig. 6. Efa6 regulates MT behaviours in GCs. A-H') Examples of primary neurons at 6HIV
which are either wild-type controls (top), Efa6-deficient (middle) or expressing Efa6-FL::GFP
(bottom); neurons were either imaged live for Eb1::GFP (green in A,D) or fixed and labelled
for Eb1 and tubulin (B,E,G; as colour-coded) or actin and tubulin (C,F,H; as colour coded;
tubulin shown as single channel image on the right); asterisks indicate cell bodies, white
arrows the tips of GCs, open arrows the tips of MT bundles and arrow heads MTs or Eb1
comets in filopodial processes; the GC in G is outlined with a white dashed line; scale bar in D
represents 5µm in all images. I-M) Quantitative analyses of MT behaviours in GCs of such
neurons as indicated above each graph (for axon shaft analyses see Fig.7). Different
genotypes are colour-coded: grey, wild-type controls; blue, different *Efa6* loss-of-function
conditions; green, neurons over-expressing Efa-FL; data are shown as single data points and
a bar indicating mean ± SEM, all representing fold-increase relative to wild-type controls
(indicated as horizontal dashed "ctrl" line); P values from Mann-Whitney tests are given above
each column, sample numbers at the bottom of each bar. The graph in L shows percentages
of neurons with (light shade) or without (dark shade) any MTs in GC filopodia; numbers above
bars indicate P values from Chi^2 tests. Throughout the figure, data obtained from live analyses
with Eb1::GFP are framed in red.

1225

Fig. 7. Loss of Efa6 promotes MT entry into shaft filopodia. Quantitative analyses of MT
behaviours in axon shafts, as indicated above each graph; bars are colour-coded: grey,
controls; blue, different *Efa6* mutant alleles; green, neurons over-expressing Efa-FL::GFP; red
outlines indicate live imaging data, all others were obtained from fixed specimens. A-C,G,H)
data all represent fold-change relative to wild-type controls (indicated as horizontal dashed
"ctrl" line) and are shown as single data points and a bar indicating mean ± SEM; P values
from Mann-Whitney tests are given above each column, sample numbers at the bottom of
each bar. D-F) Percentages of antero- / retrograde comet movements (D), percentage of
neurons harbouring at least one filopodium containing / lacking Eb1 comets (E) or containing /
lacking tubulin-stained MTs (F); fractions indicated by light / dark shades as indicated; G)
number of tubulin-stained MTs in shaft filopodia relative to axon length normalised to wildtype
controls; H) number of axon shaft filopodia containing MTs (values relative to ctrl); numbers
on top or at the upper limit of the bars indicate P values from Chi^2 tests.

1239

Fig. 8. Efa6 domain and motif requirements for MT elimination in neurons and fibroblasts. A)
Schematics of *Drosophila melanogaster (Dm)* Efa6 (isoform C, CG31158), *Caenorhabditis*
elegans (Ce) Efa6 (isoform Y55D9A.1a) and *Homo sapiens* (human) PSD(1) (isoform
201/202, NP_002770.3), illustrating the positions (numbers indicate first and last residues) of
the putative PSD95-Dlg1-ZO1 domain (PDZ), SxIP/SxLP motifs (SRIP, SQIP, SALP, SSLP),
the MT-binding domain (MTED), SEC7 domain, plekstrin homology (PH) domain and coiled-

coil domain (CC). B) Schematics on the left follow the same colour code and show the *D.m.*
Efa6 constructs used in this study (dashed red lines indicate the last/first residue
before/behind the truncation). Bar graphs on the right show the impact that transfection of
these constructs had on axon loss in primary *Drosophila* neurons (dark grey in left graph) and
on MT loss in fibroblasts (dark grey or black as indicated); for respective images see F and G
and Fig.S9) Analogous fibroblast experiments as performed with *Drosophila* constructs were
performed with full length constructs of *C. elegans* Efa6 and human PSDs (C), with N-terminal
constructs (D) or synthetic MTEDs (E) of *Dm* and *Ce*Efa6 and of human PSD(1). Throughout
this figure, construct names are highlighted in red for *Drosophila*, light blue for *C. elegans* and
yellow for *Homo sapiens*; all graph bars indicate percentages of neurons with/without axons
(light/dark grey) and of fibroblasts with normal, reduced or absent MTs (light, medium, dark
grey, respectively); numbers in the left end of each bar indicate sample numbers, on the right
end the P values from Chi^2 tests relative to GFP controls; numbers on the right of bars in B
compare some constructs to Efa6-FL::GFP, as indicated by black lines. Primary neurons
expressing Efa6-FL::GFP transgenically and stained for tubulin (asterisks, cell bodies; white
arrows, axon tips; open arrow, absent axon). G-G") Fibroblasts expressing Efa6-FL::GFP and
stained for tubulin; curved arrows indicate areas where MTs are retracted from the cell
periphery; grey dots in F-G" indicate the phenotypic categories for each neuron and
fibroblasts, as used for quantitative analyses in the graphs above. Scale bar at bottom of F
refers to 10µm in F and 25µm in G.

1266

Fig. 9. A model of the suggested roles of Efa6. **A**) The model of local axon homeostasis
(Prokop, 2016; Voelzmann et al., 2016a) states that axonal MTs (green bars) are actively kept
in bundles; for example, their continued polymerisation (1) mediated by plus end machinery
(blue circle) is guided by spectraplakins (here Shot) along cortical actin into parallel bundles
(2), or MTs are kept together through cross-linkage (brown "L"; 4) (Bettencourt da Cruz et al.,
2005; Krieg et al., 2017); here we propose that MTs accidentally leaving the bundle become
vulnerable (orange circles) through elimination (red "T") by cortically anchored Efa6. **B**) In
normal neurons, polymerising MTs in bundles (dark blue circles) are protected by Shot
(marine lines) and MT-MT cross-linkers (brown rectangles), whereas MTs approaches the
membrane (orange arrow heads) either by splaying out in GCs or leaving the bundle in the
shaft (orange arrow heads) become vulnerable (orange circles) to elimination by Efa6 (light
green/red dashes) both along the cortex and in filopodia **C**) Upon Efa6 deficiency, MTs
leaving bundles or entering GCs are no longer subjected to Efa6-mediated cortical elimination
(blue arrow heads) and can cause gradual build-up of MT disorganisation; when entering shaft
filopodia they can promote interstitial branch formation (red arrow), when entering GC
filopodia they can promote axon growth or even branching through GC splitting (yellow
arrows). **D**) Far more MTs leave the bundles in *shot* mutant neurons, but a good fraction of
them is eliminated by Efa6 (increased number of orange arrow heads). **E**) In the absence of
both Shot and Efa6, more MTs leave the bundles, but there is no compensating cortical
elimination (increased number of blue arrow heads), so that the MT disorganisation phenotype
worsens**.**

1288

Supplementary materials

1290

Fig.S1 *ato-Gal4*-driven Efa-FL::GFP expression in adult brains. Brains (region of the optic
lobe in oblique view; white dashed line indicating the lower edge of the medulla) of young (2-5

d after eclosure; top) and old flies (15-18 d; bottom) which are either from wild-type controls or
from flies driving *UAS-myr-tdTomato* and *UAS-Efa6-FL-GFP* via the *ato-Gal4* driver in dorsal
cluster neurons; specimens are stained for GFP and arrows point at stained; for myr-dtTomato
staining of brains from these experiments see Fig. 3. Scale bar in D represents 60μm in all
images.

1298

1299   **Fig. S2**. Efa6 mutant primary neurons show MT disorganisation. Quantitative analyses of MT
1300   disorganisation measured as binary score (percentage of neurons showing obvious MT
1301   disorganisation in their axons) in different *Efa6* mutant alleles (control value indicated as
1302   horizontal dashed "ctrl" line); P values from Chi^2 tests are given above the bars.

1303

Fig. S3 Phylogentic tree analysis of Efa6. Bayesian phylogenetic analysis of full length (A) or
N-termini (B) of Efa6 orthologues. Sequences were aligned using Muscle or ClustalO and
posterior probabilities and branch lengths calculated using MrBayes. Branch length scale is
indicated; blue numbers show posterior probabilities of each branch split. For the full length
tree, *Drosophila steppke* (*step*) was used as outgroup. In both full length and N-terminus
analyses, chordates (cream colour) split off very early from Efa6 versions of other species, in
line with an early speciation event separating both groups before the vertebrate multiplication
events took place. Phyla are highlighted in different colours, gene symbols and/or accession
numbers are given after the species names.

1313

Fig. S4. Efa6 N-terminal domains vary amongst different phyla. A) Domain annotations in 56
Efa6 orthologues via EMBL SMART and Uniprot. Phyla are colour-coded as in Fig.S3. Note
that there is a strong variation of lengths and domain composition in particular of the N-
terminus. The putative PDZ domain is not present in any analysed chordate orthologue, and
MTED and MTED-like sequences cannot be consistently identified in all Efa6 orthologues and
are very divergent in chordate Efa6/Psd versions. Only a subset of nematode (e.g. *C.
elegans*) and insect (e.g. *D. melanogaster*) Efa6 versions, and even fewer chordate Efa6/PSD
proteins, contain SxIP/SxLP sites flanked by positive charges (as would be expected of
functional sites) (Honnappa et al., 2009) in the N-terminal half of the protein; in mouse, alpaca
and cat SxIP/SxLP sites are flanked by negative charges. B) To determine a potential MTED
consensus sequence, 37 sequences of molluscs, nematodes, arthropods and putative MTED
sequences of mammalian PSD1-4 were grouped according to phylum; consensus sequences
were depicted using Berkley's Weblogo online server (default colour scheme). Conserved
amino acid positions are highlighted (faint yellow).
1328

1329   **Fig.S5.** Localisation of Efa6 constructs in primary neurons. Images show transfected primary
1330   *Drosophila* neurons at 18HIV stained for tubulin (magenta) and GFP (green and in greyscale
1331   below the colour image). Cell bodies are indicated by asterisks, axon tips by arrows. The
1332   transfected constructs are indicated top right following the nomenclature explained in Fig.8.
1333   Scale bar in A represents 10μm for all figures shown.

1334

Fig.S6. Efa6 constructs localisations in fibroblasts. Images show fibroblasts which are all
stained for GFP (green) and either for actin or MTs (magenta); GFP and actin/MTs shown as

single channels in greyscale below the colour image, 24hrs after transfection with control
(GFP) or Efa6-derived constructs (nomenclature as explained in Fig.8, but leaving out the
"Efa6"-prefix and "::GFP"-postfix, as indicated top right. Double chevrons indicate nuclear
localisation, arrow heads membrane localisation apparent at cell edges and curved arrows
membrane ruffles. Scale bar in A represents 10µm in all images.

1342

Fig.S7. MT elimination by Efa6-FL is concentration-dependent in fibroblasts. A-C)
Representative images of fibroblasts stained for GFP and tubulin (green and magenta,
respectively; both shown as single channels in greyscale below the colour image). Images
were taken 24hrs after transfection with *Efa6-FL::GFP*, assessed for GFP intensity (plotted on
the ordinate in D; examples for low, moderate and high expression are given in the left, middle
and right columns, respectively) and then grouped with respect to their MT phenotypes into
"MTs intact" (light grey), "mild MT defects" (medium grey) or "MTs gone" (black), as indicated
by greyscale circles in the lowest row of A-C and the abscissa of D. Arrow heads point at GFP
accumulation at membrane edges, white arrows indicate areas of the cell from where MTs
have retracted, open arrows point at retraction fibres and the double chevron indicates the
nucleus position and signs of nuclear GFP localisation. Scale bar in A represents 10µm in all
images.

1355

Fig. S8. Conserved functions of the Efa6 C-terminus in membrane ruffle formation. A-E)
Representative images of fibroblasts stained for GFP and actin (green and magenta,
respectively; both shown as single channels in greyscale below the colour image). Images
were taken 24hrs after transfection with different constructs (indicated top right): control vector
(GFP; A), Efa6-derived constructs (B-D; nomenclature as explained in Fig.8, but leaving out
the "Efa6"-prefix and "::GFP"-postfix) or PSD(1)-FL::GFP (E). To the right of each image, a
selected area is displayed with 2.5-fold magnification, showing dotted actin- and GFP-stained
ruffles in B, D and E, but not A and C; ruffle formation has been quantified and is shown as a
bar graph in F. All graph bars indicate percentages of fibroblasts with/without membrane
ruffles (dark/light grey); P values on top of bars are from Chi^2 tests relative to GFP controls;.
Scale bar in the top image of A represents 10 µM for all fibroblasts shown.

1367

Fig.S9. Representative MT phenotypes induced by the different constructs in transfected
fibroblasts. Fibroblasts 24hrs after transfection with different control (GFP) or Efa6-derived
constructs as indicated top right in each image (nomenclature as explained in Fig.8, but
leaving out the "Efa6"-prefix and "::GFP"-postfix). Cells were stained for tubulin (black; images
shown as inverted greyscale) and classified as "no MT defects" (light grey), "mild MT defects"
(medium grey) or "MTs gone" (black), as indicated by greyscale boxes bottom left of each
image. Each image represents the most prominent phenotype for each respective construct.
Scale bar at the bottom right in H represents 25µm in all images.

1376

Fig. S10. *In vitro* attempts to resolve the MT elimination mechanism of Efa6. A) To determine
whether Efa6 directly affects MT stability, we recombinantly expressed Efa6-ΔCterm::GFP in
Sf9 cells, purified the protein and observed its interaction with MTs using total internal
reflection fluorescence (TIRF) microscopy in a low ionic strength buffer (BRB20, 75mM KCl);
images on the left show three examples of kymographs of MT lattices decorated with Efa6-

∆Cterm::GFP, which displays a mixture of stationary molecules and diffusive interactions

typical of non-translocating MT-associated proteins (Helenius et al., 2006; Hinrichs et al.,

2012); bar charts (right) show quantification of the amount of interacting protein: background

signal from MTs alone, Efa6-∆Cterm::GFP (20nM) and the non-translocating kinesin

MCAK::GFP (20nM) as positive control; at the same protein concentration, over 2-fold more

molecules of MKAC typically interact with MTs; all graphs in this figure show individual data

points and bars representing mean ± SEM; numbers above bars show P values obtained from

Mann–Whitney Rank Sum statistical analyses, numbers in bars the respective sample

numbers. B) Kymographs (left) show individual fluorescently-labelled GMPCPP-stabilised MTs

in vitro (Patel et al., 2016), either alone (MTs only; basal depolymerisation rate, n=7) or in the

presence of 14 nM Efa6-∆Cterm::GFP (n=8) or 40nM MCAK::GFP (n=18); the bar chart (right)

quantifies the induced depolymerisation rates; using two different purifications of Efa6-

∆Cterm::GFP on three separate occasions, we saw no evidence of MT depolymerisation

above the basal level of depolymerisation typically observed in these assays, whereas parallel

control experiments with mitotic centromere-associated kinesin/MCAK showed MT

depolymerisation rates typical of this kinesin. C) To assess whether MT destabilisation might

require additional cytoplasmic factors, we used *Xenopus* oocyte extracts: phase contrast

images show MTs in *Xenopus* oocyte extracts (after they had been allowed to polymerise for

20 min) and then showing stills from before, 10s after and 120s after washing in 20nM

MCAK::GFP (as positive control), 20nM Efa6-∆Cterm::GFP or 40nM synthetic MTED (to

overcome potential mis-folding of the purified ∆Cterm protein); squares outlined by dashed

white lines are shown magnified in insets revealing MTs; MTs clearly vanish upon treatment

with MCAK, but counts of MTs did not reveal any obvious effects on MTs with either of the

Efa6 peptides. D) Finally, interference with MT polymerisation was analysed by adding 20nM

MTED to diluted *Xenopus* cytosol, and 20 random fields were recorded 10 min later via VE-

DIC microscopy and the numbers of MTs per field counted; data plots show that MTED does

not interfere with MT polymerisation. E,F) *RFP* controls and *Efa6-Nterm::GFP* expression

constructs were injected into *Xenopus* embryos at the 4-cell stage and analysed 24 hrs later;

only the *Efa6* construct caused a strong suppression of cell division, as indicated by the

presence of very large cells (arrows) and pigmentation defects (curved arrows) at the site of

injection, suggesting that Efa6-Nterm::GFP is functional when expressed in the *Xenopus*

context. Taken together, the very different data obtained in *Xenopus* cytosol *in vitro* or

embryos *in vivo*, suggest that either the effect of Efa6 requires a factor absent from Xenopus

extract or that Efa6-∆Cterm::GFP and MTED are non-functional due to mis-folding. The latter

is suggested by the observation of many aggregates of Efa6-∆Cterm::GFP in A. Scale bar (in

top left image) represents 3 µm in C (2.5 fold magnified insets), and 1400 µm / 350 µm / 140

µm in left / middle / right images of E and F, respectively.

Fig.1 Qu et al.

A wt 6 HIV

B *Efa6⁻/⁻*

C Efa6-FL

actin
tubulin

D Relative axon length (6 HIV)

0.002 0.024 <0.001 0.01 0.08 <0.001 <0.001

rel. fold increase

wt | Efa6RNAi | Efa6^Def | Efa6^KO#1 | Efa6^GX6[w⁻] | Efa6^GX6[w+] | Efa6-FL | Efa6-FL transfected

240 120 80 100 120 120 106 66

E wt 5 DIV

F *Efa6⁻/⁻*

actin
tubulin

G Efa6-FL

H Efa6-Nterm-CAAX

I Branch number relative to axon length (5 DIV)

0.002 0.08 <0.001

rel. fold increase

ctrl

wt | Efa6^KO#1 | Efa6-FL | Efa6-NtermCAAX

65 78 107 94

Fig. 2 Qu et al.

	control	*Efa6-RNAi*	Efa6-FL::GFP

old (2–5 d) — A (ato / myr-tdTomato), B, C

young (15–18 d) — D, E, F

G. No. of primary branches

y-axis: rel. fold increase (0 to 3)

p-values: <0.001, <0.001, <0.001, <0.001

ctrl (dashed line at 1)

Bar labels: wt 26/254, *Efa6RNAi* 31/284, *Efa6-FL* 39/118, wt 40/363, *Efa6RNAi* 21/194, *Efa6-FL* 44/385

x-axis groups: young, old

H. Normalised no. of medullas with axons

Legend: with axons (grey, light blue, green) / no axons (dark grey, blue, dark green)

y-axis: 0 to 100

p-values: 0.75, <0.001, >0.99, 0.65

Bar labels: **wt** 59, *Efa6RNAi* 37, *Efa6-FL* 67, **wt** 94, *Efa6RNAi* 30, *Efa6-FL* 57

x-axis groups: young, old

Fig. 3 Qu et al.

Fig. 4 Qu et al.

Fig. 5 Qu et al.

Fig. 6 Qu et al.

Fig. 7 Qu et al.

Fig. 8 Qu et al.

A

actin

Efa6

Shot

MTs

③

②

①

④

B wt

axon shaft

filopodium

growth cone

C *Efa6*

growth / split

branching

D *shot*

E *Efa6 shot*

MT ⬤ vulnerable + end ⋯ cortical actin ▬ Efa6

linker ⬤ extending + end ▬ actin filament ▬ Shot

Fig. 9 Qu et al.

Fig. S1 Qu et al.

MT disorganisation in *Efa6* mutants (binary data)

Legend:
- no disorganised MTs
- disorganised MTs

Values above bars: 0.019, <0.001, <0.001, <0.001, <0.001

X-axis categories: wild-type (529), *Efa6RNAi* (120), *Efa6^Def^* (80), *Efa6^KO#1^* (101), *Efa6^GX6[w-]^* (120), *Efa6^GX6[w+]^* (120)

ctrl (dashed line)

Fig. S2 Qu et al.

Fig. S3 Qu et al.

Fig. S4 Qu et al.

Fig. S5 Qu et al.

Fig. S6 Qu et al.

Fig. S7 Qu et al.

Fig. S8 Qu et al.

A GFP **B** FL **C** ΔCterm

D Nterm **E** Nterm-CAAX **F** Nterm^ΔSxIP

G Nterm^ΔMTED **H** MTED **I** ΔNterm

☐ no MT defects ☐ mild MT defects ■ MTs gone

Fig. S9 Qu et al.

A *In vitro* MT binding assay (TIRF)

Efa6-ΔCterm-GFP

GFP

5s
4µm

<0.001
<0.001

Mean GFP intensity x10³

MT only — 21
Efa6-ΔCterm — 50
MCAK — 40

B *In vitro* MT depolymerisation assay

6µm

3min

MT only   Efa6-ΔCterm   MCAK

<0.001
>0.99

Depolymerisation rate µm/min

MT only — 7
Efa6-ΔCterm — 8
MCAK — 18

C MT depolymerisation assay on *Xenopus* extract

before wash   10s after wash   120s after wash

MCAK

Efa6-ΔC.

MTBD

D MT assembly

p=0.79
p>0.99

MTs/image

control — 21
MTBD 40nM — 19
MTBD 40µM — 19

E control

normal light

fluorescence

F Efa6-Nterm

normal light

fluorescence

Fig. S10 Qu et al.