Open-Source Food: Nutrition, Toxicology, and Availability of Wild Edible Greens in the East Bay

Philip B. Stark[1][❂][*], Daphne Miller[2,3][¤], Thomas J. Carlson[4][‡], Kristen Rasmussen de Vasquez[5][¶]

1 Department of Statistics, University of California, Berkeley, California, United States of America
2 School of Public Health, University of California, Berkeley, California, United States of America
3 Department of Family and Community Medicine, University of California, San Francisco, California, United States of America
4 Department of Integrative Biology, University of California, Berkeley, California, United States of America
5 Department of Nutrition and Toxicology, University of California, Berkeley, California, United States of America

❂conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, software, supervision, visualization, writing–original draft preparation, writing–editing
¤writing–original draft preparation, writing–editing
‡conceptualization, investigation, methodology, project administration
¶investigation, methodology

* Corresponding author
E-mail stark@stat.berkeley.edu [PBS]

Abstract

Significance. Foraged leafy greens are consumed around the globe, including in urban areas, and may play a larger role when food is scarce or expensive. It is thus important to assess the safety and nutritional value of wild greens foraged in urban environments.
Methods. Field observations, soil tests, and nutritional and toxicology tests on plant tissue were conducted for three sites, each roughly 9 square blocks, in disadvantaged neighborhoods in the East San Francisco Bay Area in 2014–2015. The sites included mixed-use areas and areas with high vehicle traffic.
Results. Edible wild greens were abundant, even during record droughts. Soil at some survey sites had elevated concentrations of lead and cadmium, but tissue tests suggest that rinsed greens of the tested species are safe to eat. Daily consumption of standard servings comprise less than the EPA reference doses of lead, cadmium, and other heavy metals. Pesticides, glyphosate, and PCBs were below detection limits. The nutrient density of 6 abundant species compared favorably to that of the most nutritious domesticated leafy greens.
Conclusions. Wild edible greens harvested in industrial, mixed-use, and high-traffic urban areas in the San Francisco East Bay area are abundant and highly nutritious. Even grown in soils with elevated levels of heavy metals, tested species were safe to eat

after rinsing in tap water. This does not mean that all edible greens growing in contaminated soil are safe to eat—tests on more species, in more locations, and over a broader range of soil chemistry are needed to determine what is generally safe and what is not. But it does suggest that wild greens could contribute to nutrition, food security, and sustainability in urban ecosystems. Current laws, regulations, and public-health guidance that forbid or discourage foraging on public lands, including urban areas, should be revisited.

Introduction

Diets that include a wide array of plant-based foods are associated with lower rates of chronic disease and better health outcomes [1–3]. Edible wild and feral plants—edible weeds—are an abundant but generally overlooked food source with the potential to contribute to dietary diversity and vegetable intake, especially in areas with limited access to fresh produce.

Edible weeds, which require neither cultivation nor intentional watering, are generally abundant in farms, gardens, parks, yards, sidewalks, and medians, on both private and public land: anywhere there has not been a concerted effort to eradicate them. Some are native to their habitat, but many are nonnative feral species that were once cultivated deliberately but have since colonized the globe [4]. Due to evolutionary selection, edible weeds thrive in places where humans disrupt the soil and they are more tolerant of environmental extremes than most commercial crops [5–7]; climate change may select for weeds and for herbicide-resistant weeds [8].

Edible weeds may be a source of interesting culinary ingredients in the global north [9] and also an important source of nutrition during food shortages [10, 11]. Studies suggest that edible weeds contribute to household food supplies in cities around the globe [12–16]. Many of these plants are also recognized as having medicinal value as teas, supplements, and poultices [17, 18]. While there is little data on foraging behavior in the United States, recent surveys suggest that a noticeable percentage of urban dwellers—representing a diverse group of ethnicities, cultures, and incomes—forage and prepare edible weeds [15, 19–21]. For instance, a 2014–2015 survey of 105 foragers in Baltimore found that they collect 170 unique taxa of plants and fungi from the city and surrounding areas, with low and high income foragers collecting a greater variety of taxa and volume of plants than middle income foragers [21]. Survey respondents cited recreation, economic and health benefits, and connection to nature as their three main motivations for foraging.

Despite the growing recognition that foraged foods are a component of urban food systems and urban ecosystems, surprisingly little is known about their safety, nutritional value, or availability. Extant studies have drawn different conclusions about the safety of food—whether cultivated or "volunteers"—growing in urban environments. For instance, researchers who sampled popular edible weeds growing near urban roads and in the surrounding countryside of Bari, Italy, found that many of the collected species had high concentrations of essential vitamins and minerals, and only two species, *Amaranthus retroflexus* and *Plantago lagopus* had levels of Cd and Pb higher than the legal limit—even when growing in putatively "clean" rural areas [22]. See also [23].

There are a number of studies of the nutritional content of wild and feral foods (e.g., [24–33]). Our study contributes to understanding the potential health and nutritional impacts of urban foraging, by measuring nutrients and unhealthful contaminants (heavy metals, pesticides, herbicides, and PCBs) in six of the most abundant edible weed species harvested in highly-trafficked urban areas in the East Bay region of Northern California.

Methods ₄₄

Berkeley Open Source Food (`https://forage.berkeley.edu`), a research project at   ₄₅
the University of California, Berkeley, mapped wild and feral edible plants, primarily   ₄₆
leafy greens, in three residential areas bordering busy roadways and industrial zones in   ₄₇
Berkeley, Richmond, and Oakland, California, during 2014 and 2015. Each area   ₄₈
comprised approximately 9 square blocks; Table 1 describes them. According to the   ₄₉
USDA, the areas in Richmond and Oakland are more than a mile from any shop that   ₅₀
sells fresh produce; and the area in Berkeley is more than half a mile from such a shop.   ₅₁
All have below average income, according to the U.S. Census.   ₅₂

city	bounding streets
Berkeley	Martin Luther King Jr. Way, California St., Dwight Way, Carleton St.
Oakland	14th St., 17th St., Peralta St., Wood St.
Richmond	Carlson Blvd., Potrero Ave., S. 47th St., S. 43rd St.

Table 1. Boundaries of the study sites in Berkeley, Oakland, and Richmond, CA.

Teams of observers used the iNaturalist smartphone app with customized database   ₅₃
fields to record estimates of the number of 1/2c servings of a variety of edible weeds at   ₅₄
individual street addresses. The website for iNaturalist, an open-source citizen science   ₅₅
database of observations of plants and animals, is `https://inaturalist.org`. The   ₅₆
iNaturalist project page for the study reported here is `https://www.inaturalist.or`   ₅₇
`g/projects/berkeley-open-source-food` (last visited 19 May 2018). As of 19 May   ₅₈
2018, Berkeley Open Source Food had accumulated 631 observations of 52 species,   ₅₉
made by 13 people.   ₆₀

Observations were geo-tagged and accompanied by representative photographs to   ₆₁
support each observation—species identification and abundance. Some representative   ₆₂
"voucher" samples were submitted to the University and Jepson Herbaria at the   ₆₃
University of California, Berkeley; see table 4.   ₆₄

The number of 1/2c servings was estimated on an approximately logarithmic scale   ₆₅
intended to facilitate accurate, repeatable categorization: <1, 1–5, 6–10, 11–20, 21–50,   ₆₆
51–100, 101–500, 501–1000, and >1000. Observations were generally made by pairs of   ₆₇
observers conferring about each street address. Observers (faculty and students at the   ₆₈
University of California, Berkeley) estimated the number of "visible" and the number   ₆₉
of "available" servings. "Visible" servings are those available to a person with legal   ₇₀
access to the property. "Available" servings are those within an arm's reach of a public   ₇₁
right-of-way, such as a sidewalk or road. While we do not know the intrinsic accuracy   ₇₂
of the estimated number of servings, the measurements were reproducible: estimates   ₇₃
made by different observers were calibrated against each other on sets of ten street   ₇₄
addresses. After new observers received approximately an hour of training, inter-rater   ₇₅
concordance of the estimates across pairs of observers was essentially perfect.   ₇₆

Soil samples were taken at 28 sites in Richmond, CA, on 18 June 2014, and in West   ₇₇
Oakland, CA, on 28 August 2014. Samples were sent to the Soil and Plant Tissue   ₇₈
Testing Lab at the University of Massachusetts, Amherst, for metal assays. The   ₇₉
concentrations of Zinc (Zn), Copper (Cu), Arsenic (As), Selenium (Se), Lead (Pb),   ₈₀
Nickel (Ni), Chromium (Cr), Cadmium (Cd), and Molybdenum (Mb) were measured.   ₈₁
Soil samples were taken "per street address," homogenizing sub-samples taken at 3–4   ₈₂
locations near the road at each address tested.   ₈₃

Tissue samples, collected 12 May 2015, targeted locations where soil testing had   ₈₄
shown the concentration of metals to be highest (sample sites 28 and 29, Willow St.   ₈₅
near 16th and 17th Streets, Oakland, CA), and included samples of plants growing   ₈₆
through asphalt. Plant tissue samples were rinsed in tap water as if to make salad,   ₈₇
then dried at the University and Jepson Herbaria at the University of California,   ₈₈

Berkeley and sent to Brookside Laboratories (New Bremen, Ohio) to be assayed for   89
metals (As, Cd, Cr, Cu, Pb, Hg, Mo, Ni, Se, Zn). Voucher specimens for each tested   90
species were submitted (references Thomas Carlson 5001, 5002a/b, and 5003–5009).   91

Fresh/wet plant tissue samples collected on 21 March 2016 were rinsed in tap   92
water, then assayed by SCC Global Services (Emeryville, CA) for vitamins, minerals,   93
polyphenols, and some organic chemical contaminants, including PCBs, glyphosate,   94
and multi-residue pesticides (via QuEChERS). Oxalis (*Oxalis pes-caprae*) and dock   95
(*Rumex crispus*), two species high in oxalic acid, were assayed for oxalic acid.   96

Results   97

Soil sample locations are listed in Table 2. Soil metals test results are listed in Table 3.   98
Locations and species for dried tissue tests are listed in Table 4. Metals tests on dried   99
tissue are given in Table 5. Wet tissue sample species and locations are listed in   100
Table 6. Results of nutritional tests on wet tissue are reported per serving in Table 7   101
and per 100g in Table 8. The tables include values for kale, generally regarded as one   102
of the most nutritious cultivated leafy greens. (Values for kale are from the USDA   103
Food Composition Database, `httpa://ndb.nal.usda.gov` retrieved 15 July 2018.)   104

Metals in soils   105

Table 3 shows that a number of samples had levels of Pb approaching (in two   106
locations) or exceeding (in one location) EPA limits, and levels of Cd greater than half   107
the EPA limit in two locations. The levels of other toxic metals were generally far   108
below EPA limits.   109

Metals in plant tissue   110

The only preparation of the tissue samples was to rinse them in tapwater to remove   111
soil and dust, then to air dry them.   112

Table 5 shows that despite the elevated levels of Pb and Cd, the measured amount   113
of toxic metals in the samples was far below the US EPA maximum acceptable daily   114
dose (RfD) for children and adults, with the exception of Cd in one species. The US   115
EPA RfD for dietary Cd is 0.001mg/kg/d (the units are milligrams of cadmium per   116
kilogram of body mass per day; see `https://www.epa.gov/sites/production/files`   117
`/2016-09/documents/cadmium-compounds.pdf`, last accessed 3 June 2018). Thus, a   118
55kg (121lbs) adult would be expected to tolerate 0.055mg/d of Cd in food "likely [...]   119
without appreciable risk of deleterious noncancer effects during a lifetime." The   120
species with the highest concentration of Cd was *Lactuca ludoviciana*, a wild lettuce,   121
which had 0.678ppm Cd in dry tissue (see Table 5). While it is hard to imagine   122
someone eating 100g/d of *L. ludoviciana* (because it is intensely bitter), that would   123
translate to 0.0678mg/d, over the daily limit for a 55kg person, and equal to the limit   124
for a 68kg (150lbs) person.   125

Pesticides and other toxins in fresh (wet) plant tissue   126

All wet tissue samples were rinsed it in tapwater to remove soil and dust, as if to make   127
salad, but were not cleaned in any other way. The QuEChERS multi-pesticide residue   128
test, which covers approximately 330 pesticides and herbicides, did not detect any   129
pesticide residues, nor did specific tests for glyphosate and for PCBs.   130

Nutrition of wet plant tissue

Tables 7 and 8 show that the wild greens are highly nutritious, with particularly high
levels of Fe, Ca, K, protein, and dietary fiber.

S. 45th St., Richmond, CA	
1	807 S. 45th St.
2	819 S. 45th St.
3	825 S. 45th St.
4	831 S. 45th St.
5	845 S. 45th St.
6	800 S. 45th St.
7	814 S. 45th St.
8	822 S. 45th St.
9	826 S. 45th St.
10	832 S. 45th St.
Carlson Blvd, Richmond, CA	
11	946 Carlson Blvd
12	942 Carlson Blvd
13	934 Carlson Blvd
14	928 Carlson Blvd
15	922 Carlson Blvd
16	916 Carlson Blvd
17	908 Carlson Blvd
18	900 Carlson Blvd
West Oakland, CA	
20	1724 14th St.
21	1738 14th St.
22	1740 14th St.
23	15th St. & Wood
24	1726 15th St.
25	1725 15th St.
26	1719 15th St.
27	1715 15th St.
28	17th St. & Wood
29	16th St. & Willow

Table 2. Sites where soil samples were taken. 3–4 samples were taken along the front of each street address and homogenized.

Conclusions

A healthy and sustainable food system requires a year-round, adequately abundant
supply of nutrient-dense, safe, affordable food produced without depleting or
contaminating vital natural resources: water, air and soil. While weeds are generally
unwanted and unwelcome, our research suggests that they could be a helpful
component of sustainable food systems since all the plants we collected had a higher
concentration of most nutrients than domesticated leafy greens and—after rinsing in
water—none had detectable levels of pesticides or PCBs, and their level of heavy
metals per serving were below EPA reference doses, even though they were harvested
from high-traffic and mixed-use areas. Our conclusions about safety are limited to the
species and locations tested, but they are promising.

Element	USEPA limit (mg/kg)	1–10	11–18	20–22	23–26	27	28	29
Zn	23600	187.2	243.3	261.8	212.2	349.1	2887.2	453.1
Cu	N/A	41.4	38.6	40.8	25.6	37.8	66.8	63.8
As	25	N/A	N/A	3.4	1.7	2.8	5.1	4.1
Se	20	N/A	N/A	2.4	2.1	2.7	3.4	3.7
Pb	400	199.8	359.7	196.5	120.1	150.0	354.6	700.9
Ni	1600	32.4	30.5	32.7	23.3	32.3	73.7	40.9
Cr	230	43.7	35.5	51.3	39.1	54.9	56.7	83.6
Cd	70	1.2	0.7	25.7	21.3	30.9	58.8	41.9
Mo	N/A	N/A	N/A	0.8	0.4	1.0	0.9	0.7

Table 3. Soil metal test results. See table 2 for the locations sampled. Soil tests were performed by the Soil and Plant Nutrient Testing Laboratory at the University of Massachusetts, Amherst, Center for Agriculture, Food, and the Environment

sample	species	location	soil sample	notes
5001	*Malva sylvestris* mallow	Willow & 16th	28	
5002a	*Helminthotheca echioides* bristly ox tongue	Willow & 16th	28	
5002b	*Helminthotheca echioides* bristly ox tongue (different population)	Willow & 16th		in asphalt
5003	*Hypochaeris radicata* cat's ear	Willow between 16th & 17th	28	
5004	*Plantago lanceolata* English plantain	Willow & 17th	29	
5005	*Lactuca ludoviciana* wild lettuce	Willow & 17th	29	
5006	*Trophaleum majus* nasturtium	Willow & 15th		in asphalt
5007	*Taraxacum officinale* dandelion	Willow & 15th		in asphalt
5009	*Foeniculum vulgare* sweet fennel	Wood & 15th		in asphalt

Table 4. Species on which dried tissue tests were performed, and sites where samples were taken. All samples were collected on 12 May 2015 in Oakland, CA. Tissue samples were rinsed with tap water, then dried in botanical voucher sample driers at the University and Jepson Herbaria, University of California, Berkeley, before being sent for testing at Brookside Laboratories. Sample numbers refer to voucher specimens in the University and Jepson Herbaria, submitted by Prof. T. Carlson; the full reference for the first sample is, e.g., "Thomas Carlson 5001."

Element	5001	5002a	5002b	5003	5004-1	5004-2	5005	5006	5007	5009-1	5009-2
As	1.71	1.69	1.66	1.68	1.52	1.61	1.66	1.62	1.69	1.59	1.57
Cd	0.709	0.859	<0.383	<0.391	0.506	0.500	5.38	<0.383	<0.400	<0.392	<0.399
Cr	<0.787	<0.767	<0.767	<0.781	<0.767	<0.769	<0.781	<0.766	<0.797	<0.784	<0.799
Cu	13.9	13.6	7.97	7.87	8.87	9.04	17.5	4.79	8.51	5.27	5.07
Pb	<3.94	<3.83	<3.83	<3.91	<3.83	<3.85	<3.91	<3.83	<3.99	<3.92	<3.99
Hg	<0.0393	<0.0383	<0.0383	<0.0390	<0.0383	<0.0384	<0.0390	<0.0382	<0.0398	<0.0391	<0.0399
Mo	<3.94	<3.83	<3.83	<3.91	<3.83	<3.85	<3.91	<3.83	<3.99	<3.92	<3.99
Ni	<0.787	<0.767	<0.767	<0.781	<0.767	<0.769	<0.781	<0.766	<0.797	<0.784	<0.799
Se	<2.36	<2.30	<2.30	<2.34	<2.30	<2.31	<2.34	<2.30	<2.39	<2.35	<2.40
Zn	161.0	69.6	71.8	110.6	183.1	183.5	398.2	115.2	70.9	30.7	30.5
DWT%	24.3	13.9	12.8	12.6	19.6	19.6	12.6	15.1	12.7	19.3	19.3

Table 5. Metals tests of dry plant tissue. Results are in mg/kg (i.e., parts per million by mass). Tests were performed by Brookside Laboratories, Inc., New Bremen, OH. The concentration of lead (Pb) was below quantification limits in all samples. The repeated measurements differ by far more than the reported precision of the measurements, but with the exception of Cu, the measurements seem to be repeatable to two or three digits. Samples 5004-1 and 5004-2 are repeated tests on sample 5004; 5009-01 and 5009-02 were repeated tests on sample 5009. The bottom row, DWT%, gives the dry weight percentage for each sample. For instance, sample 5005 (a wild lettuce) weighed 36g before drying and 4.537g dry; the dry weight percentage is $100 \times 4.537/36 = 12.6\%$. The original concentration of Cd in the wet tissue of 5005 is thus estimated to be $5.383 \times 0.126 = 0.678$ppm. See table 4 for more information about the samples.

species	sample location
chickweed	Willow & 14th St.
Stellaria media	Campbell & 14th–15th St.
dandelion	Willow & 14th St.
Taraxacum officinale	
dock	Willow & 14th St.
Rumex crispus	
mallow	Willow & Wood, 14th–17th St.
Malva sylvestris	
nasturtium	Campbell & 14th–15th St.
Tropaeolum majus	
oxalis, sourgrass	Willow & 15th–17th St.
Oxalis pes caprae	

Table 6. Latin names and common names of species on which wet tissue tests were performed, and locations where samples were collected. Samples were collected in West Oakland, CA, on 21 March 2016. Samples were rinsed in tapwater before testing. Tests were performed by SCS Global Services in Emeryville, CA. Samples were assayed using the QuEChERS multi-pesticide residue test, which covers approximately 330 pesticides and herbicides; for glyphosate specifically; and for PCBs. No pesticides, glyphosate, or PCBs were detected in any of the samples. Nutritional test results for the samples is given in tables 7 and 8.

	chickweed *Stellaria media*	dandelion *Taraxacum officinale*	dock *Rumex crispus*	mallow *Malva sylvestris*	nasturtium *Tropaeolum majus*	oxalis *Oxalis pes-caprae*	kale *Brassica oleracea*
serving (g)	101	70	98	68	72	84	21
cal (Kcal)	29.51	24.40	32.85	35.36	33.78	23.21	7.0
fat cal (Kcal)	2.43	2.43	2.43	2.43	4.60	2.13	2.79
fat (g)	0.27	0.27	0.27	0.27	0.51	0.24	0.31
saturated fat (g)	0.01	0.01	0.02	0.01	0.03	0.01	0.04
TFA (g)	0	0	0	0	0	0	0
cholesterol (mg)	0	0	0	0	0	0	0
carbohydrates (g)	5.27	3.88	4.72	5.30	4.97	4.45	0.93*
dietary fiber (g)	3.69	3.68	3.34	4.88	2.23	2.52	0.90
total sugars (g)	0	0	0	0	0.27	0	0.21
protein (g)	1.45	1.59	2.59	2.78	2.33	0.83	0.61
Vitamin A (IU)	2315.32	4603.9	5311.82	3168.76	5891.04	1998.25	1011.0
Vitamin C (mg)	10.82	3.14	35.63	5.87	1.07	7.93	19.6
Na (mg)	45.83	36.64	99.46	29.07	28.78	24.33	11.0
Ca (mg)	66.92	67.13	67.40	185.39	106.89	41.07	53.0
Fe (mg)	1.56	1.91	1.29	2.27	0.85	1.58	0.34
K (mg)	446.24	308.06	305.40	242.14	214.54	108.21	73.0

Table 7. Nutritional tests of wet plant tissue (performed by SCS Global Services in Emeryville, CA) collected by Berkeley Open Source Food in West Oakland, CA, and USDA National Nutrient Database values for raw kale. Serving sizes for chickweed, dandelion, dock, and kale were 1c; serving sizes for mallow, nasturtium, and oxalis were 1/2c. Masses are listed. "cal" and "Kcal" stand for kilocalories (dietary calories) and "TFA" stands for trans fatty acids. See table 6 for sample sites. (*This number is suspiciously low—and values listed on other websites are generally 4–6g—but it is the value the USDA lists.)

	chickweed *Stellaria media*	dandelion *Taraxacum officinale*	dock *Rumex crispus*	mallow *Malva sylvestris*	nasturtium *Tropaeolum majus*	oxalis *Oxalis pes-caprae*	kale *Brassica oleracea*
cal (Kcal)	29.09	34.86	33.37	52.14	46.91	27.52	35.0
fat cal (Kcal)	2.40	3.47	2.47	3.58	6.39	2.52	13.41
fat (g)	0.27	0.39	0.27	0.40	0.71	0.28	1.49
saturated fat (g)	0.01	0.01	0.02	0.01	0.04	0.01	0.18
TFA (g)	0	0	0	0	0	0	0
cholesterol (mg)	0	0	0	0	0	0	0
carbohydrates (g)	5.19	5.55	4.79	7.81	6.90	5.27	4.42
dietary fiber (g)	3.64	5.26	3.39	7.20	3.10	2.99	4.10
total sugars (g)	0	0	0	0	0.37	0	0.99
protein (g)	1.43	2.27	2.63	4.10	3.23	0.98	2.92
Vitamin A (IU)	2282	6577	5396	4637	8182	2369	4812
Vitamin C (mg)	10.66	4.49	36.19	8.65	1.49	9.40	93.40
Na (mg)	45.17	52.34	101.04	42.87	39.97	28.85	53.0
Ca (mg)	65.96	95.90	68.47	273.39	148.46	48.69	254.0
Fe (mg)	1.54	2.73	1.31	3.35	1.18	1.87	1.60
K (mg)	439.82	440.08	310.24	357.09	297.97	128.29	348.0
total phenolics (mg/g)	0.77	0.49	2.77	1.29	2.82	1.68	NA
oxalic acid–soluble (mg/g)			0.18		10.94		
oxalic acid–total (mg/g)			0.39		15.42		

Table 8. Nutritional tests of wet plant tissue (performed by SCS Global Services in Emeryville, CA) collected by Berkeley Open Source Food in West Oakland, CA, and USDA National Nutrient Database values for raw kale. Results are per 100g of wet tissue except total phenolics and oxalic acid, which are concentrations (mg/g) See table 6 for sample locations.

Discussion

The nutrient density of the wild greens is high, and compares very favorably to commercially farmed produce. For instance, a cup of dock (*Rumex crispus*) or a half cup of Nasturtium (*Tropaeolum majus*) contains more than the adult RDA of Vitamin A (5000 IU). The amount of Ca per cup of mallow (*Malva sylvestris*) is almost 27% higher than the same volume of whole milk (370mg versus 276mg) and contains 72% of the protein (5.56g versus 7.69g). (Values for milk are from the USDA National Nutrient Database, https://ndb.nal.usda.gov/ndb/foods/show/01211, last visited 21 October 2018.) Per serving and per 100g, the wild greens are generally more nutritious than kale, although kale does stand out in its value of Vitamin C per 100g. While kale has higher concentration of Vitamin C than the species we tested, we did not test any wild greens in the same family as kale (*Brassicacaea*), such as wild mustard (*Hirschfeldia incana, Brassica nigra, Sinapis arvensis*, et al.) or wild radish (*Raphanus raphanistrum*), which are also abundant in the study areas. We suspect they would have Vitamin C levels closer to that of kale.

Laboratory tests for substances toxic to humans focused on metals and chemicals expected in the study zone; there is a possibility that other contaminants were missed by the testing, such as pathogenic microbes. Other studies, e.g., [34], suggest that appropriate washing suffices to mitigate the risk of pathogenic microbes such as *E. coli* O157:H7. We did not measure the concentration of some naturally occurring minerals and chemicals in plants (e.g., phytates) that in high concentration could have negative health consequences for some humans. We did measure oxalic acid in two species known to have high concentrations.

We did not measure soil pH. Plant uptake of toxic metals varies by species and is influenced by pH, salinity, and other soil properties; there is typically a "plateau"

effect that limits the ability of plants to accumulate metals as the concentration of ₁₇₀
metals in soil increases [35]. However, there is evidence that the phytoaccumulation of ₁₇₁
toxic metals in leafy greens is relatively unaffected by pH [36]. ₁₇₂

Measuring the year-round availability and abundance of these plants was beyond ₁₇₃
the scope of this study. However, some of the neighborhood mappings occurred in ₁₇₄
summer (August) of 2014, considered the worst drought year in California in ₁₇₅
1200 years (see `https://en.wikipedia.org/wiki/2011%E2%80%9317_California_dr` ₁₇₆
`ought#2014` Last visited 20 May 2018), a time expected to have little food, absent ₁₇₇
deliberate irrigation. Even during this low-production period, almost every street ₁₇₈
address in all three study areas had several servings of several different species, ₁₇₉
suggesting that wild edible greens are a reliable source of nutrition year-round. ₁₈₀

According to the most recent data from the USDA Economic Research Service ₁₈₁
(2015), waste-adjusted availability of vegetables in the U.S is approximately 1.72 cups ₁₈₂
per capita per day [37], somewhat less than the recommended intake of 2-3 cups ₁₈₃
daily [38]. Waste-adjusted availability is is the sum of domestically produced ₁₈₄
vegetables and imports, less the waste that occurs throughout the food chain. Our ₁₈₅
observations suggest that wild greens can potentially contribute to nutrient security by ₁₈₆
filling in the gap between recommended and available daily servings of vegetables. ₁₈₇

Many of these species volunteer on farms and in gardens, where such "accidental ₁₈₈
crops" may provide additional nutrition and income. A 2014 survey of 21 farms and ₁₈₉
gardens in the East Bay [39] found that of the 15 most most frequently reported "pest" ₁₉₀
plants, 11 are edible, and 9 are "culinary quality," namely, Plantago, oxalis, mallow, ₁₉₁
bristly ox tongue, dandelion, blackberry, calendula, purslane, and hairy bittercress. ₁₉₂
Wild foods might also contribute to a healthy ecosystem by building soil organic ₁₉₃
matter, retaining water and nutrients in the soil, and reducing erosion. Wild plants ₁₉₄
may enhance biodiversity by serving as a habitat for insects and animals and other ₁₉₅
plants. ₁₉₆

Recognizing urban foraging as a legitimate source of nutrition to promote a varied ₁₉₇
diet raises ethical, legal, and policy issues beyond the scope of this paper. [40] make a ₁₉₈
case for permitting foraging in municipal parks and public schools. Foraging is ₁₉₉
currently prohibited on most public lands in the US. For instance, the City of Berkeley ₂₀₀
Municipal Code section 12.44.020 states: ₂₀₁

> Cutting, trimming or removal–Permit and inspection required. It is ₂₀₂
> unlawful for any person to cut, trim, remove, mutilate, injure or in any way ₂₀₃
> impair the growth of any tree, shrub or plant being or growing in or on any ₂₀₄
> street, parking strip, public square, park or playground in the City, or to ₂₀₅
> cause or permit the same to be done. ...(Ord. 3380-NS §2, 1954) ₂₀₆

The East Bay Regional Park District also prohibits taking any plant material ₂₀₇
whatsoever. Despite the fact that the parks use grazing, prescribed burning, and ₂₀₈
mechanical, chemical, and biological means to control some invasive species, including ₂₀₉
a number of edible species. (See, e.g., `https://www.ebparks.org/civicax/filebank` ₂₁₀
`/blobdload.aspx?BlobID=23687`, last accessed 16 July 2018. Listed edible species ₂₁₁
include sweet fennel, artichoke thistle, and Himalayan blackberry, among others.) park ₂₁₂
users are expressly forbidden from collecting any plants, including those the District ₂₁₃
seeks to eradicate. EBRPD Ordinance 38 [41] states: ₂₁₄

> SECTION 804. PLANTS. No person shall damage, injure, collect or ₂₁₅
> remove any plant or tree or portion thereof, whether living or dead, ₂₁₆
> including but not limited to flowers, mushrooms, bushes, vines, grass, turf, ₂₁₇
> cones and dead wood located on District parklands. In addition, any ₂₁₈
> person who willfully or negligently cuts, destroys or mutilates vegetation ₂₁₉
> shall be arrested or issued a citation pursuant to Penal Code Section 384a. ₂₂₀

If foraging were permitted, what rules and norms would be needed to prevent ²²¹ over-foraging and harming existing ecosystems? To what extent can the "culture" of ²²² foragers ensure appropriate ecological stewardship? Because most of these edible ²²³ plants are invasive, it is plausible that harvesting them will improve the overall ²²⁴ ecology; however, such issues need to be addressed. Other countries where foraging is ²²⁵ an established practice, including Scandinavian countries, have national rules and ²²⁶ cultural norms governing foraging. (See, for instance, `https://en.wikipedia.org/wi` ²²⁷ `ki/Freedom_to_roam`, last accessed 16 July 2018.) ²²⁸

References

1. Farhangi MA, Jahangiry L. Dietary diversity score is associated with cardiovascular risk factors and serum adiponectin concentrations in patients with metabolic syndrome. BMC Cardiovascular Disorders. 2018;18(1):68. doi:10.1186/s12872-018-0807-3.

2. Gholizadeh F, Moludi J, Lotfi Yagin N, Alizadeh M, Mostafa Nachvak S, Abdollahzad H, et al. The relation of Dietary diversity score and food insecurity to metabolic syndrome features and glucose level among pre-diabetes subjects. Primary Care Diabetes. 2018; p. 1–7. doi:10.1016/j.pcd.2018.03.003.

3. Vadiveloo M, Dixon LB, Mijanovich T, Elbel B, Parekh N. Dietary Variety Is Inversely Associated with Body Adiposity among US Adults Using a Novel Food Diversity Index 1 – 3. Journal of Nutrition. 2015;145(3):555–564. doi:10.3945/jn.114.199067.weight.

4. Holm LG, Plucknett DL, Pancho JV, Herberger JP. The World's Worst Weeds. University Press; 1977.

5. James SW, Friel S. An integrated approach to identifying and characterising resilient urban food systems to promote population health in a changing climate. Public Health Nutrition. 2015;18(13):2498–2508. doi:10.1017/S1368980015000610.

6. Ramesh K, Matloob A, Aslam F, Florentine SK, Chauhan BS. Weeds in a Changing Climate: Vulnerabilities, Consequences, and Implications for Future Weed Management. Frontiers in Plant Science. 2017;8(February):1–12. doi:10.3389/fpls.2017.00095.

7. Richards CL, Bossdorf O, Muth NZ, Gurevitch J, Pigliucci M. Jack of all trades, master of some? On the role of phenotypic plasticity in plant invasions. Ecology Letters. 2006;9(8):981–993. doi:10.1111/j.1461-0248.2006.00950.x.

8. Matzrafi M, Seiwert B, Reemtsma T, Rubin B, Peleg Z. Climate change increases the risk of herbicide-resistant weeds due to enhanced detoxification. Planta. 2016;244(6):1217–1227. doi:10.1007/s00425-016-2577-4.

9. Rivera L. Beauty and the Feast; 2017. *The Independent*, `https://www.indepen` `dent.co.uk/life-style/food-and-drink/foraging-food-free-picking-of` `f-the-land-nettles-nordic-food-lab-the-ethicurean-a7669916.html`.

10. Vorstenbosch T, de Zwarte I, Duistermaat L, van Andel T. Famine food of vegetal origin consumed in the Netherlands during World War II. Journal of Ethnobiology and Ethnomedicine. 2017;13(1):1–15. doi:10.1186/s13002-017-0190-7.

11. Zhang L, Chai Z, Zhang Y, Geng Y, Wang Y. Ethnobotanical study of traditional edible plants used by the Naxi people during droughts. Journal of Ethnobiology and Ethnomedicine. 2016;12(1):1–16. doi:10.1186/s13002-016-0113-z.

12. Licata M, Tuttolomondo T, Leto C, Virga G, Bonsangue G, Cammalleri I, et al. A survey of wild plant species for food use in Sicily (Italy) - results of a 3-year study in four Regional Parks. Journal of Ethnobiology and Ethnomedicine. 2016;12(1). doi:10.1186/s13002-015-0074-7.

13. Maroyi A. Use of weeds as traditional vegetables in Shurugwi District, Zimbabwe. Journal of Ethnobiology and Ethnomedicine. 2013;9(1):1. doi:10.1186/1746-4269-9-60.

14. Maroyi A. Not just minor wild edible forest products: Consumption of pteridophytes in sub-Saharan Africa. Journal of Ethnobiology and Ethnomedicine. 2014;10(1):1–9. doi:10.1186/1746-4269-10-78.

15. McLain RJ, Hurley PT, Emery MR, Poe MR. Gathering "wild" food in the city: rethinking the role of foraging in urban ecosystem planning and management. Local Environment. 2014;19(2):220–240. doi:10.1080/13549839.2013.841659.

16. Sansanelli S, Tassoni A. Wild food plants traditionally consumed in the area of Bologna (Emilia Romagna region, Italy). Journal of Ethnobiology and Ethnomedicine. 2014;10(1). doi:10.1186/1746-4269-10-69.

17. Ceuterick M, Vandebroek I, Torry B, Pieroni A. Cross-cultural adaptation in urban ethnobotany: The Colombian folk pharmacopoeia in London. Journal of Ethnopharmacology. 2008;120(3):342–359. doi:10.1016/j.jep.2008.09.004.

18. Tareau MA, Palisse M, Odonne G. As vivid as a weed... Medicinal and cosmetic plant uses amongst the urban youth in French Guiana. Journal of Ethnopharmacology. 2017;203(October 2016):200–213. doi:10.1016/j.jep.2017.03.031.

19. Charnley S, McLain RJ, Poe MR. Natural Resource Access Rights and Wrongs: Nontimber Forest Products Gathering in Urban Environments. Society & Natural Resources. 2018;31(January):1–17. doi:10.1080/08941920.2017.1413696.

20. Robbins P, Emery M, Rice JL. Gathering in Thoreau's backyard: nontimber forest product harvesting as practice. Area. 2008;40(2):265– 277. doi:10.1111/j.1475-4762.2008.00794.x.

21. Synk CM, Kim BF, Davis CA, Harding J, Rogers V, Hurley PT, et al. Gathering Baltimore's Bounty. Urban Forestry & Urban Greening. 2017;2.

22. Renna M, Cocozza C, Gonnella M, Abdelrahman H, Santamaria P. Elemental characterization of wild edible plants from countryside and urban areas. Food Chemistry. 2015;177:29–36. doi:10.1016/j.foodchem.2014.12.069.

23. Säumel I, Kotsyuk I, Hölscher M, Lenkereit C, Weber F, Kowarik I. How healthy is urban horticulture in high traffic areas? Trace metal concentrations in vegetable crops from plantings within inner city neighbourhoods in Berlin, Germany. Environmental Pollution. 2012;165:124–132. doi:10.1016/j.envpol.2012.02.019.

24. Bianco VV, Santamaria P, Elia A. Nutritional value and nitrate content in edible wild species used in southern Italy. Acta Horticulturae. 1998;467(September 2016):71–87. doi:10.17660/ActaHortic.1998.467.7.

25. Cowan JW, Sakr AH, Shadarevian SB, Sabry ZI. Composition of edible wild plants of Lebanon. Journal of the Science of Food and Agriculture. 1963;14(7):484–488. doi:10.1002/jsfa.2740140707.

26. Guil JL, Rodríguez-García I, Torija E. Nutritional and toxic factors in selected wild edible plants. Plant Foods for Human Nutrition. 1997;51(2):99–107. doi:10.1023/A:1007988815888.

27. Guil-Guerrero JL, Giménez-Giménez A, Rodríguez-García I, Torija-Isasa ME. Nutritional composition of Sonchus species (S asper L, S oleraceus L and S tenerrimus L). Journal of the Science of Food and Agriculture. 1998;76(4):628–632. doi:10.1002/(SICI)1097-0010(199804)76:4¡628::AID-JSFA997¿3.0.CO;2-U.

28. Mercadante AZ, Rodriguez-Amaya DB. Carotenoid composition and vitamin A value of some native Brazilian green leafy vegetables. International Journal of Food Science & Technology. 1990;25(2):213–219. doi:10.1111/j.1365-2621.1990.tb01077.x.

29. Milovanović M, Picurić-Jovanović K, Djermanović V, Stefanović M. Antioxidant activities of the constituents of Picris echoides. Journal of the Serbian Chemical Society. 2002;67(1):7–15. doi:10.2298/JSC0201007M.

30. Phillips KM, Pehrsson PR, Agnew WW. Nutrient composition of selected traditional United States Northern Plains Native American plant foods. Journal of Food Composition and Analysis. 2014;34:136–152.

31. Sengupta SR, Pal B. Composition of edible wild greens. Journal of the Science of Food and Agriculture. 1970;21(4):215–215. doi:10.1002/jsfa.2740210412.

32. Sridhar R, Lakshminarayana G. Lipid Classes, Fatty Acids, and of Leaves of Six Edible Plant Species'. J Agric Food Chem. 1999;41:61–63. doi:10.1021/jf00025a013.

33. Zennie TM, Ogzewalla D. Ascorbic acid and Vitamin A content of edible wild plants of Ohio and Kentucky. Economic Botany. 1977;31(1):76–79. doi:10.1007/BF02860657.

34. Jensen DA, Friedrich LM, Harris LJ, Danyluk MD, Schaffner DW. Cross contamination of *Escherichia coli* O157:H7 between lettuce and washwater during home-scale washing. Food Microbiology. 2014;46:428–433. doi:10.1016/j.fm.2014.08.025.

35. Dudka S, Miller WP. Accumulation of potentially toxic elements in plants and their transfer to human food chain. Journal of Environmental Science and Health, Part B: Pesticides, Food Contaminants, and Agricultural Wastes. 1999;34(4):681–708. doi:10.1080/03601239909373221.

36. Chang CY, Yu HY, Chen JJ, Li FB, Zhang HH, Liu CP. Accumulation of heavy metals in leaf vegetables from agricultural soils and associated potential health risks in the Pearl River Delta, South China. Environmental Monitoring and Assessment. 2014;186:1547–1560. doi:doi: 10.1007/s10661-013-3472-0.

37. USDA. Total Vegetables: Fresh and Processed; 2018. `https://www.ers.usda.g ov/webdocs/DataFiles/50472/veg.xls?v=42942`.

38. USDA. All About The Vegetable Group; 2018. `https://www.choosemyplate.g ov/vegetables`.

39. Altieri M, Pallud C, Arnold J, Glettner C, Matzen S. An Agroecological Survey of Urban Agriculture Sites in the East Bay, California; 2014. `https://food.be rkeley.edu/wp-content/uploads/2014/02/BFI-2014-An-Agroecological-S urvey-of-Urban-Agriculture-Sites-in-the-East-Bay-.pdf`.

40. Dabady S, Stark PB. Urban Foraging in Municipal Parks and Public Schools: Opportunities for Policymakers; 2017. `https://forage.berkeley.edu/wp-con tent/uploads/2017/08/UrbanForagingPolicyBrief-2017-08-29.pdf`.

41. East Bay Regional Park District. Ordinance 38: Rules and Regulations; 2016. `ht tps://www.ebparks.org/civicax/filebank/blobdload.aspx?BlobID=23167`.