Two-step concentration of complex water samples for the detection of viruses

Kata Farkas [1,*], James E. McDonald [1], Shelagh K. Malham [2] and Davey L. Jones [1,3]

[1] School of Natural Sciences, Bangor University; fkata211@gmail.com; j.mcdonald@bangor.ac.uk; d.jones@bangor.ac.uk

[2] School of Ocean Sciences, Bangor University; s.mahlam@bangor.ac.uk

[3] UWA School of Agriculture and Environment, University of Western Australia, Crawley, Australia; davey.jones@uwa.edu.au

[*] Correspondence: fkata211@gmail.com; Tel.: +44-1248-383056

Abstract: The accurate detection and quantification of pathogenic viruses in water is essential to understand and reduce the risk of human infection. This paper describes a two-step method suitable for the concentration of viruses in water and wastewater samples. The method involves a tangential flow ultrafiltration step that reduces the sample volume of 1-10 l to approx. 50 ml, followed by secondary precipitation using polyethylene glycol 6000 that reduces the volume to 1-4 ml. For method validation, water samples were spiked with different concentrations of enteric viruses and viral recoveries in the concentrates exceeded 10% in all experiments. The method is suitable for water samples with high and low salinity and turbidity, allowing the accurate comparison of viral titers in a diverse range of water types. Furthermore, the method has the potential to concentrate other pathogens, e.g. bacteria or protozoa. Hence, the use of this method can improve the holistic assessment of risks associated with wastewater-contaminated environments.

Keywords: tangential flow ultrafiltration; virus precipitation; qPCR; enteric viruses; norovirus; mengovirus

1. Introduction

Enteric viruses (causing gastroenteritis) and other viral pathogens can be found in wastewater and in wastewater-contaminated surface and groundwater reservoirs. As the infective doses of these agents are low, concentration is needed to accurately quantify viruses in environmental waters and determine public health risks. A great variety of methods are available for water concentration for the recovery of viruses in wastewater and environmental water, however, many of those are not suitable and/or have not been validated for high volumes of water samples or for different water types. The most frequently used methods for the primary concentration of water samples are filtrations using electronegative (EN) or electropositive (EP) filters [1,2]. During EN and EP filtration, the water sample passes through the filter while the virus particles bind to the surface of the filter due to electrostatic forces. These methods have been shown to be suitable for the concentration of viruses in water, however, their use may be limited to low turbidity samples, due to filter clogging during filtration. Furthermore, the use of electronegative filters requires sample preconditioning (i.e. lowing sample pH), whereas electropositive filters may not be suitable for high salinity samples, and the elution of the virus particles from the filters may be difficult as well [1].

Tangential flow ultrafiltration (TFUF) has been used for the concentration of a wide range of water samples for the detection of various pathogens [1–5]. The main advantage of the TFUF approach is that during filtration the water flow takes place in parallel to the membrane, hence membrane clogging is less frequent compared to dead-end ultrafiltration and EN and EP filtration.

In general, TFUF enables 40-200x concentration, and hence secondary concentration (filtration or
precipitation) is often used to further reduce the sample volume [6–8].
This method describes an efficient, accurate and reproducible method for the concentration of
enteric viruses in surface water (fresh and seawater), and wastewater (treated and untreated) samples
(Figure 1). The recommended starting volumes are 10 l for surface water and 1 l for wastewater
samples. The first step of the method is a TFUF step using a 100 kDa cut-off modified
polyethersulfone membrane. As described by others, the efficiency of the elution of the viral particles
from the membrane and the cleaning of the system was enhanced using sodium polyphosphate [4].
The final volume of the sample after TFUF is approx. 50 ml. In order to elute viral particles attached
to solid matter in the primary concentrate, samples are further mixed with beef extract and then
centrifuged. Then polyethylene glycol 6000 (PEG 6000) is added to the supernatant and viral particles
are precipitated. These steps are based on a method for the elution and concentration of viral particles
from sediment [9,10]. The resulting pellet contains the viral particles that can be eluted in phosphate
saline buffer (PBS) and the solution can be stored at -80 °C. The concentrate may be subject to nucleic
acid filtration, viral infectivity or integrity assays.

Figure 1. The stages of the tangential flow ultrafiltration-based two-step water concentration method
for the detection and quantification of enteric viruses in water and wastewater.

2. Experimental Design

2.1. Materials

Sodium-polyphosphate (NaPP) / sodium hexametaphosphate (Sigma Aldrich, USA, Cat. No.:
305553)
Lab Lemco beef extract (Oxoid, UK, Cat. no.: LP0029)
Sodium nitrate (Sigma Aldrich, USA, Cat. no.: S8170)
Polyethylene glycol 6000 (PEG 6000) (Sigma Aldrich, USA, Cat. no.: 81255)
Sodium chloride (Sigma Aldrich, USA, Cat. no.: S7653)
Phosphate saline buffer (PBS), pH 7.4, (Gibco PBS tablets, Life Technologies, Cat. no.: 18912-
014, Life Technologies, USA)
Virkon ® solution (Lanxess, Germany)
20% ethanol (Fisher Chemical, #E/0650/17DF, Thermo Fisher Scientific, USA)
0.5 M HCl and 1 M NaOH for pH adjustment.
Optional: 30 µl mengovirus strain VMC0 (prepared according to ISO/TS150216-1:2013) solution
with approx. 10^6 mengovirus particles

2.2. Equipment

KrosFlo® Research IIi Tangential Flow Filtration System (Spectrum Labs, USA, Cat no. SYR-
U20-01N) or equivalent
100 kDa mPES MiniKros® hollow fibre filter module (Spectrum Labs, USA, Cat. no.: S02-E100-
05-N)

Silicone tubing #17 (Spectrum Labs, USA, Cat. no.: ACTU-E17-25N) or equivalent
Centrifuge (2,500 xg and 10,000 xg at 4 °C)
Pocket sized pH meter (Ishiro, Japan, Cat no.: S2K992 or equivalent)

3. Procedure

3.1 *Tangential flow ultrafiltration (Figure 2; Time of Completion: 2-4 hours)*

3.1.1 System wash
Wash system with 1 l 0.01% NaPP solution (0.1 g NaPP in 1 l deionized water) for 5 min
(permeate closed) then leave the membrane in the solution for 30-60 min. Wash the membrane
with the NaPP solution (permeate open) until the solution has been removed.
3.1.2 Sample filtration
a)   **OPTIONAL STEP** Add approx. 10 μl mengovirus solution to the sample and mix. Save
95   the rest of the mengovirus sample for control measurements.
96   b)   Filter 10 l of surface water or 1 l wastewater at 1-1.6 l/min flow at 5 psi (0.3 bar; 30 kPa)
97   pressure to achieve a permeate flow of 200 – 300 ml/min. Continue filtration until
98   approx. 5 ml sample remains in the reservoir.
99   3.1.3 Backwash, recovery
100   c)   Set the flow to 680 ml/min with no pressure applied and circulate the concentrate for 5
101   min with the permeate clamp closed.
102   d)   Stop pump, close penetrate and retentate valves.
103   e)   Inject 20 ml 0.01% NaPP solution to penetrate pressure valve. Open retentate and wash
104   with reverse flow.
105   f)   Collect the concentrate from the system by introducing air through the retentate port.
106   The final volume of the concentrate is approx. 50 ml.
107   3.1.4 Membrane wash and storage
108   g)   Wash membrane with 250 ml Virkon® solution after each sample by circulating the
109   Virkon® solution in the system (permeate closed) at low flow (400-800 ml/min). In order
110   to reuse membrane, immediately wash it with 150 ml 0.01% NaPP solution using the
111   setup used for the Virkon wash. Repeat until solution in the process reservoir is clear.
112   Leave the membrane in the solution for at least 10 min prior to reuse.
113   h)   For long-term storage, wash the membrane with 50 ml 20% ethanol solution using the
114   setup used for the Virkon wash. Repeat until solution is clear. Dissemble system and
115   store membrane in 20% ethanol solution at 4 °C.

116   3.2 *Secondary concentration (Time of Completion: 2.5 hours + overnight incubation)*

117   3.2.1 Virus elution
118   a)   Add beef extract and $NaNO_3$ to 50 ml concentrated water sample to reach final
119   concentration of 3% w/v and 2 M, respectively. Adjust the pH to 5.5 using 0.5 M HCl.
120   b)   Incubate at 50-90 rpm on ice for 30 min.
121   c)   Centrifuge at 2,500 xg for 10 min, then transfer the supernatant to a new tube. Discard
122   pellet. Adjust the pH of the solution to 7.5 using 1 M NaOH.
123   3.2.2 Virus precipitation
124   d)   Add PEG 6000 and NaCl to reach final concentration of 15% and 2% w/v, respectively.
125   Mix to dissolve PEG 6000 and incubate at 4 °C for 14-18 h. ⏸ **PAUSE STEP** The
126   solution may be stored at 4 °C for up to two days.
127   e)   Centrifuge at 10,000 xg for 30 min at 4 °C. Discard supernatant.
128   Dissolve pellet in 1-4 ml PBS (pH 7.4). The concentrate may be subject to
129   infectivity/integrity assays or nucleic acid extraction flowed by real-time PCR
130   quantification. Alternatively, viral nucleic acids can be extracted directly from the pellet.

131   **OPTIONAL STEP** For estimating method recovery percentile, RNA from 10 µl
132   mengovirus solution should be extracted and quantified.
133

Figure 2. Schematic tangential flow ultrafiltration setup

4. Expected Results

For pilot validation, 2 l of deionized water was spiked with of norovirus GII to reach the final
concentrations of 10^6, 10^5, 10^4, 10^3, 10^2 and 10^1 genome copies (gc)/l in duplicates. Samples were
concentrated using the two-step concentration method. Viral RNA was extracted from the pellet
(final volume: 50 µl) and quantified using qRT-PCR (sample volume: 2 µl) as described in Farkas et
al. [10]. High recoveries were observed in all norovirus concentrations (Table 1). The high deviations
between replicates were result of the limitations of the qPCR method used for quantification. The
limit of quantification (LOQ) was 200 gc/l and the limit of detection (LOD) was approx. 50 gc/l. The
LOD and LOQ can be further lowered by the reduction of the RNA eluent volume and the increase
of the sample volume in the qRT-PCR reaction.

Table 1. Norovirus recoveries and standard deviations (SD) in deionized water using the two-step
concentration method.

Norovirus concentration (gc/l)	Recovery percentile (SD)
5.28×10^6	78 (43)
4.72×10^5	121 (2)
4.31×10^4	99 (11)
3.03×10^3	91 (45)
2.0×10^2	100 (0)

For further validation, 10 l of surface water samples (river, estuarine and sea) in triplicates were
spiked with known concentration of human enteric viruses (norovirus GII, sapovirus GI, hepatitis A
virus and human adenovirus type 40) and with a mengovirus, which is often used as a process control
for the extraction of enteric viruses from environmental matrices. The details of validation are
described elsewhere [11]. Results showed that 10-100% recovery could be achieved in all sample

types. In the same study, the usefulness of the method for wastewater samples was also investigated.
Influent and effluent samples were taken in duplicates at four wastewater treatment plants. As the
samples were expected to contain the target viruses, samples were processed without spiking. High
viral concentrations were observed and the method showed great reproducibility as well. In
subsequent samples spiked with mengovirus, at least 10% recovery was observed. No inhibition or
cross-contamination between samples was observed. The method has also been successfully used for
viral recovery from high volumes (50 l) surface water for metagenomics applications [12].

Applications and recommendations

The method described above is suitable for the concentration of many different water samples
and hence suitable for viral surveillance, contamination source tracking and to describe viral ecology.
The TFUF method described here is suitable for the viral recovery from the mPES MiniKros® hollow
fibre filter (Spectrum Labs, USA), however, alternative recovery buffers may be used with different
membranes. Furthermore, as the membrane used for the TFUF has a 100 kDa pore size, the method
can potentially co-concentrate other microbes and protozoa as well, and hence the TFUF step of the
concentration method enables the accurate description of microbial quality of a sample. In the current
study, viral nucleic acids were directly extracted from the concentrated water samples, however, the
concentrates are suitable for viral infectivity and capsid integrity assays as well [11]. For direct nucleic
acid extraction, the resuspension of the PEG precipitate is not necessary; nucleic acids can be
extracted from the pellet. When PCR-based approaches are used for the quantification of viruses in
the concentrate, the use of robust extraction and amplification methods is recommended, as organic
matter that may interfere with the enzymes used for amplification are co-concentrated with viral
particles. The addition of process control (e.g. mengovirus) to each sample to determine method
efficiency is highly recommended.

Author Contributions: Conceptualization and Methodology, All; Validation, Investigation, K,F.; Writing-
Original Draft Preparation, K.F.; Writing-Review & Editing, All; Funding Acquisition, D.J.

Funding: This work was funded by the Natural Environment Research Council (NERC) and the Food Standards
Agency (FSA) under the Environmental Microbiology and Human Health (EMHH) Programme
(NE/M010996/1).

Acknowledgments: The authors would like to thank Lindsay Smart (Spectrum Labs, UK) for his assistance in
the method development for water concentration.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Cashdollar, J. L.; Wymer, L. Methods for primary concentration of viruses from water samples: a review
and meta-analysis of recent studies. *J. Appl. Microbiol.* **2013**, *115*, 1–11.

2. Ikner, L. A.; Gerba, C. P.; Bright, K. R. Concentration and recovery of viruses from water: a
comprehensive review. *Food Environ. Virol.* **2012**, *4*, 41–67.

3. Soule, H.; Genoulaz, O.; Gratacap-Cavallier, B.; Chevallier, P.; Liu, J.-X.; Seigneurin, J.-M. Ultrafiltration
and reverse transcription-polymerase chain reaction: an efficient process for poliovirus, rotavirus and
hepatitis A virus detection in water. *Water Res.* **2000**, *34*, 1063–1067.

4. Hill, V. R.; Polaczyk, A. L.; Hahn, D.; Narayanan, J.; Cromeans, T. L.; Roberts, J. M.; Amburgey, J. E.
Development of a rapid method for simultaneous recovery of diverse microbes in drinking water by
ultrafiltration with sodium polyphosphate and surfactants. *Appl. Environ. Microbiol.* **2005**, *71*, 6878–6884,
doi:71/11/6878.

5. Holowecky, P. M.; James, R. R.; Lorch, D. P.; Straka, S. E.; Lindquist, H. D. A. Evaluation of ultrafiltration
cartridges for a water sampling apparatus. *J. Appl. Microbiol.* **2009**, *106*, 738–747.

6. Hill, V. R.; Kahler, A. M.; Jothikumar, N.; Johnson, T. B.; Hahn, D.; Cromeans, T. L. Multistate evaluation

of an ultrafiltration-based procedure for simultaneous recovery of enteric microbes in 100-liter tap water
samples. *Appl. Environ. Microbiol.* **2007**, *73*, 4218–4225, doi:AEM.02713-06 [pii].

7.   Rhodes, E. R.; Hamilton, D. W.; See, M. J.; Wymer, L. Evaluation of hollow-fiber ultrafiltration primary
concentration of pathogens and secondary concentration of viruses from water. *J. Virol. Methods* **2011**,
176, 38–45, doi:http://dx.doi.org/10.1016/j.jviromet.2011.05.031.

8.   Kahler, A. M.; Johnson, T. B.; Hahn, D.; Narayanan, J.; Derado, G.; Hill, V. R. Evaluation of an
ultrafiltration-based procedure for simultaneous recovery of diverse microbes in source waters. *Water*
2015, *7*, 1202–1216, doi:10.3390/w7031202.

9.   Lewis, G. D.; Metcalf, T. G. Polyethylene glycol precipitation for recovery of pathogenic viruses,
including hepatitis A virus and human rotavirus, from oyster, water, and sediment samples. *Appl.*
Environ. Microbiol. **1988**, *54*, 1983–1988.

10.   Farkas, K.; Hassard, F.; McDonald, J. E.; Malham, S. K.; Jones, D. L. Evaluation of molecular methods for
the detection and quantification of pathogen-derived nucleic acids in sediment. *Front. Microbiol.* **2017**, *8*,
53, doi:10.3389/fmicb.2017.00053.

11.   Farkas, K.; Cooper, D. M.; McDonald, J. E.; Malham, S. K.; de Rougemont, A.; Jones, D. L. Seasonal and
spatial dynamics of enteric viruses in wastewater and in riverine and estuarine receiving waters. *Sci.*
Total Environ. **2018**, *634*, 1174–1183, doi:10.1016/j.scitotenv.2018.04.038.

12.   Adriaenssens, E.; Farkas, K.; Harrison, C.; Jones, D.; Allison, H. E.; McCarthy, A. J. Viromic analysis of
wastewater input to a river catchment reveals a diverse assemblage of RNA viruses. *mSystems* **2018**, *3*,
e00025-18, doi:10.1128/mSystems.00025-18.