1 **Programmed mutation of liver fluke granulin using CRISPR/Cas9 attenuates**
2 **virulence of infection-induced hepatobiliary morbidity**
3
4 ***Short title: Attenuated virulence in gene edited parasite***
5

6 **Patpicha Arunsan[1,2,¶], Wannaporn Ittiprasert[2,¶], Michael J. Smout[3,¶], Christina J.**
7 **Cochran[2,], Victoria H. Mann[2], Sujittra Chaiyadet[1], Shannon E. Karinshak[2], Banchob**
8 **Sripa[4], Neil D. Young[5], Javier Sotillo[3], Alex Loukas[3,*], Paul J. Brindley[2,*] and Thewarach**
9 **Laha[1,*]**
10
[1]Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
Thailand
[2]Department of Microbiology, Immunology and Tropical Medicine, and Research Center for
Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington
University, Washington, DC 20037, USA
[3]Centre for Biodiscovery and Molecular Development of Therapeutics, Australian Institute of
Tropical Health and Medicine, James Cook University, Cairns, QLD 4878, Australia
[4]Department of Pathology, Faculty of Medicine, Khon Kaen University, Khon Kaen 40002,
Thailand
[5]Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville,
Victoria 3010 Australia
22
23 [¶]These authors contributed equally. [*]These authors contributed equally.
24
25 [*]Correspondence: Thewarach Laha, Department of Parasitology, Faculty of Medicine, Khon
26 Kaen University, 123 Mittraparp Highway, Khon Kaen 40002 Thailand. Phone +66 43 348387.
27 E-mail: thewa_la@kku.ac.th
28
29 [*] Correspondence: Alex Loukas, Centre for Biodiscovery and Molecular Development of
30 Therapeutics, Queensland Tropical Health Alliance, Australian Institute of Tropical Health and
31 Medicine, James Cook University, Cairns Campus, Cairns, Queensland 4878, Australia; E-mail,
32 alex.loukas@jcu.edu.au
33
34 [*]Correspondence: Paul J. Brindley, Department of Microbiology, Immunology and Tropical
35 Medicine, and Research Center for Neglected Diseases of Poverty, School of Medicine & Health
36 Sciences, George Washington University, Washington, DC 20037, USA. E-mail:
37 pbrindley@email.gwu.edu

38
39
40 **KEY WORDS:** *Opisthorchis viverrini*; liver fluke; genome editing; CRISPR/Cas9;
41 opisthorchiasis; *granulin*; growth factor; cholangiocarcinoma
42
43
44
45

Abstract

Infections with several flatworm parasites represent group 1 biological carcinogens, *i.e.* definite causes of cancer. Infection with the food-borne liver fluke *Opisthorchis viverrini* causes cholangiocarcinoma (CCA). Whereas the causative agent for most cancers, including CCA in the West, remains obscure, the principal risk factor for CCA in Thailand is opisthorchiasis. We exploited this established link to explore the role of the secreted parasite growth factor termed liver fluke granulin (*Ov*-GRN-1) in pre-malignant lesions of the biliary tract. We targeted the *Ov-grn-1* gene for programmed knockout and investigated gene-edited parasites *in vitro* and in experimentally infected hamsters. Both adult and juvenile stages of the liver fluke were transfected with a plasmid encoding a guide RNA sequence specific for exon 1 of *Ov-grn-1* and the Cas9 nuclease. Deep sequencing of amplicon libraries from genomic DNA from gene-edited parasites exhibited programmed, Cas9-catalyzed mutations within the *Ov-grn-1* locus, and tandem analyses by RT-PCR and western blot revealed rapid depletion of *Ov-grn-1* transcripts and protein. Newly excysted juvenile flukes that had undergone editing of *Ov-grn-1* colonized the biliary tract, grew and developed over a period of 60 days, were active and motile, and induced a clinically relevant pathophysiological tissue phenotype of attenuated biliary hyperplasia and fibrosis in comparison to infection with wild type flukes. This is the first report of gene knock-out using CRISPR/Cas9 in a parasitic flatworm, demonstrating the activity and utility of the process for functional genomics in these pathogens. The striking clinical phenotype highlights the role in virulence that liver fluke growth factors play in biliary tract morbidity during chronic opisthorchiasis.

Introduction

Liver fluke infection caused by species of *Opisthorchis* and *Clonorchis* remains a major public health problem in East Asia and Eastern Europe. *O. viverrini* is endemic in Thailand and Laos, where ~10 million people are infected with the parasite (*1*). There is no stronger link between a human malignancy and a parasitic infection than that between CCA and infection with *O. viverrini* (*2*). In endemic regions such as Northeastern Thailand, infection causes hepatobiliary diseases including cholangitis and periductal fibrosis - a major risk factor for CCA (*3*). The north of Thailand suffers the highest incidence of CCA in the world, often exceeding 80 cases per 100,000 population, and for which up to 20,000 people annually are admitted for surgery. Unfortunately, prognosis for fluke-induced cancer is poor (*1, 4, 5*).

How and why opisthorchiasis induces cholangiocarcinogenesis is likely multi-factorial, including mechanical irritation of the biliary tract during migration and feeding of the liver fluke, inflammatory molecules released by the parasite, and nitrosamines in fermented foods that are a dietary staple in northern Thailand. To survive in the hostile host environment, parasitic helminths produce an assortment of excretory/secretory (ES) products including proteins with diverse roles at the host–parasite interface. This interaction has long been thought but not fully understood to modify cellular homeostasis and contribute to malignant transformation during chronic opisthorchiasis (*6*). Feeding activity of the liver fluke inflicts wounds in the biliary tree, lesions that undergo protracted cycles of repair and re-injury during chronic infection. The liver fluke secretes mediators that accelerate wound resolution in cholangiocytes, an outcome that can be compromised following silencing of expression of *Ov-grn-1* using RNA interference (*7, 8*). We hypothesize that proliferation of biliary epithelial cells induced by *Ov*-GRN-1 is a pivotal factor in establishing a tumorigenic microenvironment in livers of infected individuals.

Progress with development of genetic tools for functional genomic studies with platyhelminth parasites has been limited to date (*9*). The use of clustered regularly interspaced short palindromic repeats (CRISPR) associated with Cas9, an RNA-guided DNA endonuclease enzyme, has revolutionized genome editing in biomedicine, agriculture and biology at large (*10, 11*). Progress with CRISPR/Cas9 in numerous eukaryotes including the nematodes *Caenorhabditis elegans* and *Strongyloides stercoralis* has been described (*11-14*), but this form of gene editing has not been reported for flatworm parasites. Here, we deployed CRISPR/Cas9 to knockout (mutate) the *Ov-grn-1* gene and assess the virulence of gene-edited flukes *in vitro* and *in vivo* in a hamster model of opisthorchiasis.

Results and Discussion

Growth factor secreted by carcinogenic liver mutated by programmed genome ediitng

A construct termed pCas 9-*Ov-grn-1* (Fig. 1a and Supplementary Fig. 1a) was assembled following assessment of the *Ov-grn-1* locus in the annotated genome sequence of *O. viverrini* (*15*). Nucleotide sequences in pCas 9-*Ov-grn-1* were confirmed by Sanger cycle sequencing; pCas 9-*Ov-grn-1* encodes the Cas9 nuclease and a single guide RNA (sgRNA) complementary to a target sequence 5'-GATTCATCTACAAGTGTTGA within exon 1 of *Ov-grn-1*. The predicted programmed cleavage site was predicted to be at three residues upstream of a CGG proto-spacer

138 adjacent motif sequence (PAM) in exon 1 of *Ov-grn-1* (Fig. 1b; Supplementary Fig. 1b). Adult
139 *O. viverrini* flukes recovered from experimentally infected hamsters were subjected *in vitro* to
140 square wave electroporation (*8*) in the presence of pCas 9-*Ov-grn-1* DNA, and thereafter
141 maintained in culture for three weeks. The activity of CRISPR/Cas9 was evaluated by two
142 approaches.
143
First, quantitative PCR (qPCR) was employed, which relies on the inefficiency of binding of a
primer (here termed OVR-F) overlapping the target genomic sequence of the gRNA, i.e. where
mutations are expected to have occurred, compared to the binding efficiency of flanking primers,
i.e. outside the mutated region (*16, 17*) (flanking primers termed OUT and OUT-R) (Fig. 1b).
Genomic DNA (gDNA) templates were investigated by quantitative PCR to quantify the
efficiency of programmed gene-editing at the target locus; the ratio between the OVR-OUT-R
products and OUT-F-OUT-R products provided an estimate of the amplification fold-reduction
in the sample of CRISPR/Cas9-edited compared to gDNA from control, wild type liver flukes at
the target sequence of the sgRNA, i.e. the annealing site for the OVR primer. A reduction in
relative fold amplification of 2.7% was detected in gDNA from the Cas9-treated worms (Fig. 1e;
Supplementary Fig. 1c).
155
Second, to identify and quantify the mutations that arose in the genome of *Ov-grn-1*-edited
(termed Δ*Ov-grn-1*) flukes, we used an amplicon-sequencing approach. A targeted (amplicon)
sequence library was constructed from genomic DNA from some of the flukes. A 173 bp
fragment spanning the predicted double stranded break site of *Ov-grn-1* was amplified from the
gDNAs primed with oligonucleotides flanking 1496-1668 nt of *Ov-grn-1*. Adaptors and barcodes
were ligated into these amplicon libraries. The MiSeq libraries were undertaken by Illumina
MiSeq-based deep sequencing. Insertion-deletion (INDEL)/mutation profiles in the sequence
reads were compared in multiple sequence alignment with wild type reference template sequence
(1496-1668 nt) of *Ov-grn-1*. Each amplicon was sequenced on the MiSeq Illumina platform and
quantified the gene editing frequency by the CRISPResso pipeline (*18, 19*). More than 2 million
sequenced reads were aligned against the reference sequence, which predicted the presence of
27, 616 non-homologous end joining (NHEJ) reads, specifically 170 reads with insertions
(0.6%), 193 reads with deletions (0.7%) and 27,277 reads with substitutions (98.7%). At large,
1.25% of the sequenced reads exhibited NHEJ mutations (Fig. 1c). Among these NHEJ reads,
there were >100 forms exhibiting mutations that would disrupt the coding sequencing of *Ov-grn-*
1. Four representatives of the INDEL-bearing traces aligned with the WT allele are presented in
Supplementary Fig. 1b. The Illumina sequencing reads are available as GenBank accessions
SRR5764463-5764618, at https://www.ncbi.nlm.nih.gov/Traces/study/?acc=SRP110673,
Bioproject, www.ncbi.nlm.nih.gov/bioproject/PRJNA385864.
175
Effects of gene editing on transcription and protein expression were investigated. Levels of both
Ov-grn-1 mRNA transcripts as determined by RT-PCR and of granulin, as detected by western
blot with anti-*Ov*-GRN-1 sera, fell significantly from days 1 and 2 after transfection, respectively
($P \leq 0.0001$; Fig. 1d, e; Supplementary Fig. 1c). Expression levels of two reference genes, the
actin transcript (Supplementary Fig. 1c) and the *Ov*-TSP-2 tegument protein (Fig. 1d), were not
influenced by the programmed mutation of *Ov-grn-1*.

Mutaion of liver fluke granulin attenuates infection-induced biliary tract morbidity

To investigate whether gene editing of *Ov-grn-1* impacted *in vitro* indicators of pathogenesis, the
capacity of ES products from WT, mock-transfected and gene-edited flukes to drive proliferation
and scratch wound repair of the H69 human primary cholangiocyte cell line was assessed. ES
from WT and mock-transfected adult flukes stimulated cell proliferation and wound closure
whereas an equivalent amount of ES products from Δ*Ov-grn-1* flukes resulted in significantly
reduced cell proliferation over the six day course of the assay ($P \leq 0.0001$; Fig. 2a, b;
Supplementary Fig. 2a, b) and significantly reduced *in vitro* wound closure over 36 hours ($P \leq$
0.0001; Fig. 2c; Supplementary Fig. 2c, d), consistent with a reduction in the amount of *Ov-*
GRN-1 secreted from the gene-edited liver flukes.

Notwithstanding the noteworthy effects observed with gene-edited, adult developmental forms,
the metacercaria (MC) (Fig. 3a) is the infective stage of *O. viverrini* for humans. Accordingly,
we investigated gene knockout in MC. No effect was apparent on *Ov-grn-1* transcript levels in
MC, (Supplementary Fig. 3), suggesting that delivery of the pCas 9-*Ov-grn-1* by electroporation
through the MC cyst wall was ineffective. Exposure to bile acids and gastric enzymes results in
excystation of *O. viverrini* MC in the duodenum of the infected mammalian host. We mimicked
this process *in vitro* using trypsin to release the newly excysted juvenile worms (NEJ; Fig. 3b),
after which NEJ were subjected to electroporation with CRISPR/Cas9 constructs as described for
adult flukes. Marked depletion of *Ov-grn-1* transcripts ($P < 0.0001$) followed this manipulation
(Fig. 3c).
202
In parallel, hamsters were infected, using gastric gavage, with 100 Δ*Ov-grn-1* NEJ or WT NEJ
immediately after electroporation. At necropsy of the hamsters two to three weeks later, it was
clear that Δ*Ov-grn-1* flukes had colonized the bile ducts in similar numbers to WT flukes, and
were similarly motile. Strikingly, the infection with Δ*Ov-grn-1* parasites failed to induce the
marked hyperplasia of the bile duct epithelia characteristic of infection with WT flukes.
Specifically, infection with WT flukes had induced markedly disordered, hyperplasic growth of
the epithelium adjacent to the parasites; at day 14, five times more than in uninfected controls (P
≤ 0.0001) (Fig. 3d) whereas infection with the Δ*Ov-grn-1* flukes (Fig. 3e) provoked 11-fold less
($P \leq 0.0001$) biliary hyperplasia than WT fluke infected livers (day 14, 145% thickening
compared to uninfected controls; $P \leq 0.01$) (Fig. 3g). Indeed, the bile ducts from hamsters
infected with the Δ*Ov-grn-1* flukes generally resembled those of the control, uninfected hamsters
(Fig. 3f).

To assess long-term survival of Δ*Ov-grn-1* NEJ in hamsters and associated chronic biliary
morbidity, hamsters were infected with Δ*Ov-grn-1* and WT NEJ, and adult flukes were
recovered and counted from the livers 60 days post-infection. Similar numbers of worms were
recovered from both control and gene-edited liver fluke-infected hamsters (Fig. 4a). To assess
the impact of infection with Δ*Ov-grn-1* on markers of chronic opisthorchiasis such as biliary
fibrosis, liver sections from infected hamsters were stained with Picro-Sirius Red to detect
collagen bundles in the biliary tract (Fig. 4b). Minimal deposits of collagen were seen in the
periductal regions of the biliary tract of the control, uninfected hamsters whereas hamsters
infected with WT flukes had thick bands of collagen surrounding enlarged bile ducts in the
vicinity of the flukes. Significantly less collagen (28%) was detected in periductal sites of the
biliary tract infected with Δ*Ov-grn-1* flukes ($P < 0.001$) compared to livers of hamsters infected
with WT flukes (Fig. 4b, c). To further assess fibrosis in hepatobiliary tract, we stained for
smooth muscle actin (ACTA2), an established marker of hepatic fibrosis (*20*), by probing thin

tissue sections with anti-ACTA2 antibody. Livers of hamsters infected with WT flukes showed
densely packed collagen fibrils that stained for ACTA-2 in periductal regions proximal to the
parasites. In contrast, livers from hamsters infected with Δ*Ov-grn-1* flukes displayed an irregular
distribution of less dense collagen fibrils with less ACTA2-specific fluorescence (Fig. 4d;
Supplementary Fig. 4). The intensity of ACTA2-specific fluorescence was quantified by
measuring fluorescence intensity; livers from Δ*Ov-grn-1* fluke-infected hamsters showed 94%
reduction (P <0.01) of median fluorescence values relative to controls infected with WT flukes
(Fig. 4e).

Despite reaching high levels of significance, the data range was high for both markers of
collagen deposition assessed. We hypothesized that this was likely due to inconsistencies in the
electroporation-mediated delivery of the CRISPR/Cas9 constructs between individual flukes. We
therefore collected Δ*Ov-grn-1* and WT flukes from hamster bile ducts 60 days post-infection and
assessed *Ov-grn-1* gene expression from individual flukes (Fig. 4f). We then separated flukes
into three groups based on *Ov-grn-1* mRNA expression levels: >100% relative to WT average;
10-100% relative to WT average; <10% relative to WT average. We pooled genomic DNA from
flukes to form the three groups described above and assessed mutation frequencies. In line with
the *Ov-grn-1* mRNA expression profiles, Δ*Ov-grn-1* flukes where gene editing appeared to be
relatively inefficient (i.e. >100% mRNA expression relative to WT average) had an average
mutation frequency of just 0.7%, whereas Δ*Ov-grn-1* flukes where gene editing appeared to be
moderately efficient (10-100% mRNA expression relative to WT average) had an average
mutation frequency of 3.2%; Δ*Ov-grn-1* flukes where gene editing appeared to be highly
efficient (<10% mRNA expression) had an average mutation frequency of 4.6% (Fig. 4f). The
combined mutation frequency was 2.7% (Fig. 4f), notably similar to the 2.7% mutation rate
detected when Δ*Ov-grn-1* flukes were cultured *in vitro* for 21 days (Fig 1e and Supplementary
Fig. 1c).

In overview, we describe the first example of successful somatic gene editing of a parasitic
flatworm using CRISPR/Cas9. The results revealed that gene editing induced disruption of
expression of *Ov-grn-1* in liver flukes, which in turn revealed a pathologically relevant
phenotype in the mammalian biliary tract. Following programmed gene editing, the lesion was
apparently repaired by non-homologous end joining. The bacterial Type II Cas9 system is active
in this liver fluke, and we conjecture that Cas9-catalyzed gene editing will be active in
trematodes and parasitic platyhelminths at large. As noted, whereas the causative agent for many
cancers, including CCA in the West, remains obscure, the principal risk factor for CCA in
Thailand and Laos has long been established - infection with *O. viverrini*. Cas9-based gene
editing and the hamster model of human opisthorchiasis utilized herein, including genetic
manipulation of the infective NEJ parasite, together provide a facile, functional genomics system
to interrogate parasite pathogenicity and carcinogenicity in an informative rodent model of liver
fluke infection-induced malignancy.
266
267 **Materials and Methods**
268
269 *Opisthorchis viverrini* **parasite preparation**
270
Metacercariae (MC) of *O. viverrini* were isolated from the naturally infected cyprinid fish by
pepsin digestion as previously described (*21*). In brief, fishes were minced with an electric

blender, and then minced tissues were digested with 0.25% porcine pepsin, 1.5% HCl in 150 mM
NaCl at 37ºC in for 2 hours. After tissue digestion, the tissue mixture was filtered sequentially
through 1,100, 350, 250, and 140 μm diameter pore size sieves; the final filtrate sedimented by
gravity and the aqueous supernatant was discarded. The *O. viverrini* MC enriched-sediment was
washed once with 150 mM NaCl (normal saline solution, NSS), and the identity of *O. viverrini*
MC confirmed using a stereomicroscope. The *O. viverrini* MCs were stored in NSS at 4ºC until
used. The newly excysted-juvenile (NEJs) were prepared from MC in 0.25% trypsin in 1×PBS
containing 2× 200U/ml penicillin, 200 μg/ml streptomycin (Gibco) (2× Pen/Strep) for 5 min at
RT prior to separate the cyst walls from juvenile parasite by insulin needle (*7, 22*). NEJ were
transferred to RPMI medium containing 1% glucose, 2 g/L $NaHCO_3$, 2× Pen/Strep and 1 μM E-
64 protease inhibitor (Thermo Fisher Scientific) and maintained at 37ºC, 5% CO_2 for 60 min
before use.
285
To obtain adult developmental stages of the liver fluke, Syrian golden hamsters (*Mesocricetus*
auratus) were infected at 6-8 weeks of age with 50 MC per hamster by intragastric tube (*23*).
The hamsters were maintained at the Animal Facility of Faculty of Medicine, Khon Kaen
University, Khon Kaen, Thailand. Sixty days after infection, hamsters were euthanized and the
liver flukes collected, as described (*22, 23*). The study protocol was reviewed and approved by
the Animal Ethics Committee of Khon Kaen University. The study adhered to standard
guidelines of the Ethics of Animal Experimentation of the National Research Council of
Thailand (approval number ACUC-KKU-61/60).
294
295 **Vector and guide RNA targeting exon 1 of *Ov-grn-1***
296
To edit the gene *Ov-grn-1* that encodes *O. viverrini* granulin-1 (6287 bp, mRNA GenBank
accession FJ436341.1), CRISPR online tools including CRISPR design (http://crispr.mit.edu/)
and ChopChop (http://chopchop.cbu.uib.no/) were employed to design a guide RNA (gRNA)
targeting exon 1 *Ov-grn-1* gene at nucleotide position 1589-1608,
GATTCATCTACAAGTGTTGA (Fig. 1a and 1b). A CRISPR/Cas9 encoding vector encoding
the above gRNA under the control of the mammalian U6 promoter and encoding Cas9 (with
nuclear localization signal 1 and 2) driven by the CMV promoter was assembled (GeneArt
CRISPR Nuclease Vector Kit, Thermo Fisher), and termed pCas-*Ov-grn-1* (Supplementary Fig.
1a). *Escherichia coli* TOP-10 competent cells were transformed with pCas-*Ov-grn-1* and vector
plasmid recovered from cultures of a positive clone (NucleoBond Xtra Midi, Macherey-Nagel
GmbH, Germany). The nucleotide sequence of pCas-*Ov-grn-1* was confirmed by Sanger direct
sequencing.
309
310 **Transfection of liver flukes with pCas-*Ov-grn-1***
311
Twenty mature flukes were transfected with 10 μg pCas-*Ov-grn-1* pDNA in ~500 μl RPMI-1640
(Sigma) by electroporation. The electroporation was performed in 4 mm cuvettes (Bio-Rad) with
a single square wave pulse of 125 volts for 20 ms using a Gene Pulser Xcell (Bio-Rad). Flukes
were then washed several times with NSS and an additional 5 times with RPMI-1640 containing
2× Pen/Strep. Flukes were cultured in RPMI-1640 containing 2× Pen/Step at 37°C in 5% CO_2
atmosphere (*7, 24*). Two control groups were included: wild type (WT) mature flukes and
'mock' control flukes which were exposed to identical electroporation conditions with RPMI-

1640 and 1× Pen/Strep in the absence of plasmid DNA. The adult flukes were observed and
collected after 1, 2, 3, 5, 7, 14 and 21 days of culture following pCas-*Ov-grn-1* transfection.
RNA and protein was extracted from flukes and *Ov-grn-1* mRNA expression was assessed by
RT-qPCR and *Ov*-GRN-1 protein expression was assessed by western blot. Mutations and/or
insertions-deletions (INDELs) resulting from CRISPR/Cas were analyzed by CRISPR efficiency
estimation (*17, 25*) and MiSeq Next Generation Sequencing (NGS).
325
326 MC and NEJ (750 parasites per cuvette) were subjected to square wave electroporation in the
327 presence of pCas-*Ov-grn-1* pDNA as described above for adult flukes. The parasites were
328 washed as above and cultured in RPMI complete medium (10% FBS, 2× Pen/Strep) at 37°C in
329 5% CO_2 in air. The parasites were collected on days 1, 2, 3 and 5 after transfection and *Ov-grn-1*
330 transcript levels were ascertained by RT-qPCR, as above.
331
332 **Extraction of nucleic acids**
333
RNA was extracted from pooled or individual flukes using TRIZOL (Invitrogen) according to
the manufacturer's recommendations. Concentration of RNA was estimated by 260 nm by
NanoVue spectrophotometer. Genomic DNA from individual or pooled (25 worms) flukes was
extracted using the QIAamp DNA Mini Kit (Qiagen, Valencia, CA). The dual RNA and DNA
extraction technique was used for some experiments with individual worms using RNAzol®RT
(Molecular Research Center, Inc.) and DNAzol (Molecular Research Center, Inc.) (*26, 27*).In
brief, each worm was homogenized in RNAzol®RT using a motorized pestle, the DNA and
protein from the lysate was precipitated using DNAse-RNAse-free water. The aqueous top
solution was transferred into isopropanol to precipitate the RNA. The DNA/protein pellet was
resuspended in DNAzol® RT, and DNA extracted as per the manufacturer's instructions.
Individual RNA samples were evaluated for *Ov-grn-1* expression. To assess variations in *Ov-*
grn-1 transcript levels, individual flukes were assigned into 3 groups; <10% of *Ov-grn-1*
transcript fold change (fc) compared to WT, 10-70% *Ov-grn-1* transcript fc compared to WT,
and 100-120% *Ov-grn-1* transcript fc compared to WT. Genomic DNA from individual worms
was used to assess CRISPR efficiency and estimate mutation levels (*25, 28*). Estimates of
mutation efficiency positively correlated with transcripts levels of *Ov*-*grn-1* transcript level (Fig.
4f). Pooled genomic DNA samples from low to high *Ov-grn-1* transcript levels and high to low
percent mutations from experimental groups were prepared for next-generation Illumina
sequencing.
Quantitative Real-time PCR
354
Complementary DNA (cDNA) was synthesized from parasite total RNA using an iScript cDNA
synthesis kit (Thermo Fisher Scientific) prior to proceeding with quantitative real-time PCR (RT-
qPCR). RT-qPCR was performed with biological triplicate samples using a SYBR Green kit
(TAKARA Perfect Real-time Kit) according to the manufacturer's recommendation in a Light
Cycler 480 II thermal cycler (Roche). Each RT-qPCR reaction consisted of 7.5 μl SYBR Green
Master Mix, 0.5 μl (10 μM) each of specific forward and reverse primers for *Ov-grn-1* (Fig. 1b)
(forward primer, *Ov-grn-1*-RT-F: 5′-GGGATCGGTTAGTCTAATCTCC and reverse primer,
Ov-grn-1-RT-R: 5′-GATCATGGGGGTTCACTGTC), amplifying 359 base pairs (bp) of the
product (position 7-365 nt of *O. viverrini* granulin-1 mRNA GenBank accession FJ436341.1), 2
μl of cDNA and distilled water to a final volume of 15 μl. The thermal cycle was a single

initiation cycle at 95°C for 3 min followed by 40 cycles of denaturation at 95°C for 30 sec,
annealing at 55°C for 30 sec, extension at 72°C for 45 sec and a final extension at 72°C for 10
min. The endogenous actin gene (GenBank accession AY005475) was used as a housekeeping
control (*7, 24*) (forward primer, *Ov*-actin-F: 5′-AGCCAACCGAGAGAAGATGA and reverse
primer *Ov*-actin-R: 5′-ACCTGACCATCAGGCAGTTC). The *Ov-grn-1* transcript fold change
was calculated by $2^{(-\Delta\Delta Ct)}$ method using *Ov-actin* for normalization (*7, 24, 29*). Means and
standard deviations were calculated using Graph Pad Prism software; two-way ANOVA.
372
373   **Rabbit anti-*Ov*-GRN-1 antiserum and immunoblot analysis**
374
One milligram of adjuvanted recombinant *Ov*-GRN-1 protein (*30*) was used to subcutaneously
inject an outbred New Zealand White rabbit. The rabbit was boosted twice with 500 μg of
adjuvanted protein and two weeks after the last booster the rabbit was sacrificed for blood
collection via cardiac puncture. The Animal Ethics Committee of Khon Kaen University
approved the protocols used for animal experimentation, based on the guidelines of the National
Research Council of Thailand for Ethics of Animal Experimentation (ACUC-KKU-61/60). *Ov*-
GRN-1 protein levels were determined by western blot using rabbit anti-recombinant *Ov*-GRN-1
antiserum. The adult flukes from either WT or Δ*Ov-grn-1* groups were collected individually at
days 1, 2, 3, 5, 7, 14 and 21 after electroporation (3 flukes per group). Groups of 3 flukes were
homogenized by sonication (Sonics & Materials) in 1× PBS with alternating pulses of 5 sec
duration (with 5 sec pause between pulses) for 45 sec at 4°C. The homogenate was centrifuged at
13,000 *g* at 4°C for 30 min and the supernatant collected and stored at -20°C. Protein
concentration of fluke homogenates was measured using the Bradford assay and homnogenates
were electrophoresed on 15% SDS-PAGE gels. Proteins were transblotted onto nitrocellulose
membrane using a Mini Trans-Blot Cell (Bio-Rad). Membrane strips containing 2 μg of total
protein were washed with 0.5% Tween-20 in PBS (PBST) then blocked for 1 hour with 5%
skimmed milk in PBST. Strips were incubated with rabbit anti-*Ov*-GRN-1 serum or pre-
immunization serum diluted 1:50 with 1% skimmed milk in PBST and incubated with shaking
for 2 h. Strips were washed then incubated for 1 hour with horseradish peroxidase (HRP)-goat
anti-rabbit IgG (Invitrogen) (diluted 1:1,000 in antibody buffer). The strips were washed again
and color reactions were detected by enhanced chemiluminescence (ECL) substrate (GE
Healthcare Life Sciences) and imaged using an Image Quant LAS 4000 mini (GE Healthcare
Life Sciences). As a control protein also derived from the tegument of *O. viverrini* flukes, we
assessed the protein expression levels of *Ov*-TSP-2 by western blot using a specific antibody
raised to the recombinant protein (*31*). Relative protein expression levels from western blots
were measured by densitometry using Image J (https://imagej.nih.gov/ij/download.html). Protein
expression levels were compared statistically between groups using independent *t*-tests.
402
403   **CRISPR/Cas efficiency and mutation levels estimated by quantitative SYBR green PCR**
404
Adult flukes were collected on days 1, 2, 3, 5, 7, 14 and 21 after pCas-*Ov-grn-1* transfection.
Each individual fluke DNA was investigated for mutation(s) around the expected double
stranded break (DSB) site by using three-primers; 2 forward primers and 1 reverse primer; *Ov-*
grn-1-OUT-F, *Ov-grn-1*-OVR-F and *Ov-grn-1*-reverse, respectively. The primer pair of *Ov-*
grn-1-OUT-F and *Ov-grn-1*-reverse was used for amplify the fragment flanking (1496-2312 nt)
the DSB, while another primer pair (*Ov-grn-1*-OVR-F and *Ov-grn-1*-reverse) was amplify

overlap of DSB site (1599-2312) (Fig. 1b). Both primer pairs exhibited equivalent amplification
efficiency with genomic DNA form WT flukes. The OUT and OVR amplicons were 450 and 347
bp, respectively, using PCR conditions using 7.5 μl of SYBR Green Master Mix (TAKARA
Perfect Real-time Kit), 0.5 μl (0.4 μM) of each primer, 10 ng/μl of gDNA and distilled water to
15 μl. The thermal cycles included initiation for one cycle at 95°C for 3 min followed by 40
cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, extension at 72°C for 45 s,
and a final extension at 72°C for 10 min. The SYBR green signal was read every annealing cycle
and reported as threshold cycle (Ct). Efficiency of programmed CRISPR/Cas editing was
estimated as the ratio of $Ct_{OUT}:Ct_{OVR}$ from experimental (A) compared with $Ct_{OUT}:Ct_{OVR}$ of
control group (B) as described (*17*). The $Ct_{OUT}:Ct_{OVR}$ ratio from control group would equal '1'
(CRISPR efficiency = 0) since differences were not seen in Ct value from OUT and OVR
primers. By contrast, the OVR primer is anticipated to be less efficient than the OUT primer for
the experimental group, and hence the $Ct_{OUT}:Ct_{OVR}$ is less than '1'. Here, we calculated %
mutation indirectly by subtraction of the CRISPR/Cas9 efficiency value from '1', as
indicated(*17, 32*).

$$\text{Efficiency (F)} = \frac{\text{Average Ct OUT}}{\text{Average Ct OVR}}$$

$$\text{CRISPR/Cas9 efficiency} = \frac{F_{\Delta Ov\text{-}grn\text{-}1}}{F_{Control}}$$

$$\text{Mutation rate} = 100\% - \text{CRISPR Cas9 efficiency}$$

Illumina-based deep sequencing

The MiSeq NGS library was constructed from pooled DNA samples of *Ov-grn-1* gene-edited
adult liver flukes at days 14 and 21 post-transfection. The 173 bp amplicon flanking the DSB
was amplified from MiSeq-F (position 1496-1514 nt): 5' TTCGAGATTCGGTCAGCCG-3' and
MiSeq-R (position 1649-1668 nt): 5'GCACCAACTCGCAACTTACA-3' primers (Fig. 1b). The
amplicon was purified (Agencourt AMPure XP beads, Beckman) and ligated with Gene Read
Adaptors Set A (Qiagen) and Illumina compatible adaptor(s) and barcode(s) using QIAseq 1-step
Amplicon library kit (Qiagen). The library was quantified using a GeneRead Library Quant Kit
(Qiagen) with Illumina index/barcode specific primers included in the kit. The concentration of
the MiSeq library was read against known standard libraries provided with the Library Quant
Kit. NGS was performed at GENEWIZ, South Plainland, NJ). The MiSeq NGS reads were
trimmed of index/adaptor and primer out sequences prior to further analysis for programmed
mutations by SnapGene software (GSL Biotech LLC) and CRISPResso analysis platform (*18,
19*) against the target amplicon of the reference (WT) *Ov-grn-1* gene.

Illumina-based deep sequencing

The MiSeq NGS library was constructed from pooled DNA samples of *Ov-grn-1* gene-edited
adult liver flukes at days 14 and 21 post-transfection. The 173 bp amplicon flanking the DSB
was amplified from *Ov-grn-1* MiSeq-F (position 1496-1514 nt): 5'

TTCGAGATTCGGTCAGCCG-3' and *Ov-grn-1* MiSeq-R (position 1649-1668 nt):
5'GCACCAACTCGCAACTTACA-3' primers (Fig. 1b). The amplicon was purified using
Agencourt AMPure XP beads (Beckman) and ligated with index/barcoded adapters compatible
with the Illumina system using Gene Read Adaptors Set A with the QIAseq 1-step Amplicon
library kit (Qiagen). The library was quantified using a GeneRead Library Quant Kit (Qiagen)
with Illumina index/barcode specific primers supplied in the kit. The concentration of the MiSeq
library was compared with known standard libraries provided with the Library Quant Kit. The
MiSeq reads (NGS performed by GENEWIZ, South Plainfield, NJ) were trimmed of
index/adaptor and primer out sequences before downstream analysis for programmed mutations
by SnapGene software (GSL Biotech LLC) and the CRISPResso platform (*18, 19*) against the
target amplicon from the reference WT *Ov-grn-1* gene.
468
469   **Cell proliferation and *in vitro* wound healing assay**
470
To evaluate the effect of *Ov-grn-1* gene editing on liver fluke-driven proliferation of human
cholangiocytes, motile WT or Δ*Ov-grn-1* adult flukes were co-cultured with cell of the human
cholangiocyte cell line H69 in 24-well Trans-well plates (3 wells per groups) (*7*) containing a 4
μm pore size membrane separating the upper and lower chambers (Corning).In brief, 15,000 H69
cells were seeded into the lower chamber of the plate and cultured with complete medium
containing DMEM/F12 supplemented with 1× antibiotic, 10% fetal bovine serum, 25 μg/ml
adenine, 5 μg/ml insulin, 1 μg/ml epinephrine, 8.3 μg/ml holo-transferrin, 0.62 μg/ml
hydrocortisone, 1.36 μg/ml T3 and 10 ng/ml EGF (*33*) for 24 hours, after which the cells were
fasted for 4-6 hours in medium supplemented with only one twentieth of the growth factor
content of complete medium. Five viable *O. viverrini* adult flukes that had been transfected (or
not) with CRISPR/Cas9 *Ov-grn-1* plasmid in a total of 500 μl of RPMI (or medium alone) were
placed into the upper chamber of each well. The number of cells in each well was determined at
days 1, 2 and 3 using 1× PrestoBlue cell viability reagent (Invitrogen) (*34*) and added to cells at
37°C for up to 1 hour. Cell number was determined at 570 nm and calculated from a standard
curve before transforming into relative growth compared to control groups. Cell proliferation
assay was carried out in triplicate.
487
To assess the effect of *Ov-grn-1* knockout on *in vitro* wound healing, 3×10^5 H69 cholangiocytes
in monolayer were grown in 6-well Trans-well plates with a 4 μm pore size. H69 cells were
cultured in complete media for 2 days at 37°C then transferred to incomplete media overnight.
Monolayers in each well were scratched using a sterile 200 μl autopipette tip (*7, 8, 35*) and
washed with PBS twice to remove disconnected cells or debris. Ten transfected adult or control
flukes were added to the upper chamber of the Trans-well plate containing the wounded cell
monolayer in the lower chamber. The migration rate of cell wound closure was measured at 0,
12, 24 and 36 hours, respectively. Trans-well plates were imaged using an inverted microscope
(Nikon) and images of all groups were captured at all-time points quantitatively using Adobe
Photoshop CS6. The distances between different sides of the scratch were measured by drawing
a line in the middle of the scratch on the captured image (*7, 8, 35, 36*). The analysis of wound
healing was carried three times.
500
501   **Infection of hamsters with *Ov-grn-1* gene-edited NEJs and assessment of hepatobiliary**
502   **histopathology**

503
Thirty male Syrian golden hamsters, 6-8 weeks of age, were obtained from the Animal Unit,
Faculties of Medicine, Khon Kaen University (approval number ACUC-KKU-61/60). The
hamsters were randomly divided into three groups of 10 per group: uninfected control, infected
with WT flukes and infected with Δ*Ov-grn-1* flukes. Each hamster was infected with 100 active
NEJs through intragastric intubation; the uninfected control group was fed normal saline solution
instead of NEJ (*23*). Hamsters (5 animals per cage) were contained under conventional
conditions and fed a stock diet (C.P. Ltd., Thailand) and water *ad libitum* until they were
euthanized (*23*). Following euthanasia, five hamsters from each group were necropsied for
histopathological assessment of the hepatobiliary tract at day 14 and at day 60 post-infection
(*23*). The hamsters were euthanized by overdose of anesthesia with diethyl ether. Subsequently,
blood was obtained by cardiac puncture and the livers were removed. Fluke numbers were
counted from two livers of both, WT and Δ*Ov-grn-1* groups at day 60 after infection of hamsters
and compared with unpaired two-tailed *t*-test. The left and right lobes of the liver from five
hamsters were dissected, cross-sectioned and each lobe was divided into three parts. The liver
fragments were fixed in 10% buffered formalin and stored overnight at 4°C before processing.
Formalin-fixed liver was dehydrated through an ethanol series (70, 95 and 100%), cleared in
xylene and embedded in paraffin. Paraffin embedded sections of 4 μm thickness, cut by
microtome, were stained with hematoxylin and eosin (H&E) or Picro-Sirius Red, or probed with
anti-ACTA2 antibodies, and analyzed for pathologic changes (below).
523
524 **Biliary hyperplasia**
525
H&E staining was used to assess pathological changes. The sections were deparaffinized in
100% xylene, rehydrated with descending series of alcohol, stained with H&E for 5 min,
dehydrated with an ascending series of alcohol, cleared with 100% xylene, mounted on a slide
with permount media, and slides were dried at 37°C overnight and photographed under light
microscopy. Images (200×) from H&E stained liver sections from 5 untreated control hamsters
or 5 hamsters per treatment group (WT and Δ*Ov-grn-1* groups) were assessed. Thickness (width)
of the bile duct epithelium from each thin liver section was measured with ImageJ at eight
equidistant positions around the bile duct. To compensate for outliers the median width for each
bile duct was used for graphing and analysis. The two-way ANOVA Holm-Sidak multiple
comparison test was used to compare each group at each time point.
536
537 **Fibrosis**
538
Two stains were used separately to assess biliary fibrosis. First, sections were stained with Picro-
Sirius Red (Abcam, Cambridge Science Park, UK). Sufficient Picro-Sirius Red solution was
applied to completely cover the tissue sections on the slide, the stained slide was incubated at
ambient temperature for 60 min, rinsed in two changes of acetic acid solution, and dehydrated
through two changes of absolute ethanol. Slides were cleared with 100% xylene, mounted in Per-
mount, dried at 37°C overnight, and examined and photographed by light microscopy to
document collagen surrounding the bile ducts. ImageJ was used to auto-color balance the images
using the macro written by Vytas Bindokas; Oct 2006, Univ. of Chicago
(https://digital.bsd.uchicago.edu/docs/imagej_macros/_graybalancetoROI.txt) followed by
application of the MRI fibrosis tool to determine percentage area of red-stained fibrosis at default

settings (red 1: 0.148, green 1: 0.772, blue 1: 0.618, red 2: 0.462, green 2: 0.602, blue 2: 0.651,
red 3: 0.187, green 3: 0.523, blue 3: 0.831) (*37*). Twenty discrete images (200×) stained with
Picro-Sirius Red from each animal (five hamsters per treatment group) were assessed (in total,
100 images per group). Kruskal-Wallis with Dunn's multiple comparison test was used to
compare the findings due to the range of data points among the groups.
554
Fibrosis was also assessed via levels of smooth muscle alpha-actin (ACTA2). Liver sections
from hamsters at day 60 post-infection were deparaffinized three times with 100% xylene, 5 min
each. Sections were rehydrated with ascending series of ethanol; 100%, 3 times, 3 min each,
95% 3 times, 3 min each, 70% for 3 min, followed by thorough washing in tap water for 5 min,
distilled water for 5 min, and PBS for 5 min. Thereafter, slides were incubated in citrate buffer
pH 6.0 (citric acid (anhydrous) 1.92 g and Tween 20 0.5 ml in total volume of DW 1000 ml) at
110°C for 5 min, allowed to cool for 20 min, and then washed in PBS 3 times, 5 min each.
Sections were then blocked with 5% bovine serum albumen (BSA) for 30 min in a humidified
chamber and washed in PBS 3 times, 3 min each with occasional shaking. The slides were
probed with Alexa Fluor 594 –labeled anti-ACTA2 antibody (Abcam) diluted 1:200 in 1% BSA
in PBST, 18 hours at 4°C in a humidified atmosphere. Lastly, slides were washed in PBS 3
times, 3 min each with occasional shaking, mounted in glycerol diluted 1:4 with PBS and
examined under bright and fluorescence light (Zeiss Axio Observer, with AxioVision SE64 Rel.
4.9.1 software, Jena, Germany). Images with a bile duct containing a fluke were selected and
ImageJ was used to manually select a narrow strip surrounding the bile duct epithelial layer that
excluded potential blood vessels (any enclosed curved oval-like shape). Three liver regions not
including the bile duct or blood vessels were manually selected for background fluorescence
measurements, each comprising 5-10% of the image. The fluorescence intensity of the bile duct
epithelial strip was measured and blanked against the average of the three background regions
and reported as average intensity per cm^2 at 300 PPI (pixels per inch). A total of 25-30 distinct
bile duct images per group (3 animals) were assessed. Zero values from the uninfected group
were deemed to have a value of 1 to allow a log axis. The groups were compared with one-way
ANOVA with Holm-Sidak multiple comparison test.
578
579   **ACKNOWLEDGMENTS**
580

We thank Suwit Balthaisong for technical assistance. This study was supported by the Thailand
Research Fund through the Royal Golden Jubilee PhD Program (grant no. PHD/0111/2557 to PA
and TL), the National Cancer Institute (NCI), US National Institutes of Health (NIH) award
R01CA164719 (AL, TL, PJB,) and the National Health and Medical Research Council (NHMRC
award APP1085309 to AL, JS and TL, and senior principal research fellowship APP1117504 to
AL). The content is solely the responsibility of the authors and does not necessarily represent the
official views of the NIAID, NCI, NIH or NHMRC. These studies were supported in part by
Wellcome Trust Strategic Award number 107475/Z/15/Z (Karl F. Hoffmann, principal
investigator; PJB, co-investigator).

590
591   **AUTHOR CONTRIBUTIONS**
592

593   T. Laha, P.J. Brindley, W. Ittiprasert , P. Arunsan, N. D. Young, A. Loukas, J. Sotillo, M. Smout,
594   S. Chaiyadet, C. Cochran, and V. Mann conceived and designed the research; P. Arunsan, M.

601
602 The authors declare no competing interests.

641
642   (*6*)**Figure legends**
643
Figure 1 CRISPR/Cas9-mediated gene editing strategy to knock-out *Ov-grn-1* in adult
***Opisthorchis viverrini* liver flukes. a**, Schematic depiction of *Ov-grn-1* gene with CRISPR/Cas9
target site in first exon marked with green arrow. **b**, The exon1, 2 and 3 location of *Ov-grn-1*
gene, size of 6,267 bp and gRNA targeting exon1: gRNA sequence is blue color and PAM is red
in bracket. Primer pairs were used to detect mRNA expression levels by using RT-qPCR assay
(*Ov-grn-1* RT- forward or RT-F and *Ov-grn-1* RT-reverse or RT-R), Tri-primers were used to
detect the % relative fold amplicon or mutations (outside-forward or OUT-F, overlap-forward or
OVR-F and reverse primer or OUT/OVR-R) and MiSeq forward and reverse (MiSeq-F and
MiSeq-R) primers were used to prepare the NGS amplicon. **c**, CRISPR/Cas9-induced adult
fluke genomic insertion (red bars) and deletion (black bars) mutations (INDELs) detected in the
Ov-grn-1 gene; CRISPR target site is denoted by the green arrow. Average mutation length is
plotted against *Ov-grn-1* gene amplicon position in base pairs (bp). **d**, Somatic tissues of
individual adult worms (in triplicate per time per group) were solubilized, electrophoresed in
SDS-PAGE gels, transferred to nitrocellulose membrane and probed with anti-*Ov*-GRN-1 rabbit
antibody. WT: wild type control fluke tissues; D1-21: Δ*Ov-grn-1* fluke tissues sampled the arrow
highlighting the ~9 kDa *Ov*-GRN-1 band at different days post-transfection and Δ*Ov-grn-1*
flukes show similar *Ov*-TSP-2 protein (control antibody) expression levels. D1-21 = protein
products from flukes day 1-21 post Δ*Ov-grn-1* treatment. Western blot panels of probed with
anti-*Ov*-TSP-2 rabbit antibody, the arrow highlighting the ~24 kDa *Ov*-TSP-2 band. **e**, Reduced
expression of *Ov-grn-1* mRNA and *Ov*-GRN-1 protein after transfection of adult flukes with *Ov-*
grn-1 CRISPR/Cas9 construct using quantitative real-time PCR (mRNA) and densitometry
conversion of Western blot signals (protein). Data are plotted relative to wild type (WT) fluke
values (100%) as an average percentage from 3 replicates with SD error bars. **** = $P<0.0001$
compared to WT fluke protein (black) or RNA (pink) at each time point with two-way ANOVA
Holm-Sidak multiple comparison test.

Figure 2 Δ*Ov-grn-1* adult fluke ES products induce less *in vitro* cell proliferation and
wound repair. a, Representative cell proliferation images of H69 immortalized human
cholangiocyte cell line co-cultured with flukes in Trans-well plates; mock transfected (top) and
Δ*Ov-grn-1* (bottom) groups are shown at day 3. **b**, Reduced cell proliferation induced by Δ*Ov-*
grn-1 fluke ES products, as shown in panel a, quantified from days 1-6. Data is plotted as
average relative percentage to cells cultured with "no flukes". **c**, Representative image of H69
cholangiocyte scratch wound repair when cells were co-cultured in Trans-well plates with flukes.
Mock transfected (top) and Δ*Ov-grn-1* (bottom) groups are shown at 0 and 36 h post-scratch
wounding. Dotted line shows the edge of the wound. **d**, Scratch wound repair assay quantified
from 0 to 36 hours, reveling diminished healing in the Δ*Ov-grn-1* group. Panels **b** and **d**: mean ±
SD, three replicates; ****$P<0.0001$ compared to wild type flukes with two-way ANOVA Holm-
Sidak multiple comparison test.
681
Figure 3 Δ*Ov-grn-1* newly excysted juveniles can infect hamsters and drive reduced short-
term pathology. Images of infective metacercariae (**a**) and newly excysted juvenile flukes
(NEJ) (**b**). **c**, *Ov-grn-1* mRNA expression in mock-transfected and Δ*Ov-grn-1* flukes over 5 days
as quantified with qPCR and plotted relative to the wild type (WT) untreated group; mean ± SD,
three replicates; **** = $P<0.0001$ compared to WT flukes with two-way ANOVA Holm-Sidak

multiple comparison test at each time point. **d**, Representative image of 400x magnification of
H&E stained thin section from hamster liver at 14 days after infection with WT flukes. **e**,
Representative image of H&E stained thin sections showing hamster liver 14 days after infection
with Δ*Ov-grn-1* flukes. **f**, Representative image of H&E stained thin section of control,
uninfected hamster liver, which shows healthy, organized pavement-like profile of the cells of
the biliary epithelium (BE) enclosing the lumen of the bile duct (BD) near a blood vessel (BV)
within the liver (L). Infection by WT flukes (panel **d**) revealed thickened, disordered epithelium
adjacent to the parasite (*Ov*). Infection with Δ*Ov-grn-1* flukes (panel **e**) revealed a bile duct
epithelium more closely resembling the normal, uninfected hamster. **g**, Epithelium
width/hyperplasia (green bracket) was quantified using ImageJ and plotted as the mean ± SD of
five biological replicates (hamsters). Significant difference were apparent when compared to the
uninfected group using the two-way ANOVA with Holm-Sidak multiple comparison test:
***P<0.01* and ********P<0.0001*, and wild-type compared to Δ*Ov-grn-1*, [#] *P<0.05* and
[####]*P<0.0001*.
701
Figure 4. Diminished level of fibrosis in hamsters during chronic infection with gene-edited
Δ*Ov-grn-1* liver flukes. Panel **a**, Adult fluke numbers were counted from the livers at day 60
post-infection and shown as the average and range for two hamsters per group . Fluke counts
showed minimal (12%; *ns*) difference between the wild type (WT) and Δ*Ov-grn-1* groups. **b,**
Representative images of Sirius red stained hamster liver (L) sections from uninfected animals
show minimal collagen fibrosis (fb) deposition, stained red surrounding the blood vessel (BV)
and bile ducts (BD) with normal biliary epithelia (BE). Livers from hamsters infected with WT
flukes (*Ov*) show heavy collagen deposition with elongated BE cells adjacent to resident flukes. .
Δ*Ov-grn-1* fluke infected livers showed substantial collagen deposition compared to uninfected
liver sections, yet showed far less than the livers infected with WT flukes. **c**, Liver fibrosis
quantified with ImageJ MRI-fibrosis plugin shown as a violin plot. 100 images containing bile
ducts from 20 sections (5 animals) per group showing SD as a vertical line with the median
indicated by the central black dot. Δ*Ov-grn-1* group showed a 23% reduction in median fibrosis
compared to WT group. The width of the "violin-shape" represents measurement frequency.
Kruskal-Wallis with Dunn's multiple comparison test used to compare groups. Against
uninfected: *********P<0.0001*, and Δ*Ov-grn-1* against WT: ### *P<0.001*. **d**, Representative
immunofluorescence/bright field overlay images of liver sections probed with anti-ACTA2
antibody with fluorescence intensity shown on a blue/green/red scale. ACTA2 protein is always
detected in my fibroblasts surrounding blood vessels, but not near normal bile ducts. When
detected near the BE layer it is suggestive of myofibroblast generation as a response to BE
damage. The upper panels (Overlay) show a wide view of the liver sections with a boxed region
that highlights a section of interest that is magnified in the lower panels (Zoom). Hamster liver
sections exhibited intense fluorescence (arteries: red/green, veins: blue/green) surrounding BVs
whereas only minimal fluorescence was seen near BD in livers of uninfected hamsters The
highlighted magnified (Zoom) panels show WT infected animal livers with weak (blue) but
consistent ACTA2 staining surrounding thickened BE layer (inner and outer cell edge marked
with orange dotted line) around BDs with WT flukes. Livers from hamsters infected with Δ*Ov-*
grn-1 flukes show weak and irregular ACTA2 staining surrounding BD. **e**, Quantified levels of
ACTA2 surrounding BDs from hamster liver sections. Violin plot with reverse log2 Y-axis
showing the ACTA2 intensity (per cm^2 at 300 PPI) surrounding BE from 25-30 discrete BD
images per group (three hamsters) assessed with ImageJ. Zero values from uninfected group

were deemed to have a value of 1 so that a log axis could be plotted. SD indicated as a line with
mean value marked as a central black dot and the violin-shape width representing measurement
frequency. ACTA2 staining shows 94% median reduction in Δ*Ov-grn-1* fluke-infected livers
compared to WT fluke-infected animals. One-way ANOVA with Holm-Sidak multiple
comparison test, ****P<0.0001 compared to uninfected and ##P<0.001 compared to Δ*Ov-grn-1*
flukes against WT flukes. **f**, *Ov-grn-1* mRNA expression levels are reduced in Δ*Ov-grn-1* flukes
compared to WT flukes. Reverse log10 Y-axis shows the qPCR $2^{\wedge}(-\Delta\Delta Ct)$ results from flukes 60
days after programmed CRISPR/Cas9 gene editing and hamster infection plotted relative to
mean value of the WT infection. The WT group showed a broad level of expression; the mean
expression level for the Δ*Ov-grn-1* flukes was 19.4% of the WT group; Mann-Whitney
nonparametric test, **P<0.01. While significantly reduced overall, the Δ*Ov-grn-1* fluke group
ranged from no apparent effect to markedly diminished expression of *Ov-grn-1*. Mutation
frequency was assessed by assigning the Δ*Ov-grn-1* group of flukes into three sub-groups based
on CRISPR/Cas9 mutation frequency. Eight flukes with highly effective CRISPR gene knock-
out (Δ*Ov-grn-1*: *Ov-grn-1*<10% expression), seven flukes with modest levels of transcript
knock-out (Δ*Ov-grn-1*: 10-70% *Ov-grn-1*) and 10 flukes with little or no effect (Δ*Ov-grn-1*: *Ov-*
grn-1 100-120%). The mean mutation frequency amongst the three Δ*Ov-grn-1* sub-groups was
2.7%.

779
780 **Supporting information**
781
Supplementary Figure 1. CRISPR/Cas9 targeting *Ov-grn-1* design and flukes transfection.
a, *Ov-grn-1* guide RNA insertion and Cas9 cassette. **b**, The original *Ov-grn-1* sequence with the
target (red text) and PAM (blue text) sites highlighted. The dark green arrow indicates the
specific CRISPR target site. Deep sequencing is used to determine specific mutations and the
four sequences below the original sequence show examples of successfully mutated *Ov-grn-1*
genes with variations with red boxed bases highlighted mutations induced by the CRISPR/Cas9
transfection. All four examples show a range of insertion, deletion or mismatch mutations boxed
in red. The "+2 insertion" with a dotted light green box highlights two alanine insertions near the
target site. The "-6 deletion" highlights six absent base pairs with a light blue box which include
the target site. **c**, The mutation rates were calculated at each time point from Δ*Ov-grn-1 in vitro*
cultured adult flukes with qPCR and the averages and standard deviation were determined . RT-
qPCR electrophoresis gel (lower image) shows *Ov-grn-1* RNA levels are reduced in Δ*Ov-grn-1*
adult flukes.
795
Supplementary Figure 2. Extended image set showing Δ*Ov-grn-1* adult fluke ES products
induce less *in vitro* cell proliferation and wound repair. a, Representative cell proliferation
images of H69 immortalized human cholangiocyte cell line co-cultured with flukes in Trans-well
plates; shown at 24 and 72hrs. **b**, Comparing the difference in cell proliferation values from Fig.
2b between ES products from the *ΔOv-grn-1* flukes and ES from WT flukes revealed a linear
correlation of reduction of cell proliferation that increases over time. **c**, Representative image of
scratch wound repair of cultured cholangiocytes when H69 cells were co-cultured in Trans-well
plates with flukes; time points from 0 - 36 hours post-scratch. Dotted line indicates the margin of
the wound. **d**, Scratch wound repair assay quantified values from Fig. 2d showed an increasing
linear correlation over time of reduced wound closure when plotting the difference between the
Δ*Ov-grn-1* and the WT fluke groups. Panels **b** and **d**: mean ± SD, three biological replicates.
807
Supplementary Figure 3. CRISPR/Cas9 ineffective at silencing gene expression in
metacercariae (MC). Expression of *Ov-grn-1* mRNA in wild type (WT), mock-transfected, and
Δ*Ov-grn-1* MC over five days as quantified with qPCR and plotted relative to the WT untreated
group; mean ± SD from three replicates. ****$P<0.0001$; Δ*Ov-grn-1* compared to WT control
group, $P>0.05$; non-significant (ns) difference seen between mock (control) group using the
two-way ANOVA Holm-Sidak multiple comparison test.
814
815 **Supplementary Figure 4. Representative wide-angle view of anti-ACTA2**
816 **immunofluorescence and bright field liver sections.** Thin liver sections presented in bright
817 field or under fluorescent light following probing with anti-ACTA2 antibody; fluorescence
818 intensity shown on a blue/green/red scale. These images were used to create the bright
819 field/immunofluorescence overlay and zoomed images in Fig. 4d.

825
826 **References**
827

1. B. Sripa *et al.*, Opisthorchiasis and Opisthorchis-associated cholangiocarcinoma in
Thailand and Laos. *Acta Trop* **120 Suppl 1**, S158-168 (2011).
2. J. S. Pagano *et al.*, Infectious agents and cancer: criteria for a causal relation. *Semin*
Cancer Biol **14**, 453-471 (2004).
3. T. Laha *et al.*, Gene discovery for the carcinogenic human liver fluke, Opisthorchis
viverrini. *BMC genomics* **8**, 189 (2007).
4. N. Khuntikeo *et al.*, Cohort profile: cholangiocarcinoma screening and care program
(CASCAP). *BMC Cancer* **15**, 459 (2015).
5. N. Khuntikeo, W. Loilome, B. Thinkhamrop, N. Chamadol, P. Yongvanit, A
Comprehensive Public Health Conceptual Framework and Strategy to Effectively
Combat Cholangiocarcinoma in Thailand. *PLoS Negl Trop Dis* **10**, e0004293 (2016).
6. P. J. Brindley, A. Loukas, Helminth infection-induced malignancy. *PLoS Pathog* **13**,
e1006393 (2017).
7. A. Papatpremsiri *et al.*, Suppression of Ov-grn-1 encoding granulin of Opisthorchis
viverrini inhibits proliferation of biliary epithelial cells. *Exp Parasitol* **148**, 17-23 (2015).
8. M. J. Smout *et al.*, Carcinogenic Parasite Secretes Growth Factor That Accelerates
Wound Healing and Potentially Promotes Neoplasia. *PLoS Pathog* **11**, (2015).
9. K. F. Hoffmann, P. J. Brindley, M. Berriman, Medicine. Halting harmful helminths.
Science **346**, 168-169 (2014).
10. P. D. Hsu, E. S. Lander, F. Zhang, Development and applications of CRISPR-Cas9 for
genome engineering. *Cell* **157**, 1262-1278 (2014).
11. J. D. Sander, J. K. Joung, CRISPR-Cas systems for editing, regulating and targeting
genomes. *Nat Biotechnol* **32**, 347-355 (2014).
12. S. Waaijers, M. Boxem, Engineering the Caenorhabditis elegans genome with
CRISPR/Cas9. *Methods* **68**, 381-388 (2014).
13. J. B. Lok, H. Shao, H. C. Massey, X. Li, Transgenesis in Strongyloides and related
parasitic nematodes: historical perspectives, current functional genomic applications and
progress towards gene disruption and editing. *Parasitology* **144**, 327-342 (2017).
14. S. S. Gang *et al.*, Targeted mutagenesis in a human-parasitic nematode. *PLoS Pathog* **13**,
e1006675 (2017).
15. N. D. Young *et al.*, The Opisthorchis viverrini genome provides insights into life in the
bile duct. *Nat Commun* **5**, 4378 (2014).
16. A. N. Shah, C. F. Davey, A. C. Whitebirch, A. C. Miller, C. B. Moens, Rapid reverse
genetic screening using CRISPR in zebrafish. *Nat Methods* **12**, 535-540 (2015).
17. C. Yu, Y. Zhang, S. Yao, Y. Wei, A PCR based protocol for detecting indel mutations
induced by TALENs and CRISPR/Cas9 in zebrafish. *PLoS One* **9**, e98282 (2014).
18. M. C. Canver *et al.*, Integrated design, execution, and analysis of arrayed and pooled
CRISPR genome-editing experiments. *Nat Protoc* **13**, 946-986 (2018).
19. L. Pinello *et al.*, Analyzing CRISPR genome-editing experiments with CRISPResso. *Nat*
Biotechnol **34**, 695-697 (2016).
20. M. Guido, M. Rugge, G. Leandro, I. M. Fiel, S. N. Thung, Hepatic stellate cell
immunodetection and cirrhotic evolution of viral hepatitis in liver allografts. *Hepatology*
26, 310-314 (1997).

21. P. Sithithaworn, V. Pipitgool, T. Srisawangwong, D. B. Elkins, M. R. Haswell-Elkins,
Seasonal variation of Opisthorchis viverrini infection in cyprinoid fish in north-east
Thailand: implications for parasite control and food safety. *Bull World Health Organ* **75**,
125-131 (1997).
22. A. Papatpremsiri, Effect of radiation attenuation of metacercariae and gene silencing
targeting granulin on infection with Opisthorchis viverrini, Doctor of Philosophy in
biomedical science, Khon Kaen University, Thailand, (2014).
23. B. Sripa, S. Kaewkes, Gall bladder and extrahepatic bile duct changes in Opisthorchis
viverrini-infected hamsters. *Acta Tropica* **83**, 29-36 (2002).
24. S. Piratae *et al.*, Molecular characterization of a tetraspanin from the human liver fluke,
Opisthorchis viverrini. *PLoS Negl Trop Dis* **6**, e1939 (2012).
25. H. Yang, J. J. Wu, T. Tang, K. D. Liu, C. Dai, CRISPR/Cas9-mediated genome editing
efficiently creates specific mutations at multiple loci using one sgRNA in Brassica napus.
Sci Rep **7**, 7489 (2017).
26. S. N. Chan, N. Abu Bakar, M. Mahmood, C. L. Ho, N. A. Shaharuddin, Molecular
cloning and characterization of novel phytocystatin gene from turmeric, Curcuma longa.
Biomed Res Int **2014**, 973790 (2014).
27. H. Chen, M. Rangasamy, S. Y. Tan, H. Wang, B. D. Siegfried, Evaluation of five
methods for total DNA extraction from western corn rootworm beetles. *PLoS One* **5**,
e11963 (2010).
28. J. J. Vasquez, C. Wedel, R. O. Cosentino, T. N. Siegel, Exploiting CRISPR-Cas9
technology to investigate individual histone modifications. *Nucleic Acids Res*, (2018).
29. T. D. Schmittgen, K. J. Livak, Analyzing real-time PCR data by the comparative C(T)
method. *Nat Protoc* **3**, 1101-1108 (2008).
30. O. Strannegard, A. Yurchision, Formation of rabbit reaginic antibodies to protein and
hapten-protein conjugates. *Immunology* **16**, 387-397 (1969).
31. S. Chaiyadet *et al.*, Suppression of mRNAs encoding CD63 family tetraspanins from the
carcinogenic liver fluke Opisthorchis viverrini results in distinct tegument phenotypes.
Sci Rep **7**, 14342 (2017).
32. M. F. Sentmanat, S. T. Peters, C. P. Florian, J. P. Connelly, S. M. Pruett-Miller, A Survey
of Validation Strategies for CRISPR-Cas9 Editing. *Sci Rep* **8**, 888 (2018).
33. K. Ninlawan *et al.*, Opisthorchis viverrini excretory/secretory products induce toll-like
receptor 4 upregulation and production of interleukin 6 and 8 in cholangiocyte. *Parasitol*
Int **59**, 616-621 (2010).
34. R. J. Tynan *et al.*, A comparative examination of the anti-inflammatory effects of SSRI
and SNRI antidepressants on LPS stimulated microglia. *Brain Behav Immun* **26**, 469-479
(2012).
35. C. C. Liang, A. Y. Park, J. L. Guan, In vitro scratch assay: a convenient and inexpensive
method for analysis of cell migration in vitro. *Nat Protoc* **2**, 329-333 (2007).
36. M. J. Smout *et al.*, Infection with the carcinogenic human liver fluke, Opisthorchis
viverrini. *Mol Biosyst* **7**, 1367-1375 (2011).
37. M. A. Pereira, Tratamento com células derivadas do fígado embrionário retarda a
progressão da fibrose hepática em ratos, Tese de Doutorado, Universidade de São Paulo,
Brazil, (2016).
915

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4