Evolution of phenotypic plasticity in response to ethanol between sister species with different ecological

histories (*Drosophila melanogaster* and *D. simulans*)

Sarah A. Signor[1*], Sergey V. Nuzhdin[1]

1. Molecular and Computational Biology, Dornsife College of Letters, Arts and Sciences, University of

Southern California.

* Communicating author: ssignor@usc.edu

Running title: Evolution of plasticity to ethanol

Please note: Supplemental Files 1-3 are currently available at: https://github.com/signor-

molevol/simulans-ethanol.git

Abstract

The role of phenotypic plasticity in evolution is contentious, in part because different types of plasticity –

adaptive, neutral, or non-adaptive, are often not distinguished. Adaptive plasticity is expected to facilitate

expansion into new environments, while non-adaptive plasticity will result in a mean phenotype further

from the adaptive optimum and/or an increase in variance due to the expression of variation that was

neutral and shielded from selection in the prior environment. We explore these patterns here by exposing

Drosophila melanogaster and *D. simulans* to high ethanol concentrations, with the knowledge that *D.*

melanogaster associates with high concentrations of ethanol in nature while *D. simulans* does not. Using

changes in gene expression and splicing we find that in *D. simulans* there is a large genotype-specific

response to ethanol, orders of magnitude larger than any effect that is not genotype-specific. In *D.*

melanogaster the response to ethanol is limited, and it is concordant among different genotypes. This

response to ethanol in *D. simulans* is enriched for non-protein coding nested genes that do not have

orthologs in *D. melanogaster*, suggesting rapid evolution of transcription. Sequence variation in ethanol-

implicated genes is consistent with balancing selection or selection relaxation in *D. simulans*, and they are

more divergent from *D. melanogaster* than the genome average. Overall, these patterns are consistent

with a maladaptive passive response to ethanol-induced stress in *D. simulans*, while in *D. melanogaster*

the reduced response and lack of genotype-specific variation suggests selection for an optimal response.

Keywords: phenotypic plasticity, *Drosophila*, ethanol, alternative splicing

Introduction

Genetic variation occurs when there are differences in a phenotype due to genotypic differences among individuals within a population (Figure 1A). In contrast, plasticity – i.e. environment specific adjustment in phenotype - can occur without genetic variation between individuals and is a change in a given phenotype in response to an external stimulus (Figure 1A). Plasticity is a ubiquitous property of organisms, and it may be adaptive, neutral, or maladaptive with regard to the fitness of a plastic individual (Ghalambor *et al*. 2007; Lande 2009; Marais *et al*. 2013). Genetic variation in plasticity describes variation between genotypes for the response to the environment (Figure 1A). Quantifying genetic variation, plasticity, and genetic variation for plasticity is important for understanding the genotype to phenotype map and adaptation. For example, the evolution of a reduced plastic response can reconfigure the relationship between fitness and the environment, redrawing the genotype to phenotype map. In another example, genetic variation in plasticity can make the evolution of locally adapted populations unpredictable in response to environmental change.

It is often assumed that plasticity is adaptive, however phenotypic plasticity can be adaptive, neutral, or maladaptive. This depends in part on the evolutionary history of exposure to the agent causing plasticity. Were such exposure infrequent, plasticity could arise as a passive consequence of environmental stress. This is likely an important, or perhaps prevalent, form of phenotypic plasticity (Ghalambor *et al*. 2007; 2015). One potential prediction arising from stress inducing passive phenotypic plasticity is that within population variation for performance is expected to be greater, as stress breaks down buffering mechanisms that might normally reduce between individual variation (Joshi & Thompson 1997). Thus under lower stress conditions, populations will exhibit less individual and genetic variation compared to high stress, rare conditions (Ghalambor *et al*. 2007). The novel or rare high stress environment is essentially uncovering cryptic variation that was allowed to neutrally accumulate in lower stress environments where selection was unable to shape the reaction norm (Rutherford 2000).

When a population colonizes a new habitat, this could manifest as a transient increase in genotype

by environment interactions, followed by a reduction in genetic variation for plasticity, assuming that

there is selection for a single optimal phenotype in response to the environment (Via & Lande 1985;

Guntrip & Sibly 1998; Lande 2009; Matzkin 2012; Huang & Agrawal 2016). The maintenance of

plasticity, or loss of plasticity, would then depend upon the frequency with which the population

continues to experience multiple environments, or the cost of maintaining the plastic response (Joshi &

Thompson 1997; Lande 2009; Scoville & Pfrender 2010; Lee *et al*. 2011; Hodgins-Davis *et al*. 2012).

Thus if phenotypic plasticity is costly to maintain, or the organism no longer experiences the previous

environment, the expectation is that phenotypic plasticity will transition to local adaptation.

In this manuscript we will examine plasticity and genetic variation for plasticity in response to

ethanol in *D. simulans* and *D. melanogaster*, two species with divergence adaptive histories with respect

to ethanol. *D. melanogaster* and *D. simulans* are thought to have originated in or around Africa (*D.*

melanogaster, southern-central Africa, *D. simulans*, Madagascar) followed by an out-of-Africa expansion

~10,000 years ago, however *D. melanogaster* may have colonized Europe and America earlier than *D.*

simulans (David and Capy 1988; Lachaise et al. 1988; Baudry et al. 2004; Li and Stephan 2006; Thornton

and Andolfatto 2006). *Drosophila simulans* is thought to have colonized the Americas in the last 100

years, creating a large and more recent population bottleneck than in *D. melanogaster* (Sturtevant 1920;

Signor *et al.* 2017c).

D. melanogaster and *D. simulans* are cosmopolitan species and human commensals, however

while *D. melanogaster* is commonly found inside houses, breweries, and wineries, *D. simulans* is more

often observed in orchards and parks (though their niches overlap significantly and they are often found

on the same patches) (David & Bocquet 1976; Parsons & King 1977; McKenzie & McKechnie 1979;

Parsons & Spence 1981; Dickinson *et al.* 1984; Gibson & Wilks 1988; Thomson *et al.* 1991; Chakir *et al.*

1993; Mercot *et al.* 1994; Joshi & Thompson 1997). *D. melanogaster* is considerably more ethanol

tolerant than *D. simulans*, and is found regularly feeding upon and ovipositing in resources with ethanol

concentrations greater than 8% (Hoffmann & McKechnie 1991; Fry 2014; Zhu & Fry 2015). At

concentrations of 4% ethanol *D. simulans* shows reductions in survivorship and increased development

time relative to *D. melanogaster* (Joshi & Thompson 1997). However, at low concentrations of ethanol

(0.5 - 3.0%) adult *D. simulans* show increases in longevity (less so than adult *D. melanogaster*, which

increases longevity from 0.5 - 9.0%) (Parsons *et al.* 1979). During multi-generation exposure to ethanol-

rich substrates in the lab, *D. simulans* evolves tolerance (Joshi & Thompson 1997; Lefèvre *et al.* 2012). It

has been demonstrated that ethanol tolerance comes as a trade-off that reduces the efficacy of processing

other biochemical targets, thus it is possible that in *D. simulans* this trade-off is under different selective

pressure (Chakraborty & Fry 2016). However, ethanol is by no means a novel resource for *D. simulans.*

In general genetic adaptation and phenotypic plasticity are considered as alternative adaptive

strategies in response to the environment (Schlichting & Pigliucci, 1998). For a variety of traits *D.*

simulans has been observed to be 'plastic' while *D. melanogaster* is described as being locally adapted

(David & Bocquet 1975; Hyytia *et al.* 1985; Watada *et al.* 1986; Singh 1989) . For example, for

temperature tolerance *D. simulans* has been shown to exhibit less difference between populations and to

produce 'optimal' phenotypes under a wider range of environments (consistent with adaptive plasticity)

(Austin & Moehring 2013). This includes less differentiation between populations, in that populations

from different temperatures show similar levels of plasticity across temperatures. Population genetic

surveys generally corroborate these phenotypic observations, where *D. melanogaster* also shows greater

differences between populations, and more evidence of clinal variation than in *D. simulans* (Singh 1989;

Weeks *et al.* 2002; Hoffmann & Weeks 2006; Schmidt & Paaby 2008; Machado *et al.* 2015; Sedghifar *et*

al. 2016; Bergland *et al.* 2016; Signor *et al.* 2017c). However, *D. simulans* has generally been observed to

harbor more within population diversity than *D. melanogaster* (Machado *et al.* 2015; Sedghifar *et al.*

2016; Signor *et al.* 2017c).

Gene expression is both affected by the environment and mediates an organisms response to the

environment. It has been shown to be plastic in response to many environmental variables such as heat

stress, ethanol, and the presence of predators (Hodgins-Davis & Townsend 2009; Levine *et al*. 2011;

Yampolsky *et al*. 2012; Hodgins-Davis *et al*. 2012). While many studies have quantified the effect of the

environment on gene expression, far fewer have performed these experiments in multiple genotypes

within a common garden to obtain the interaction between genotype and the environment for gene

expression (DeBiasse & Kelly 2015). Indeed, most work on gene expression plasticity has not quantified

the interaction between genotype and the environment, and has found inconsistent results with regard to

theoretical expectations for the maintenance of plasticity (Cheviron *et al*. 2008; McCairns & Bernatchez

2009; Hodgins-Davis *et al*. 2012; DeBiasse & Kelly 2015; Heckel *et al*. 2016; Mathur & Schmidt 2016).

Furthermore, many plastic responses are time dependent, and if performed in a common garden without

sampling across time points may not detect early plastic responses that usher in 'maintenance'

phenotypes, or dynamic responses to the environment (Aubin-Horth & Renn 2009; Lewis *et al*. 2014). In

addition, alternative splicing has been shown to respond to changes in the environment, and it is thought

to diverge more rapidly than gene expression between lineages (Barbosa-Morais *et al*. 2012; Merkin *et al*.

2012; Gueroussov & Gonatopoulos-Pournatzis 2015; Jakšić & Schlötterer 2016; Wang *et al*. 2017; Pajoro

et al. 2017; Singh *et al*. 2017; Calixto *et al*. 2018). As such it may be an important component of

plasticity, necessary for adaptation to rapidly changing and heterogeneous environments (Marais *et al*.

2013; Preußner *et al*. 2017; Price *et al*. 2018; Calixto *et al*. 2018). Furthermore, it has been previously

implicated in the response to ethanol, and in *D. melanogaster* may be a more important component of the

response than gene expression changes (Oomizu *et al*. 2003; Newton *et al*. 2004; Pietrzykowski *et al*.

2008; Sasabe & Ishiura 2010; Hemby 2012; Zaharieva *et al*. 2012; Robinson & Atkinson 2013; Signor &

Nuzhdin 2018). However, alternative splicing is rarely accounted for in studies of plasticity.

140   Overall, we compare plasticity in response to ethanol, including genetic variation for plasticity

over time and between genotypes, among *Drosophila* species with different patterns of ethanol utilization.

We measure plasticity and variation in expression and splicing in response to ethanol treatment, and we

observe that in *D. melanogaster*, which regularly exploits ethanol-rich substrates, there is essentially no

genetic variation for plasticity within the tested population, while *D. simulans* exhibits an order of

magnitude larger effect of genotype by environment compared to other components of variance. This

suggests the following scenario: for *D. simulans* high ethanol concentrations are essentially a novel

stressful environment that reveals cryptic variation for phenotypic plasticity, while *D. melanogaster* does

not experience high stress under 15% ethanol and cryptic variation is not revealed. Furthermore, *D.*

melanogaster does not experience high stress as it is adapted to ethanol and genetic variation has been

removed in favor of an optimal, and reduced, plastic response

<u>Methods</u>

Fly lines

Male flies for *D. melanogaster* originated from six genotypes collected from an orchard in Winters,

California in 1998 and made isogenic by 40 generations of full sibling inbreeding (Yang & Nuzhdin

2003; Campo *et al*. 2013; Signor *et al*. 2017a; b). Male flies for *D. simulans* came from six genotypes

collected from the Zuma Organic Orchard in the winter of 2012 and made isogenic by 15 generations of

full sib inbreeding (Signor *et al*. 2017b; c). Residual heterozygosity is similar between these lines of *D.*

melanogaster and *D. simulans* (Campo *et al*. 2013; Signor *et al*. 2017c). In natural conditions flies will

not by homozygous at all loci, thus each inbred genotype was crossed to a white eyed 'tester' strain to

create the F1 flies used in the ethanol exposure assays (*D. melanogaster*, w^{1118}, Bloomington stock number

3605; *D. simulans*, w^{501}, Cornell species stock number 14021-0251.011). This design allows us to replicate

observations of gene expression because the flies are identical twins, while also maintaining

heterozygosity such that they more closely resemble wild type flies. Rearing occurred on a standard

medium at 25 °C with a 12-h light/12-h dark cycle. Several measures were taken to standardize offspring

quality: F1 flies were produced from females of the same age, these females were held at the same density

(10 individuals per sex/vial), males used for ethanol exposure assays were collected as virgins and reared

in single sex vials, and males used for assays were held at a standard density (24-30 per vial).

During the ethanol exposure assay a female was included as a stimulus, but was not collected for

RNA-seq (Signor *et al*. 2017a; b). The females were from yw^{1} mutants for both *D. simulans* and *D.*

melanogaster (*D. melanogaster*, Bloomington stock number 1495, *D. simulans*, Cornell species stock

number 14021-0251.013). Quality of the females was controlled in the same manner as previously

described for males, and both males and females were aged three to four days before being used in the

ethanol exposure assay.

Experiment setup

Ethanol exposure took place in a circular arena, each of which was part of a larger chamber containing 12

arenas each with a diameter of 2.54 cm (VWR cat. no. 89093-496). Prior to the assay the flies were

sedated through exposure at 4 °C for ten minutes, to avoid the confounding effects of CO_2 exposure on

behavior and gene expression. Once the flies were sedated they were placed in the arenas with a

paintbrush (two males and one female per arena) and the chambers were secured with two small pieces of

cryogenic tape. It is standard to allow organisms to recover in a new location for ten minutes prior to

beginning behavioral assays, which was a portion of the goal of this study (see Signor *et al*. 2017a; b); so

the chambers were left upside down for ten minutes while the flies regained consciousness and oriented

themselves. The bottom of each chamber contained a standard amount of either grapefruit medium or

medium in which 15% of the water has been replaced with ethanol. During ethanol administration the

flies were recorded using VideoGrabber (http://code.google.com/p/video-grabber/), and set-up of the

assays was facilitated with FlyCapture (PointGrey, Canada). In order to better standardize the

transcriptional response, these videos were also used to determine if any male/female pairs mated during

the assay or any flies were damaged during setup, and flies in those chambers were not collected for

RNA-seq.

Assays were conducted for ten, 20, or 30 minutes for three replicates of each of the two conditions.

Flies are most active during the hours following dawn, thus to standardize behavior and circadian rhythms

all assays were conducted within a two-hour window after dawn. Replicates for both species were

conducted randomly under standardized conditions (25 °C, 70% humidity). At the conclusion of the assay

the chambers were flash frozen in liquid nitrogen, allowed to freeze through, transferred to dry ice, and all

of the males were collected for RNA-seq. For both *D. melanogaster* and *D. simulans* the expectation is

that intoxication is occurring through inhalation of ethanol vapors, and evidence of the behavioral effects

in both species and the efficacy of this approach have been published previously (Signor *et al*. 2017a; b).

Sample preparation and RNA sequencing

Sample preparation has been described previously, and will be briefly summarized here (Signor & Nuzhdin

2018). Flash-frozen flies were freeze dried and ten to 12 heads were placed into a 96-tube plate (Axygen

MTS-11-C-R). mRNA purification, cDNA synthesis and library preparation were carried out by RAPiD

GENOMICS (http://rapid-genomics.com) using a robot. mRNA was purified using Dynabeads mRNA

DIRECT Micro kit (Invitrogen # 61021) with slight modifications. To fragment the RNA mRNA-beads

were resuspended in 10 uL 2X first strand buffer (Invitrogen # 18064-014), incubated at 80 C for two

minutes and placed on ice, then the supernatant was collected after five minutes on magnetic stand. First

strand synthesis was performed using standard protocols for Superscript II (Invitrogen #18640-014) and

reverse transcription (25 C 10 min, 42 C for 50 min, 70 C for 15 min, 10 C hold). Second strand synthesis

was carried out using standard protocols with DNA Pol I and incubated at 16 C for 2.5 hours. cDNA was

purified with 1.8 volume of AMPure XP following manufactures instructions (Beckman Coulter A63880).

Illumina RNAseq libraries were prepared by Rapid Genomics (http://rapid-genomics.com/home/) using

dual barcodes.   Sequencing was performed using the Illumina HiSeq 2500 as both 2×150 bp or 2×50 bp

reads. The two run lengths (and runs) were intended to provide extra coverage, and all replicates were

sequenced in both runs.

Gene expression analysis

It is common in organisms with alternative splicing for exons from different isoforms of a single gene to

overlap with one another, or be shared between all or most isoforms (Figure 1B). Short read data

fundamentally cannot resolve these exons to individual isoforms, however, one approach is to quantify each

exon separately and decompose exons overlapping between isoforms into those which are shared and

unique. When the differences between overlapping exons are less than 10 bp, there is no appreciable

amount of information loss in not decomposing overlapping exons, and this approach has been taken in the

past (Dalton *et al*. 2013; Graze *et al*. 2014; Newell *et al*. 2016; Fear *et al*. 2016). However, in many cases

the differences in exon overlap are much larger than this, so to address this issue we use a classification

scheme where reads may be assigned to exons, exonic regions, or exon fragments (Signor & Nuzhdin 2018)

(Figure 1B). Here, an *exon* does not overlap any other exons (as opposed to *exonic regions* or *fragments*),

and it may be unique if it is found in a single isoform, common if it is shared between some but not all

isoforms, or constitutive if it is found in all isoforms of a gene. For example, if a gene has one transcript,

all gene regions are unique constitutive exons (Figure 1B). If exons from different isoforms overlap, they

are grouped into an *exonic region*. Exonic regions are made up of exons that overlap from different

isoforms, thus they are not unique, but they may be common to several isoforms or constitutive and found

in all isoforms – they may also be formed of overlapping exons from neighboring genes (Figure 1B). *Exon*

fragments are classified by decomposing *exonic regions* based on the 5' and 3' positions of the exons within

the region. Thus all *exon fragments* are subregions of *exonic regions*. Exon boundaries were determined

using the *D*. *melanogaster* FlyBase 6.17 genome features file and the *D*. *simulans* 2.02 genome features

file. Alignment was performed using BWA-MEM version 0.7.15 and BED files were used to count reads

in each region and obtain the length adjusted read count (reads in region divided by the length of region),

and the APN (average per nucleotide) (Li 2015).

The APN was summed for technical replicates of the same read length then averaged between read

lengths to handle the mixture of read lengths for each sample (2×150 bp and 2×50 bp). If the APN was

greater than zero in at least half of all samples per condition it was considered detected. While we

considered several approaches to normalize coverage counts upper-quartile normalization with log-

transformation and median centering within time × treatment × genotype were selected due to better

performance of the residuals (Bullard *et al*. 2010; Dillies *et al*. 2013).

To test the significance of components of expression variation, the log APN for each exonic region was

modeled as

$$Y_{ijkl} = \mu + g_i + t_j + (gt)_{ik} + m_k + (gm)_{ik} + (tm)_{ijk} + \epsilon_{ijkl}$$

for the ith genotype (g_i), jth treatment (t_j; j = ethanol or no ethanol), kth time point (m_k; k = 10 min, 20 min, 30

min), and lth replicate. For the interaction between treatment and time point, the log APN for each exonic

region was modeled as

$$Y_{ij} = \mu + t_j + \epsilon_{ij}$$

or the ith condition (time × treatment) and jth replicate. Contrasts to compare treatments within time point

(ethanol versus no ethanol, for ten min, 20 min and 30 min) were conducted. Residuals were evaluated for

conformation with normality assumptions, and assumptions were met in excess of 95% of the models.

To evaluate whether there was evidence for splicing differences among times or treatments, exons

and exonic regions for each gene and for each sample were ranked and the most expressed region ranked

as one, the least expressed region as three and all others as two. Exon ranks for each gene were modeled as

$$Y_{ijk} = \mu + r_i + t_j + (rt)_{ij} + \epsilon_{ijkl}$$

where Y_{ijk} is the exon rank (1,2,3) of the ith exonic region of the gene, jth condition (time × treatment), and

the kth replicate; r_i is the exonic region of the gene; t_j is condition; and $(rt)_{ij}$ is the interaction between

exonic region and condition. We took this approach, rather than what may be viewed as a more traditional

GLM approach, due to a lack of normality in the distribution of model residuals. Accordingly, a non-

parametric test must be relied upon to look for changes in exon or exonic region representation between

exons of a gene and we used of a rank test to summarize changes in exon representation (Supplemental File

3). F-tests for the significance of the mean square attributed to the effect tested versus the mean square

attributed to error, or the appropriate interaction term, were used. The false discovery rate was controlled

using the Benjamini-Hochberg procedure, with a significance cutoff of $\alpha = 0.05$ (Benjamini & Hochberg

1995).

GO Analysis

When a gene in *D. simulans* had more than one ortholog in *D. melanogaster* only one ortholog was

included for the GO analysis, so as not to inflate the number of genes involved in a given process. This

does presume that orthologs will be annotated with the same GO terms, and this is generally the case. For

example, in *D. simulans* there is only *AOX4,* while in *D. melanogaster* there is *AOX3* and *AOX4,* but the

GO terms for each are the same. However, as *D. simulans* genes are generally not independently

annotated, especially those without *D. melanogaster* orthologs, if there was no *D. melanogaster* ortholog

the gene was not included in the enrichment analysis. This is a significant fraction of the overall genes

that were involved in the response to treatment, treatment by genotype, and genotype by treatment by

time, however there is no viable alternative. Lists of significant genes were tested for GO enrichment

using the PANTHER classification system (Mi *et al.* 2017). They were corrected for multiple testing and

a *p*-value of 0.01 was required for significance.

Functional class enrichment

To test for functional class enrichment multigene exonic regions were not included. Every test of

functional class enrichment compared the frequency of a given subcategory among all exons and exonic

regions detected in the dataset compared to the frequency within a significant list of exons and exonic

regions. A χ^2 test was performed in R to test the significance of the enrichment of each of these

categories.

D. melanogaster polymorphism and divergence

If the response to ethanol has been under selection in either *D. simulans* or *D. melanogaster*, the genes

which respond to ethanol exposure may show indicative population genetic signatures. To compare

polymorphism between the significant sets of genes and genome-wide averages we calculated Tajima's D

genome-wide for the source populations of *D. melanogaster* and *D. simulans*. We obtained the VCF files

from the Winters population sequenced by (Campo *et al.* 2013), which includes all six of the genotypes

assayed here as well as 29 other inbred lines sampled from the same population at the same time. The

coordinates of the genes implicated for exons, exonic regions, and exon fragments were converted from

the current assembly coordinates (v6) to those used in the previous study (v5). For *D. simulans* we used

data previously obtained from 170 individuals from this population (Signor *et al.* 2017c). To obtain

estimates that were consistent with our dataset (which is gene regions) we extracted regions from

genome-wide VCFs that corresponded to genes as annotated in the latest assembly. Note that we are

considering gene regions here rather than trying to include regulatory regions for a number of reasons: 1)

Gene regions includes introns, which will include some regulatory regions and splice sites. 2) The

location of regulatory regions for these genes is not well established, much less so between species where

we do not know if they may have shifted locations. This approach is more conservative than including

arbitrary amounts of putative upstream regulatory regions. 3) If there has been recent selection on a

regulatory region it may still be linked to the gene region and show the same differences in polymorphism

frequency. We excluded regions of reduced recombination near the centromeres, either 1 MB or more if

there were significant reductions in diversity for a broader region, calculated as extended negative

Tajima's D or values of $\pi < \frac{1}{2}$ the chromosomal average (Sedghifar *et al.* 2016; Signor *et al.* 2017c).

SNPs within these coordinates were separated using bedtools intersect (v2.26.0). We excluded regions

from the fourth chromosome and unassembled scaffolds for the analysis of population genetic differences.

In both species this is a trivial number of significant regions. Alternate SNPs that were present in > 99%

of the mapping population were excluded, as were SNPs with more than 10% missing data. Tajima's D

was calculated in windows of 1 KB using VCFtools v1.12a. Windows of Tajima's D that overhung the

ends of genes were included in the analysis.

We calculated per-gene D_{XY} for the significant exons, exonic regions, and genes with significant

changes in rank abundance, to determine if there were any unusual patterns of divergence for these

subsets. To calculate divergence comparable regions of the genome need to be identified, and there has

been considerable evolution of transcription start sites between *D. melanogaster* and *D. simulans* (Main *et*

al. 2013). Comparing regions annotated as the gene regions in each species introduces large and

unexpected gaps in the start of the alignments. While the evolution of transcription start sites is of

interest, if unaccounted for it will artificially inflate estimates of D_{XY}. Thus for genes implicated in

differences in *D. melanogaster* the regions annotated as genes were blasted using ncbi blastn (v2.4.0) to

the *D. simulans* assembly and then back to the *D. melanogaster* assembly and used for divergence

statistics, and vice versa for *D. simulans* genes. This means that for different genes small non-coding

regions may be included depending upon the direction of evolution of transcription start sites, or small

portions of coding regions may be excluded. Furthermore, for genes from *D. simulans* with more than one

ortholog in *D. melanogaster* the top blast hit was used, and while there may have been orthologous

sequence in *D. melanogaster* (or *D. simulans*) if there was no annotated ortholog it was excluded from the

analysis. For example, some of the non-protein coding genes from *D. simulans* have very conserved

BLAST hits in *D. melanogaster* that are not annotated as containing genes. The sequences were aligned

using the R package DECIPHER (Wright 2016) and if necessary they were reverse complimented using

Biostrings (Pages *et al.* 2017, for example, in the region on 3R in *D. melanogaster* that is an inversion).

Following alignment the *D. simulans* and *D. melanogaster* reads were output separately in aligned fasta

format and D_{XY} was calculated using an R script courtesy of Dr. Emily Delaney, which incorporates

commands from the R packages ape and pegas (Supplemetnal file 4, Paradis *et al.* 2004; Paradis 2010)) .

This was done for every orthologous gene in the genome from *D. melanogaster* and *D. simulans*, and

each significant subset from each component of variance, to compare distributions of divergence.

<u>Results</u>

Gene expression and isoform usage

It is difficult to decouple alternative isoform usage from gene expression, given that many exons are

shared between isoforms or overlap other exons. To infer isoforms from short read data, one must rely

upon unique junctions or regions of individual isoforms and extrapolate to shared regions. This requires

accurate isoform annotation (knowing that any given exon/junction is found in combination with other

exons/junctions) and in general can be very noisy. Accordingly, we subscribe to a simpler but more robust

approach and summarize the abundance of different exons and exonic regions separately. We detail

overall abundance of exons, exonic regions, and exon fragments, and changes in the rank abundance of

exons within a gene (Figure 1 C&D). As illustrated in Figure 1 C&D, these approaches summarize

different features of expression and splicing – for example differences in the expression level of exon

fragments from *cabut* suggest both that there are overall expression differences for the gene between

environments, and that in both *D. simulans* and *D. melanogaster* one of the alternative isoforms is more

abundant with ethanol (Figure 1C). While exons within *Drat* change their expression in response to

ethanol by time, and between environments at 30 minutes, that doesn't capture the fact that the third exon

is most abundant in ethanol while the first is most abundant without it, suggesting differences in isoform

abundance that may also belong to unannotated isoforms (Figure 1D). In the following sections we will

first summarize changes in exon, exonic region, and exon fragment abundance followed by differences in

rank abundance between environments.

Exons, exonic regions, and exon fragments

1994 exons, 406 exonic regions, and 608 exon fragments changed their expression in response to

genotype in *D. simulans*, while in *D. melanogaster* 1445 exons, 631 exonic regions, and 1135 exon

fragments altered their representation (Table 1). Note that the results for *D. melanogaster* are summarized

in (Signor & Nuzhdin 2018), but are included here for comparison. 76 exons, 23 exonic regions, and 18

exon fragments changed expression in response to ethanol in *D. simulans*, compared to 15 exons, 13

exonic regions, and 21 exon fragments in *D. melanogaster* (Table 1; Signor & Nuzhdin 2018). Seven

exons and exonic regions were shared between species for treatment, *Drat*, *cabut*, *CG11741*, *CG32512*,

CG4607, *Pino*, and *sugarbabe.* For all discussion of shared genes, the particular exon or exonic region

may or may not be the same, as well as the direction or nature of the change in expression. The

complexity of this comparison is shown in Figure 1C, where in *D. melanogaster cabut* has three

annotated transcripts and four exon fragments (one exonic region), while in *D. simulans cabut* there are

two transcripts, one exon, and two exon fragments (one exonic region). In addition, in *D. melanogaster*

cabut is nested within the gene *ush*, while in *D. simulans* it is not nested. In *D. melanogaster* an area

annotated as an intron in *D. simulans* is unique to its third isoform, and this is more frequent with ethanol

than without; in *D. simulans* there is one unique exon fragment belonging to one isoform which is more

frequent with ethanol than without, suggesting that isoform is more common in ethanol environments

(Figure 1C). In *D. melanogaster* the exon fragment which is unique in *D. simulans* is not unique, and only

increased frequency of its unique third isoform can be inferred (Figure 1C). The interaction between

genotype and treatment was significant for 387 exons, 98 exonic regions, and 81 exon fragments in *D.*

simulans and three exons, no exonic regions, and no fragments in *D. melanogaster* (Table 1; Figure 2

A&B, Signor & Nuzhdin 2018). No fusions or fragments were shared between *D. simulans* and *D.*

melanogaster.

24 exons, seven exonic regions, and six exon fragments were significantly different in response to

the interaction between ethanol and time in *D. simulans*, compared to 22 exons, eight exonic regions, and

12 fragments in *D. melanogaster* (Table 1; Signor & Nuzhdin 2018). Three genes were shared between

exons and exonic regions for these species, *Drat*, *cabut*, and *CG43366*. In *D. simulans* 1158 exons, 299

exonic regions, and 225 exon fragments were significantly different for the interaction between genotype,

ethanol, and time, while in *D. melanogaster* no exons, two exonic regions, and four exon fragments were

significantly different (Table 1; Figure 2 A&B, Signor & Nuzhdin 2018). No exon fusions or fragments

were shared between species for this component of variance. At ten minutes nine exons, three exonic

regions, and four exon fragments are significantly different between treatments, compared to two exons,

one exonic region, and three exon fragments in *D. melanogaster* (Table 1; Figure 2A). *cabut* is shared

between exons and exonic regions in *D. melanogaster* and *D. simulans*. At 20 minutes three exons, one

exonic region, and two exon fragments were different in *D. simulans*, two of which belong to *cabut* in

both exons/exonic regions and exon fragments. In *D. melanogaster* one exon, four exonic regions, and

five exonic fragments are different at this time point, and the gene *cabut* is shared between species for

both categories. At 30 minutes 29 exons, four exonic regions, and six fragments in *D. simulans* were

different between treatments, all of which have *D. melanogaster* orthologs (Table 1). In *D. melanogaster*

46 exons, 24 exonic regions, and 20 exon fragments were significantly different between treatments at 30

minutes (Signor & Nuzhdin 2018). Among exons and exonic regions *Drat*, *CG32103*, *sugarbabe*, *cabut*,

Pinocchio, *CG32512*, and *CG4607* are shared between species. Among exon fragments *cabut* and

Pinocchio are shared.

Shared and unique exons, exonic regions, and fragments

Comparisons between *D. simulans* and *D. melanogaster* are shown in Figure 2C for the three components

of variance in which enough genes are implicated in *D. melanogaster* to make the frequency of different

categories meaningful. When an exon is unique/constitutive it belongs to the only annotated transcript for

that gene, and as such no exonic regions are unique/constitutive. However, the combined counts are

shown for the common and constitutive categories. However, in *D. melanogaster* constitutive exons and

exonic regions are much more common than in *D. simulans*, where unique/constitutive exons are

overwhelmingly implicated. This could potentially be explained by differences in annotation, for example

if fewer genes have multiple transcripts annotated in *D. simulans*. While this does appear to be the case

(Supplemental Figure 1), it is unclear if it is enough to explain the discrepancy between species. It is also

possible that in response to the environment genes with alternative splicing are more important in *D.*

melanogaster compared to *D. simulans*. The proportion of genes in each category in *D. simulans* is

consistent between components of variance, including ethanol by genotype and ethanol by genotype by

time, again suggesting the possibility that differences in annotation are responsible. If annotation

differences are responsible this does not affect the overall results, for example the number of exons

implicated, but may make comparison between species for the number of unique versus constitutive

differences not meaningful. It also underlines the importance of not relying upon isoform annotation

when trying to understand differences in expression and alternative splicing.

GO enrichment analysis

We report here only the results of enrichment for exons and exonic regions (Table 2). In *D. simulans* the

response to treatment, treatment by time, and the three time points were not enriched for any GO terms.

Much of the lack of enrichment is likely due to annotation issues – for example 9% of the exons

implicated in treatment over time could not be resolved to a single gene, and of the remaining genes 14%

do not have a *D. melanogaster* ortholog. Of those with annotated orthologs, 20% do not have any gene

ontology terms associated with them. A summary of significant GO terms for the remaining components

of variance is shown in Table 2. In *D. melanogaster* exons and exonic regions were not significantly

enriched for any category of genes. In general, the small number of genes implicated for many categories

precludes any conclusion of enrichment.

Changes in rank abundance

In response to treatment 54 genes showed changes in rank abundance, and 94 genes change the rank

abundance of their exons for the interaction between treatment and time. In *D. melanogaster* 71 genes

changed the rank abundance of their exons in response to ethanol and 145 changed for treatment by time

(Signor & Nuzhdin 2018). No genes were shared between species. For the interaction between treatment

and time *lola*, *Mhc*, and *Prm* were shared between species. Figure 1D illustrates an example of a change

in rank abundance for the gene *Drat* in *D. simulans*, where with ethanol the third exon is most abundant,

and without ethanol the first exon is most abundant. While differences in exon abundance summarized

above capture some of this variation, as *Drat* is significant for some exons for ethanol by time and 30

minutes, the change in rank abundance highlights potential differences in exon inclusion between

environments.

GO enrichment analysis

Changes in rank abundance were not enriched for any GO terms in *D. simulans*, with the exception of

peroxiredoxin activity in response to ethanol, albeit slightly above the more conservative cut-off used in

our other tests ($p = .017$). Peroxiredoxin activity has been associated with protection against alcohol

induced liver damage (Bae *et al.* 2011; Chattopadhyay *et al.* 2015). In *D. melanogaster* genes implicated

in changes in rank abundance in response to ethanol by time were enriched for cellular components actin

cytoskeleton, again somewhat above our more conservative cutoff ($p = 0.018$) (Signor & Nuzhdin 2018).

In D. simulans genotype-specific reactions to the environment are abundant

In *D. melanogaster* components of variance for interaction terms have very few significant genes, with

the largest category being exons and exonic regions that respond to ethanol and that are different at 30

minutes. *D. simulans* is roughly comparable for the majority of these categories. However, many more

exons and exonic regions are significant for components of variance that interact with genotype: 1457 for

the interaction between genotype, treatment and time, and 486 for the interaction between genotype and

treatment, compared to two and three exons and exonic regions respectively in *D. melanogaster* (Table 1;

Figure 2 A&B). This suggests that in *D. simulans* interactions with genotype are a far more important

component of the response to ethanol than in *D. melanogaster*. It is also worth noting that in *D. simulans*

the response to ethanol (plasticity) is three times as large (28 exons and exonic regions in *D.*

melanogaster compared to 99), though this is many order of magnitudes less of a difference than for

genetic variation for plasticity.

Genotype-specific responses are enriched for non-coding genes in D. simulans

A large number of non-protein coding genes were implicated in gene expression changes in *D. simulans*

in these analyses, so we applied a χ^2 test to understand if our gene lists were enriched for this functional

category. The number of non-protein coding genes that were significant for ethanol, ethanol by genotype,

and ethanol by genotype by time, were more than would be expected by chance in *D. simulans* (13, $\chi =$

$49.598, p < 0.0005$; $68, \chi = 235.21, p < 2.2 \; x \; 10^{-16}$; $196, \chi = 727.17, p < 2.2 \; x \; 10^{-16}$). Other than

the pseudogenes, these are long non-coding RNAs (lncRNAs) as the shortest is annotated at 713 bp, and

the majority are over 4,000 bp (long non-coding RNAs being any non-protein coding genes over 200 nt).

The interaction between treatment and time, and the differences in expression at ten, 20, and 30 minutes

were not enriched for non-protein coding genes, nor were changes in rank abundance.

D. melanogaster is not enriched for non-protein coding genes in response to ethanol, ethanol by

time, ethanol by genotype, ethanol by genotype by time, ten or 20 minutes, exons or exonic regions

expressed only in one environment, or genes implicated in changes in rank abundance. However, at 30

minutes *D. melanogaster* is enriched for non-protein coding genes, though this concerns far fewer genes

than in *D. simulans* $(5, \chi = 12.831, p = 0.005)$.

Non-coding genes that respond to the environment in D. simulans do not have D. melanogaster orthologs

Given that a considerable number of non-protein coding genes were implicated in this analysis for *D.*

simulans, we were suspicious that this may be an annotation issue in *D. melanogaster*. Using the exons

associated with every transcript from each of these non-protein coding genes as a reference we did not

find that *D. melanogaster* reads mapped to these exons across a range of relaxed mapping parameters

allowing for mismatches and gaps (using bwa mem, from defaults to -B 2 -O 3; using both the *D.*

simulans exons or existing homologous regions in *D. melanogaster*). In several cases, such as *CG30377*,

an annotated protein coding gene in *D. melanogaster* is annotated as a pseudogene in *D. simulans*, with

no noted relationship between them. Therefore there may be a small number of cases in which an ortholog

exists but is not recognized, but in general these non-protein coding genes appear to be unique to *D.*

simulans. It has been noted previously that lncRNAs can share high sequence similarity between related

species, but be expressed in only one (Ulitsky 2016). Determining the orthology of these genes and the

dynamics of non-protein coding gene evolution on the *Drosophila* phylogeny is an interesting question

that will require additional future research.

The response to the environment is enriched for nested non-protein coding genes in D. simulans

In analyzing the non-protein coding genes identified in *D. simulans* we noted that many of them were nested in the introns of other genes. We investigated the frequency with which nested genes were implicated in our analysis, and nested non-protein coding genes, to understand the possibility that differences in expression could also be changes in intron retention in the parental genes (as 'nestedness' generally refers to exons or entire genes found within the introns of other genes). Using the criteria that an exon nested in an intron must overlap the intron by at least 80 bp or 10%, we found that in *D. melanogaster* 9.2% of exons were nested within introns, while in *D. simulans* 9.7% were nested, similar to what has been previously reported (Lee & Chang 2013). This was reflected in our data, where for both *D. melanogaster* and *D. simulans* 6.8% of exons and exonic regions were nested within introns (lower because multi-gene exons were excluded). However, among significant exons and exonic regions in *D. melanogaster* 18.6% were nested, while in *D. simulans* 33.5% were nested. This is a significant enrichment of exons that are nested within other introns, for both *D. melanogaster* ($\chi = 12.344, p < .0004$) and *D. simulans* ($\chi = 1721.1, p < 2.2 \ x \ 10^{-16}$).

The remaining question then is whether nested genes are more likely to be non-protein coding, or whether the dataset is enriched for both. Indeed, compared to the total number of nested genes that are noncoding within the dataset, the number that are significant for the response to ethanol is enriched for nested, non-protein coding genes in *D. simulans* ($\chi = 237.49, p < 2.2 \ x \ 10^{-16}$), but not in *D. melanogaster*. Overall *D. melanogaster* has more annotated non-protein coding genes (2963) than *D. simulans* (1675), and similarly more exons from noncoding genes are nested within introns (1772 *D. melanogaster*, 1066 *D. simulans*), suggesting that this pattern is not reflective of annotation issues between the two species. However, the question remains as to why nested genes are so much more likely to be non-protein coding than non-nested genes – for example in the total dataset of *D. simulans* 9.7% of exons are nested, while of the annotated non-coding genes in *D. simulans* 63% are nested. Furthermore the importance of nested non-protein coding genes for the response to the environment, or potentially delays in splicing that are specific to certain components of variance, is unclear.

Nested non-protein coding genes involved in the response to ethanol are regulated independently of their

parental gene

In *D. simulans* 328 significant nested genes (375 nested exons and exonic regions) are on the opposite

strand as their parental gene, 75% of the total significant nested genes. Of non-protein coding genes, this

bias is stronger, with 83% having the opposite strandedness as their parental gene. In *D. melanogaster* 13

significant genes are nested (16 total exons and exonic regions) and of these nine are on the opposite

strand as their parent gene (69%). Only one non-protein coding gene shares strandedness with its parental

gene, *CR44660/drp1*. It has been observed that many nested non-protein coding genes require expression

of the parental gene, share strandedness, and depend on splicing out of the parental intron for activation

(miRNAs, (Boivin *et al.* 2018). In both *D. simulans* and *D. melanogaster*, however, we do not observe

non-protein coding genes as being more likely to share strandedness with their parental gene among our

significant nested exons and exonic regions. This suggests that whatever the reason for the observed

enrichment in nested non-protein coding genes, it is likely not because the parental genes are being

expressed and the nested non-protein coding genes are being spliced out of the introns. We could not test

the *D. melanogaster* dataset for correlation between the expression of parental and nested genes as it is

too small. However, in *D. simulans* genes that were both non-coding and shared strandedness with their

parental gene had highly correlated expression differences between treatments (0.83, 14 genes). All other

categories (i.e. nested non-protein coding, opposite strand) were essentially uncorrelated (0.09-0.27). We

note that of the parental genes only three are also represented as significant in the main dataset, *sima*,

mira, and *bru-3* (significant in response to ethanol or genotype by ethanol). Thus it is unlikely that overall

the observed differences in expression are due to changes in the expression of the parental gene, as this

would likely result in differences in expression detected at both loci.

Polymorphism and divergence in D. melanogaster and D. simulans

We were interested to determine if the patterns of polymorphism and divergence in the genes implicated

in the response to ethanol suggested that they were evolving under a particular selection regime or were

unusually diverged compared to background levels of polymorphism or divergence. If the genes

implicated in expression differences in *D. melanogaster* had unusually low Tajima's *D*, for example, this

could suggest that directional selection for adaptation to ethanol is responsible for the observed

expression differences. We compare polymorphism in genes with expression differences in D. simulans

in both *D. simulans* and *D. melanogaster*, because it is possible that it is those same genes that were under

selection for adaptation to ethanol in *D. melanogaster*. This might help to clarify whether the observed

expression differences are adaptive or show evidence of adaptation. For both polymorphism and

divergence in the following sections we will compare the genome-wide background distribution of

polymorphism and divergence to differences in expression and rank abundance as a whole, rather than by

components of variance, for two reasons. Firstly, for the majority of categories other than genotype

interaction terms in *D. simulans*, the number of independent genes on each list is ~50% confounding any

attempt to separate out components of variance. Second, in *D. melanogaster* there are too few genes that

are significant for many components of variance and they cannot be considered separately.

Polymorphism

We calculated Tajima's *D* in windows of 1 kb for both all annotated gene regions in the genome and the

subset of genes implicated in differences in ethanol response in both species, and consider anything in the

top ±2.5% to be an outlier. Outliers for *D. simulans* and *D. melanogaster*, separated between the X and

the autosome, are shown in Table 3, and the percentage of outliers for each category are graphed in Figure

3A.

D. melanogaster

For changes in abundance of exons and exonic regions on the X and autosome, *D. melanogaster* is not

enriched for outliers of Tajima's *D* but there are more outliers than expected in *D. simulans*

(Autosomes: $\chi = 5.26, p <\ 0.02, \text{X: } \chi = 29.361, p <\ 0.0005$). In *D. melanogaster* these intervals

correspond to 27 genes, while in *D. simulans* they cover a total of 25, including both the X and

autosomes. Between the two lists there is considerable overlap in the location of outlier intervals, at

CG43366, AOX4, dsb, fumble, CG31875, Mical, Ada2b, forked, CG1986, and *CG42749.*

For genes implicated in changes in rank abundance in *D. melanogaster*, there is also no

significant enrichment for outliers of Tajima's *D* relative to the genome-wide frequency in *D.*

melanogaster or *D. simulans.* In *D. melanogaster* on the autosome these intervals cover 59 genes, while

in *D. simulans* they cover 56. *14-3-3ε, AcCoAs, boi, CG31522, CG34398, eff, elF4EHP, fru, grh, kay,*

lds, lola, mRpL21, osp, siz, Syp, Tep2, and *tweek* are shared in containing outlier intervals for Tajima's *D*

in both *D. simulans* and *D. melanogaster.* In *D. simulans* on the X chromosome these intervals cover

four genes, while in *D. melanogaster* they cover nine, and two are in common between the species

(*CG34417, Ten-a*).

D. simulans

Many more genes were implicated in the response to ethanol in *D. simulans*, and for exons and exonic

regions that changed in abundance there was an excess of outliers for Tajima's *D* for the autosomes and

the X chromosome (Table 3; Figure 3; Autosomes: $\chi = 9.41, p = 0.0022, \text{X}: X = 14.03, p = 0.00018$).

These intervals correspond to 428 unique genes, and many outlier intervals overlap the same gene region.

For example, every 1 kb interval that overlaps the gene *Cubilin* is an outlier, with an average of -2.41.

Polymorphisms in the human ortholog of *Cubilin*, *CUBN*, have been associated with lifetime rates of

heavy drinking (Hamidovic *et al.* 2013). It is worth noting here that the outliers that were significantly

enriched in the *D. melanogaster* dataset for *D. simulans* were all in the positive direction, while these are

both positive and negative. Given the tendency towards positive Tajima's *D* in this population of *D.*

simulans, negative values of Tajima's *D* may be a more suggestive measure of selection, either purifying

or directional (Signor *et al.* 2017c). In *D. melanogaster* the genes that were implicated in changes in the

expression of exons and exonic in response to ethanol in *D. simulans* were less enriched for outliers on

the X chromosome than expected, and not enriched on the autosomes (X: $\chi = 6.38, p = .0012$).

For genes implicated in changes in rank abundance in *D. simulans* there was a significant

enrichment compared to genome-wide frequency for both the autosomes and the X (Autosomes: $\chi =$

$19.78, p < 8.7 \ x \ 10^{-6}$, X: $\chi = 458.73, p < 2.2 \ x \ 10^{-16}$). This includes intervals in 44 unique genes

including 16 intervals in the gene *Mondo*. In *D. melanogaster* this did not constitute an enrichment

relative to genome-wide. Outlier intervals that occur within shared genes between the two species include

CG7881, *lola*, *mbf1* (a stress response gene), *Mhc*, *PI31*, and *Pka-1* (which has been implicated in the

response to ethanol previously (Chen *et al.* 2008)). Note that for changes in the abundance of exons and

exonic regions in *D. simulans*, if regions of low recombination are excluded from the candidate regions

the enrichment is no longer significant, but there is no change among the other tests.

Divergence

We calculated D_{XY} for orthologous genes on the autosomes (10,435 genes) and the X (1,943 genes)

between *D. melanogaster* and *D. simulans* (Figure 3B). D_{XY} was similar to that previously reported, with

an average of 0.52 on the autosomes and 0.55 on the X (compared to 0.048 on the autosomes, 0.054 on

the X (Nolte *et al.* 2013)). For the subsets of genes implicated in expression differences the mean was

0.047 on the autosomes and 0.057 on the X for expression differences in *D. melanogaster* and 0.053 on

the autosomes and 0.06 on the X in *D. simulans*. The genes implicated in changes in rank abundance in *D.*

melanogaster had a mean D_{XY} of 0.047 on the autosomes and 0.05 on the X. In *D. simulans* this was

0.049 on the autosomes and 0.059 on the X. To determine if the distribution of D_{XY} for any of the

significant subsets of genes differed compared to the genome-wide distribution we used a Kolmogorov-

Smirnov two-sample test, separately for genes on the X and the autosomes. In *D. melanogaster* exons and

exonic regions that changed abundance in response to ethanol were not significantly different on the

autosomes or the X. However, exons and exonic regions with significant differences in rank abundance

did differ in distribution on the autosomes, but not the X (Autosomes: D = 0.14, *p* = 0.002). The genes

implicated in expression differences on the autosomes in *D. simulans* differed significantly from the

genome-wide distribution, but not on the X (Autosomes: D = 0.061, p = 0.005). However, those

implicated in changes in rank abundance did not, on the autosomes or the X. A difference in distribution

does not indicate the direction of effect, and in fact in *D. melanogaster* the mean D_{XY} for changes in rank

abundance is lower than genome-wide, while for expression differences in *D. simulans* it is slightly

higher. This suggests that the genes implicated in changes in rank abundance in response to ethanol in *D.*

melanogaster are somewhat more conserved than expected based on genome-wide distributions, while

those implicated in expression differences in *D. simulans* are slightly less conserved. Note that because

we can only include genes with orthologs here this does not include any of the abundant non-protein

coding genes implicated in *D. simulans*, likely to be among the more rapidly evolving genes.

Population genetic patterns – or a lack thereof- among genes that respond to ethanol are difficult

to interpret, in part because plastic responses may be due to a single regulatory change at an upstream

gene. As the number of regulatory (or coding) changes involved in the plastic response are not known, it

is difficult to interpret a lack of enrichment, as it is possible, for example, that only one of the genes

whose expression changed could be under selection. Furthermore, it is not known whether the response to

ethanol is adaptive – while it is presumed to be so in *D. melanogaster* due to its relative fitness on ethanol

compared to *D. simulans*, functional links between gene expression differences and the phenotype were

not established.

Discussion

In this study both *D. melanogaster* and *D. simulans* have a plastic response to the environment, but *D.*

melanogaster is lacking in genetic variation for plasticity. Given the observed patterns, it appears most

likely that for *D. simulans* 15% ethanol is a novel environment, and that the marked G x E is a passive

stress response. This maladaptive plasticity would allow for the expression of previously cryptic variation

that had accumulated in the absence of selection. In *D. melanogaster* this environment may not be

stressful or novel, resulting in past selection removing genetic variation for plasticity that in *D. simulans*

was allowed to neutrally accumulate. In this sense *D. melanogaster* may be more 'locally adapted', as it

has evolved a reduced plastic response compared to *D. simulans*. However, it could also be that because

D. melanogaster colonized the Americas a few thousand years prior to *D. simulans*, *D. simulans* has not

yet reached an optimal level of plasticity (a nonequilibrium situation). Also, the frequency with which

they encounter ethanol rich environments may vary for *D. simulans* and *D. melanogaster*, resulting in

different selection pressures for developing the optimal level of plasticity – particularly if there is a trade-

off with traits that are beneficial to the semi-domestic habitat of *D. simulans*. Trade-offs among

genetically correlated traits may mean that a genotype with optimal phenotypic plasticity in one

environment is constrained from evolving the optimal phenotype in another environment, resulting in the

maintenance of variation for phenotypic plasticity.

Joshi and Thompson (1997) found that in *D. simulans* there was greater between family variation

for the phenotypic response to ethanol substrate prior to selection for tolerance to ethanol. Compared to

D. melanogaster and control populations, after selection in an ethanol environment the greatest change

was a reduction in variation for plasticity between *D. simulans* families. Joshi and Thompson (1997)

quantified phenotypes such as development time, which are not easily generalizable to expression

patterns, but it is suggestive. However, in studies of other phenotypes in presumably adapted and non-

adapted populations of *D. melanogaster*, no increase in genotype by environment response were observed

in non-adapted populations (Heckel *et al.* 2016). This could be due to differences in the degree of stress or

novelty of different environments, or it is possible that the response in *D. simulans* is due to something

other than stress induced maladaptive plasticity.

663   The abundance of lncRNAs which are involved in genotype by environment interactions

in *D. simulans* is perhaps not surprising, given that they have been implicated previously in the response

to stress (Valadkhan & Valencia-Hipólito 2016). It is also possible they are more frequently involved

because lncRNAs are often less conserved, as there is no requirement for the maintenance of ORFs and

codon synonymy (Chodroff *et al.* 2010; Ulitsky *et al.* 2011; Quinn *et al.* 2016; Ulitsky 2016). It has been

observed previously that transcription evolves more quickly than sequences, and lncRNA are commonly

homologous to non-transcribed sequences in other species, however these species are typically more

diverged than *D. melanogaster* and *D. simulans* (Main *et al.* 2013; Ulitsky 2016). If neutral variation was

allowed to accumulate without selection, and then uncovered in a stressful environment, preferential

accumulation within less constrained sequences would be expected. *D. melanogaster* is also enriched for

lncRNAs for the response to ethanol at 30 minutes, which could be explained by increasing ethanol stress

over time. While it is possible that lncRNAs are indicative of stress-induced maladaptive plasticity this

cannot be separated from a more general involvement in the stress response, which would not necessarily

occur as a result of maladaptive plasticity.

Why the lncRNAs are preferentially nested in *D. simulans* is less easily envisioned, though four

potential scenarios for an increase in nested lncRNAs are depicted in Figure 4. The simplest explanation

is only that there has been a change in the expression of nested lncRNAs, and perhaps nested lncRNAs

are more commonly involved in less essential processes than other lncRNAs and are therefore less

constrained. It is also possible that ethanol causes a change in the intron stability of the spliced transcript,

causing increased (or decreased) detection of the nested lncRNA (Figure 4). It is also possible that a

change in the processing of the parental gene occurred, causing a change in the number of reads mapping

to unspliced introns, or that the parental gene simply changed expression. These latter two explanations

would predict correlation between the parental gene expression and nested gene expression, which was

only observed for the very small fraction of lncRNAs that shared strandedness with their parental gene.

Thus it is more likely that either nested lncRNAs are less constrained, or ethanol alters the stability of

introns during or after the process of splicing.

The population genetic patterns observed in these two species are not easily interpreted. *D.*

melanogaster does not show any increase in outliers for Tajima's *D* in any direction for either changes in

rank abundance or expression (with the exception of fewer than expected on the X chromosome for *D.*

simulans genes implicated in expression differences). *D. simulans* is an outlier in the positive direction for

D. melanogaster genes implicated in differential expression, which could indicate that the genes involved

in expression differences in *D. melanogaster* are subject to relaxed purifying or balancing selection in *D.*

simulans. As these are all in the positive direction, it is worth noting that genome-wide *D. simulans* is

biased towards positive values of Tajima's D, likely due to recent population contraction (Signor *et al.*

2017c). While they are still more positive than expected due to background levels of Tajima's D, caution

is also warranted in interpreting these patterns as due to selection. Among genes with expression

differences and changes in rank abundance in *D. simulans*, there is an enrichment of outliers for

expression differences on the autosome and X. This is due in large part to negative Tajima's D outliers,

which given genome-wide patterns is overall more suggestive of selection, in this case directional

selection. It may be that *D. simulans* phenotypic plasticity is currently not in equilibrium, and there is

selection for an optimal phenotypic response. This can be true whether or not the observed response is

due to maladaptive plasticity – diversity of the passive stress response to ethanol implies that some

responses will be more beneficial than others, and there may be selection against the less adaptive stress

responses.

Between species divergence (D_{XY}) suggests that in *D. melanogaster* the genes involved in

changes in rank abundance are less diverged than expected compared to background levels of divergence.

In *D. simulans* the genes implicated in expression differences for exons and exonic regions are more

diverged than expected based on background levels of divergence, which combined with being outliers

for largely negative Tajima's D could indicate that they have been important for adaptation in *D.*

simulans. However, given that the number of nucleotide differences involved in the response to the

environment is unknown – for example all the observed patterns could be due to a single *trans* variant, it

is difficult to interpret the results of Tajima's D and D_{XY} in terms of selection.

Inferring that gene expression differences are adaptive or non-adaptive remains a major challenge

in the study of gene expression reaction norms, given the lack of direct correlation between gene

expression phenotypes and organismal phenotypes. However, the patterns observed in *D. simulans* do

suggest maladaptive plasticity in response to ethanol exposure. In this scenario abundant genotype by

environment interactions are expected to have accumulated neutrally and become uncovered in response

to environment stress. In contrast, in *D. melanogaster* this ethanol environment is not novel and

maladaptive plasticity has been selected out in favor of an adaptive phenotypic response. lncRNAs are

preferentially differentially expressed in *D. simulans* in response to ethanol either because they are less

constrained and can accumulate more neutral variation, or because they are involved in the general stress

response. It is also possible that environmental heterogeneity has caused *D. simulans* to maintain balanced

polymorphisms for plasticity in a way that has not occurred in *D. melanogaster*, though we believe there

is less evidence in favor of this interpretation. In the future comparing African, non-ethanol adapted

populations of *D. melanogaster* to cosmopolitan populations may be a way of discerning between these

hypotheses.

Acknowledgements

The authors would like to thank Jeremy Newman and Lauren McIntyre for contributions to the manuscript.

The authors would like to thank our undergraduate researchers for assistance in producing this data: N.

Shadman, V. Paterson, Z. Polonus, K. Cortez, L. Hassanzadeh, L. Cline, A. Khokhar, E. Lee, K.L. Yee, M.

Ling, S. Sarva, O. Akintonwa, A. Gupta, R. Manson, P. Hassanzadeh, and K. Kavoussi. Lastly, the authors

would like to thank B. & C. Emery for thoughtful commentary.

Agrawal A, Hinrichs AL, Dunn G *et al.* (2008) Linkage scan for quantitative traits identifies new regions
of interest for substance dependence in the Collaborative Study on the Genetics of Alcoholism
(COGA) sample. *Drug and Alcohol Dependence*, **93**, 12–20.
Aubin-Horth N, Renn SCP (2009) Genomic reaction norms: using integrative biology to understand
molecular mechanisms of phenotypic plasticity. *Molecular Ecology*, **18**, 3763–3780.
Austin CJ, Moehring AJ (2013) Optimal temperature range of a plastic species, *Drosophila simulans* (S
Plaistow, Ed,). *Journal of Animal Ecology*, **82**, 663–672.
Bae SH, Sung SH, Cho EJ *et al.* (2011) Concerted action of sulfiredoxin and peroxiredoxin I protects
against alcohol-induced oxidative injury in mouse liver. *Hepatology*, **53**, 945–953.
Barbosa-Morais NL, Irimia M, Pan Q *et al.* (2012) The evolutionary landscape of alternative splicing in
vertebrate species. *Science*, **338**, 1587–1593.
Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach
to multiple testing. *Journal of the Royal Statistical Society, Series B*, **57**, 289–300.
Bergland AO, Tobler R, González J, Schmidt P, Petrov D (2016) Secondary contact and local adaptation
contribute to genome-wide patterns of clinal variation in *Drosophila melanogaster*. *Molecular
Ecology*, **25**, 1157–1174.
Boivin V, Deschamps-Francoeur G, Scott MS (2018) Protein coding genes as hosts for noncoding RNA
expression. *Seminars in Cell & Developmental Biology*, **75**, 3–12.
Bullard JH, Purdom E, Hansen KD, Dudoit S (2010) Evaluation of statistical methods for normalization
and differential expression in mRNA-Seq experiments. *BMC Bioinformatics*, **11**, 94–13.

Calixto CPG, Guo W, James AB *et al.* (2018) Rapid and dynamic alternative splicing impacts the Arabidopsis cold response transcriptome. *The Plant Cell*, tpc.00177.2018–45.

Campo D, Lehmann K, Fjeldsted C *et al.* (2013) Whole-genome sequencing of two North American *Drosophila melanogaster* populations reveals genetic differentiation and positive selection. *Molecular Ecology*, **22**, 5084–5097.

Chakir M, Peridy O, Capy P, Pla E, David JR (1993) Adaptation to alcoholic fermentation in *Drosophila*: A parallel selection imposed by environmental ethanol and acetic acid. *Proceedings of the National Academy of Sciences*, **90**, 3621–3625.

Chakraborty M, Fry JD (2016) Evidence that environmental heterogeneity maintains a detoxifying enzyme polymorphism in *Drosophila melanogaster*. *Current Biology*, **26**, 219–223.

Chattopadhyay A, Pinkaew D, Doan HQ *et al.* (2015) Fortilin potentiates the peroxidase activity of Peroxiredoxin-1 and protects against alcohol-induced liver damage in mice. *Scientific Reports*, 1–16.

Chen J, Zhang Y, Shen P (2008) A protein kinase C activity localized to neuropeptide Y–like neurons mediates ethanol intoxication in *Drosophila melanogaster*. *Neuroscience*, **156**, 42–47.

Cheviron ZA, Whitehead A, Brumfield RT (2008) Transcriptomic variation and plasticity in rufous-collared sparrows (*Zonotrichia capensis*) along an altitudinal gradient. *Molecular Ecology*, **17**, 4556–4569.

Chodroff RA, Goodstadt L, Sirey TM *et al.* (2010) Long noncoding RNA genes: conservation of sequence and brain expression among diverse amniotes. *Genome Biology*, **11**, R72.

Dalton JE, Fear JM, Knott S *et al.* (2013) Male-specific Fruitless isoforms have different regulatory roles conferred by distinct zinc finger DNA binding domains. *BMC Genomics*, **14**, 659.

David J, Bocquet C (1976) Compared toxicities of different alcohols for two *Drosophila* sibling species: *D. melanogaster* and *D. simulans*. *Comparative biochemistry and physiology. C: Comparative pharmacology*, **54**, 71–74.

David JR, Bocquet C (1975) Similarities and differences in latitudinal adaptation of two Drosophila sibling species. *Nature*, **257**, 588–590.

Davies MN, Krause L, Bell JT *et al.* (2014) Hypermethylation in the ZBTB20 gene is associated with major depressive disorder. *Genome Biology*, **15**, R56–12.

DeBiasse MB, Kelly MW (2015) Plastic and evolved responses to global Change: What can we learn from comparative transcriptomics? *The Journal of Heredity*, **107**, 71–81.

Dickinson WJ, Rowan RG, Brennan MD (1984) Regulatory gene evolution: adaptive differences in expression of *alcohol dehydrogenase* in *Drosophila melanogaster* and *Drosophila simulans*. *Heredity*, **52 (Pt 2)**, 215–225.

Dillies MA, Rau A, Aubert J *et al.* (2013) A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. *Briefings in Bioinformatics*, **14**, 671–683.

Fear JM, León-Novelo LG, Morse AM *et al.* (2016) Buffering of genetic regulatory networks in *Drosophila melanogaster*. *Genetics*, **203**, 1177–1190.

Fry JD (2014) Mechanisms of naturally evolved ethanol resistance in *Drosophila melanogaster*. *Journal of Experimental Biology*, **217**, 3996–4003.

Ghalambor CK, Hoke KL, Ruell EW *et al.* (2015) Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature*, **525**, 372–375.

Ghalambor CK, McKay JK, Carroll SP, Reznick DN (2007) Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*, **21**, 394–407.

Gibson JB, Wilks AV (1988) The alcohol dehydrogenase polymorphism of *Drosophila melanogaster* in relation to environmental ethanol, ethanol tolerance and alcohol dehydrogenase activity. *Heredity*, **60 (Pt 3)**, 403–414.

Graze RM, McIntyre LM, Morse AM *et al.* (2014) What the X has to do with it: Differences in regulatory variability between the sexes in *Drosophila simulans*. *Genome Biology and Evolution*, **6**, 818–829.

Gueroussov S, Gonatopoulos-Pournatzis T (2015) An alternative splicing event amplifies evolutionary differences between vertebrates. *Science,* **349**, 868-73.

Guntrip J, Sibly RM (1998) Phenotypic plasticity, genotype-by-environment interaction and the analysis
of generalism and specialization in *Callosobruchus maculatus*. *Heredity*, **81**, 198–204.
Hamidovic A, Goodloe RJ, Young TR *et al.* (2013) Genetic analysis of a population heavy drinking
phenotype identifies risk variants in whites. *Journal of Clinical Psychopharmacology*, **33**, 206–210.
Heckel von K, Stephan W, Hutter S (2016) Canalization of gene expression is a major signature of
regulatory cold adaptation in temperate *Drosophila melanogaster*. *BMC Genomics*, 1–14.
Hemby SE (2012) Alternative splicing of AMPA subunits in prefrontal cortical fields of cynomolgus
monkeys following chronic ethanol self-administration. *Frontiers in Psychiatry*, **3**, 72.
Hodgins-Davis A, Townsend JP (2009) Evolving gene expression: from G to E to G×E. *Trends in*
Ecology & Evolution, **24**, 649–658.
Hodgins-Davis A, Adomas AB, Warringer J, Townsend JP (2012) Abundant gene-by-environment
interactions in gene expression reaction norms to copper within *Saccharomyces cerevisiae*. *Genome*
Biology and Evolution, **4**, 1061–1079.
Hoffmann AA, McKechnie SW (1991) Heritable variation in resource utilization and response in a
winery population of *Drosophila melanogaster*. **45**, 1000.
Hoffmann AA, Weeks AR (2006) Climatic selection on genes and traits after a 100 year-old invasion: a
critical look at the temperate-tropical clines in *Drosophila melanogaster* from eastern Australia.
Genetica, **129**, 133–147.
Huang Y, Agrawal AF (2016) Experimental evolution of gene expression and plasticity in alternative
selective regimes. *PLoS Genetics*, **12**, e1006336.
Hyytia P, Capy P, David JR, Singh RS (1985) Enzymatic and quantitative variation in European and
African populations of *Drosophila simulans*. *Heredity*, **54**, 209–217.
Jakšić AM, Schlötterer C (2016) The interplay of temperature and genotype on patterns of alternative
splicing in *Drosophila melanogaster*. *Genetics*, **204**, 315-325.
Joshi A, Thompson JN (1997) Adaptation and specialization in a two-resource environment in
Drosophila species. **51**, 846.
Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic plasticity and
genetic assimilation. *Journal of Evolutionary Biology*, **22**, 1435–1446.
Lee CE, Kiergaard M, Gelembiuk GW, Eads BD, Posavi M (2011) Pumping ions: Rapid parallel
evolution of ionic regulation following habitat invasions. *Evolution*, **65**, 2229–2244.
Lee YCG, Chang HH (2013) The evolution and functional significance of nested gene structures in
Drosophila melanogaster. *Genome Biology and Evolution*, **5**, 1978–1985.
Lefèvre T, de Roode JC, Kacsoh BZ, Schlenke TA (2012) Defense strategies against a parasitoid wasp in
Drosophila: fight or flight? *Biology Letters*, **8**, 230–233.
Levine MT, Eckert ML, Begun DJ (2011) Whole-genome expression plasticity across tropical and
temperate *Drosophila melanogaster* populations from Eastern Australia. *Molecular Biology and*
Evolution, **28**, 249–256.
Lewis JA, Broman AT, Will J, Genetics AG, 2014 Genetic architecture of ethanol-responsive
transcriptome variation in *Saccharomyces cerevisiae* strains. *Genetics*, *198*, 369-382.
Li H (2015) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997.
Liu G, Zhou L, Zhang H *et al.* (2017) Regulation of hepatic lipogenesis by the zinc finger protein Zbtb20.
Nature Communications, **8**, 1–13.
Machado HE, Bergland AO, O'Brien KR *et al.* (2015) Comparative population genomics of latitudinal
variation in *D. simulans* and *D. melanogaster*. *Molecular Ecology*, **25**, 723-740.
Main BJ, Smith AD, Jang H, Nuzhdin SV (2013) Transcription start site evolution in *Drosophila*.
Molecular Biology and Evolution, **30**, 1966–1974.
Marais Des DL, Hernandez KM, Juenger TE (2013) Genotype-by-environment interaction and plasticity:
Exploring genomic responses of plants to the abiotic environment. *Annual Review of Ecology,*
Evolution, and Systematics, **44**, 5–29.

Mathur V, Schmidt PS (2016) Adaptive patterns of phenotypic plasticity in laboratory and field
environments in Drosophila melanogaster. *Evolution*, **71**, 465–474.
Matzkin LM (2012) Population transcriptomics of cactus host shifts in Drosophila mojavensis. *Molecular*
Ecology, **21**, 2428–2439.
McCairns RJS, Bernatchez L (2009) Adaptive divergence between freshwater and marine sticklebacks:
insights into the role of phenotypic plasticity from an integrated analysis of candidate gene
expression. *Evolution*, **64**, 1029–1047.
McKenzie JA, McKechnie SW (1979) A comparative study of resource utilization in natural populations
of *Drosophila melanogaster* and *D. simulans*. *Oecologia*, **40**, 299-309.
Mercot H, Defaye D, Capy P, Pla E, David JR (1994) Alcohol tolerance, ADH activity, and ecological
niche of *Drosophila* species. **3**, 746–757.
Merkin J, Russell C, Chen P, Burge CB (2012) Evolutionary dynamics of gene and isoform regulation in
Mammalian tissues. *Science*, **338**, 1593–1599.
Mi H, Huang X, Muruganujan A *et al.* (2017) PANTHER version 11: expanded annotation data from
Gene Ontology and Reactome pathways, and data analysis tool enhancements. *Nucleic Acids*
Research, **45**, D183–D189.
Newell NR, New FN, Dalton JE *et al.* (2016) Neurons that underlie *Drosophila melanogaster*
reproductive behaviors: detection of a large male-bias in gene expression in *fruitless*-expressing
neurons. *Genes, Genomes, Genetics*, **6**, 2455–2465.
Newton PM, Tully K, McMahon T *et al.* (2004) Chronic ethanol exposure induces an N-type calcium
channel splice variant with altered channel kinetics. *FEBS Letters*, **579**, 671–676.
Nolte V, Pandey RV, Kofler R, Schlötterer C (2013) Genome-wide patterns of natural variation reveal
strong selective sweeps and ongoing genomic conflict in *Drosophila mauritiana*. *Genome Research*,
23, 99–110.
Oomizu S, Boyadjieva N, Sarkar DK (2003) Ethanol and estradiol modulate alternative splicing of
dopamine D2 receptor messenger RNA and abolish the inhibitory action of bromocriptine on
prolactin release from the pituitary gland. *Alcoholism: Clinical and Experimental Research*, **27**, 975–
980.
Pages H, Aboyoun P, Gentleman R, Version SDRP (2017) Biostrings: Efficient manipulation of
biological strings.
Pajoro A, Severing E, Angenent GC, Immink RGH (2017) Histone H3 lysine 36 methylation affects
temperature-induced alternative splicing and flowering in plants. **18**, 102.
Paradis E (2010) pegas: an R package for population genetics with an integrated-modular approach.
Bioinformatics, **26**, 419–420.
Paradis E, Claude J, Strimmer K (2004) APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics, **20**, 289–290.
Parsons PA, King SB (1977) Ethanol: Larval discrimination between two *Drosophila* sibling species.
Experientia, **33**, 898–899.
Parsons PA, Spence GE (1981) Ethanol utilization: threshold differences among three *Drosophila*
species. *The American Naturalist*, **117**, 568–571.
Parsons PA, Stanley SM, Spence GE (1979) Environmental ethanol at low concentrations: Longevity and
development in the sibling species *Drosophila melanogaster* and *D. simulans*. *Australian Journal of*
Zoology, **27**, 747–754.
Pietrzykowski AZ, Friesen RM, Martin GE *et al.* (2008) Posttranscriptional regulation of BK channel
splice variant stability by miR-9 underlies neuroadaptation to alcohol. *Neuron*, **59**, 274–287.
Preußner M, Goldammer G, Neumann A *et al.* (2017) Body temperature cycles control rhythmic
alternative splicing in mammals. *Molecular Cell*, **67**, 433–446.e4.
Price J, Harrison MC, Hammond RL *et al.* (2018) Alternative splicing associated with phenotypic
plasticity in the bumble bee *Bombus terrestris*. *Molecular Ecology*, **27**, 1036–1043.
Quinn JJ, Zhang QC, Georgiev P *et al.* (2016) Rapid evolutionary turnover underlies conserved lncRNA-
genome interactions. *Genes & development*, **30**, 191–207.

Robinson BG, Atkinson NS (2013) Is alcoholism learned? Insights from the fruit fly. *Current Opinion in*
Neurobiology, **23**, 529–534.
Rutherford SL (2000) From genotype to phenotype: buffering mechanisms and the storage of genetic
information. *BioEssays*, **22**, 1095–1105.
Sasabe T, Ishiura S (2010) Alcoholism and alternative splicing of candidate genes. *International Journal*
of Environmental Research and Public Health, **7**, 1448–1466.
Schmidt PS, Paaby AB (2008) Reproductive diapause and life-history clines in North American
populations of *Drosophila melanogaster*. *Evolution*, **62**, 1204–1215.
Scoville AG, Pfrender ME (2010) Phenotypic plasticity facilitates recurrent rapid adaptation to introduced
predators. *Proceedings of the National Academy of Sciences*, **107**, 4260–4263.
Sedghifar A, Saelao P, Begun DJ (2016) Genomic patterns of geographic differentiation in *Drosophila*
simulans. *Genetics*, **202**, 1229-1240.
Shi J, Li W, Ding X (2018) Assessment of the association between ZBTB20 rs9841504 polymorphism
and gastric and esophageal cancer susceptibility: A meta-analysis. *The International Journal of*
Biological Markers, **32**, 96–101.
Signor S, Nuzhdin S (2018, *in press*). Dynamic changes in gene expression and alternative splicing
mediate the response to acute alcohol exposure in *Drosophila melanogaster*. *Heredity*.
Signor SA, Abbasi M, Marjoram P, Nuzhdin SV (2017a) Social effects for locomotion vary between
environments in *Drosophila melanogaster* females. *Evolution*, **71**, 1765–1775.
Signor SA, Abbasi M, Marjoram P, Nuzhdin SV (2017b) Conservation of social effects (Ψ) between two
species of *Drosophila* despite reversal of sexual dimorphism. *Ecology and Evolution*, **7**, 10031–
10041.
Signor SA, New FN, Nuzhdin S (2017c) A large panel of *Drosophila simulans* reveals an abundance of
common variants. *Genome Biology and Evolution*, **10**, 189–206.
Singh P, Börger C, More H, Sturmbauer C (2017) The role of alternative splicing and differential gene
expression in cichlid adaptive radiation. *Genome Biology and Evolution*, **9**, 2764–2781.
Singh RS (1989) Population genetics and evolution of species related to *Drosophila melanogaster*.
Annual Review of Genetics, **23**, 425–453.
Thomson MS, Jacobson JW, Laurie CC (1991) Comparison of alcohol dehydrogenase expression in
Drosophila melanogaster and *D. simulans*. *Molecular Biology and Evolution*, **8**, 31–48.
Ulitsky I (2016) Evolution to the rescue: using comparative genomics to understand long non-coding
RNAs. *Nature Reviews Genetics*, **17**, 601–614.
Ulitsky I, Shkumatava A, Jan CH, Sive H, Bartel DP (2011) Conserved function of lincRNAs in
vertebrate embryonic development despite rapid sequence evolution. *Cell*, **147**, 1537–1550.
Valadkhan S, Valencia-Hipolito A (2016) lncRNAs in stress response. *Current Topics in Microbiology*
and Immunity, **394**, 203-236.
Via S, Lande R (1985) Genotype-environment interaction and the evolution of phenotypic plasticity.
Evolution, **39**, 505–522.
Wang G, Weng L, Li M, Xiao H (2017) Response of gene expression and alternative splicing to distinct
growth environments in tomato. *International Journal of Molecular Sciences*, **18**, 475–19.
Watada M, Tobari YN, Ohba S (1986) Genetic differenciation in Japanese populations of *Drosophila*
simulans and *D. melanogaster*. *The Japanese Journal of Genetics*, **61**, 253–269.
Weeks AR, McKechnie SW, Hoffmann AA (2002) Dissecting adaptive clinal variation: markers,
inversions and size/stress associations in *Drosophila melanogaster* from a central field population.
Ecology letters, **5**, 756–763.
Wright E (2016) Using DECIPHER v2. 0 to analyze big biological sequence data in R. *The R Journal*, **8**,
352–359.
Yampolsky LY, Glazko GV, Fry JD (2012) Evolution of gene expression and expression plasticity in
long-term experimental populations of *Drosophila melanogaster* maintained under constant and
variable ethanol stress. *Molecular Ecology*, **21**, 4287–4299.

Yang H-P, Nuzhdin SV (2003) Fitness costs of *Doc* expression are insufficient to stabilize its copy
number in *Drosophila melanogaster. Molecular Biology and Evolution*, **20**, 800–804.
Zaharieva E, Chipman JK, Soller M (2012) Alternative splicing interference by xenobiotics. *Toxicology*,
296, 1–12.
Zhu J, Fry JD (2015) Preference for ethanol in feeding and oviposition in temperate and tropical
populations of *Drosophila* melanogaster. *Entomologia Experimentalis et Applicata, 155*, 64-70.
966
967

968 Data Accessibility
969 *D. melanogaster* sequence data have been submitted to Bioproject: accession number PRJNA482662. *D.*
970 *simulans* sequence data will be made available upon acceptance.
971

972 Competing interests
973 The authors declare that they have no competing interests.
974

975 Funding
976 This work was supported by grants GM102227 and MH09156.

978 Author's contributions
979 S.S. performed the experiments, analyzed the dataset, and wrote the manuscript. S.V.N. conceived of the
980 experiment and assisted in writing the manuscript.
981
982

D. simulans

	G	ETOH	ETOH x G	ETOH x T	ETOH x G x T	10	20	30
Exons								
Unique	96	6	16	2	75	1	0	6
Unique/Constitutive	1547	58	324	19	943	5	1	16
Common	52	2	3	0	19	0	0	1
Constitutive	299	10	44	3	121	3	2	6
Sum	**1994**	**76**	**387**	**24**	**1158**	**9**	**3**	**29**
Exonic Regions								
Common	119	9	22	1	76	1	0	2
Constitutive	287	14	76	6	223	2	1	2
Sum	**406**	**23**	**98**	**7**	**299**	**3**	**1**	**4**
Fragments								
Unique	101	2	5	2	26	1	1	2
Unique/Constitutive	169	7	45	3	93	0	0	1
Common	147	6	6	1	32	1	0	2
Constitutive	191	3	25	0	74	2	1	1
Sum	**608**	**18**	**81**	**6**	**225**	**4**	**2**	**6**

D. melanogaster*

	G	ETOH	G x ETOH	ETOH x T	G x ETOH x T	10	20	30
Exons								
Unique	79	3	0	0	0	0	0	4
Unique/Constitutive	630	1	1	12	0	1	0	17
Common	89	2	2	3	0	0	0	4
Constitutive	647	9	0	7	0	1	1	21
Sum	**1445**	**15**	**3**	**22**	**0**	**2**	**1**	**46**
Exonic Regions								
Common	84	2	0	1	0	0	1	2
Constitutive	547	11	0	7	2	1	3	22
Sum	**631**	**13**	**0**	**8**	**2**	**1**	**4**	**24**
Exon Fragments								
Unique	191	6	0	5	1	0	1	10
Unique/Constitutive	99	0	0	1	1	0	0	2
Common	301	4	0	1	1	1	1	7
Constitutive	544	11	0	5	1	2	3	1
Sum	**1135**	**21**	**0**	**12**	**4**	**3**	**5**	**20**

*reproduced from Signor and Nuzhdin 2018

984 **Table 1**: A summary of the exons, exonic regions, and exon fragments that are significantly differently
985 expressed for each component of variance. Abbreviations are as follows: (G) Genotype, (ETOH)
986 Ethanol, and (T) Time.

		Biological Process		Cellular Component		Molecular Function	
D. simulans	Ethanol x genotype						
				axonemal dynein complex	3×10^{-3}	dynein light chain binding	3.3×10^{-3}
	Ethanol x genotype x time						
		sensory perception of taste	6.6×10^{-3}	nucleplasm	6.5×10^{-3}	RNA binding	7.8×10^{-3}
				cytoskeletal part	8.4×10^{-3}		
		detection of chemical stimulus	8.2×10^{-3}	ciliary part	4.6×10^{-3}		

989 **Table 2**: A summary of the significant GO enrichment terms for *D. simulans*. Two components of
990 variance (genotype x time; genotype x ethanol x time) have significant terms, the cutoff is $p < .01$.
991

	Threshold	Abundance: # of outliers	Rank abundance: # of outliers		Threshold	Abundance: # of outliers	Rank abundance: # of outliers
D. melanogaster **candidate gene regions**				*D. simulans* **candidate gene regions**			
D. melanogaster				*D. melanogaster*			
X-linked	-1.49	5	10	X-linked	-1.49	**13**	11
	2.29	4	9		2.29	**35**	19
Autosomal	-1.5	18	68	Autosomal	-1.5	191	1
	1.9	13	49		1.9	225	4
D. simulans				*D. simulans*			
X-linked	-1.89	**0**	10	X-linked	-1.89	77	77
	3.25	**15**	6		3.25	**46**	**4**
Autosomal	-1.09	**0**	37	Autosomal	-1.09	318	72
	3.06	**35**	67		3.06	267	15

992
993 **Table 3**: A summary of the threshold values for Tajima's *D* and the number of intervals in each species
994 which fall beyond this cutoff. X-linked and autosomal genes are shown separately, outlier categories
995 which are significantly enriched are shown in bold. The same numbers are shown as percentages of the
996 total number of intervals in Figure 4.
997

Figure 1: **A**. An illustration, for a single gene, of hypothetical gene expression changes for different genotypes with different main and interaction effects. For main effect genotype, no difference is seen in expression between environments but different genotypes have different expression levels. In response to the environment (ethanol), gene expression level changes between environments but for every genotype it changes in the same way. Ethanol by time is complicated by the fact that the x-axis is now time, and there are separate lines for ethanol (solid) and non-ethanol (dashed) environment. Here again there is a component of time for gene expression changes, but every genotype changes in the same manner. For genotype by environment interactions in the first panel different genotypes respond to ethanol differently, and in the second panel different genotypes respond to ethanol differently over time. **B**. An illustration of the classification scheme for exons, exonic regions, and exon fragments. Exons either do not overlap other exons in different isoforms (exons, shown in blue), or are fused regions consisting of a set of overlapping exons (exonic regions, shown in shades of orange and red). Exonic regions can be decomposed into exon fragments, depending upon their overlap between different isoforms. Exons may be unique (found in a single isoform), common (found in some isoforms but not all), or constitutive (present in all isoforms). **C.** An example of the information about gene expression and isoform abundance that can be inferred from measures of differential exon abundance. *cabut* is a relatively simple example, having few isoforms and exons, and yet it is still a complicated inter-species comparison given that *D. melanogaster* has more annotated isoforms that *D. simulans*. Below each gene model the frequency of different exons and exon fragments in each environment is shown, with the most frequent in dark blue. In

D. simulans the first of two isoforms shown has an exon fragment that is more abundant in the presence
of ethanol, suggesting that that isoform is more abundant. In *D. melanogaster* only the second of the three
isoforms has a unique fragment, which is also more abundant in ethanol and suggests differential isoform
usage. **D.** An illustration of a change in the rank abundance of a the gene *Drat*. Only the example of *D.*
simulans is shown. *Drat* has two annotated isoforms in *D. simulans*, with two constitutive exons and two
exons unique to each of the two isoforms. With ethanol, the exon unique to the first annotated transcript is
the most abundant, while without ethanol the first constitutive exon is most abundant. In addition, the
second constitutive exon is significant for changes in abundance for the interaction between treatment and
time, and the two constitutive exons and exon unique to the first transcript have significant differences in
expression for ethanol, and between environments at 30 minutes. The dynamic changes in the expression
of different exons are captured in different ways by these three approaches to quantifying changes in
expression and splicing, where even within one gene there can be four complicated effects on expression
characterized by two approaches.

1032
Figure 2: A. Components of variance for *D. melanogaster* and *D. simulans*. The number of significant
exons and exonic regions for each component of variance are shown. Please note that in cases where so
few exons and exonic regions are significant that no color can be seen, *D. melanogaster* is always on the
left. Many more exons and exonic regions are significant for *D. simulans* than *D. melanogaster*, but only
for components of variance that contain an interaction with genotype. **B.** *D. melanogaster* and *D.*
simulans differ markedly in their genotype-specific reactions to ethanol, thus shown here is the relative
expression in ethanol (*y*-axis) against the relative expression without ethanol (*x*-axis) for all genes with an
interaction term with genotype. Too few genes are significant in *D. melanogaster* to make anything but a
qualitative comparison, but the graphs serve to illustrate the magnitude and nature of this contrast. **C.** The
proportion of exons and exonic regions that are either unique, unique/constitutive, common, or
constitutive for the three components of variance for which a reasonable number of genes are significant
in *D. melanogaster*. Note that while species names are only indicated for the first component of variance,
D. melanogaster is on the left for every comparison. *D. melanogaster* has far more 'constitutive' exons
and exonic regions compared to *D. simulans*, where unique/constitutive is more common. This is likely

due to differences in the annotation of isoforms between the species, where *D. melanogaster* has more
isoforms per gene annotated. It is also possible that alternatively spliced genes are more important for
expression differences in *D. melanogaster*.
1050
1051

1052
Figure 3: **A.** The percentage of intervals that were considered outliers in *D. melanogaster* and *D.*
simulans, split between the autosomes and the X. The percentage of intervals that are outliers are shown
separately for positive and negative, with the percent negative shown as a negative number. Note that the
percentage is not actually negative but rather serves to illustrate the distinction between positive and
negative. Enrichment for outlier intervals was determined as a composite of positive and negative
intervals. Significant enrichment is indicated by a star, significant under-enrichment is illustrated with a
minus sign. *D. melanogaster* is always on the left. The genome-wide cutoff for outliers was ±2.5%, thus
enrichment is anything significantly more than or less than 5% for the positive and negative intervals
together. **B**. The distribution of D_{xy} for *D. simulans* and *D. melanogaster*, compared to the genome-wide
background distribution of D_{xy}. The genes are split between the autosomes (top) and X-linked (bottom),
and the genome-wide background distribution is shown in blue. For *D. melanogaster* and *D. simulans* the
color corresponding to changes in the abundance of exons or exonic regions is shown first, while that
corresponding to changes in rank abundance is shown second , i.e. *D. melanogaster* pink – changes in
overall abundance; green – changes in rank abundance.
1067
1068

1069
Figure 4: An illustration of the possible explanations for an enrichment in nested genes, and nested non-
protein coding genes, in *D. simulans*. Note that in each illustration there are many possible details for
each scenario – for example gene expression could increase or decrease in the presence of ethanol, but we
illustrate here an increase in scenario **A.** This does not assume that this particular change is more likely,
only that illustrating both an increase and a decrease will not aid in the readers comprehension. A. In the
first scenario the nested gene changes its expression without any relationship with the expression of its
parental gene, as shown by the number of squiggly lines representing RNA. Generic exons are
represented by colored squares, introns by lines. The parental gene is shown in mustard yellow, and the
nested gene in magenta, with RNA belonging to either in matching colors **B.** It is possible that ethanol
alters the stability of introns after they are spliced out of the maturing transcript from the parental gene,
potentially altering the expression level of the non-protein coding gene within the intron. Shown here and
in the remaining scenarios is an extra step, with an unspliced RNA transcript from the parental gene
preceding a spliced transcript. We do not attempt to accurately depict the process or machinery of
splicing, only to illustrate that while expression of the nested gene may occur independently, some
expression may also occur from introns as they are spliced out of the parental gene. **C.** Expression of the
nested gene could also be affected if there were changes in the number of unspliced transcripts, for
example through ethanol-induced alterations in the rate of splicing. **D.** The expression of the nested gene
could also appear to change, or actually change through changes in the expression of the parental gene.
Note that **C-D** would all predict some amount of correlation between parental gene expression and nested
gene expression – such that if a change in nested gene expression is detected a change in parental gene
expression would generally be expected as well.
1091
1092