Evolution of phenotypic plasticity in response to ethanol between sister species with different ecological histories (*Drosophila melanogaster* and *D. simulans*)

Sarah A. Signor[1,2*], Sergey V. Nuzhdin[1]

1. Molecular and Computational Biology, University of Southern California.

2. Integrative Biology, University of California, Berkeley

* Communicating author: sasignor@berkeley.edu

Running title: Evolution of plasticity in response to ethanol

Please note: Supplemental Files 1-3 are currently available at: https://github.com/signor-molevol/simulans-ethanol.git

Acknowledgements
The authors would like to thank Jeremy Newman and Lauren McIntyre for contributions to the manuscript. The authors would like to thank our undergraduate researchers for assistance in producing this data: N. Shadman, V. Paterson, Z. Polonus, K. Cortez, L. Hassanzadeh, L. Cline, A. Khokhar, E. Lee, K.L. Yee, M. Ling, S. Sarva, O. Akintonwa, A. Gupta, R. Manson, P. Hassanzadeh, and K. Kavoussi. Lastly, the authors would like to thank B. & C. Emery for thoughtful commentary.

Data Accessibility
D. melanogaster sequence data have been submitted to Bioproject: accession number PRJNA482662. *D. simulans* sequence data will be made available upon acceptance.

Competing interests
The authors declare that they have no competing interests.

Funding
This work was supported by grants GM102227 and MH09156.

Author's contributions
S.S. performed the experiments, analyzed the dataset, and wrote the manuscript. S.V.N. conceived of the experiment and assisted in writing the manuscript.

Abstract

The contribution of phenotypic plasticity to adaptation is contentious, with contradictory empirical

support for its role in evolution. Phenotype plasticity has been posited as being important for adaptation to

novel environments, invasion success, and survival in response to climate change, as it could potentially

increase the likelihood of initial persistence. Here we investigate the possibility that phenotype plasticity

has contributed to adaptation to a novel resource. If phenotype plasticity contributes to adaptation, it is

thought to evolve in a process termed genetic accommodation. Under this model, the initial response to

the environment is widely variable due to cryptic genetic variation, which is then refined by selection to a

single adaptive response. We examine the role of phenotypic plasticity in adaptation here by comparing

two species of *Drosophila* that differ in their adaptation to ethanol (*Drosophila melanogaster* and *D.*

simulans). Both species are human commensals with a recent cosmopolitan expansion, but only *D.*

melanogaster is adapted to ethanol exposure. We measure phenotype plasticity in response to ethanol

with gene expression and an approach that combines information about expression and alternative

splicing. We find evidence for adaptation to ethanol through genetic accommodation, suggesting that the

evolution of phenotype plasticity contributed to the ability of *D. melanogaster* to exploit a novel resource.

We also find evidence that alternative splicing may be more important for the adaptive response to

ethanol.

Keywords: phenotypic plasticity, *Drosophila*, ethanol, genetic accommodation

Introduction

The contribution of phenotypic plasticity to adaptation remains controversial despite considerable empirical and theoretical investigation. This includes whether phenotypic plasticity itself evolves, and how frequently the evolution of plasticity is important for adaptive evolution. Recent work has suggested that the evolution of plasticity may be important for adaptation to changing environments (Lande 2009; Scoville and Pfrender 2010; Chevin et al. 2012; Oostra et al. 2018; Li et al. 2018). For example, in a novel environment phenotypic plasticity may increase the probability that a species can persist, which can be followed by adaptation towards local optima (Baldwin 1902; Simpson 1953; (Miner et al. 2005; Ghalambor et al. 2007; Lande 2009; Murren et al. 2015; Velotta et al. 2018). If plasticity itself is adaptive, then selection would proceed by favoring the reaction norm that is most adaptive, a process referred to as genetic accommodation (in contrast to simply fixing a different phenotype, known as genetic assimilation) (Via et al. 1995; West-Eberhard 2005). This could potentially accelerate adaptation or divergence because organisms already possess the ability to form alternative phenotypes, rather than requiring the production of a novel phenotype. This is thought to be an important mechanism of adaptation for invasive species, and potentially to be increasingly important in human modified landscapes and climate.

When exposed to novel environments, the predicted path for evolution by genetic accommodation is one in which initially there is increased variance, or genotype-specific differences – in the reaction norm of a trait, followed by refinement by selection for a single adaptive response (West-Eberhard 2005; Ghalambor et al. 2007; Robinson 2013; Schlichting and Wund 2014; Morris et al. 2014). This is because if a trait only exists as an environmentally induced variant, and is therefore not exposed to selection (or infrequently so) then genetic variation should accumulate in the response of that trait to a novel environment (Schlichting 2008; Gibson 2009; Hayden et al. 2011; Paaby and Rockman 2014). After exposure to the novel environment, this cryptic genetic variation is uncovered, manifesting as greater phenotypic variation (Rutherford 2000; Gibson and Dworkin 2004; Hermisson and Wagner 2004;

Schlichting 2008). As lineages adapt to the novel environment, selection on this variation in reaction

norm should result in the loss of variation for this trait as genetic accommodation occurs (Baldwin 1896;

Waddington 1953; West-Eberhard 2003). As long as the environment is heterogeneous the expectation is

that it will remain a plastic trait, rather than evolving to be stably expressed. This would manifest as a

transient increase in genotype by environment interactions, followed by reduced genotype by

environment interactions but persistent environmental response, a pattern which is consistent evidence of

adaptation (Via and Lande 1985; Guntrip and Sibly 1998; Lande 2009; Matzkin 2012; Huang and

Agrawal 2016).

While changes in heritability or variance have been previously shown between populations

occupying ancestral and novel habitats, inbred lines have not been used previously to specifically measure

genotype by environment interactions in adapted and non-adapted populations. Furthermore, in general

the evolution of phenotypic plasticity is investigated with regards to an ancestral and a novel

environment, rather than a heterogeneous environment where plasticity is expected to be maintained. In

this manuscript we will test the predictions from genetic accommodation using inbred lines in

Drosophila, an approach which will allow direct quantification of the genotype by environment

interaction. We will also investigate a heterogeneous environmental variable, alcohol, for which different

species of *Drosophila* have different adaptive histories.

D. simulans and *D. melanogaster* have divergent adaptive histories with respect to ethanol. *D.*

melanogaster is considerably more ethanol tolerant than *D. simulans*, and is found regularly feeding upon

and ovipositing in resources with ethanol concentrations greater than 8% (Hoffmann and McKechnie

1991; Fry 2014; Zhu and Fry 2015). At concentrations of 4% ethanol *D. simulans* shows reductions in

survivorship and increased development time relative to *D. melanogaster* (Joshi and Thompson 1997).

Furthermore*, D. melanogaster* and *D. simulans* are both cosmopolitan species and human commensals,

however while *D. melanogaster* is commonly found inside houses, breweries, and wineries, *D. simulans*

is more often observed in orchards and parks (though their niches overlap significantly and they are often

found on the same patches) (David and Bocquet 1976; Parsons and King 1977; McKenzie and

McKechnie 1979; Parsons and Spence 1981; Dickinson et al. 1984; Gibson and Wilks 1988; Thomson et

al. 1991; Chakir et al. 1993; Mercot et al. 1994; Joshi and Thompson 1997). Therefore, the expectation is

that *D. simulans* will show a non-adapted phenotype manifested as considerable genotype by environment

interaction. *D. melanogaster* will retain an interaction with the environment, but it will not be genotype-

specific, which is consistent with adaptation to a heterogeneous environment.

The trait we measure to test this prediction is gene expression. Phenotypic plasticity is often

achieved by dynamic changes in gene expression across environments, and current evidence suggests that

it plays an important role in regulating evolutionarily important phenotypes. (Hodgins-Davis and

Townsend 2009; Levine et al. 2011; Yampolsky et al. 2012; Hodgins-Davis et al. 2012). Gene by

environment interactions also appear to be a common attribute of gene expression, and yet while many

studies have quantified the effect of the environment on gene expression, far fewer have performed these

experiments in multiple genotypes to obtain the interaction between genotype and the environment for

gene expression (DeBiasse and Kelly 2015). Indeed, most work on gene expression plasticity has not

quantified the interaction between genotype and the environment (Cheviron et al. 2008; McCairns and

Bernatchez 2009; Hodgins-Davis et al. 2012; DeBiasse and Kelly 2015; Mathur and Schmidt 2016;

Heckel et al. 2016).

Short read sequencing fundamentally cannot separate the contributions of alternative splicing and

gene expression to changes in abundance of an exon, thus we approach our analysis in a way that includes

both phenomenon (gene expression and alternative splicing, see Methods). In addition, because of the

potential for alternative splicing to respond more rapidly to environmental differences than gene

expression *sensu stricto*, it may be one of the earliest mediators of environmental response. This is

because it does not require de novo transcription or protein production. It is thought to diverge more

rapidly than gene expression between lineages, and it has been linked to stress responses and changes in

transcriptome complexity in response to environmental differences (Merkin et al. 2012; Barbosa-Morais

et al. 2012; Gueroussov and Gonatopoulos-Pournatzis 2015; Jakšić and Schlötterer 2016; Singh et al.

2017; Wang et al. 2017; Pajoro et al. 2017; Calixto et al. 2018). Furthermore, it has been previously

implicated in the response to ethanol, and in *D. melanogaster* may be a more important component of the

response than gene expression changes (Oomizu et al. 2003; Newton et al. 2004; Pietrzykowski et al.

2008; Sasabe and Ishiura 2010; Hemby 2012; Zaharieva et al. 2012; Robinson and Atkinson 2013). Yet it

remains the topic of few investigations into environmental response.

Overall, we compare plasticity in response to ethanol, including genetic variation for plasticity

over time and between genotypes, among *Drosophila* with different patterns of ethanol utilization. We

measure plasticity and variation in expression and splicing in response to ethanol treatment. We address

three questions from theoretical work on phenotypic plasticity: Is cryptic genetic variation uncovered in

the non-adapted species (*D. simulans*)? Do we observe maintenance of an environmental response and

removal of genotype by environment interactions in the adapted species (*D. melanogaster*)? What

contribution does alternative splicing have to adaptation? We find that in *D. melanogaster*, which

regularly exploits ethanol-rich substrates, there is maintenance of an environmental response of similar

magnitude to *D. simulans*. However, there is essentially no genetic variation for plasticity within the

tested *D. melanogaster* population, while *D. simulans* exhibits a much larger effect of genotype by

environment compared to other components of variance (cryptic variation). This suggests the following

scenario: In *D. simulans* high ethanol concentrations are essentially a novel environment that reveals

cryptic variation for phenotypic plasticity. In *D. melanogaster*, genetic variation in phenotypic plasticity

has been removed by selection in favor of a single environmental response through a process of genetic

accommodation. This single environmental response involves more alternative splicing than observed in

D. simulans, suggesting it is an important component of adaptation.

Methods

Fly lines

Six genotypes from each species were used in these experiments. The genotypes used for male flies from

D. melanogaster were collected from an orchard in Winters, California in 1998 and made isogenic by 40

generations of full sibling inbreeding (Yang and Nuzhdin 2003; Campo et al. 2013; Signor et al. 2017b,c).

The genotypes used for male flies from *D. simulans* were collected from the Zuma Organic Orchard in the

winter of 2012 and made isogenic by 15 generations of full sib inbreeding (Signor et al. 2017a,b).

Residual heterozygosity is similar between these lines of *D. melanogaster* and *D. simulans* (Campo et al.

2013; Signor et al. 2017a). In natural conditions flies will not by homozygous at all loci, thus each inbred

genotype was crossed to a white eyed 'tester' strain to create the F1 flies used in the ethanol exposure

assays (*D. melanogaster*, w^{1118}, Bloomington stock number 3605; *D. simulans*, w^{501}, Cornell species stock

number 14021-0251.011). Rearing occurred on a standard medium at 25 °C with a 12-h light/12-h dark

cycle. In order to control for maternal effects and variation in offspring quality, female parents were

collected as virgins, aged one day, and then density matched with male flies (10 per sex). The F1

offspring were then collected as virgins, reared in single sex vials with standardized density (24-30 flies if

male, 12-15 if female), and aged for three to four days prior to the ethanol exposure assay.

A portion of this experiment was intended to analyze behavioral differences in ethanol, thus during

the ethanol exposure a female was included as a stimulus, but was not collected for RNA-seq (Signor et al.

2017b,c). The females were from yw^1 mutants for both *D. simulans* and *D. melanogaster* (*D. melanogaster*,

Bloomington stock number 1495, *D. simulans*, Cornell species stock number 14021-0251.013). The day

prior to the ethanol exposure these females were mated to flies from a standard genotype, such that in the

final assay each chamber contained one mated female fly from the yw^1 mutant lines and two virgin males

from the target F1 cross, of which only the latter were collected for RNA-seq.

Experimental setup

In this experiment intoxication occurs through inhalation of ethanol vapors, and evidence of the

behavioral effects in both species and the efficacy of this approach have been published previously

(Signor et al. 2017b,c). Ethanol exposure took place in a circular arena, each of which was part of a

larger chamber containing 12 arenas each with a diameter of 2.54 cm (VWR cat. no. 89093-496). The

bottom of each chamber contained a standard amount of either grapefruit medium or medium in which

15% of the water has been replaced with ethanol. Replicates for both species were conducted randomly

under standardized conditions (25 °C, 70% humidity). Prior to the assay the flies were sedated through

exposure at 4 °C for ten minutes, to avoid the confounding effects of CO_2, then placed in the chambers

which were left then upside down at room temperature for ten minutes while the flies oriented themselves

(see Signor et al. 2017a; b; Signor & Nuzhdin 2018). After ten minutes the timing of the assay began, and

were conducted for ten, 20, or 30 minutes for three replicates of each of the two conditions. Flies are most

active during the hours following dawn, thus to standardize behavior and circadian rhythms all assays

were conducted within a two-hour window after dawn. At the conclusion of the assay the chambers were

flash frozen in liquid nitrogen, allowed to freeze through, transferred to dry ice, and all of the males were

collected for RNA-seq.

Sample preparation and RNA sequencing

Sample preparation has been described previously, and will be briefly summarized here (Signor and

Nuzhdin 2018). Flash-frozen flies were freeze dried and ten to 12 heads were placed into a 96-tube plate

(Axygen MTS-11-C-R). mRNA purification, cDNA synthesis and library preparation were carried out by

RAPiD GENOMICS (http://rapid-genomics.com) using a robot. mRNA was purified using Dynabeads

mRNA DIRECT Micro kit (Invitrogen # 61021) with slight modifications. To fragment the RNA mRNA-

beads were resuspended in 10 uL 2X first strand buffer (Invitrogen # 18064-014), incubated at 80 C for two

minutes and placed on ice, then the supernatant was collected after five minutes on a magnetic stand. First

strand synthesis was performed using standard protocols for Superscript II (Invitrogen #18640-014) and

reverse transcription (25 C 10 min, 42 C for 50 min, 70 C for 15 min, 10 C hold). Second strand synthesis

was carried out using standard protocols with DNA Pol I and incubated at 16 C for 2.5 hours. cDNA was

purified with 1.8 volume of AMPure XP following manufactures instructions (Beckman Coulter A63880).

Sequencing was performed using the Illumina HiSeq 2500 as both 2 × 150 bp or 2 × 50 bp reads, resulting

in two technical replicates per sample. The two run lengths (and runs) were intended to provide extra

coverage, and all replicates were sequenced in both runs.

Exon expression analysis

It is common in organisms with alternative splicing for exons from different isoforms of a single gene to

overlap with one another, or be shared between all or most isoforms (Figure 1A). Short read data

fundamentally cannot resolve these exons to individual isoforms, however, one approach is to quantify each

exon separately and decompose exons overlapping between isoforms into those which are shared and

unique. When the differences between overlapping exons are less than 10 bp, there is no appreciable

amount of information loss in not decomposing overlapping exons, and this approach has been taken in the

past (Dalton et al. 2013; Graze et al. 2014; Fear et al. 2016; Newell et al. 2016). However, in many cases

the differences in exon overlap are much larger than this, so to address this issue we use a classification

scheme where reads may be assigned to *exons*, *exonic regions*, or *exon fragments*, and then compare the

abundance of each exon/exonic region/exon fragment in each condition (i.e. the abundance of an exon with

and without ethanol) (Signor and Nuzhdin 2018) (Figure 1A). Here, an *exon* does not overlap any other

exons. If exons from different isoforms overlap, they are grouped into an *exonic region*. *Exon fragments*

are classified by decomposing *exonic regions* based on the 5' and 3' positions of the exons within the region.

Thus, all *exon fragments* are subregions of *exonic regions*. Exon boundaries were determined using the *D.*

melanogaster FlyBase 6.17 genome features file and the *D. simulans* 2.02 genome features file. Alignment

was performed using BWA-MEM version 0.7.15 (which has been shown to perform better than split read

mappers such as STAR (Newman et al. 2018)) and BED files were used to count reads in each region and

obtain the length adjusted read count (reads in region divided by the length of region), and the APN (average

per nucleotide) (Li 2015).

The APN was calculated separately for each read length and then combined between read lengths

to handle the mixture of read lengths for each sample (2×150 bp and 2×50 bp). If the APN was greater

than zero in at least half of all samples per condition it was considered detected. While we considered

several approaches to normalize coverage counts upper-quartile normalization with log-transformation and

median centering within time × treatment × genotype were selected due to better performance of the

residuals (Bullard et al. 2010; Dillies et al. 2013).

To test the significance of components of expression variation, the log APN for each exonic region was

modeled as

$$Y_{ijkl} = \mu + g_i + t_j + (gt)_{ik} + m_k + (gm)_{ik} + (tm)_{ijk} + \epsilon_{ijkl}$$

for the ith genotype (g_i), jth treatment (t_j; j = ethanol or no ethanol), kth time point (m_k; k = 10 min, 20 min, 30

min), and lth replicate (Supplemental File 1-2). For the interaction between treatment and time point, the log

APN for each exonic region was modeled as

$$Y_{ij} = \mu + t_j + \epsilon_{ij}$$

or the ith condition (time × treatment) and jth replicate. Contrasts to compare treatments within time point

(ethanol versus no ethanol, for ten, 20 and 30 minutes) were conducted (Supplemental Files 1-2). Residuals

were evaluated for conformation with normality assumptions, and assumptions were met in excess of 95%

of the models.

In contrast to exon abundance, we evaluated alternative splicing specifically by comparing the

abundance of all exons within a gene to each other, as this is more direct evidence of a change in isoform

abundance than a change in exon/exonic region/exon fragment abundance; Figure 2A). For example, a

change in relative abundance might be detected if the last exon of a gene was the most abundant compared

to all other exons/exonic regions/exon fragments without ethanol and the least abundant with ethanol.

Exons and exonic regions for each gene and for each sample were ranked and the most expressed region

ranked as one, the least expressed region as three and all others as two (Figure 2A). Exon ranks for each

gene were modeled as

$$Y_{ijk} = \mu + r_i + t_j + (rt)_{ij} + \epsilon_{ijkl}$$

where Y_{ijk} is the exon rank (1,2,3) of the iᵗʰ exonic region of the gene, jᵗʰ condition (time × treatment), and

the kᵗʰ replicate; r_i is the exonic region of the gene; t_j is condition; and $(rt)_{ij}$ is the interaction between

exonic region and condition. More traditional GLM approaches can only be taken if their assumptions are

met, and in this case they are not due to a lack of normality in the distribution of model residuals.

Accordingly, a non-parametric test must be relied upon to look for changes in exon or exonic region

representation between exons of a gene and we used a rank test to summarize changes in exon representation

(Supplemental File 3). F-tests for the significance of the mean square attributed to the effect tested versus

the mean square attributed to error, or the appropriate interaction term, were used. The false discovery rate

was controlled using the Benjamini-Hochberg procedure, with a significance cutoff of $\alpha = 0.05$ (Benjamini

& Hochberg 1995).

GO Analysis

When a gene in *D. simulans* had more than one ortholog in *D. melanogaster* only one ortholog was

included for the GO analysis, so as not to inflate the number of genes involved in a given process. This

does presume that orthologs will be annotated with the same GO terms, and this is generally the case. For

example, in *D. simulans* there is only *AOX4*, while in *D. melanogaster* there is *AOX3* and *AOX4*, but the

GO terms for each are the same. However, as *D. simulans* genes are generally not independently

annotated, especially those without *D. melanogaster* orthologs, if there was no *D. melanogaster* ortholog

the gene was not included in the enrichment analysis. Lists of significant genes were tested for GO

enrichment using the PANTHER classification system (Mi et al. 2017). They were corrected for multiple

testing and a *p*-value of 0.01 was required for significance.

Functional class enrichment

We tested the significant sets of exons and exonic regions for enrichment with non-protein coding genes.

Multigene exonic regions were not included, meaning exons that belong to more than one gene, as they

often do not correspond to the same functional class of gene. Every test of functional class enrichment

compared the frequency of a given subcategory among all exons and exonic regions detected in the

dataset compared to the frequency within a significant list of exons and exonic regions. A χ^2 test was

performed in R to test the significance of the enrichment of each of these categories.

Results

Exon expression and isoform usage

It is difficult to decouple alternative isoform usage from gene expression, given that many exons are

shared between isoforms or overlap other exons. To infer isoforms from short read data, one must rely

upon unique junctions or regions of individual isoforms and extrapolate to shared regions. This requires

accurate isoform annotation (knowing that any given exon/junction is found in combination with other

exons/junctions) and in general can be very noisy. Accordingly, we subscribe to a simpler but more robust

approach and summarize the abundance of different exons and exonic regions separately (Dalton et al.

2013; Graze et al. 2014; Newell et al. 2016; Signor & Nuzhdin 2018). We detail overall abundance of

exons, exonic regions, and exon fragments, and changes in the relative abundance of exons within a gene

(Figure 1 A-C; Figure 2 A&B).

Changes in exon expression in D. melanogaster and D. simulans

Table 1 summarizes the number of exons, exonic regions, and exon fragments which alter their

expression in response to ethanol, but note that the table for *D. melanogaster* is reproduced from Signor

& Nuzhdin 2018. For treatment seven exons and exonic regions were shared between species: *Drat*,

cabut, *CG11741*, *CG32512*, *CG4607*, *Pinocchio*, and *sugarbabe* (Table 1; Figure 1 A-C, Signor &

Nuzhdin 2018). For the interaction between ethanol and time three genes were shared between exons and

exonic regions for these species, *Drat*, *cabut*, and *CG43366*. At ten and twenty minutes *cabut* is shared

between exons and exonic regions in *D. melanogaster* and *D. simulans*. At 30 minutes *Drat*, *CG32103*,

sugarbabe, *cabut*, *Pinocchio*, *CG32512*, and *CG4607* are shared between species. This suggests that

Drat, *cabut*, *sugarbabe*, *Pinocchio*, *CG32512*, and *CG407* are important for the response to ethanol, as

they are shared between species for multiple components of variance and have been implicated in the

response to ethanol previously (Morozova et al. 2006; Kong et al. 2010; Awofala et al. 2012).

GO enrichment analysis of differentially expressed exons in D. melanogaster and D. simulans

D. simulans is enriched for several GO terms in response to interactions with genotype, including several

terms relating to cilia, such as the axonemal dynein complex and ciliary part (Table 2). GO terms were

also enriched for the sensory perception of taste and detection of chemical stimulus. Many components of

variance in *D. simulans* are not enriched for GO terms due in part to the frequency with which non-

protein coding genes were implicated – for example in response to ethanol 15% of the implicated genes

do not have a *D. melanogaster* ortholog, and of those only two are protein coding . In *D. melanogaster*

exons and exonic regions were not significantly enriched for any category of genes. In general, the small

number of genes implicated for many categories precludes any conclusion of enrichment.

Changes in the relative abundance of exons within a gene in D. melanogaster and D. simulans

In response to treatment in *D. simulans* 54 genes showed changes in the relative abundance of their

constitutive exons/exonic regions, and 94 genes change the relative abundance of their exons for the

interaction between treatment and time (Figure 2B). In *D. melanogaster* 71 genes changed the relative

abundance of their exons in response to ethanol and 145 changed for treatment by time (Signor and

Nuzhdin 2018). For the interaction between treatment and time *lola*, *Mhc*, and *Prm* were shared between

species.

GO enrichment analysis of genes with changes in relative abundance in D. melanogaster and D. simulans

Changes in relative abundance were not enriched for any GO terms in *D. simulans*, with the exception of

peroxiredoxin activity in response to ethanol, albeit slightly above the more conservative cut-off used in

our other tests ($p = .017$). Peroxiredoxin activity has been associated with protection against alcohol

induced liver damage (Bae et al. 2011; Chattopadhyay et al. 2015). In *D. melanogaster* genes implicated

in changes in relative abundance in response to ethanol by time were enriched for cellular components

actin cytoskeleton, again somewhat above our more conservative cutoff ($p = 0.018$) (Signor and Nuzhdin

2018).

In D. simulans genotype-specific reactions to the environment are abundant

In *D. simulans* many exons and exonic regions are significant for components of variance that interact

with genotype: 1457 for the interaction between genotype, treatment and time, and 486 for the interaction

between genotype and treatment, compared to two and three exons and exonic regions respectively in *D.*

melanogaster (Table 1; Figure 1 A-C). For other components of variance, the number of exons and

exonic regions implicated in expression differences in roughly comparable between species.

In D. simulans interactions with genotype are enriched for nested non-protein coding genes

A large number of the genes implicated in differences due to interaction with genotype were non-protein

coding genes. The number of non-protein coding genes that were significant for ethanol, ethanol by

genotype, and ethanol by genotype by time, were more than would be expected by chance in *D. simulans*

($\chi^2 = 49.598, p < 0.0005; \chi^2 = 235.21, p < 2.2 \times 10^{-16}; \chi^2 = 727.17, p < 2.2 \times 10^{-16}$). These are

generally long non-coding RNAs (lncRNAs) as the shortest is annotated at 713 bp, and the majority are

over 4,000 bp (long non-coding RNAs being any non-protein coding genes over 200 nt), with some

pseudogenes. *D. melanogaster* is not enriched for non-protein coding genes in response to ethanol,

ethanol by time, ethanol by genotype, ethanol by genotype by time, ten or 20 minutes, exons or exonic

regions expressed only in one environment, or genes implicated in changes in relative abundance.

However, at 30 minutes *D. melanogaster* is enriched for non-protein coding genes, though this concerns

far fewer genes than in *D. simulans* ($\chi^2 = 12.831, p = 0.005$).

Many of the non-protein coding genes implicated in interactions with genotype are nested in the

introns of other genes, thus we sought to determine if it was more than expected by chance. First we must

determine the overall frequency of gene nesting, using the criteria that an exon nested in an intron must

overlap the intron by at least 80 bp or 10%. We found that in *D. melanogaster* 9.2% of exons were nested

within introns, while in *D. simulans* 9.7% were nested, similar to what has been previously reported (Lee

and Chang 2013). This was reflected in our data, where for both *D. melanogaster* and *D. simulans* 6.8%

of exons and exonic regions were nested within introns (lower because multi-gene exons were excluded).

However, among significant exons and exonic regions in *D. melanogaster* 18.6% were nested, while in *D.*

simulans 33.5% were nested. This is a significant enrichment of exons that are nested within other

introns, for both *D. melanogaster* ($\chi^2 = 12.344, p < .0004$) and *D. simulans* ($\chi^2 = 1721.1, p <$

2.2×10^{-16}). In addition, compared to the total number of nested genes that are noncoding within the

dataset, the number that are significant for the response to ethanol is enriched for nested, non-protein

coding genes in *D. simulans* ($\chi^2 = 237.49, p < 2.2 \times 10^{-16}$), but not in *D. melanogaster*. Of these

nested non-protein coding genes, 83% are on the opposite strand as their parental genes, suggesting that

these genes are regulated independently of their parental genes (and their expression is uncorrelated, see

Supplemental File 4). Overall *D. melanogaster* has more annotated non-protein coding genes (2963) than

D. simulans (1675), and similarly more exons from noncoding genes are nested within introns (1772 *D.*

melanogaster, 1066 *D. simulans*). The importance of nested non-protein coding genes for the response to

the environment, or potentially delays in splicing that are specific to certain components of variance, is

unclear.

<u>Discussion</u>

The prediction from theory on genetic accommodation was that in *D. simulans* there would be abundant

cryptic genetic variation in response to ethanol exposure, while in *D. melanogaster* there would be a

response to the environment that was not genotype-specific. This is because *D. simulans* does not exploit

ethanol-rich resources, thus environmentally induced variants are not exposed to selection and accumulate

as cryptic genetic variation. This manifests as greater variation between genotypes in the novel

environment (Rutherford 2000; Gibson and Dworkin 2004; Hermisson and Wagner 2004; Schlichting

2008). In *D. melanogaster* ethanol is used as a resource, however the environment is patchy and therefore

response to ethanol is expected to be selected upon for the optimal response, but to remain plastic. This

would manifest as an environmental response, but without extensive genotype-specific variation for that

response (Via and Lande 1985; Guntrip and Sibly 1998; Lande 2009; Matzkin 2012; Huang and Agrawal

2016). The use of inbred lines is a unique opportunity to assess these predictions, as the contribution of

interactions with genotype can be directly assessed.

Here we show that these expectations are met in *D. melanogaster* and *D. simulans*. In *D.*

melanogaster adaptation to ethanol has occurred through genetic accommodation – selection on plasticity

to reduce genetic variation and produce a single plastic response. This has facilitated expansion to a novel

resource in *D. melanogaster*, ethanol rich substrates. In *D. simulans* this is not the case, and lack of

selection in ethanol environments has allowed for the accumulation of cryptic genetic variation for

phenotype plasticity manifested as extensive genotype by environment interactions. Joshi and Thompson

(1997) found that in *D. simulans* there was a reduction in variation between families for the phenotypic

response to ethanol substrate after selection for tolerance to ethanol. This is consistent with the observed

scenario in D. *melanogaster* and *D. simulans* – adaptation to a novel environment results in a reduction in

variation for environmental response and while retaining a single response to the environment. This is

also consistent with genetic accommodation as a mechanism of adaptation to ethanol, where initial

responses to environmental differences are a mix of adaptive and maladaptive, which are subsequently

honed by selection.

Previous literature on the response to novel environments is equivocal concerning the expectation

that plasticity evolves from a starting point of increased genetic variance due to cryptic genetic variation.

For example, early estimates of heritability found reduced expression of genetic variation under 'stressful'

conditions – though stressful does not necessarily imply novel (Charmantier and Garant 2005; Chevin and

Hoffmann 2017). Work in *Drosophila* has been similarly ambiguous – in *D. mojavensis* a similar

reduction in genotype by environment interactions was detected in adaptation to a novel cactus host,

however in *D. melanogaster* differences in genotype by environment interactions were not detected

between cold-adapted and non-cold adapted populations (Matzkin 2012; Heckel et al. 2016). These

differences may be due to difficulty in defining novelty, and in equating novelty with stress. As posited

by Chevin and Hoffman (2017), it may be that genetic variance increased in novel environments, but that

favorable conditions are rare – meaning that 'stressful' environments are commonly experienced resulting

in a reduction in genetic variation.

The abundance of lncRNAs which are involved in genotype by environment interactions in *D.*

simulans is perhaps not surprising, given that they have been implicated previously in dynamic responses

(Valadkhan & Valencia-Hipólito 2016). It is also possible they are more frequently involved because

lncRNAs are often less conserved, as there is no requirement for the maintenance of ORFs and codon

synonymy (Chodroff et al. 2010; Ulitsky et al. 2011; Ulitsky 2016; Quinn et al. 2016). If neutral variation

was allowed to accumulate without selection, and then uncovered in a novel environment (cryptic genetic

variation), preferential accumulation within less constrained sequences would be expected.

Alternative splicing has the potential to be an important component of adaptation and response to

the environment, as it does not require de novo transcription or protein production. In *D. melanogaster* it

is a more important component of the response to ethanol than in *D. simulans*, as evidenced by

differences in the frequency of significant changes in relative abundance of exons. The response to

ethanol in *D. melanogaster* is adaptive, thus it can be deduced that alternative splicing is important for

adaptation to ethanol. Alternatively, the fact that it is less important in *D. simulans* suggests that perhaps

there is selection against the accumulation of cryptic genetic variation for splicing patterns. This is not

surprising – previous studies have linked alternative splicing to the response to the environment (Merkin

et al. 2012; Barbosa-Morais et al. 2012; Gueroussov and Gonatopoulos-Pournatzis 2015; Jakšić and

Schlötterer 2016; Singh et al. 2017; Wang et al. 2017; Pajoro et al. 2017; Calixto et al. 2018), as well as

important adaptive differences (Singh et al. 2017; Dong et al. 2017; Evantal et al. 2018; Shi et al. 2019).

Furthermore, it has been previously implicated in the response to ethanol, and in *D. melanogaster* may be

a more important component of the response than gene expression changes (Oomizu et al. 2003; Newton

et al. 2004; Pietrzykowski et al. 2008; Sasabe and Ishiura 2010; Hemby 2012; Zaharieva et al. 2012;

Robinson and Atkinson 2013).

In general, the GO terms implicated in this study relate to known effects of ethanol, for example,

alcohol-induced ciliary dysfunction is a known consequence of alcohol exposure (Yang et al. 2015; Price

et al. 2018). Furthermore, the genes implicated in both species (*Drat*, *cabut*, *sugarbabe*, *Pinocchio*,

CG32512, and *CG407*) have generally been implicated in the response to ethanol in other studies. *cabut*

has been previously observed as being upregulated in response to ethanol, and it is in general responsive

to changes in metabolic conditions (Awofala et al. 2012; Havula and Hietakangas 2012; Bartok et al.

2015). *sugarbabe* and *cabut* are both downstream of the *Mondo-Mlx* sugar sensing pathway (Mattila et al.

2015), which has been linked to severe obesity, high circulating triglycerides, and tumorigenesis (Kooner

et al. 2008; Kathiresan et al. 2008; Herman et al. 2012; Carroll et al. 2015). *Drat* has been implicated in

ethanol-related cell death, though it is not as well studied as *cabut* (Chen et al. 2012). Specifically with

regard to other studies of ethanol exposure in *Drosophila* overlap is low between significant datasets,

however this may in large part be due to differences in methodology (Signor and Nuzhdin 2018).

Previous studies used targeted qPCR (which requires candidate genes) (Morozova et al. 2011), and

microarrays (Kong et al. 2010), as well as different assays and exposure lengths (inebriometer

(Morozova 2011, 2015), booz-o-mat (Kong et al. 2010). What we are primarily interested in here is the

overall pattern rather than the individual genes.

Inferring that gene expression differences are adaptive or non-adaptive remains a major challenge

in the study of gene expression reaction norms, given the lack of direct correlation between gene

expression phenotypes and organismal phenotypes. However, the patterns observed in *D. simulans*

suggest that abundant genotype by environment interactions have accumulated neutrally and become

uncovered in response to a novel environment. In contrast, in *D. melanogaster* this ethanol environment is

not novel and variation in plasticity has been selected out in favor of an adaptive phenotypic response.

lncRNAs are preferentially differentially expressed in *D. simulans* in response to ethanol likely because

they are less constrained and can accumulate more neutral variation. This is an excellent illustration of

genetic accommodation, where phenotypic plasticity has encouraged adaptation to a novel resource

within a patchy environment.

Acosta, G., Freidman, D. P., Grant, K. A., and Hemby, S. E. 2012. Alternative splicing of AMPA
subunits in prefrontal cortical fields of cynomolgus monkeys following chronic ethanol self-
administration. Front. Psychiatry. 3:72.

Awofala, A. A., J. A. Davies, and S. Jones. 2012. Functional roles for redox genes in ethanol sensitivity
in *Drosophila*. Funct. Integr. Genomics 12:305–315.

Bae, S. H., S. H. Sung, E. J. Cho, S. K. Lee, H. E. Lee, H. A. Woo, D.-Y. Yu, I. S. Kil, and S. G. Rhee.
2011. Concerted action of sulfiredoxin and peroxiredoxin I protects against alcohol-induced oxidative
injury in mouse liver. Hepatology 53:945–953.

Baldwin, J. M. 1896. A new factor in evolution. Am. Nat. 30:441-451.

Barbosa-Morais, N. L., M. Irimia, Q. Pan, H. Y. Xiong, S. Gueroussov, L. J. Lee, V. Slobodeniuc, C.
Kutter, S. Watt, R. Çolak, T. Kim, C. M. Misquitta-Ali, M. D. Wilson, P. M. Kim, D. T. Odom, B. J.
Frey, and B. J. Blencowe. 2012. The evolutionary landscape of alternative splicing in vertebrate species.
Science 338:1587–1593.

Bartok, O., M. Teesalu, R. Ashwall-Fluss, V. Pandey, M. Hanan, B. M. Rovenko, M. Poukkula, E.
Havula, A. Moussaieff, S. Vodala, Y. Nahmias, S. Kadener, and V. Hietakangas. 2015. The transcription
factor Cabut coordinates energy metabolism and the circadian clock in response to sugar sensing. The
EMBO Journal 34:1538–1553.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach
to multiple testing. J. Royal Stat. Soc., Series B, 57, 289–300.

Bullard, J. H., E. Purdom, K. D. Hansen, and S. Dudoit. 2010. Evaluation of statistical methods for
normalization and differential expression in mRNA-Seq experiments. BMC Bioinformatics 11:94–13.

Calixto, C. P. G., W. Guo, A. B. James, N. A. Tzioutziou, J. C. Entizne, P. E. Panter, H. Knight, H.
Nimmo, R. Zhang, and J. W. S. Brown. 2018. Rapid and dynamic alternative splicing impacts the
Arabidopsis cold response transcriptome. The Plant Cell. 40:1424-1444.

Campo, D., K. Lehmann, C. Fjeldsted, T. Souaiaia, J. Kao, and S. V. Nuzhdin. 2013. Whole-genome
sequencing of two North American *Drosophila melanogaste*r populations reveals genetic differentiation
and positive selection. Mol. Ecol. 22:5084–5097.

Carroll, P. A., D. Diolaiti, L. McFerrin, H. Gu, D. Djukovic, J. Du, P. F. Cheng, S. Anderson, M. Ulrich,
J. B. Hurley, D. Raftery, D. E. Ayer, and R. N. Eisenman. 2015. Deregulated Myc Requires MondoA/Mlx
for metabolic reprogramming and tumorigenesis. Cancer Cell 27:271–285.

Chakir, M., O. Peridy, P. Capy, E. Pla, and J. R. David. 1993. Adaptation to alcoholic fermentation in
Drosophila: A parallel selection imposed by environmental ethanol and acetic acid. Proc. Nat. Acad. Sci.
USA 90:3621–3625.

Charmantier, A., and D. Garant. 2005. Environmental quality and evolutionary potential: lessons from
wild populations. Proc. R. Soc. B 272:1415–1425.

Chattopadhyay, A., D. Pinkaew, H. Q. Doan, R. B. Jacob, S. K. Verma, H. Friedman, A. C. Peterson, M.
N. Kuyumcu-Martinez, O. M. McDougal, and K. Fujise. 2015. Fortilin potentiates the peroxidase activity
of Peroxiredoxin-1 and protects against alcohol-induced liver damage in mice. Sci. Rep. 1–16.

Chen, P., X. Tu, F. Akdemir, S. K. Chew, A. Rothenfluh, and J. M. Abrams. 2012. Effectors of alcohol-
induced cell killing in *Drosophila*. Cell Death Differ. 19:1655–1663.

Chevin, L.-M., and A. A. Hoffmann. 2017. Evolution of phenotypic plasticity in extreme environments.
Philos. Trans. R. Soc. B 372:20160138.

Chevin, L.-M., S. Collins, and F. Lefèvre. 2012. Phenotypic plasticity and evolutionary demographic
responses to climate change: taking theory out to the field. Funct. Ecol. 27:967–979.

Cheviron, Z. A., A. Whitehead, and R. T. Brumfield. 2008. Transcriptomic variation and plasticity in
rufous-collared sparrows (*Zonotrichia capensis*) along an altitudinal gradient. Mol. Ecol. 17:4556–4569.

Chodroff, R. A., L. Goodstadt, T. M. Sirey, P. L. Oliver, K. E. Davies, E. D. Green, Z. Molnár, and C. P.
Ponting. 2010. Long noncoding RNA genes: conservation of sequence and brain expression among
diverse amniotes. Genome Biology 11:R72.

Dalton, J. E., J. M. Fear, S. Knott, B. S. Baker, L. M. McIntyre, and M. N. Arbeitman. 2013. Male-
specific Fruitless isoforms have different regulatory roles conferred by distinct zinc finger DNA binding
domains. BMC Genomics 14:659.

David, J., and C. Bocquet. 1976. Compared toxicities of different alcohols for two *Drosophila* sibling
species: *D. melanogaster* and *D. simulans*. Comp. Biochem. Physiol. C, Comp. Pharmacol. 54:71–74.

DeBiasse, M. B., and M. W. Kelly. 2015. Plastic and evolved responses to global change: What can we
learn from comparative transcriptomics? J. Hered. 107:71–81.

Dickinson, W. J., R. G. Rowan, and M. D. Brennan. 1984. Regulatory gene evolution: adaptive
differences in expression of *alcohol dehydrogenase* in *Drosophila melanogaster* and *Drosophila*
simulans. Heredity 52 (Pt 2):215–225.

Dillies, M. A., A. Rau, J. Aubert, C. Hennequet-Antier, M. Jeanmougin, N. Servant, C. Keime, G. Marot,
D. Castel, J. Estelle, G. Guernec, B. Jagla, L. Jouneau, D. Laloe, C. Le Gall, B. Schaeffer, S. Le Crom, M.
Guedj, F. Jaffrezic, on behalf of The French StatOmique Consortium. 2013. A comprehensive evaluation
of normalization methods for Illumina high-throughput RNA sequencing data analysis. Briefings in
Bioinformatics 14:671–683.

Dong, W.-X., J.-L. Ding, Y. Gao, Y.-J. Peng, M.-G. Feng, and S.-H. Ying. 2017. Transcriptomic insights
into the alternative splicing-mediated adaptation of the entomopathogenic fungus *Beauveria bassianato*
host niches: autophagy-related gene 8 as an example. Environ. Microbiol. 19:4126–4139.

Evantal, N., A. M. Anduaga, O. Bartok, I. L. Patop, R. Weiss, and S. Kadener. 2018. Thermosensitive
alternative splicing senses and mediates temperature adaptation in *Drosophila*. bioRxiv, 1–63.

Fear, J. M., L. G. León-Novelo, A. M. Morse, A. R. Gerken, K. Van Lehmann, J. Tower, S. V. Nuzhdin,
and L. M. McIntyre. 2016. Buffering of genetic regulatory networks in *Drosophila melanogaster*.
Genetics 203:1177–1190.

Fry, J. D. 2014. Mechanisms of naturally evolved ethanol resistance in *Drosophila melanogaster*. J. Exp.
Biol. 217:3996–4003.

Ghalambor, C. K., J. K. Mckay, S. P. Carroll, and D. N. Reznick. 2007. Adaptive versus non-adaptive
phenotypic plasticity and the potential for contemporary adaptation in new environments. Funct. Ecol.
21:394–407.

Gibson, G. 2009. Decanalization and the origin of complex disease. Nat. Rev. Genet. 10:134–140.

Gibson, G., and I. Dworkin. 2004. Uncovering cryptic genetic variation. Nat. Rev. Genet. 5:681–690.

Gibson, J. B., and A. V. Wilks. 1988. The alcohol dehydrogenase polymorphism of *Drosophila*
melanogaster in relation to environmental ethanol, ethanol tolerance and alcohol dehydrogenase activity.
Heredity 60 (Pt 3):403–414.

Graze, R. M., L. M. McIntyre, A. M. Morse, B. M. Boyd, S. V. Nuzhdin, and M. L. Wayne. 2014. What
the X has to do with it: Differences in regulatory variability between the sexes in *Drosophila simulans*.
Genome Biol. Evol. 6:818–829.

Gueroussov, S., and T. Gonatopoulos-Pournatzis. 2015. An alternative splicing event amplifies
evolutionary differences between vertebrates. Science 349: 868-73.

Guntrip, J., and R. M. Sibly. 1998. Phenotypic plasticity, genotype-by-environment interaction and the
analysis of generalism and specialization in *Callosobruchus maculatus*. Heredity 81:198–204.

Havula, E., and V. Hietakangas. 2012. Glucose sensing by ChREBP/MondoA–Mlx transcription factors.
Semin. Cell Dev. Biol. 23:640–647.

Hayden, E. J., E. Ferrada, and A. Wagner. 2011. Cryptic genetic variation promotes rapid evolutionary
adaptation in an RNA enzyme. Nature 474:92–95.

Heckel von, K., W. Stephan, and S. Hutter. 2016. Canalization of gene expression is a major signature of
regulatory cold adaptation in temperate *Drosophila melanogaster*. BMC Genomics 1–14.

Herman, M. A., O. D. Peroni, J. Villoria, M. R. Schön, N. A. Abumrad, M. Blüher, S. Klein, and B. B.
Kahn. 2012. A novel ChREBP isoform in adipose tissue regulates systemic glucose metabolism. Nature
484:333–338.

Hermisson, J., and G. P. Wagner. 2004. The population genetic theory of hidden variation and genetic
robustness. Genetics 168:2271–2284.

Hodgins-Davis, A., A. B. Adomas, J. Warringer, and J. P. Townsend. 2012. Abundant gene-by-
environment interactions in gene expression reaction norms to copper within *Saccharomyces cerevisiae*.
Genome Biol. Evol. 4:1061–1079.

Hodgins-Davis, A., and J. P. Townsend. 2009. Evolving gene expression: from G to E to G×E. Trends in
Ecol. Evol. 24:649–658.

Hoffmann, A. A., and S. W. McKechnie. 1991. Heritable variation in resource utilization and response in
a winery population of *Drosophila melanogaster*. 45:1000.

Huang, Y., and A. F. Agrawal. 2016. Experimental evolution of gene expression and plasticity in
alternative selective regimes. PLoS Genet. 12:e1006336.

Jakšić, A. M., and C. Schlötterer. 2016. The interplay of temperature and genotype on patterns of
alternative Splicing in *Drosophila melanogaster*. Genetics, 204:209-217.

Joshi, A., and J. N. Thompson. 1997. Adaptation and specialization in a two-resource environment in
Drosophila species. 51:846.

Kathiresan, S., O. Melander, C. Guiducci, A. Surti, N. P. Burtt, M. J. Rieder, G. M. Cooper, C. Roos, B.
F. Voight, A. S. Havulinna, B. Wahlstrand, T. Hedner, D. Corella, E. S. Tai, J. M. Ordovas, G. Berglund,
E. Vartiainen, P. Jousilahti, B. Hedblad, M.-R. Taskinen, C. Newton-Cheh, V. Salomaa, L. Peltonen, L.
Groop, D. M. Altshuler, and M. Orho-Melander. 2008. Six new loci associated with blood low-density
lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. Nature Genet.
40:189–197.

Kong, E. C., Allouche, L., Chapot, P. A., Vranizan, K., Moore M. S., Heberlein U., et al. 2010. Ethanol-
regulated genes that contribute to ethanol sensitivity and rapid tolerance in *Drosophila* . Alcohol Clin.
Exp. R 34: 302–316.

Kooner, J. S., J. C. Chambers, C. A. Aguilar-Salinas, D. A. Hinds, C. L. Hyde, G. R. Warnes, F. J. Gómez
Pérez, K. A. Frazer, P. Elliott, J. Scott, P. M. Milos, D. R. Cox, and J. F. Thompson. 2008. Genome-wide
scan identifies variation in MLXIPL associated with plasma triglycerides. Nature Genet. 40:149–151.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity and
genetic assimilation. J. Evol. Biol. 22:1435–1446.

Lee, Y. C. G., and H. H. Chang. 2013. The evolution and functional significance of nested gene structures
in *Drosophila melanogaster*. Genome Biol. Evol. 5:1978–1985.

Levine, M. T., M. L. Eckert, and D. J. Begun. 2011. Whole-genome expression plasticity across tropical
and temperate *Drosophila melanogaster* populations from Eastern Australia. Mol. Biol. Evol. 28:249–
256.

Li H (2015) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997.

Li, L., A. Li, K. Song, J. Meng, X. Guo, S. Li, C. Li, P. De Wit, H. Que, F. Wu, W. Wang, H. Qi, F. Xu,
R. Cong, B. Huang, Y. Li, T. Wang, X. Tang, S. Liu, B. Li, R. Shi, Y. Liu, C. Bu, C. Zhang, W. He, S.
Zhao, H. Li, S. Zhang, L. Zhang, and G. Zhang. 2018. Divergence and plasticity shape adaptive potential
of the Pacific oyster. Nat. Ecol. Evol. 2, 1751-1760.

Mathur, V., and P. S. Schmidt. 2016. Adaptive patterns of phenotypic plasticity in laboratory and field
environments in *Drosophila melanogaster*. Evolution 71:465–474.

Mattila, J., E. Havula, E. Suominen, M. Teesalu, I. Surakka, R. Hynynen, H. Kilpinen, J. Väänänen, I.
Hovatta, R. Käkelä, S. Ripatti, T. Sandmann, and V. Hietakangas. 2015. Mondo-Mlx mediates organismal
sugar sensing through the Gli-similar transcription factor *sugarbabe*. Cell Reports 13:350–364.

Matzkin, L. M. 2012. Population transcriptomics of cactus host shifts in *Drosophila mojavensis*. Mol.
Ecol. 21:2428–2439.

McCairns, R. J. S., and L. Bernatchez. 2009. Adaptive divergence between freshwater and marine
sticklebacks: insights into the role of phenotypic plasticity from an integrated analysis of candidate gene
expression. Evolution 64:1029–1047.

McKenzie, J. A., and S. W. McKechnie. 1979. A comparative study of resource utilization in natural
populations of *Drosophila melanogaster* and *D. simulans*. Oecologia 40:299-309.

Mercot, H., D. Defaye, P. Capy, E. Pla, and J. R. David. 1994. Alcohol tolerance, ADH activity, and
ecological niche of *Drosophila* species. 3:746–757.

Merkin, J., C. Russell, P. Chen, and C. B. Burge. 2012. Evolutionary dynamics of gene and isoform
regulation in mammalian tissues. Science 338:1593–1599.

Mi, H., X. Huang, A. Muruganujan, H. Tang, C. Mills, D. Kang, and P. D. Thomas. 2017. PANTHER
version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis
tool enhancements. Nucleic Acids Res. 45:D183–D189.

Miner, B. G., S. E. Sultan, S. G. Morgan, D. K. Padilla, and R. A. Relyea. 2005. Ecological consequences
of phenotypic plasticity. Trends in Ecol. Evol. 20:685–692.

Morozova, T. V., Huang, W., Pray V. A., Whitham, T., Anholt, R. R. H., Mackay, T. F. C. 2015.
Polymorphisms in early neurodevelopmental genes affect natural variation in alcohol sensitivity in adult
Drosophila. BMC Genomics: 1–16.

Morozova, T. V., Mackay, T. F. C., Anholt R. R. H. (2011). Transcriptional networks for alcohol
sensitivity in *Drosophila melanogaster*. *Genetics* 187: 1193–1205.

Morris, M. R. J., R. Richard, E. H. Leder, R. D. H. Barrett, N. Aubin-Horth, and S. M. Rogers. 2014.
Gene expression plasticity evolves in response to colonization of freshwater lakes in threespine
stickleback. Mol. Ecol. 23:3226–3240.

Murren, C. J., J. R. Auld, H. Callahan, C. K. Ghalambor, C. A. Handelsman, M. A. Heskel, J. G.
Kingsolver, H. J. Maclean, J. Masel, H. Maughan, D. W. Pfennig, R. A. Relyea, S. Seiter, E. Snell-Rood,

U. K. Steiner, and C. D. Schlichting. 2015. Constraints on the evolution of phenotypic plasticity: limits
and costs of phenotype and plasticity. Heredity 115:293–301.

Newell, N. R., F. N. New, J. E. Dalton, L. M. G. Genes, Genomes, 2016. 2016. Neurons that underlie
Drosophila melanogaster reproductive behaviors: detection of a large male-bias in gene expression in
fruitless-expressing neurons. G3 6:2455–2465.

Newton, P. M., K. Tully, T. McMahon, J. Connolly, J. Dadgar, S. N. Treistman, and R. O. Messing. 2004.
Chronic ethanol exposure induces an N-type calcium channel splice variant with altered channel kinetics.
FEBS Letters 579:671–676.

Oomizu, S., N. Boyadjieva, and D. K. Sarkar. 2003. Ethanol and estradiol modulate alternative splicing of
dopamine D2 receptor messenger RNA and abolish the inhibitory action of bromocriptine on prolactin
release from the pituitary gland. Alcohol Clin. Exp. R. 27:975–980.

Oostra, V., M. Saastamoinen, B. J. Zwaan, and C. W. Wheat. 2018. Strong phenotypic plasticity limits
potential for evolutionary responses to climate change. Nat. Comm. 9:1005.

Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden substrate. Nat.
Rev. Genet. 15:247–258.

Pajoro, A., E. Severing, G. C. Angenent, and R. G. H. Immink. 2017. Histone H3 lysine 36 methylation
affects temperature-induced alternative splicing and flowering in plants. Genome Biology 18: 1021.

Parsons, P. A., and G. E. Spence. 1981. Ethanol utilization: threshold differences among three *Drosophila*
species. Am. Nat. 117:568–571.

Parsons, P. A., and S. B. King. 1977. Ethanol: Larval discrimination between two *Drosophila* sibling
species. Experientia 33:898–899.

Pietrzykowski, A. Z., R. M. Friesen, G. E. Martin, S. I. Puig, C. L. Nowak, P. M. Wynne, H. T.
Siegelmann, and S. N. Treistman. 2008. Posttranscriptional regulation of BK channel splice variant
stability by miR-9 underlies neuroadaptation to alcohol. Neuron 59:274–287.

Price J, Harrison MC, Hammond RL et al. (2018) Alternative splicing associated with phenotypic
plasticity in the bumble bee *Bombus terrestris*. Mol. Ecol. 27:1036–1043.

Quinn, J. J., Q. C. Zhang, P. Georgiev, I. A. Ilik, A. Akhtar, and H. Y. Chang. 2016. Rapid evolutionary
turnover underlies conserved lncRNA-genome interactions. Genes Dev. 30:191–207.

Robinson, B. G., and N. S. Atkinson. 2013. Is alcoholism learned? Insights from the fruit fly. Curr. Opin.
Neurobiol. 23:529–534.

Robinson, B. W. 2013. Evolution of growth by genetic accommodation in Icelandic freshwater
stickleback. Proc. R. Soc. B 280:2197.

Rutherford, S. L. 2000. From genotype to phenotype: buffering mechanisms and the storage of genetic
information. BioEssays 22:1095–1105.

Sasabe, T., and S. Ishiura. 2010. Alcoholism and alternative splicing of candidate genes. IJERPH 7:1448–
1466.

Schlichting, C. D. 2008a. Hidden reaction norms, cryptic genetic variation, and evolvability. Ann. N. Y.
Acad. Sci. 1133:187–203.

Schlichting, C. D., and M. A. Wund. 2014. Phenotypic plasticity and epigenetic marking: an assessment
of evidence for genetic accommodation. Evolution 68:656–672.

Scoville, A. G., and M. E. Pfrender. 2010. Phenotypic plasticity facilitates recurrent rapid adaptation to
introduced predators. Proc. Nat. Acad. Sci. USA 107:4260–4263.

Shi, Y., Z. Su, H. Yang, W. Wang, G. Jin, G. He, A. N. Siddique, L. Zhang, A. Zhu, R. Xue, and C.
Zhang. 2019. Alternative splicing coupled to nonsense-mediated mRNA decay contributes to the high-
altitude adaptation of maca (*Lepidium meyenii*). Gene 694:7–18.

Signor, S. A., F. N. New, and S. Nuzhdin. 2017a. A large panel of *Drosophila simulans* reveals an
abundance of common variants. Genome Biol. Evol. 10:189–206.

Signor, S. A., M. Abbasi, P. Marjoram, and S. V. Nuzhdin. 2017b. Conservation of social effects (Ψ)
between two species of *Drosophila* despite reversal of sexual dimorphism. Ecol. Evol. 7:10031–10041.

Signor, S. A., M. Abbasi, P. Marjoram, and S. V. Nuzhdin. 2017c. Social effects for locomotion vary
between environments in *Drosophila melanogaster* females. Evolution 71:1765–1775.

Signor, S., and S. Nuzhdin. 2018. Dynamic changes in gene expression and alternative splicing mediate
the response to acute alcohol exposure in *Drosophila melanogaster*.

Singh P, Börger C, More H, Sturmbauer C 2017. The role of alternative splicing and differential gene

expression in cichlid adaptive radiation. Genome Biol. Evol. 9:2764–2781.

Thomson, M. S., J. W. Jacobson, and C. C. Laurie. 1991. Comparison of alcohol dehydrogenase
expression in *Drosophila melanogaster* and *D. simulans*. Mol. Biol. Evol. 8:31–48.

Ulitsky, I. 2016. Evolution to the rescue: using comparative genomics to understand long non-coding
RNAs. Nat. Rev. Genet. 17:601–614.

Ulitsky, I., A. Shkumatava, C. H. Jan, H. Sive, and D. P. Bartel. 2011. Conserved function of lincRNAs
in vertebrate embryonic development despite rapid sequence evolution. Cell 147:1537–1550.

Valadkhan, S. and Valencia-Hipolito, A. 2016. lncRNAs and stress response. Curr. Topics Microbiol.
Immunol. 394:203-236.

Velotta, J. P., C. M. Ivy, C. J. Wolf, G. R. Scott, and Z. A. Cheviron. 2018. Maladaptive phenotypic
plasticity in cardiac muscle growth is suppressed in high-altitude deer mice. Evolution 72:2712–2727.

Via, S., and R. Lande. 1985. Genotype-environment interaction and the evolution of phenotypic plasticity.
Evolution 39:505–522.

Via, S., R. Gomulkiewicz, G. De Jong, S. M. Scheiner, C. D. Schlichting, and P. H. Van Tienderen. 1995.
Adaptive phenotypic plasticity: consensus and controversy. Trends in Ecol. Evol. 10:212–217.

Waddington, C. H. 1953. Genetic assimilation of an acquired character. Evolution 7:118-126.

Wang, G., L. Weng, M. Li, and H. Xiao. 2017. Response of gene expression and alternative splicing to
distinct growth environments in tomato. IJMS 18:475–19.

West-Eberhard, M. J. 2005. Developmental plasticity and the origin of species differences. Proc. Nat.
Acad. Sci. USA 102:6543–6549.

West-Eberhard, M. J. 2003. Developmental plasticity and evolution. Oxford Univ. Press, New York.

Yampolsky, L. Y., G. V. Glazko, and J. D. Fry. 2012. Evolution of gene expression and expression
plasticity in long-term experimental populations of *Drosophila melanogaster* maintained under constant
and variable ethanol stress. Mol. Ecol. 21:4287–4299.

Yang, H.-P., and S. V. Nuzhdin. 2003. Fitness costs of *Doc* expression are insufficient to stabilize its
copy number in *Drosophila melanogaster*. Mol. Biol. Evol. 20:800–804.

Zaharieva, E., J. K. Chipman, and M. Soller. 2012. Alternative splicing interference by xenobiotics.
Toxicology 296:1–12.

Zhu, J., and J. D. Fry. 2015. Preference for ethanol in feeding and oviposition in temperate and tropical
populations of *Drosophila melanogaster*. Entomol. Exp. Appl. 155:64-70.

721
722
723

D. simulans

	G	ETOH	G x ETOH	ETOH x T	G x ETOH x T	10	20	30
Exons	1994	76	387	24	1158	9	3	29
Exonic Regions	406	23	98	7	299	3	1	4
Fragments	608	18	81	6	225	4	2	6

D. melanogaster*

	G	ETOH	G x ETOH	ETOH x T	G x ETOH x T	10	20	30
Exons	1445	15	3	22	0	2	1	46
Exonic Regions	631	13	0	8	2	1	4	24
Exon Fragments	1135	21	0	12	4	3	5	20

*reproduced from Signor and Nuzhdin 2018

725 **Table 1**: A summary of the exons, exonic regions, and exon fragments that are significantly differently
726 expressed for each component of variance. Abbreviations are as follows: (G) Genotype, (ETOH)
727 Ethanol, and (T) Time. Ethanol in this case is the environment.

D. simulans

	Biological Process		Cellular Component		Molecular Function	
G x ETOH			axonemal dynein complex	3×10^{-3}	dynein light chain binding	3.3×10^{-3}
G x ETOH x T	sensory perception of taste	6.6×10^{-3}	nucleplasm	6.5×10^{-3}	RNA binding	7.8×10^{-3}
			cytoskeletal part	8.4×10^{-3}		
	detection of chemical stimulus	8.2×10^{-3}	ciliary part	4.6×10^{-3}		

730 **Table 2**: A summary of the significant GO enrichment terms for *D. simulans*. The abbreviations are the
731 same as in Table 1. *p*-values are listed adjacent to the significant term, the significance cutoff is $p < .01$.
732
733

Figure 1: **A**. An illustration of the classification scheme for exons, exonic regions, and exon fragments. Exons either do not overlap other exons in different isoforms (exons, shown in blue), or are fused regions consisting of a set of overlapping exons (exonic regions, shown in shades of orange and red). Exonic regions can be decomposed into exon fragments, depending upon their overlap between different isoforms. **B.** An example of the information about gene expression and isoform abundance that can be inferred from measures of differential exon abundance. *cabut* is a relatively simple example, having few isoforms and exons, and yet it is still a complicated inter-species comparison given that *D. melanogaster* has more annotated isoforms that *D. simulans*. Below each gene model the frequency of different exons and exon fragments in each environment is shown, with the most frequent in dark blue. In *D. simulans* the first of two isoforms shown has an exon fragment that is more abundant in the presence of ethanol, suggesting that that isoform is more abundant. In *D. melanogaster* only the second of the three isoforms has a unique fragment, which is also more abundant in ethanol and suggests differential isoform usage. **C.** The number of exons or exonic regions that are significant for each component of variance in *D. melanogaster* and *D. simulans*. Many more exons and exonic regions are significant for *D. simulans* than *D. melanogaster*, but only for components of variance that contain an interaction with genotype

757 **Figure 2: A.** An illustration of a change in the relative abundance of the gene *Drat*. Only the example of
758 *D. simulans* is shown. With and without ethanol different exons are the most abundant compared to the
759 other exons within the gene. As one of these exons is not shared among all isoforms, this suggests a
760 change in isoform usage between environments. **B.** The number of genes that have a significant change in
761 relative abundance for each component of variance in *D. melanogaster* and *D. simulans*.