Infection of the *Biomphalaria glabrata vector snail* by *Schistosoma mansoni*

parasites drives snail microbiota dysbiosis.

Anaïs Portet [1], Eve Toulza [1], Ana Lokmer [2], Camille Huot [1], David Duval [1], Richard Galinier [1] and

Benjamin Gourbal [1, §]

[1] IHPE, Univ. Montpellier, CNRS, Ifremer, Univ. Perpignan Via Domitia, Perpignan France

[2] Department Coastal Ecology, Wadden Sea Station Sylt, Alfred Wegener Institute, Helmholtz

Centre for Polar and Marine Research, List/Sylt, Germany.

AP: ap2133@cam.ac.uk, ET: eve.toulza@univ-perp.fr, AL: Ana.Lokmer@awi.de, CH:

camillehuot86@gmail.com, DD: david.duval@univ-perp.fr, RG: richard.galinier@univ-perp.fr,

BG: benjamin.gourbal@univ-perp.fr

[§] Corresponding author :

Benjamin GOURBAL

UMR5244 IHPE, Université de Perpignan

58 avenue Paul Alduy, 66860 Perpignan

benjamin.gourbal@univ-perp.fr

Running title: Biomphalaria glabrata snail microbiota dysbiosis

Summary

Host-associated microbiota can affect the fitness of its host in a number of ways, including the

modification of host-parasite interactions and thus the outcome of disease. *Biomphalaria*

glabrata is the vector snail of the trematode *Schistosoma mansoni*, the agent of human

schistosomiasis, causing hundreds of thousands of deaths every year. Here, we present the first

study of the snail bacterial microbiota in response to *Schistosoma* infection. To examine the

interplay between *B. glabrata, S. mansoni* and snail microbiota, snails were infected and the

microbiota composition was analysed by massive 16S rDNA amplicon sequencing approach. We

characterized the *Biomphalaria* bacterial microbiota at the individual level in both naïve and

infected snails. Sympatric and allopatric strains of parasites were used for infections and re-

infections to analyse the modification or dysbiosis of snail microbiota in different host-parasite

co-evolutionary contexts. Concomitantly, using RNAseq data, we investigated the link between

bacterial microbiota dysbiosis and snail anti-microbial peptide immune response. This work

paves the way for a better understanding of snail/schistosome interaction, and would have

critical consequences in terms of snail control strategies for fighting schistosomiasis disease in

the field.

Key words: microbiota; bacteria; *Biomphalaria* snail; *Schistosoma* infection; immune response;

dysbiosis

Introduction

Several studies have shown that the microbiota interacts with pathogens and/or host immunity in invertebrates. For example, the midgut microbiota of the mosquito *Aedes sp.* elicits a basal immune activity of the immune system (Ramirez et al., 2012) mostly by activating Toll pathways (Xi et al., 2008). In *Anopheles gambiae*, some members of its microbiota can limit malaria transmission, by inducing a wide antimicrobial immune response (Gendrin, 2017). More precisely, bacteria of the genus *Enterobacter*, found in the mosquito microbiota, were shown to produce reactive oxygen species (ROS) that inhibit *Plasmodium* development (Cirimotich et al., 2011). Furthermore, the endosymbiotic bacteria *Wolbachia* induce expression of immune genes, like TEP1 (Thioester-containing protein 1), LRIM1 (Leucine-Rich Immune Molecule 1) or defensin 1 (Joshi et al., 2017). In *Drosophila*, the bacterial microbiota is necessary to produce the Pvf2, a PDGF/VEGF-like growth factor that restricts enteric viral infection (Sansone et al., 2015). The microbiota may also play a role in immune priming. For example, in *Anopheles gambiae*, hemocyte priming during the *Plasmodium* infection is naturally induced by the gut microbiota, whose absence results in more severe infections and re-infections (Rodrigues et al., 2010). Similarly, the gut microbiota is necessary for immune priming in *Tribolium castaneum* and efficient response against *Bacillus thuringiensis for Tenebrionis* (Futo et al., 2015). Finally, host immunity can influence the tolerance or control the microbiota. In the tick *Ixodes scapularis*, the protein PIXR, secreted by the tick's gut, inhibits the bacterial biofilm formation and supports gut eubiosis (healthy gut microbiota), its inactivation leading to dysbiosis. This change in gut microbiota facilitates the colonization by *Borrelia burgdorferi*, the Lyme disease agent (Narasimhan et al., 2017). In *Drosophila*, for example, the intestinal homeobox gene Caudal regulates the commensal-gut mutualism by repressing nuclear factor kappa B-

dependent antimicrobial peptide genes (Ryu et al., 2008). In Hydra, the innate immune sensors

and effectors protect not only against pathogens but also control microbiota homeostasis

(Augustin et al., 2010). Moreover, host specific antimicrobial peptides (Fraune et al., 2010;

Franzenburg et al., 2012) and a rich repertoire of pattern recognition receptors (Bosch, 2013)

are involved in maintaining homeostasis between the host and the resident microbiota.

Similarly, bactericidal permeability-increasing proteins (BPIs) shape bacterial communities in

the light organ of the squid *Euprymna scolopes* and prevent their spill over to other tissues

(Chen et al., 2017).

Biomphalaria glabrata is a freshwater snail (Lophotrochozoa, Planorbidae), living in inter-

tropical regions of Latin America, in rivers, ponds, waterways and other freshwater

environments. *B. glabrata* snails have important medical and epidemiological impacts due to

their role as the main vector of *Schistosoma mansoni* (Lophotrochozoa, Platyhelminthes,

Trematoda), the agent of intestinal schistosomiasis. Schistosomiasis is the second most

widespread human parasitic disease after malaria, affecting over 200 million people worldwide

and causing 200 000 deaths annually (WHO, 2002). The *Schistosoma* adult parasites mate in the

human host venous system. Female worms produce eggs that cross endothelial mesenteric

vessels and intestinal epithelium to reach faeces and finally the aquatic environment. Once in

the water, eggs hatch and release miracidia, the free-living snail-infective parasite stage. At the

following step of the life cycle, the miracidium needs to infect the freshwater snail *B. glabrata*.

Intensive asexual multiplication in the snail tissues leads to the continuous production of

hundreds of generations of cercariae, the free-living human-infective parasite stage.

At present, there is no effective vaccine against schistosomiasis and the treatment relies on a

single chemotherapeutic treatment, the Praziquantel (Doenhoff et al., 2009), against which

resistance has been observed (Fallon and Doenhoff, 1994). Molluscicides have been also used

to impair *Schistosoma* transmission in the field. However, the dramatic effects of molluscicides

on natural environments prompt us to seek new ways to prevent and / or control this disease in

the field (Tennessen et al., 2015). We propose that a better understanding of ecological

interactions between, *Schistosoma mansoni, Biomphalaria glabrata* snails and its associated

microbiota might represent an interesting approach in search for alternative control strategies.

In this context, we have conducted numerous studies on immunological interactions

between the snail and the parasite (Mitta et al., 2005; Bouchut et al., 2008; Moné et al., 2011;

Coustau et al., 2015; Galinier et al., 2017; Mitta et al., 2017). We have demonstrated that the

nature of the snail immune response depends on the strain/species of host or of parasite

considered. Depending on both the host and the parasite intrinsic capacities, parasites either

develop normally in snail tissue (compatible interaction) or are encapsulated by hemocytes (the

snail immune cells) (incompatible interaction) (Mone et al., 2010; Mitta et al., 2012; Coustau et

al., 2015) (Fig. 1). Depending on the ecological context, totally different immunobiological

interactions have been described in *B. glabrata/S. mansoni* interactions. If immunosuppression

occurs during a sympatric interaction, an activation of the immune response is observed during

an allopatric interaction (Portet et al., 2019). Moreover, following primo-infection and

successive challenges, a shift from a cellular immune response toward a humoral immune

response has been described (Pinaud et al., 2016) (Fig. 1).

However, in the present host-parasite model, only few studies examined the bacterial

microbiota of *Biomphalaria glabrata* so far (Ducklow et al., 1979; Silva et al., 2013). The study

of aerobic heterotrophic cultivable flora of 200 snails (Ducklow et al., 1979) revealed five

predominant bacterial genera including *Pseudomonas, Acinetobacter, Aeromonas, Vibrio* and

Enterobacter. In a more recent study, the *Biomphalaria glabrata* bacterial microbiota has been

characterized using 16S rRNA sequences (Silva et al., 2013) of cultured isolates, identifying six

additional bacteria genera: *Citrobacter, Cupriavidus, Rhizobium, Stenotrophomonas, Klebsiella*

and *Sphingomonas*. In addition, diversity and composition of bacterial microbiota differed

between resistant and susceptible *Biomphalaria* phenotypes, with differences in relative

abundances of *Gemmatimonas aurantiaca* and *Micavibrio aeruginosavorus*. This observation

supports the potential link between the resistance against *Schistosoma* parasite and the

structure of snail-associated microbiota (Allan et al., 2018). However, no studies have examined

changes in bacterial microbiota of *Biomphalaria* snails by culture-independent methods nor

following Schistosoma infections so far.

Taking into account all the peculiarities of this system, the aim of the present study was to

investigate variation of microbiota in snail populations as well as to assess the influence of

varying immune response against *Schistosoma* on the host microbiota composition and

dynamics. To reach this goal, we first characterized the bacterial microbiota of naive

Biomphalaria glabrata snails using 16S rDNA metabarcoding. Then, we analysed changes in

microbiota composition following, (i) primo-infections with sympatric and allopatric *S. mansoni*

parasite isolates displaying the same prevalence and intensity phenotypes and, (ii) subsequent

secondary challenges with homologous or heterologous parasite strains. As snail immune

response strongly differs between such diverse snail/parasite combinations, we concomitantly

analysed the snail immune response using a massive sequencing transcriptomic approach

(RNAseq) in order to link it with the observed changes in the composition and diversity of the

microbiota.

Results

Characterization of healthy B. glabrata *microbiota*

To examine the stability of naive BgBRE snail microbiota, we first inspected the microbiota

diversity and composition in the control naive snails at day 0 and 25 of the experiment. We

found no significant differences between the B0 and B25 snails in any of the alpha diversity

indices (Table S3, Table S4).

Naive snails showed little inter-individual variability and a stable composition at the

phylum level over time (Fig. 2A, Table S3, Table S4), with Proteobacteria, Bacteroidetes,

Cyanobacteria and Planctomycetes phyla being the most represented (Fig 2A). Moreover, the

bacterial microbiota of naïve BgBRE snails displayed a considerable temporal and inter-

individual stability at the family level (Fig. 2B). The stability of naïve snail microbiota

composition over time (Fig. 2) could be related to the laboratory rearing conditions and

laboratory environmental abiotic variables (water composition, temperature, pH, food) that

were tidily controlled and that did thus not result in changes in host snail physiology or

metabolism.

In terms of composition, we observed that 67 (69%) and 86 (89%) out of 97 identified

families were shared by all individuals of B0 or B25 naive snails respectively (Fig. 2B). Those

results were used to determine the core microbiota. We defined core microbiota as the families

that were present in 100% of the naive snails. Applying this definition, we identified 62 out of

97 families found in all naive individual snails (B0 and B25 altogether), and thus constituting the

B. glabrata core-microbiota (Table S7 and S8).

Microbiota dynamics following B. glabrata *infections by* S. mansoni

After establishing the stability of naive *Biomphalaria* bacterial microbiota, we investigated

whether *Schistosoma mansoni* infections affected the microbiota structure and dynamics.

To investigate the influence of parasite infections on the bacterial microbiota, we analysed

microbiota dynamics following sympatric or allopatric primo-infections and homologous or

heterologous challenges (Fig. 1).

We did not observe any significant changes in alpha diversity during the course of primo-

infection compared to naive snails, excepted a decrease in the Shannon's H diversity index at

the fourth day after sympatric infection (Fig. 3, Table S3, Mann-Whitney U test: Table S4).

Conversely, all indices changed significantly following homologous or heterologous challenges

(Fig. 3, Table S3, Table S4). Indeed, the observed species richness (Mann-Whitney U test,

p=0.002), the Chao 1 richness index (Mann-Whitney U test, p=0.006), Shannon diversity index

(p=0.0004) and Pielou evenness index (Mann-Whitney U test, p=0.009) were significantly

reduced following homologous or heterologous challenges compared to naive and primo-

infected snails (Fig. 3, Table S3, Table S4). However, the observed drop in alpha diversity

disappeared by the fourth day after heterologous challenge (Fig. 3, Table S4). In addition, alpha

diversity was mainly affected by the challenge, regardless of its type (homologous or

heterologous). The primo-infection did not significantly affect alpha diversity except for the

sympatric combination at day 4 after infection (Table S4).

Regarding the beta diversity, Principal Coordinate Analysis (PCoA) of Bray-Curtis dissimilarities

revealed that BB25 samples grouped together with the naive snail samples (B0 and B25) and

were separated from the infected-snail samples along both axes (Fig. 4A, Table S5). However,

the BB25 samples were significantly different from the naive snail samples (Table S5). The fact

that, BB25 grouped with naive snails rather than with infected-snail samples suggests that the

snail microbiota is resilient to infection, with a tendency to recover between day 4 and day 25

after the primo-infection (Fig. 4A). Analysis of Bray-Curtis dissimilarity index revealed a

significant difference between naive and primo-infected samples (p=0.001) and also between

naive and challenged samples (p=0.001) (Table S5). Concerning the infected-snail samples, all

experimental samples were significantly different from each other with the exception of BB1

versus BB4 and BBB1 versus BBV1 (Table S5).

In addition, the second PcoA axis separated the samples according to the course of infection

(i.e. the day 1 from the day 4 samples) (Fig. 4A), whereas the day 1 after-challenge (BBB1,

BBV1), the day 1 after-primo-infection (BB1, BV1) and all of the day 4 infection samples (BB4,

BV4, BBB4, BBV4) were separated along the first axis (Fig. 4A). Interestingly, day 1 after-

challenge samples (BBB1, BBV1) were more different from the naive microbial communities

than the day 1 primo-infection samples (BB1, BV1). Moreover, the second axis separated the

sympatric primo-infection (BB1) from the allopatric primo-infection (BV1) (Fig. 4A). Finally, all

day 4 samples grouped together reflecting the similarity between these samples regardless of

experimental infection conditions (Fig. 4A). Even if the day of infection appears as the main

explaining factor, these results indicate that the snail microbiota response to the infection also

depends on the type of infection (sympatric vs. allopatric or primo-infection vs. challenge) and

suggest the existence of predictable dynamics.

We further investigated the microbial community dynamics following infection, and we

observed some consistent changes in response to experimental infection type (Fig. 4B). The

relative abundance of Tenericutes increased after infection, particularly at day 1 after challenge

(BB25 vs BBB1, student T test P = 0.0114; BB25 vs BBV1 student T test P = 0.0209). Increase in

Verrucomicrobia following allopatric primo-infection (B0/B25 vs BV4, student T test P = 6.53 e-

06) and homologous / heterologous challenge (B0/B25 vs BBB4/BBV4, student T test P =

0.0012) was most prominent at day 4 after infection (Fig. 4B). Interestingly, high relative

abundance of Verrucomicrobia in BB25 represents the main difference with naive snail

microbial community composition (B0/B25 vs BB25, student T test P = 1.83 e -05) (Fig. 4B). We

further observed decrease in Planctomycetes following sympatric primo-infection (B0/B25 vs

BB1/BB4, student T test P = 5.7 e-07) and a decrease in Cyanobacteria following allopatric

primo-infection (B0/B25 vs BV1/BV4, student T test P = 0.00254). Interestingly, the relative

abundance of both phyla decreased following the challenge regardless of the challenge type

(Planctomycetes B0/B25/BB25 vs BBB1/BBB4/BBV1/BBV4, student T test P = 0.0017 and

Cyanobacteria B0/B25/BB25 vs BBB1/BBB4/BBV1/BBV4, student T test P = 0.000568) (Fig. 4B).

Bacteroidetes decreased 25 days after sympatric primo-infection (B0/B25 vs BB25, student T

test P = 0.0467) but increased 4 days after heterologous challenge (BB25 vs BBV4, student T

test P = 0.00577). To conclude, depending on the phylum, we observed the largest

modifications in microbiota composition at day 1 or 4 after primo-infections and challenges.

Whereas the microbiota responded differently to sympatric and allopatric primo-infections, we

found no significant differences between the homologous or heterologous challenge (see BBB1

vs BBV1, Fig. 4) (Table S5).

Finally, we examined the core microbiota dynamics during infection. Similarly, to the complete

microbiota, the core-microbiota, consisting of 62 families, was affected by the type of infection

(naive vs primo-infection (p=0.003) and naive *vs* challenge (p=0.003)), by the time of infection

(early (1 day) vs late (4 days)) (p=0.006) and also by the strain of parasite used for primo-

infection (SmBRE vs SmVEN) (p=0.016) (Table S6). Principal Coordinate Analysis (PCoA) based

on Bray-Curtis dissimilarities between the core microbiotas (Fig. S1) yielded very similar results

to those based on the entire dataset (Fig. 4A).

In addition, we observed that 69.4% (43) of the core microbiota families were significantly

affected by infection. Among those, 6.5% (4 families) were affected regardless of the infection

type (Table S7). Interestingly, these families belong to the most abundant ones (Table S8).

Nineteen families (30.6%) were never affected by infection (Table S7). Those families belong to

the less represented ones, with the exception of *Xanthomonadaceae,* which was the 7[th] most

represented family (see Table S8). Further 32.6% (14 families) of the core microbiota responded

solely to the primo-infection and 4.7% (2 families) to the challenge (Table S7). Similarly, 20.9%

(9 families) of the core microbiota was affected early after infection (1 day) and 7% later (3

families, day 4). Finally, 23.3% (10 families) were affected by the SmBRE infection and 9.3% (4

families) by the SmVEN infection (Table S7).

Link between the microbiota dysbiosis and B. glabrata *antimicrobial immune response*

The expression level of transcripts encoding antimicrobial peptides and antimicrobial proteins

(AMP) was investigated following sympatric or allopatric primo-infection and homologous or

heterologous challenges using RNAseq data (Fig. 5 and Fig. S2). Two achacin genes, 5

lipopolysaccharide-binding protein / bactericidal permeability-increasing protein (LBP/PBI)

genes, and 5 biomphamacin genes were selected based on the *Biomphalaria glabrata* genome

annotation (Adema et al., 2017). In comparison with the naive snails, the LBP/BPI 3.1 and 3.2

were over-expressed at day 25 after infection and following the challenges, while all other

genes of this family were under-expressed in all conditions (Fig. S2). The achacins were under-

expressed following primo-infection in the sympatric combination only, and at day 25 after

primo-infection and following both challenges (Fig. S2). No changes in expression were

observed following allopatric primo-infection compared to the naive snails (Fig. S2). Finally, the

AMP biomphamacins 1, 4, 5 and 6 were over-expressed throughout the infection stages, except

for BV1 and BB25 (Fig.6). The biomphamacin 3 was mainly under-expressed except at BV4 and

BBB, where no differential expressions have been detected compared with naive snails (Fig. 5).

Based on our observations of microbial community shifts, changes in expression of

antimicrobial molecules were expected to occur following primo-infections and challenges and

not for BB25, where resilience of microbial communities has been observed (Fig. 4 and Fig. S1).

In this context, the LBP/BPI and achacin seem not to be linked with dysbiosis, as both remained

highly down regulated even at BB25 when communities had already recovered (Fig. S2). All

biomphamacins, except the biomphamacin 3, were over-expressed after infection, from day 1

in BB infection and day 4 in BV infection and for homologous or heterologous challenges (BBB

and BBV). Finally, only a subset of biomphamacins (1, 4, 5, 6) changed in the course of infection

but were not differentially expressed at BB25 compared with the naive snails (Fig. 5),

suggesting their possible link with dysbiosis.

Discussion

Fine-tuned interactions between microbiota, host immunity and pathogens have been

observed in many vertebrate and invertebrate models (Rodrigues et al., 2010; Ramirez et al.,

2012; Futo et al., 2015; Sansone et al., 2015; Chen et al., 2017). Following an infection by a

pathogen, a host displays an innate or adaptive immune response against the intruder. The

activation of such immune response may in turn affect the microbial community structure.

Herein, we investigated the interactions between the host immune system, parasite and the

bacterial microbiota in an invertebrate model - the gastropod snail *Biomphalaria glabrata and*

its trematode parasite *Schistosoma mansoni*. Depending on the past evolutionary history

between snails and schistosomes, different immune responses against *S. mansoni* have been

observed. We showed recently that in a sympatric interaction, the parasite that coevolved with

its host induced a strong immunosuppression, whereas allopatric interaction resulted in a

strong host cellular immune response (Portet et al., 2019). Moreover, a cellular immune

response was observed following primo-infection, but a humoral immune response was

observed following homologous or heterologous challenges (Pinaud et al., 2016). So using

appropriate parasite-host combinations, we have the opportunity to modulate the host

immune response type and to test changes in bacterial community composition and diversity

(dysbiosis). First, studying the global bacterial microbiota community of *Biomphalaria* snails, we

showed that the bacterial alpha diversity was not modified following primo-infection whatever

the parasite strain or the time point of infection (Fig. 3, S3 Table). A decrease in alpha diversity

is observed exclusively following challenge infections, as reflected by multiple indices (Fig. 3, S3

Table). Conversely, primo-infection and challenge strongly affected the bacterial OTU

composition (Fig. 4). Moreover, differences in immunobiological interactions

(immunosuppression, immune cellular response or immune humoral response) resulted in

different microbiota dysbiosis reflected by specific changes in the snail microbial community

(Fig. 4). Interestingly, homologous and heterologous challenges activated similar humoral

immune response (Pinaud et al., 2016; Portet et al., 2019) that resulted in a similar change of

microbiota alpha diversity and composition (Fig. 3, S3 Table, Fig. 4). Then, four days after

infection, regardless of its type, the microbiota was still very different from the one in the naïve

snails, but the differences between the primo-infection and challenge disappeared, as apparent

from the grouping of BB4, BV4, BBB4 and BBV4 in the PCoA (see Fig. 4). Furthermore, most of

the OTUs affected by the primo-infection, returned to their initial state by the day 25, indicating

that the snail microbiota is resilient against the infection (Fig. 4). However, some differences

persisted: the Verrucomicrobia phylum remained highly represented in the infected snails at

day 25 after primo-infection (Fig. 4B). Interestingly, some Verrucomicrobia species have been

recently proposed as a hallmark of healthy gut due to its anti-inflammatory and immune-

stimulant properties and its ability to improve gut barrier function (Fujio-Vejar et al., 2017).

High abundance of Verrucomicrobia in the recovered snails could thus potentially reflect their

role in community restoration (Fig. 4). Moreover, the expansion of a Verrucomicrobia species

*Akkermansia muciniphila*has been demonstrated in the gut of *S. mansoni*-infected mice,

suggesting a potentially functional role of Verrucomicrobia in *Schistosoma* infection processes

in both final and intermediate hosts (Jenkins et al., 2018). We also paid a particular attention to

the core-microbiota. Given the various ways to define core microbiota, we considered

exclusively the persistent occurrence in the bacterial community in naive *Biomphalaria glabrata*

snails (Astudillo-García et al., 2017). Similarly, to the entire microbiota, the core microbiota was

affected by the type of infection (naive *vs* challenge), by the time of infection (day 1 vs day 4) as

well as by the parasite strain (SmBRE vs SmVEN) (Table S6). Core microbiota seemed to be

affected by immunosuppression or the activation of the immune response in a similar way than

the whole microbiota (Fig. S1 and Fig. 4A). Understanding the shifts in core microbiota

following infection is important as the long-term stability and persistent occurrence of

beneficial microbes and their associated functions may contribute to host health and well-being

to maintain host functionality and fitness toward changing ecological environment or

environmental stress (Nyholm and McFall-Ngai, 2004; Lozupone et al., 2012; McFall-Ngai et al.,

2013).

325   Given that infections affect the total and core microbiota structure, we explored the

antimicrobial immune response following infection to find potential molecular mechanisms

involved in the observed microbiota dysbiosis (Fig. 5 and Fig. S2). We analysed antimicrobial

peptides (AMP) and antimicrobial proteins and our results suggest that only the AMP belonging

to biomphamacin family may potentially affect the microbial communities in response to

Schistosoma infection. Indeed, even though antimicrobial families can be involved in complex

process of regulation or of control of microbiota communities, solely the biomphamacins AMP

family members specifically displayed an expression pattern that can be linked with the

observed dysbiosis of *Biomphalaria* bacterial communities (Fig. 5). Even if AMPs is often

considered as the main immune pathway responsible for microbiota regulation (Onchuru and

Kaltenpoth, 2019), an overall modification of the immune system may also be considered as a

cause of dysbiosis. The potential role of immune recognition or ROS/NOS pathways as to be

considered (Wang et al., 2014; Budachetri and Karim, 2015; Yang et al., 2015; Yang et al., 2016)

and this will deserve further investigations in the present biological experimental model.

It has been shown that the immune system is a key determinant of host-associated bacterial

communities in many biological systems. Two mechanisms have been proposed to explain host-

microbiota interactions through crosstalk with the host innate immune system. The first one

proposes that the host immune system exerts constant pressure on the microbiota in order to

maintain homeostasis (Hooper et al., 2012), the host immune system can thus control the

composition of the resident microbiota (Zhang et al., 2015). According to this hypothesis, any

changes in host immune response to infection would potentially affect the resident microbiota

diversity and composition. The second mechanism proposes that the immune system would be

tolerant to weak and continuous antigenic immune stimulations experienced during a lifespan

(Pradeu and Eric, 2014), and thus that host immune system would not exert any pressure on

the resident microbiota. In this case, any changes in host immune response to infection would

potentially not affect the resident microbiota diversity and composition.

Interestingly, numerous studies demonstrate a direct control of microbial communities by the

host immune system. As an example, species-specific antimicrobial peptides can shape species-

specific bacterial associations in *Hydra* (Franzenburg et al., 2013). Other immune pathways

have been also demonstrated to regulate or control the microbiota communities, like the

intestinal homeo-box gene Caudal in *Drosophila* (Ryu et al., 2008), or neuropeptides with

antibacterial activity which are secreted to shape the microbiome on the body surface of *Hydra*

(Augustin et al., 2017), even host lectins were demonstrated to stabilize microbiota

communities (Dinh et al., 2018).

Thus, if biotic stress (i.e. an infection) modifies the expression of antimicrobial peptides or

other immune related pathways, an effect on the microbial communities can be expected. Our

results indicate that bacterial communities could indeed be shaped by the immune system of *B.*

glabrata. Based on the hypothesis proposed by Hooper and collaborators (Hooper and

Macpherson, 2010; Hooper et al., 2012; Dinh et al., 2018), it seems that the immune system of

B. glabrata snails maintains or controls the microbial communities permanently. Therefore,

following an infection, the immune system is diverted from its function of managing the

microbiota and consequently the bacterial communities evade its regulation, resulting in

changes of their composition and diversity. When the immune response returns to a basal level,

the microbiota is again under-control and the microbiota shows its resilience by returning to its

"healthy" state (Hooper and Macpherson, 2010; Hooper et al., 2012), as observed in the

present study (Fig. 4). In other words, the immune system is likely no longer able to maintain

the microbiota homeostasis after infection (resulting in dysbiosis), which, may in turn affect

host homeostasis or fitness (Contijoch et al., 2019; Warne et al., 2019) (Contijoch, Britton et al.

2019, Warne, Kirschman et al. 2019).

Changes in microbiota composition may result from shifts in the abundance of specific bacterial

groups participating in anti-pathogen response, or just be a collateral effect of the immune

response activation against metazoan parasite infection. This question will deserve further

investigation, by testing *S. mansoni* prevalence and intensity in experimental infections of

Biomphalaria glabrata snails following antibiotic treatment or microbiota transplantation.

In order to fully understand Schistosomiasis transmission and to develop new ways to

control the expansion of this widespread human parasitic disease in the field, it will be crucial

to determine if the snail-associated bacterial communities affect the parasite transmission. For

example, expanding knowledge on *Biomphalaria* snail microbiota is an essential step for

developing paratransgenetic solutions to the spread of Schistosomiasis, involving the use of

transgenic bacteria expressing foreign gene products (i.e., schistosomicidal compounds) that

can reduce host competence and block pathogen development or transmission when

introduced into the microbiota of vector snail field populations (Aksoy et al., 2008; Coutinho-

Abreu et al., 2010; Gilbert et al., 2016).

Experimental procedures

Ethical statements

Our laboratory holds permit # A66040 for experiments on animals, which was obtained from

the French Ministry of Agriculture and Fisheries and the French Ministry of National Education,

Research, and Technology. The housing, breeding and care of the utilized animals followed the

ethical requirements of our country. The experimenter possesses an official certificate for

animal experimentation from both of the above-listed French ministries (Decree # 87–848,

October 19, 1987). The various protocols used in this study have been approved by the French

veterinary agency of the DRAAF Languedoc-Roussillon (Direction Régionale de l'Alimentation,

de l'Agriculture et de la Forêt), Montpellier, France (authorization # 007083).

Biological material

In this study, we used a single strain of the host snail, the albino Brazilian strain of *Biomphalaria*

glabrata (BgBRE) and two strains of the trematode parasite *Schistosoma mansoni*: a Brazilian

strain (SmBRE, for sympatric infection) and a Venezuelan strain (SmVEN, for allopatric

infection). BgBRE and SmBRE strains originate from the locality of Recife, Brazil; the SmVEN

parasite strain was recovered from the locality of Guacara, Venezuela. All host and parasite

strains were maintained in the laboratory on their respective sympatric snail hosts (SmBRE on

BgBRE and SmVEN on BgVEN). The snails were reared at constant temperature of 26°C and fed

only with washed salad every 3 days.

Experimental Infections (Fig. 1)

In order to decipher the inter-individual structure of snail microbiota as well as investigate the

influence of different snail immune stimulations on its microbiota structure and dynamics, we

applied a two-step experimental infection protocol (Fig. 1).

Briefly, BgBRE snails were primo-infected with one of the two parasite strains (SmBRE

sympatric or SmVEN allopatric). The snails were then sampled 1 and 4 days following primo-

infection by both parasite strains and 25 days after primo-infection for SmBRE infection

condition. Then, the BgBRE snails primo-infected with SmBRE were challenged (25 days after

the primo-infection) with SmBRE or SmVEN parasite strains for homologous or heterologous

challenges respectively. The challenged snails were then sampled 1 and 4 days following

challenge (Fig. 1). For all experimental infections, the snails were individually exposed for 12h

to 10 miracidia in 5mL of pond water, thereafter snails were returned to the water tanks and

separated according to the parasite strain and infection conditions.

Infections and sampling for microbiota analysis

We performed a primo-infection of 63 BgBRE snails with either SmBRE (sympatric, 49 snails) or

SmVEN (allopatric, 14 snails) strains. Then, 25 days after the primo-infection, we challenged a

subset (28 snails) of SmBRE-primo-infected snails with either SmBRE (14 snails) (homologous

challenge) or SmVEN (14 snails) (heterologous challenge).

To examine the effect of primo-infection on the snail microbiota, we sampled 7 whole snails for

each infection combination on day one (named BB1 and BV1; the first letter refers to the

Biomphalaria strain BgBRE and the second letter refers to the origin of *S. mansoni* strains used

for primo-infection, SmBRE or SmVEN), on day four (BB4 and BV4), and on day twenty-five

(BB25, no SmVEN-infected snails were sampled on day 25) after the primo-infection. This BB25

sample is used as control for microbiota changes observed following challenge infections. Thus

then, we sampled 7 whole-snails on day one after the challenge (BBB1 and BBV1; the third

letter refers to the origin of the *S. mansoni* strain used for the challenge, SmBRE or SmVEN) and

on day four (BBB4 and BBV4) after the challenge. In addition, we used 6 naive snails collected at

the beginning of the experiment (B0) and 6 naive snails collected at the time of challenge (i.e.

day 25, named B25) as controls to assess the variability and stability of the BgBRE snail's

bacterial communities in our breeding and rearing laboratory conditions. Snails were not fed 24

hours before the DNA extraction for metabarcoding analysis.

Infection and sampling for host antimicrobial immune response

We performed a primo-infection of 180 BgBRE snails with either SmBRE (sympatric, 140 snails)

or SmVEN (allopatric, 40 snails). Then, 25 days after infection, we challenged a subset (80

snails) of SmBRE-infected snails with either SmBRE (40 snails) (homologous challenge) or

SmVEN (40 snails) (heterologous challenge) (Portet et al., 2019).

To study the primo-infection transcriptomic immune response of the snail, we took 20 snails

pooled for each infection type on day one (BB1 and BV1), four (BB4 and BV4), and twenty-five

(BB25). After the challenge with SmBRE or SmVEN, we sampled 20 snails in each condition at

days 1 and 4. Then samples of day 1 and day 4 were mixed together into a single sample

referring to BBB or BBV. In addition, we used 2 pools of 30 naive snails (B0.1 and B0.2) to

establish the basal snail transcriptomic activity (Portet et al., 2019).

Extraction and sequencing

DNA extraction and 16S rDNA sequencing

Immediately after sampling, snail shells were cleaned with alcohol and removed, whole snails

were then frozen in liquid nitrogen and grounded. The total DNA was extracted with DNeasy

Blood and Tissue Kit (Qiagen) according to the manufacturer's protocol. The DNA quantification

was performed by Qubit 2.0 Fluorometer, using dsRNA BR Assay kit. Individual 16S rDNA

amplicon libraries were generated using the 341F (CCTACGGGNGGCWGCAG) and 805R

(GACTACHVGGGTATCTAATCC) primers targeting the variable V3-V4 loops (Klindworth et al.,

2013). Paired-end sequencing with 250 bp read length was performed on the Illumina MiSeq

sequencing system (Genome Québec, Montréal, Québec, Canada) using the v2 chemistry

according to the manufacturer's protocol.

RNA extraction and transcriptomic sequencing

Immediately after sampling, snail shells were cleaned with alcohol and removed, then snails

were pooled according to infection type. Total RNA was extracted using TRIZOL (Sigma Life

Science, USA) according to the manufacturer's instructions. For BBB and BBV equimolar

amounts of RNA extracted from molluscs challenged at both 1 and 4 days were mixed together

into a single sample. mRNAs were sequenced in paired-end 72-bp read lengths, using the

mRNA-Seq kit for transcriptome sequencing on Illumina Genome Analyzer II (MGX-Montpellier

GenomiX, Montpellier, France).

Microbiota analysis

Data analysis of 16S sequences

The FROGS pipeline (Find Rapidly OTU with Galaxy Solution) implemented on a galaxy instance

(http://sigenae-workbench.toulouse.inra.fr/galaxy/) was used for data processing (Escudié et

al., 2017). Briefly, paired reads were merged using FLASH (Magoc and Salzberg, 2011). After

denoising and primer/adapters removal (Didion et al., 2017), de novo clustering was done using

SWARM, which uses a local clustering threshold, with aggregation distance d=3 after denoising.

Chimeras were removed using VSEARCH (Rognes et al., 2016). We filtered out the singletons

and performed taxonomic assignment using Blast+ against the Silva database (release 128).

All statistical analyses were done using R v3.3.1 (R: a language and environment for

statistical computing, 2008; R Development Core Team, R Foundation for Statistical Computing,

Vienna, Austria [http://www.R-project.org]). We used the phyloseq R package for community

composition analysis (McMurdie and Holmes, 2013) to infer alpha diversity metrics as well as

beta diversity (between sample distance). Beta diversity was examined by Principal Coordinate

Analysis (PCoA) using the Bray-Curtis distance matrices. We performed a Mann Whitney U test

(p<0.05) to compare alpha diversity and one-way PERMANOVA with a Benjamini & Hochberg

post-hoc to analyze beta diversity between the experimental groups. For all analyses, the

threshold significance level was set at 0.05.

<u>Analysis of core-microbiota</u>

We defined the core-microbiota as the set of bacterial families that were present in 100% of

the naive individuals excluding unknown or multi-affiliations at lower taxonomic ranks (Table

S1). We then used the abundances of OTUs belonging to these families to examine the

composition of the core-microbiota.

To check if the core-microbiota is affected by infection, we compared the abundance of core

families between infected snails and naive conditions with a one-way PERMANOVA with a

Benjamini & Hochberg post-hoc correction. Moreover, a frequency test was performed to

determine which specific families were affected during infection. The number of significantly

differentially represented families at each sampling day (1, 4 and 25 days after primo infections

and 1, 4 days after challenges) was calculated to assess the temporal variability during the

course of infection.

Transcriptome analysis of antimicrobial immune response

<u>Antimicrobial response</u>

An antimicrobial transcriptome was built from transcripts known to be involved in *Biomphalaria*

immune response against bacteria (antimicrobial peptides: biomphamacin or antimicrobial

proteins: LBP/BPI and achacin; see Table S2 for details). The full-length sequences of these

transcripts were recovered from GenBank and the *Biomphalaria* genome (Adema et al., 2017)

and joined in a subset that represents the antimicrobial transcriptome of *B. glabrata*. This

antimicrobial transcriptome was then concatenated with a de-novo assembled transcriptome

of *Biomphalaria* available in our laboratory (see (Dheilly et al., 2015; Pinaud et al., 2016;

Galinier et al., 2017) for details) and uploaded on the Galaxy server. Before concatenation, a

blastn (70 % identity and 90% coverage) was conducted to identify redundant transcripts across

the transcriptomes. Redundant transcripts were then discarded using CDhitEst to avoid

mapping errors and bias in read counts when using Bowtie2 package.

<u>Differential expression analysis</u>

High-quality reads (Phred score >29) were aligned to the concatenated transcriptome using

Bowtie2 (v.2.0.2), which was run locally on a Galaxy server. The DESeq2 (v2.12) R package was

used to identify differential expression levels between uninfected (B0.1 and B0.2) and infected

conditions (p value < 0.05).

Acknowledgments

BG was supported by ANR JCJC INVIMORY (number ANR 13-JSV7-0009) from the French

National Research Agency (ANR). The authors want to thanks the DHOF program of the

UMR5244/IHPE (http://ihpe.univ-perp.fr/en/ihpe-transversal-holobiont/) for partial financial

support. This study is set within the framework of the "Laboratoires d'Excellences (LABEX)"

TULIP (ANR-10-LABX-41).

Reference

Adema, C.M., Hillier, L.W., Jones, C.S., Loker, E.S., Knight, M., Minx, P. et al. (2017) Whole

genome analysis of a schistosomiasis-transmitting freshwater snail. *Nature communications* **8**:

15451.

Aksoy, S., Weiss, B., and Attardo, G. (2008) Paratransgenesis applied for control of tsetse

transmitted sleeping sickness. *Adv Exp Med Biol* **627**: 35-48.

Allan, E.R.O., Tennessen, J.A., Sharpton, T.J., and Blouin, M.S. (2018) Allelic Variation in a Single

Genomic Region Alters the Microbiome of the Snail Biomphalaria glabrata. *J Hered* **109**: 604-

609.

Astudillo-García, C., Bell James, J., Webster Nicole, S., Glasl, B., Jompa, J., Montoya Jose, M.,

and Taylor Michael, W. (2017) Evaluating the core microbiota in complex communities: A

systematic investigation. *Environmental Microbiology* **19**: 1450-1462.

Augustin, R., Fraune, S., and Bosch, T.C.G. (2010) How Hydra senses and destroys microbes.

Seminars in Immunology **22**: 54-58.

Augustin, R., Schröder, K., Murillo Rincón, A.P., Fraune, S., Anton-Erxleben, F., Herbst, E.-M. et

al. (2017) A secreted antibacterial neuropeptide shapes the microbiome of Hydra. *Nature*

communications **8**: 698.

Bosch, T.C. (2013) Cnidarian-microbe interactions and the origin of innate immunity in

metazoans. *Annu Rev Microbiol* **67**: 499-518.

Bouchut, A., Roger, E., Gourbal, B., Grunau, C., Coustau, C., and Mitta, G. (2008) The

compatibility polymorphism in invertebrate host/trematode interactions: research of molecular

determinants. *Parasite* **15**: 304-309.

Budachetri, K., and Karim, S. (2015) An insight into the functional role of thioredoxin reductase,

a selenoprotein, in maintaining normal native microbiota in the Gulf Coast tick (Amblyomma

maculatum). *Insect Mol Biol* **24**: 570-581.

Chen, F., Krasity, B.C., Peyer, S.M., Koehler, S., Ruby, E.G., Zhang, X., and McFall-Ngai, M.J.

(2017) Bactericidal Permeability-Increasing Proteins Shape Host-Microbe Interactions. *mBio* **8**:

e00040-00017.

Cirimotich, C.M., Dong, Y., Clayton, A.M., Sandiford, S.L., Souza-Neto, J.A., Mulenga, M., and

Dimopoulos, G. (2011) Natural microbe - mediated refractoriness to Plasmodium infection in

Anopheles gambiae. *Science* **332**: 855-858.

Contijoch, E.J., Britton, G.J., Yang, C., Mogno, I., Li, Z., Ng, R. et al. (2019) Gut microbiota density

influences host physiology and is shaped by host and microbial factors. *Elife* **8**: pii: e40553.

Coustau, C., Gourbal, B., Duval, D., Yoshino, T.P., Adema, C.M., and Mitta, G. (2015) Advances in

gastropod immunity from the study of the interaction between the snail Biomphalaria glabrata

and its parasites: A review of research progress over the last decade. *Fish & Shellfish*

Immunology **46**: 5-16.

Coutinho-Abreu, I.V., Zhu, K.Y., and Ramalho-Ortigao, M. (2010) Transgenesis and

paratransgenesis to control insect-borne diseases: current status and future challenges.

Parasitol Int **59**: 1-8.

Dheilly, N.M., Duval, D., Mouahid, G., Emans, R., Allienne, J.-F., Galinier, R. et al. (2015) A family

of variable immunoglobulin and lectin domain containing molecules in the snail Biomphalaria

glabrata. *Developmental & Comparative Immunology* **48**: 234-243.

Didion, J.P., Martin, M., and Collins, F.S. (2017) Atropos: specific, sensitive, and speedy

trimming of sequencing reads. *PeerJ* **5**: e3720.

Dinh, C., Farinholt, T., Hirose, S., Zhuchenko, O., and Kuspa, A. (2018) Lectins modulate the

microbiota of social amoebae. *Science* **361**: 402-406.

Doenhoff, M.J., Hagan, P., Cioli, D., Southgate, V., Pica-Mattoccia, L., Botros, S. et al. (2009)

Praziquantel: its use in control of schistosomiasis in sub-Saharan Africa and current research

needs. *Parasitology* **136**: 1825-1835.

Ducklow, H.W., Boyle, P.J., Maugel, P.W., Strong, C., and Mitchell, R. (1979) Bacterial flora of

the schistosome vector snail Biomphalaria glabrata. *Applied and Environmental Microbiology*

38: 667-672.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K. et al. (2017) FROGS:

Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics*: btx791.

Fallon, P.G., and Doenhoff, M.J. (1994) Drug-Resistant Schistosomiasis: Resistance to

Praziquantel and Oxamniquine Induced in Schistosoma Mansoni in Mice is Drug Specific. *The*

American Journal of Tropical Medicine and Hygiene **51**: 83-88.

Franzenburg, S., Fraune, S., Künzel, S., Baines, J.F., Domazet-Lošo, T., and Bosch, T.C.G. (2012)

MyD88-deficient Hydra reveal an ancient function of TLR signaling in sensing bacterial

colonizers. *Proceedings of the National Academy of Sciences of the United States of America*

109: 19374-19379.

Franzenburg, S., Walter, J., Künzel, S., Wang, J., Baines, J.F., Bosch, T.C.G., and Fraune, S. (2013)

Distinct antimicrobial peptide expression determines host species-specific bacterial

associations. *Proceedings of the National Academy of Sciences of the United States of America*

110: E3730-E3738.

Fraune, S., Augustin, R., Anton-Erxleben, F., Wittlieb, J., Gelhaus, C., Klimovich, V.B. et al. (2010)

In an early branching metazoan, bacterial colonization of the embryo is controlled by maternal

antimicrobial peptides. *Proceedings of the National Academy of Sciences of the United States of*

America **107**: 18067-18072.

Fujio-Vejar, S., Vasquez, Y., Morales, P., Magne, F., Vera-Wolf, P., Ugalde, J.A. et al. (2017) The

Gut Microbiota of Healthy Chilean Subjects Reveals a High Abundance of the Phylum

Verrucomicrobia. *Front Microbiol* **8**: 1221.

Futo, M., Armitage, S.A., and Kurtz, J. (2015) Microbiota Plays a Role in Oral Immune Priming in

Tribolium castaneum. *Front Microbiol* **6**: 1383.

Galinier, R., Roger, E., Moné, Y., Duval, D., Portet, A., Pinaud, S. et al. (2017) A multistrain

approach to studying the mechanisms underlying compatibility in the interaction between

Biomphalaria glabrata and Schistosoma mansoni. *PLOS Neglected Tropical Diseases* **11**:

e0005398.

Gendrin, M. (2017) A Swiss Army Knife to Cut Malaria Transmission. *Cell Host & Microbe* **22**:

577-579.

Gilbert, J.A., Medlock, J., Townsend, J.P., Aksoy, S., Ndeffo Mbah, M., and Galvani, A.P. (2016)

Determinants of Human African Trypanosomiasis Elimination via Paratransgenesis. *PLoS Negl*

Trop Dis **10**: e0004465.

Hooper, L.V., and Macpherson, A.J. (2010) Immune adaptations that maintain homeostasis with

the intestinal microbiota. *Nat Rev Immunol* **10**: 159–169.

Hooper, L.V., Littman, D.R., and Macpherson, A.J. (2012) Interactions between the microbiota

and the immune system. *Science (New York, N.Y.)* **336**: 1268-1273.

Jenkins, T.P., Peachey, L.E., Ajami, N.J., MacDonald, A.S., Hsieh, M.H., Brindley, P.J. et al. (2018)

Schistosoma mansoni infection is associated with quantitative and qualitative modifications of

the mammalian intestinal microbiota. *Sci Rep* **8**: 12072.

Joshi, D., Pan, X., McFadden, M.J., Bevins, D., Liang, X., Lu, P. et al. (2017) The Maternally

Inheritable Wolbachia wAlbB Induces Refractoriness to Plasmodium berghei in Anopheles

stephensi. *Frontiers in Microbiology* **8**: 366.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glockner, F.O.

(2013) Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. *Nucleic Acids Res* **41**: e1.

Lozupone, C.A., Stombaugh, J.I., Gordon, J.I., Jansson, J.K., and Knight, R. (2012) Diversity,

stability and resilience of the human gut microbiota. *Nature* **489**: 220-230.

Magoc, T., and Salzberg, S.L. (2011) FLASH: fast length adjustment of short reads to improve

genome assemblies. *Bioinformatics* **27**: 2957-2963.

McFall-Ngai, M., Hadfield, M.G., Bosch, T.C., Carey, H.V., Domazet-Loso, T., Douglas, A.E. et al.

(2013) Animals in a bacterial world, a new imperative for the life sciences. *Proc Natl Acad Sci U*

S A **110**: 3229-3236.

McMurdie, P.J., and Holmes, S. (2013) phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. *PLoS One* **8**: e61217.

Mitta, G., Adema, C.M., Gourbal, B., Loker, E.S., and Theron, A. (2012) Compatibility

polymorphism in snail/schistosome interactions: From field to theory to molecular

mechanisms. *Dev Comp Immunol* **37**: 1-8.

Mitta, G., Gourbal, B., Grunau, C., Knight, M., Bridger, J.M., and Théron, A. (2017) Chapter

Three – The Compatibility Between Biomphalaria glabrata Snails and Schistosoma mansoni: An

Increasingly Complex Puzzle. *Advances in Parasitology* **97**: 111-145.

Mitta, G., Galinier, R., Tisseyre, P., Allienne, J.F., Girerd-Chambaz, Y., Guillou, F. et al. (2005)

Gene discovery and expression analysis of immune-relevant genes from Biomphalaria glabrata

hemocytes. *Developmental and comparative immunology* **29**: 393-407.

Mone, Y., Gourbal, B., Duval, D., Du Pasquier, L., Kieffer-Jaquinod, S., and Mitta, G. (2010) A

large repertoire of parasite epitopes matched by a large repertoire of host immune receptors in

an invertebrate host/parasite model. *PLoS Negl Trop Dis* **4**: pii: e813.

Moné, Y., Ribou, A.-C., Cosseau, C., Duval, D., Théron, A., Mitta, G., and Gourbal, B. (2011) An

example of molecular co-evolution: Reactive oxygen species (ROS) and ROS scavenger levels in

Schistosoma mansoni/Biomphalaria glabrata interactions. *International Journal for Parasitology*

41: 721-730.

Narasimhan, S., Schuijt, T.J., Abraham, N.M., Rajeevan, N., Coumou, J., Graham, M. et al. (2017)

Modulation of the tick gut milieu by a secreted tick protein favors Borrelia burgdorferi

colonization. *Nature communications* **8**: 184.

Nyholm, S.V., and McFall-Ngai, M.J. (2004) The winnowing: establishing the squid-vibrio

symbiosis. *Nat Rev Microbiol* **2**: 632-642.

Onchuru, T.O., and Kaltenpoth, M. (2019) Established cotton stainer gut bacterial mutualists

evade regulation by host antimicrobial peptides. *Appl Environ Microbiol* **pii: AEM.00738-19.** .

Pinaud, S., Portela, J., Duval, D., Nowacki, F.C., Olive, M.-A., Allienne, J.-F. et al. (2016) A Shift

from Cellular to Humoral Responses Contributes to Innate Immune Memory in the Vector Snail

Biomphalaria glabrata. *PLOS Pathogens* **12**: e1005361.

Portet, A., Pinaud, S., Chaparro, C., Galinier, R., Dheilly, N.M., Portela, J. et al. (2019) Sympatric

versus allopatric evolutionary contexts shape differential immune response in Biomphalaria /

Schistosoma interaction. *PLoS Pathog* **15**: e1007647.

Pradeu, T., and Eric, V. (2014) The discontinuity theory of immunity. *Sci Immunol* **1**: 1-9.

Ramirez, J.L., Souza-Neto, J., Cosme, R.T., Rovira, J., Ortiz, A., Pascale, J.M., and Dimopoulos, G.

(2012) Reciprocal tripartite interactions between the Aedes aegypti midgut microbiota, innate

immune system and dengue virus influences vector competence. *PLOS Neglected Tropical*

Diseases **6**: 1-11.

Rodrigues, J., Brayner, F.A., Alves, L.C., Dixit, R., and Barillas-mury, C. (2010) Hemocyte

Differentiation Mediates Innate Immune Memory in Anopheles gambiae Mosquitoes. *Science*

329: 1353-1355.

Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahe, F. (2016) VSEARCH: a versatile open

source tool for metagenomics. *PeerJ* **4**: e2584.

Ryu, J.-H., Kim, S.-H., Lee, H.-Y., Bai, J.Y., Nam, Y.-D., Bae, J.-W. et al. (2008) Innate Immune

Homeostasis by the Homeobox Gene Caudal and Commensal-Gut

Mutualism in Drosophila. *Science* **319**: 777.

Sansone, C.L., Cohen, J., Yasunaga, A., Xu, J., Osborn, G., Subramanian, H. et al. (2015)

Microbiota-dependent priming of antiviral intestinal immunity in Drosophila. *Cell Host and*

Microbe **18**: 571-581.

Silva, T.M., Melo, E.S., Lopes, A.C.S., Veras, D.L., Duarte, C.R., Alves, L.C., and Brayner, F.A.

(2013) Characterization of the bacterial microbiota of Biomphalaria glabrata (Say, 1818)

(Mollusca: Gastropoda) from Brazil. *Letters in Applied Microbiology* **57**: 19-25.

Tennessen, J.A., Théron, A., Marine, M., Yeh, J.-Y., Rognon, A., and Blouin, M.S. (2015)

Hyperdiverse Gene Cluster in Snail Host Conveys Resistance to Human Schistosome Parasites.

PLoS Genetics **11**: e1005067.

Wang, X.W., Xu, J.D., Zhao, X.F., Vasta, G.R., and Wang, J.X. (2014) A shrimp C-type lectin

inhibits proliferation of the hemolymph microbiota by maintaining the expression of

antimicrobial peptides. *J Biol Chem* **289**: 11779-11790.

Warne, R.W., Kirschman, L., and Zeglin, L. (2019) Manipulation of gut microbiota during critical

developmental windows affects host physiological performance and disease susceptibility

across ontogeny. *J Anim Ecol* **Epub ahead of print**.

WHO (2002) TDR Strategic Direction for Research: Schistosomiasis. *World Health Organization*

Information.

Xi, Z., Ramirez, J.L., and Dimopoulos, G. (2008) The Aedes aegypti toll pathway controls dengue

virus infection. *PLOS Pathogens* **4(7):e1000098**.

Yang, H.T., Yang, M.C., Sun, J.J., Shi, X.Z., Zhao, X.F., and Wang, J.X. (2016) Dual oxidases

participate in the regulation of intestinal microbiotic homeostasis in the kuruma shrimp

Marsupenaeus japonicus. *Dev Comp Immunol* **59**: 153-163.

Yang, H.T., Yang, M.C., Sun, J.J., Guo, F., Lan, J.F., Wang, X.W. et al. (2015) Catalase eliminates

reactive oxygen species and influences the intestinal microbiota of shrimp. *Fish Shellfish*

Immunol **47**: 63-73.

Zhang, H., Sparks, J.B., Karyala, S.V., Settlage, R., and Luo, X.M. (2015) Host adaptive immunity

alters gut microbiota. *ISME J* **9**: 770-781.

Legends to Figures

Fig. 1: Experimental protocol

Fig. 2: *Biomphalaria glabrata* microbiota characterization

Characterisation of the *Biomphalaria* bacterial microbiota of six naive snails recovered at the

start of experimentation (B0.1; B0.2; B0.3; B0.4; B0.5 and B0.6) and 6 naive snails recovered 25

days after the start of experimentation (B25.1; B25.2; B25.3; B25.4; B25.5 and B25.6). A.

Phylum level composition of the 20 most abundant OTUs phyla among the 12 naive snails. B.

The Venn diagram represents the number of the 97 OTUs families which shared between the 6

naive snails at B0 (left Venn diagram), and between the 6 naive snails at B25 (right Venn

diagram).

Fig. 3: Microbiota alpha Diversity

Boxplots of Chao1 and Shannon indices for all samples. For the Naive condition, B0 and B25

snails were pooled; BB: primo infection of BgBRE by SmBRE; BV: primo infection of BgBRE by

SmVEN; BBB: primo infection of BgBRE by SmBRE and challenge by SmBRE; BBV: primo

infection of BgBRE by SmBRE and challenge by SmVEN. The time point is mentioned with 1, 4 or

25 corresponding to the day after primo-infection or challenge. The differences between naive

and infected conditions were tested with a Mann-Whitney U test and significant differences

mentioned with "*".

Fig. 4: Beta diversity and bacterial communities following *Biomphalaria* infection

Dynamics of the bacterial microbiota of *Biomphalaria glabrata* following *Schistosoma* primo-

infection and challenge. A. Functional diversity comparisons of *Biomphalaria* microbiota along

infection. Principal coordinate analysis of pairwise Bray-Curtis distances between all infection

type and time samples. Axes represent the two synthetic variables explaining the greatest

proportion of variation in the data set. The sample name indicated in the figure corresponds to

the centroid of all the biological replicates points of the respective experimental sample. B.

Phylum level composition of the 20 most abundant OTUs among all points of the kinetic. In this

representation, the replicate naive snails were pooled for more readability.

Fig. 5: Differential gene expression of Biomphamacin antimicrobial peptides

Log2FC (fold change) of antimicrobial immune transcripts between naive and infected snails

inferred from previous RNAseq analysis on the same experiment. A positive Log2 fold-change

indicates over-expression in infected snails compared to the naive snails. Antimicrobial peptide

families included 6 Biomphamacins (macin-like AMPs) consisting of 6 genes (shade of green).

Fig. S1: Core Microbiota analysis

Modifications of the bacterial core microbiota of *Biomphalaria glabrata* following *Schistosoma*

primo-infection and challenge. After a one-way Permanova with a Benjamini & Hochberg post-

hoc, a frequency test was realised to determine which specific families were affected during

infection. A. Proportion of OTU Families affected by infection are represented by colour circles.

The Venn diagram shows more precisely the specific impact of infections on the OTU families

partially influenced. B. Heatmap corresponding to all core microbiota with clustering on Familiy

modification patterns between conditions. The scale of red correspond to ratio between naive

condition and each point of infection.

Fig. S2: Differential gene expression of LBP/BPI and Achacin antimicrobial protein families

Log2FC (fold change) of antimicrobial immune transcripts between naive and infected snails

inferred from previous RNAseq analysis on the same experiment. A positive Log2 fold-change

indicates over-expression in infected snails compared to the naive snails. Antimicrobial protein

families: LBP/BPIs (*Lipopolysaccharide-binding protein / Bactericidal Permeability-Increasing*

protein) consisting of 5 genes (in blue) and achacins consisting of 2 genes (in orange).

Legend (right of panel A):
- Acidobacteria
- Actinobacteria
- Armatimonadetes
- Bacteroidetes
- Chlamydiae
- Chlorobi
- Chloroflexi
- Cyanobacteria
- Deinococcus-Thermus
- Firmicutes
- Gemmatimonadetes
- Nitrospirae
- Parcubacteria
- Planctomycetes
- Proteobacteria
- Saccharibacteria
- Spirochaetae
- Tenericutes
- TM6 (Dependentiae)
- Verrucomicrobia

A

B

Naive snails	Primo-Infection			Challenge	
	Sympatry	*Allopatry*	*Late Sympatry*	*Homologous*	*Heterologous*

Acidobacteria
Actinobacteria
Armatimonadetes
Bacteroidetes
Chlamydiae
Chlorobi
Chloroflexi
Cyanobacteria
Deinococcus-Thermus
Firmicutes
Fusobacteria
Gemmatimonadetes
Nitrospirae
Parcubacteria
Planctomycetes
Proteobacteria
Saccharibacteria
Spirochaetae
Tenericutes
TM6 (Dependentiae)
Verrucomicrobia