Geographically Opposing Responses of Sea Urchin Recruitment to Changes in Ocean Climate

Daniel K. Okamoto[*1,2], Daniel C. Reed[†3], and Stephen Schroeter[‡3]

[1]Department of Biological Science, Florida State University, Tallahassee, Florida 32303
[3]Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106 USA
[3]Marine Science Institute, University of California Santa Barbara, Santa Barbara, CA 93106 USA

Corresponding Author
Daniel K Okamoto
Department of Biological Science
The Florida State University
Tallahassee, Florida 32303
Phone: (850) 645-9532
emaildokamoto@bio.fsu.edu

Author contributions
DKO, SS, and DR designed research. DKO designed, built and conducted analyses and wrote the initial manuscript. DKO, DCR and SS managed data. SS initiated and oversaw data collection and collaborated on all analyses. All authors contributed to revisions.

---
[*]corresponding author: dokamoto@bio.fsu.edu
[†]email: dan.reed@lifesci.ucsb.edu
[‡]email: schroete@lifesci.ucsb.edu

Abstract

Sea urchins are voracious herbivores that influence the ecological structure and function of
nearshore ecosystems throughout the world. Urchin population growth rates may be particularly
sensitive to climate change because adult reproduction and larval development can vary greatly
with food availability and temperature, and the transport of their larvae, which spend months
feeding in the plankton, are affected by changes in ocean currents. Yet how climate alters sea
urchin populations in space and time by modifying larval recruitment and year-class strength
remains untested. Using an unprecedented spatially replicated 27-year dataset we illustrate how
ocean temperature and climate oscillations differentially affect larval recruitment of the purple
sea urchin (*Strongylocentrotus purpuratus*) and give rise to geographic asynchrony between
northern California (positive) vs southern California (negative). Importantly, we found a strong
correlation between larval recruitment and regional year class strength suggesting that
recruitment variation plays an important role in controlling population dynamics. These results
are the first to show that climatic fluctuations shape broad-scale patterns of sea urchin larval
recruitment and are likely to control dynamics of both populations and marine ecosystems that
vary over the geographical range of their distribution.

Introduction

Climate forecasts project large changes in ocean temperature, biogeochemistry and the nature
and frequency of disruptive events that affect ocean circulation and primary productivity (e.g.
Cai *et al.* 2014; Yeh *et al.* 2009). Such shifts are expected to impose wide-reaching ecological
impacts, in part by altering animal recruitment in space and time (Sydeman *et al.* 2015). As a
result, characterizing how climate shapes variability in recruitment can provide substantial
benefits to the conservation and management of marine resources. For species with planktonic
larvae, significant challenges remain in achieving such objectives due to the substantial effort
needed to characterize spatial and temporal variation in larval abundance and the numerous
sensitive vital rates that contribute to it.

For species like sea urchins, understanding causes and consequences of recruitment
variability has both ecological and economic implications. Sea urchin grazing can alter the
structure of some of the world's most diverse and productive marine ecosystems, including coral
reefs (e.g. Edmunds & Carpenter 2001; Mumby *et al.* 2007), seagrass meadows (reviewed by
Valentine & Heck Jr 1999) and kelp forests (reviewed by Filbee-Dexter & Scheibling 2014). In
addition, sea urchins form the basis of important nearshore fisheries in many regions of the world
(e.g. Andrew *et al.* 2003; Kato & Schroeter 1985). As a result, climate-driven changes in sea
urchin populations have the potential to profoundly affect both marine ecosystems and associated
fisheries. Much of the ecological research on sea urchins has focused on the roles of predation

and disease in controlling adult abundance and their cascading influence on community structure
(e.g. Burt *et al.* 2018; Estes & Duggins 1995; Filbee-Dexter & Scheibling 2014; Lafferty 2004;
Mann & Breen 1972). However, despite the widespread recognition of the importance of
recruitment variation in controlling population fluctuations in many marine species (Shelton &
Mangel 2011), relatively few studies have examined the biotic and abiotic processes controlling
the supply of sea urchin larvae, and the degree to which they affect the abundance and dynamics
of older life stages.

Changes in climate can affect spatial and temporal patterns of larval supply by influencing
the production of larvae by benthic adults and the survival of larvae in the plankton. Increases in
ocean temperature and storm-related disturbances can impact larval production by (1) increasing
adult mortality via physical disturbance (Ebeling *et al.* 1985) and the spread of water-borne
pathogens (Behrens & Lafferty 2004; Lafferty 2004), and (2) reducing adult fecundity by
altering food availability (Foster *et al.* 2015; Okamoto 2014) and inhibiting gametogenesis
(Basch & Tegner 2007; Cochran & Engelmann 1975). Because sea urchins produce feeding
larvae that spend weeks to months in the plankton, increases in ocean temperature can also affect
larval development, growth and survival, either directly, or indirectly by altering the availability
of their phytoplankton food source (Bertram & Strathmann 1998; Byrne *et al.* 2009; Hoegh-
Guldberg & Pearse 1995; Meyer *et al.* 2007; Strathmann 1987). Finally, climate related changes
in patterns of ocean circulation can affect the transport of larvae from source to destination
(Siegel *et al.* 2008; but see Morgan 2014). Thus, the effects of climatic change on sea urchin
recruitment represent cumulative impacts on adult abundance and reproduction, current patterns
that transport larvae, and larval development and survival. Because patterns of ocean
temperature, circulation and upwelling can vary dramatically in space, the effects of climate
change on sea urchin recruitment potentially vary over large spatial scales. A dearth of long-
term, high frequency, spatially extensive data has prevented characterizing temporal and spatial
variability in larval recruitment in sea urchins, the degree to which it is explained by different
sources of environmental variation, and the relative importance of these drivers in accounting for
fluctuations in population size.

Here we used a unique 27-year weekly to biweekly time series of newly metamorphosed
larvae (which we refer to as "larval recruits"), collected on artificial substrates placed at seven
locations distributed across >1000 km of coast in California to: (1) quantify variation in larval
recruitment across different temporal and spatial scales; (2) evaluate whether larval recruitment
on artificial substrates predicts year-class strength in natural populations; and (3) determine the
relative importance of factors contributing to the observed variation in larval recruitment, namely
those that affect reproductive output (such as benthic adult abundance and per-capita fecundity)

and oceanographic conditions that affect planktonic larval survival and transport. Due to the

correlative nature of our analyses our primary aim was to reveal potential causal relationships

that merit further evaluation as drivers of sea urchin recruitment dynamics at spatial and

temporal scales relevant to natural populations.

Study system

Populations of the purple sea urchin (*Strongylocentrotus purpuratus*) occupy shallow

subtidal and intertidal rocky substrata from at least 27°N on the western coast of the Baja

Peninsula (Olivares-Bañuelos *et al.* 2008) to at least 59°N on the Kenai Peninsula in Alaska

(Field & Walker 2003). Purple urchins are broadcast spawners and the seasonality of their

spawning in California is generally thought to be driven by photoperiod and temperature

(Cochran & Engelmann 1975; Gonor 1973; Pearse *et al.* 1986). Fertilized zygotes develop into

planktonic echinoplutei and are obligate planktivores that consume phytoplankton (Strathmann

1987). After spending several weeks to months in the plankton individuals begin final

metamorphosis and settle to the benthos (Strathmann 1978) at a size of ~500 μm in diameter

(Okamoto unpublished data). Larval recruitment varies dramatically among locations at both

small and large spatial scales (Ebert 2010).

Collection of newly settled urchins along the coast of California, USA (1990-2016)

Recruitment of newly metamorphosed purple sea urchins was sampled in three major regions

along the California coast from 1990 through 2016, with a total of 54,588 observations.

Sampling regions (from south to north) included two sites in San Diego (Scripps Pier and Ocean

Beach Pier), four sites in the Santa Barbara Channel (Anacapa Island, Stearns Wharf, Ellwood

Pier and Gaviota Pier) and one site at Fort Bragg (Figure 1). San Diego and the Santa Barbara

Channel lie within the Southern California Bight and Fort Bragg is in northern California. At

each site, urchins were collected using nylon-bristled scrub brushes (2.5 cm long bristles and a 6

x 9 cm wooden base) suspended 1 to 2 m from the benthos (Ebert *et al.* 1994). The majority of

deployments included 4-8 replicate brushes collected weekly at each site from 1990 to 2003, and

biweekly thereafter through 2016. Upon collection brushes were transported to the laboratory in

plastic bags and rinsed through a 350 μm mesh sieve. Purple urchins were then identified to

species, sorted from other organisms, counted and preserved see (Ebert *et al.* 1994) for further

details.

We conducted three primary analyses using these data (see Methods for full details): (1) we

examined biweekly, seasonal and interannual trends and spatial correlations in *S. purpuratus*

larval recruitment using a Bayesian multivariate time series model; (2) we tested whether trends

in larval recruitment corresponded to year-class strength using a generalized linear mixed effects
model; and (3) we nested a linear regression into the Bayesian time series model from (1) to
evaluate the degree to which various biotic and abiotic factors explained patterns of sea urchin
larval recruitment in northern and southern California. In this latter analysis we linked larval
recruitment data to long-term data on sea surface temperature, chlorophyll a (larval food), kelp
coverage (adult food), regional upwelling indices, regional adult abundance indices and global
climate indices.

Results
(1) Estimated Patio-temporal Trends in Purple Urchin Larval Recruitment

Substantial interannual variability in larval recruitment of *S. purpuratus* was observed with
years of poor recruitment (e.g. 1995, 1998, 2005, 2016) that were, on average 1-2 orders of
magnitude lower than neighboring years (Figure 2, Figure S1). Larval recruitment was highly
synchronous among sites within each of the two regions in the Southern California Bight
indicated by mean interannual correlations of r = 0.73 and 0.85 for sites within the Santa Barbara
Channel and San Diego, respectively (Figure 2b, c). Within the Santa Barbara Channel, pairwise
correlations in interannual trends involving Gaviota, Ellwood and Stearns Wharf were higher (r =
0.86 to 0.90) those involving Anacapa (r = 0.46 to 0.71). This resulted in part because different
rates of interannual declines from high larval recruitment in 2011 to a common low value in
2016. Anacapa began to decline in 2012 while the other sites, while the declines at the other
sites did not begin until 2014 (Figure 2b, Supplementary Figure 1 b-e). While Santa Barbara
Channel sites all exhibited a period of strong increases following the low in 2005, San Diego
sites remained below average for the remainder of the time series (Figure 2 c, Supplementary
Figure 1 f-g). In contrast, interannual trends in recruitment at Fort Bragg were asynchronous
with sites in the Santa Barbara Channel (Figure 2a vs. 2b, mean r = -0.23) and uncorrelated to
sites in San Diego (Figure 2a vs. 2c; mean r = 0.12).

Larval recruitment in southern California was highly seasonal, with similar patterns among
sites (Figure 3, Figure S2). On average 90% of recruitment occurred from March to July with a
single peak in late April/early May (Figure 3a). By contrast recruitment at Fort Bragg in
northern California extended over a longer period of time (90% occurred, between January and
September) and often included two peaks per year (a large consistent peak around July and a
smaller peak in February and March of some years; Figure 3a, S3). The seasonal peaks in
recruitment in southern California coincided with the peaks in sea surface chlorophyll (Figure 3a
vs. 3b) and troughs in sea surface temperature (Figure 3a vs. 3c). In northern California the
primary peak occurred slightly after the peak in chlorophyll a.

(2) Relationship Between larval recruitment and subsequent juvenile year-class-strength.

Recruitment of juvenile purple urchins at shallow subtidal reefs in the Santa Barbara Channel
exhibited a significant, positive correlation with larval recruitment to brushes two years prior
(Figure 4, $\chi^2_{df=1}$ = 9.48, p = 0.002). Years with high larval recruitment corresponded with a
nearly three-fold increase in the mean density of juvenile urchins two summers later.

(3) Relationships with Environmental Conditions

Larval recruitment of purple urchins showed strong correlations with local sea surface
temperature (SST, Figure 5 a-g, Figure 6 a), and climate indices, specifically ENSO - measured
as the multivariate El Niño Southern Oscillation Index (MEI - Figure 5 h-n, Figure 6 f) and the
North Pacific Gyre Oscillation (NPGO –Figure 5 o-u, Figure 6 g). The sign of the correlation
was opposite between Northern California (Fort Bragg) and Southern California (Santa Barbara
Channel and San Diego). The Pacific Decadal Oscillation showed a correlation with larval
recruitment only at Fort Bragg (Figure 6 h), but the estimate was highly collinear with that of the
MEI leading to inflated variance of both coefficients (Supplemental Figure 3). No correlation
was observed between larval recruitment and adult urchin density or canopy kelp (data that were
only available for southern California sites; Figure 6 b, e), or the Bakun Upwelling Index (Figure
6c). Sea surface chlorophyll showed no correlation with larval recruitment at Fort Bragg and in
the Santa Barbara Channel and a negative correlation in San Diego – the opposite of the
hypothesized relationship that more food in the plankton would lead to higher abundances of
larvae and higher numbers of larval recruits (Figure 6 d). The network of correlations among
these variables are depicted in (Supplemental Figure 4).

In southern California recruitment was orders of magnitude lower during warm, El Niño
conditions and during the negative phase of the NPGO with the more southern sites in San Diego
responding more strongly to temperature than those further north in the Santa Barbara Channel
(Figure 6 a). In contrast, Fort Bragg, the northern California site, responded positively to
warmer El Niño conditions and negatively to the NPGO. Importantly, sites in southern California
showed opposite relationships from those at Fort Bragg in northern California (Figure 6 f, g).
[Note that we fit models with local variables (SST, adults, kelp coverage, chlorophyll, and Bakun
upwelling index) and ocean-scale indices (ENSO/NPGO/PDO) separately because of strong
functional dependence of the latter with SST]. The correlations between recruitment and both
SST and ENSO occurred on roughly 3-5 year cycles, while the correlation with NGPO occurred
on decadal scales.

Discussion

Climate change poses significant threats to marine ecosystems. Yet determining how many
species and communities will respond to shifting temperature and ocean storm regimes remains a

difficult task. Using a multi-decadal, high frequency, spatially extensive time series we show
how large-scale climatic variation can give rise to geographically different responses in larval
sea urchin recruitment that were synchronous within regions and asynchronous between them.
Larval recruitment of purple sea urchins varied by orders of magnitude among years and
responded oppositely to climate in northern versus southern California. Our finding that larval
recruitment was a significant factor in determining year class strength in natural populations
indicates that large fluctuations in larval recruitment of this prominent herbivore could have far
reaching ecological impacts that resonate throughout the regional marine ecosystems (Pearse
2006).

Climate related changes in the supply of sea urchin larvae are thought to be a primary factor
contributing to increases in sea urchin populations that have caused the dramatic transformation
of benthic communities (Fabricius *et al.* 2010; Hernández *et al.* 2010; Ling *et al.* 2008). Yet
until now, no long-term data have corroborated whether urchin larval supply, regulated by
climate, can alter year class strength. Demonstrating this link has historically proved challenging
because "larval recruitment" can be attenuated by early post-settlement mortality (Connell 1985;
Rowley 1989). Our analysis indicates that the year class strength of purple sea urchins in a
region is significantly influenced by larval supply. This result points to the possibility that future
oceanographic changes may reduce purple urchin recruitment in southern California, but increase
it in northern California with potentially cascading effects on benthic ecosystems. The fact that
our study shows similar correlations between oceanographic factors and recruitment over a 27-
year period, which included multiple ENSO events, suggests that such climatic associations will
continue.

While many recruitment-environment correlations break down over time (Myers 1998), our
study shows a strong association between urchin larval recruitment and climate over a nearly 3
decade time scale. Although sea urchin larval recruitment data prior to 1990 were not available
in the fine temporal scale characterized by our study, there is evidence of ENSO-related
recruitment failures in populations in southern California since 1960s. Between 1969 and the
early 1980's, recruitment of juvenile sea urchins was anomalously low in El Niño years (Ebert
1983; Tegner & Dayton 1991). Between 1969 and 1977, juvenile recruitment was lowest during
the three El Niño years of 1970, 1973, and 1977 at False Point, California near San Diego (Ebert
1983; Tegner & Dayton 1991) and at nearby Point Loma, low recruitment followed the 1982-83
El Niño event (Ebert 1983; Tegner & Dayton 1991). How patterns of larval recruitment may
change in the future depends largely on the process underlying their associations with El Niño.

Although climate related changes to both the benthic and planktonic stages of purple sea
urchins can affect larval recruitment via their effects on the production of larvae, our analysis
shows no significant correlations between larval recruitment and adult abundance, adult food
(kelp) or larval food (phytoplankton) – (Figure 6 and Supplemental Figure 3). This narrows
down the range to two plausible hypotheses to the regionally varying effects of ENSO and
temperature on larval supply and recruitment: (1) temperature-related effects on larval
production and survival, and 2) ENSO-related changes in currents that vary between southern
and northern California, which affect larval delivery. While adult purple urchins can tolerate
temperatures up to 25°C (Farmanfarmaian & Giese 1963), temperatures above 17°C during
ENSO events in southern California may adversely affect larval supply via effects on gamete
production (Basch & Tegner 2007; Cochran & Engelmann 1975), fertilization (Schroeder &
Battaglia 1985) larval development and gene expression (Padilla-Gamiño *et al.* 2013; Runcie *et*
al. 2012; Wong *et al.* 2018), and larval survival (Azad *et al.* 2012). Such responses to high
temperatures may, in part, explain why recruitment at Fort Bragg, where sea surface
temperatures are rarely above 16°C, was not negatively correlated with temperature.

Like variation in temperature, regional circulation is hypothesized to be an important driver
of larval recruitment for many species, including those in the Southern California Bight
(Blanchette *et al.* 2006; Broitman *et al.* 2005; McManus & Woodson 2012; Woodson *et al.*
2012). ENSO can cause major changes in patterns of ocean circulation along the US west coast,
with warm El Niño years often characterized by anomalously strong poleward flow and
decreased local retention (Lynn & Bograd 2002; Mitarai *et al.* 2009). Importantly, these effects
differ between northern and southern California and may partly explain the divergences in
climatic responses in recruitment between these two regions. In northern California, El Niño
events are associated with relaxed upwelling and downwelling Kelvin waves (Chavez *et al.*
2002) which hypothetically improves larval retention, but is thought to have a more nuanced
impact in the Southern California Bight (Lynn & Bograd 2002; Mitarai *et al.* 2009). Although
we cannot definitively partition out the effects of climate associated changes in current patterns
and temperature on larval recruitment, our observations on larval recruitment and subsequent
year-class strength point strongly to broader-scale ocean climate effects.

Our study demonstrates a consistent and geographically variable relationship between
climate and larval recruitment and year-class strength of purple sea urchins over a three-decade
time period. These unique time series provide an unprecedented illustration of climate related
impacts on larval recruitment dynamics that are notoriously difficult to investigate. We show
that fluctuations in larval recruitment can influence year-class strength in an herbivore that is
known to severely alter the composition of an entire benthic ecosystem. Although these results

are based on correlations, they were observed consistently over three decades (with further

anecdotal evidence spanning six decades) that have included multiple ENSO events. Moreover,

our findings provide important insights into the environmental drivers of larval recruitment,

which is essential for understanding and predicting the influence of climate on regional

population and community dynamics of marine species. Oceans are experiencing simultaneous

shifts in temperature, water chemistry, productivity and circulation; thus future investigations

aimed at determining the specific biotic and abiotic processes that regulate larval recruitment

should provide much needed knowledge of how climate change is influencing regional

population dynamics to alter the dynamics and structure marine ecosystems.

Methods

(1) Estimated patio-temporal Trends in Purple Urchin Larval Recruitment

We estimated annual, seasonal, and spatial trends in larval recruitment using an integrated

spatio-temporal model that accounted for both the intercorrelated and heterogeneous nature of

the multivariate time series. Specifically, we used a Bayesian hierarchical model with

spatiotemporal correlations in the underlying annual and seasonal scale trends, and spatial

correlations in the process error terms. We used this model to simultaneously estimate biweekly,

seasonal and interannual trends in recruitment, and spatial synchrony while accounting for

observation noise.

Using this model, we calculated the log-scale biweekly mean trend in larval recruitment as

the sum of the estimated annual trend, seasonal trend and process error in the statistical model.

The annual trend was estimated using a separable spatio-temporal covariance prior where the

spatial covariance is unstructured (i.e. each element estimated) and the temporal covariance

determined by a Matérn 3/2 prior. Functionally, we constructed the covariance matrix for the

annual trend using Kronecker matrix-vector operations with the Cholesky decompositions of the

spatial and temporal correlation functions independently for efficiency (Steeb & Hardy 2011).

The seasonal trend within each site was estimated using a periodic temporal covariance (MacKay

1998). The process error (deviations from the sum of smoothed annual and seasonal trends) was

modeled with an unstructured spatial covariance, assuming temporal independence. We used a

Poisson likelihood to link the expectation (biweekly mean trend) to the observed counts of *S.*

purpuratus larval recruits.

Occasionally we encountered samples with sufficiently large numbers of larval recruits

(hundreds to thousands of urchins per sample - 1,070 out of 53,478 samples or ~2%), which

prevented efficient identification of all urchins to species.   In these cases, *S. purpuratus* were

identified from a homogenized subsample of 30-100 individuals. We accounted for the
uncertainty in *S. purpuratus* abundance introduced by this subsampling routine by using a beta
prior with parameters determined by the subsampled *S. purpuratus* counts and the non-*S.*
purpuratus counts. This prior was derived as the analytical posterior distribution of the
proportion of the true, latent fraction that was *S. purpuratus* given a Jeffrey's prior on a binomial
model of the observed counts.

All equations for the model are given in Supplementary Table 1 and parameters and variables
defined in Supplementary Table 2. We estimated the models using Stan (Carpenter *et al.* 2016;
Hoffman & Gelman 2014; Stan Development Team 2016b, a) via R (R Core Team 2017) with
three 1000 iteration chains after 1000 iteration burn-in.   Stan code for the model is provided in
the appendix.

(2) Relationship Between larval recruitment and subsequent juvenile year-class-strength.
We tested whether the mean of our estimates of larval recruitment in the Santa Barbara
Channel (mean of Anacapa, Stearns Wharf, Ellwood Pier and Gaviota Pier) was predictive of
subsequent juvenile recruitment on natural reefs in the region. We calculated juvenile densities
from the Channel Islands Kelp Forest Monitoring (KFM) Program (Kushner *et al.* 2013),
including only sites whose time series extend from 1990 through 2016. To test for correlations,
we used a Bayesian generalized linear mixed effects model (GLMM) with a negative binomial
likelihood and survey site as a random effect.

The KFM program surveys purple sea urchins at the Santa Barbara Channel Islands by
counting the total number of individuals in defined areas and measuring approximately 100
individuals at each site. Thus, to model density we used the number of juveniles measured
divided by the total number of individuals measured, multiplied by the total number of
individuals counted, divided by the area surveyed.

$$Juvenile\ Density_{i,t}\ =\ \frac{Juveniles\ Measured_{i,t}}{Total\ Meas_{i,t}}\ \frac{Total\ Count_{i,t}}{Total\ Area_{i,t}}$$

We modeled density in the GLM using the number of measured juvenile urchins (2.5 cm in
test diameter- the approximate cutoff size for reproduction; ref Kenner & Lares 1991). Let $\mu_{i,t}$
represent expected juvenile density at site *i* in year *t*. We constructed the regression as:

$$\ln \mu_{i,t} = \alpha_i + \beta \ln Larval\ recruitment_t$$

We allowed the intercept (α_i) to vary by site nested within each island because of overall
differences in mean juvenile density among sites and islands. We used a negative binomial (NB)
with the using direct mean and variance parameterization form:

$Juveniles\ Measured_{i,t} \sim \text{NB} \left(\ln \mu_{i,t} - \ln \left[\frac{1}{Tot.Measured_{i,t}} \frac{Tot.Count_{i,t}}{Tot.Area_{i,t}} \right], \sigma^2 \right)$

We constructed the model in this format (i.e. with the density denominator in the left hand
side of the equation) to maintain the sample size and integer nature of the data while modeling
the mean density. We tested the hypothesis of no correlation using a likelihood ratio test.
Models were estimated using glmmTMB (Magnusson *et al.* 2017) in R.

(3) Relationships with Environmental Conditions

We tested how larval recruitment trends correlated with physical and biological variables
using two core analyses. Covariates included the following:

(i) Oceanographic climate indices (monthly, 1990-2016, all sites): We used three major
global indices of oceanographic climate. The multivariate El Niño Southern Oscillation Index
(MEI) provides a metric of the intensity of El Niño/La Niña fluctuations and is derived from
several metrics of sea surface temperature, surface winds, sea level pressure, and cloudiness of
the sky (Wolter & Timlin 1993; Wolter & Timlin 1998). The Pacific Decadal Oscillation (PDO)
is the 1st empirical orthogonal function analysis mode of sea surface temperature variability and
height in the Northeast Pacific (Mantua & Hare 2002) and the North Pacific Gyre Oscillation
(NPGO) is the 2nd mode (Di Lorenzo *et al.* 2008).

(ii) Coastal upwelling index (monthly, 1997-2016, all sites): The Bakun index (Bakun 1973)
provides an index of coastal upwelling and specifically describes the volume of water that is
transported offshore from Ekman transport (http://www.pfel.noaa.gov/). Negative values indicate
downwelling and surface waters that moved onshore, while positive values indicate upwelling
and surface waters moving offshore. This index has been used in relation to delivery of larvae
onshore and advection from shore in addition to its value as a metric of coastal productivity. For
the Southern California Bight sites, we used data from 33N-119W, while in Northern California
we used data from 39N-125W.

(iii) Sea surface chlorophyll (monthly, 1997-2016, all sites): Satellite imagery of sea surface
chlorophyll provides a spatially and temporally resolved estimate of phytoplankton biomass that
is not available from in situ sampling. Thus, as a metric of planktonic food availability we used
version 3.1 of the OC-CCI merged ocean color time series (Sathyendranath *et al.* 2018) from
SeaWIFs (O'Reilly *et al.* 1998), MERIS (Rast *et al.* 1999), MODIS (Esaias *et al.* 1998) and
VIIRS (Schueler *et al.* 2002) that allows for the temporal and spatial coverage required for this
study. Because sea surface chlorophyll concentration data are spatial grids of time series, we

included averaged cells meeting particular geographic conditions for each site. For mainland
sites, all cells had to lay within 10 km offshore of the mainland coastline that stretched 150 km in
coastline length from the larval recruitment collection site or within 5 km of any island coastline
of any island within a 150 km radius. For the site at Anacapa Island, we included any point
within a 150 km radius and within 10 km of any coastline. We used 150 km because that is the
average Lagrangian estimate for dispersal distances for species with a planktonic larval duration
(PLD) of 30-days (Siegel *et al.* 2003). However, this is potentially an overestimate for a 30-day
PLD as behavior and complex current patterns may reduce this distance substantially (Shanks
2009). Figure 1 depicts the aerial extent of chlorophyll data used in the analysis.

(*iv*) *Sea surface temperature (monthly, 1997-2016, all sites):* We used sea surface
temperature data derived from Pathfinder AVHRR (advanced very high resolution radiometer)
that were optimally interpolated on daily and 0.25 degree latitude/longitude resolution (Reynolds
et al. 2007). To produce monthly scale time series, we ran data through a 30-day moving
average filter with a rectangular, backwards looking window in order to capture general temporal
trends in the data. We spatially aggregated data using the same buffers as for chlorophyll indices.
Figure 1 illustrates mean spatial trend in winter sea surface temperature for the entire study
region.

(*v*) *Fall kelp canopy biomass (annual, 1996-2015, Santa Barbara Channel & San Diego*
only): The regional biomass of giant kelp *Macrocystis pyrifera* can fluctuate dramatically from
year to year (Bell et al. 2015). Giant kelp is a preferred food and a major constituent of *S.*
purpuratus diets in southern California (Foster *et al.* 2015). Given that fecundity in *S. purpuratus*
can vary by orders of magnitude as a result of food supply (Okamoto 2014), these fluctuations in
M. pyrifera may in part influence urchin reproduction. Thus, biomass of *M. pyrifera* was derived
from Landsat Thematic Mapper satellite imagery (Bell 2017). Data were aggregated for the
Santa Barbara Channel (including islands and mainland from Point Conception to Santa Monica
Bay through Ventura County) or the San Diego region (mainland coast from the US-Mexico
through Orange County including Santa Catalina and San Clemente Islands). We used the 3-
month running mean of kelp biomass during the period leading into the spawning season (Sept-
Nov) because marked declines in reproductive capacity require several months of consistently
low food supply (Okamoto 2014).

(*vi*) *Adult density in the Channel Islands (annual, 1997-2016, Santa Barbara Channel only):*
As an index of regional adult reproductive density of sea urchins in the Santa Barbara Channel,
we used the geometric mean of adult biomass density using survey data from The Channel
Islands KFM Program (Kushner et al. 2013). We used the same sites and data as described
above (see relationship between urchin larval recruitment and benthic juvenile recruitment) with
the exception that instead of juvenile density we used adult biomass density. This was calculated
as the biomass of adults based on: (1) the mean relationship between size and individual

biomass, (2) the size frequency of adults at a given site in a given year, (3) the total number of
urchins, and 4) the total area surveyed.

$$\text{Adult Biomass Density}_t \; = \frac{1}{S} \sum_{i=1}^{S} \left[\frac{\text{Adults Measured}_{i,t}}{\text{Tot Measured}_{i,t}} \; \frac{\text{Tot Count}_{i,t}}{\text{Tot Area}_{i,t}} \; \frac{1}{N_{i,t}} \sum_{k=1}^{N_{i,t}} f(\text{size}_{k,i,t}) \right]$$

To assess which variables exhibited correlations with larval recruitment of purple sea
urchins, we nested multiple regression analyses within a time-series modeling framework, where
coefficients varied among major regions (Fort Bragg, Santa Barbara, and San Diego sites). The
regression analyses included either all global covariates (ENSO, PDO, NPGO) or all local
covariates (temperature, chlorophyll, adult densities, kelp coverage, and the Bakun upwelling
index). We inserted a centered and standardized regression covariate matrix into the time series
model of sea urchin recruitment, replacing the explicit estimates of the interannual trends
(Supplementary Table 1). The regression coefficients were given a regularized horseshoe prior
(Carvalho *et al.* 2010; Piironen & Vehtari 2017) that biases the model coefficients towards zero
without affecting coefficients of relevant variables.

Acknowledgements

Funding was provided by the California Urchin Commission, the South Bay Cable
Committee, the California Department of Fish and Game Director's Sea Urchin Advisory
Committee, and the National Science Foundation's support of the Santa Barbara Coastal Long
Term Ecological Research program. We are indebted to Tom Ebert and John Dixon, who
initiated larval recruitment time series with SS. John Richards, Clint Nelson, Shannon Harrer,
Jenny Wolf and many students assisted with field collections, laboratory sampling and data
compilation. Sally Holbrook, Russ Schmitt and Cherie Briggs contributed discussions during
initial development of analyses. Kyle Cavanaugh, Dave Siegel and Tom Bell provided valuable
discussions about giant kelp canopy data. Kyle Cavanaugh and Rachel Simons gave important
comments on previous manuscript drafts.

Figure Captions

Figure 1: Map of larval recruitment collection locations in California. Colors represent the
average spatial gradient of sea surface. The hatched white lines show the shoreline buffer used to
constrain sea surface temperature and sea surface irradiance (the chlorophyll index) to local
larval recruitment observations.

Figure 2: Standardized annual scale trends (log-scale, standardized to one standard deviation) of
larval recruitment of purple urchins (*S. purpuratus*) from 1990 through 2016 for: (a) Fort Bragg,
(b) sites in the Santa Barbara Channel, and (c) sites in San Diego. Colors within each panel
represent individual sites associated with the legend. Interannual trends are estimated using a 3/2
Matérn Gaussian process covariance within a Bayesian multivariate state-space model
accounting for spatiotemporal correlations.

Figure 3: Seasonal trends at each site for (a) standardized larval recruitment of *S. purpuratus*, (b)
biweekly mean sea surface chlorophyll a (chla) and (c) biweekly mean sea surface temperature
(SST). The seasonal trend in larval recruitment at each site was standardized by the mean
maximum seasonal value. Chla and SST were measured using satellite imagery within 25 km of
the coastline and 150 alongshore km of the location where recruitment was sampled. See
Methods for details.

Figure 4: The relationship between the summer density of juvenile purple sea urchins on reefs in
the Santa Barbara Channel and the density of recruits on larval collectors during March- July of
the previous year (i.e., 12-20 months prior). Points and error bars represent annual means (+/-
SE) averaged over all sites. The line with the band represents the estimated relationship and
95% uncertainty interval.

Figure 5: Correlation between purple urchin recruitment at each site and (a-g) sea surface
temperature, (h-n) the multivariate ENSO index, and (m-u) the North Pacific Gyre Oscillation
(NPGO). Small grey points represent biweekly means and black points and lines represent
annual means.

Figure 6: Standardized Bayesian multiple regression coefficients for relationships between *S.*
purpuratus and local covariates (a-e) or global climatic indices (f-h). Models with global and
local variables were fit separately. Coefficients were estimated using a multiple regression
nested within a Bayesian multivariate time series model. Coefficients are a-priori biased towards
zero using a horseshoe prior. For details see Methods.

Supplementary Tables:

Eq. No	Description	Equation
Eq. 1	Biweekly estimate	$\widehat{\boldsymbol{\mu}} = \exp[\mathbf{X}\beta + \mathbf{Annual} + \mathbf{Seasonal} + \boldsymbol{\varepsilon}]^{\S}$ $\widehat{\boldsymbol{\mu}} = \exp[\mathbf{X}\beta + \mathbf{Seasonal} + \boldsymbol{\varepsilon}]^{\ddagger}$
Eq. 2	Subsampling prior	$p_{t,s,b} \sim \text{beta}\left(N_{t,s,b} + 0.5, NO_{t,s,b} + 0.5\right)$ (if subsampling was performed for brush b at time t at site s)
Eq. 3	Total count likelihood	$N_{t,s,b} \sim \text{Poisson}\left(\hat{\mu}_{t,s} \times D_{t,s,b}\right) \; if$ sample t, s, b is not subsampled $Tot_{t,s,b} \sim \text{Poisson}\left(\hat{\mu}_{t,s} \times D_{t,s,b}/p_{t,s,b}\right) \; otherwise$
Eq. 4	Prior annual temporal correlation function	$\bar{r}_{sp\langle t_i,t_j\rangle} = \left(1 + 3^2\left|t_i - t_j\right|\right)e^{-3^2\left|t_i-t_j\right|}$ (Periodic Kernel - MacKay 1998)
Eq. 5	Prior seasonal temporal correlation function	$\bar{\bar{r}}_{sp\langle t_i,t_j\rangle} = \exp\left[-\frac{2}{1.5^2}\left(0.5 - 0.5\cos\left(\pi\left|t_i - t_j\right|\right)\right)\right]$ (Matérn 3/2 Kernel - Rasmussen & Williams 2006)
Eq. 6[§]	Prior annual spatiotemporal covariance function (separable)	$\overline{\boldsymbol{\Sigma}} = \overline{\boldsymbol{\Omega}} \otimes [\overline{\boldsymbol{\sigma}}\ \mathbf{R}]$
Eq. 7	Prior seasonal spatiotemporal covariance function (separable)	$\overline{\overline{\boldsymbol{\Sigma}}} = \overline{\overline{\boldsymbol{\Omega}}} \otimes [\overline{\overline{\boldsymbol{\sigma}}}\ \overline{\overline{\mathbf{R}}}]$
Eq. 8[§]	Matrix of annual trends	$\mathbf{Annual}_{sp} \sim \text{MVN}(0, \overline{\boldsymbol{\Sigma}})$
Eq. 9	Vector of seasonal trends	$\boldsymbol{Seasonal}_{s,sp} \sim \text{MVN}\left(0, \overline{\overline{\boldsymbol{\Sigma}}}\right)$
Eq. 10	Matrix of process error covariance (spatial)	$\boldsymbol{\varepsilon}_{t,sp} \sim \text{MVN}(0, \textstyle\sum)$

457   Supplementary Table 1: Equations for the multivariate spatiotemporal models. Parameters
458   and priors are described in Supplementary Table 2.

459   [§]only included in the trend estimation model.

460   [‡]only included in the Bayesian regression model.

Symbol	Description	Type	Prior/Input
$\overline{\mathbf{\Omega}}$	Annual scale temporal correlation matrix	Input	Eq. 4
$\overline{\overline{\mathbf{\Omega}}}$	Seasonal scale temporal correlation matrix	Input	Eq. 5
$\overline{\mathbf{R}}$	Annual scale spatial correlation matrix	Estimated (via cholesky factor)	LKJ prior: scale = 2 (Lewandowski *et al.* 2009)
$\overline{\overline{\mathbf{R}}}$	Vector of site-specific annual standard deviations	Estimated	Half-Cauchy: scale = 2.5 (Polson & Scott 2012)
$\overline{\mathbf{\sigma}}$	Vector of site-specific seasonal standard deviations	Estimated	Half-Cauchy: scale = 2.5
$\overline{\overline{\mathbf{\sigma}}}$	Spatial error covariance matrix	Estimated (via cholesky factor)	LKJ prior[1]: scale = 2
Σ	Site & species specific intercepts and covariate coefficients	Estimated	Intercepts: Student-t: scale= 2 Coefficients: Horseshoe prior (Carvalho *et al.* 2010; Piironen & Vehtari 2017)
β	Site & species specific first order autoregressive parameter	Estimated	Uniform -0.95 to 0.95
ϕ_s	Total number urchins counted at time *t*, site *s,* on brush *b*	Data	
$N_{t,s,b}$	Number of subsampled urchins at time *t*, site *s,* on brush *b*	Data	
$Tot_{t,s,b}$	Purple urchins counted from the subsample $n_{t,s,b}$	Data	
$NO_{t,s,b}$	Sampling duration of brush *b* at time *t* and site *s*	Data	
X	Covariate Design Matrix	Data	

462   Supplementary Table 2: Table of parameters and data, descriptions, and priors (if estimated).

References

1. Andrew N., Agatsuma Y., Ballesteros E., Bazhin A., Creaser E., Barnes D., Botsford L., Bradbury A., Campbell A. & Dixon J. (2003) Status and management of world sea urchin fisheries. *Oceanography and Marine Biology-An Annual Review*, 40, 343-425

2. Azad A.K., Pearce C.M. & McKinley R.S. (2012) Influence of stocking density and temperature on early development and survival of the purple sea urchin, *Strongylocentrotus purpuratus* (Stimpson, 1857). *Aquac Res*, 43, 1577-1591

3. Bakun A. 1973. Coastal upwelling indices, west coast of North America, 1946-71. NOAA Tech. Rep. pp. 103.

4. Basch L.V. & Tegner M.J. (2007) Reproductive responses of Purple sea urchin (*Strongylocentrotus purpuratus*) populations to environmental conditions across a coastal depth gradient. *Bull Mar Sci*, 81, 255-282

5. Behrens M.D. & Lafferty K.D. (2004) Effects of marine reserves and urchin disease on southern Californian rocky reef communities. *Mar Ecol Prog Ser*, 279, 129-139

6. Bell T., K. Cavanaugh and D. Siegel, (2017) SBC LTER: Time series of quarterly NetCDF files of kelp biomass in the canopy from Landsat 5, 7 and 8, 1984 - 2016 (ongoing). Santa Barbara Coastal Long Term Ecological Research Project. doi:10.6073/pasta/817d2c24ebd78621869e17d94ba0df0c

7. Bell T.W., Cavanaugh K.C., Reed D.C. & Siegel D.A. (2015) Geographical variability in the controls of giant kelp biomass dynamics. *J Biogeogr*, 42, 2010-2021

8. Bertram D.F. & Strathmann R.R. (1998) Effects of maternal and larval nutrition on growth and form of planktotrophic larvae. *Ecology*, 79, 315-327

9. Blanchette C.A., Broitman B.R. & Gaines S.D. (2006) Intertidal community structure and oceanographic patterns around Santa Cruz Island, CA, USA. *Mar Biol*, 149, 689-701

10. Broitman B.R., Blanchette C.A. & Gaines S.D. (2005) Recruitment of intertidal invertebrates and oceanographic variability at Santa Cruz Island, California. *Limnol Oceanogr*, 50, 1473-1479

11. Burt J.M., Tinker M.T., Okamoto D.K., Demes K.W., Holmes K. & Salomon A.K. (2018) Sudden collapse of a mesopredator reveals its complementary role in mediating rocky reef regime shifts. *Proc R Soc B*, 285

12. Byrne M., Ho M., Selvakumaraswamy P., Nguyen H.D., Dworjanyn S.A. & Davis A.R. (2009) Temperature, but not pH, compromises sea urchin fertilization and early development under near-future climate change scenarios. *Proc R Soc B*, 276, 1883-1888

13. Cai W., Borlace S., Lengaigne M., Van Rensch P., Collins M., Vecchi G., Timmermann A., Santoso A., McPhaden M.J. & Wu L. (2014) Increasing frequency of extreme El Niño events due to greenhouse warming. *Nature climate change*, 4, 111

14. Carpenter B., Gelman A., Hoffman M., Lee D., Goodrich B., Betancourt M., Brubaker M.A., Guo J., Li P. & Riddell A. (2016) Stan: A probabilistic programming language. *J Stat Softw*, 20

15. Carvalho C.M., Polson N.G. & Scott J.G. (2010) The horseshoe estimator for sparse signals.
Biometrika, 97, 465-480

16. Chavez F., Pennington J., Castro C., Ryan J., Michisaki R., Schlining B., Walz P., Buck K.,
McFadyen A. & Collins C. (2002) Biological and chemical consequences of the 1997–
1998 El Niño in central California waters. *Progress in Oceanography*, 54, 205-232

17. Cochran R.C. & Engelmann F. (1975) Environmental regulation of the annual reproductive
season of *Strongylocentrotus purpuratus* (Stimpson). *Biol Bull*, 148, 393-401

18. Connell J.H. (1985) The consequences of variation in initial settlement vs. post-settlement
mortality in rocky intertidal communities. *J Exp Mar Biol Ecol*, 93, 11-45

19. Di Lorenzo E., Schneider N., Cobb K.M., Franks P., Chhak K., Miller A.J., McWilliams J.C.,
Bograd S.J., Arango H. & Curchitser E. (2008) North Pacific Gyre Oscillation links
ocean climate and ecosystem change. *Geophysical Research Letters*, 35

20. Ebeling A., Laur D. & Rowley R. (1985) Severe storm disturbances and reversal of
community structure in a southern California kelp forest. *Mar Biol*, 84, 287-294

21. Ebert T. (1983) Recruitment in echinoderms. *Echinoderm Stud*, 1, 169-203

22. Ebert T., Schroeter S., Dixon J. & Kalvass P. (1994) Settlement patterns of red and purple
sea urchins (*Strongylocentrotus franciscanus* and *S. purpuratus*) in California, USA. *Mar
Ecol Prog Ser*, 111, 41-52

23. Ebert T.A. (2010) Demographic patterns of the purple sea urchin *Strongylocentrotus
purpuratus* along a latitudinal gradient, 1985-1987. *Mar Ecol Prog Ser*, 406, 105-120

24. Edmunds P.J. & Carpenter R.C. (2001) Recovery of Diadema antillarum reduces macroalgal
cover and increases abundance of juvenile corals on a Caribbean reef. *Proceedings of the
National Academy of Sciences*, 98, 5067-5071

25. Esaias W.E., Abbott M.R., Barton I., Brown O.B., Campbell J.W., Carder K.L., Clark D.K.,
Evans R.H., Hoge F.E. & Gordon H.R. (1998) An overview of MODIS capabilities for
ocean science observations. *IEEE Transactions on Geoscience and Remote Sensing*, 36,
1250-1265

26. Estes J.A. & Duggins D.O. (1995) Sea otters and kelp forests in Alaska: generality and
variation in a community ecological paradigm. *Ecol Monogr*, 65, 75-100

27. Fabricius K., Okaji K. & De'Ath G. (2010) Three lines of evidence to link outbreaks of the
crown-of-thorns seastar Acanthaster planci to the release of larval food limitation. *Coral
Reefs*, 29, 593-605

28. Farmanfarmaian A. & Giese A. (1963) Thermal tolerance and acclimation in the western
purple sea urchin, *Strongylocentrotus purpuratus*. *Physiol Zool*, 36, 237-243

29. Field C.M. & Walker C. 2003. *A Site Profile of the Kachemak Bay Research Reserve, a Unit
of the National Estuarine Research Reserve System*. Kachemak Bay Research Reserve
Homer

30. Filbee-Dexter K. & Scheibling R.E. (2014) Sea urchin barrens as alternative stable states of
collapsed kelp ecosystems. *Mar Ecol Prog Ser*, 495, 1-25

31. Foster M.C., Byrnes J.E. & Reed D.C. (2015) Effects of five southern California macroalgal
diets on consumption, growth, and gonad weight, in the purple sea urchin
Strongylocentrotus purpuratus. *PeerJ*, 3, e719

32. Gonor J.J. (1973) Reproductive cycles in oregon populations of the echlnoid,
StrongylocentrotUs purpuratus (Stimpson). I. Annual gonad growth and ovarian
gametogenic cycles. *J Exp Mar Biol Ecol*, 12, 45-64

33. Hernández J.C., Clemente S., Girard D., Pérez-Ruzafa Á. & Brito A. (2010) Effect of
temperature on settlement and postsettlement survival in a barrens-forming sea urchin.
Mar Ecol Prog Ser, 413, 69-80

34. Hoegh-Guldberg O. & Pearse J.S. (1995) Temperature, food availability, and the
development of marine invertebrate larvae. *Am Zool*, 35, 415-425

35. Hoffman M.D. & Gelman A. (2014) The No-U-Turn Sampler: Adaptively Setting Path
Lengths in Hamiltonian Monte Carlo. *J Mach Learn Res*, 15, 1593-1623

36. Kato S. & Schroeter S.C. (1985) Biology of the red sea urchin, *Strongylocentrotus
franciscanus*, and its fishery in California. *Mar. Fish. Rev*, 47, 1-20

37. Kenner M. & Lares M. (1991) Size at first reproduction of the sea urchin *Strongylocentrotus
purpuratus* in a central California kelp forest. *Mar Ecol Prog Ser*, 76, 303-306

38. Kushner D.J., Rassweiler A., McLaughlin J.P. & Lafferty K.D. (2013) A multi-decade time
series of kelp forest community structure at the California Channel Islands: Ecological
Archives E094-245. *Ecology*, 94, 2655-2655

39. Lafferty K.D. (2004) Fishing for lobsters indirectly increases epidemics in sea urchins. *Ecol
Appl*, 14, 1566-1573

40. Lewandowski D., Kurowicka D. & Joe H. (2009) Generating random correlation matrices
based on vines and extended onion method. *Journal of multivariate analysis*, 100, 1989-
2001

41. Ling S.D., Johnson C., Frusher S. & King C. (2008) Reproductive potential of a marine
ecosystem engineer at the edge of a newly expanded range. *Global Change Biology*, 14,
907-915

42. Lynn R. & Bograd S. (2002) Dynamic evolution of the 1997–1999 El Niño–La Niña cycle in
the southern California Current system. *Progress in Oceanography*, 54, 59-75

43. MacKay D.J. (1998) Introduction to Gaussian processes. *NATO ASI Series F Computer and
Systems Sciences*, 168, 133-166

44. Magnusson A., Skaug H.J., Nielsen A., Berg C.W., Kristensen K., Maechler M., Bentham
K.J.v., and B.M.B. & Brooks M.E. (2017) glmmTMB: Generalized Linear Mixed Models
using Template Model Builder. R package version 0.1.3. Available at
https://github.com/glmmTMB

45. Mann K. & Breen P. (1972) The relation between lobster abundance, sea urchins, and kelp
beds. *Journal of the Fisheries Board of Canada*, 29, 603-605

46. Mantua N.J. & Hare S.R. (2002) The Pacific decadal oscillation. *Journal of Oceanography*,
58, 35-44

47. McManus M.A. & Woodson C.B. (2012) Plankton distribution and ocean dispersal. *J Exp*
Biol, 215, 1008-1016

48. Meyer E., Green A., Moore M. & Manahan D. (2007) Food availability and physiological
state of sea urchin larvae (*Strongylocentrotus purpuratus*). *Mar Biol*, 152, 179–191

49. Mitarai S., Siegel D., Watson J., Dong C. & McWilliams J. (2009) Quantifying connectivity
in the coastal ocean with application to the Southern California Bight. *J Geophys Res*,
114

50. Morgan S.G. (2014) Behaviorally mediated larval transport in upwelling systems. *Advances*
in Oceanography, 2014

51. Mumby P.J., Hastings A. & Edwards H.J. (2007) Thresholds and the resilience of Caribbean
coral reefs. *Nature*, 450, 98-101

52. Myers R.A. (1998) When do environment–recruitment correlations work? *Rev Fish Biol*
Fish, 8, 285-305

53. O'Reilly J.E., Maritorena S., Mitchell B.G., Siegel D.A., Carder K.L., Garver S.A., Kahru M.
& McClain C. (1998) Ocean color chlorophyll algorithms for SeaWiFS. *Journal of*
Geophysical Research: Oceans, 103, 24937-24953

54. Okamoto D.K. 2014. *The role of fluctuating food supply on recruitment, survival and*
population dynamics in the sea. University of California, Santa Barbara

55. Olivares-Bañuelos N.C., Enríquez-Paredes L.M., Ladah L.B. & De La Rosa-Vélez J. (2008)
Population structure of purple sea urchin Strongylocentrotus purpuratus along the Baja
California peninsula. *Fisheries science*, 74, 804-812

56. Padilla-Gamiño J.L., Kelly M.W., Evans T.G. & Hofmann G.E. (2013) Temperature and
CO2 additively regulate physiology, morphology and genomic responses of larval sea
urchins, *Strongylocentrotus purpuratus*. *Proc R Soc B*, 280

57. Pearse J.S., Pearse V.B. & Davis K.K. (1986) Photoperiodic Regulation of Gametogenesis
and Growth in the Sea Urchin *Strongylocentrotus purpuratus*. *J Exp Zool*, 237, 107-118

58. Pearse J.S. (2006) Ecological role of purple sea urchins. *Science*, 314, 940-941

59. Piironen J. & Vehtari A. (2017) Sparsity information and regularization in the horseshoe and
other shrinkage priors. *Electronic Journal of Statistics*, 11, 5018-5051

60. Polson N.G. & Scott J.G. (2012) On the half-Cauchy prior for a global scale parameter.
Bayesian Analysis, 7, 887-902

61. R Core Team (2017) R: A language and environment for statistical computing. Available at
http://www.r-project.org/

62. Rasmussen C.E. & Williams C.K.I. 2006. *Gaussian Processes for Machine Learning*. The
MIT Press, Cambridge, Massachusetts

63. Rast M., Bezy J. & Bruzzi S. (1999) The ESA Medium Resolution Imaging Spectrometer
MERIS a review of the instrument and its mission. *Int J Remote Sens*, 20, 1681-1702

64. Reynolds R.W., Smith T.M., Liu C., Chelton D.B., Casey K.S. & Schlax M.G. (2007) Daily
high-resolution-blended analyses for sea surface temperature. *J Clim*, 20, 5473-5496

65. Rowley R. (1989) Settlement and recruitment of sea urchins (*Strongylocentrotus spp.*) in a
sea-urchin barren ground and a kelp bed: are populations regulated by settlement or post-
settlement processes? *Mar Biol*, 100, 485-494

66. Runcie D.E., Garfield D.A., Babbitt C.C., Wygoda J.A., Mukherjee S. & Wray G.A. (2012)
Genetics of gene expression responses to temperature stress in a sea urchin gene network.
Mol Ecol, 21, 4547-4562

67. Sathyendranath S., Grant M., Brewin R.J.W., Brockmann C., Brotas V., Chuprin A., Doerffer
R., Dowell M., Farman A., Groom S., Jackson T., Krasemann H., Lavender S., Martinez
Vicente V., Mazeran C., Mélin F., Moore T.S., Müller D., Platt T., Regner P., Roy S.,
Steinmetz F., Swinton J., Valente A., Zühlke M., Antoine D., Arnone R., Balch W.M.,
Barker K., Barlow R., Bélanger S., Berthon J.-F., Beşiktepe Ş., Brando V.E., Canuti E.,
Chavez F., Claustre H., Crout R., Feldman G., Franz B., Frouin R., García-Soto C., Gibb
S.W., Gould R., Hooker S., Kahru M., Klein H., Kratzer S., Loisel H., McKee D.,
Mitchell B.G., Moisan T., Muller-Karger F., O'Dowd L., Ondrusek M., Poulton A.J.,
Repecaud M., Smyth T., Sosik H.M., Taberner M., Twardowski M., Voss K., Werdell J.,
Wernand M. & Zibordi G. (2018) ESA Ocean Colour Climate Change Initiative
(Ocean_Colour_cci): Version 3.1 Data. Centre for Environmental Data Analysis

68. Schroeder T. & Battaglia D. (1985) "Spiral asters" and cytoplasmic rotation in sea urchin
eggs: induction in *Strongylocentrotus purpuratus* eggs by elevated temperature. *J Cell
Biol*, 100, 1056-1062

69. Schueler C.F., Clement J.E., Ardanuy P.E., Welsch C., DeLuccia F. & Swenson H. (2002)
NPOESS VIIRS sensor design overview. In: *Earth Observing Systems VI*, pp. 11-24.
International Society for Optics and Photonics

70. Shanks A.L. (2009) Pelagic larval duration and dispersal distance revisited. *The Biological
Bulletin*, 216, 373-385

71. Shelton A.O. & Mangel M. (2011) Fluctuations of fish populations and the magnifying
effects of fishing. *Proceedings of the National Academy of Sciences*, 108, 7075-7080

72. Siegel D., Kinlan B., Gaylord B. & Gaines S. (2003) Lagrangian descriptions of marine
larval dispersion. *Mar Ecol Prog Ser*, 260, 83-96

73. Siegel D., Mitarai S., Costello C., Gaines S., Kendall B., Warner R. & Winters K. (2008) The
stochastic nature of larval connectivity among nearshore marine populations.
Proceedings of the National Academy of Sciences, 105, 8974-8979

74. Stan Development Team (2016a) Stan: A C++ library for probability and sampling. Version
2.12.0. . Available at http://mc-stan.org/

75. Stan Development Team (2016b) RStan: the R interface to Stan. Version 2.10.1. . Available
at http://mc-stan.org/rstan.html

76. Steeb W.-H. & Hardy Y. 2011. *Matrix Calculus and Kronecker Product: A Practical
Approach to Linear and Multilinear Algebra Second Edition*. World Scientific Publishing
Company

77. Strathmann M.F. 1987. *Reproduction and development of marine invertebrates of the*
northern Pacific coast: data and methods for the study of eggs, embryos, and larvae.
Univ. of Wash. Press, Seattle

78. Strathmann R. (1978) Length of pelagic period in echinoderms with feeding larvae from the
northeast Pacific. *J Exp Mar Biol Ecol*, 34, 23-27

79. Sydeman W.J., Poloczanska E., Reed T.E. & Thompson S.A. (2015) Climate change and
marine vertebrates. *Science*, 350, 772-777

80. Tegner M. & Dayton P. (1991) Sea urchins, El Niños, and the long term stability of Southern
California kelp forest communities. *Mar Ecol Prog Ser*, 77, 49-63

81. Valentine J.F. & Heck Jr K. (1999) Seagrass herbivory: evidence for the continued grazing of
marine grasses. *Mar Ecol Prog Ser*, 291-302

82. Wolter K. & Timlin M. (1993) Monitoring ENSO in COADS with a seasonally adjusted
principal component index. In: *Proc. of the 17th Climate Diagnostics Workshop*, pp. 52-
57, NOAA/NMC/CAC, NSSL, Oklahoma Clim. Survey, CIMMS and the School of
Meteor., Univ. of Oklahoma, Norman, OK

83. Wolter K. & Timlin M.S. (1998) Measuring the strength of ENSO events: How does 1997/98
rank? *Weather*, 53, 315-324

84. Wong J.M., Johnson K.M., Kelly M.W. & Hofmann G.E. (2018) Transcriptomics reveal
transgenerational effects in purple sea urchins exposed to upwelling conditions, and the
response of their progeny to differential pCO2 levels. *Mol Ecol*

85. Woodson C.B., McManus M.A., Tyburczy J.A., Barth J.A., Washburn L., Caselle J.E., Carr
M.H., Malone D.P., Raimondi P.T., Menge B.A. & Palumbi S.R. (2012) Coastal fronts
set recruitment and connectivity patterns across multiple taxa. *Limnol Oceanogr*, 57, 582-
596

86. Yeh S.-W., Kug J.-S., Dewitte B., Kwon M.-H., Kirtman B.P. & Jin F.-F. (2009) El Niño in a
changing climate. *Nature*, 461, 511

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Information - Appendices for "Geographically Opposing Responses of Sea Urchin Recruitment to Changes in Ocean Climate"

Daniel K. Okamoto[*1,2], Daniel C. Reed[†3], and Stephen Schroeter[‡3]

[1]Department of Biological Science, Florida State University, Tallahassee, Florida 32303
[3]Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106 USA
[3]Marine Science Institute, University of California Santa Barbara, Santa Barbara, CA 93106 USA

[*]corresponding author: dokamoto@bio.fsu.edu
[†]email: dan.reed@lifesci.ucsb.edu
[‡]email: schroete@lifesci.ucsb.edu
[*]corresponding author: dokamoto@bio.fsu.edu
[†]email: dan.reed@lifesci.ucsb.edu
[‡]email: schroete@lifesci.ucsb.edu

Figure S.1: Observed and estimated purple urchin (*S. purpuratus*) settlement trends from 1990-2016 for each site. Large points represent empirical biweekly means (non-zero means across all brushes and collection times), lines on the axis represent biweekly mean values equal to zero, small points represent counts on individual brushes (non-zero only), red lines represent biweekly estimates, lines with blue 95% uncertainty intervals represent model estimates of the interannual trends. The model estimates the seasonal trend and the interannual trend at each site simultaneously. For reference, interannual trends for *S. purpuratus* shown here are the unstandardized versions of those shown in Figure 2.

Figure S.2: Seasonal settlement trends from 1990-2016 with the mean estimated seasonal trend for each site. Large points represent empirical biweekly means (non-zero means across all brushes and collection times), lines on the axis represent biweekly mean values equal to zero, small points represent counts on individual brushes (non-zero only), black lines represent the posterior mean and bands represent the 95% uncertainty interval. The model estimates the seasonal trend and the interannual trend at each site simultaneously. For reference, seasonal trends for *S. purpuratus* shown here are the unstandardized versions of those shown in Figure 3.

Figure S.3: Correlation between coefficients for ENSO and PDO at Fort Bragg creating variance inflation. Contours represent the 95%, 90% and 80% quantile estimates from 3000 posterior samples. Note that when one is at or near zero, the other is centered away from zero.

Variable Correlations

To illustrate the intercorrelated nature of the covariates and responses, we estimated a network model for Gaviota Pier that includes the independent model estimates of *S. purpuratus* settlement and its seasonality. To do so, we first constrained network structure by excluding all nonsensical interactions (i.e. chlorophyll does not cause ENSO events and is thus eliminated a priori) and including all known directional interactions (i.e. estimated seasonality affecting settlement for is forced into the network). Learning of the network skeleton is achieved via the Hill Climbing algorithm using the bnlearn[1] package in R. Note that strong collinearity can result in the weaker correlation being ignored (e.g. ENSO over SST with Larval Recruitment).

Figure S.4: Bayesian network model structure inferred for the Santa Barbara Channel. Numbers represent the partial correlation coefficients accounting for all other variables pointing to that node. All correlations shown are significant at alpha= 0.05

---

[1]Marco Scutari (2010). Learning Bayesian Networks with the bnlearn R Package. Journal of Statistical Software, 35(3), 1-22. URL http://www.jstatsoft.org/v35/i03/

```
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
\\ Stan Code Required to Estimate Trends      \\
\\ Author: D.K. Okamoto                        \\
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\

data {
  // input information
  int NO; // number of observations
  int NS; // number of sites
  int NM; // total number of periods per site
  int NYM; // number of periods within a seasonal trend
  int NSUB; // number of observations with subsamples
  // input data
  vector[NSUB] P1; // SP counts +0.5
  vector[NSUB] P2; // Other urchin counts + 0.5
  int SUB[NSUB]; // indicator variable for which observations are subsampled
  int N[NO]; // total count (on the original scale)
  int OBS[NO]; // continuous month-site observation index
  int PRED_MONTH[NM]; // month-site observation index associated with MA
  vector[NO] D; // number of days
  cholesky_factor_cov[NYM] D_seas;   // seasonal cholesky covariance factor
  cholesky_factor_cov[NM] D_ann;   // annual cholesky covariance factor
}

parameters {
  // mean volatility
  vector<lower=1e-7>[NS] sigma_S_mu;

  // spatial volatility correlation matrix
  cholesky_factor_corr[NS] L_Omega_S; //
  cholesky_factor_corr[NS] L_Omega_Spatial;

  // seasonal and annual variance
  vector<lower = 0>[NS] sigma;
  vector<lower = 0>[NS] sigma2;

  // mean site settlement
  row_vector<lower= -10, upper= 10>[NS] mu_S;

  // iid errors for seasonality
  matrix<lower= -10, upper= 10>[NYM,NS] z_s;

  // iid errors for annual trends
  matrix<lower= -10, upper= 10>[NS,NM] w_z;

  // iid errors for residuals
  matrix<lower= -4, upper= 4>[NS,NM] e_z;

  // proportion of samples that are purps
  vector<lower= 0,upper= 1>[NSUB] theta;
}

transformed parameters {
  matrix[NYM,NS] LS;  // site specific seasonal trends
  matrix[NM,NS] S;    // estimated abundance
  vector[NO] S_exp;   // normal scale estimated sp for each observation
  vector[NO] theta2;  // fraction that is SP
  matrix[NM,NS] w;    // spatially and temporally correlated, standardized annual
      trends
```

```
  matrix[NM,NS] e;     // spatially correlated process errors

  LS  = D_seas*z_s;
  w = D_ann*transpose(L_Omega_Spatial*w_z);
  e = transpose(diag_pre_multiply(sigma_S_mu,L_Omega_S)*e_z);

  theta2 = rep_vector(1.0,NO);
  theta2[SUB] = theta;

  // caculate expected values
  S= rep_matrix(mu_S,NM)+diag_post_multiply(LS[PRED_MONTH,],sigma)+
     diag_post_multiply(w,sigma)+e;
  S_exp = exp(to_vector(S)[OBS]) .*D ./theta2;
}

model {
  // site SDs for annual, seasonal trends and process errors
  sigma ~ cauchy(0, 0.5);
  sigma2 ~ cauchy(0, 0.5);
  sigma_S_mu ~ cauchy(0,2.5);

  // spatial correlations for annual and seasonal trends
  L_Omega_S ~ lkj_corr_cholesky(2);
  L_Omega_Spatial ~ lkj_corr_cholesky(2);

  for(i in 1:NS){
    z_s[i]~ normal(0,1);
    mu_S[i] ~ normal(0,5); // site specific log-scale means
  }

  // uncertainty on proportions of purples in the sample
  // jeffrey's prior given subsampling
  theta~beta(P1, P2);

  // observation likelihood
  N ~ poisson( S_exp );
}

generated quantities {
  corr_matrix[NS] Omega_S;
  corr_matrix[NS] Omega_Spatial;
  Omega_S = multiply_lower_tri_self_transpose(L_Omega_S);
  Omega_Spatial = multiply_lower_tri_self_transpose(L_Omega_Spatial);
}
```