Effects of Ocean Climate on Spatiotemporal Variation in Sea Urchin Settlement and Recruitment

Daniel K. Okamoto[*1,2], Stephen Schroeter[†3], and Daniel C. Reed[‡3]

[1]Department of Biological Science, Florida State University, Tallahassee, Florida 32303
[2]Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106 USA
[3]Marine Science Institute, University of California Santa Barbara, Santa Barbara, CA 93106 USA

[*]corresponding author: dokamoto@bio.fsu.edu
[†]email: schroete@lifesci.ucsb.edu
[‡]email: dan.reed@lifesci.ucsb.edu

Abstract

Sea urchins are voracious herbivores that influence the ecological structure and function of nearshore ecosystems throughout the world. Like many species that produce planktonic larvae, their recruitment is thought to be particularly sensitive to climatic fluctuations in temperature that directly or indirectly affect adult reproduction and larval transport and survival. Yet how climate alters sea urchin populations in space and time by modifying larval recruitment and year-class strength on the time-scales that regulate populations remains understudied. Using an unprecedented, spatially replicated weekly-biweekly dataset spanning 27 years and 1100 km of coastline, we characterized seasonal, interannual, and spatial patterns of larval settlement of the purple sea urchin (*Strongylocentrotus purpuratus*). We show that large spatial differences in temporal patterns of larval settlement were associated with different responses to fluctuations in ocean temperature and climate. Importantly, we found a strong correlation between larval settlement and regional year class strength suggesting that such temporal and spatial variation in settlement plays an important role in controlling population dynamics. These results provide strong evidence over extensive temporal and spatial domains that climatic fluctuations shape broad-scale patterns of larval settlement and subsequent population structure of an important marine herbivore known to control the productivity and function of marine ecosystems.

Introduction

Large scale climate oscillations (e.g., El Niño/Southern Oscillation, North Atlantic Oscillation) lead to changes in ocean temperature, biogeochemistry and the severity and frequency of disruptive events that affect ocean circulation, upwelling and primary productivity (Cai *et al.* 2014; Mantua *et al.* 1997). Such shifts impose wide-reaching ecological impacts, in part by altering animal recruitment and food web structure in space and time (Sydeman *et al.* 2015). Hence, understanding how climate variability alters the recruitment of marine species is particularly important for effective conservation and management of the ocean's resources. Yet for recruitment variability, climatic fluctuations give rise to shifts in numerous factors that shape both adult reproduction and larval supply, including primary productivity, temperature, and advection and transport. Thus, significant challenges remain in achieving such understanding for benthic species with planktonic larvae due to the substantial effort needed to characterize spatial and temporal variation in larval settlement and the numerous sensitive vital rates that contribute to it.

For benthic species like sea urchins, understanding causes and consequences of recruitment variability has both ecological and economic implications. Sea urchin grazing can alter the structure of some of the world's most diverse and productive marine ecosystems, including coral reefs (Edmunds & Carpenter 2001), seagrass meadows (reviewed by Valentine & Heck Jr 1999) and kelp forests (reviewed by Filbee-Dexter & Scheibling 2014). In addition, sea urchins form the basis of important nearshore fisheries in many regions of the world (e.g. Andrew *et al.* 2003; Kato & Schroeter 1985). As a result, climate-driven changes in sea urchin

populations have the potential to profoundly affect the functioning of marine ecosystems and

the economic value of the fisheries that they support. Much of the research on controls of sea

urchin population dynamics has focused on the roles of predation and disease in controlling

adult abundance and their cascading influence on community structure (e.g. Burt *et al.* 2018;

Estes & Duggins 1995; Filbee-Dexter & Scheibling 2014; Lafferty 2004). Yet short-term

empirical studies (months to a few years) suggest that environmentally regulated larval supply

is likely an important driver of adult urchin dynamics (Hernández *et al.* 2010; Ling *et al.* 2009).

Despite the widespread recognition of the importance of recruitment variation in controlling

population fluctuations in many marine species (Shelton & Mangel 2011), relatively few studies

have examined the biotic and abiotic processes controlling the supply of sea urchin larvae in

nature (but see Hernández *et al.* 2010; Ling *et al.* 2009), and the degree to which they affect the

abundance and dynamics of older life stages over time scales that impact population dynamics.

Fluctuations in climate can affect spatial and temporal patterns of larval supply by

influencing the production of larvae by benthic adults, transport of larvae to adjacent habitats,

and the survival of larvae in the plankton. Increases in ocean temperature can impact larval

production by: (1) increasing adult mortality via the spread of water-borne pathogens

(reviewed by Feehan & Scheibling 2014), and (2) reducing adult fecundity and inhibiting

gametogenesis by altering food quantity and quality (Basch & Tegner 2007; Cochran &

Engelmann 1975; Foster *et al.* 2015; Okamoto 2014). Because sea urchins produce feeding

larvae that spend weeks to months in the plankton, increases in ocean temperature can also

affect larval development, growth and survival, either directly, or indirectly by altering the

availability of their phytoplankton food source (Bertram & Strathmann 1998; Byrne *et al.* 2009;

Hoegh-Guldberg & Pearse 1995; Strathmann 1987). Finally, climate related changes in patterns

of ocean circulation can affect the transport of larvae from source to destination (but see

Morgan 2014; Siegel *et al.* 2008). Thus, the effects of climatic change on sea urchin recruitment

represent cumulative impacts on adult abundance and reproduction, complex current patterns

that transport larvae, behavioral responses of larvae, and larval development and survival.

Because patterns of ocean temperature, circulation and upwelling can vary dramatically in

space, the effects of climate oscillations on sea urchin recruitment potentially vary over large

spatial scales. A dearth of long-term, high frequency, spatially extensive data has prevented

characterizing temporal and spatial variability in larval settlement in sea urchins, the degree to

which it is explained by different sources of environmental variation, and the relative

importance of these drivers in accounting for fluctuations in population size.

Here we analyzed a unique 27-year weekly to biweekly time series of the recruitment of

newly metamorphosed larvae (hereafter referred to as larval settlement) at sites distributed

across 1100 km of coast in California to investigate sources of spatial and temporal variability in

larval settlement of the purple sea urchin *Strongylocentrotus purpuratus*. Our objectives were

to: (1) quantify variation in larval settlement across different temporal and spatial scales; (2)

evaluate whether larval settlement on artificial substrates predicts year-class strength in

natural populations; and (3) determine the relative importance of adult abundance, larval and

adult food supply, sea surface temperature and broad-scale fluctuations in ocean climate in

contributing to the observed variation in larval settlement

 Study system

Populations of the purple sea urchin (*Strongylocentrotus purpuratus*) occupy shallow

subtidal and intertidal rocky substrata from at least 27°N on the western coast of the Baja

Peninsula (Olivares-Bañuelos *et al.* 2008) to at least 59°N on the Kenai Peninsula in Alaska (Field

& Walker 2003). Purple urchins are broadcast spawners and the seasonality of their spawning

is generally thought to be driven by spring photoperiod and temperature (Cochran &

Engelmann 1975; Gonor 1973; Pearse *et al.* 1986). Results from the field and laboratory

indicate that the thermal upper limit to completion of gametogenesis may lie around 17°C

(Basch & Tegner 2007; Cochran & Engelmann 1975). Fertilized zygotes develop into planktonic

echinoplutei larvae that feed exclusively on phytoplankton (Strathmann 1987). After spending

several weeks to months in the plankton, individuals begin final metamorphosis and settle to

the benthos (Strathmann 1978) at a size of ~500 μm in diameter (Okamoto unpublished data).

Larval settlement varies dramatically among locations at both small and large spatial scales

(Ebert 2010). Once settled, fully competent individuals occupy cobble and other complex

substrata and between 12-24 months become visible in benthic surveys at 1-2 cm diameter.

Collection of newly settled urchins along the coast of California, USA (1990-2016)

Settlement of newly metamorphosed purple sea urchins was sampled in three major

regions along the California coast from 1990 through 2016, with a total of 54,588 replicate

observations. Sampling regions (from south to north) included two sites in San Diego County

(Scripps Pier and Ocean Beach Pier), four sites in the Santa Barbara Channel (Anacapa Island,

Stearns Wharf, Ellwood Pier and Gaviota Pier) and one site at Point Cabrillo in Fort Bragg (Figure

1). San Diego and the Santa Barbara Channel lie within the Southern California Bight and Fort

Bragg is in northern California. At each site, newly settled urchins were collected using nylon-

bristled scrub brushes (2.5 cm long bristles and a 6 x 9 cm wooden base) suspended 1 to 2 m

from the benthos (Ebert *et al.* 1994). The majority of deployments included 4-8 replicate

brushes collected weekly at each site from 1990 to 2003, and biweekly thereafter through

2016. Upon collection brushes were transported to the laboratory in plastic bags and rinsed

through a 350 μm mesh sieve. Purple urchins were sorted from other species, counted and

preserved. See (Ebert *et al.* 1994) for further details.

Methods

Spatio-temporal trends in larval settlement

We estimated seasonal, annual, and spatial patterns in larval settlement using an

integrated spatiotemporal Bayesian model that accounted for the spatially and temporally

intercorrelated and heterogeneous nature of the multivariate time series. We used a Bayesian

state-space formulation for the model for several reasons. First, episodic periods of low

replication, low observation numbers, missing data, or slight variation in the sampling interval

mean that the true number of settlers may not be reflected in the empirical mean value from

brush data (i.e. the true value is not always observable). Second, when brushes contained

hundreds or more individual urchins, species identification consisted of subsampling urchins to

estimate proportions of individual species (in this case *S. purpuratus* versus the more rare red

urchin, *Mesocentrotus franciscanus*). This subsampling routine requires accounting for the

uncertainty in the sampled species ratios in the estimation of settler abundance, which we did

by incorporating a Bayesian prior in the form of a Beta distribution with hyperparameters as the

observed ratio and total subsample count (the Jeffrey's prior for ratios from binomial counts).

Third, initial examination of the data indicated that there were strong temporal components

(interannual and seasonal trends, substantial and multiplicative among-sample variability

within each period, and serial autocorrelation) and spatial components that needed to be

simultaneously accounted for.

To account for the above described nuances of the data we estimated biweekly

settlement density ($\hat{\mu}_{t,l}$) as the sum of a site-specific mean ($\beta_{0,l}$), log annual trend

($Annual\_Trend_{s,t}$), log seasonal trend ($Seas\_Trend_{t,l}$), and a spatially correlated lognormal

process error ($\varepsilon_{t,l}$);

Eq. 1 $\ln \hat{\mu}_{t,l} = \beta_{0,l} + \ln Annual\_Trend_{t,l} + \ln Seasonal\_Trend_{t,l} + \varepsilon_{t,l}$ (Trend Equation)

We estimated trends on the log-scale so that process errors, and seasonal and interannual

trends were multiplicative and strictly positive on the original scale. We used a Poisson

observation likelihood to link the biweekly mean estimate from the trend equation ($\hat{\mu}_{t,l}$ -see

Tables 1 and 2) to observed counts of larval settlers (N) for each brush (b) within each site

within each year .

The annual trends were assumed to be correlated in both space and time, where spatial

and temporal covariances were assumed to be independent. The annual spatial covariance was

assumed to be unstructured (i.e. no formal distance decay structure) because we only had

seven sites. The annual temporal covariance was determined by a Matérn 3/2 kernel because it

provided sufficient flexibility to capture interannual trends. The seasonal trend (log-scale)

within each site was estimated using a seasonal periodic temporal kernel within the model to

capture the cyclical nature of seasonality. The process error ($\varepsilon_{t,l}$) was modeled with an

unstructured spatial covariance to account for correlations among sites in their deviations from

model expectations. We estimated the model posteriors using Stan (Carpenter *et al.* 2016) with

three 1000 iteration chains after a 1000 iteration burn-in. Stan model code is provided in the

supplement. For model equations and symbology see Tables 1 and 2.

Relationship between benthic year-class strength and larval settlement.

If year class strength of sea urchins is limited by larval supply, then we should see

increases (decreases) in the abundance of juveniles in years following anomalously high (low)

larval settlement. We therefore tested whether larval settlement in the Santa Barbara Channel

(mean of Anacapa, Stearns Wharf, Ellwood Pier and Gaviota Pier) was predictive of subsequent

juvenile recruitment (hereafter referred to as "year-class strength") on natural reefs in the

region. We calculated densities of juveniles (<2cm test diameter) from the Channel Islands Kelp

Forest Monitoring (KFM) Program (Kushner *et al.* 2013), including only the XXX sites with time

series extending from 1990 through 2016. We examined the relationship between larval

settlement and year-class strength using a generalized linear mixed effects model (GLMM) with

a negative binomial likelihood (a Poisson likelihood indicated overdispersion) and survey site as

a random effect. We tested the hypothesis of no correlation using a likelihood ratio test

comparing the model with versus without settlement as a covariate. Models were estimated

using glmmTMB (Magnusson *et al.* 2017) in R (see Electronic Supplement for further details).

Relationships between larval settlement and biotic and abiotic conditions

We estimated the strength of relationships between biweekly larval settlement and

various hypothesized physical and biological drivers using an integrated regression model that

included a regularized regression within a Bayesian time-series modelling framework (Fig. 2).

Specifically, the model considered estimated log biweekly density ($\ln \hat{\mu}_{t,l}$) as a function of the

centered and standardized covariates (denoted by the vector $x_{t,l}$) while directly estimating and

accounting for seasonal trends (Seasonal_Trend$_{t,l}$), and separable spatially and temporally

correlated, multivariate lognormal process error ($\varepsilon_{t,l}$):

Eq. 2 $\ln \hat{\mu}_{t,l} = \beta_{0,l} + [x_{t,l}]'\beta_l + \ln \text{Seasonal\_Trend}_{t,l} + \varepsilon_{t,l}$ (Regression Equation)

We used a Poisson observation likelihood to link the latent trends $\hat{\mu}_{t,l}$ to the observations (see

Tables 1, 2) and we assumed the temporal correlation in the process error followed a first order

autoregressive model.

Because these analyses were correlative, we focused on unbiased parameter estimation

rather than hypothesis testing or model comparison per se. We estimated two separate

models: one included all locally measured (i.e. for a given site) environmental variables and the

other included a common set of composite indices of oceanographic climate ("global

covariates" – ENSO Index, Pacific Decadal Oscillation, North Pacific Gyre). Because inclusion of

numerous covariates can cause overfitting that leads to bias and uncertainty in the explanatory

power of the covariates, we erred on the side of sparsity and assigned the vector of regression

coefficients (β_l) a regularized horseshoe prior (i.e. the Finnish Horseshoe). Sparsity in this case

assumed that only a few of the covariates were meaningful without a priori knowledge of which

covariates were relevant and which were not. A sparse regression encoded this assumption to

allow the data to inform which covariates were relevant and how they were correlated with the

response variable. The Finnish horseshoe prior is a Bayesian version of the lasso (Carvalho *et al.*

2010; Piironen & Vehtari 2017) that produced a data-driven reduction in the influence of

weaker covariates by regularization of those coefficients given the data. For full details and

model equations, see Tables S.1 and S.2.

Below are the covariates used in the integrated regression model:

(i) Oceanographic climate indices (monthly, 1990-2016, all sites): We used three major

global indices of oceanographic climate. The multivariate El Niño Southern Oscillation Index

(MEI), the Pacific Decadal Oscillation (PDO - Mantua & Hare 2002) and the North Pacific Gyre

Oscillation (NPGO - Di Lorenzo *et al.* 2008). The MEI provides a metric of the intensity of El

Niño Southern Oscillation (ENSO) fluctuations, which persist for 6-18 months and explains much

of the oceanographic variability in the tropics (Di Lorenzo *et al.* 2013; Wolter & Timlin 1993;

Wolter & Timlin 1998). In contrast, the PDO and NPGO exhibit longer-period fluctuations,

explain much of the oceanographic variability in higher latitudes, and the low frequency

variability of these metrics are shaped through ENSO forcing of the Aleutian Low and the North

Pacific Oscillation, respectively (Di Lorenzo *et al.* 2013).

(ii) Coastal upwelling index (monthly, 1997-2016, all sites): The Bakun index (Bakun

1973) provides an index of large-scale coastal upwelling and specifically describes the volume of

water that is transported offshore from Ekman transport (http://www.pfel.noaa.gov/ - sites =

33N-119W and 39N-125W. This index has been used as a large-scale proxy for processes that

may favor larval retention and advection from shore in addition to its value as a predictor of

coastal productivity (Menge *et al.* 2011; Menge & Menge 2013; Shkedy & Roughgarden 1997;

Wing *et al.* 2003).   We emphasize that this index is not a location specific metric nor does it

consider fine-scale nearshore hydrodynamic processes that are also likely to affect larval

retention (Fisher *et al.* 2014; Morgan *et al.* 2009; Morgan *et al.* 2016; Morgan *et al.* 2018;

Shanks & Shearman 2009; Shanks & Eckert 2005; Shanks & Morgan 2018; Shanks *et al.* 2017)

which remain outside the scope of this study but worthy of investigation as drivers of

settlement trends. Rather we include the Bakun index here as an index of broader scale coastal

upwelling that some argue affect trends in larval supply.

(iii) Sea surface chlorophyll (monthly, 1997-2016, all sites): Satellite imagery of sea

surface chlorophyll a provides a spatially and temporally resolved estimate of phytoplankton

biomass (mg m^{-3}) that is not available from in situ sampling. We used version 3.1 of the OC-CCI

merged ocean color time series (Sathyendranath *et al.* 2018) that combines SeaWIFs, MERIS,

MODIS and VIIRS to provide the temporal and spatial coverage required for this study. For each

larval settlement collection site, we aggregated data into a 30-day moving average (30 days

prior to brush collection) within a buffer of 10 km from shore that stretched 150 km alongshore

(the average Lagrangian estimate for dispersal distances for species with a planktonic larval

duration (PLD) of 30-days (but see Shanks 2009; Siegel *et al.* 2003)). For the site at Anacapa

Island, we included any point within a 150 km radius and within 10 km of any coastline. We

used 30-day moving averages for chlorophyll and temperature (below) because larvae were

exposed to these conditions for at least 30 days prior to settlement, and because averaging

over 30 days minimized the effects of serial autocorrelation in the data.

242   (*iv) Sea surface temperature (monthly, 1997-2016, all sites):* We used the 30-day moving

243   average of sea surface temperature, derived from Pathfinder AVHRR (Reynolds *et al.* 2007)

244   (advanced very high resolution radiometer) that was optimally interpolated at daily and 0.25

245   degree latitude/longitude resolution. We spatially aggregated data in the same way as sea

246   surface chlorophyll.

248   *(v) Fall kelp canopy biomass (annual, 1996-2015, Santa Barbara Channel & San Diego*

only): Giant kelp (*Macrocystis pyrifera*) is a preferred food and a major constituent of *S.*

purpuratus diets in southern California (Harrold & Reed 1985). The regional biomass of giant

kelp *Macrocystis pyrifera* can fluctuate dramatically from year to year (Bell *et al.* 2015) and

cause orders of magnitude variations in *S. purpuratus* fecundity (Okamoto 2014). We estimated

intra and interannual variability in the biomass of giant kelp from Landsat Thematic Mapper

satellite imagery (Bell 2017). Data were aggregated for the Santa Barbara Channel (including

islands and mainland from Point Conception to Santa Monica Bay through Ventura County) or

the San Diego region (mainland coast from the US-Mexico through Orange County including

Santa Catalina and San Clemente Islands). We used the 3-month running mean of kelp canopy

biomass during the period leading into the spawning season (Sept-Nov) because marked

declines in reproductive capacity require several months of consistently low food supply

(Okamoto 2014).

(vi) Adult sea urchin density in the Channel Islands (annual, 1997-2016, Santa Barbara

Channel only): The abundance of adult sea urchins can also fluctuate over time, which can

affect larval production and supply. We used the spatial geometric mean of adult biomass

density (kg m^{-2}) of purple sea urchins (calculated from surveys of density and size structure

from the Channel Islands Kelp Forest Monitoring program and a size-biomass model from the

Santa Barbara Coastal LTER) as an index of the density of adult purple sea urchins in the Santa

Barbara Channel. We used the geometric mean (exponentiated log-scale mean) as it accounts

for spatial differences in overall mean abundance among sites. We use a biomass index that

incorporates size structure and density instead of justdensity because individual maximum

fecundity increases with size. For details on calculation of biomass from the KFM data see the

Electronic Supplement.

We allowed regression coefficients to vary among major regions (Fort Bragg, Santa

Barbara, and San Diego sites). We used this level of inference because the covariates are

aggregated over spatial regions that encompass all sites within a region due to the long

planktonic larval duration of sea urchins. The regression analyses included either all global

covariates (ENSO, PDO, NPGO) or all local covariates (temperature, chlorophyll, adult densities,

giant kelp biomass, and the Bakun upwelling index). We conducted separate analyses for these

two groups of covariates because global indices were directly correlated with local covariates,

or in the case of temperature were partially derived from them. Thus, our analyses provided

non-mechanistic explanatory variables for global covariates and mechanistic explanatory

variables for local covariates.

Results

Spatio-temporal trends in larval settlement

Substantial interannual variability in larval settlement of *S. purpuratus* was observed

among years (Figure 3, Appendix Figure S1). Fluctuations in larval settlement were highly

synchronous among sites within each of the two regions in the Southern California Bight (r =

0.73 and 0.85 for sites within the Santa Barbara Channel and San Diego, respectively; Figure 3b,

c). Within the Santa Barbara Channel, pairwise correlations in interannual trends involving

Gaviota, Ellwood and Stearns Wharf were higher (r =0.86 to 0.90) than those involving Anacapa

(r = 0.46 to 0.71). Anacapa began to decline in 2012, while the declines at the other sites did

not begin until 2014 (Figure 3b, Appendix Figure S.1 b-e). While all of the Santa Barbara

Channel sites except Anacapa exhibited eight continuous years of above average settlement

following the low in 2005, San Diego sites remained mostly below average after 2003(Figure 3c,

Appendix Figure S.1 f-g). This trend produced modest synchrony among San Diego and Santa

Barbara sites from 1991 through 2007 (mean r = 0.68), but none thereafter (mean r= -0.06). In

contrast, interannual trends in larval settlement at Fort Bragg were largely uncorrelated

throughout the time series with sites in the Santa Barbara Channel (Figure 3a vs. 3b, mean r = -

0.23) and sites in San Diego (Figure 3a vs. 3c; mean r = 0.12).

Larval settlement in southern California (San Diego and the Santa Barbara Channel) was

highly seasonal, with similar patterns among sites (Figure 4, Appendix Figure S2). On average

90% of recruitment occurred from March to July with a single peak in late April/early May

(Figure 4a). By contrast, recruitment at Fort Bragg in northern California extended over a

longer period of time (90% occurred, between January and September) and typically included

two peaks per year (a large peak around July and a smaller peak in February and March; Figure

4a). The seasonal peaks in settlement in southern California coincided with peaks in sea surface

chlorophyll (Figure 4a vs. 4b.) and troughs in sea surface temperature (Figure 4a vs. 4c). At Fort

Bragg in northern California the primary peak occurred slightly after the peak in chlorophyll a

(Figure 4a. vs. 4b).

Relationship between benthic year-class strength and larval settlement.

Recruitment of juvenile purple urchins at shallow subtidal reefs in the Santa Barbara

Channel exhibited a significant, positive correlation with larval settlement to brushes two years

prior (Figure 5**,** negative binomial GLMM - $\chi^2_{df=1}$ = 9.48, p = 0.002). Years with the highest larval

settlement corresponded with a nearly three-fold increase in the mean density of juvenile

urchins two summers later (Figure 5). Over the time series, average settlement densities in the

Santa Barbara Channel varied by more than three orders of magnitude among years. This

interannual variation in settlement (averaged across the sites in the Santa Barbara Channel)

corresponded to, on average, more than a three-fold increase in the density of benthic

juveniles.

Relationships between larval settlement and biotic and abiotic conditions

Larval settlement of purple urchins showed strong correlations with local sea surface

temperature (SST, Figure 6 a-g, Figure 7 a) as well as the major climate indices (e.g. ENSO - Fig 6

h-n, Figure 7f). For temperature and ENSO, the sign of the correlation at Fort Bragg was

opposite of that at sites in the Santa Barbara Channel and San Diego. The correlation between

settlement and temperature varied from strongly negative at sites in San Diego and Santa

Barbara to positive at the Fort Bragg site. Correlations between larval settlement and adult

urchin density, upwelling, chlorophyll and kelp biomass were either not different from zero or

were opposite of that expected at all sites (Figure 7 b-e). Larval settlement and sea surface

chlorophyll were uncorrelated at Fort Bragg and in the Santa Barbara Channel and negatively

correlated in San Diego – the opposite of the hypothesized relationship that more food in the

plankton would lead to higher larval settlement (Figure 7 d). The network of correlations

among these variables are depicted in (Appendix Figure S.3).

Larval settlement in southern California was orders of magnitude lower during warm, El

Niño conditions and during the negative phase of the NPGO with the more southern sites in San

Diego responding more strongly to temperature than those further north in the Santa Barbara

Channel (Figure 7 a). In contrast, Fort Bragg, the northern most site, responded positively to

warmer El Niño conditions, negatively to the NPGO, and positively to the PDO. Importantly,

relationships observed between larval settlement and ENSO and temperature in southern

California were opposite of those observed at Fort Bragg (Figure 7 f, g). The correlations

between larval settlement and SST and ENSO occurred on roughly 3-5 year cycles, while the

correlation with NGPO and PDO occurred on decadal scales.

Discussion

Climate variability can significantly impact marine ecosystems by affecting recruitment

which in turn influences the dynamics of populations and communities. Yet determining which

species and communities will respond to shifting climate remains a difficult task. Using a multi-

decadal, high frequency, and spatially extensive time series we show how large-scale climatic

variation can give rise to spatially different responses in the settlement of larval sea urchins.

We found that settlement patterns were synchronous within Santa Barbara and San Diego and

synchronous among regions until the late 2000s. Importantly, these sites responded negatively

to elevated temperature and ENSO events. This contrasted sharply with settlement at our

northern California site (Fort Bragg) which was positively correlated to temperature and ENSO.

This difference was most obvious during the two strongest ENSO events in 1998 and 2014,

when settlement responded dramatically in opposite directions.

While many recruitment-environment correlations often break down over time (Myers

1998), the correlations between oceanographic factors and larval settlement in our study were

evident over a 27-year period that included multiple major ENSO events and several minor

ones. We observed repeatable patterns of diminished larval settlement in response to

temperature and ENSO events in southern California with opposite responses at Fort Bragg in

northern California. Others have found larval settlement in sea urchins to be positively or

negatively related to temperature depending on the species (Ebert 1983; Hernández *et al.*

2010) (Ebert 1983, Himmelman 1986 Hernandez et al. 2010). Our study is among the first to

show that the relationship between larval settlement and temperature can vary dramatically

within a species depending on location. Although sea urchin larval settlement data at the fine

temporal scale of our study are not available prior to 1990, there is a history of evidence for

ENSO-related recruitment failures in sea urchin populations off southern California. Between

1969 and the early 1980's, recruitment of juvenile sea urchins was anomalously low during El

Niño years (Ebert 1983; Tegner & Dayton 1991). Between 1969 and 1977, juvenile recruitment

was lowest during the El Niño years of 1970, 1973, and 1977 at False Point, California near San

Diego (Ebert 1983; Tegner & Dayton 1991) and at nearby Point Loma, low recruitment followed

the 1982-83 El Niño event (Ebert 1983; Tegner & Dayton 1991). How patterns of larval

settlement change in the future will undoubtedly depend on the processes underlying their

associations with ENSO events.

Climate related fluctuations in the supply of larvae have been hypothesized to affect

coastal species around the world (Caley *et al.* 1996; Grosberg & Levitan 1992; Underwood &

Fairweather 1989). In the case of sea urchins, larval supply has been linked to dramatic

changes in the state of benthic communities (Estes & Duggins 1995; Hernández *et al.* 2010; Ling

et al. 2009). Yet empirically demonstrating such links has historically proved challenging

because recruitment can be attenuated by high mortality of newly settled larvae (Connell 1985;

Rowley 1989), increasing the need for high frequency data of larval settlement spanning

multiple years. Data such as ours that meet these criteria are rare.

The potential factors affecting larval supply that we examined other than temperature

and climate showed no meaningful correlations with larval settlement. Of particular note was

the lack of a correlation with ocean chlorophyll revealed by our analyses. This finding differed

from that observed for the tropical sea urchin *Diadema aff. antillarum* (Hernández *et al.* 2010)

and highlights a need for exploring mechanistically how spatial and temporal variability in

ocean circulation and phytoplankton productivity affect patterns of larval settlement. Like

recent studies (Shanks & Shearman 2009; Shanks & Morgan 2018), we found no relationship

between large-scale fluctuations in upwelling and larval settlement, which runs counter to

hypotheses that large-scale coastal upwelling shapes patterns of larval supply by broadly

altering productivity and retention (Menge & Menge 2013; Roughgarden *et al.* 1991). The

major ocean currents in our study system (the Southern California Counter Current and the

California Current) are characterized by stochastic and spatially variable eddies, fronts,

filaments and bores (e.g. Bassin *et al.* 2005; Davis *et al.* 2008; DiGiacomo & Holt 2001), whereas

flows closer to shore tend to promote larval retention along wave exposed shores (Morgan *et*

al. 2016; Shanks *et al.* 2017). As a result, offshore transport and retention off the coast of

California is spatially nuanced and temporally inconsistent (Shanks & Eckert 2005). Such

heterogeneity not only influences larval transport, but also when and where larvae encounter

productive food environments. Consequently, if larval food limitation plays a role in shaping

the dynamics of urchin settlement at our sites, then it likely arises out of more complex

dynamics than our spatially aggregated composite metrics were able to detect. For example,

processes that shape stratification, front formation and spatially isolated phytoplankton blooms

in combination with larval behavior may play a more important role than predicted by

simplifications of the higher or lower spatially averaged phytoplankton production or greater or

lesser coastal upwelling. Integrating quantitative measurements of these processes was

beyond the scope of our study, yet such approaches are needed to develop a mechanistic

understanding of processes that control settlement dynamics across large gradients in

oceanographic settings.

Although climate related changes can alter larval supply via their effects on the

production of larvae by adults, we found no relationship between larval settlement and

regional adult abundance or adult food (kelp). While fisheries research relies heavily on stock-

recruit dynamics, there is continual debate about whether adult dynamics actually control

recruitment patterns (Gilbert 1997; Szuwalski *et al.* 2015). The lack of a positive correlation

between adults and larval settlement that we found for purple sea urchins adds to this debate

by showing that high abundances of adult urchins and kelp averaged across large spatial scales

did not translate into high larval supply. We note that spatial structure in the dynamics of kelp

and adult urchins may have affected this outcome. For example, modest but uniform densities

of adult sea urchins in food rich kelp forests can provide very different outcomes for the

production of embryos compared to mosaics of dense adults in barren patches interspersed

with forested patches having few adults (Okamoto 2014; Okamoto 2016). Indeed, previous

work has demonstrated that assessing population productivity in spatially structured

populations requires a spatially structured analysis to account for processes of population

regulation and competition (Chesson 1996, 1998; Chesson *et al.* 2005; Thorson *et al.* 2015).

That said, we found no evidence that synchronous fluctuations in settlement arose from broad-

scale collapses and proliferations in populations of adult sea urchins or kelp at the appropriate

lags and time-scales.

Temperature-related effects on larval production and survival and ENSO-related

changes in currents or oceanographic features that transport larvae represent two possible

explanations for the spatially opposing correlations between larval settlement and temperature

and ENSO that we observed. Many species have upper temperature thresholds beyond which

reproduction is impaired that are well below the lethal limits for adults. Temperatures in

southern California kelp forests routinely exceed 17°C during ENSO events (Reed et al. 2016),

which may adversely affect larval supply of purple sea urchins by limiting gamete production

(Basch & Tegner 2007; Cochran & Engelmann 1975), fertilization (Schroeder & Battaglia 1985)

larval development and gene expression (Padilla-Gamiño *et al.* 2013; Runcie *et al.* 2012; Wong

et al. 2018), and larval survival (Azad *et al.* 2012). Such effects may in part explain why larval

settlement at Fort Bragg, where sea surface temperatures rarely exceed 16°C (Figure 6a) was

not negatively correlated with temperature.

Regional circulation is considered to be an important driver of larval supply for many

species, including those in the Southern California Bight (Blanchette *et al.* 2006; Broitman *et al.*

2005; McManus & Woodson 2012; Woodson *et al.* 2012). ENSO events can cause major

changes in patterns of ocean circulation along the US west coast. For example, the 1997/98 El

Niño caused anomalously strong poleward flow and decreased local retention in coastal areas

off southern California (Lynn & Bograd 2002; Mitarai *et al.* 2009). Yet El Niño events also create

stochastic small-scale patterns of water motion in the form of fronts, eddies, and bores (Pineda

et al. 2018). Spatial variability in such patterns could explain the opposing climatic responses in

larval settlement between Fort Bragg and our sites in southern California. El Niño events in

southern California are associated lower stratification and reduced internal waves that promote

the onshore transport of larvae (Pineda 1994; Pineda *et al.* 2018; Shanks 1983). In contrast, El

Nino events in northern California are often associated with relaxed upwelling, downwelling

Kelvin waves and increased stratification (Chavez *et al.* 2002). Although we cannot definitively

partition out the effects of climate associated changes in temperature and circulation on

recruitment, our observations of larval settlement and subsequent year-class strength point

strongly to broader-scale ocean climate effects.

Our finding that larval settlement was a good predictor of year class strength in natural

populations indicates that large fluctuations in larval settlement of this prominent herbivore

could have far reaching ecological impacts that resonate throughout marine ecosystems across

a broad geographic region (Pearse 2006). For example, the high settlement that we observed in

Fort Bragg in 2013-2015 coupled with a period of anomalously warm water may explain the

marked increase in the abundance of purple urchins and coincident loss of canopy forming

kelps reported for northern California beginning in 2014 (Catton *et al.* 2016). Our results

suggest that predicted changes in ocean climate that lead to more frequent or severe marine

heatwaves (Frölicher and Laufkötter 2018) will reduce purple urchin recruitment in southern

California, but increase it in northern California with potentially cascading effects on benthic

ecosystems similar to those recently observed in northern California.

Oceans are experiencing simultaneous shifts in temperature, water chemistry,

productivity and circulation, which have profound consequences for the ecological structure

and function of marine systems and the services that they provide (Wong *et al.* 2014). Our

findings provide valuable insights into the ecological consequences of climate related effects on

patterns of larval settlement of an important reef herbivore whose distribution spans most of

the Pacific coast of North America. Future investigations aimed at determining the specific

biotic and abiotic processes that regulate larval settlement in ecological important species such

as *S. purpuratus* will undoubtedly improve our understanding of how climate fluctuations affect

regional population dynamics to alter the structure and functions of marine ecosystems.

Acknowledgements

Funding was provided by the California Urchin Commission, the South Bay Cable Committee,

the California Department of Fish and Wildlife Director's Sea Urchin Advisory Committee, and

the National Science Foundation's support of the Santa Barbara Coastal Long Term Ecological

Research program. We are indebted to Tom Ebert and John Dixon, who initiated the larval

settlement time series with SS. The Channel Islands Naturalist Corps volunteers, Channel

Islands National Park Staff, and Channel Islands long term Kelp Forest Monitoring Program were

and remain instrumental in collecting settlement data from Anacapa and benthic data in the

Channel Islands. John Richards, Clint Nelson, Shannon Harrer, Jenny Wolf, Peter Kalvass,

Brigitte Bondoux and many students assisted with field collections, laboratory sampling and

data compilation. Sally Holbrook, Russ Schmitt and Cherie Briggs contributed discussions

during initial development of analyses. Kyle Cavanaugh, Dave Siegel and Tom Bell provided

valuable discussions about giant kelp canopy data. Kyle Cavanaugh and Rachel Simons gave

important comments on previous manuscript drafts.

the coastline and 150 alongshore km of the location where settlement was sampled (hatched

areas in Figure 1). See Methods for details.

Figure 5: The relationship between the summer density of juvenile purple sea urchins on reefs

in the Santa Barbara Channel and the density of recruits on larval collectors during March- July

of the previous year (i.e., 12-20 months prior). Points and error bars represent annual means

(+/- SE) averaged over all sites. The line with the band represents the estimated relationship

and 95% uncertainty interval.

Figure 6: Standardized coefficients for the relationships between larval settlement of purple sea

urchins at each site and: (a-g) sea surface temperature, (h-n) the multivariate ENSO index, and

(m-u) the North Pacific Gyre Oscillation (NPGO). Small grey points represent biweekly means

and black points and lines represent annual means.

Figure 7: Standardized Bayesian multiple regression coefficients for relationships between *S.*

purpuratus and local covariates (a-e) or global climatic indices (f-h). Models with global and

local variables were fit separately. Coefficients were estimated using a multiple regression

nested within a Bayesian multivariate time series model. Coefficients are *a priori* biased

towards zero using a regularized horseshoe prior for variable selection. For details see

Methods.

References

References

Andrew N., Agatsuma Y., Ballesteros E., Bazhin A., Creaser E., Barnes D., Botsford L.,

Bradbury A., Campbell A. & Dixon J. (2003) Status and management of world sea urchin

fisheries. *Oceanography and Marine Biology-An Annual Review*, 40, 343-425

Azad A.K., Pearce C.M. & McKinley R.S. (2012) Influence of stocking density and temperature

on early development and survival of the purple sea urchin, *Strongylocentrotus*

purpuratus (Stimpson, 1857). *Aquac Res*, 43, 1577-1591

Bakun A. 1973. Coastal upwelling indices, west coast of North America, 1946-71. US

Department of Commerce, National Oceanic and Atmospheric Administration, National

Marine Fisheries Service pp. 103.

Basch L.V. & Tegner M.J. (2007) Reproductive responses of Purple sea urchin

(*Strongylocentrotus purpuratus*) populations to environmental conditions across a coastal

depth gradient. *Bull Mar Sci*, 81, 255-282

Bassin C.J., Washburn L., Brzezinski M. & McPhee-Shaw E. (2005) Sub-mesoscale coastal

eddies observed by high frequency radar: A new mechanism for delivering nutrients to

kelp forests in the Southern California Bight. *Geophysical Research Letters*, 32

Bell T., K. Cavanaugh and D. Siegel, (2017) SBC LTER: Time series of quarterly NetCDF files

of kelp biomass in the canopy from Landsat 5, 7 and 8, 1984 - 2016 (ongoing). Santa

Barbara Coastal Long Term Ecological Research Project.

doi:10.6073/pasta/817d2c24ebd78621869e17d94ba0df0c

Bell T.W., Cavanaugh K.C., Reed D.C. & Siegel D.A. (2015) Geographical variability in the

controls of giant kelp biomass dynamics. *J Biogeogr*, 42, 2010-2021

Bertram D.F. & Strathmann R.R. (1998) Effects of maternal and larval nutrition on growth and

form of planktotrophic larvae. *Ecology*, 79, 315-327

Blanchette C.A., Broitman B.R. & Gaines S.D. (2006) Intertidal community structure and

oceanographic patterns around Santa Cruz Island, CA, USA. *Mar Biol*, 149, 689-701

Broitman B.R., Blanchette C.A. & Gaines S.D. (2005) Recruitment of intertidal invertebrates

and oceanographic variability at Santa Cruz Island, California. *Limnol Oceanogr*, 50,

1473-1479

Burt J.M., Tinker M.T., Okamoto D.K., Demes K.W., Holmes K. & Salomon A.K. (2018)

Sudden collapse of a mesopredator reveals its complementary role in mediating rocky

reef regime shifts. *Proc R Soc B*, 285

Byrne M., Ho M., Selvakumaraswamy P., Nguyen H.D., Dworjanyn S.A. & Davis A.R. (2009)

Temperature, but not pH, compromises sea urchin fertilization and early development

under near-future climate change scenarios. *Proc R Soc B*, 276, 1883-1888

Cai W., Borlace S., Lengaigne M., Van Rensch P., Collins M., Vecchi G., Timmermann A.,

Santoso A., McPhaden M.J. & Wu L. (2014) Increasing frequency of extreme El Niño

events due to greenhouse warming. *Nature climate change*, 4, 111

Caley M., Carr M., Hixon M., Hughes T., Jones G. & Menge B. (1996) Recruitment and the

local dynamics of open marine populations. *Annu Rev Ecol Syst*, 27, 477-500

Carpenter B., Gelman A., Hoffman M., Lee D., Goodrich B., Betancourt M., Brubaker M.A.,

Guo J., Li P. & Riddell A. (2016) Stan: A probabilistic programming language. *J Stat

Softw*, 20

Carvalho C.M., Polson N.G. & Scott J.G. (2010) The horseshoe estimator for sparse signals.

Biometrika, 97, 465-480

Catton C., L. , Bennett R. & Amrhein. A. (2016) Perfect storm decimates northern California

kelp forests. California Department of Fish and Wildlife Marine Management News.

https://cdfwmarine.wordpress.com/2016/03/30/perfect-storm-decimates-kelp/

Chavez F., Pennington J., Castro C., Ryan J., Michisaki R., Schlining B., Walz P., Buck K.,

McFadyen A. & Collins C. (2002) Biological and chemical consequences of the 1997–

1998 El Niño in central California waters. *Progress in Oceanography*, 54, 205-232

Chesson P. 1996. Matters of scale in the dynamics of populations and communities. In: *Frontiers*

of population ecology (eds. RB Floyd, AW Sheppard & PJ de Barro). CSIRO, Victoria,

Australia. pp. 353-368.

Chesson P. (1998) Spatial scales in the study of reef fishes: a theoretical perspective. *Aust J Ecol*,

23, 209-215

Chesson P., Donahue M.J., Melbourne B.A. & Sears A.L. 2005. Scale transition theory for

understanding mechanisms in metacommunities. In: *Metacommunities: spatial dynamics*

and ecological communities (eds. M Holyoak, MA Leibold & RD Holt). University of

Chicago Presspp. 279-306.

Cochran R.C. & Engelmann F. (1975) Environmental regulation of the annual reproductive

season of *Strongylocentrotus purpuratus* (Stimpson). *Biol Bull*, 148, 393-401

Connell J.H. (1985) The consequences of variation in initial settlement vs. post-settlement

mortality in rocky intertidal communities. *J Exp Mar Biol Ecol*, 93, 11-45

Davis R.E., Ohman M.D., Rudnick D.L. & Sherman J.T. (2008) Glider surveillance of physics

and biology in the southern California Current System. *Limnol Oceanogr*, 53, 2151-2168

Di Lorenzo E., Schneider N., Cobb K.M., Franks P., Chhak K., Miller A.J., McWilliams J.C.,

Bograd S.J., Arango H. & Curchitser E. (2008) North Pacific Gyre Oscillation links

ocean climate and ecosystem change. *Geophysical Research Letters*, 35

Di Lorenzo E., Combes V., Keister J.E., Strub P.T., Thomas A.C., Franks P.J., Ohman M.D.,

Furtado J.C., Bracco A. & Bograd S.J. (2013) Synthesis of Pacific Ocean climate and

ecosystem dynamics. *Oceanography*, 26, 68-81

DiGiacomo P.M. & Holt B. (2001) Satellite observations of small coastal ocean eddies in the

Southern California Bight. *Journal of Geophysical Research: Oceans*, 106, 22521-22543

Ebert T. (1983) Recruitment in echinoderms. *Echinoderm Stud*, 1, 169-203

Ebert T., Schroeter S., Dixon J. & Kalvass P. (1994) Settlement patterns of red and purple sea

urchins (*Strongylocentrotus franciscanus* and *S. purpuratus*) in California, USA. *Mar*

Ecol Prog Ser, 111, 41-52

Ebert T.A. (2010) Demographic patterns of the purple sea urchin *Strongylocentrotus purpuratus*

along a latitudinal gradient, 1985-1987. *Mar Ecol Prog Ser*, 406, 105-120

Edmunds P.J. & Carpenter R.C. (2001) Recovery of Diadema antillarum reduces macroalgal

cover and increases abundance of juvenile corals on a Caribbean reef. *Proceedings of the*

National Academy of Sciences, 98, 5067-5071

Estes J.A. & Duggins D.O. (1995) Sea otters and kelp forests in Alaska: generality and variation

in a community ecological paradigm. *Ecol Monogr*, 65, 75-100

Feehan C.J. & Scheibling R.E. (2014) Effects of sea urchin disease on coastal marine

ecosystems. *Mar Biol*, 161, 1467-1485

Field C.M. & Walker C. 2003. *A Site Profile of the Kachemak Bay Research Reserve, a Unit of*

the National Estuarine Research Reserve System. Kachemak Bay Research Reserve

Homer

Filbee-Dexter K. & Scheibling R.E. (2014) Sea urchin barrens as alternative stable states of

collapsed kelp ecosystems. *Mar Ecol Prog Ser*, 495, 1-25

Fisher J.L., Peterson W.T. & Morgan S.G. (2014) Does larval advection explain latitudinal

differences in recruitment across upwelling regimes? *Mar Ecol Prog Ser*, 503, 123-137

Foster M.C., Byrnes J.E. & Reed D.C. (2015) Effects of five southern California macroalgal

diets on consumption, growth, and gonad weight, in the purple sea urchin

Strongylocentrotus purpuratus. *PeerJ*, 3, e719

Gilbert D. (1997) Towards a new recruitment paradigm for fish stocks. *Can J Fish Aquat Sci*, 54,

969-977

Gonor J.J. (1973) Reproductive cycles in oregon populations of the echlnoid, *StrongylocentrotUs*

purpuratus (Stimpson). I. Annual gonad growth and ovarian gametogenic cycles. *J Exp*

Mar Biol Ecol, 12, 45-64

Grosberg R.K. & Levitan D.R. (1992) For adults only? Supply-side ecology and the history of

larval biology. *Trends Ecol Evol*, 7, 130-133

Harrold C. & Reed D.C. (1985) Food availability, sea urchin grazing, and kelp forest community

structure. *Ecology*, 66, 1160-1169

Hernández J.C., Clemente S., Girard D., Pérez-Ruzafa Á. & Brito A. (2010) Effect of

temperature on settlement and postsettlement survival in a barrens-forming sea urchin.

Mar Ecol Prog Ser, 413, 69-80

653 Hoegh-Guldberg O. & Pearse J.S. (1995) Temperature, food availability, and the development of

654   marine invertebrate larvae. *Am Zool*, 35, 415-425

655 Kato S. & Schroeter S.C. (1985) Biology of the red sea urchin, *Strongylocentrotus franciscanus*,

656   and its fishery in California. *Mar. Fish. Rev*, 47, 1-20

657 Kushner D.J., Rassweiler A., McLaughlin J.P. & Lafferty K.D. (2013) A multi-decade time

658   series of kelp forest community structure at the California Channel Islands: Ecological

659   Archives E094-245. *Ecology*, 94, 2655-2655

660 Lafferty K.D. (2004) Fishing for lobsters indirectly increases epidemics in sea urchins. *Ecol*

661   *Appl*, 14, 1566-1573

662 Ling S., Johnson C., Ridgway K., Hobday A. & Haddon M. (2009) Climate-driven range

663   extension of a sea urchin: inferring future trends by analysis of recent population

664   dynamics. *Global Change Biology*, 15, 719-731

665 Lynn R. & Bograd S. (2002) Dynamic evolution of the 1997–1999 El Niño–La Niña cycle in the

666   southern California Current system. *Progress in Oceanography*, 54, 59-75

667 Magnusson A., Skaug H.J., Nielsen A., Berg C.W., Kristensen K., Maechler M., Bentham K.J.v.,

668   and B.M.B. & Brooks M.E. (2017) glmmTMB: Generalized Linear Mixed Models using

669   Template Model Builder. R package version 0.1.3. Available at

670   https://github.com/glmmTMB

671 Mantua N.J., Hare S.R., Zhang Y., Wallace J.M. & Francis R.C. (1997) A Pacific interdecadal

672   climate oscillation with impacts on salmon production. *Bulletin of the american*

673   *Meteorological Society*, 78, 1069-1080

674 Mantua N.J. & Hare S.R. (2002) The Pacific decadal oscillation. *Journal of Oceanography*, 58,

675   35-44

McManus M.A. & Woodson C.B. (2012) Plankton distribution and ocean dispersal. *J Exp Biol*,
215, 1008-1016

Menge B.A., Gouhier T.C., Freidenburg T. & Lubchenco J. (2011) Linking long-term, large-
scale climatic and environmental variability to patterns of marine invertebrate
recruitment: Toward explaining "unexplained" variation. *J Exp Mar Biol Ecol*, 400, 236-
249

Menge B.A. & Menge D.N. (2013) Dynamics of coastal meta-ecosystems: the intermittent
upwelling hypothesis and a test in rocky intertidal regions. *Ecol Monogr*, 83, 283-310

Mitarai S., Siegel D., Watson J., Dong C. & McWilliams J. (2009) Quantifying connectivity in
the coastal ocean with application to the Southern California Bight. *J Geophys Res*, 114

Morgan S.G., Fisher J.L., Miller S.H., McAfee S.T. & Largier J.L. (2009) Nearshore larval
retention in a region of strong upwelling and recruitment limitation. *Ecology*, 90, 3489-
3502

Morgan S.G. (2014) Behaviorally mediated larval transport in upwelling systems. *Advances in*
Oceanography, 2014

Morgan S.G., Shanks A.L., Fujimura A.G., Reniers A.J., MacMahan J., Griesemer C.D., Jarvis
M. & Brown J. (2016) Surfzone hydrodynamics as a key determinant of spatial variation
in rocky intertidal communities. *Proc R Soc B*, 283, 20161017

Morgan S.G.M., Miller S.H., Robart M.J. & Largier J.L. (2018) Nearshore larval retention and
cross-shelf migration of benthic crustaceans at an upwelling center. *Frontiers in Marine*
Science, 5, 161

Myers R.A. (1998) When do environment–recruitment correlations work? *Rev Fish Biol Fish*, 8,
285-305

Okamoto D.K. 2014. *The role of fluctuating food supply on recruitment, survival and population*

dynamics in the sea. University of California, Santa Barbara

Okamoto D.K. (2016) Competition among eggs shifts to cooperation along a sperm supply

gradient in an external fertilizer. *Am Nat*, 187, E129-E142

Olivares-Bañuelos N.C., Enríquez-Paredes L.M., Ladah L.B. & De La Rosa-Vélez J. (2008)

Population structure of purple sea urchin Strongylocentrotus purpuratus along the Baja

California peninsula. *Fisheries science*, 74, 804-812

Padilla-Gamiño J.L., Kelly M.W., Evans T.G. & Hofmann G.E. (2013) Temperature and CO2

additively regulate physiology, morphology and genomic responses of larval sea urchins,

Strongylocentrotus purpuratus. *Proc R Soc B*, 280

Pearse J.S., Pearse V.B. & Davis K.K. (1986) Photoperiodic Regulation of Gametogenesis and

Growth in the Sea Urchin *Strongylocentrotus purpuratus*. *J Exp Zool*, 237, 107-118

Pearse J.S. (2006) Ecological role of purple sea urchins. *Science*, 314, 940-941

Piironen J. & Vehtari A. (2017) Sparsity information and regularization in the horseshoe and

other shrinkage priors. *Electronic Journal of Statistics*, 11, 5018-5051

Pineda J. (1994) Internal tidal bores in the nearshore: Warm-water fronts, seaward gravity

currents and the onshore transport of neustonic larvae. *Journal of Marine Research*, 52,

427-458

Pineda J., Reyns N. & Lentz S.J. (2018) Reduced barnacle larval abundance and settlement in

response to large-scale oceanic disturbances: Temporal patterns, nearshore thermal

stratification, and potential mechanisms. *Limnol Oceanogr*, 63, 2618-2629

Reynolds R.W., Smith T.M., Liu C., Chelton D.B., Casey K.S. & Schlax M.G. (2007) Daily

high-resolution-blended analyses for sea surface temperature. *J Clim*, 20, 5473-5496

Roughgarden J., Pennington J., Stoner D., Alexander S. & Miller K. (1991) Collisions of

upwelling fronts with the intertidal zone: the cause of recruitment pulses in barnacle

populations of central California. *Acta Oecol*, 12, 35-51

Rowley R. (1989) Settlement and recruitment of sea urchins (*Strongylocentrotus spp.*) in a sea-

urchin barren ground and a kelp bed: are populations regulated by settlement or post-

settlement processes? *Mar Biol*, 100, 485-494

Runcie D.E., Garfield D.A., Babbitt C.C., Wygoda J.A., Mukherjee S. & Wray G.A. (2012)

Genetics of gene expression responses to temperature stress in a sea urchin gene network.

Mol Ecol, 21, 4547-4562

Sathyendranath S., Grant M., Brewin R.J.W., Brockmann C., Brotas V., Chuprin A., Doerffer R.,

Dowell M., Farman A., Groom S., Jackson T., Krasemann H., Lavender S., Martinez

Vicente V., Mazeran C., Mélin F., Moore T.S., Müller D., Platt T., Regner P., Roy S.,

Steinmetz F., Swinton J., Valente A., Zühlke M., Antoine D., Arnone R., Balch W.M.,

Barker K., Barlow R., Bélanger S., Berthon J.-F., Beşiktepe Ş., Brando V.E., Canuti E.,

Chavez F., Claustre H., Crout R., Feldman G., Franz B., Frouin R., García-Soto C., Gibb

S.W., Gould R., Hooker S., Kahru M., Klein H., Kratzer S., Loisel H., McKee D.,

Mitchell B.G., Moisan T., Muller-Karger F., O'Dowd L., Ondrusek M., Poulton A.J.,

Repecaud M., Smyth T., Sosik H.M., Taberner M., Twardowski M., Voss K., Werdell J.,

Wernand M. & Zibordi G. (2018) ESA Ocean Colour Climate Change Initiative

(Ocean_Colour_cci): Version 3.1 Data. Centre for Environmental Data Analysis.

http://dx.doi.org/10.5285/9c334fbe6d424a708cf3c4cf0c6a53f5

Schroeder T. & Battaglia D. (1985) "Spiral asters" and cytoplasmic rotation in sea urchin eggs:

induction in *Strongylocentrotus purpuratus* eggs by elevated temperature. *J Cell Biol*,

100, 1056-1062

Shanks A. & Shearman R. (2009) Paradigm lost? Cross-shelf distributions of intertidal

invertebrate larvae are unaffected by upwelling or downwelling. *Mar Ecol Prog Ser*, 385,

189-204

Shanks A.L. (1983) Surface slicks associated with tidally forced internal waves may transport

pelagic larvae of benthic invertebrates and fishes shoreward. *Marine ecology progress*

series. Oldendorf, 13, 311-315

Shanks A.L. & Eckert G.L. (2005) Population persistence of California Current fishes and

benthic crustaceans: a marine drift paradox. *Ecol Monogr*, 75, 505-524

Shanks A.L. (2009) Pelagic larval duration and dispersal distance revisited. *The Biological*

Bulletin, 216, 373-385

Shanks A.L., Morgan S.G., MacMahan J. & Reniers A.J. (2017) Alongshore variation in

barnacle populations is determined by surf zone hydrodynamics. *Ecol Monogr*, 87, 508-

532

Shanks A.L. & Morgan S.G. (2018) Testing the intermittent upwelling hypothesis: upwelling,

downwelling, and subsidies to the intertidal zone. *Ecol Monogr*, 88, 22-35

Shelton A.O. & Mangel M. (2011) Fluctuations of fish populations and the magnifying effects of

fishing. *Proceedings of the National Academy of Sciences*, 108, 7075-7080

Shkedy Y. & Roughgarden J. (1997) Barnacle recruitment and population dynamics predicted

from coastal upwelling. *Oikos*, 487-498

765 Siegel D., Kinlan B., Gaylord B. & Gaines S. (2003) Lagrangian descriptions of marine larval

766   dispersion. *Mar Ecol Prog Ser*, 260, 83-96

767 Siegel D., Mitarai S., Costello C., Gaines S., Kendall B., Warner R. & Winters K. (2008) The

768   stochastic nature of larval connectivity among nearshore marine populations.

769   *Proceedings of the National Academy of Sciences*, 105, 8974-8979

770 Strathmann M.F. 1987. *Reproduction and development of marine invertebrates of the northern*

771   *Pacific coast: data and methods for the study of eggs, embryos, and larvae*. Univ. of

772   Wash. Press, Seattle

773 Strathmann R. (1978) Length of pelagic period in echinoderms with feeding larvae from the

774   northeast Pacific. *J Exp Mar Biol Ecol*, 34, 23-27

775 Sydeman W.J., Poloczanska E., Reed T.E. & Thompson S.A. (2015) Climate change and marine

776   vertebrates. *Science*, 350, 772-777

777 Szuwalski C.S., Vert-Pre K.A., Punt A.E., Branch T.A. & Hilborn R. (2015) Examining common

778   assumptions about recruitment: a meta-analysis of recruitment dynamics for worldwide

779   marine fisheries. *Fish and Fisheries*, 16, 633-648

780 Tegner M. & Dayton P. (1991) Sea urchins, El Niños, and the long term stability of Southern

781   California kelp forest communities. *Mar Ecol Prog Ser*, 77, 49-63

782 Thorson J.T., Skaug H.J., Kristensen K., Shelton A.O., Ward E.J., Harms J.H. & Benante J.A.

783   (2015) The importance of spatial models for estimating the strength of density

784   dependence. *Ecology*, 96, 1202-1212

785 Underwood A. & Fairweather P. (1989) Supply-side ecology and benthic marine assemblages.

786   *Trends Ecol Evol*, 4, 16-20

Figure 1: Map of larval settlement collection sites in California. Colors represent the average spatial gradient of sea surface temperature from 1991-2017. The hatched white lines show the shoreline buffer used to constrain sea surface temperature and sea surface chlorophyll index to local larval settlement observations.

Figure 2: Diagram of the model structure using either a Matérn temporal covariance (trend estimation model) or a horseshoe regression (correlation estimation model) to estimate seasonal anomalies. For details see Tables 1 and 2. *Bayesian Regression Model Only. † Trend Estimation Model Only

Figure 3: Standardized annual scale trends (log-scale, standardized to one SD) of larval settlement S. purpuratus from 1990 through 2016 for: (a) Fort Bragg, (b) sites in the Santa Barbara Channel, and (c) sites in San Diego. Colors within represent individual sites associated with the legend. Interannual trends are estimated using a 3/2 Matrn Gaussian process covariance within a Bayesian multivariate state-space model accounting for spatiotemporal correlations.

Figure 4: Seasonal trends at each site for: (a) standardized larval settlement of S. purpuratus, (b) biweekly mean sea surface chlorophyll a (chl a) and (c) biweekly mean sea surface temperature (SST). The seasonal trend in larval settlement at each site was standardized by the mean maximum seasonal value. Chl a and SST were measured using satellite imagery within 25 km of the coastline and 150 alongshore km of the location where settlement was sampled (hatched areas in Figure 1). See Methods for details.

Figure 5: The relationship between the summer density of juvenile purple sea urchins on reefs in the Santa Barbara Channel and the density of recruits on larval collectors during March- July of the previous year (i.e., 12-20 months prior). Points and error bars represent annual means (+/- SE) averaged over all sites. The line with the band represents the estimated relationship and 95% uncertainty interval.

Figure 6: Relationships between larval settlement of purple sea urchins at each site and: (a-g) sea surface temperature, (h-n) the multivariate ENSO index, and (m-u) the North Pacific Gyre Oscillation (NPGO). Small grey points represent biweekly means and black points and lines represent annual means.

Figure 7: Standardized Bayesian multiple regression coefficients for relationships between *S. purpuratus* settlement and local covariates (a-e) or global climatic indices (f-h). Models with global and local variables were fit separately. Coefficients were estimated using a multiple regression nested within a Bayesian multivariate time series model. Coefficients are a priori biased towards zero using a regularized horseshoe prior for variable selection. For details see Methods, Figure 2 and appendix for details.

Table 1: Equations for the multivariate spatiotemporal models. Variables, parameters and priors are described in Table 2. Note that there are two primary model forms: a trend estimation model, and a Bayesian regression model. Equations apply to both model forms except where specified.

Eq.	Description	Equation
3	total count likelihood	$\begin{cases} Tot_{t,s,b} \sim \text{Poisson}(\hat{\mu}_{t,s} \times D_{t,s,b}/p_{t,s,b}) & \text{when subsampled} \\ N_{t,s,b} \sim \text{Poisson}(\hat{\mu}_{t,s} \times D_{t,s,b}) & \text{otherwise} \end{cases}$
4	subsampling prior	$p_{t,s,b} \sim \text{beta}(N_{t,s,b} + 0.5, NO_{t,s,b} + 0.5)$ when subsampled
5	prior seasonal temporal correlation function	$\bar{r}_{<t_i,t_j>} = \exp\left[-\frac{2}{1.5^2}(0.5 - 0.5\cos(\pi|t_i - t_j|))\right]$ Periodic Kernel (MacKay 1998)
6[†]	prior annual temporal correlation function	$\bar{\bar{r}}_{<t_i,t_j>} = (1 + 3^2|t_i - t_j|)e^{-3^2|t_i - t_j|}$ Matérn 3/2 Kernel (Rasmussen & Williams 2006)
7	prior seasonal spatiotemporal covariance function (separable)	$\bar{\boldsymbol{\Sigma}} = \bar{\boldsymbol{\Omega}} \otimes [\bar{\boldsymbol{\sigma}}\bar{\boldsymbol{R}}]$
8[†]	prior annual spatiotemporal covariance function (separable)	$\bar{\bar{\boldsymbol{\Sigma}}} = \bar{\bar{\boldsymbol{\Omega}}} \otimes [\bar{\bar{\boldsymbol{\sigma}}}\bar{\bar{\boldsymbol{R}}}]$
9	matrix of seasonal trends	$\textbf{Seasonal} \sim \text{MVN}(0, \bar{\boldsymbol{\Sigma}})$
10[†]	matrix of annual trends	$\textbf{Annual} \sim \text{MVN}(0, \bar{\bar{\boldsymbol{\Sigma}}})$
11a[*]	first order autoregressive model	$\boldsymbol{\epsilon}_t = \phi\boldsymbol{\epsilon}_{t-1} + \boldsymbol{e}_t\sqrt{(1 - \phi^2)}$
11b	spatially correlated process error	$\boldsymbol{e}_t \sim \text{MVN}(0, \boldsymbol{\sigma\Omega})$

[*] Bayesian Regression Model Only
[†] Trend Estimation Model Only

Table 2: Definition of parameters, variables, and priors for equations described in Table 1

Symbol	Description	Type	Prior/Input
$\hat{\mu}_{t,s}$	expected settlement density (per brush, per day) of purple urchins, defined in Eqs. 1 or 2 in the main text for each model.	estimated	Eqs. 1, 2
$N_{t,s,b}$	number of purple urchins counted at time t, site s and brush b. When urchin species within brushes are subsampled, data produced are a total urchin count ($Tot_{t,s,b}$), a total subsample count ($NO_{t,s,b}$, and a purple urchin subsample count ($N_{t,s,b}$).	input	data
$Tot_{t,s,b}$	total number of urchins counted at time t, site s and brush b	input	data
$p_{t,s,b}$	estimated proportion of purple urchins in the counts given the subsample (when urchin counts in brushes are subsampled).	input	data
$D_{t,s,b}$	exact number of days of the brush deployment	input	data
$\mathbf{X}$	covariate design matrix - for estimation model represents only site indices, for regression model includes covariates	input	data
$\boldsymbol{\beta}$	vector of site specific scale parameters and covariate coefficients	estimated	**Scale parameters**: Student-t: scale=2; **Coefficients**: regularized horseshoe prior (see methods).
$\boldsymbol{\phi}$	vector of site specific first order autoregressive parameters & estimated (regression model only)	estimated	Uniform -0.95 to 0.95
$\boldsymbol{\Omega}, \bar{\boldsymbol{\Omega}}, \bar{\bar{\boldsymbol{\Omega}}}$	spatial correlation matrices for process error, annual trends, and seasonal trends (estimated through the cholesky factor)	estimated	LKJ prior:scale= 2 (Lewandowski *et al.* 2009)
$\bar{\boldsymbol{R}}, \bar{\bar{\boldsymbol{R}}}$	temporal correlation matrices for annual and seasonal gaussian processes	input	eqs 5 & 6
$\boldsymbol{\sigma}, \bar{\boldsymbol{\sigma}}, \bar{\bar{\boldsymbol{\sigma}}}$	site specific standard deviation for process error, annual trends, and seasonal trends	estimated	half-Cauchy: scale = 2.5 (Polson & Scott 2012)

Supplementary Information - Appendices for
"Geographically Opposing Responses of Sea Urchin Recruitment to Changes in Ocean Climate"

Daniel K. Okamoto[*1,2], Stephen Schroeter[†3], and Daniel C. Reed[‡3]

[1]Department of Biological Science, Florida State University, Tallahassee, Florida 32303
[2]Ecology, Evolution and Marine Biology, University of California Santa Barbara, Santa Barbara, CA 93106 USA
[3]Marine Science Institute, University of California Santa Barbara, Santa Barbara, CA 93106 USA

[*]corresponding author: dokamoto@bio.fsu.edu

[†]email: schroete@lifesci.ucsb.edu

[‡]email: dan.reed@lifesci.ucsb.edu

[*]corresponding author: dokamoto@bio.fsu.edu

[†]email: schroete@lifesci.ucsb.edu

[‡]email: dan.reed@lifesci.ucsb.edu

Estimation of juvenile urchin densities

The KFM program surveys purple sea urchins at the Santa Barbara Channel Islands by counting the total number of individuals in defined areas and measuring approximately 100 individuals at each site. Thus, to model density we used the number of juveniles measured divided by the total number of individuals measured, multiplied by the total number of individuals counted, divided by the area surveyed:

$$Juvenile \ \ Density_{i,t} = \frac{Juveniles \ \ Measured_{i,t}}{Total \ \ Measured_{i,t}} \frac{Total \ \ Count_{i,t}}{Total \ \ Area_{i,t}}$$

We modeled density in the GLM using the number of measured juvenile urchins (2.5 cm in test diameter- the approximate cutoff size for reproduction - Kenner & Lares 1991). Let represent expected juvenile density at site i in year t. We constructed the regression as:

$$\ln\mu_{i,t} = \alpha_i + \beta \ \ \ln Larval \ \ recruitment_t$$

We allowed the intercept (α_i) to vary by site nested within each island because of overall differences in mean juvenile density among sites and islands. We used a negative binomial (NB) with the using direct mean and variance parameterization form:

$$Juveniles \ \ Measured_{i,t} \sim \text{NB}(\ln\mu_{i,t} - \ln[\frac{1}{TotalMeasured_{i,t}} \frac{TotalCounted_{i,t}}{TotalArea_{i,t}}], \sigma^2)$$

We constructed the model in this format (i.e. with the density denominator in the left hand side of the equation) to maintain the sample size and integer nature of the data while modeling the mean density.

Figure S.1: Observed and estimated purple urchin (*S. purpuratus*) settlement trends from 1990-2016 for each site. Large points represent empirical biweekly means (non-zero means across all brushes and collection times), lines on the axis represent biweekly mean values equal to zero, small points represent counts on individual brushes (non-zero only), red lines represent biweekly estimates, lines with blue 95% uncertainty intervals represent model estimates of the interannual trends. The model estimates the seasonal trend and the interannual trend at each site simultaneously. For reference, interannual trends for *S. purpuratus* shown here are the unstandardized versions of those shown in Figure 2.

Figure S.2: Seasonal settlement trends from 1990-2016 with the mean estimated seasonal trend for each site. Large points represent empirical biweekly means (non-zero means across all brushes and collection times), lines on the axis represent biweekly mean values equal to zero, small points represent counts on individual brushes (non-zero only), black lines represent the posterior mean and bands represent the 95% uncertainty interval. The model estimates the seasonal trend and the interannual trend at each site simultaneously. For reference, seasonal trends for *S. purpuratus* shown here are the unstandardized versions of those shown in Figure 3.

Variable Correlations

To illustrate the intercorrelated nature of the covariates and responses, we estimated a network model for Gaviota Pier that includes the independent model estimates of *S. purpuratus* settlement and its seasonality. To do so, we first constrained network structure by excluding all nonsensical interactions (i.e. chlorophyll does not cause ENSO events and is thus eliminated a priori) and including all known directional interactions (i.e. estimated seasonality affecting settlement for is forced into the network). Learning of the network skeleton is achieved via the Hill Climbing algorithm using the bnlearn (Scutari 2010) package in R. Note that strong collinearity can result in the weaker correlation being ignored (e.g. ENSO over SST with Larval Recruitment).

Figure S.3: Bayesian network model structure inferred for the Santa Barbara Channel. Numbers represent the partial correlation coefficients accounting for all other variables pointing to that node. All correlations shown are significant at alpha= 0.05

References

Kenner M. & Lares M. (1991) Size at first reproduction of the sea urchin *Strongylocentrotus purpuratus* in a central California kelp forest. Mar Ecol Prog Ser, 76, 303-306

Lewandowski D., Kurowicka D. & Joe H. (2009) Generating random correlation matrices based on vines and extended onion method. J Multivar Anal, 100, 1989-2001

MacKay D.J. (1998) Introduction to Gaussian processes. NATO ASI Series F Computer and Systems Sciences, 168, 133-166

Polson N.G. & Scott J.G. (2012) On the half-Cauchy prior for a global scale parameter. Bayesian Analysis, 7, 887-902

Rasmussen C.E. & Williams C.K.I. 2006. Gaussian Processes for Machine Learning. The MIT Press, Cambridge, Massachusetts

Scutari M. (2010). Learning Bayesian Networks with the bnlearn R Package. Journal of Statistical Software, 35(3), 1-22. URL http://www.jstatsoft.org/v35/i03/

```
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
\\ Stan Code Required to Estimate Trends      \\
\\ Author: D.K. Okamoto                        \\
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\

data {
  // input information
  int NO; // number of observations
  int NS; // number of sites
  int NM; // number of months
  int NYM; // number of months
  int NSUB; // number of observations with subsamples
  // input data
  //  Beta priors for uncertainty in subsamples
  vector[NSUB] P1; // SP observations (on the original scale)+0.5
  vector[NSUB] P2; // SP observations - subsample count (on the original scale) +
      0.5
  int SUB[NSUB];

  int N[NO]; // total count (on the original scale)
  int OBS[NO]; // continuous month-site observation index
  int PRED_MONTH[NM]; // month-site observation index associated with MA
  vector[NO] D; // number of days
  cholesky_factor_cov[NYM] D_seas;   // seasonal cholesky covariance factor
  cholesky_factor_cov[NM] D_ann;   // annual cholesky covariance factor
}

parameters {
  // mean volatility
  vector<lower=1e-7>[NS] sigma_S_mu;

  // spatial volatility correlation matrix
  cholesky_factor_corr[NS] L_Omega_S; //
  cholesky_factor_corr[NS] L_Omega_Spatial;

  // seasonal and annual variance
  vector<lower = 0>[NS] sigma;
  vector<lower = 0>[NS] sigma2;

  // mean site settlement
  row_vector<lower= -10, upper= 10>[NS] mu_S;

  // iid errors for seasonality
  matrix<lower= -10, upper= 10>[NYM,NS] z_s;

  // iid errors for annual trends
  matrix<lower= -10, upper= 10>[NS,NM] w_z;

  // iid errors for residuals
  matrix<lower= -4, upper= 4>[NS,NM] e_z;

  // proportion of samples that are purps
  vector<lower= 0,upper= 1>[NSUB] theta;
}

transformed parameters {
  matrix[NYM,NS] LS; // log scale expected sp per brush, per day for every month

  matrix[NM,NS] S; // estimated abundance
```

```
    vector[NO] S_exp; // normal scale estimated sp for each observation

    vector[NO] theta2; // fraction that is SP

    matrix[NM,NS] w;

    // correlated errors
    matrix[NM,NS] e;

    LS  = D_seas*z_s;
    w = D_ann*transpose(L_Omega_Spatial*w_z);

    theta2 = rep_vector(1.0,NO);
    theta2[SUB] = theta;
    // put it all together
    e = transpose(diag_pre_multiply(sigma_S_mu,L_Omega_S)*e_z);
    S= rep_matrix(mu_S,NM)+diag_post_multiply(LS[PRED_MONTH,],sigma2)+
        diag_post_multiply(w,sigma)+e;
    S_exp = exp(to_vector(S)[OBS]) .*D ./theta2;
}

model {
  sigma ~ cauchy(0, 0.5);
  sigma2 ~ cauchy(0, 0.5);

  L_Omega_S ~ lkj_corr_cholesky(2);
  L_Omega_Spatial ~ lkj_corr_cholesky(2);

  for(i in 1:NS){
    z_s[i]~ normal(0,1);
    mu_S[i] ~ normal(0,5);
  }

  sigma_S_mu ~ cauchy(0,2.5);

  // uncertainty on proportions of purples in the sample
  // jeffrey's prior given subsampling
  theta~beta(P1, P2);

  // observation likelihood
  N~poisson(S_exp);
}
```

```
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
\\ Stan Code Required to Estimate Bayesian Regression \\
\\ Author: D.K. Okamoto                               \\
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
data {
  // input information
  int NO; // number of observations
  int NS; // number of sites
  int NM; // number of months
  int NYM; // number of months
  int NSUB; // number of observations with subsamples
  int NSP; // number of subsample counts
  int NP; // number of variables
  // input data
  //  Beta priors for uncertainty in subsamples
  vector[NSP] P1; // SP observations (on the original scale)+0.5
  vector[NSP] P2; // NON SP observations (on the original scale) + 0.5
  int SUB[NSUB];
  int ID[NO];
  int OBS[NO]; // continuous month-site observation index
  // input data
  int N[NO]; // total count (on the original scale)
  int OBS_MONTH[NO]; // continuous month-site observation index associated with MA
  int OBS_SITE[NO]; // month-site observation index associated with MA
  int PRED_MONTH[NM]; // month-site observation index associated with MA
  matrix[NM*NS,NP] X; // covariate matrix
  int xind[NM,NS]; // covariate matrix indices
  vector[NO] D; // number of days
  cholesky_factor_cov[NYM] D_seas;   // seasonal cholesky covariance factor
  real<lower= 0> m0;
  real<lower= 0> sigma_scale;
  real<lower= 0> slab_scale;
}

parameters {
  // mean volatility
  vector<lower=0>[NS] sigma_S_mu;

  // spatial volatility correlation matrix
  cholesky_factor_corr[NS] L_Omega_S; //
  cholesky_factor_corr[NS] L_Omega_Spatial_seas;

  // iid errors for seasonality
  matrix[NYM,NS] z_s;

  // seasonal and annual variance
  real<lower = 0> sigma[NS];

   // mean site settlement
  row_vector[NS] mu_S;

  // iid errors for residuals
  matrix<lower= -4,upper=4>[NS,NM] e_z;

  // AR1 parameters
  vector<lower= -0.9,upper= 0.9>[NS] phi;

  // proportion of samples that are purps
  vector<lower= 0,upper= 1>[NSP] theta;
```

```
  // iid errors for seasonal gaussian process
  vector[NP] beta_tilde;
  vector<lower=0>[NP] lambda;
  real<lower=0> c2_tilde;
  real<lower=0> tau_tilde;
  real alpha;
}

transformed parameters {
  matrix[NYM,NS] LS; // log scale expected sp per brush, per day for every month
  matrix[NM,NS] S; // estimated abundance
  vector[NO] S_exp; // normal scale estimated sp for each observation

  vector[NO] N_mu; // expected total count
  vector[NO] q_mu; // expected fraction of N that is SP
  vector[NS] sigma_star;

  vector[NM*NS] s_hat;

  // process variables
  vector[NO] theta2; // fraction that is SP

  // correlated errors
  matrix[NM,NS] e;
  matrix[NM,NS] e_hat;

  vector[NP] beta;
  vector[NP] lambda_tilde;

  real tau0 = (m0/(NP-m0))*(mean(sigma_S_mu)*sqrt(1.0*NM*NS));
  real slab_scale2 = square(slab_scale);
  real half_slab_df = 0.5 * 25;

  real tau = tau0 * tau_tilde;
  real c2 = slab_scale2 * c2_tilde;
  lambda_tilde  = sqrt( c2 * square(lambda) ./ (c2 + square(tau) * square(lambda))
      );
  beta = tau * lambda_tilde .* beta_tilde;

  // regression model
  s_hat = X*beta;

  // seasonal gaussian process model
  LS  = D_seas*z_s;

  // process error
  for(i in 1:NS){
    sigma_star[i] = sigma_S_mu[i]*sqrt(1.0-phi[i]^2.0);
  }
  e = transpose(diag_pre_multiply(sigma_star,L_Omega_S)*e_z);
  // put it all together

  for(j in 1:NS){
    e_hat[1,j] = e[1,j];
    S[1,j]=mu_S[j]+LS[PRED_MONTH[1],j]*sigma[j]+s_hat[xind[1,j]]+e_hat[1,j]-5;
    for(i in 2:NM){
      e_hat[i,j] =  phi[j]*e_hat[i-1,j]+e[i,j];
      S[i,j]=mu_S[j]+LS[PRED_MONTH[i],j]*sigma[j]+s_hat[xind[i,j]]+e_hat[i,j]-5;
```

```
    }
  }

  theta2 = rep_vector(1.0,NO);
  theta2[SUB] = theta[ID[SUB]];

  S_exp = exp(to_vector(S)[OBS]) .*D ./theta2;
}

model {
  sigma ~ normal(0.0, 2);

  L_Omega_S ~ lkj_corr_cholesky(2.0);

  for(i in 1:NS){
    z_s[i]~ normal(0.0,1.0);
    e_z[i]~ normal(0.0,1.0);
    mu_S[i] ~ cauchy(0.0,10);
  }

  sigma_S_mu ~ normal(0.0,sigma_scale);

  beta_tilde ~ normal(0, 1);
  lambda ~ cauchy(0, 1);
  tau_tilde ~ cauchy(0, 1);
  c2_tilde ~ inv_gamma(half_slab_df, half_slab_df);

  // jeffrey's prior given subsampling
  theta~beta(P1, P2);

  // observation likelihood
  N~poisson(S_exp);
}
```