Genome-wide association study, replication, and mega-analysis using a dense marker panel in a multi-generational mouse advanced intercross line

Xinzhu Zhou[1], Celine L. St. Pierre[2], Natalia M. Gonzales[3], Riyan Cheng[4], Apurva Chitre[4], Greta Sokoloff[5], Abraham A. Palmer[4,6*]

[1] Biomedical Sciences Graduate Program, University of California San Diego, La Jolla, CA, USA;

[2] Department of Genetics, Washington University School of Medicine, St. Louis, MO, USA

[3] Department of Human Genetics, University of Chicago, Chicago, IL, USA

[4] Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

[5] Department of Psychological & Brain Sciences, University of Iowa, Iowa City, IO, USA

[6] Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA

*Corresponding author

Email: aapalmer@ucsd.edu (AAP)

Author contributions

XZ imputed genotypes, performed SNP and individual QC, and conducted GWAS in F_{34} and $F_{39\text{-}43}$ AILs under supervision of AAP. AAP also provided computational resources for the analyses in this paper. CS prepared GBS libraries for sequencing, as well as organizing portions of the $F_{39\text{-}43}$ phenotypes. NMG de-multiplexed GBS sequencing results and performed alignment and variant calling. RC helped with kinship relatedness matrix calculated from AIL pedigree. AC provided technical support for running programs and scripts. GS collected $F_{39\text{-}43}$ phenotypes, respectively. XZ co-wrote the manuscript with AAP, who designed the study and oversaw data collection.

Data Availability

All relevant data are within the paper and its Supporting Information files. The GeneNetwork accession numbers for genotypes and phenotypes of F_{34}, $F_{39\text{-}43}$, and mega-analysis cohort of AIL are currently pending but will be secured prior to final acceptance/publication.

Abstract

Genome-wide association studies (**GWAS**) in multigenerational outbred populations offer improved mapping precision compared to traditional populations such as F_2s. Advanced intercross lines (**AILs**) are the simplest possible multigenerational intercross; AILs are produced by crossing two inbred strains and then breeding unrelated offspring for additional generations. New recombinations accumulate with each successive generation; these new recombinations provide increased mapping resolution. We used genotyping-by-sequencing (**GBS**) to re-genotype a cohort of sparsely genotyped F_{34} LG/J x SM/J AIL mice that were the subject of several prior publications as well as to obtain genotypes for new cohort of AIL mice from the $F_{39\text{-}43}$ generations. The denser set of GBS markers allowed us to identify 110 significant loci, 36 of which were novel, for 79 behavioral and physiological traits in the 428 F_{34} mice. Genetic correlations between F_{34} and $F_{39\text{-}43}$ were high, though $F_{39\text{-}43}$ AILs showed systematically lower SNP-heritability estimates. We explored replication of loci identified in either F_{34} or $F_{39\text{-}43}$ in the other cohort for the traits measured in both F_{34} and $F_{39\text{-}43}$: locomotor activity, body weight, and coat color. While coat color loci were robustly replicated, we observed only partial replication of associations for locomotor activity and body weight. We then performed a mega-analysis of locomotor activity, body weight by combining F_{34} and $F_{39\text{-}43}$ mice (N=1028), which identified four novel loci. Finally, we showed that imputation using the old, sparse marker set and the newer dense marker set had little impact on our results, emphasizing the need of the denser GBS genotypes. The present study provides empirical insights into replication, the utility of denser genotyping and identifies new candidate loci that can be explored in future studies.

Author summary

Using a sufficiently dense marker set is essential for genome-wide association studies. The required SNP density is a function of population history. When considering populations derived from a cross between two or more inbred strains, more recombinations will require a correspondingly greater number of markers. We performed several genome-wide association studies using different marker sets - sparse SNPs from a previously publish dataset, dense SNPs obtained using genotyping-by-sequencing, and imputed versions of both the sparse and dense SNPs - to compare their association results in 34th and 39-43rd generations of an LG/J x SM/J advanced intercross line (**AIL**). We found that the dense SNP set substantially increased the number of significant results, while the imputed sets did not. To achieve maximum power, we performed a mega-analysis using all the mice and the dense SNP set; this analysis identified several novel loci. Finally, we explored the replicability of associated loci discovered in either cohort by examining the same SNP in the other cohort. We have made these data publicly available on www.genenetwork.org, providing the community with a unique reference dataset.

Introduction

The use of multi-parental crosses and commercially available outbred populations for genome wide association studies (**GWAS**) in model organisms such as mice [1–17], rats [18], chickens [19,20], zebrafish [21,22], fruit flies [23–27], *C. elegans* [28] and various plant species [29–31] has become increasingly popular over the last decade. These mapping populations can further be categorized as multi-parental crosses, which are created by interbreeding two or more inbred strains, and commercially available outbred populations, in which the founders are of unknown provenance. An F_2 cross between two inbred strains is the prototypical mapping population; however, F_2s provide poor mapping resolution [32]. To improve mapping resolution, Darvasi and Soller [33] proposed the creation of advanced intercross lines (**AILs**), which are produced by intercrossing F_2 mice for additional generations. AILs accumulate additional crossovers with every successive generation, leading to a population with shorter linkage disequilibrium (**LD**) blocks, which improves mapping precision, albeit at the expense of power [32,34].

The longest running mouse AIL was generated by crossing LG/J and SM/J inbred strains, which were selectively bred for large and small body size. We obtained this AIL in 2006 at generation 33 from Dr. James Cheverud (Jmc: LG,SM-G_{33}, Washington University in St. Louis). Since then, we have collected genotype and phenotype information from multiple generations, including F_{34} [16,35–38], F_{39}-F_{43} and F_{50-56} [39]. Our previous publications using the F_{34} generation used a custom Illumina Infinium genotyping microarray to obtain genotypes for 4,593 SNPs [35,36]. Although not previously published, we also collected phenotype information from the F_{39}-F_{43} generations, including body weight, fear conditioning, locomotor activity in response to methamphetamine, and the light dark test for anxiety.

In the present study, we used genotyping-by-sequencing (**GBS**), which is a reduced-

representation sequencing method [40–42], to obtain a much denser set of SNPs in the F_{34} and

to genotype mice from the F_{39}-F_{43} generations for the first time. With this denser set of SNPs,

we attempted to identify novel loci in the F_{34}s that were not detected using the sparse SNPs. We

also performed a GWAS on using the mice from the F_{39}-F_{43} AILs. Because F_{39}-F_{43} AILs are

descendants of the F_{34}, they are uniquely suited to be a replication population for GWAS in the

F_{34} generation. Moreover, F_{39}-F_{43} may provide improved resolution and allow for the discovery

of novel loci not detected in the F_{34} generation. We also performed a mega-analysis of F_{34} mice

and F_{39}-F_{43} mice to identify loci that were not identified in either individual dataset. Finally, we

explored whether imputation from the array SNPs could have provided the additional coverage

we obtained using the denser GBS genotypes. This study investigates the effect of SNP density

on association analyses, addresses rarely explored questions about replication in outbred

rodent populations, and identifies novel loci associated with a range of physiological and

behavioral traits in mice.

Results

We used 214 males and 214 females from generation F_{34} (Aap:LG,SM-G34) and 305

males and 295 females from generations F_{39-43}. We performed association studies on 79 traits in

F_{34} AIL and 49 traits in F_{39-43} AIL (S1 Table). F_{34} mice had been previously genotyped on a

custom SNP array [35,36]. The average minor allele frequency (**MAF**) of those 4,593 array

SNPs was 0.388 (Fig 1). To obtain a denser set of SNP markers, we used genotyping-by-

sequencing in F_{34} and F_{39-43} AIL mice. Since data about the F_{39-43} AIL mice had been collected

over the span of approximately two years, we carefully considered the possibility of sample

contamination and sample mislabeling [43]. We removed samples based on four major features:

heterozygosity distribution, number of reads aligned to sex chromosomes, discrepancies

between pedigree and genetic kinship relatedness, and coat color genotype to phenotype

mismatch (see Methods; S1 and S2 Figs). The final SNP sets included 60,392 GBS-derived

SNPs in 428 F_{34} AIL mice, 59,790 GBS-derived SNPs in 600 F_{39-43} AIL mice, and 58,461 GBS-

derived SNPs that existed in both F_{34} and F_{39-43} AIL mice (S2 Table). The MAF for the GBS

SNPs was 0.382 in F_{34}, 0.358 in F_{39-43}, and 0.370 in F_{34} and F_{39-43} (Fig 1). There were 66 SNPs

called from our GBS data that were also present on the genotyping array. The genotype

concordance rate, which reflects the sum of errors from both sets of genotypes, was 95.4% (S3

Fig). We found that LD decay rates using F_{34} array, F_{34} GBS, F_{39-43} GBS, and F_{34} and F_{39-43} GBS

genotypes were generally similar to one another, though levels of LD using the GBS genotypes

appear to be slightly reduced in the later generations of AILs (S4 Fig).

GBS genotypes produced more significant associations than array genotypes in

F_{34}

We used a linear mixed model (**LMM**) as implemented in GEMMA [44] to perform

GWAS. We used the leave-one-chromosome-out (**LOCO**) approach to address the problem of

proximal contamination, as previously described [39,45–47]. We performed GWAS using both

the sparse array SNPs and the dense GBS SNPs to determine whether additional SNPs would

produce additional genome-wide significant associations. Autosomal and X chromosome SNPs

were included in all GWAS. We obtained adjusted significance threshold for each SNP set using

MultiTrans and SLIDES [48,49].To select independently associated loci ("lead loci"), we used a

LD-based clumping method implemented in PLINK to group SNPs that passed the adjusted

genome-wide significance thresholds over a large genomic region flanking the index SNP [50].

Applying the most stringent clumping parameters (r^2 = 0.1 and sliding window size = 12,150kb,

S3 Table), we identified 110 significant lead loci in 49 out of 79 F_{34} phenotypes using the GBS

SNPs. In contrast, we identified 83 significant lead loci in 45 out of 79 F_{34} phenotypes using the

sparse array SNPs. Among F_{34} lead loci, 36 were uniquely identified in the GBS GWAS,

whereas 11 were unique to the array GWAS. GBS SNPs consistently yielded more significant

lead loci compared to array SNPs regardless of the clumping parameter values (S3 Table),

indicating that a dense marker panel was able to detect more association signals compared to a

sparse marker panel. Significant lead loci in F_{34} GBS and array are summarized in Table S4.

We considered whether the disparity between the numbers of loci identified by the two

SNP sets could be resolved by imputation. We used LG/J and SM/J whole genome sequencing

data as reference panels [51] and performed imputation on array and GBS SNPs using Beagle

v4.1 [52]. After QC filtering, we obtained 4.3M SNPs imputed from the array SNPs and 4.1M

SNPs imputed from the GBS SNPs. More imputed GBS SNPs were filtered out because GBS

SNPs were called from genotype probabilities, thus introducing uncertainty in imputed SNPs.

We found that imputed array genotypes and imputed GBS genotypes did not meaningfully

increase the number of new loci discoveries (Fig S5), indicating that the greater number of

genome-wide significant associations obtained using the GBS SNPs could not be overcome by

imputation.

Under a polygenic model where a large number of additive common variants contribute

to a complex trait, heritability estimates could be higher when more SNPs are considered [53].

Given that there were more GBS SNPs than array SNPs, we used autosomal SNPs to examine

whether GBS SNPs would generate higher SNP heritability estimates compared to the sparse

array SNPs. Heritability estimates were similar for the two SNP sets, with the exception of

agouti coat color, which showed marginally greater heritability for the GBS SNPs (S6 Fig; S5

Table)**.** Our results show that while the denser GBS SNP set was able to identify more genome-

wide significant loci, greater SNP density did not improve the polygenic signal.

Genetic correlation and SNP heritability of traits in F_{34} and F_{39-43}

One goal of collecting phenotypic information in the F_{39-43} generations was to evaluate whether or not the genetic signal driving the phenotypes are consistent across cohorts. To this end, we used two approaches to evaluate replication between F_{34} and F_{39}-F_{43}. First, we used autosomal SNPs to calculate genetic correlations between the F_{34} and F_{39-43} generations for body weight, coat color, and locomotor activity phenotypes (S6 Table). Locomotor activity on days 1 and 2, albino and agouti coat color, and body weight were highly genetically correlated (r_Gs >0.7). In contrast, locomotor activity on day 3 showed a significant but weaker genetic correlation (r_G=0.577), perhaps reflecting variability in the quality of the methamphetamine injection, which was not given on days 1 and 2. Overall, these results suggest that genetic influences on these traits were broadly similar in the two cohorts. Despite the relatively robust genetic correlations, SNP heritability was consistently lower in the F_{39-43} compared to the F_{34}, possibly a result of increased experimental variance introduced by our extended phenotype collection period (Fig 2; S7 Table).

Replication of loci detected in F_{34} using F_{39-43}

Using the dense GBS SNP set, we identified 27 genome-wide significant loci in 21 out of 49 F_{39-43} phenotypes. A subset of those traits, including coat color, body weight, and locomotor activity, were also phenotyped in the F_{34} AILs (S8 Table). To assess replication, we determined whether loci that were significant in one cohort (either F_{34} or F_{39-43}) would also be significant in the other. We termed the cohort in which a locus was initially discovered as its "discovery set" and the cohort we attempted replication in as the "replication set" (Table 1). Coat color phenotypes (both albino and agouti) are Mendelian traits and thus served as positive control. As expected, all body weight loci replicated. All three body weight loci identified in the F_{34} were

replicated at nominal levels of significance ($p<0.05$) in F_{39-43}; similarly, the only body weight

locus identified in F_{39-43} was replicated in F_{34}. However, none of the five locomotor activity

associated loci were replicated in the reciprocal (replication) cohorts. We also used the "sign

test" to determine whether the directions of the effect (beta) of the loci were in the same

direction between the discovery and replication cohorts. We found that 11 of 13 loci passed this

much less stringent test of replication. The two loci that did not pass the sign test were the two

locomotor loci "discovered" in F_{39-43} (Table 1).

Table 1. Replication of significant SNPs between F_{34} and $F_{39\text{-}43}$ AIL association analyses. "Discovery set" indicates the AIL generation that significant SNPs were identified. "Replication set" shows the association p-value, β estimates, etc. of the "discovery set" significant SNPs in the replication AIL generation. SNPs that replicated (p<0.05, same sign for the beta) between F_{34} and $F_{39\text{-}43}$ are highlighted in bold italics. Genetic correlations for phenotypes measured in both F_{34} and $F_{39\text{-}43}$ are listed (see also Supplementary Table 6).

			Discovery set				Replication set			
Phenotype	rG(*s.e.*)(*p<0.05)	SNP	P F34 GBS	af	beta	se	P F3943 GBS replicate	af	beta	se
Body weight		*chr4.66414508*	*8.06E-08*	0.419	-2.50E-01	4.56E-02	*1.56E-02*	0.406	-1.05E-01	4.33E-02
	0.711(0.25)*	*chr6.81405109*	*5.89E-06*	0.497	2.06E-01	4.51E-02	*2.36E-02*	0.518	9.48E-02	4.18E-02
		chr14.79312393	*7.53E-06*	0.514	-2.01E-01	4.45E-02	*1.44E-02*	0.566	-1.04E-01	4.26E-02
Coat color, albino	0.967(0.04)*	*chr7.87642045*	*5.00E-106*	0.432	-5.81E-01	1.91E-02	*2.85E-163*	0.387	-6.07E-01	1.55E-02
Coat color, agouti	0.971(0.04)*	*chr2.154464466*	*9.43E-191*	0.129	9.39E-01	1.25E-02	*5.70E-93*	0.207	7.20E-01	2.57E-02
Locomotor test day 1, total distance travelled in 30min	0.968(0.24)*	chr19.21812298	9.28E-07	0.461	-6.90E+02	1.39E+02	7.72E-01	0.52	-1.94E-02	6.74E-02
Locomotor test day2, total distance travelled in 30min	0.988(0.19)*	chr7.45084416	1.12E-05	0.246	6.77E+02	1.53E+02	3.40E-01	0.217	7.38E-02	7.77E-02
		chr8.17410225	4.96E-06	0.171	7.91E+02	1.71E+02	4.03E-01	0.192	7.04E-02	8.43E-02
			F3943 GBS				F34 GBS replicate			
Body weight	0.711(0.25)*	*chr14.82586326*	*2.63E-06*	0.658	-2.09E-01	4.43E-02	*2.87E-05*	0.575	-1.89E-01	4.50E-02
Coat color, albino	0.967(0.04)*	*chr7.87255156*	*5.91E-167*	0.389	-6.24E-01	1.57E-02	*7.80E-97*	0.444	-5.75E-01	2.07E-02
Coat color, agouti	0.971(0.04)*	*chr2.155091628*	*1.78E-115*	0.218	7.42E-01	2.17E-02	*1.51E-185*	0.135	8.98E-01	1.33E-02
Locomotor test day 2, total distance travelled in 30min	0.988(0.19)*	chr15.67235072	3.21E-06	0.47	3.14E-01	6.63E-02	1.69E-01	0.522	-1.72E+02	1.25E+02
Locomotor test day 3, total distance travelled in 30min	0.577(0.22)*	chr7.113250866	5.88E-06	0.389	3.29E-01	7.20E-02	8.27E-01	0.483	-1.21E+02	5.45E+02

Mega-analysis

200   Due to the high genetic correlations between cohorts (S6 Table), we suspected that a

201   mega-analysis using the combined sample set would allow for the identification of additional

202   loci. This analysis identified four novel genome-wide significant associations that were not

203   discovered in either individual cohort (Fig 3; Table 2; S9 Table).

Table 2. Lead SNPs in F_{34} and F_{39-43} (N=1028) mega-analysis in locomotor activity (day 1 – 3) and body weight. Significant SNPs are clumped using parameters r_2=0.1, 12150kb. Rows in bold indicate SNPs that were discovered in the mega-analysis but not in individual F_{34} or F_{39-43} AILs.

Phenotype	SNP	P	af	Beta	se	Discovery set
Locomotor test day 1, total distance travelled in 30min	**chr1.122479820**	**8.84E-06**	**0.292**	**-2.03E-01**	**4.56E-02**	**Combined set**
Locomotor test day 1, total distance travelled in 30min	**chr10.104988207**	**1.94E-07**	**0.373**	**2.16E-01**	**4.13E-02**	**Combined set**
Locomotor test day 2, total distance travelled in 30min	**chr1.40907532**	**2.65E-07**	**0.664**	**2.27E-01**	**4.38E-02**	**Combined set**
Locomotor test day 2, total distance travelled in 30min	**chr1.123997667**	**3.53E-08**	**0.289**	**-2.53E-01**	**4.56E-02**	**Combined set**
Body weight	**chr2.157502044**	**1.10E-05**	**0.42**	**-1.35E-01**	**3.06E-02**	**Combined set**
Body weight	chr4.66866758	1.64E-09	0.403	-1.84E-01	3.01E-02	F_{34} array, F_{34} GBS, combined set
Body weight	chr6.81267890	8.61E-07	0.507	1.40E-01	2.83E-02	F_{34} GBS, combined set
Body weight	chr14.82672838	2.06E-10	0.623	-1.90E-01	2.96E-02	F_{34} array, F_{34} GBS, F_{39-43} GBS, combined set

Discussion

Outbred model organism populations have become important resources for genetics

studies. We used F_{34} and F_{39-43} generations of a LG/J x SM/J AIL to investigate how SNP

density influences GWAS results, heritability, and genetic correlations. Using the ~60K SNPs in

the F_{34} GBS set, we found 110 loci, of which 36 were unique to the GBS genotypes. Notably,

F_{34} array genotypes were able to detect the majority of loci that were identified in F_{34} GBS SNP

set, indicating that many but not all loci were successfully 'tagged' by the sparse array SNPs.

Although the genetic correlations between traits measured in F_{34} and F_{39-43} were high,

replication was incomplete, with the locomotor traits showing especially poor replication. Finally,

imputation to reference panels increased the number of SNPs available for analysis but did not

meaningfully enhance the number of loci we discovered because it did not improve our ability to

capture recombination events.

Previous work from our lab used the array genotypes described in this study to map

QTLs for various behavioral and physiological traits in 688 F_{34} AILs. With F_{34} GBS SNPs, which

were only available for 428 of the initial 688 AIL mice, we were able to replicate many of our

previously published loci. For instance, using the array SNPs in the F_{34} generation in

conjunction with the LMM software QTLRel, Cheng et al. [35] found a significant locus on

chromosome 8 for locomotor day 2 activity that contained only one gene: *Csmd1* (CUB and

sushi multiple domains 1). Gonzales et al. [39] replicated this finding in F_{50-56} AIL and identified a

cis-eQTL mapped to the same region. *Csmd1* mutant mice showed increased locomotor activity

compared to wild-type and heterozygous mice, indicating that *Csmd1* is likely a causal gene for

locomotor and related traits [39]. We replicated this locus in our own analysis of the F_{34} array

and GBS SNP, using GEMMA rather than QTLRel (S7 Fig). We also replicated a locus on

chromosome 17 for distance traveled in the periphery in the open field test (Fig 4; [36,39]), three

loci on chromosomes 4, 6, and 14 for body weight (Supplementary Fig 7; [54]), one locus on

chromosome 7 for mean corpuscular hemoglobin concentrations (MCHC, complete blood count;

S7 Fig; [55]), and numerous loci on chromosome 4, 6, 7, 8, and 11 for muscle weights

(Supplementary Fig 7; [37]). We noticed that while using the same set of F_{34} array genotypes,

some previously published loci were not detected in our GWAS. The most likely explanation is

that we had only 428 of the 688 mice used in the previous publications in this study.

Methodological differences between prior studies and the current study, such as the use of

QTLRel rather than GEMMA and the choice of pedigree rather than genotypes in relatedness

calculation, may also lead to lack of complete replication [56].

F_{39-43} AILs replicated some, but not all, significant loci identified in F_{34}, despite generally

high genetic correlations between the two cohorts. Significant loci for coat color, which are

monogenic and viewed as positive controls, were consistent between the two AIL cohorts. Loci

for body weight were fully replicated between F_{34} and F_{39-43} using the p<0.05 cutoff for the

replication cohort, whereas loci for locomotor activity were not. Nevertheless, the beta estimates

for all but two loci shared the same sign, which provides modest evidence for replication. The

circumstances under which the F_{39-43} data were collected were not ideal, likely contributing to

their lower heritability and the limited replication. Unlike the F_{34} dataset, the F_{39-43} used multiple

technicians to conduct the behavioral tests and occurred over a prolonged period of time in

which numerous environmental factors may have changed.

We performed a mega-analysis using F_{34} and F_{39-43} AIL mice. The combined dataset

increased our power and allowed us to identify four novel genome-wide significant associations

that were not detected in either the F_{34} or the F_{39-43} cohorts. For example, the mega-analysis

identified a locus for body weight on chromosome 2 (Fig S7). Parker et al. [54] identified the

same locus using an integrated analysis of LG/J x SM/J F_2 and F_{34} AILs.

Many GWAS use a 1.0~2.0 LOD support interval to approximate the size of the

association region (see [57,58]). The LOD support interval, proposed by Conneally et al. [59]

and Lander & Botstein [60], is a simple confidence interval method involving converting the p-

value of the peak locus into a LOD score, subtracting "drop size" from the peak locus LOD

score, and finding the two physical positions to the left and to the right of the peak locus location

that correspond to the subtracted LOD score. Although Mangin et al. [61] showed via simulation

that the boundaries of LOD support intervals depend on effect size, others observed that a 1.0 ~

2.0 LOD support interval accurately captures ~95% coverage of the true location of the loci

when using a dense set of markers [60,62,63]. In the present study, we considered using LOD

support intervals but found that the sparse array SNPs produced misleadingly large support

intervals. Various methods have been proposed for calculating confidence intervals in

analogous situations (e.g. [12,64]). Rather than adopting a formal method, we compared

LocusZoom plots of the same locus region between array SNPs and the GBS SNPs (S7 Fig;

[65]). For example, the benefit of the denser SNP coverage is easily observed in the locus on

chromosome 7 for "complete blood count, repeat measure of retic parameters, cell hemoglobin

concentration mean" (Fig S7). Thus, there are advantages of dense SNP sets that go beyond

the ability to discover additional loci (Table 1).

Our study has notable limitations. First, not all F_{34} and F_{39-43} animals that were

phenotyped were later genotyped by GBS due to missing tissue samples, which in turn lowered

our sample size and reduced the power of association analyses. Second, F_{39-43} traits have been

collected by different technicians over the span of several years, which introduced noise and

diminished trait heritability (Fig 2).

The present study explored the role of marker density and imputation in GWAS.

Furthermore, the combination of denser marker coverage and the addition of 600 F_{39-43} AIL mice

allowed us to identify novel loci for locomotor activity, open field test, fear conditioning, light dark

test for anxiety, complete blood count, iron content in liver and spleen, and muscle weight. Our

findings add to the large body of phenotypic/genotypic data available for the LG/SM AIL, which

can be found on GeneNetwork [http://www.genenetwork.org].

Materials and Methods

Animals

All mice used in this study were members of the advanced intercross line (**AIL**) between

LG/J and SM/J that was originally created by Dr. James Cheverud (Washington University, St.

Louis, MO). The AIL line has been maintained in the Palmer laboratory since generation F_{33}.

Age and exact number of animals tested in each phenotype are described in S1 Table. Several

previous publications [16,35–38,54,55] have reported on association analyses of the F_{34} mice

(N=428). No prior publications have described the $F_{39\text{-}43}$ generations (N=600). The sample size

of F_{34} mice reported in this study (N=428) is smaller than that in previous publications of F_{34}

(N=688) because we only sequenced a subset of F_{34} animals using GBS. With the exception of

coat color and locomotor activities, we quantile normalized all phenotypes. Coat color traits

were coded in binary numbers (albino: 1 = white, 0 = non-white; agouti: 1 = tan, 0 = black,

NA=white). Locomotor activity traits were not quantile transformed in order to follow the

guideline described in Cheng et al. [35] for direct comparison.

F_{34}, $F_{39\text{-}43}$ Phenotypes

We have previously described the phenotyping of F_{34} animals for locomotor activity [35],

fear conditioning [36], open field [36], coat color, body weight [54], complete blood counts [55],

heart and tibia measurements [37], muscle weight [37]. Iron content in liver and spleen, which

have not been previously reported in these mice, was measured by atomic absorption

spectrophotometry, as described in Gardenghi et al. [66] and Graziano, Grady and Cerami [67].

Although the phenotyping of F_{39-43} animals has not been previously reported, we used method

that were identical to those previously reported for locomotor activity [35], open field [36], coat

color, body weight [54], and light/dark test for anxiety [15].

F_{34} AIL Array Genotypes

316   F_{34} animals had been genotyped on a custom SNP array on the Illumina Infinium

317 platform [35,36], which yielded a set of 4,593 SNPs on autosomes and X chromosome that we

318 refer to as 'array SNPs'.

F_{34} and F_{39-43} GBS Genotypes

F_{34} and F_{39-43} animals were genotyped using genotyping-by-sequencing (**GBS**), which is

a reduced-representation genome sequencing method [1,39]. We used the same protocol for

GBS library preparation that was described in Gonzales et al [39]. We called GBS genotype

probabilities using ANGSD [68]. GBS identified 1,667,920 autosomal and 43,015 X-

chromosome SNPs. To fill in missing genotypes at SNPs where some but not all mice had calls,

we ran within-sample imputation using Beagle v4.1, which generated hard call genotypes as

well as genotype probabilities [52]. After imputation, only SNPs that had dosage $r^2 > 0.9$ were

retained. We removed SNPs with minor allele frequency < 0.1 and SNPs with $p < 1.0 \times 10^{-6}$ in the

Chi-square test of Hardy–Weinberg Equilibrium (**HWE**) (S2 Table). All phenotype and GBS

genotype data are deposited in GeneNetwork (http://www.genenetwork.org).

QC of individuals

We have found that large genetic studies are often hampered by cross-contamination between samples and sample mix-ups. We used four features of the data to identify problematic samples: heterozygosity distribution, proportion of reads aligned to sex chromosomes, pedigree/kinship, and coat color. We first examined heterozygosity across autosomes and removed animals where the proportion of heterozygosity that was more than 3 standard deviations from the mean (S1 Fig). Next, we sought to identify animals in which the recorded sex did not agree with the sequencing data. We compared the ratio of reads mapped to the X and Y chromosomes. The 95% CI for this ratio was 196.84 to 214.3 in females and 2.13 to 2.18 in males. Twenty-two F_{34} and F_{39-43} animals were removed because their sex (as determined by reads ratio) did not agree with their recorded sex; we assumed this discrepancy was due to sample mix-ups. To further identify mislabeled samples, we calculated kinship coefficients based on the full AIL pedigree using QTLRel. We then calculated a genetic relatedness matrix (**GRM**) using IBDLD, which estimates identity by descent using genotype data. The comparison between pedigree kinship relatedness and genetic kinship relatedness identified 7 pairs of animals that showed obvious disagreement between kinship coefficients and the GRM. Lastly, we excluded 14 F_{39-43} animals that showed discordance between their recorded coat color and their genotypes at markers flanking *Tyr*, which causes albinism in mice. The numbers of animals filtered at each step are listed in S2 Table. Some animals were detected by more than one QC step, substantiating our belief that these samples were erroneous.

At the end of SNP and sample filtering, we had 59,561 autosomal and 831 X chromosome SNPs in F_{34}, 58,966 autosomal and 824 X chromosome SNPs in F_{39-43}, and 57,635 autosomal and 826 X chromosome SNPs in the combined F_{34} and F_{39-43} set (S2 Table). GBS genotype quality was estimated by examining concordance between the 66 SNPs that were present in both the array and GBS genotyping results.

LD decay

358   Average LD (r^2) was calculated using allele frequency matched SNPs (MAF difference <

359   0.05) within 100,000bp distance, as described in Parker et al. [1].

Imputation to LG/J and SM/J reference panels

F_{34} array genotypes (N=428) and F_{34} GBS genotypes (N=428) were imputed to LG/J and

SM/J whole genome sequence data [51] using BEAGLE. For F_{34} array imputation, we used a

large window size (100,000 SNPs and 45,000 SNPs overlap). Imputation to reference panels

yielded 4.3 million SNPs for F_{34} array and F_{34} GBS imputed sets. Imputed SNPs with $DR^2 > 0.9$,

MAF > 0.1, HWE p value > 1.0×10^{-6} were retained, resulting in 4.1M imputed F_{34} GBS SNPs

and 4.3M imputed F_{34} array SNPs.

Genome-wide association analysis (GWAS)

We used the linear mixed model, as implemented in GEMMA [44], to perform a GWAS

that accounted for the complex familial relationships among the AIL mice [35,39]. We used the

leave-one-chromosome-out (**LOCO**) approach to calculate genetic relatedness matrix, which

effectively circumvented the problem of proximal contamination [46]. Separate GWAS were

performed using F_{34} array genotypes, F_{34} GBS genotypes, and F_{39-43} GBS genotypes. Apart

from coat color (binary trait) and locomotor activity, raw phenotypes were quantile normalized

prior to analysis. Locomotor activity was not quantile normalized because the trait was

reasonably normally distributed already and because we wanted our analysis to match those

performed in Cheng et al [35]. Because F_{34} AIL had already been studied using array genotypes

[35] and mapped using QTLRel [56], we used the same covariates as described in Cheng et al.

[35] in order to examine whether our array and GBS GWAS would replicate their findings. We

included sex and body weight as covariates for locomotor activity traits (see covariates used in

[35])and sex, age, and coat color as covariates for fear conditioning and open field test in F_{34}

AILs (see covariates used in [36]). We used sex and age as covariates for all other phenotypes.

Covariates for each analysis are shown in S1 Table. Finally, we performed mega-analysis of F_{34}

and F_{39-43} animals (N=1,028) for body weight, coat color, and locomotor activity, since these

traits were measured in the same way in both cohorts. For the mega-analyses, locomotor

activity was quantile normalized after the combination of the two datasets to ensure that data

were normally distributed across generations.

Identifying suspicious SNPs

Some significant SNPs in F_{34} GWAS and in the mega-analysis of F_{34} and F_{39-43} were

suspicious because nearby SNPs, which would have been expected to be in high LD (a very

strong assumption in an AIL), did not have high -log10 values. We only examined SNPs that

obtained significant p-values; these examinations reveled that these SNPs had suspicious ratios

of heterozygotes to homozygotes calls and had corresponding HWE p-values that were close to

our 1.0×10^{-6} threshold (S10 and S11 Tables). To avoid counting these as novel loci, we

removed those SNPs prior to summarizing our results as they likely reflected genotyping errors.

Selecting independent significant SNPs

To identify independent "lead loci" among significant GWAS SNPs that surpass the

significance threshold, we used the LD-based clumping method in PLINK v1.9. We empirically

chose clumping parameters (r^2 = 0.1 and sliding window size = 12,150kb) that gave us a

conservative set of independent SNPs (S3 Table). For the coat color phenotypes, we found that

multiple SNPs remained significant even after LD-based clumping, presumably due to the

extremely significant associations at these Mendelian loci. In these cases, we used a stepwise

model selection procedure in GCTA [69] and performed association analyses conditioning on

the most significant SNPs.

Significance thresholds

We used MultiTrans and SLIDE to set significance thresholds for the GWAS [48,49].

MultiTrans and SLIDE are methods that assume multivariate normal distribution of the

phenotypes, which in LMM models, contain a covariance structure due to various degrees of

relatedness among individuals. We were curious to see whether MultiTrans/SLIDE produces

significance thresholds drastically different from the threshold we obtained from a standard

permutation test ('naïve permutation' as per Cheng et al. [46]). We performed 1,000

permutations using the F_{34} GBS genotypes and the phenotypic data from locomotor activity

(days 1, 2, and 3). We found that the 95th percentile values for these permutations were 4.65,

4.79, and 4.85, respectively, which were very similar to 4.85, the threshold obtained from

MultiTrans using the same data. Thus, the thresholds presented here were obtained from

MultiTrans but are similar (if anything slightly more conservative) than thresholds we would have

obtained had we used permutation. Because the effective number of tests depends on the

number of SNPs and the specific animals used in GWAS, we obtained a unique adjusted

significance threshold for each SNP set in each animal cohort (S12 Table).

Genetic correlation and heritability estimates between F_{34} and F_{39-43} phenotypes

427   Locomotor activity, body weight, and coat color had been measured in both F_{34} and F_{39-}

428 $_{43}$ populations. We calculated both SNP heritability and genetic correlations between F_{34} and

429 F_{39-43} animals using GCTA bivariate GREML analysis [69]. Because F_{39-43} day 1 locomotor

430 activity data were not normally distributed, we quantile normalized locomotor activity data when

431 estimating SNP heritabilities and genetic correlations.

LocusZoom Plots

434 LocusZoom plots were generated using the standalone implementation of LocusZoom [65],

435 using LD scores calculated from PLINK v.1.9 --ld option and mm10 gene annotation file

436 downloaded from USCS genome browser.

Ethics Statement

439 All procedures were approved by the Institutional Animal Care and Use Committee (IACUC

440 protocol: S15226) Euthanasia was accomplished using CO_2 asphyxiation followed by cervical

441 dislocation

442   .

Acknowledgements

444 We would like to recognize Jackie Lim and Kaitlin Samocha for collecting F_{34} AIL phenotype

445 data and Ryan Walters for collecting F_{39-43} AIL phenotype data. We wish to acknowledge Alex

446 Gileta for input on a draft of this manuscript.

References

1. Parker CC, Gopalakrishnan S, Carbonetto P, Gonzales NM, Leung E, Park YJ, et al. Genome-wide association study of behavioral, physiological and gene expression traits in outbred CFW mice. Nat Genet. 2016;48(8):919.

2. Gatti DM, Svenson KL, Shabalin A, Wu L-Y, Valdar W, Simecek P, et al. Quantitative trait locus mapping methods for diversity outbred mice. G3 Genes Genomes Genet. 2014;4(9):1623–1633.

3. Chesler EJ. Out of the bottleneck: the Diversity Outcross and Collaborative Cross mouse populations in behavioral genetics research. Mamm Genome. 2014;25(1–2):3–11.

4. Churchill GA, Gatti DM, Munger SC, Svenson KL. The diversity outbred mouse population. Mamm Genome. 2012;23(9–10):713–718.

5. Consortium CC. The genome architecture of the Collaborative Cross mouse genetic reference population. Genetics. 2012;190(2):389–401.

6. Talbot CJ, Nicod A, Cherny SS, Fulker DW, Collins AC, Flint J. High-resolution mapping of quantitative trait loci in outbred mice. Nat Genet. 1999;21(3):305.

7. Demarest K, Koyner J, McCaughran J, Cipp L, Hitzemann R. Further characterization and high-resolution mapping of quantitative trait loci for ethanol-induced locomotor activity. Behav Genet. 2001;31(1):79–91.

8. Valdar W, Solberg LC, Gauguier D, Burnett S, Klenerman P, Cookson WO, et al. Genome-wide genetic association of complex traits in heterogeneous stock mice. Nat Genet. 2006;38(8):879.

9. Ghazalpour A, Doss S, Kang H, Farber C, Wen P-Z, Brozell A, et al. High-resolution mapping of gene expression using association in an outbred mouse stock. PLoS Genet. 2008;4(8):e1000149.

10. Svenson KL, Gatti DM, Valdar W, Welsh CE, Cheng R, Chesler EJ, et al. High-resolution genetic mapping using the Mouse Diversity outbred population. Genetics. 2012;190(2):437–447.

475   11.   Yalcin B, Willis-Owen SA, Fullerton J, Meesaq A, Deacon RM, Rawlins JNP, et al. Genetic dissection
476   of a behavioral quantitative trait locus shows that Rgs2 modulates anxiety in mice. Nat Genet.
477   2004;36(11):1197.

478   12.   Nicod J, Davies RW, Cai N, Hassett C, Goodstadt L, Cosgrove C, et al. Genome-wide association of
479   multiple complex traits in outbred mice by ultra-low-coverage sequencing. Nat Genet.
480   2016;48(8):912.

481   13.   Carbonetto P, Cheng R, Gyekis JP, Parker CC, Blizard DA, Palmer AA, et al. Discovery and
482   refinement of muscle weight QTLs in B6$\times$ D2 advanced intercross mice. Physiol Genomics.
483   2014;46(16):571–582.

484   14.   Coyner J, McGuire JL, Parker CC, Ursano RJ, Palmer AA, Johnson LR. Mice selectively bred for High
485   and Low fear behavior show differences in the number of pMAPK (p44/42 ERK) expressing neurons
486   in lateral amygdala following Pavlovian fear conditioning. Neurobiol Learn Mem. 2014;112:195–
487   203.

488   15.   Parker CC, Cheng R, Sokoloff G, Palmer AA. Genome-wide association for methamphetamine
489   sensitivity in an advanced intercross mouse line. Genes Brain Behav. 2012;11(1):52–61.

490   16.   Samocha KE, Lim JE, Cheng R, Sokoloff G, Palmer AA. Fine mapping of QTL for prepulse inhibition in
491   LG/J and SM/J mice using F2 and advanced intercross lines. Genes Brain Behav. 2010;9(7):759–
492   767.

493   17.   Hernandez Cordero AI, Carbonetto P, Riboni Verri G, Gregory JS, Vandenbergh DJ, P Gyekis J, et al.
494   Replication and discovery of musculoskeletal QTLs in LG/J and SM/J advanced intercross lines.
495   Physiol Rep. 2018;6(4).

496   18.   Baud A, Guryev V, Hummel O, Johannesson M, Hermsen R, Stridh P, et al. Genomes and phenomes
497   of a population of outbred rats and its progenitors. Sci Data. 2014;1:140011.

498   19.   Besnier F, Wahlberg P, Rönnegard L, Ek W, Andersson L, Siegel PB, et al. Fine mapping and
499   replication of QTL in outbred chicken advanced intercross lines. Genet Sel Evol. 2011;43(1):3.

500   20.   Johnsson M, Henriksen R, Höglund A, Fogelholm J, Jensen P, Wright D. Genetical genomics of
501   growth in a chicken model. BMC Genomics. 2018;19(1):72.

502   21.   Guryev V, Koudijs MJ, Berezikov E, Johnson SL, Plasterk RH, Van Eeden FJ, et al. Genetic variation in
503   the zebrafish. Genome Res. 2006;16(4):491–497.

504   22.   Patowary A, Purkanti R, Singh M, Chauhan R, Singh AR, Swarnkar M, et al. A sequence-based
505   variation map of zebrafish. Zebrafish. 2013;10(1):15–20.

506   23.   Mackay TF, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The Drosophila
507   melanogaster genetic reference panel. Nature. 2012;482(7384):173.

508   24.   Vonesch SC, Lamparter D, Mackay TF, Bergmann S, Hafen E. Genome-wide analysis reveals novel
509   regulators of growth in Drosophila melanogaster. PLoS Genet. 2016;12(1):e1005616.

25.   King EG, Macdonald SJ, Long AD. Properties and power of the Drosophila Synthetic Population
Resource for the routine dissection of complex traits. Genetics. 2012;genetics–112.

26.   Kislukhin G, King EG, Walters KN, Macdonald SJ, Long AD. The genetic architecture of
methotrexate toxicity is similar in Drosophila melanogaster and humans. G3 Genes Genomes
Genet. 2013;g3–113.

27.   Marriage TN, King EG, Long AD, Macdonald SJ. Fine-mapping nicotine resistance loci in Drosophila
using a multiparent advanced generation inter-cross population. Genetics. 2014;198(1):45–57.

28.   Doitsidou M, Jarriault S, Poole RJ. Next-generation sequencing-based approaches for mutation
mapping and identification in Caenorhabditis elegans. Genetics. 2016;204(2):451–474.

29.   Diouf IA, Derivot L, Bitton F, Pascual L, Causse M. Water Deficit and Salinity Stress Reveal Many
Specific QTL for Plant Growth and Fruit Quality Traits in Tomato. Front Plant Sci. 2018;9:279.

30.   Cockram J, Mackay I. Genetic Mapping Populations for Conducting High-Resolution Trait Mapping
in Plants. 2018;

31.   Rishmawi L, Bühler J, Jaegle B, Hülskamp M, Koornneef M. Quantitative trait loci controlling leaf
venation in Arabidopsis. Plant Cell Environ. 2017;40(8):1429–1441.

32.   Parker CC, Palmer AA. Dark matter: are mice the solution to missing heritability? Front Genet.
2011;2:32.

33.   Darvasi A, Soller M. Advanced intercross lines, an experimental population for fine genetic
mapping. Genetics. 1995;141(3):1199–1207.

34.   Gonzales NM, Palmer AA. Fine-mapping QTLs in advanced intercross lines and other outbred
populations. Mamm Genome. 2014;25(7–8):271–292.

35.   Cheng R, Lim JE, Samocha KE, Sokoloff G, Abney M, Skol AD, et al. Genome-wide association
studies and the problem of relatedness among advanced intercross lines and other highly
recombinant populations. Genetics. 2010;

36.   Parker CC, Carbonetto P, Sokoloff G, Park YJ, Abney M, Palmer AA. High-resolution genetic
mapping of complex traits from a combined analysis of F2 and advanced intercross mice. Genetics.
2014;198(1):103–116.

37.   Lionikas A, Cheng R, Lim JE, Palmer AA, Blizard DA. Fine-mapping of muscle weight QTL in LG/J and
SM/J intercrosses. Physiol Genomics. 2010;42(1):33–38.

38.   Carroll AM, Cheng R, Collie-Duguid ESR, Meharg C, Scholz ME, Fiering S, et al. Fine-mapping of
genes determining extrafusal fiber properties in murine soleus muscle. Physiol Genomics.
2017;49(3):141–150.

39.   Gonzales NM, Seo J, Hernandez-Cordero AI, Pierre CLS, Gregory JS, Distler MG, et al. Genome wide
association study of behavioral, physiological and gene expression traits in a multigenerational
mouse intercross. bioRxiv. 2017 Dec 8;230920.

40. Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A robust, simple
genotyping-by-sequencing (GBS) approach for high diversity species. PloS One. 2011;6(5):e19379.

41. Davey JW, Hohenlohe PA, Etter PD, Boone JQ, Catchen JM, Blaxter ML. Genome-wide genetic
marker discovery and genotyping using next-generation sequencing. Nat Rev Genet.
2011;12(7):499.

42. Fitzpatrick CJ, Gopalakrishnan S, Cogan ES, Yager LM, Meyer PJ, Lovic V, et al. Variation in the form
of Pavlovian conditioned approach behavior among outbred male Sprague-Dawley rats from
different vendors and colonies: sign-tracking vs. goal-tracking. PloS One. 2013;8(10):e75042.

43. Toker L, Feng M, Pavlidis P. Whose sample is it anyway? Widespread misannotation of samples in
transcriptomics studies. F1000Research (Internet). 2016 Sep 30 (cited 2018 Jul 11);5. Available
from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5034794/

44. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies. Nat
Genet. 2012;44(7):821.

45. Listgarten J, Lippert C, Kadie CM, Davidson RI, Eskin E, Heckerman D. Improved linear mixed
models for genome-wide association studies. Nat Methods. 2012;9(6]:525.

46. Cheng R, Parker CC, Abney M, Palmer AA. Practical considerations regarding the use of genotype
and pedigree data to model relatedness in the context of genome-wide association studies. G3
Genes Genomes Genet. 2013;g3–113.

47. Yang J, Zaitlen NA, Goddard ME, Visscher PM, Price AL. Advantages and pitfalls in the application of
mixed-model association methods. Nat Genet. 2014;46(2):100.

48. Joo JWJ, Hormozdiari F, Han B, Eskin E. Multiple testing correction in linear mixed models. Genome
Biol. 2016;17(1):62.

49. Han B, Kang HM, Eskin E. Rapid and accurate multiple testing correction and power estimation for
millions of correlated markers. PLoS Genet. 2009;5(4):e1000456.

50. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for
whole-genome association and population-based linkage analyses. Am J Hum Genet.
2007;81(3):559–575.

51. Nikolskiy I, Conrad DF, Chun S, Fay JC, Cheverud JM, Lawson HA. Using whole-genome sequences
of the LG/J and SM/J inbred mouse strains to prioritize quantitative trait genes and nucleotides.
BMC Genomics. 2015;16(1):415.

52. Browning SR, Browning BL. Rapid and accurate haplotype phasing and missing-data inference for
whole-genome association studies by use of localized haplotype clustering. Am J Hum Genet.
2007;81(5):1084–1097.

53. Yang J, Zeng J, Goddard ME, Wray NR, Visscher PM. Concepts, estimation and interpretation of
SNP-based heritability. Nat Genet. 2017 Sep;49(9):1304–10.

54.   Parker CC, Cheng R, Sokoloff G, Lim JE, Skol AD, Abney M, et al. Fine-mapping alleles for body
581   weight in LG/J × SM/J F$_2$ and F$_{34}$ advanced
582   intercross lines. Mamm Genome. 2011 Oct 1;22(9–10):563.

583   55.   Bartnikas TB, Parker CC, Cheng R, Campagna DR, Lim JE, Palmer AA, et al. QTLs for murine red
584   blood cell parameters in LG/J and SM/J F2 and advanced intercross lines. Mamm Genome. 2012
585   Jun;23(5–6):356–66.

586   56.   Cheng R, Abney M, Palmer AA, Skol AD. QTLRel: an R package for genome-wide association studies
587   in which relatedness is a concern. BMC Genet. 2011 Jul 27;12:66.

588   57.   Logan RW, Robledo RF, Recla JM, Philip VM, Bubier JA, Jay JJ, et al. High-precision genetic mapping
589   of behavioral traits in the diversity outbred mouse population. Genes Brain Behav.
590   2013;12(4):424–437.

591   58.   Cervino AC, Li G, Edwards S, Zhu J, Laurie C, Tokiwa G, et al. Integrating QTL and high-density SNP
592   analyses in mice to identify Insig2 as a susceptibility gene for plasma cholesterol levels. Genomics.
593   2005;86(5):505–517.

594   59.   Conneally PM, Edwards JH, Kidd KK, Lalouel J-M, Morton NE, Ott J, et al. Report of the committee
595   on methods of linkage analysis and reporting. Cytogenet Genome Res. 1985;40(1–4):356–9.

596   60.   Lander ES, Botstein D. Mapping mendelian factors underlying quantitative traits using RFLP linkage
597   maps. Genetics. 1989;121(1):185–199.

598   61.   Mangin B, Goffinet B, Rebai A. Constructing confidence intervals for QTL location. Genetics.
599   1994;138(4):1301–1308.

600   62.   Dupuis J, Siegmund D. Statistical methods for mapping quantitative trait loci from a dense set of
601   markers. Genetics. 1999;151(1):373–386.

602   63.   Manichaikul A, Dupuis J, Sen S, Broman KW. Poor performance of bootstrap confidence intervals
603   for the location of a quantitative trait locus. Genetics. 2006;

604   64.   Sequencing RG, Baud A, Hermsen R, Guryev V, Stridh P, Graham D, et al. Combined sequence-
605   based and genetic mapping analysis of complex traits in outbred rats. Nat Genet. 2013;45(7):767.

606   65.   Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, et al. LocusZoom: regional
607   visualization of genome-wide association scan results. Bioinformatics. 2010;26(18):2336–2337.

608   66.   Gardenghi S, Marongiu MF, Ramos P, Guy E, Breda L, Chadburn A, et al. Ineffective erythropoiesis
609   in β-thalassemia is characterized by increased iron absorption mediated by down-regulation of
610   hepcidin and up-regulation of ferroportin. Blood. 2007;109(11):5027–5035.

611   67.   Graziano JH, Grady RW, Cerami A. The identification of 2, 3-dihydroxybenzoic acid as a potentially
612   useful iron-chelating drug. J Pharmacol Exp Ther. 1974;190(3):570–575.

613   68.   Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: analysis of next generation sequencing data.
614   BMC Bioinformatics. 2014;15(1):356.

615   69.   Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis.
616   Am J Hum Genet. 2011 Jan 7;88(1):76–82.

Main figure legends

Fig 1. Minor allele frequency distributions for F_{34} array, F_{34} GBS, F_{39}-F_{43} GBS, and F_{34} and
F_{39}-F_{43} GBS SNP sets. MAF distributions are highly comparable between AIL generations.

Fig 2. Chip-heritability estimates in F_{34} and $F_{39\text{-}43}$ AILs. All heritability estimates are highly
significant (p < 1.0E-05; see S7 Table).

Fig 3. Manhattan plots comparing F_{34} GBS, $F_{39\text{-}43}$ GBS, and mega-analysis on locomotor
day 1 test using 57,170 shared SNPs in all AIL generations. Mega-analysis identified a locus
on chromosome 10 (chr10.104988207) that was not detected in the F_{34} or F_{39}-F_{43} alone,
suggesting that mega-analysis enhanced power to detect some loci.

Fig 4. Significant loci on chromosome 17 for open field, distance traveled in periphery in
F_{34} AIL. (**a**) Genome-wide Manhattan plots highlighting the significant loci on chromosome 17
associated with open field, distance traveled in periphery in F_{34} array SNP set (chr17:27427358)
and F_{34} GBS SNP set (chr17:27130383). (**b**) As exemplified in this pair of LocusZoom plots,
GBS SNPs defined the boundaries of the loci much more precisely than array SNPs. GBS
SNPs that are in high LD (r_2 > 0.8, red dots) with lead SNP chr17:27130383 resides between 27
~ 28.2 Mb. In contrast, too few SNPs are present in the array plot to draw any definitive
conclusion about the boundaries or LD pattern in this region. This locus has also been observed
in GWAS for a variety of other traits, including locomotor activity, startle and prepulse inhibition
[39] and open field test for anxiety [36]. LocusZoom plots for all loci identified in this paper are in
Fig S7.

Supporting information

S1 Fig. Autosomal heterozygosity distribution in F_{34}, F_{39-43} AILs. Animals with excessive or insufficient heterozygosity (3 *s.d.* away from mean) were removed from further analysis. As controls, we have sequenced two F_2s of LG and SM, four LG mice and four SM mice (see annotated data points with 1 and 0 heterozygosity).

S2 Fig. Kinship coefficients in F_{34} and F_{39-43} AILs calculated from pedigree against genetic relatedness matrix calculated using IBDLD [49]. Each circle represents a pair of animals, which their genetic kinship relatedness on the x-axis and pedigree kinship relatedness on the y-axis. Color signifies relatedness based on AIIL pedigree. Blue circles represent identical twins, red full siblings, yellow parent-offspring pairs, grey other relationships. Seven animal pairs that deviate from the pedigree relationship clusters were excluded (see black arrows).

S3 Fig. Heatmap showing F_{34} array and F_{34} GBS genotype concordance in percentages, using 66 shared SNPs. "A" codes for the LG/J allele, and "B" codes for the SM/J allele. "AA" genotype concordance between array and GBS is 24.54%, "AB" 43.23%, "BB" 27.60%.

S4 Fig. LD decay in F_{34} array, F_{34} GBS, F_{39-43} GBS, and F_{34} and F_{39-43} GBS SNP sets.

S5 Fig. Manhattan plots comparing 4,593 F_{34} array, 60.3K F_{34} GBS, 4.3M imputed F_{34} array, and 4.1M imputed F_{34} GBS (N=428) SNPs on day 2 locomotor activity. Adjusted significance thresholds for imputed array and GBS SNPs were estimated using LD pruned SNPs (r2=0.1, window size=20kb; PLINK v1.9). Notice that even though the imputed sets have more SNPs (the two right panels), they are frequently blocks of many SNPs with almost identical position and LD=1, therefore making it hard to visualize the additional SNPs.

S6 Fig. SNP heritability using F_{34} GBS and F_{34} array SNPs (slope=1).

S7 Fig. LocusZoom for F_{34} array, F_{34} GBS, F_{39-43} GBS, and mega-analysis QTLs.

S1 Table. List of phenotypes used in GWAS.

S2 Table. SNP and individual QC filter table. Numbers of animals and SNPs remained after each step of filtering are shown per GBS SNP set.

S3 Table. Effect of PLINK v1.9 clump-based pruning parameters on number of independent SNPs remained. At all r_2 values examined, a sliding window size of 12150kb was the first smallest window that yield the most stringent number of clumped SNPs in both array and GBS GWAS.

S4 Table. Lead QTL in F_{34} GBS and F_{34} array GWAS studies across phenotypes. Significant SNPs are clumped using parameters r_2=0.1, 12150kb.

S5 Table. F_{34} GBS and array SNP heritability estimates.

S6 Table. F_{34} and F_{39-43} genetic correlations in locomotor activity, coat color, and body weight.

S7 Table. SNP-heritability comparison between F_{34} and F_{39-43} GBS.

694
695 **S8 Table. Lead QTL in F$_{39-43}$ N=600 GBS GWAS studies across phenotypes.** Significant
696 SNPs are clumped using parameters r_2=0.1, 12150kb.
697
698 **S9 Table. Lead QTL in F$_{34}$ and F$_{39-43}$ (N=1028) mega-analysis across phenotypes.**
699 Significant SNPs are clumped using parameters r_2=0.1, 12150kb.
700
701 **S10 Table. SNPs in F$_{34}$ GBS set with HWE p-values close to 1.0×10-6 cutoff threshold.**
702 These SNPs are removed from QTL summary tables.
703
704 **S11 Table. SNPs in F$_{34}$ and F$_{39-43}$ mega-analysis GBS set with HWE p values close to**
705 **1.0×10-6 cutoff threshold.** These SNPs are removed from QTL summary tables.
706
707 **S12 Table. Adjusted significance threshold for each SNP set and GWAS cohort.**

Phenotype