Superior dispersal ability leads to persistent ecological dominance by *Candida*

pseudoglaebosa in the *Sarracenia purpurea* metacommunity

Primrose J. Boynton[1*], Celeste N. Peterson[2], Anne Pringle[3]

1) Environmental Genomics Working Group, Max-Planck Institute for Evolutionary

Biology, Plön, Germany

2) Department of Biology, Suffolk University, Boston, MA, USA

3) Departments of Botany and Bacteriology, University of Wisconson-Madison,

Madison, WI, USA

* To whom correspondence should be addressed, pboynton@evolbio.mpg.de

Abstract

A large number of descriptive surveys document changes in microbial communities

over time, but direct evidence for the ecological processes mediating succession or

causing ecological dominance remains rare. Differential dispersal may be a key

mechanism. We surveyed microbial diversity within a metacommunity of pitchers of

the model carnivorous plant *Sarracenia purpurea* and discovered the yeast *Candida*

pseudoglaebosa as ecologically dominant. Its frequency in the metacommunity

increased over the growing season, and it was not replaced by other taxa. We next

measured its competitive ability in a manipulative laboratory experiment and tracked

its dispersal over time in nature. Despite its dominance, *C. pseudoglaebosa* is not a

superior competitor. Instead, it is a superior disperser: it arrives in pitchers earlier, and

disperses into more pitchers, than other taxa. Differential dispersal across the spatially

structured metacommunity of individual pitchers emerges as a key driver of the

continuous dominance of *C. pseudoglaebosa* during succession.

Introduction

Primary microbial succession occurs when a microbial community colonizes and develops on a newly available substrate (Fierer *et al.* 2010). The advent of high-throughput sequencing has revolutionized observational studies of microbial succession, enabling researchers to describe the development of microbial communities in fine detail (Copeland *et al.* 2015, Boynton & Greig 2016, Koenig *et al.* 2011, Cutler *et al.* 2014). A variety of successional patterns have been observed. For example, taxon diversity can increase, decrease, or randomly vary with successional time (Cutler *et al.* 2014, Zumsteg *et al.* 2012, Redford & Fierer 2009). However, researchers commonly observe replacement of early-successional taxa by late-successional taxa (*e.g.,* Boynton & Greig 2016, Wolfe *et al.* 2014). An ongoing challenge in microbial ecology is to connect observed ecological patterns to the ecological processes responsible for the patterns.

The development of ecological dominance by one or a few microbes over successional time is a particularly intriguing phenomenon (Boynton & Greig, 2016, Findley *et al.* 2013, Jiao *et al.* 2016), in part because of its obvious parallels to plant and animal ecology (Hillebrand *et al.* 2008). Ecological dominance is apparent when one or a few species comprise most of the individuals or biomass in a community (Hillebrand *et al.* 2008). Environmental filtering, superior competitive ability of the dominant species, and ecosystem engineering can all cause dominance (Wolfe *et al.* 2014, Albergaria *et al.* 2010, Goddard 2008, Nissen & Arneborg 2003, Williams *et al.* 2015). However, the dynamics of microbial dominance have been primarily studied in domesticated systems, and the most frequently reported mechanisms may not be responsible for dominance in all, or even most, natural microbial communities. For

example, environmental filtering and competition may be more important in systems

where human beings have designed environments to favor domesticated microbes

(Wolfe & Dutton 2015) and less important in natural environments with

heterogeneous environmental conditions.

An overlooked mechanism driving ecological dominance in natural systems

may be dispersal. When primary succession occurs on a sterile substrate, all members

of the microbial community must first disperse onto the new substrate before

establishing in the community. A good disperser may prevent the establishment of

other community members through priority effects if it arrives in a habitat first, either

by pre-empting or modifying available niches (Fukami 2015). Additionally, good

dispersers in nearby communities can impact a microbial community during

succession by producing propagules that then disperse into the developing community

(Wilson & McTammany 2016).

Dispersal may emerge as a key driver of ecological dominance in microbial

metacommunities. Metacommunities are physically structured groups of communities

in which individual communities are spatially isolated from one another but linked

through dispersal (Logue *et al.* 2011, Holyoak *et al.* 2005). Community assembly in

metacommunities is a function of dispersal among communities and ecological

processes occurring inside each community. Ecological theory explains how dispersal

can interact with intracommunity processes to maintain metacommunity diversity

(Leibold *et al.* 2004, Holyoak *et al.* 2005). For example, populations occupying low-

quality environmental patches can be maintained by dispersal from high-quality

patches ("mass effects"), or fitness trade-offs between competitive ability and

dispersal ability can mediate species diversity ("patch dynamics"). Theory predicts

that dispersal and competition interact during succession in the individual component

communities of a metacommunity and the result is a hump-shaped relationship

between successional time and species richness (Moquet *et al.* 2003, Sferra *et al.*

2017). Species richness is predicted to be low early in a component community's age,

to increase as more species disperse into the community, and then to decrease as

competitive interactions remove species from the community. Although rarely

documented, it is also possible that a particularly good disperser will become

dominant and remain dominant (and therefore decreases community diversity) in a

metacommunity solely as a result of its dispersal ability.

We investigated the contribution of dispersal to ecological dominance over the

course of natural microbial succession in pitchers of the carnivorous pitcher plant

Sarracenia purpurea. *S. purpurea* is a perennial plant native to bogs and savannas in

northern and eastern North America (Buckley *et al.* 2003). Each *S. purpurea* plant

produces modified vase-shaped leaves, or pitchers (Figure 1a), annually. At first,

developing pitchers are entirely closed, sterile chambers (Peterson *et al.* 2008). Once

mature, the top portion of each pitcher opens, and open pitchers accumulate rainwater

to form small pools of water (phytotelmata). Potential prey (ants and other small

insects) are attracted to pitchers (Bennett & Ellison 2009); some prey fall into pitchers

and drown, and are then shredded, decomposed, and mineralized by a food web of

microorganisms and invertebrates (reviewed in Ellison *et al.* 2003). The pitcher

microbial community includes bacteria, algae, and fungi, including culturable yeasts

(Cochran-Stafira & von Ende 1998). Pitchers are individual communities within a

metacommunity of other pitchers on the same and on different plants.

Figure 1: Example *S. purpurea* plant and study location. (A) One of the study pitcher plants at the edge of a *Sphagnum* island. This photograph was taken early in the growth season and both opened and unopened pitchers are visible. (B) The location of Harvard Pond in Massachusetts, USA. (C) Locations of the 43 pitchers sampled for this study. Each white circle represents one pitcher. Note that some pitchers were close enough to one another that the white circles overlap. Maps were created using ArcMap™ Version 9.2 (ESRI 2006); map data are from the Office of Geographic Information, Commonwealth of Massachusetts Information Technology Division (2000) and the National Atlas of the United States (2006).

We followed fungal succession within individual pitchers in a *Sphagnum* bog

in central Massachusetts, where the pitcher growing season lasts for about three

months (although pitchers persist for longer and can overwinter, Judd 1959). We first

documented that a single yeast species, *Candida pseudoglaebosa*, was numerically

dominant in the metacommunity throughout the growing season, and its frequency

increased between early and later-successional stages of individual pitchers. We next

investigated the ecological processes leading to *C. pseudoglaebosa* dominance.

Unlike dominant yeasts in many other systems, *C. pseudoglaebosa* was not an

especially good competitor against other tested yeasts. However, it was an especially

good disperser. It is one of the first fungi to arrive in pitchers and it maintained high

frequency in the pitcher plant metacommunity over the course of the season, even as

its frequency within individual pitchers ranged from completely absent to over 90% of

sequences. The apparently superior dispersal ability of *C. pseudoglaebosa* leads it to

dominate the pitcher fungal metacommunity, demonstrating that dispersal ability, like

competitive ability, is an important contributor to ecological dominance in microbial

communities.

Results

Fungal communities in pitchers change over successional time

To understand how fungal communities develop in pitchers, we first

sequenced entire fungal communities in seventeen pitchers over the course of a

growing season. Before selecting pitchers to track for sequencing, we identified 43

unopened pitchers on *Sphagnum* islands in Harvard Pond, located in Petersham,

Massachusetts (Figure 1). We recorded the opening date of each pitcher and collected

water from each pitcher at four days, seven days ("one week"), 34-42 days ("one

month"), and 66-74 days ("two months") after opening. At the two month time point,

insect herbivores, including moth larvae (Atwater *et al.* 2006), had destroyed ten of

the original 43 pitchers, and we could only sample water from 33 pitchers. In the

sampled pitchers, the presence of fungal DNA was assayed using the ITS1F/ITS4

primer pair (Gardes & Bruns 1993, White *et al.* 1990) Fungi were detectable starting

from the first measured timepoint at four days (in 33% of sampled plants), and were

widespread after one week, one month, and two months (in 91%, 95%, and 73% of

sampled plants, respectively). Seventeen of the pitchers contained detectable fungal

DNA at every time point from one week to two months. We sequenced fungal DNA

from these seventeen pitchers at all available time points, including four days if

available, using 454 sequencing of PCR amplicons of the internal transcribed spacer

(ITS) ribosomal region.

Fungal succession varied among pitchers. While community composition

changed significantly with time (Figure 2A), only a small amount of variation in

Figure 2: Non-metric Multidimensional Scaling (NMDS) plots of pitcher plant community similarities.
Similarities of community OTU compositions were calculated using the Jaccard metric (Jaccard 1912). Purple triangles represent four-day-old communities, orange circles represent week-old communities, red squares represent approximately one-month old communities, and blue diamonds represent approximately two-month-old communities. (A) NMDS plot with similarities among time points highlighted. Ellipses depict 95% confidence intervals of the centroid of each time point. No ellipse is depicted for four-day-old communities because only two were measured. (B) NMDS plot with individual pitchers highlighted. Lines connect measurements for each pitcher. Fungal communities in pitcher group 1 are represented with black lines and fungal communities in pitcher group 2 are represented with gray lines. All points are located at the same coordinates in (A) and (B).

community composition was due to variation in time (distance-based redundancy

analysis adjusted $R^2 = 0.03$, F = 1.96, df = 1,27, p = 0.011). This small influence of

time on community composition was likely a result of high variation among pitchers.

Succession followed two trajectories—five pitchers (hereafter referred to as "pitcher

group 1") followed a different successional trajectory from the other twelve ("pitcher

group 2")—and there was considerable variation among communities within each

trajectory (Figure 2B). Distance among pitchers did not explain significant variation

in community composition (partial mantel test of community composition on space

controlling for time, mantel statistic r = -0.04, significance = 0.715).

Despite the observed variation in fungal community composition, diversity

decreased, on average, in pitchers between four days and two months (Figure 3). To

determine a sample's diversity, we calculated Hill numbers of orders 0 to 2 (^0D to ^2D)

for each sample after rarefaction to 1140 sequences per sample. Hill numbers of

different orders give community diversity with an emphasis on rare species (low

orders) or common species (high orders) (Hill 1973, Chao *et al.* 2014). ^0D, ^1D, and ^2D

are equal to operational taxonomic unit (OTU) richness, the exponent of Shannon

diversity, and the inverse of Simpson's index, respectively. Diversity as indicated by

all calculated Hill numbers decreased between early and late timepoints: on average,

^{0}D declined significantly from 42.5 within the first week (including four day and one

week timepoints) to 23.1 after two months (t=-3.6, df=27, p=0.001); ^{1}D declined from

14.2 to 5.2 (t=-3.8, df=27, p=0.0008); and ^{2}D declined from 8.9 to 3.2 (t=3.3, df=27,

p=0.003).

C. pseudoglaebosa *is the dominant fungal taxon in pitchers throughout succession*

C. pseudoglaebosa, in the class Saccharomycetes, was the numerically

dominant taxon in the metacommunity, but was not dominant in every pitcher (Figure

4). In the metacommunity, *C. pseudoglaebosa* was more frequent than any other

taxon at every time point and its frequency increased between early- and late-

successional timepoints. Its metacommunity frequency increased from 22% of total

sequences at four days to 58% at two months (Figure 4A). However, its frequency did

not increase over time in every pitcher. Depending on the pitcher, *C.*

pseudoglaebosa's within-pitcher frequency increased over the season, decreased over

the season, peaked midway through the season, or dipped midway through the season

(Figure 4B). We cannot say whether these increases or decreases in *C.*

pseudoglaebosa sequence frequency reflect changes in the total cell numbers because

we did not measure cell numbers or fungal biomass in pitchers. Pitcher group 1 never

contained appreciable *C. pseudoglaebosa*: each pitcher in group 1 contained less than

1.7% *C. pseudoglaebosa* sequences regardless of the sampled time point (Figure 4B).

C. pseudoglaebosa *is not a superior competitor, but has complex interactions with*

other yeasts

Figure 4: Taxon diversity in pitchers over time. Proportions are reported based on non-rarefied OTU assignments. (A) Taxon diversity in the entire bog metacommunity. Colored bars represent proportions of total sequences for each fungal class (or phylum for basal fungal lineages). The hatched area represents total *C. pseudoglaebosa* frequency for each time point. Note that *C. pseudoglaebosa* is in the class Saccharomycetes and represents over 99% of Saccharomycetes sequences at the one and two month time points. (B) *C. pseudoglaebosa* sequence frequency in individual pitcher communities. Data points for communities in the same pitcher are connected with lines. Black lines connect points for pitcher group 1 and gray lines connect points for pitcher group 2. Data for this figure is non-rarefied OTU data; for rarefied data, see Figure 4 Figure Supplement 1.

To better understand how interactions with other yeasts might influence the

frequency of *C. pseudoglaebosa*, we grew *C. pseudoglaebosa* and other potentially

interacting pitcher yeasts in laboratory microcosms. We followed the strategy

advocated by Goldberg and Werner (1983), who suggested determining interacting

species' effects on one another by measuring organism performance as the number of

interacting individuals increases. We inoculated microcosms with all possible pairs of

three culturable pitcher yeasts (*C. pseudoglaebosa*, *Rhodotorula babjevae*, and

Moesziomyces aphidis). *C. pseudoglaebosa* represented 41%, *R. babjevae* represented

2%, and *M. aphidis* represented 0.06% of total sequences in the sequencing dataset.

Each low nutrient microcosm contained a focal yeast, which was inoculated as a fixed

number of cells, and an interactor, which was inoculated as a varying number of cells.

We then let the pairs of yeasts grow in the microcosms and investigated the effects of

interactors on each focal yeast using regressions. We evaluated interaction qualities

based on the direction (increasing or decreasing focal yeast yield with more

interacting cells) and shape (linear or polynomial) of each regression, and we

evaluated differences between interactor yeasts based on whether adding interactor

yeast identity to each regression improved its fit.

Interactions between pitcher plant yeasts ranged from facilitation to

competition, depending on the identities of the yeasts and the number of interactor

cells present. Under microcosm conditions, interactions between focal yeasts and

interactors were polynomial when the focal yeast was *C. pseudoglaebosa* or *M.*

aphidis (Figures 5A, B, Figure 5 figure supplemental tables 1-4): both yeasts were

facilitated by small numbers of co-inoculated cells, but their growth was impeded by

larger numbers of co-inoculated cells. Note that we detected facilitation of *C.*

pseudoglaebosa by *R. babjevae* when few *R. babjevae* cells were inoculated, but we

Figure 5: Influence of interacting species on (A) *C. pseudoglaebosa*, **(B)** *M. aphidis*, **and (C)** *R. babjevae*. **The plots depict the yield of each focal species as a function of the number of cells of an interacting species co-inoculated with the focal species. Interacting species are coded by color: red =** *C. pseudoglaebosa*, **yellow =** *M. aphidis*, **and blue =** *R. babjevae*.

did not inoculate *M. aphidis* in small enough numbers to confirm *M. aphidis*

facilitation of *C. pseudoglaebosa* (Figure 5A). At high numbers of co-inoculated cells,

M. aphidis had a more detrimental impact on *C. pseudoglaebosa* than *R. babjevae*

(F=6.79, df=1,50, p = 0.012, Figure 5, figure supplemental table 2). In contrast, the

two interactors of *M. aphidis* had similar effects on its yield: at low and intermediate

inoculum sizes, both *R. babjevae* and *C. pseudoglaebosa* promoted *M. aphidis*

growth, but at high inoculum sizes, both interactors inhibited *M. aphidis* growth

(Figure 5B). Interactions between *R. babjevae* and interactor yeasts were linear

(Figure 5C, figure supplemental tables 5-6): *R. babjevae* yield was impeded by

interactors regardless of the number of interactor cells present, and *C. pseudoglaebosa*

had a more detrimental impact on *R. babjevae* than *M. aphidis* did (F=86.07, df=1,58,

p < 0.001, Figure 5, figure supplemental table 6).

C. pseudoglaebosa *is an early disperser in pitchers*

To investigate whether dispersal influences *C. pseudoglaebosa* dominance in

pitchers, we observed the arrival times each of the three yeasts mentioned above in

pitchers over the *S. purpurea* growth season in Harvard Pond. We surveyed the

presence or absence of each yeast in each of the 43 sampled pitchers using taxon-

specific PCR primers (Table 1) to determine when each yeast arrived in a pitcher and

whether it persisted throughout the season. The three yeasts appeared in pitchers

sequentially (Figure 6): *C. pseudoglaebosa* first arrived in pitchers within four days

after the pitchers opened; *R. babjevae* arrived between four days and one week after

pitchers opened; and *M. aphidis* arrived one week to one month after pitchers opened.

Once a yeast colonized a pitcher, it either persisted in or disappeared from that pitcher

later in the season, but it did not disappear from the broader metacommunity.

Table 1: Taxon-specific PCR primer sequences used to detect pitcher yeasts

yeast	forward sequence	Tm	reverse sequence	Tm	product length
C. glaebosa	5'-CTGCGGAAGGATCATTACAGT-3'	54.6	5'-TGTTCAGACAACACTGTTCA-3'	51.8	466
R. glutinis	5'-AAGTCGTAACAAGGTTTCCG-3'	52.8	5'-CCCAACTCGGCTCTAGTAAA-3'	53.9	527
P. aphidis	5'-GGTAATGCGGTCGTCTAAAA-3'	52.6	5'-CTCTTCCAAAGAAGCGAGG-3'	53.1	467

| Four days | One week | One month | Two months |

= *C. pseudoglaebosa*   = *R. babjevae*   = *M. aphidis*

Figure 6: Presences and absences of each of three yeasts in 43 pitchers over time. Each large gray shape represents the Harvard Pond at one of four time points. Circles represent pitchers: completely white circles represent pitchers in which none of the three yeasts were detected, and circles containing colored pie slices represent pitchers in which one or more of the three assayed yeasts were detected. Pie slices are colored by detected yeast: red = *C. pseudoglaebosa*, yellow = *M. aphidis*, and blue = *R. babjevae*. Circles indicate the approximate locations of pitchers, and are offset to make all data visible; see Figure 1C for accurate pitcher locations.

Discussion

C. pseudoglaebosa *is a pitcher metacommunity dominant*

C. pseudoglaebosa was the dominant fungal taxon throughout succession in

the *S. purpurea* pitcher metacommunity, although it did not dominate every individual

pitcher community. It was the most frequent taxon in the metacommunity at every

sampled timepoint, and its frequency in the metacommunity increased after initial

colonization (Figure 4). Dominance in the pitcher metacommunity was in part a result

of the metacommunity structure itself: while *C. pseudoglaebosa* was not dominant in

every pitcher, it was dominant in enough pitchers (in some cases with a frequency

above 90% of all OTUs) to dominate the metacommunity as a whole (Figure 4). The

poor competitive performance of *C. pseudoglaebosa* relative to other yeasts in

microcosms suggests that overall competitive superiority was not the cause of its

dominance (Figure 5). Instead, *C. pseudoglaebosa*'s early dispersal is a more likely

cause (Figure 6).

Our best explanation for *C. pseudoglaebosa*'s dominance throughout

succession in the pitcher metacommunity is that it pre-empted other community

members by dispersing and establishing in pitchers before other taxa. Early dispersal

likely gave *C. pseudoglaebosa* a numerical advantage by providing the opportunity to

begin exponential growth before other members of the fungal community could arrive

and begin growing (Scheffer *et al.* 2017). Additionally, once *C. pseudoglaebosa* was

established in a pitcher, facilitation by low-frequency interacting taxa may have

helped it maintain dominance (Figure 5A). Once established early in the

metacommunity, *C. pseudoglaebosa* continued to disperse into new pitchers from one

week to one month into the approximately two month long growing season (Figure 6).

Although *C. pseudoglaebosa* was the numerically dominant fungal taxon in

the metacommunity of pitchers, chance events, dispersal, and interactions among

fungi determined whether it was the dominant taxon inside any given pitcher. We

expect that interactions with taxa that arrive, by chance, at different times in different

pitchers caused the variety of *C. pseudoglaebosa* relative frequency changes observed

(*i.e.,* increasing, decreasing, or nonmonotonic, Figure 4B) because interacting yeasts

have qualitatively different impacts on *C. pseudoglaebosa* performance depending on

the number of interacting cells present (Figure 5A). Individual pitchers experienced

priority effects because the timing of taxon arrival in each pitcher (*e.g.,* early arrival

of *C. pseudoglaebosa* in a pitcher) determined later community composition (*e.g., C.*

pseudoglaebosa dominance in the pitcher) (Fukami 2015). Despite the influence of

chance events on *C. pseudoglaebosa* dominance in any given pitcher, the early and

frequent dispersal of *C. pseudoglaebosa* compared to other yeasts enabled its overall

dominance in the metacommunity (Figure 4A, 6).

Ecological patterns and processes during pitcher metacommunity succession

In the metacommunity, and in many individual pitchers, *C. pseudoglaebosa*

remained dominant through decreases in fungal taxon richness and diversity (Figures

3, 4). We did not observe a hump-shaped relationship between pitcher age and species

richness, as previously predicted (Figure 3A) (Moquet *et al.* 2003). Species richness

instead decreased, even as *R. babjevae* and *M. aphidis* were first dispersing into

pitchers late in the season (Figures 3A, 6). It is likely that *C. pseudoglaebosa*

repression of taxa through priority effects has a larger influence on species richness

than increasing diversity through new dispersal does.

Previous studies have also documented biotic and abiotic successional changes

in pitchers; while we did not measure the same parameters as these previous studies,

we assume that similar changes occurred in our pitcher metacommunity and that *C.*

pseudoglaebosa maintained its dominance through these changes. For example,

previous studies documented decreasing pH with increasing pitcher age, an early peak

in prey insect capture during pitchers' life spans (Fish & Hall 1978), and a variety of

changes in bacterial, protist, and invertebrate community compositions over time

(Gray *et al.* 2012, Miller & terHorst 2012). In bacterial, protist, and invertebrate

communities, the identities of dominant taxa changed as succession progressed. In

contrast, *C. pseudoglaebosa* remained the dominant fungus throughout succession. *C.*

pseudoglaebosa appears to be a classical early-successional taxon (Connell & Slatyer

1977, Fierer *et al.* 2010) because it disperses early and frequently (Figure 6), but it is

not replaced by late-successional taxa.

C. pseudoglaebosa dominance throughout succession may be enabled by the

short lifespans of pitchers in Harvard Pond; *i.e.,* pitchers may not live long enough to

enable late-succession fungal taxa to dominate the metacommunity. We sampled

pitchers that were up to 66-74 days old, and stopped sampling at this age because 23%

of pitchers had been destroyed by moths. However, pitchers in northern *S. purpurea*

populations can survive intact through winter conditions (Judd 1959), and pitchers can

be active for over a year in the southern United States (Miller & terHorst 2012). We

speculate that fungal succession more closely resembles classical successional

patterns and the patterns observed for other pitcher guilds (*e.g.,* bacteria,

invertebrates) in longer-lived pitchers. For example, it is possible that a strong

competitor such as *M. aphidis* could replace *C. pseudoglaebosa* in southern *S.*

purpurea metacommunities where pitchers are active for many months. However,

consistent dominance of a single taxon over succession may be common in microbial

habitats that, like northern *S. purpurea* pitchers, have short lifespans but repeatedly

become available.

Conclusions

In the model pitcher plant metacommunity, taxon dispersal ability has a

profound influence on community structure. In particular, *C. pseudoglaebosa*'s ability

to disperse into pitchers before other fungal taxa enables it to persist as the dominant

taxon in the pitcher metacommunity, even as intertaxon interactions and the

stochasticity of individual dispersal events prevent its dominance in every pitcher. It

is likely that dispersal ability leads to persistent dominance in a variety of other

natural succeeding microbial communities and metacommunities, especially when

early dispersal allows a taxon to prevent establishment of other taxa.

Future studies of microbial succession should explicitly include

metacommunity structure when investigating ecological processes. In the pitcher

metacommunity, overall taxon composition changed little over time, with *C.*

pseudoglaebosa dominant throughout succession (Figure 4A). However, individual

pitchers followed a variety of trajectories (Figures 2, 4B, 6). Studies of succession that

do not take a metacommunity's structure into account may miss community

heterogeneity and the diversity of ecological processes, especially dispersal ability, in

play among communities.

Materials and Methods

Study site and field collections

Observations were made on *Sphagnum* islands in Harvard Pond, adjacent to

Tom Swamp, a 50 ha *Sphagnum* bog located in Petersham, Massachusetts at 42°30'N,

72°12'W (Figure 1; Swan & Gill 2007). The *C. pseudoglaebosa* and *M. aphidis*

isolates used in the microcosm study were collected from pitchers in Harvard Pond,

and the *R. babjevae* isolate was collected from a pitcher in Swift River Bog, a 2 ha

kettlehole bog located 75 km south of Tom Swamp in Belchertown, MA at 42°16'N,

72°20'W (Ellison *et al.* 2002). These three yeast isolates were collected in the

summer of 2006 and identified by comparing their ribosomal sequences, amplified

using the ITS1F/ITS4 and LS1/LR5 primer pairs (Gardes & Bruns 1993, White *et al.*

1990, Hausner *et al.* 1993 Vilgalys & Hester 1990), to sequences in the NCBI BLAST

database (Zhang *et al.* 2000). We chose *C. pseudoglaebosa*, *R. babjevae*, and *M.*

aphidis in part because they were all easily cultured from pitchers, and in part because

they formed colonies with different morphologies on agar plates: *C. pseudoglaebosa*

forms smooth white colonies; *M. aphidis* forms wavy white colonies; and *R. babjevae*

forms smooth pink colonies.

All *S. purpurea* pitcher water samples for PCR and 454 sequencing were

collected in the spring and summer of 2009. In May of 2009, we identified 43

unopened *S. purpurea* pitchers on 32 *Sphagnum* islands in Harvard Pond. Pitchers

ranged from less than 1 m to 908 m in distance to other pitchers (Figure 1C). We

visited each pitcher daily until it opened, and counted pitcher age from the date it

opened. For each pitcher water collection, the water inside a pitcher was first mixed

by pipetting up and down with a sterile plastic transfer pipette. We then removed

about 0.25 ml pitcher water and mixed it with 0.25 ml of CTAB buffer (100 mM Tris

pH 8.0, 1.4 M sodium chloride, 20 mM EDTA, 2% CTAB). To the best of our ability,

we avoided collecting insect prey or macrofauna in these samples, although any

protists and microscopic animals present in our samples were included; collected

pitcher water contained no large animal parts and appeared as a cloudy liquid. All

samples were flash-frozen in liquid nitrogen within five hours of collection and stored

at -20 or -80°C before DNA extraction.

PCR assay

We assayed each pitcher water sample for amplifiable DNA from all fungi,

using the ITS1F/ITS4 primer pair, and for each of the three yeasts in the microcosm

experiment, using the primers in Table 1. Primers to selectively amplify portions of

each microcosm yeast's ITS sequence were designed using the NCBI BLAST primer

tool (Rozen & Skaletzky 2000). We chose primer sequences to reliably amplify as

much of the ITS sequence of each yeast species as possible, while not amplifying

other sequences in the BLAST database.

370   To extract DNA from each pitcher water sample before the PCR assay, we

first thawed and centrifuged frozen samples at 16.1 g for 10 min and removed the

supernatant from each pellet. We then suspended each pellet in 200 μL of breaking

buffer (2% Triton X-100, 1% sodium dodecyl sulfate, 100 mM sodium chloride, 10

mM Tris, and 1 mM EDTA) (Hoffman 1997). We mixed each suspension with about

200 μL of 0.5 mm glass beads and 200 μL 25:24:1 chloroform:phenol:isoamyl

alcohol. We vortexed each mixture for 2 min, and then centrifuged it for 5 min at 16.1

g. After centrifugation, we removed the aqueous layer and mixed it with 2.5 volumes

of 95% ethanol and 0.1 volume of 3M sodium acetate (Sambrook & Russell 2001);

we incubated each aqueous layer mixture at -20°C for at least three hours. Next, we

centrifuged each aqueous layer mixture for 15 min at 16.1 g, and removed the

supernatant. Finally, we washed each pellet with 0.5 ml 70% ethanol, centrifuged

each mixture for 10 min at 16.1 g, removed the supernatant, and resuspended each

pellet in 50 μl water.

384   We then assayed each DNA extract for the presence of each Fungal taxon, or

any Fungal DNA in the case of the ITS1F/ITS4 primer pair, using PCR. Each PCR

reaction was composed of 7.9 μL water, 0.1 μL GoTaq® Flexi polymerase (Promega),

5 μL Flexi buffer with green dye added, 5 μL 5x CES (combinatorial PCR enhancer

solution: 2.7 M betaine, 6.7 mM dithiothreitol, 6.7% dimethyl sulfoxide, 55 μg/mL

bovine serum albumin) (Ralser *et al.* 2006), 5 μL nucleotide mix, 2 μL magnesium

chloride, 1 μL of 10 μM of each primer, and 1 μL undiluted template DNA extract.

All reactions were cycled on a Biorad iCycler or myCycler using denaturing,

annealing, and extension temperatures of 95, 55, and 72 °C, respectively. We

denatured for 85 s, then ran 13 cycles of 35 s denaturing, 55 s annealing, and 45 s

extension, followed by 13 cycles that were identical but had a 2 min extension, and

finally 9 cycles with a 3 min extension. We ran a subsequent 10 min extension. Two

μL of each PCR product were visualized on 1% agarose gels stained with SYBR Safe

dye (Invitrogen) and photographed using a U:genius gel documenting system

(Syngene) and a Stratagene transilluminator. Photographs of gels were scored for

presence or absence of a band. Bands that were too faint to reliably score were run a

second time with 6 μL of PCR product per well. Presence of a band on a gel indicated

the presence of detectable fungal or yeast species DNA in a water sample.

To confirm that primers only amplified sequences from the target yeasts, we

randomly selected nine PCR products generated from the *C. pseudoglaebosa* and *R.*

babjevae primer pairs for sequencing. The primer pair that targets *M. aphidis* only

amplified DNA from six pitcher water extracts, and we sequenced all six PCR

products for this primer pair. Sequences were identical to or within one base of

cultured isolate sequences.

Pitcher water fungal DNA amplification and 454 sequencing

We extracted and amplified fungal DNA for fungal community amplicon

sequencing using the protocols described above, with the following changes. Gotaq®

Hotstart polymerase (Promega) was used instead of Flexi polymerase, and we used 50

μM instead of 10 μM of the reverse primer. The forward primer consisted of (in order

from 5' to 3') the 454 "A" primer (CCATCTCATCCCTGCGTGTCTCCGACTCAG)

concatenated with a 10-bp multiplex tag (454 Life Sciences Corporation 2009), and

ITS4; the reverse primer consisted of the 454 "B" primer

(CCTATCCCCTGTGTGCCTTGGCAGTCTCAG) concatenated with ITS1F.

Multiplex tags were unique to each sample. Reactions were cycled at 95 °C for 15

min; 30 cycles of 95 °C for 1 min, 51 °C for 1 min, 72 °C for 1 min; and a final

extension of 72 °C for 8 min.

Products were purified using Agencourt® *AMPure*® XP (Beckman Coulter)

and quantitated using a Qubit® dsDNA HS Assay (Invitrogen) according to the

manufacturers' instructions. We combined equimolar concentrations of the products

of each of three separate PCR reactions from each DNA extract. The sequencing pool

consisted of pooled equimolar concentrations of each pooled PCR product. The pool

was sequenced on one-eighth of a 454 Titanium sequencing run by the Duke Genome

Sequencing & Analysis Core Resource.

454 sequence processing

We processed sequences using QIIME 1.3.0 (Caporaso *et al.* 2010). Low

quality sequences were removed and the remaining sequences were assigned to

mulitplex barcodes using the default quality filtering settings. Primers and barcodes

were trimmed from each sequence and sequences shorter than 200 and longer than

1000 bp were removed from each dataset. Sequences were denoised using the QIIME

denoiser. We reduced chimeric sequences by trimming the 5.8s and ITS1 portions

from all sequences using the Fungal ITS Extractor (Nilsson *et al.* 2010a), and only

analyzing the ITS2 portion. The 5.8s ribosomal region lies between the ITS1 and

ITS2 spacers, and is conserved among fungi relative to the spacers. We expected most

chimeric sequences to form in the 5.8s region and to be composed of ITS1 and ITS2

sequences from different templates (Nilsson *et al.* 2010b). We chose operational

taxonomic units (OTUs) using the uclust method in QIIME, at 97% similarity (Edgar

2010). We discarded all OTUs composed of a single sequence (singleton OTUs)

because we assumed that they resulted from sequencing errors. The longest sequence

in each remaining cluster was retained as a representative sequence. Of the 141,424

total sequences produced, 27,632 were discarded for having lengths less than 200 or

more than 1000 bp and 10,938 were discarded because they had low quality or did not

have a matching barcode. Sixty-six sequences were discarded because the ITS2

subunit could not be extracted and 139 sequences were discarded because they

represented singleton OTUs. In total, we retained 102,649 sequences for further

analysis. Each pitcher water sample produced between 253 and 4365 sequences. We

will deposit the 454 sff files and corresponding FASTA files in the NCBI Small Read

Archive database (accession numbers to be added after manuscript acceptance).

Sequence taxonomy assignments

To assign genus-level taxonomy to OTUs, we performed a MEGAN analysis

of OTU representative sequences on the top ten hits from the NCBI BLAST

nucleotide database extracted using BLAST 2.2.25+ and default MEGAN settings

(Huson *et al*. 2011, Zhang *et al.* 2000). We discarded OTUs matching organisms not

in the kingdom Fungi (plant, animal, and protist sequences), but we assumed that

OTUs with no BLAST matches or matching unassigned fungal environmental

sequences were fungal sequences not yet identified in the NCBI database. We

retained unassigned OTUs for diversity measurements, but did not include them in

taxonomy summaries. We reviewed the taxon assignments output by MEGAN

manually, and filled in higher-level classifications (*e.g.,* order or class) using Index

Fungorum (http://www.indexfungorum.org/) when an OTU was assigned to a genus

but not higher-level classifications. We attempted to assign all taxa to genera; if it was

not possible to assign a taxon to a genus, we assigned it to the most specific

taxonomic group possible. OTUs assigned to the genus *Candida* were further assigned

to the species *C. pseudoglaebosa* if they matched either *C. pseudoglaebosa* or its

close relative *C. glaebosa* in a BLAST search. When we compared these seven OTUs

against sequences representing the type strains of the two *Candida* species (accession

numbers KY102342.1 and KY102112.1, respectively), all seven OTUs aligned to *C.*

pseudoglaebosa better than they did *C. glaebosa*. In total, we detected 553 OTUs, of

which 379 were assigned to fungal taxa; fifteen were discarded because they matched

non-fungal sequences; and 159 were not assigned. Of the 379 fungal taxa in the

temporal data set, 52% were Basidiomycota, 43% Ascomycota, and 5% basal fungal

lineages. Sequences, metadata, and OTU tables including taxonomic assignments in

QIIME 1.3.0 format are included in the supplementary datasets 2-3.

Microcosm interaction assays

Interactions between yeasts were assayed in microcosms designed to mimic

pitchers simultaneously colonized by different numbers of two yeast species. We

grew microcosms in low-nutrient media designed to mimic natural conditions in

pitchers. Microcosms contained sterile yeast extract media (YEM) composed of 1g/L

yeast extract in local tap water (Cambridge, MA, USA). Tap water was used instead

of deionized water because we wanted the media to include micronutrients present in

local rainwater that may be important for pitcher plant yeast growth. The tap water

supply in Cambridge, MA, where this experiment was conducted, comes from three

Massachusetts reservoirs (Waldron & Bent 2001), and we expected it to have similar

inputs as rainwater in Harvard Pond pitchers. Each microcosm was inoculated with a

target yeast species and an interactor in 200 μL of liquid yeast media. Each target

yeast was inoculated with about 1000 cells per microcosm, and each interactor yeast

493   was inoculated at zero, low, medium, and high cell numbers (0 and approximately

494   100, 1000, and 10,000 cells).

495   Eighteen treatments of yeast mixtures were prepared, with ten replicates each,

for a total of 180 microcosms. Before inoculation, yeasts were grown in liquid YEM

for 48 hours. Inoculation sizes were measured after inoculation using counts of

colony-forming units (CFUs) on solid YEM (YEM plus 1.5% agar). Microcosms

were arranged in sterile 96-well polystyrene flat bottom cell culture plates and

incubated between 25 and 27 °C, shaking at 700 rpm for 48 hours. After incubation

32 microcosms were discarded because of suspected contamination. We diluted each

remaining microcosm $1:10^3$ or $1:10^4$ in sterile water, plated it to solid YEM, and

counted CFUs on plates containing at least 30 total CFUs. When no CFUs of an

inoculated yeast were present on a plate, we conservatively assumed that the yeast

was present in the microcosm in numbers just below our detection limit. We

calculated total cell numbers assuming one instead of zero CFUs for these yeasts

absent from plates. CFU counts before and after incubation are included in

supplementary dataset 4.

Statistical analyses

511   OTU datasets rarefied to 1140 sequences were used to produce Non-metric

Multidimensional Scaling (NMDS) plots and to compare community similarities, and

alpha diversity indices among pitchers. Eight samples contained fewer than 1140

sequences and were discarded. Proportions of samples assigned to taxonomic groups

were calculated based on the full non-rarefied dataset. Community similarities over

time were compared using partial distance-based redundancy analysis (db-RDA) of

Jaccard dissimilarity (Jaccard 1912) between each pair of samples with pitcher age as

the explanatory variable, conditioned on pitcher identity. A correlation between

geographic distance and community similarity was conducted using a partial Mantel

test conditioned on pitcher age. Hill numbers of order $q=0$ or 2 (^{q}D) were calculated

as $^qD = \left(\sum_{i=1}^{S} p_i^q\right)^{1/(1-q)}$, where S is the total number of OTUs and p_i is the relative

abundance of OTU i; ^{1}D was calculated as the exponent of Shannon diversity (Hill

1973, Chao $et\ al.$ 2014). Changes in Hill numbers were modeled over time using

repeated-measures linear models controlled for pitcher identity; ^{1}D and ^{2}D were log-

transformed before analyses to homogenize variances among timepoints, and ^{0}D was

not transformed.

We modeled the impact of interactor yeasts on focal yeasts in microcosms

using multiple linear and polynomial regressions. Separate regressions were

conducted for each focal yeast. For each regression, focal yeast yield was the

dependent variable, and both the number of co-inoculated interactor yeast cells and

the identity of the interactor yeast were independent variables. We modeled both

linear and quadratic relationships between the number of co-inoculated interactor

yeast cells and the dependent variable because the relationship did not always appear

linear when plotted. Before constructing the regressions, we square-root-transformed

focal yeast yield to homogenize variances for the focal yeasts $R.\ babjevae$ and $C.$

$pseudoglaebosa$, but left yield untransformed for the focal yeast $M.\ aphidis$. We also

transformed competitor inoculum size by $\log_{10}(x+1)$ because interactor inoculum size

was varied on a log scale in the experiment. When comparing the influences of

competitor species, we randomly assigned treatments with no interacting yeast

inoculum to one of the two interacting yeasts. When selecting the best-fitting

regression model, we first established the best-fitting relationship (linear, quadratic, or

both) between log-transformed interactor inoculum size and focal yeast yield, and

then determined whether adding interactor identity or interactions between interacting

yeast identity and inoculum size to the model improved it. The best-fitting model was

the one with the lowest Akaike Information Criterion (AIC).

All statistical analyses and index calculations were conducted using R version

3.3.1 (R Development Core Team, 2016) and the packages vegan, fields, nlme, and

GUniFrac (Chen 2018, Pinheiro *et al.* 2016, Oksanen *et al.* 2016, Nychka *et al.* 2016)

plots were made using ggplot2 (Wickham 2016)

Acknowledgments

Thank you to Aaron Ellison for help throughout this project, and especially for helpful

comments on this manuscript. We would also like to thank staff and REUs of Harvard

Forest for help in the field; members of the Marx lab for advice on analyzing

competition data; Chris Hittinger for advice on *Candida* taxonomy; and Tadashi

Fukami, and the Pringle lab for helpful comments on the manuscript. We would like

to thank Harvard Forest and the Massachusetts Department of Fish and Wildlife for

fieldwork permissions. Funding was provided by Harvard University and a National

Science Foundation Doctoral Dissertation Improvement Grant (award ID number

0909694).

Figure Legends

Figure 1: Example *S. purpurea* plant and study location. (A) One of the study pitcher

plants at the edge of a *Sphagnum* island. This photograph was taken early in the

growth season and both opened and unopened pitchers are visible. (B) The location of

Harvard Pond in Massachusetts, USA. (C) Locations of the 43 pitchers sampled for

this study. Each white circle represents one pitcher. Note that some pitchers were

close enough to one another that the white circles overlap. Maps were created using

ArcMap™ Version 9.2 (ESRI 2006); map data are from the Office of Geographic

Information, Commonwealth of Massachusetts Information Technology Division

(2000) and the National Atlas of the United States (2006).

Figure 2: Non-metric Multidimensional Scaling (NMDS) plots of pitcher plant

community similarities. Similarities of community OTU compositions were

calculated using the Jaccard metric (Jaccard 1912). Purple triangles represent four-

day-old communities, orange circles represent week-old communities, red squares

represent approximately one-month old communities, and blue diamonds represent

approximately two-month-old communities. (A) NMDS plot with similarities among

time points highlighted. Ellipses depict 95% confidence intervals of the centroid of

each time point. No ellipse is depicted for four-day-old communities because only

two were measured. (B) NMDS plot with individual pitchers highlighted. Lines

connect measurements for each pitcher. Fungal communities in pitcher group 1 are

represented with black lines and fungal communities in pitcher group 2 are

represented with gray lines. All points are located at the same coordinates in (A) and

(B).

Figure 3: Hill numbers of orders A) 0, B) 1, and C) 2 in pitchers over time. Data

points for communities in the same pitcher are connected with lines. Black lines

connect points for pitcher group 1 and gray lines connect points for pitcher group 2.

Figure 4: Taxon diversity in pitchers over time. Proportions are reported based on

non-rarefied OTU assignments. (A) Taxon diversity in the entire bog metacommunity.

Colored bars represent proportions of total sequences for each fungal class (or phylum

for basal fungal lineages). The hatched area represents total *C. pseudoglaebosa*

frequency for each time point. Note that *C. pseudoglaebosa* is in the class

Saccharomycetes and represents over 99% of Saccharomycetes sequences at the one

and two month time points. (B) *C. pseudoglaebosa* sequence frequency in individual

pitcher communities. Data points for communities in the same pitcher are connected

with lines. Black lines connect points for pitcher group 1 and gray lines connect

points for pitcher group 2. Data for this figure is non-rarefied OTU data; for rarefied

data, see Figure 4 Figure Supplement 1.

Figure 4 Figure Supplement 1: Taxon diversity in pitchers over time; rarefied

sequence data. The OTU table was rarefied to 1140 sequences per sample. (A) Taxon

diversity in the entire bog metacommunity. Colored bars represent proportions of total

sequences for each fungal class (or phylum for basal fungal lineages). The hatched

area represents total *C. pseudoglaebosa* frequency for each time point. Note that *C.*

pseudoglaebosa is in the class Saccharomycetes and represents over 99% of

Saccharomycetes sequences at the one and two month time points. (B) *C.*

pseudoglaebosa sequence frequency in individual pitcher communities. Data points

for communities in the same pitcher are connected with lines. Black lines connect

points for pitcher group 1 and gray lines connect points for pitcher group 2. Note that

data for some pitchers are missing at some time points because these samples were

discarded during sample rarefaction.

Figure 5: Influence of interacting species on (A) *C. pseudoglaebosa*, (B) *M. aphidis*,

and (C) *R. babjevae*. The plots depict the yield of each focal species as a function of

the number of cells of an interacting species co-inoculated with the focal species.

Interacting species are coded by color: red = *C. pseudoglaebosa*, yellow = *M. aphidis*,

and blue = *R. babjevae*.

Figure 6: Presences and absences of each of three yeasts in 43 pitchers over time.

Each large gray shape represents the Harvard Pond at one of four time points. Circles

represent pitchers: completely white circles represent pitchers in which none of the

three yeasts were detected, and circles containing colored pie slices represent pitchers

in which one or more of the three assayed yeasts were detected. Pie slices are colored

by detected yeast: red = *C. pseudoglaebosa*, yellow = *M. aphidis*, and blue = *R.*

babjevae. Circles indicate the approximate locations of pitchers, and are offset to

make all data visible; see Figure 1C for accurate pitcher locations.

Citations

454 Life Sciences Corporation (2009) *Using multiplex identifier (MID) adaptors for*
the GS FLX Titanium Chemistry—extended MID set. Genome Sequencer FLX System
Technical Bulletin No. 005-2009. Roche Applied Science, Basel, Switzerland.
636
637   Albergaria H, Francisco D, Gori K, *et al.* (2010) *Saccharomyces cerevisiae* CCMI
638   885 secretes peptides that inhibit the growth of some non-*Saccharomyces* wine-
639   related strains. *Applied Microbiology and Biotechnology* 86:965-972.
640
641   Atwater DZ, Butler JL, Ellison AM (2006) Spatial distribution and impacts of moth
642   herbivory on northern pitcher plants. *Northeastern Naturalist* 13:43-56.
643
644   Bennett KT, Ellison AM (2009) Nectar, not colour, may lure insects to their death.
645   *Biology Letters* 5:469-472.
646
647   Boynton PJ, Greig D (2016) Species richness influences wine ecosystem function
648   through a dominant species. *Fungal Ecology* 22:61-72.
649
650   Buckley HL, Miller TE, Ellison AM, Gotelli NJ (2003) Reverse latitudinal trends in
651   species richness of pitcher-plant food webs. *Ecology Letters* 6:825-829.
652

653 Caporaso JG, Kuczynski J, Stombaugh J *et al.* (2010) QIIME allows analysis of high-
654 throughput community sequencing data. *Nature Methods* 7:355-336.
655
656 Chao A, Gotelli NJ, Hsieh TC, *et al.* (2014) Rarefaction and extrapolation with Hill
657 numbers: A framework for sampling and estimation in species diversity studies.
658 *Ecological Monographs* 84:45-67.
659
660 Chen J (2018) GUniFrac: Generalized UniFrac Distances. R package version 1.1.
661 https://CRAN.R-project.org/package=GUniFrac
662
663 Cochran-Stafira DL, von Ende CN (1998) Integrating bacteria into food webs: Studies
664 with *Sarracenia purpurea* inquilines. *Ecology* 79:880-898.
665
666 Connell JH, Slatyer RO (1977) Mechanisms of succession in natural communities and
667 their role in community stability and organization. *American Naturalist* 111:1119-
668 1144.
669
670 Copeland JK, Yuan L, Layeghifard M, *et al.* (2015) Seasonal community succession
671 of the phyllosphere micriobiome. *Molecular Plant-Microbe Interactions* 28:274-285.
672
673 Cutler NA, Chaput DL, van der Gast CJ (2014) Long-term changes in soil microbial
674 communities during primary succession. *Soil Biology & Biochemistry* 69:359-370.
675
676 Edgar RC (2010) Search and clustering orders of magnitudes faster than BLAST.
677 *Bioinformatics* 26:2460-2461.
678
679 Ellison AM, Farnsworth EH, Gotelli NJ (2002) Ant diversity in pitcher-plant bogs of
680 Massachusetts. *Northeastern Naturalist* 9:267-284.
681
682 Ellison AM, Gotelli NJ, Brewer JS *et al.* (2003) The evolutionary ecology of
683 carnivorous plants. *Advances in Ecological Research* 33:1-74.
684
685 ESRI (Environmental Systems Resource Institute) (2006) *Arcmap 9.2*. ESRI.
686
687 Fierer N, Nemergut D, Knight R, Craine JM (2010) Changes through time: integrating
688 microorganisms into the study of succession. *Research in Microbiology* 161:635-642.
689
690 Findley K, Oh J, Conlan S, *et al.* (2013) Topographic diversity of fungal and bacterial
691 communities in human skin. *Nature* 498:367-370.
692
693 Fish D, Hall DW (1978) Succession and stratification of aquatic insects inhabiting the
694 leaves of the insectivorous pitcher plant, *Sarracenia purpurea. American Midland
695 Naturalist* 99:172-183.
696
697 Fukami T (2015) Historical contingency in community assembly: Integrating niches,
698 species pools, and priority effects. *Annual Review of Ecology, Evolution, and
699 Systematics* 46:1-23.
700

701 Gardes M, Bruns T (1993) ITS primers with enhanced specificity for Basidiomycetes
702 —application to the identification of mycorrhizae and rusts. *Molecular Ecology*
703 2:113-118.
704
705 Goddard MR (2008) Quantifying the complexities of *Saccharomyces cerevisiae*'s
706 ecosystem engineering via fermentation. *Ecology* 89:2077-2082.
707
708 Goldberg DE, Werner PA (1983) Equivalence of competitors in plant communities: A
709 null hypothesis and a field experimental approach. *American Journal of Botany*
710 70:1098-1104.
711
712 Gray SM, Akob DM, Green SJ, Kostka JE (2012) The bacterial composition within
713 the *Sarracenia purpurea* model system: Local scale differences and the relationship
714 with the other members of the food web. *PLOS One* 12:e50969.
715
716 Hausner G, Reid J, Klassen G (1993) On the subdivision of *Ceratocystis* SL based on
717 partial ribosomal DNA-sequences. *Candian Journal of Botany* 71:52-63.
718
719 Hill MO (1973) Diversity and evenness: A unifying notation and its consequences.
720 *Ecology* 54:427-432.
721
722 Hillebrand H, Bennett DM, Cadotte MW (2008) Consequences of dominance: A
723 review of evenness effects on local and regional ecosystem processes. *Ecology*
724 89:1510-1520.
725
726 Hoffman, CS (1997) Preparations of yeast DNA, RNA, and proteins. *Current*
727 *Protocols in Cell Biology Supplement* 39:13.11.1-13.11.4.
728
729 Holyoak M, Leibold MA, Mouquet N, *et al.* (2005) Metacommunities: a framework
730 for large-scale community ecology. In: Holyoak M, Leibold MA, Holt RD (eds).
731 *Metacommunities: spatial dynamics and ecological communities*. The University of
732 Chicago Press. 1-31.
733
734 Huson DH, Mitra S, Ruscheweyh H, *et al.* (2011) Integrative analysis of
735 environmental sequences using MEGAN 4. *Genome Research* 21:1552-1560.
736
737 Jaccard P (2012) The distribution of the flora in the alpine zone. *The New Phytologist*
738 11:37-50.
739
740 Jiao S, Chen W, Wang E, *et al* (2016) Microbial succession in response to pollutants
741 in batch-enrichment culture. *Scientific Reports* 6:21791.
742
743 Judd WW (1959) Studies of the Byron Bog in southwestern Ontario X. Inquilines and
744 victims of the pitcher-plant, *Sarracenia purpurea* L. *The Canadian Entomologist*
745 XCI:171-180.
746
747 Koenig JE, Spor A, Scalfone N, *et al.* (2011) Succession of microbial consortia in the
748 developing infant gut microbiome. *Proceedings of the National Academy of Sciences*
749 108:4578-4585.
750

751 Leibold MA, Holyoak M, Mouquet N *et al.* (2004) The metacommunity concept: a
752 framework for multi-scale community ecology. *Ecology Letters* 7:601-613.
753
754 Logue JB, Mouquet N, Peter H, *et al.* (2011) Empirical approaches to
755 metacommunities: a review and comparison with theory. *Trends in Ecology and*
756 *Evolution* 26:482-491.
757
758 Miller TE, terHorst C (2012) Testing successional hypotheses of the stability,
759 heterogeneity, and diversity in pitcher-plant inquiline communities. *Oecologia*
760 170:243-251.
761
762 Moquet N, Munguia P, Kneitel JM, Miller TE (2003) Community assembly time and
763 the relationship between local and regional species richness. *OIKOS* 103:618-626.
764
765 National Atlas of the United States (2006) *State Boundaries of the United States*.
766 National Atlas of the United States. Available online at http://www.nationalatlas.gov/.
767
768 Nilsson RH, Abarenkov K, Veldre V *et al.* (2010a) An open source chimera checker
769 for the fungal ITS region. *Molecular Ecology Resources* 10:1076-1081.
770
771 Nilsson RH, Veldre V, Hartmann M et al. (2010b) An open source software package
772 for automated extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-
773 throughput community assays and molecular ecology. *Fungal Ecology* 3:284-287.
774
775 Nissen P, Arneborg N (2003) Characterization of early deaths of non-*Saccharomyces*
776 yeasts in mixed cultures with *Saccharomyces cerevisiae*. *Archives of Microbiology*
777 180:257-263.
778
779 Nychka D, Furrer R, Paige J, Sain S (2016) fields: Tools for spatial data. doi:
780 10.5065/D6W957CT, R package version 9.0. www.image.ucar.edu/fields
781
782 Office of Geographic Information (MassGIS), Commonwealth of Massachusetts
783 Information Technology Division (2000) 1:5,000 Black and White Digital Orthophoto
784 Images. http://www.mass.gov/mgis/massgis.htm.
785
786 Oksanen J, Blanchet FG, Friendly M, *et al.* (2016) vegan: Community Ecology
787 Package. R package version 2.4-1. https://CRAN.R-project.org/package=vegan
788
789 Peterson CN, Day S, Wolfe BE *et al.* (2008) A keystone predator controls bacterial
790 diversity in the pitcher-plant *Sarracenia purpurea* microecosystem. *Environmental*
791 *Microbiology* 10:2257-2266.
792
793 Pinheiro J, Bates D, DebRoy S, *et al.* (2016) nlme: Linear and Nonlinear Mixed
794 Effects Models. R package version 3.1-128. http://CRAN.R-
795 project.org/package=nlme
796
797 R Development Core Team (2016) *R: A Language and Environment for Statistical*
798 *Computing*. R Foundation for Statistical Computing: Vienna, Austria.
799

800 Ralser M, Querfurth R, Warnatz *et al.* (2006) An efficient and economic enhancer
801 mix for PCR. *Biochemical and Biophysical Research Comunications* 347:747-751.
802
803 Redford AJ, Fierer N (2009) Bacterial succession on the leaf surface: A novel system
804 for studying successional dynamics. *Microbial Ecology* 58:189-198.
805
806 Rozen S, Skaletsky HJ (2000) In: Misener S, Krawetz SA (eds) *Bioinformatics*
807 *Methods and Protocols: Methods in Molecular Biology.* Humana Press. 365–386.
808
809 Sambrook J, Russell DW (2001) *Molecular Cloning, a laboratory manual.* Cold
810 spring harbor laboratory press. A8.12-A8.15.
811
812 Scheffer M, van Bavel B, van de Leemput IA, van Nes EH (2017) Inequality in nature
813 and society. *Proceedings of the National Academy of Sciences* 114:13154-13157.
814
815 Sferra CO, Hart JL, Howeth JG (2017) Habitat age influences metacommunity
816 assembly and species richness in successional pond ecosystems. *Ecosphere* 8:e01871.
817
818 Swan JMA, Gill AM (1970) The origins, spread, and consolidation of a floating bog
819 in Harvard Pond, Petersham, Massachusetts. *Ecology* 51:829-840.
820
821 Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of
822 enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal*
823 *of Bacteriology* 172:4238-4246.
824
825 Waldron MC, Bent GC (2001) Factors affecting reservoir and stream-water quality in
826 the Cambridge, Massachusetts, drinking-water source area and implications for
827 source-water protection. *U.S. Department of the Interior, U.S. Geological Survey.*
828 Water-Resources Investigations Report 00-4262.
829
830 White TJ, Bruns T, Lee S, Taylor J (1990) In: Innis MA, Gelfand DH, Sninsky JJ,
831 White TJ (eds) *PCR Protocols: a Guide to Methods and Applications.* Academic
832 Press. 315-322.
833
834 Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag.
835
836 Williams KM, Liu P, Fay JC (2015) Evolution of ecological dominance of yeast
837 species in high-sugar environments. *Evolution* 69:2079-2093.
838
839 Wilson MJ, McTammany ME (2016) Spatial scale and dispersal influence
840 metacommunity dynamics of benthic invertebrates in a large river. *Freshwater*
841 *Science* 35:738-747.
842
843 Wolfe BE, Dutton RJ (2015) Fermented foods as experimentally tractable microbial
844 ecosystems. *Cell* 161:49-55.
845
846 Wolfe BE, Button JE, Santarelli M, Dutton RJ (2014) Cheese rind communities
847 provide tractable systems for *in situ* and *in vitro* studies of microbial diversity. *Cell*
848 158:422-433.
849

850 Zhang Z, Schwartz S, Wagner L, Miller W (2000) A greedy algortihm for aligning
851 DNA sequences. *Journal of Computational Biology* 7:203-214.
852
853 Zumsteg A, Luster J, Göransson H, *et al.* (2012) Bacterial, archaeal, and fungal
854 succession in the forfield of a receding glacier. *Environmental Microbiology* 63:552-
855 564.
856