1 **TITLE**
2
3 Expanding Parkinson's disease genetics: novel risk loci, genomic context, causal insights and
4 heritable risk.
5
6 **AUTHORS**
7

Mike A. Nalls[1,2,CA]*, Cornelis Blauwendraat[1]*, Costanza L. Vallerga[3,4]*, Karl Heilbron[5]*, Sara
Bandres-Ciga[1]*, Diana Chang[6]*, Manuela Tan[7], Demis A. Kia[7], Alastair J. Noyce[7,8], Angli Xue[3,4],
Jose Bras[9,10], Emily Young[11], Rainer von Coelln[12], Javier Simón-Sánchez[13,14], Claudia
Schulte[13,14], Manu Sharma[15], Lynne Krohn[16,17], Lasse Pihlstrom[18], Ari Siitonen[19,20], Hirotaka
Iwaki[1,2,21], Hampton Leonard[1,2], Faraz Faghri[1,22], J. Raphael Gibbs[1], Dena G. Hernandez[1], Sonja
W. Scholz[23,24], Juan A. Botia[7,25], Maria Martinez[26], Jean-Christophe Corvol[27], Suzanne Lesage[27],
Joseph Jankovic[11], Lisa M. Shulman[11], The 23andMe Research Team[5], System Genomics of
Parkinson's Disease (SGPD) Consortium, Margaret Sutherland[28], Pentti Tienari[29,30], Kari
Majamaa[19,20], Mathias Toft[18,31], Ole A. Andreassen[32,33], Tushar Bangale[6], Alexis Brice[27], Jian
Yang[3,4], Ziv Gan-Or[16,17,34], Thomas Gasser[13,14], Peter Heutink[13,14], Joshua M Shulman[11,35,36],
Nicolas Wood[7], David A. Hinds[5], John A. Hardy[7], Huw R Morris[37,38], Jacob Gratten[3,4], Peter M.
Visscher[3,4], Robert R. Graham[6], Andrew B. Singleton[1] for the International Parkinson's Disease
Genomics Consortium.
21
22 **AFFILIATIONS**
23
[1] Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health,
Bethesda, MD, 20892 USA
[2] Data Tecnica International, Glen Echo, MD, 20812 USA
[3] Queensland Brain Institute, The University of Queensland, Brisbane, QLD 4072 Australia
[4] Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD 4072
Australia
[5] 23andMe, Inc., Mountain View, California 94041 USA
[6] Department of Human Genetics, Genentech, South San Francisco, 94080, CA, USA
[7] Department of Molecular Neuroscience, UCL Institute of Neurology, London, UK
[8] Preventive Neurology Unit, Wolfson Institute of Preventive Medicine, Queen Mary University of
London, London, UK
[9] Center for Neurodegenerative Science, Van Andel Research Institute, Grand Rapids,
Michigan, USA
[10] Department of Neurodegenerative Diseases, UCL Institute of Neurology, University College
London, London, UK
[11] Department of Neurology, Baylor College of Medicine, Houston, USA
[12] Department of Neurology, University of Maryland School of Medicine, Baltimore, MD, USA
[13] Department for Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research,
University of Tübingen, Tübingen, Germany
[14] German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany.

[15] Centre for Genetic Epidemiology, Institute for Clinical Epidemiology and Applied Biometry,
University of Tubingen, Germany
[16] Department of Human Genetics, McGill University, Montreal, Quebec, Canada
[17] Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada
[18] Department of Neurology, Oslo University Hospital, Oslo, Norway
[19] Institute of Clinical Medicine, Department of Neurology, University of Oulu, Oulu, Finland
[20] Department of Neurology and Medical Research Center, Oulu University Hospital, Oulu,
Finland
[21] The Michael J. Fox Foundation, New York, New York, 10036 USA
[22] Department of Computer Science, University of Illinois Urbana-Champaign, Champaign, IL,
61820, USA
[23] Neurodegenerative Diseases Research Unit, National Institute of Neurological Disorders and
Stroke, National Institutes of Health, Bethesda, MD 20892, USA
[24] Department of Neurology, Johns Hopkins University Medical Center, Baltimore, MD 21287,
USA
[25] Departamento de Ingeniería de la Información y las Comunicaciones, Universidad de Murcia,
Spain
[26] INSERM UMR 1220; and Paul Sabatier University, Toulouse, France
[27] INSERM U1127, CNRS UMR 7225, Sorbonne Université UMR S1127, APHP, Institut du
Cerveau et de la Moelle épinière, ICM, Paris F-75013, France
[28] National Institute on Neurological Diseases and Stroke, National Institutes of Health,
Bethesda, MD 20892 USA
[29] Clinical Neurosciences, Neurology, University of Helsinki, Helsinki, Finland
[30] Helsinki University Hospital, Helsinki, Finland
[31] Institute of Clinical Medicine, University of Oslo, Oslo, Norway
[32] Jebsen Centre for Psychosis Research, University of Oslo, Oslo, Norway
[33] Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway
[34] Department of Neurology & Neurosurgery, McGill University, Montreal, Quebec, Canada
[35] Departments of Molecular & Human Genetics and Neuroscience, Baylor College of Medicine,
Houston, USA
[36] Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston
[37] Department of Clinical Neuroscience, UCL Institute of Neurology, London UK
[38] UCL Movement Disorders Centre, UCL Institute of Neurology, London, UK
*denotes shared first authorship.
[CA]denotes corresponding author, mike[at]datatecnica[dot]com
79
80 Full consortia membership (PubMed indexed) is available in the supplemental materials (Text
81 S1).

82
83
84 **ACKNOWLEDGEMENTS AND FUNDING**
85

87
88 **ABSTRACT**

89
We performed the largest genome-wide association study of PD to date, involving the analysis
of 7.8M SNPs in 37.7K cases, 18.6K UK Biobank proxy-cases, and 1.4M controls. We identified
90 independent genome-wide significant signals across 78 loci, including 38 independent risk
signals in 37 novel loci. These variants explained 26-36% of the heritable risk of PD. Tests of
causality within a Mendelian randomization framework identified putatively causal genes for 70
risk signals. Tissue expression enrichment analysis suggested that signatures of PD loci were
heavily brain-enriched, consistent with specific neuronal cell types being implicated from single
cell expression data. We found significant genetic correlations with brain volumes, smoking
status, and educational attainment. In sum, these data provide the most comprehensive
understanding of the genetic architecture of PD to date by revealing many additional PD risk
loci, providing a biological context for these risk factors, and demonstrating that a considerable
genetic component of this disease remains unidentified.
102
103 **INTRODUCTION**
104
Parkinson's disease (PD) is a neurodegenerative disorder, affecting up to 2% of the population
older than 60 years, an estimated 1 million individuals in the United States alone. PD patients
suffer from a combination of progressive motor and non-motor symptoms that increasingly
impair daily function and quality of life. There are no treatments that delay or alter PD[1]. As the
global population continues to age, the prevalence of PD is projected to double in some age
groups by 2030, creating a substantial burden on healthcare systems.[1,2,3]
111
112 Early investigations into the role of genetic factors in PD focused on the identification of rare
113 mutations underlying familial forms of the disease,[4–6] but over the past decade there has been a
114 growing appreciation for the important contribution of genetics in sporadic disease[7,8]. Genetic
115 studies of sporadic PD have altered the foundational view of disease etiology as much of
116 sporadic disease was formerly thought to be environmental.
117
With this in mind, we executed a series of experiments to further explore the genetics of PD
(summarized in Figure 1). We performed the largest-to-date GWAS for PD, including 7.8M
SNPs, 37.7K cases, 18.6K UK Biobank (UKB) "proxy-cases" and 1.4M controls. We identified
putatively causal genes for PD, providing valuable targets for therapeutic research. We
assessed the function of these putatively causal genes on a larger scale than in previous
studies of PD via Mendelian randomization (MR), expression enrichment, and protein-protein
interaction network analysis [9,10,11]. We estimated PD heritability, developed a polygenic risk
score that predicted a substantial proportion of this heritability, and leveraged these results to
inform future studies of PD genetics. Finally, we identified putative PD biomarkers and risk
factors using genetic correlation and Mendelian randomization.
128
129 **METHODS**
130
131 **See Supplementary Methods**
132

RESULTS

Novel loci and multiple signals in known loci identified

To maximize our power for locus discovery we used a single stage design, meta-analyzing all available GWAS summary statistics. In support of this design, we found strong genetic correlations between GWAS using PD cases ascertained by clinicians compared to 23andMe self-reported cases (rG = 0.85, SE = 0.06) and UKB proxy cases (rG = 0.84, SE = 0.134).

We identified a total of 90 independent genome-wide significant association signals through our meta-analysis and conditional analyses of 37,688 cases, 18,618 UKB proxy-cases and 1,417,791 controls at 7,784,415 SNPs (Figure 2, Table 1, Supplementary Appendices, Table S1, Table S2). Of these, 38 signals are new and more than 1MB from loci described in a previous report by Chang *et al.* 2017 (Table S3).

In an attempt to detect multiple independent signals within loci we implemented conditional and joint analysis (GCTA-COJO, http://cnsgenomics.com/software/gcta/) with a large study-specific reference genotype series, as well as a participant-level conditional analysis using 23andMe data [12]. We considered independent risk signals from conditional analyses to share the same locus if they were within 250kb of each other. We detected 10 loci containing more than one independent risk signal (22 risk SNPs in total across these loci), of which nine had been identified by previous GWAS, including multi-signal loci in the vicinity of *GBA, NUCKS1 / RAB29, GAK / TMEM175, SNCA* and *LRRK2*. The novel multi-signal locus comprised independent risk variants rs2269906 (*UBTF / GRN*) and rs850738 (*FAM171A2*). Detailed summary statistics on all nominated loci can be found in Table S2.

Refining heritability estimates and determining extant genetic risk

To quantify how much of the genetic liability we have explained and what direction to take with future PD GWAS we calculated updated heritability estimates and polygenic risk scores (PRS). Using LD score regression (LDSC) on a meta-analysis of all 11 clinically-ascertained datasets from our GWAS and estimated the liability-scale narrow-sense heritability of PD as 0.22 (95% CI 0.18 - 0.26), only slightly lower than a previous estimate derived using GCTA (0.27, 95% CI 0.17 - 0.38)[10,13,14]. This may be because LDSC is known to be more conservative than GCTA, however, our LDSC heritability estimate does fall within the 95% confidence interval of the GCTA estimate.

Next, we sought to determine the proportion of SNP-based heritability explained by our PD GWAS results using polygenic risk scores (PRSs). We utilized a two-stage design for our PRS analyses, with variant selection and training in the NeuroX-dbGaP dataset (5,851 cases and 5,866 controls) and then validation in the Harvard Biomarker Study (HBS, 527 cases and 472 controls). We focused on the NeuroX-dbGaP and HBS cohort as both of these clinically characterized cohorts were genotyped on the same PD-focused array (NeuroX) and have been used in previous studies of PRSs [8,15–18]. In addition, both of these studies directly genotyped

177 larger effect, rare variants within *LRRK2* (rs34637584, G2019S) and *GBA* (rs76763715, N370S)
178 of great interest in previous PRS analyses.
179
In order to prevent bias, we estimated the effect size of each SNP contributing to the PRS using
a meta-analysis of all PD GWAS datasets except NeuroX-dbGAP and HBS. Using permutation
testing in the NeuroX-dbGAP training cohort, we found that the optimal *P* threshold for variant
inclusion was 1.35E-03, which included 1809 variants. Two PRSs were tested in HBS, one
limited to 88 of the 90 genome-wide significant variants (two variants failed to pass quality
control in the HBS study), and the other incorporating 1805 variants from the training phase
(four variants failed to pass quality control in HBS due to low imputation quality). The 88 variant
PRS had an area under the curve (AUC) of 0.651 (95% CI 0.617 - 0.684), while the 1805 variant
PRS had an AUC of 0.692 (95% CI 0.660 - 0.725). The AUCs from our 88 variant PRS in both
the NeuroX-dbGAP cohort and the HBS cohort were significantly larger than the AUCs in those
same cohorts using a published PRS (Chang *et al.* 2017, AUC = 0.624, P < 0.002 from
DeLong's test). Although the HBS cohort was used to discover the 90 PD GWAS risk variants,
therefore potentially biasing our 88 variant PRS, all 90 variants remained genome-wide
significant in a meta-analysis of all GWAS datasets excluding the HBS study. Extended results
for all included studies at all P-value thresholds can be found in the Supplementary Appendix.
195
196 Using equations from Wray *et al.* 2010 and our current heritability estimates, the 88 variant PRS
197 explained approximately 16% of the genetic liability of PD assuming a global prevalence of
198 0.5%[13,19]. The 1805 variant PRS explained roughly 26% of PD heritability. In a high-risk
199 population with a prevalence of 2%, the 1805 variant PRS explained 36% of PD heritable risk
200 [13,19] (Table S4).
201
202 We then attempted to quantify strata of risk in our more inclusive PRS. Compared to individuals
203 with PRS values in the lowest quartile , the PD odds ratio for individuals with PRS values in the
204 highest quartile was 3.74 (95% CI = 3.35 - 4.18) in the NeuroX-dbGaP cohort and 6.25 (95% CI
205 = 4.26 - 9.28) in the HBS cohort (Table 2, Figure 3, Figure S1).
206
Variants in the range of 5E-08 < P < 1.35E-03 (used in the 1805 variant PRS) were rarer and
had smaller effect estimates than variants reaching genome-wide significance. These sub-
significant variants had a median minor allele frequency of 21.3% and a median effect estimate
(absolute value of the log odds ratio of the SNP parameter from regresion) of 0.047. Genome-
wide significant risk variants were more common with a median minor allele frequency of 25.1%,
and had a median effect estimate of 0.081. We performed power calculations to forecast the
number of additional PD cases needed to achieve genome-wide significance at 80% power for a
variant with a minor allele frequency of 21.3% and an effect estimate of 0.047 [20]. Assuming that
all incoming data is well harmonized with current data and that disease prevalence is 0.5%, we
estimated that we would need a total of ~99K cases, ~2.3 times as many as our current
analysis. Variant discovery at this point will help us work towards the maximum achievable AUC
for a genetic predictor in PD (estimated 85%). Past this point it is possible that effect estimates
get too marginal, variants get too rare and they are no longer useful in predictions or in
estimating heritability [21].

Functional causal inferences via Quantitative Trait Loci (QTL)

There were 305 genes within the 78 GWAS loci. We sought to identify the causal gene in each locus in order to help direct future high-throughput functional studies. Specifically, we used large QTL datasets and summary-data-based Mendelian randomization (SMR) to test whether the expression or methylation of these genes led to a causal change in PD risk (Table 3, Table S5, Table S6)[11]. This method allows for functional inferences between two datasets to be made within a similar framework as a randomized controlled trial, treating the genotype as the randomizing factor.

We used four QTL datasets: a large meta-analysis of mRNA expression across brain tissues, mRNA expression in the substantia nigra, mRNA expression in blood, and methylation in blood [22–26]. Of the 305 genes under linkage disequilibrium (LD) peaks around our risk variants of interest, 237 were possibly associated with at least one QTL and were therefore testable via SMR (Supplementary Methods, Table S6). The expression or methylation of 151 of these 237 genes (63.7%) was significantly linked to a causal change in PD risk.

Of the 90 PD GWAS risk variants, 70 were in loci containing at least one of these putatively causal genes after multiple test correction (Table 3 summarizes top QTL per gene). For 53 out of these 70 PD GWAS hits (75.7%), the gene nearest to the sentinel SNP was a putatively causal gene (Table S2). Most loci tested contained multiple putatively causal genes. Interestingly, the nearest putatively causal gene to the rs850738 / *FAM171A2* GWAS risk signal is *GRN*, a gene known to be associated with frontotemporal dementia (FTD)[27]. Mutations in this gene (*GRN*) have also been shown to be connected with another lysosomal storage disorder, neuronal ceroid lipofuscinosis[28].

Rare coding variant burden analysis

As an orthogonal approach for nominating putatively causal genes, we also carried out rare coding variant burden analyses. While the main GWAS analysis was limited to MAF ≥ 1% (except for known coding risk variants), we carried out rare variant burden analyses in a subset of studies (Supplementary Methods). We performed kernel-based burden tests on the 113 genes in our PD GWAS loci that contained two or more rare coding variants (MAF< 5% or MAF < 1%). After Bonferroni correction for 113 genes, we identified 7 significant putatively causal genes: *LRRK2*, *GBA*, *CATSPER3* (rs11950533/*C5orf24* locus), *LAMB2* (rs12497850/*IP6K2* locus), *LOC442028* (rs2042477/*KCNIP3* locus), *NFKB2* (rs10748818/*GBF1* locus), and *SCARB2* (rs6825004 locus). These results suggest that some of the risk associated with these loci may be due to rare coding variants. The *LRRK2* and *NFKB2* associations at MAF < 1% remained significant after correcting for all ~20,000 genes in the human genome (P = 2.15E-10 and P = 4.02E-07, Table S7, Table S5).

Tissue and cell specific expression enrichment plus protein-protein interactions

In order to better understand the function of the genes highlighted by this study, we tested
whether these genes were enriched in 10,651 biological pathways. We tested for gene
expression and pathway enrichment in PD loci using Functional Mapping and Annotation of
Genome-Wide Association Studies (FUMA) and webgestaltR, respectively [9,29]. We found 10
significantly enriched pathways (false discovery rate [FDR]-adjusted P < 0.05, Table S8),
including four related to vacuolar function and three related to known drug targets (calcium
transporters: ikeda_mir1_targets_dn and ikeda_mir30_targets_up, kinase signaling:
kim_pten_targets_dn). Known pathways of interest relating to lysosomal function, endocytosis,
and dopamine metabolism were significantly enriched when using a more lenient *P* value (FDR-
adjusted *P* < 0.1). At least three candidate genes within novel loci are involved in lysosomal
storage disorder (*GUSB*, *GRN,* and *NEU1)*, a pathway of interest in recent PD research [30].
276

Next, we sought to determine the tissues and cell types most relevant to PD etiology using
FUMA[9]. We tested whether the genes highlighted by our PD GWAS were enriched for
expression in 53 tissues from across the body. We found 13 significant tissues, all of which
were brain-derived (Figure S2A), in contrast to what has been seen in Alzheimer's disease
which shows a strong bias towards blood, spleen, lungs and microglial enrichments [31]. To
further disentangle the enrichment in brain tissues, we tested whether our PD GWAS genes
were enriched for expression in 88 brain cell types using single cell RNA sequencing reference
data from DropViz (http://dropviz.org)[32]. After false discovery rate correction we found seven
significant brain cell types, all of which were neuronal (Figure S2B). The strongest enrichment
was for neurons in the substantia nigra (SN) at P = 1.0E-06, with additional significant results at
P < 5.0E-4 for the globus pallidus (GP), thalamus (TH), posterior cortex (PC), frontal cortex
(FC), hippocampus (HC) and entopeduncular nucleus (ENT).
289

290   Finally, we analyzed protein-protein interaction networks using webgestaltR[29] and found that the
291   genes highlighted by our PD GWAS were enriched in six functional ontological networks (FDR-
292   adjusted *P* < 0.1). The majority of these networks were related to chemical signaling pathways
293   or response to some type of stressor. The most significant protein-protein interaction was
294   related to response to interferon-gamma (Table S9, Figure S3A, Figure S3B).
295

296   **Genetic correlations and Mendelian randomization across phenotypes**
297

Next, we used cross-trait genetic correlation and Mendelian randomization to identify putative
PD biomarkers and risk factors. We estimated the cross-trait genetic correlation between our
PD GWAS and 757 other GWAS datasets curated by LD hub [33]. We found four significant
genetic correlations (FDR-adjusted *P* < 0.05, Table 4, Table S10) including positive correlations
with intracranial volume (rG = 0.351, SE = 0.077, P = 4.64E-06) and putamen volume (rG =
0.248, SE = 0.064, P = 9.55E-05, respectively)[34], and negative correlations with current tobacco
use (rG = -0.134, SE = 0.034, P = 7.92E-05) and "academic qualifications: National Vocational
Qualifications (NVQ) or Higher National Diploma (HND) or Higher National Certificate (HNC) or
equivalent" (rG = -0.169, SE 0.045, P = 2.00E-04)[35]. The negative association with one's
academic qualifications suggests that individuals without a college education may be at less risk

of PD than individuals with higher levels of education. The correlation between PD and smoking
status may not be independent from the correlation between PD and education as smoking
status and years of education were significantly correlated (rG = -0.361, SE = 0.064, P = 1.64E-
08) [36].
312
We used Mendelian randomization to assess whether there was evidence of a causal
relationship between PD and five phenotypes related to academic qualification, smoking, and
brain volumes described above (Figure S4). Cognitive performance had a large, significant
causal effect on PD risk (MR effect = 0.213, SE = 0.041, Bonferroni-adjusted P = 8.00E-07),
while PD risk did not have a significant causal effect on cognitive performance (Bonferroni-
adjusted P = 0.125). Educational attainment also had a significant causal effect on PD risk (MR
effect = 0.162, SE = 0.040, Bonferroni-adjusted P = 2.06E-04), but PD risk also had a weak but
significant causal effect on educational attainment (MR effect = 0.007, SE = 0.002, Bonferroni-
adjusted P = 7.45E-3). There was no significant causal relationship between PD and current
smoking status (forward analysis: MR effect = -0.069, SE = 0.031, Bonferroni-adjusted P =
0.125; reverse analysis: MR effect = 0.004, SE = 0.010, Bonferroni-adjusted P = 1). Smoking
initiation (the act of ever starting smoking) did not have a causal effect on PD risk (MR effect = -
0.063, SE = 0.034, Bonferroni-adjusted P = 0.315), whereas PD had a small, but significantly
positive causal effect on smoking initiation (MR effect = 0.027, SE = 0.006, Bonferroni-adjusted
P = 1.62E-05). Intracranial volume could not be tested because its GWAS did not contain any
genome-wide significant risk variants. There was no significant causal relationship between PD
and putamen volume (P > 0.05 in both the forward and reverse directions).
330
331   **DISCUSSION**
332
Our work marks a significant step forward in our understanding of the genetic architecture of PD
and provides a genetic reference set for the broader research community. We identified 90
independent common genetic risk factors for PD, nearly doubling the number of known PD risk
variants. We re-evaluated the cumulative contribution of genetic risk variants, both genome-
wide significant and not-yet discovered, in order to refine our estimates of heritable Parkinson's
disease risk. We also nominated likely causal genes at each locus for further follow-up using
QTL analyses and rare variant burden analyses. Our work has highlighted the pathways,
tissues, and cell types involved in PD etiology. Finally, we identified intracranial and putaminal
volume as potential PD biomarkers, and cognitive performance as a PD risk factor. Altogether,
the data presented here has significantly expanded the resources available for future
investigations into potential PD interventions.
344
345   Using a PRS constructed from our GWAS results, we were able to explain up to 36% of PD
346   heritability. Power estimates suggest that expansions of case numbers to 99K cases will
347   continue to reveal additional insights into PD genetics. While these yet-to-be defined risk
348   variants will have relatively small effects, cumulatively they will improve our ability to predict PD
349   and will help to further expand our knowledge of the genes and pathways that drive PD risk.
350

Population-wide screening for individuals who are likely to develop PD is currently not feasible
using our 1805 variant PRS alone. There would be roughly 14 false positives per true positive
assuming a prevalence of 0.5%. While large-scale genome sequencing and non-linear machine
learning methods will likely improve these predictive models, we have previously shown that we
will need to incorporate other data sources (*e.g.* smell tests, family history, age, sex) in order to
generate algorithms that are useful for population-wide screening [18].

Evaluating these results in the larger context of pathway, tissue, and cellular functionality
revealed that genes near PD risk variants showed enrichment for expression in the brain,
contrasting with previous work in Alzheimer's disease. Notably, we showed that the expression
enrichment of genes at PD loci occured exclusively in neuronal cell types. We also found that
PD genes were enriched in chemical signaling pathways and pathways involving the response
to a stressor. These observations may be informative for disease modeling efforts, highlighting
the importance of disease modeling in neurons and of incorporating a cellular stress
component. This will help inform and focus stem cell derived therapeutic development efforts
that are currently underway.

Using cross-trait LD score regression, we found four phenotypes that were genetically
correlated with PD. Putamen and intracranial volumes may prove to be valuable PD biomarkers.
Our bi-directional GSMR results suggest a complex etiological connection between smoking
initiation and PD that will require further follow-up and should be viewed with some caution. One
of the implications of this work is that PD trials of nicotine or other smoking-related compound(s)
may be less likely to succeed. The strong causal effect of cognitive performance on PD is
supported by observational studies [37].

While this study marks major progress in assessing genetic risk factors for PD, there remains a
great deal to be done. No defined external validation dataset was used, which may be seen as a
limitation. Simulations have suggested that without replication variants with *P* values between
5E-08 and 5E-9 should be interpreted with greater caution [38,39]. We found 16 risk variants in this
P value range, including two known variants near *WNT3* (proximal to the *MAPT* locus) and
BIN3. To a degree, the fact that we filtered our variants with a secondary random-effects meta-
analysis may make our 90 PD GWAS hits somewhat more robust due to the conservative
nature of random-effects. Secondly, this study focused on PD risk in individuals of European
ancestry. Adding datasets from non-European populations would be helpful to further improve
our granularity in association testing and ability to fine-map loci through integration of more
variable LD signatures while also evaluating population specific associations. Additionally, large
ancestry-specific PD LD reference panels, such as those for Ashkenazi Jewish patients, will
help us further unravel the genetic architecture of loci such as GBA and LRRK2. This may be
particularly crucial at these loci where LD patterns may be quite variable within European
populations, accentuating the possible influence of LD reference series on conditional analyses
in some cases [40]. Finally, our work utilized state-of-the-art QTL datasets to nominate candidate
genes, but many QTL associations are hampered by both small sample size and low *cis*-SNP
density. Larger QTL studies and PD-specific network data from large scale cellular screens
would allow us to build a more robust functional inference framework.

As the field moves forward there are some critical next steps that should be prioritized. First,
allowing researchers to share participant-level data in a secure environment would facilitate
inclusiveness and uniformity in analyses while maintaining the confidentiality of study
participants. Our work suggests that GWASes including up to 99,000 cases will continue to
provide useful biological insights into PD. In addition to studies of the genetics of PD risk,
studies of disease onset, progression, and subtype will be important and will require large series
of well-characterized patients. We also believe that work across diverse populations is
important, not only to be able to best serve these populations but also to aid in fine mapping of
loci. Notably, the use of genome sequencing technologies could further improve discovery by
capturing rare variants and structural variants, but with the caveat that very large samples sizes
will be required. While there is still much left to do, we believe that our current work represents a
significant step forward and that the results and data will serve as a foundational resource for
the community to pursue this next phase of PD research.
409

WORKS CITED

1. Gasser, T. Genetics of parkinson's disease. *Ann. Neurol.* **44**, S53–S57 (1998).

2. Dorsey, E. R. *et al.* Projected number of people with Parkinson disease in the most

populous nations, 2005 through 2030. *Neurology* **68**, 384–386 (2007).

3. Dorsey, E. R., Ray Dorsey, E. & Bloem, B. R. The Parkinson Pandemic—A Call to Action.

JAMA Neurol. **75**, 9 (2018).

4. Polymeropoulos, M. H. *et al.* Mapping of a gene for Parkinson's disease to chromosome

4q21-q23. *Science* **274**, 1197–1199 (1996).

5. Singleton, A. B. *et al.* alpha-Synuclein locus triplication causes Parkinson's disease.

Science **302**, 841 (2003).

6. Parkinson, J. An Essay on the Shaking Palsy. *J. Neuropsychiatr.* **14**, 223–236 (2002).

7. Fung, H.-C. *et al.* Genome-wide genotyping in Parkinson's disease and neurologically

normal controls: first stage analysis and public release of data. *Lancet Neurol.* **5**, 911–916

(2006).

8. Chang, D. *et al.* A meta-analysis of genome-wide association studies identifies 17 new

Parkinson's disease risk loci. *Nat. Genet.* **49**, 1511–1516 (2017).

9. Wang, J., Vasaikar, S., Shi, Z., Greer, M. & Zhang, B. WebGestalt 2017: a more

comprehensive, powerful, flexible and interactive gene set enrichment analysis toolkit.

Nucleic Acids Res. **45**, W130–W137 (2017).

10. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity

in genome-wide association studies. *Nat. Genet.* **47**, 291–295 (2015).

11. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex

trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).

12. Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics

identifies additional variants influencing complex traits. *Nat. Genet.* **44**, 369–75, S1–3

(2012).

13. Keller, M. F. *et al.* Using genome-wide complex trait analysis to quantify 'missing

heritability' in Parkinson's disease. *Hum. Mol. Genet.* **21**, 4996–5009 (2012).

14. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide

complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).

15. Nalls, M. A. *et al.* Large-scale meta-analysis of genome-wide association data identifies six

new risk loci for Parkinson's disease. *Nat. Genet.* **46**, 989–993 (2014).

16. Nalls, M. A. *et al.* NeuroX, a fast and efficient genotyping platform for investigation of

neurodegenerative diseases. *Neurobiol. Aging* **36**, 1605.e7–12 (2015).

17. Nalls, M. A. *et al.* Genetic risk and age in Parkinson's disease: Continuum not stratum.

Mov. Disord. **30**, 850–854 (2015).

18. Nalls, M. A. *et al.* Diagnosis of Parkinson's disease on the basis of clinical and genetic

classification: a population-based modelling study. *Lancet Neurol.* **14**, 1002–1009 (2015).

19. Wray, N. R., Yang, J., Goddard, M. E. & Visscher, P. M. The genetic interpretation of area

under the ROC curve in genomic profiling. *PLoS Genet.* **6**, e1000864 (2010).

20. Skol, A. D., Scott, L. J., Abecasis, G. R. & Boehnke, M. Joint analysis is more efficient than

replication-based analysis for two-stage genome-wide association studies. *Nat. Genet.* **38**,

209–213 (2006).

21. Kim, H., Grueneberg, A., Vazquez, A. I., Hsu, S. & de los Campos, G. Will Big Data Close

the Missing Heritability Gap? *Genetics* **207**, 1135 (2017).

22. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet.* **45**, 580–

585 (2013).

23. Qi, T. *et al.* Identifying gene targets for brain-related traits using transcriptomic and

methylomic data from blood. *Nat. Commun.* **9**, 2282 (2018).

24. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex

trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).

25. Wu, Y. *et al.* Integrative analysis of omics summary data reveals putative mechanisms

underlying complex traits. *Nat. Commun.* **9**, 918 (2018).

26. Võsa, U. *et al.* Unraveling the polygenic architecture of complex traits using blood eQTL

meta-analysis. *bioRxiv* 447367 (2018). doi:10.1101/447367

27. Cruts, M. *et al.* Null mutations in progranulin cause ubiquitin-positive frontotemporal

dementia linked to chromosome 17q21. *Nature* **442**, 920–924 (2006).

28. Smith KR, E. al. Strikingly different clinicopathological phenotypes determined by

progranulin-mutation dosage. - PubMed - NCBI. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/22608501. (Accessed: 10th January 2019)

29. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and

annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).

30. Robak, L. A. *et al.* Excessive burden of lysosomal storage disorder gene variants in

Parkinson's disease. *Brain* **140**, 3191–3203 (2017).

31. Jansen, I. E. *et al.* Genome-wide meta-analysis identifies new loci and functional pathways

influencing Alzheimer's disease risk. *Nat. Genet.* (2019). doi:10.1038/s41588-018-0311-9

32. Saunders A, E. al. Molecular Diversity and Specializations among the Cells of the Adult

Mouse Brain. - PubMed - NCBI. Available at:

https://www.ncbi.nlm.nih.gov/pubmed/30096299. (Accessed: 31st October 2018)

33. LD Hub. Available at: http://ldsc.broadinstitute.org. (Accessed: 26th June 2018)

34. Hibar, D. P. *et al.* Common genetic variants influence human subcortical brain structures.

Nature **520**, 224–229 (2015).

35. Rapid GWAS of thousands of phenotypes for 337,000 samples in the UK Biobank. *Neale*

lab Available at: http://www.nealelab.is/blog/2017/7/19/rapid-gwas-of-thousands-of-

phenotypes-for-337000-samples-in-the-uk-biobank. (Accessed: 24th June 2018)

36. Bulik-Sullivan, B. *et al.* An atlas of genetic correlations across human diseases and traits.

Nat. Genet. **47**, 1236–1241 (2015).

37. Valdés, E. G. *et al.* Occupational complexity and risk of Parkinson's disease. *PLoS One* **9**,

e106676 (2014).

38. Wu, Y., Zheng, Z., Visscher, P. M. & Yang, J. Quantifying the mapping precision of

genome-wide association studies using whole-genome sequencing data. *Genome Biol.* **18**,

86 (2017).

39. Pulit, S. L., de With, S. A. J. & de Bakker, P. I. W. Resetting the bar: Statistical significance

in whole-genome sequencing-based association studies of global populations. *Genet.*

Epidemiol. **41**, 145–151 (2017).

40. Rivas, M. A. *et al.* Insights into the genetic epidemiology of Crohn's and rare diseases in

the Ashkenazi Jewish population. *PLoS Genet.* **14**, e1007329 (2018).

Table 1: Novel loci associated with Parkinson's disease.
Summary statistics for all 90 genome-wide significant PD variants. Columns include single
nucleotide polymorphism ID (SNP), chromosome (CHR), base pair position (BP), nearest gene
annotation for the variant, effect allele designation and frequency, as well as metrics for the
odds ratio (OR), regression coefficient (beta), and standard error of the beta for the SNP from
fixed-effects meta-analysis as well as the index of heterogeneity (I2). We also include four p-
values from: fixed-effects meta-analyses, random-effects meta-analyses, standard conditional
analyses in 23andMe, and a conditional joint analysis approach (COJO).
505
Table 2: Summary of genetic predictive model performance.
These are estimates of performance for predictive models including single study estimates,
estimates from meta-analyses across studies, as well as a two stage design. Here the best P

threshold column denotes the filtering value for SNP inclusion to achieve the maximal pseudo

(Nagelkerke's) R2. The odds ration (OR) colum is the exponent of the regression coefficient

(beta) from logistic regression of the polygenic risk score (PRS) on case status, with the

standard error (SE) representing the precision of these estimates. These same metrics are

derived across array types and datasets using random-effects meta-analyses. The area under

the curve (AUC) is included as the most common metric for predictive model performance. In

the table, * denotes R2 approximation adjusted for an estimated prevalence of 0.5%, equivalent

to roughly half of the unadjusted R2 estimates for the PRS. All calculations and reported

statistics include only the PRS and no other parameters after adjusting for principal components

1-5, age and sex at variant selection in the NeuroX-dbGaP dataset.

Table 3: Summary of significant functional inferences from QTL associations via

Mendelian randomization for nominated genes of interest.

Multi-SNP eQTL Mendelian randomization results focusing only on the most significant

association per nearest genes to PD risk loci after Bonferroni correction. If a locus was

significantly associated with both brain and blood QTLs after multiple test correction, we opted

to show the most signficant brain tissue derived association here after filtering for possible

polygenicity (HEIDI P > 0.01). All tested QTL summary statistics can be found in Supplementary

Table S6. Effect estimates represent the change in PD odds ratio per one standard deviation

increase in gene expression or methylation.

Table 4: Significant cross-trait genetic correlations.

The genetic correlations between PD and four significantly-associated traits from LD Hub. An

extended version of this table included in the Supplementary Materials (Supplementary Table

S10) showing data for all tested correlations. In this table, h2 represents the heritability estimate

of the trait.

FIGURES

Figure 1: Workflow and rationale summary.

This figure describes study design and rationale behind the analyses included in this report.

Figure 2: Manhattan plot.

The nearest gene to each of the 90 significant variants are labeled in green for previously-

identified loci and in blue for novel loci. –log10P values were capped at 40. Variant points are

color coded red and orange, with orange representing significant variants at P 5E-08 and 5E-9

and red representing significant variants at P < 5E-9. The X axis represents the base pair

position of variants from smallest to largest per chromosome (1-22).

Figure 3: Predictive model details.

A. The odds ratio of developing PD for each quartile of polygenic risk score (PRS) compared to

the lowest quartile of genetic risk. B. PRS receiver-operator curves for each array type and

sampling design.

SUPPLEMENTARY MATERIALS

Supplementary Methods: Detailed methods section.

Supplemental Appendix: This appendix is split into four sections detailing: first comparisons of effect estimates across GWAS cohorts (beta~beta plots), second forest plots for each significant variant, thirdly locus plots showing regional GWAS results, and QTL and burden associations for each variant, finally the fourth section including extended PRS results. Beta~beta plots compare the regression coefficients for up to 90 of the significant variants in one study to a meta-analysis of all others via linear regression. Forest plots communicate similar sensitivity analyses, for each of the 92 variants of interest. In the forest plots, box size indicates relative sample size for that study, and the width of the diamond representing the meta-analysis effect estimates indicate the 95% confidence interval. The locus plots are a zoomed-in version of Figure 2 for each of the 90 significant variants. These plots are truncated at a -log10 P value of 50 for display purposes and include the most significant burden test and QTL analysis results per gene denoted by label color-coding in each figure. In each locus plot, R2 is measured in our in-house LD reference dataset and shows the correlation between the most significant local SNP and all other proximal SNPs. Additional detailed PRS results for a subset of cohorts are available in the appendix summarizing PRS estimates at varied P thresholds. Each cohort specific PRS in the appendix is based on meta-analyses excluding that cohort when calculating SNP weights. A smaller table summarizing PRS associations at the P threshold with the highest r^2 is also included. Column headers in the PRS section of the appendix mirror that of Table 2.

Figure S1: The odds ratio of developing PD for each decile of PRS, comparing each decile to all others for all samples in this analysis.

Figure S2: Results of FUMA analysis for tissue and cell type specific expression enrichment. A. Tissue enrichment. B. Cell type-specific enrichment. Red bars indicate levels of significance surpassing multiple test correction.

Figure S3: Panel A: Gene ontology term connectivity within protein-protein networks. This panel shows network of gene ontology (GO) terms from pathway analyses. Most significant GO terms are shown in green. Panel B: Gene level connectivity within protein-protein networks. This panel shows connectivity between genes across enriched pathways.

Figure S4: Comparison of regression coefficients in Mendelian randomization analyses across traits. Each cross represents a SNP, with the dashed lines representing the trend across all variants. Axes position are regression coefficients from GWAS for significant SNPs from either GWAS. Panel A includes results for cognitive performance, panel B includes results for educational attainment, panel C includes results for putamen volume, panel D includes results for smoking initiation and panel E includes results for current smoking status.

Table S1: Descriptive statistics and quality control summaries for meta-analyzed genome-wide association studies. ! denotes age at exam for both cases and controls. $ denotes age at death,

597 onset not available. * based on 599 PD cases and 715 controls. ^ denotes samples checked for
598 overlap across datasets as per Nalls et al. 2014 and Chang et al. 2017, ^^ denotes checked for
599 overlap within IPDGC sample series, ^^^denotes a combination of both workflows for identifying
600 sample overlap.
601
Table S2: Summary statistics for all nominated risk variants, known and novel. For binary
variables, 0 = negative and 1 = positive. Some specific notes include: delineations of all studies,
new studies and previous studies as discussed in the methods section. Betas and standard
errors (StdErr) refer to effect estimates per SNP from logistic regression or fixed-effects meta-
analyses.I2 is the index of heterogeneity. QTL Nominated Gene = genes which represent the
nearest cis-QTL for that locus significant in MR.
608
609 **Table S3:** Comparison with novel results from Chang et al., 2017. This table summarizes
610 linkage disequilibrium estimates between Chang et al., 2017 novel loci and variants passing
611 quality control in this report.
612
613 **Table S4:** Estimates of genetic liability explained in different scenarios.
614 Here we compare how different AUC estimates and prevalence rates change the amount of
615 genetic liability explained by GWAS.
616
617 **Table S5:** SNPs of interest tagging genes for functional inferences and networks analysis.
618 Nominated genes and SNPs for follow-up analyses based on minimum r^2 > 0.5 within +/- 1MB of
619 one of our 90 risk loci.
620
621 **Table S6:** Complete summary statistics for QTL Mendelian randomization. Output from the
622 SMR package for all QTLs of interest. Additional columns include QTL reference dataset,
623 dataset-level Bonferroni corrected P values and a binary indicator if a candidate association
624 passed multiple test correction. All columns prefixed by SMR indicate multi-SNP SMR results.
625
626 **Table S7:** Rare coding variant burden analyses for genes under GWAS peaks. Detailed results
627 of burden tests for genes proximal to risk loci. This includes variant counts, test statistics (rho, q,
628 P, adjusted P) for each gene of interest.
629
630 **Table S8:** FUMA expression pathway enrichment analysis results.
631 Pathway enrichment from collapsed GWAS summary statistics.
632
633 **Table S9:** Protein network analysis for linked genes under association peaks. Gene ontology
634 terms passing false discovery rate adjustment.
635
636 **Table S10:** Bivariate LDscores. Default output from LD Hub. Abbreviations defined in main text
637 and methods section.
638
639 **Text S1:** Authors and affiliations.
640

641   **Text S2:** Acknowledgements and Funding.
642
643
644
645   **AUTHOR CONTRIBUTIONS**
646
647   **Study level analysis**
648   MAN, CB, CLV, KH, SB-C, DC, MT, DK, LR, JS-S, LK, LP, ABS
649
650   **Additional analysis and data management**
651   MAN, CB, SB-C, AJN, AX, JY, JG, PMV, ABS
652
653   **Design and funding**
654   MAN, CB, CLV, KH, SB-C, LP, MS, KM, MT, AB, JY, ZG-O, TG, PH, JMS, NW, DAH, JH, HRM,
655   JG, PMV, RRG, ABS
656
657   **Critical review and writing the manuscript**
658   MAN, CB, CLV, KH, SB-C, DC, MT, DAK, AJN, AX, JB, EY, RvC, JS-S, CS, MS, LK, LP, AS,
659   HI, HL, FF, JRG, DGH, SWS, JAB, MM, J-CC, SL, JJ, LMS, MS, PT, KM, MT, AB, JY, ZG-O,
660   TG, PH, JMS, NW, DAH, JH, HRM, JG, PMV, RRG, ABS
661
662   **DATA ACCESS**
663
GWAS summary statistics for 23andMe datasets (post-Chang and data included in Chang et al.
2017 and Nalls et al. 2014) will be made available through 23andMe to qualified researchers
under an agreement with 23andMe that protects the privacy of the 23andMe participants.
Please visit research.23andme.com/collaborate/#publication for more information and to apply
to access the data. An immediately accessible version of the summary statistics is available
here excluding Nalls et al. 2014, 23andMe post-Chang et al. 2017 and Web-Based Study of
Parkinson's Disease (PDWBS) but including all analyzed SNPs. After applying with 23andMe,
the full summary statistics including all analyzed SNPs and samples in this GWAS meta-
analysis will be accessible to the approved researcher(s). Underlying participant level IPDGC
data is available to potential collaborators, please contact ipdgc.contact@gmail.com.
674

SNP	CHR	BP	Nearest Gene	Effect	Other	Effect allele	OR	Beta	SE	P, fixed-effects	P, COJO	P, conditional	P, random-effects	I2, %
rs76949143	1	161469054	FCGR2A	c	g	0.501	1.07	0.065	0.009	6.10E-12	4.69E-12	1.38E-05	3.71E-05	40.2
rs11950533	1	171719769	VAMP4	t	c	0.195	0.93	-0.070	0.012	4.47E-09	4.45E-09	2.63E-03	1.09E-07	5.1
rs77351827	2	18147848	KCNS3	a	t	0.904	1.12	0.110	0.019	1.27E-08	1.27E-08	3.75E-07	1.27E-08	0
rs34025766	2	96000943	KCNIP3	a	t	0.242	0.94	-0.066	0.012	1.38E-08	1.48E-08	3.49E-05	1.38E-08	0
rs61169879	3	28705690	LINC00693	t	c	0.379	1.07	0.066	0.010	8.09E-12	7.18E-12	8.84E-05	8.09E-12	0
rs55961674	3	122196892	KPNA1	t	c	0.172	1.09	0.086	0.013	9.98E-12	8.30E-12	2.80E-06	9.98E-12	0
rs11578699	3	151108965	MED12L	a	t	0.367	0.94	-0.063	0.010	1.13E-10	1.02E-10	2.66E-04	1.77E-07	10.9
rs4771268	3	161077630	SPTSSB	a	g	0.674	0.94	-0.062	0.010	5.01E-10	4.90E-10	3.51E-04	2.27E-05	24.6
rs2042477	4	17968811	LCORL	a	t	0.159	0.92	-0.084	0.013	2.87E-10	2.82E-10	7.43E-06	2.87E-10	0
rs9261484	4	170583157	CLCN3	a	g	0.326	0.94	-0.064	0.012	2.00E-10	1.77E-10	5.10E-05	2.17E-05	21.3
rs2248244	5	102365794	PAM	c	g	0.703	1.06	0.062	0.010	1.57E-09	1.65E-09	6.00E-03	2.36E-07	7.9
rs12528068	5	134199105	C5orf24	a	c	0.102	0.91	-0.092	0.016	7.16E-09	6.73E-09	5.08E-04	2.68E-08	1.9
rs2904880	6	30108683	TRIM40	t	c	0.245	0.94	-0.064	0.011	1.62E-08	1.43E-08	1.26E-06	1.62E-08	0
rs62333164	6	72487762	RIMS1	t	c	0.284	1.07	0.066	0.010	1.63E-10	1.79E-10	9.80E-06	1.63E-10	0
rs11707416	6	112243291	FYN	a	g	0.805	1.07	0.071	0.012	1.84E-09	1.97E-09	2.61E-05	1.84E-09	0
rs6808178	6	133210361	RPS12	t	c	0.967	0.80	-0.221	0.034	1.04E-10	9.67E-11	1.09E-06	1.04E-10	0
rs7134559	7	66009851	GS1-124K5.11	a	t	0.051	0.87	-0.143	0.025	1.43E-08	1.51E-08	5.47E-09	2.04E-06	12.3
rs12147950	8	130901909	FAM49B	t	c	0.723	0.94	-0.061	0.011	1.81E-08	1.57E-08	6.07E-06	1.81E-08	0
rs11610045	9	34046391	UBAP2	t	c	0.734	0.94	-0.062	0.011	6.58E-09	6.56E-09	2.74E-04	6.58E-09	0
rs1941685	10	104015279	GBF1	a	g	0.851	0.92	-0.079	0.013	1.05E-09	1.23E-09	7.47E-06	1.05E-09	0
rs6658353	11	10558777	RNF141	a	g	0.878	1.09	0.087	0.015	2.12E-09	1.97E-09	2.17E-07	2.12E-09	0
rs8087969	12	46419086	SCAF11	t	c	0.404	0.95	-0.054	0.010	3.96E-08	3.80E-08	1.69E-02	1.84E-05	25.2
rs6500328	12	133063768	FBRSL1	a	g	0.490	1.06	0.060	0.009	1.77E-10	1.62E-10	3.57E-05	8.79E-07	19.5
rs850738	13	49927732	CAB39L	t	c	0.740	1.06	0.062	0.011	1.15E-08	1.11E-08	4.29E-06	2.46E-04	21.4
rs12600861	13	97865021	MBNL2	t	c	0.230	1.07	0.068	0.011	1.45E-09	1.67E-09	1.41E-04	1.45E-09	0
rs2269906	14	37989270	MIPOL1	t	c	0.438	0.95	-0.053	0.010	3.54E-08	3.58E-08	1.06E-03	3.54E-08	0
rs1450522	14	75373034	RPS6KL1	a	c	0.787	1.07	0.071	0.012	1.92E-09	2.08E-09	2.22E-06	8.18E-06	24.8
rs26431	16	28944396	CD19	c	g	0.309	0.94	-0.065	0.011	7.87E-10	8.68E-10	1.39E-05	7.87E-10	0
rs2086641	16	50736656	NOD2	a	g	0.599	1.06	0.059	0.010	1.82E-09	1.53E-09	1.43E-03	1.82E-09	0
rs6476434	17	7355621	CHRNB1	a	c	0.648	0.95	-0.057	0.010	1.01E-08	1.15E-08	5.10E-03	1.01E-08	0
rs9568188	17	42294337	UBTF	a	c	0.653	1.07	0.063	0.010	6.24E-10	8.63E-09	1.17E-05	6.24E-10	0
rs3742785	17	42434630	FAM171A2	a	g	0.606	0.93	-0.071	0.011	1.29E-11	3.55E-10	4.18E-04	2.17E-07	17
rs997368	17	59917366	BRIP1	t	c	0.164	1.09	0.082	0.013	9.28E-10	9.40E-10	9.07E-07	6.21E-06	16.4
rs666463	17	76425480	DNAH17	a	t	0.833	1.08	0.076	0.013	3.20E-09	2.90E-09	1.62E-05	4.17E-04	41
rs10748818	18	31304318	ASXL3	t	g	0.498	1.05	0.053	0.009	1.69E-08	1.61E-08	1.64E-08	1.69E-08	0
rs7938782	18	48683589	MEX3C	t	g	0.550	0.94	-0.058	0.010	1.41E-08	1.46E-08	1.09E-04	1.41E-08	0
rs76116224	20	6006041	CRLS1	t	c	0.128	1.08	0.080	0.014	8.87E-09	7.94E-09	1.84E-05	4.38E-07	11.2
rs75859381	21	38852361	DYRK1A	a	g	0.283	1.07	0.071	0.011	2.74E-11	2.51E-11	6.31E-05	8.78E-06	34.3

Study	Max P threshold	from PRS*	OR	Beta	SE	P	quartile PRS	quartile PRS	N SNPs	N samples	AUC	95% CI (DeLong)	Sensitivity	Specificity	value (PPV)	value (NPV)	accuracy
Training dataset: IPDGC - Neurox	1.35E-03	0.029	1.74	0.553	0.022	8.99E-135	3.74	3.35 - 4.18	1809	11,243	0.640	0.630 - 0.650	0.569	0.632	0.591	0.611	0.601
Test dataset: HBS	4.00E-02	0.054	2.03	0.709	0.072	8.28E-23	6.25	4.26 - 9.28	1805	999	0.692	0.660 - 0.725	0.628	0.686	0.691	0.623	0.657

	Probe	CHR	Probe, BP	Top SNP, BP	Top SNP	N SNPs	QTL reference	Effect	SE	Odds ratio	P	Bonferroni adjusted P
VAMP4	ENSG00000117533	1	171,690,343	#########	rs10913587	98	Võsa et al. 2018 - blood expression	-0.272	0.05	0.762	5.67E-07	1.19E-04
KCNIP3	ENSG00000115041	2	96,007,438	95,989,766	rs3772034	14	Qi et al. 2018 - brain expression	-0.161	0.04	0.851	1.12E-05	1.15E-03
MAP4K4	ENSG00000071054	2	102,410,880	#########	rs6733355	3	Võsa et al. 2018 - blood expression	1.119	0.24	3.063	2.32E-06	4.87E-04
TMEM163	ENSG00000152128	2	135,344,950	#########	rs598668	28	Qi et al. 2018 - brain expression	0.074	0.02	1.077	3.55E-07	3.65E-05
KPNA1	ENSG00000114030	3	122,187,294	#########	rs73190142	110	Võsa et al. 2018 - blood expression	0.310	0.05	1.363	1.56E-06	3.28E-04
GAK	ENSG00000178950	4	884,612	906,131	rs11248057	1	Qi et al. 2018 - brain expression	0.508	0.10	1.663	7.47E-07	7.69E-05
CAMK2D	ENSG00000145349	4	114,527,635	#########	rs115671064	146	Võsa et al. 2018 - blood expression	-0.006	0.05	0.994	5.74E-06	1.21E-03
PAM	ENSG00000145730	5	102,228,247	#########	rs2432162	679	Võsa et al. 2018 - blood expression	-0.031	0.01	0.970	2.08E-06	4.36E-04
LOC100131289	cg21339923	6	27,636,378	27,636,378	rs78149975	2	Qi et al. 2018 - brain methylation	-0.094	0.02	0.911	1.53E-06	3.06E-04
TRIM40	cg01641092	6	30,094,300	30,094,315	rs9261443	8	Qi et al. 2018 - brain methylation	0.072	0.01	1.075	6.15E-06	1.23E-03
HLA-DRB5	cg26036029	6	32,552,443	32,570,311	rs34039593	8	Qi et al. 2018 - brain methylation	-0.153	0.02	0.858	7.53E-10	1.51E-07
GPNMB	ENSG00000136235	7	23,295,156	23,294,668	rs858274	74	Qi et al. 2018 - brain expression	0.090	0.01	1.094	2.73E-21	2.81E-19
CTSB	ENSG00000164733	8	11,713,495	11,699,279	rs4631423	33	Qi et al. 2018 - brain expression	-0.150	0.04	0.861	4.37E-09	4.50E-07
BIN3	ENSG00000147439	8	22,502,296	22,456,517	rs71513892	32	Qi et al. 2018 - brain expression	0.046	0.01	1.047	1.43E-06	1.48E-04
SH3GL2	ENSG00000107295	9	17,688,103	17,684,784	rs10756899	15	Qi et al. 2018 - brain expression	0.252	0.05	1.287	5.83E-08	6.00E-06
ITGA8	ENSG00000077943	10	15,659,036	15,548,925	rs7910668	6	Qi et al. 2018 - brain expression	-0.201	0.05	0.818	6.13E-05	6.32E-03
RNF141	ENSG00000110315	11	10,548,001	10,553,355	rs4910153	120	Võsa et al. 2018 - blood expression	-0.054	0.05	0.947	6.25E-07	1.31E-04
IGSF9B	cg25790212	11	133,800,774	#########	rs11223626	1	Qi et al. 2018 - brain methylation	-0.172	0.04	0.842	3.24E-06	6.48E-04
FBRSL1	cg03621470	12	133,137,479	#########	rs10781619	16	Qi et al. 2018 - brain methylation	-0.057	0.01	0.944	6.35E-05	1.27E-02
CAB39L	ENSG00000102547	13	49,950,524	49,918,175	rs35214871	30	Qi et al. 2018 - brain expression	0.097	0.02	1.102	3.51E-08	3.62E-06
GCH1	ENSG00000131979	14	55,339,148	55,348,837	rs3825611	6	Qi et al. 2018 - brain expression	0.113	0.03	1.120	2.76E-04	2.85E-02
SYT17	ENSG00000103528	16	19,229,472	19,273,554	rs727747	4	Qi et al. 2018 - brain expression	0.177	0.05	1.193	1.54E-04	1.58E-02
SETD1A	ENSG00000099381	16	30,982,526	30,950,352	rs7206511	34	Võsa et al. 2018 - blood expression	-0.710	0.09	0.492	2.75E-13	5.77E-11
CHRNB1	ENSG00000170175	17	7,354,703	7,373,595	rs60488855	18	Qi et al. 2018 - brain expression	0.115	0.03	1.122	1.67E-05	1.72E-03
UBTF	ENSG00000108312	17	42,290,697	42,297,631	rs113844752	34	Võsa et al. 2018 - blood expression	-0.466	0.09	0.628	5.68E-06	1.19E-03
MAPT	ENSG00000186868	17	44,038,724	44,862,347	rs199502	6	Qi et al. 2018 - brain expression	0.265	0.03	1.304	7.13E-24	7.35E-22
WNT3	ENSG00000108379.5	17	44,875,148	44,908,263	rs9904865	2	GTEx v7 - substantia nigra brain expression	-0.082	0.02	0.921	4.01E-06	4.81E-05
DNAH17	cg09006072	17	76,425,972	76,427,732	rs589582	3	Qi et al. 2018 - brain methylation	0.100	0.02	1.106	2.44E-05	4.88E-03
MEX3C	ENSG00000176624	18	48,722,797	48,731,131	rs12458916	40	Võsa et al. 2018 - blood expression	-0.291	0.05	0.748	5.28E-05	1.11E-02

Trait of interest	PMID	correlation, RG	SE, RG	Z, RG	P, RG	adjusted	Observed H2	SE, H2	H2 intercept	SE, H2 intercept	Cross trait intercept	SE, Cross trait intercept
Intracranial volume	25607358	0.351	0.077	4.580	4.64E-06	3.51E-03	0.166	0.045	1.003	0.007	-0.013	0.005
Current tobacco smoking	Not available (UKB)	-0.134	0.034	-3.947	7.92E-05	2.41E-02	0.055	0.003	1.014	0.010	0.004	0.007
Mean Putamen	25607358	0.248	0.064	3.902	9.55E-05	2.41E-02	0.282	0.047	0.952	0.007	-0.007	0.006
Qualifications: NVQ or HND or HNC or equivalent	Not available (UKB)	-0.169	0.045	-3.726	2.00E-04	3.79E-02	0.015	0.002	1.011	0.007	0.005	0.005

General workflow and rationale

Previously published GWAS

Nalls et al. 2014
13,708 cases
95,282 controls

IPDGC-NeuroX
5,851 cases
5,866 controls

PDWBS
(Chang et al. 2017)
6,476 cases
302,042 controls

+

New and unpublished GWAS

UK Biobank
18,618 proxies
436,419 controls

SGPD
1,169 cases
968 controls

IPDGC
8,036 cases
5,803 controls

Post-Chang, 23andMe
2,448 cases
571,411 controls

=

Risk loci discovery

37.7K cases
18.6K proxies
1.4M controls
7.8M SNPs

90 independent risk signals (38 novel).
305 candidate genes under LD peaks.

➡

Heritability and polygenic risk estimates

LDSC to quantify heritability.

Predictive models using polygenic risk scoring.

Quantifying heritable risk detected here and estimates for future, larger GWAS.

Gene-level analyses

Mendelian randomization of QTLs.

Rare coding variant burden analysis.

Gene set analyses

WebGestaltR to investigate biological pathway enrichment in GWAS loci.

FUMA to investigate tissue and cell type enrichment in GWAS loci.

Shared heritability

LD Hub survey of 757 curated GWAS datasets.

Significant results from LDSC were followed up using Mendelian randomization to infer causal associations.

Fixed effects meta-analysis plus conditional analyses to identify PD risk loci using GWAS data

Additional analyses of GWAS loci investigating genetic risk at various levels from single variants to genes, then pathways and finally across phenotypes