The genetics and genome-wide screening of regrowth loci, a key component of perennialism

in *Zea diploperennis*

Anjun Ma[1,*], Yinjie Qiu[1,*,†], Tajbir Raihan[1], Bimal Paudel[1], Subha Dahal[1,‡], Yongbin Zhuang[1,§],

Aravid Galla[1,¶], Donald Auger[1] & Yang Yen[1,#]

Author Affiliation:

[1]Department of Biology and Microbiology, South Dakota State University, Brookings, SD

57007, USA

*: These authors made the equal contributions to this research

†:Current address: Department of Horticultural Science, University of Minnesota, St. Paul, MN

55108, USA

‡: Current address: Department of Molecular Genetics, University of Toronto, Toronto, ON M5S

1A8, Canada;

§:Current address: College of Agronomy, Shandong Agricultural University, Taian, Shandong,

271018, China

¶: Current address: Department of Entomology, University of Arkansas, Fayetteville, AR 72701,

USA

Corresponding Author: Dr. Yang Yen, SNP 249C, Box 2140D, Brookings, SD 57007, USA;

+1-605-688-4567; yang.yen@sdstate.edu

Classification:

Major: Biological Science

Minor: Plant Biology

Short Title: Regrowth loci in *Zea diploperennis*

Key Words: Perennial maize, teosinte, *Zea mays*, corn, genetics, perenniality, GBS

Abstract

Perennialism is common among the higher plants, yet little is known about its

inheritance. To address this, six hybrids were made by reciprocally crossing perennial *Zea*

diploperennis Iltis, Doebley & R. Guzman with inbred lines B73 and Mo17 and Rhee Flint, a

heirloom variety, of Z. *mays* L. ssp. *mays*. All the F_1 plants demonstrated several cycles of

growth, flowering, senescence and regrowth into normal flowering plants, indicating a dominant

effect of the Z. *diploperennis* alleles. The regrowability (i.e. the plants' ability to regrow after

senescence) was stably transmitted to progeny of the hybrids, so we focused on this trait.

Segregation ratios in the F_2 generations are consistent with the trait controlled by two dominant,

complementary loci, but do not exclude the influence of other modifiers or environment.

Genome-wide screening with genotyping-by-sequencing (GBS) indicated two major regrowth

loci, *regrowth 1* and *regrowth 2*, were on chromosomes 2 and 7, respectively. These findings lay

the foundation for further exploration of the molecular mechanism of regrowth in Z.

diploperennis.

Significance Statement: This study contributes to the general understanding of inheritance

of perennialism in the higher plants. Previous genetic studies of the perennialism in *Zea* have

yielded contradictory results. We take a reductionist approach by specifically focusing on one

trait, the plant's ability to restart a new life cycle after senescence on the same body. While traits,

such as rhizome formation, tillering and dormancy may be important to converting Z. *mays* to

becoming truly perennial, understanding the conditions for regrowth after senescence will be

substantial first step. Importantly, our data indicate that there is no major barrier to transferring

this trait into maize or other grass crops for perennial crop development with proper technology,

which enhances sustainability of grain crop production in an environmentally friendly way.

Introduction

Perennialism is the phenomenon that a plant can live for more than two years; the ability

of doing so is termed perenniality. Plants typically have a life cycle of growth, reproduction

(sexual and/or vegetative) and senescence. Annuals and biennials have only one such cycle in

their life, leaving behind seeds, bulbs, tubers, etc. to initiate another life cycle. Perennials

maintain juvenile meristematic tissues capable of regrowth after senescence to start a new life

cycle on the same body. How perennials do so remains as a mystery. Subterranean stems (such

as rhizomes), polycarpy and tuberous roots are often cited as the means by which plants achieve

perenniality. However, none of these traits is absolutely required by perennials. For instance,

bamboos are essentially monocarpic perennial that regrow from rhizomes. Many perennial

temperate grasses, such as switchgrass[1], cordgrass[2] and eastern gamagrass[3], regrow from the

crowns instead of rhizomes. On the other hand, some annual/biennial plants, such as radish

(*Raphanus sativus*), grow tuberous roots.

Although perennialism is common among higher plants, the study of its genetics and

molecular biology is sporadic. So far, the only published research in molecular mechanism of

plant perennialism was conducted in Arabidopsis. Melzer et al. successfully mutated this annual

herb to show some perennial habits, such as increased woody fiber in the stem, by down-

regulating two flowering genes coding for MADS-box proteins, SUPPRESSOR OF

OVEREXPRESSION OF CONSTANT 1 and FRUITFUL[4]. Unfortunately, this woody mutant

was sterile, and no follow-up research was reported. Perennial-related genes and quantitative loci

(QTL) have been reported in other species. Major QTL controlling rhizome development,

regrowth and tiller number have been mapped on sorghum linkage groups C (chromosome 1)

and D (chromosome 4)[5,6], which are homoeologous to regions of maize chromosomes 1, 4, 5 and

9, respectively[7]. Hu at al. mapped two dominant, complementary QTL *Rhz2* (*Rhizomatousness 2*)

and *Rhz3* that control rhizome production on rice chromosomes 3 and 4 at the loci homoeologous

to the sorghum QTL[6]. Tuberous roots in a wild perennial mungbean (*Vigna radiate* ssp.

sublobata) are conditioned by two dominant, complementary genes[8]. However, after years of

effort these perennialism-related genes have yet to be cloned from any of the species despite that

mapping data and complete genomic sequences of rice and sorghum are readily available.

Therefore, no further research has been reported about these perennialism-related loci/genes.

Recently, Ryder and the associates reported a set of 98 expressed contigs in Johnsongrass (*S.*

halepense) that are likely associated with rhizome development [9].

In the genus *Zea* L., most species, including maize (*Z. mays* ssp. *mays*), are annual.

However, two closely related species, tetraploid *Z. perennis* [Hitchc.] Reeves and Mangelsdorf

and diploid *Z. diploperennis* Iltis, Doebley & R. Guzman, are perennial. Perenniality of these

two teosintes is manifested as regrowth after seed production and senescence, which includes

developing juvenile basal axillary buds and rhizomes, under favorable environment. A fertile F_1

hybrid between *Z. mays* and *Z. perennis* was made by Emerson[10] in the 1920s, and *Z. mays*'

hybrids with *Z. diploperennis* were also obtained soon after the diploid perennial teosinte was

discovered in the 1990s[11]. Evergreen stalks, bulbils (highly-condensed rhizomes), basal shoot

development, stiff stalk and robust root system have all been cited as phenotypic features of

perennialism in *Z. diploperennis*[12-14]. For example, evergreen stalks, which was proposed as a

component of perennialism in Z. *diploperennis*[12], appears to be linked to *sugary 1* on the short

arm of chromosome 4[15].

Conflicting conclusions have been reached in various studies on how perennialism is

inherited in *Zea*. Shaver, who worked with tetraploid *Z. perennis*, proposed that a triple

homozygous recessive genotype is needed for the perenniality in *Zea*[16]. In this model, *pe*

(*perennialism*), interacting with *gt* (*grassy tillers*) and *id* (*indeterminate*), plays a key role in

conferring totipotency to the basal axillary buds and rhizomes in the perennial teosintes[16, 17]. The

nature of *pe* remains unknown and the *Z. perennis*-derived genotype from which *pe* was

identified by Shaver[16] was lost and never recovered despite decades of intensive efforts (Shaver,

personal communication). Mangelsdorf and Dunn mapped *Pe*-d*, the maize allele of the *pe*

homologue in *Z. diploperennis*, to the long arm of maize chromosome 4[18]. The *gt* gene (aka *gt1*),

located on the short arm of maize chromosome 1, encodes a class I homeodomain leucine zipper

that promotes lateral bud dormancy and suppresses elongation of lateral ear branches[15]. It appears

that *gt1* depends on the activity of a major maize domestication gene, *teosinte branched 1* (*tb1*),

and is inducible by shading[19]. The *id* gene (aka *id1*) alters maize's ability to flower[20]. Both *tb1*

and *id1* are located on the long arm of maize chromosome 1 and both encode transcription

factors with zinc finger motifs[19, 21]. Singleton believed that *id1* inhibits plantlet generation at the

upper nodes of a maize stalk[20]. Mangelsdorf and the associates proposed that one or two

dominant genes control annual growth habit in their *Z. diploperennis*-popcorn hybrid[22].

In contrast to the recessive inheritance model, Galinat proposed that perennialism in *Z.*

diploperennis is at least partially controlled by two dominant complementary genes[15]. Also, Ting

and Yu obtained three perennial F₁ hybrids by pollinating three Chinese field corn varieties with

Z. diploperennis[23], which indicate that perennial factors are dominant. Unfortunately, there is no

further report about these hybrids or their derivatives.

Westerbergh and Doebley regarded perennialism in *Z. diploperennis* as a quantitative

trait and identified a total of 38 QTL for eight perennial-habit traits from a *Z. diploperennis* x *Z.*

mays ssp. *parviglumis* (annual) mapping population[12]. Intriguingly, they did not identify any

QTL that shows a singularly large effect. Murray and Jessup indicated that non-senescence and

rhizomatousness are essential traits in their perennial maize breeding practice[24].

Perennialism appears to be a complex trait, strongly influenced by genetic and

environmental factors. A perennial plant in one environment usually cannot survive in another

due to the lack of the required adaptability. For example, *Z. diploperennis*, which is perennial in

the highlands of Mexico, cannot survive the harsh winter in the American Midwest. The various

criteria for what constitutes perennialism in *Zea* may have contributed to contradictory

observations. Traits such as rhizome formation, evergreen stalks, and dormancy are important

adaptive features that support the viability of various perennial plants in various environments. In

this study, we take a reductionist approach and specifically focus on a plant's regrowability (i.e.

the ability to maintain some juvenile meristematic tissues after each life cycle that can initiate a

new life cycle). Although this trait by itself is insufficient for functional perenniality, it appears

to be an essential component of perenniality in *Zea* L. Here we report the results of our genetic

analysis and genome-wide screening of the regrowth trait with genotyping-by-sequencing (GBS)

technology.

Results and Discussion

The production and growth of the hybrids

We made reciprocal crosses of *Z. diploperennis* (Zd, hereafter in a cross combination)

with the following three maize lines: B73, Mo17 and Rhee Flint (RF, hereafter in a cross

combination). B73 and Mo17 are inbred lines and Rhee Flint is an heirloom maize variety. The

first F[1] was made with Rhee Flint in a greenhouse. Rhee flint is small, fast-growing and usually

has tillers, which affords serial plantings with an increased opportunity of a plant simultaneously

flowering with *Z. diploperennis*. Because Rhee Flint is an open-pollinated variety, later F_1s were

made with B73 and Mo17 to facilitate molecular analysis. All the F_1 plants are fertile and

completed multiple cycles of growth, reproduction and senescence (Fig. 1; Supplementary Fig.

S1). Regrowth (as opposed to accidental replanting from seed) of F_1 plants was insured by

inspection that new shoots were attached to the base of the F_1 and confirmed by the

heterozygosity of polymorphic PCR markers (examples shown in Supplementary Fig. S2).

Regrowth of these F_1s originates from basal axillary buds after stem senescence in all the crosses

(Figs. 1D, 1E, 1F).

Some of the basal regrowth immediately developed into a female (Fig. 2A) or a male

(Fig. 2B) inflorescence, or a forest of them (Fig. 2D). These F_1 plants with abnormal regrowth

most often can later undergo normal regrowth in an alternative environment, such as being

moved from the greenhouse to the field, etc., which suggests a strong environmental influence.

No such abnormal growth has been seen in advanced generations. Sometimes, plantlets also can

develop at the upper nodes of some hybrid plants when B13 and Mo17 were used as the parent

(Fig. 2C). The plantlets developed at the upper nodes, however, can only survive if transplanted

into soil. This indicates that the senescent stalks do not function to provide the necessary

nutrients to the plantlets.

Because the F_1 plants and their perennial derivatives are not winter hardy, the

regeneration cycles were alternated between the greenhouse and the field (Supplementary Figs.

S1 & S3). Interestingly, the ears and kernels of the F_1s of the six crosses all were more teosinte-

like (i.e. two rows of oppositely positioned spikelets with paired kernels encased by woody

rachides and glumes) when grown in greenhouse but were more maize-like (i.e. multiple rows of

naked kernels with short soft glumes and rachides around a silica-filled soft core) when grown in

the field (Fig. 3). In the F_2 and higher generations, ear morphology segregated even under

greenhouse condition (Fig. 3). These observations suggest that environmental factors are

important in the preferential expression of the teosinte or the maize alleles of the genes

influencing ear morphogenesis in the hybrids. These observations also indicate that it should be

possible to breed regrowable maize with maize-like ears and kernels.

The contrast between our observations and those of some previous reports is remarkable.

While we focus on a single trait, regrowth after senescence, previous studies were interested in

perennialism generally using varying criteria. Conclusions that perennialism in *Zea* is recessive

might have resulted from the hypothesis that traits such as tiller number at tasseling (TNT) or

rhizome development are indispensable components of perennialism in *Zea*. Indeed, other

studies have used rhizome development as an indicator of perennialism in *Zea*[11, 16, 17, 22, 25] and we

have not observed rhizomes in any of our F_1s or the derived plants. When regrowth occurs, it is

always from an axillary bud. Indeed, it is also our observation that the regrowth of *Z.*

diploperennis is mainly from basal axillary buds, and only occasionally from rhizomes. The *Z.*

perennis - 4X maize F_1s made by R. A. Emerson also were all "weekly perennial" under the

environmental conditions with few or no rhizomes[10].

Other possible explanations for contrasting results is that the perennial teosinte plants

used in those studies were polymorphic for one or more regrowth genes, that the experimental

environments were unfavorable for regrowth to happen, or that some plants needed more time to

break up their dormancy. Shaver[16] and Camara-Hernandez and Mangelsdorf[25] observed that some

of their F_1 plants eventually regrew from basal axillary buds after a period of dormancy. Indeed,

some of our F_2 plants need about two months of dormancy before regrowth. This observation

reinforces the view that even regrowth is a complex trait that is modified by genetics and

environment.

TNT has been associated with perennialism in several studies[14, 17, 26, 27], so we investigated

the relationship of TNT with regrowth in the Zd-RF F_2s. One-way ANOVA of TNT by regrowth

(Supplementary Table S1), however, revealed no significant difference of TNT ($F = 0.897$, $p =$

0.353) between the regrowth and the non-regrowth F_2s. Indeed, we observed regrowth from

several single-stalked hybrid derivatives (Fig. 4A) and non-regrowth of some multi-stalked

plants (Fig. 4B). These results suggest that TNT is not essential to regrowth in *Zea*.

The genetics of regrowth

All our F_1 plants regrew and underwent several life cycles alternating between the

greenhouse and the field. This indicates that, with our materials and in our environment,

regrowth is a dominant trait. Although this contrasts with some reports, regrowable F_1 hybrids of

maize with perennial teosintes were previously obtained by Emerson[9], Shaver[16], Galinat[15], Ting

and Yu[24] and Camara-Hernandez and Mangelsdorf[25]. Brewbaker suggested cytoplasm may

contribute to perennialism[28], but our reciprocal F_1s performed similarly.

To analyze the genetics of regrowth further, 134 B73-Zd F_2s (derived from several F_1

plants where B73 was the female) and 159 Zd-RF F_2s (derived from a single F_1 plant where Zd

was the female) were tested. Among the 134 B73-Zd F_2s, 81 regrew and 53 did not (Table 1).

Similarly, among the 159 Zd-RF F_2s, 90 regrew after senescence and 69 did not (Table 2). One

B73-Zd F_3 population (Supplementary Table S1) and three Zd-RF F_3 populations (Table 2), each

of which was derived from a single regrowth F_2 plant, were also evaluated for their regrowth.

A chi square (χ^2) goodness-of-fit test suggests that both of the F_2 populations and one Zd-

RF F_3 population (Zd-RF F_3-5) we tested best fit a 9:7 regrowth to non-regrowth ratio (Table 3),

and the B73-Zd F_3 population and the remaining two Zd-RF F_3 populations best fit a 3:1 ratio

(Table 3). The simplest model that explains these results is that regrowth in the F_1s and their

derivatives is mainly controlled by two dominant, complementary *regrowth* (*reg*) loci. A two

dominant, complementary gene model parallels what has been found in other species, such as

rice (*Oryza sativa*)[6], Johsongrass (*Sorghum halepense*)[5,6,29], basin wildrye (*Leymus cinereus*)[30] and

wild mungbean (*Vigna radiate* ssp. *sublobata*)[8], for perennialism-related traits.

The Zd-RF F_1 was also backcrossed to each parental line. All plants from the F_1-to-Zd

backcross regrew, showing dominant effect of the Zd alleles. However, only one of the 20 plants

from the F_1-to-RF backcross showed regrowth. Therefore, alternative models, such as one or

three dominant complementary genes, two major genes with a few minor modifiers, or that

regrowth is a complex trait controlled by many QTL, are not eliminated by this genetic analysis,

but are less probable.

The number of regrowth plants observed in any generation might be understated, because

some plants initially recorded as non-regrowth eventually regrew after about two months of

dormancy. Therefore, some plants recorded as non-regrowth and discarded to open up

greenhouse space may have possessed the ability to regrow. Furthermore, transplanting from the

field to the greenhouse and *vise versa* is very stressful so that some regrowable plants may have

been killed this way.

The F_2 and F_3 plants afforded an opportunity to investigate whether some factors

previously implicated in perennialism may contribute to the regrowth trait. The rice

rhizomatousness gene *Rhz2* has been mapped to rice chromosomes 3[6] and sorghum chromosome

1[5,6,29], which are both homoeologous to parts of maize chromosome 1[7]. Additionally, *gt1* and *id1*,

which have been implicated with perenniality in *Zea*[8], and *tb1*, which controls *gt1*[19], are all on

chromosome 1 in Zea[19,21]. Therefore, we investigated the allele compositions of these three genes

in the B73-Zd F_2s (Table 1), and 26 Zd-RF F_2 plants and the three Zd-RF F_3 populations

(Supplementary Table S2), and assayed their association with regrowth. Of the 134 regrowth

hybrid derivatives we examined, 5, 33 and 115 were homozygous for the maize $gt1$, $tb1$ or $id1$

alleles, respectively (Table 1 and Supplementary Table S2). Zd-RF F_3 family Zd-RF F_3-5 is

homozygous for the $gt1$ allele of $Z. diploperennis$ (Supplementary Table S2) but segregates

approximately 9:7 for regrowth and non-regrowth (Table 2). Therefore, our results are

inconsistent with the model of Shaver[16], and show that $gt1$ and $id1$ do not control regrowth in our

F_1s and their derivatives. $Z. diploperennis$'s $gt1$ allele may be helpful to regrowth because the

majority of the plants that regrew had at least one copy, but it is not indispensable because some

plants regrew without it.

Interestingly, we observed no heterozygosity for $id1$ and very low heterozygosity for $tb1$

in all the hybrid derivatives that were examined, regardless of regrowth (Tables 1;

Supplementary Table S2). Of 134 B73-Zd F_2 plants, only 16 had the $Z. diploperennis$ $id1$ allele

(Table 1). Similar phenomena were observed in the derivatives of the Zd-RF cross

(Supplementary Table S2). It seems that the maize chromosome fragment that carries $id1$ was

preferentially transmitted to the hybrid derivatives. Excess homozygosity of the maize $id1$ allele

indicates some sort of selection. It could be that a deficiency or other rearrangement adjacent to

the teosinte $id1$ allele causes it not to transmit efficiently, or it could be that the teosinte $id1$ or

tightly linked allele causes the plant not to grow well or flower in our experimental conditions.

Identifying regrowth loci with genotyping-by-sequencing assay

A genome-wide mining of single nucleotide polymorphisms (SNPs) was conducted in a

randomly selected sub-population of 94 (55 regrowth and 39 non-regrowth) B73-Zd F_2 plants

with GBS technology (Table 1). We conducted the GBS assays to identify QTL for the regrowth

trait. To prepare for these assays, a total of 2,204,834 (85.14%) Illumina sequencing tags that

passed routine quality control filtrations were aligned with the B73_v4 reference genome. A total

of 714,158 SNPs, covering all ten chromosomes with an average of 71,416 SNPs per

chromosome, were then called from 83 (46 regrowth and 37 non-regrowth, labeled in bold in

Table 1) of the 94 F_2 plants using TASSEL 3 pipeline[31, 32]. SNP-calling for the excluded 11 plants

probably failed due to an error in barcode addition before sequencing. These SNPs were first

subjected to a two-step filtration in TASSEL 3 (MergeDuplicateSNPsPlugin and

GBSHapMapFiltersPlugin) to remove SNPs with poor qualities (see the Materials & Methods

section and Supplementary Table S3 for the details), which resulted in 37,925 SNPs (Table 4, 1st

filtration). The 37,925 SNPs were then manually filtered with high missing data rate > 20%,

which resulted in 10,432 remaining SNPs among all ten chromosomes (Table 4, 2nd filtration).

To explore which chromosomal regions may control the regrowth phenotype, the 10,432

SNPs from the 2nd filtration step were pooled for QTL analysis using R/qtl (version 1.42-8)

(Supplementary Table S4). The result is shown in Figure 5A. A total of 126 SNPs (104 real sites

and 22 simulated sites) showed LOD scores higher than 3.00. A permutation test of 1,000 with

the p-value of 0.05 resulted in a significant LOD score of 5.23 (Fig. 5A). This significance

threshold revealed two major QTL with one at 33,041,409 bp (the nucleotide position in the

B73_v4 reference genome sequence) on chromosome 2 with a LOD score of 5.46 and one at

4,284,633 bp on chromosome 7 with a LOD score of 5.53, respectively.

To test if the two strongest QTL correspond to two dominant and complementary loci

suggested by the genetic analysis, we applied a χ^2 test with 9:7 allele segregation ratio model to

the 10,431 SNPs to investigate if the observed and the expected genotypes are significantly

different ($p \leq 0.05$). This is based on our hypothesis that, if a SNP is associated with a *reg* locus,

the teosinte allele of the SNP should be carried by all the regrowable F2s but only by three

sevenths non-regrowable F2s. This step kept 946 SNPs (Table 4, 3rd filtration). Finally, to

simplify the mapping effort, the 946 SNPs were filtered once more by collapsing immediately

neighboring SNPs that share the same haplotypes into one cluster. This step resulted in 597 SNP

cluster with each being represented by the first SNP in the cluster (Table 4, 4th filtration). Locus

mapping was then conducted with a threshold of $\text{LOD}_{95\%} = 4.17$. This analysis revealed two

significant loci with one on chromosome 2 in the interval from 24,244,192 bp to 28,975,747 bp

with the peak at 27,934,739 bp and another on chromosome 7 in the interval from 2,862,253 bp

to 6,681,861 bp with the peak at 5,060,739 bp (Fig. 5B). These two loci were mapped closely to

the two major QTL indicated by the mapping without imputation. These results are consistent

with the two-factor model, which warrants further investigation. To that end we are naming the

factor underlying the chromosome 2 QTL *regrowth 1* (*reg1*) and the factor underlying the

chromosome 7 QTL as *regrowth 2* (*reg2*).

Our LOD analysis located two minor peaks on chromosome 1 that are associated with

regrowth (Fig. 5B). We wanted to know if these two loci are related to *gt1* and *id1*, respectively.

The SNPs at the peak of these loci are at 82,273,951 bp and 177,235,112 bp, far away from *id1*

(around 243,201,405 bp) and *gt1* (around 23,625,801 bp) (Fig. 6). These observations further

indicate that *id1* and *gt1* are not related to regrowth. Also, previous studies reported that *Z.*

diploperennis carried perennialism-related *Pe*-d* and an evergreen gene on chromosome 4[15, 19].

However, our data could not support these observations since no SNP on chromosome 4 is

significantly associated with regrowth (Figs. 5 and 6).

In summary, the results presented here indicate that regrowth in Zea is inherited

dominantly in our experimental conditions. Both the genetic and GBA analyses support a model

where the regrowth trait is mainly controlled by two major regrowth loci, *reg1* and *reg2* on

chromosomes 2 and 7, respectively. Even so, the data do not eliminate more complex models.

Identification and functional study of the candidate genes for *reg1* and *reg2* and their possible

modifiers will initiate an understanding about the molecular mechanism of perenniality in *Zea* L.

We recognize that adaptability is very important for a plant to realize perennialism in a certain

environment. However, this issue can be addressed separately after we understand molecularly

how *Z. diploperennis* is regrowable.

Materials and Methods

<u>Plant materials and phenotyping</u>

Zea diploperennis (PI 462368) and *Z. mays* cv. Rhee Flint (PI 213764) were obtained

from the USDA North Central Region Plant Introduction Station, Ames, IA. B73 and Mo17

inbreds were from the collection of D. Auger and are traceable back to the Maize Genetics

Cooperation Stock Center, Urbana/Champaign, IL. In our designations of F_1s and their

derivatives, the female parental is shown first. All the plants used in this study were grown in the

greenhouse during the winter and in the field during the summer in Brookings, SD. Controlled

pollinations were done by covering tassels and ears with paper bags before and after the

pollination was made. In the greenhouse, plants were maintained with a 16 h-light/8 h-dark cycle

and 20/16 °C day/night temperature except that two-month old *Z. diploperennis* and its hybrid

plants were treated with a 10 h light/14 h dark cycle for four weeks to induce the floral transition.

Plants were scored as regrowth if they produced shoots from the basal axillary buds after

the original stalks finished flowering and senesced. Rhizome and tuber development were

visually investigated on plants that were dug from the soil after senescence. TNT was

investigated by counting numbers of tillers per plant when the tassel had fully emerged. Ear and

kernel morphology was visually examined and photographed.

PCR assay

DNA samples were isolated from young leaves using the CTAB procedure[33] and used for

PCR-based marker assay. Table 5 lists all the PCR primers used in this study. PCR assays were

done using GoTaq Green Master Mix (Catalog# M7505, Promega, Madison, WI) at the

following conditions: 95°C, 35 cycles of 95°C for 45 sec, 55~62°C (primer dependent, see Table

6 for detail) for 1 min and 72°C for 1 min, and 72°C for 10 min. The annealing temperatures

were determined using a 1°C-touchdown PCR step starting from 65°C.

SNP discovery

The GBS assay was conducted according to Elshire and the associates[31]. The preparation

and sequencing of the library were conducted by the University of Wisconsin Biotechnology

Center (UWBRC). Generally, DNA samples were digested with *ApeK*I restriction enzyme (RE),

and unique barcodes were annealed to each DNA fragments. A single-end 100 bp (1x100bp)

sequencing run was carried out on an Illumina HiSeq 2500 platform. The raw data were pooled

as a single fastq file and downloaded from UWBRC.

The TASSEL (Trait Analysis by Association, Evolution and Linkage) 3 pipeline was

used under the guidance of TASSEL manual[32] for the discovery of SNPs between *Z.*

diploperennis and *Z. mays* (Supplementary Table S3). The barcoded sequence reads were

collapsed into a set of unique sequence tags with counts. The tag count files were filtered for a

minimum count threshold and merged into the master tag count file. The B73_RefGen_V4

reference genome sequence was downloaded from MaizeGDB and processed with Bowtie2 for

alignment[34]. Master tags were aligned to the B73 reference genome to generate a "Tags On

Physical Map" (TOPM) file, which contains the genomic position of each tag with the best

unique alignment. The occupancies of tags for each taxon were observed from barcodes

information in the original FASTQ files. Tag counts were stored in a "Tags by Taxa" (TBT) file.

The TOPM and TBT files were used to call SNPs at the tag locations on the genome. The SNPs

were filtered by minimum tag counts of 5, genotype mismatch rate of 0.1, minimum taxon

coverage of 0.01, minimum site coverage of 0.2 and minimum minor allele frequency of 0.01.

Fastq files containing sequences of chromosomes 1 to 10 were merged by FASTX_Toolkit and

indexed. All commands for SNP discovery were executed in Ubuntu 16.04 LTS platform.

SNPs resulted from TASSEL filters plugin with a minimum minor allele frequency of

0.01 were filtered again by removing sites that had missing data in more than 20% of the B73-Zd

F_2 plants. For those SNPs that have missing data in less than 20% of the B73-Zd F_2 plants, the

missing data were imputed by treating them as heterozygote since both two alleles can be

embodied and considered to be moderate. The SNPs from the 2^{nd} filtration identification were

used for QTL mapping. To understand the relationship between the mapped QTL with the

genetic factors revealed by the genetic analysis, the SNPs from the 2^{nd} filtration were further

filtered (the 3^{rd} filtration) with χ^2 ($p < 0.05$). For each regrowth-associated SNP, we expected, in

the regrowable subpopulation, 33.3% plants to carry the homozygous Zd alleles (AA) and 66.6%

to have the Zd-B73 heterozygous allele combination (AB) and none with the B73 homozygous

alleles (BB) and, in the non-regrowable subpopulation, 14% of the plants with AA, 28.6% with

AB and 57.1% with BB. Altogether, a χ^2 contingency table were generated with expected

$\chi^2_{0.05,4} = 9.49$. Any SNPs with $\chi^2 < 9.49$ were kept fitting the 9:7 segregation model. The 4^{th} SNP

filtration was performed to keep the first SNP and remove the rest in a cluster in which all the

SNPs in the range of 100 bp share the same haplotypes. In the locus analysis with the chi-square

imputation, such a cluster of SNPs was treated as one locus. Removing the redundant SNPs

makes the locus analysis more precise because repeated SNP sites would affect the LOD value

and influence the interval estimation.

The SNPs after the 2nd and the 4th filtrations were used for candidate locus/QTL

estimation, respectively. The locus analysis was executed by a standard QTL procedure in *R*

using the *R*/qtl package (version 1.40-8)[35] to better observe the contribution of each SNP and its

neighbors. The *R* codes used for the analyses are listed in Supplementary Table S4. Position

simulation was drawn with a maximum distance of 1.0 cM and an error probability of 1×10^{-4}. The

conditional genotype probability (calc.genoprob), as well as simulated genotypes (sim.geno with

n.draw=32), were calculated. The "haldane" function was used to convert genetic distances into

recombination fractions. Genome scan with a single locus model (scanone) was performed with a

binary model using the expectation-maximization algorithm[35]. A permutation test with 1000

replicates was performed in scanone to visualize the LOD thresholds. We determined a locus

interval by selecting the first and last SNP sites with significant LOD value. Genes within the

intervals were identified by searching the corresponding region on the Gramene website.

<u>Statistical analyses</u>

For statistical analyses, all genotypes and phenotypes were transformed into numeric

values. For phenotypes, the regrowth plants were scored as "1" and the non-regrowth plants were

scored as "2". For genotypes, the plants that were homozygous to the *Z. diploperennis* allele

were scored as "1"; those that were homozygous to the B73 or Rhee Flint allele were scored as

"2"; and those that were heterozygous were scored as "3". When conducting locus analysis,

genotype "1" was transformed to "AA", "2" to "BB" and "3" to "AB".

A chi square goodness-of-fit test was used to find the best-fit model or linkage in the

genetic analysis and reveal candidate loci on chromosomes. To determine if TNT has any

correlation with regrowth, a One-Way ANOVA of TNT by regrowth was performed in JMP

(JMP® 11.2.0).

<u>Sequencing Data availability</u>

All raw fastq data from this study are available at NCBI data deposition site

(https://www.ncbi.nlm.nih.gov/bioproject/) with accession number PRJNA477673.

Acknowledgement   This research was partially supported by funds from USDA-NIFA via

South Dakota Experiment Station and Department of Biology and Microbiology, South Dakota

State University. We greatly appreciate Dr. Frank M. You of Agriculture and Agri-Food Canada

for his help in statistics.

Author Contributions   Y.Y. designed and supervised this project and all the experiments,

and drafted the manuscript; Y.Q., A.M, T.R., B.P., A.G., Y.Z., Y.Y. & D.A. performed the

experiments and collected data; Y.Q., A.M., T.R., D.A. & Y.Y. analyzed the data; all authors

discussed the results and communicated on and approved the final manuscript.

Competing financial interests   The authors declare no competing financial interests.

References

1. Haferkamp MR, Copeland TD (1984) Shoot growth and development of Alamo switchgrass

as influenced by mowing and fertilization. *J Range Management* 37: 406-412.

2. Boe A, Owens V, Gonzalez-Hernandez J, Stein J, Lee DK,Koo BC. (2009) Morphology and

biomass production of prairie cordgrass on marginal lands. *GCB Bioenergy* 1:240-250.

3. Jackson LL, Dewald CL (1994) Predicting evolutionary consequences of greater

reproductive effort in *Tripsacum dactyloides*, a perennial grass. *Ecology* 75:627-641.

4.   Melzer S, Lens F, Gennen J, Vanneste S, Rohde A, Beeckman T (2008) Flowering-time

genes modulate meristem determinacy and growth form in *Arabidopsis thaliana*. *Nat Genet*

40:1489-1492.

5.   Paterson AH, Schertz KF, Lin Y, Liu S, Chang Y (1995) The weediness of wild plants:

Molecular analysis of genes influencing dispersal and persistence of johnsongrass, *Sorghum*

halepense (L.) Pers. *PNAS* 92:6127-6131.

6.   Hu FY, Tao DY, Sacks E, Fu BY, Xu P, Li j, Yang Y, McNally K, Khush GS, Paterson AH,

Li ZK (2003) Convergent evolution of perenniality in rice and sorghum. *PNAS* 100:4050-

4054.

7.   Wei F, Coe E, Nelson W, Bharti AK, Engler F, Butler E, et al. (2007) Physical and genetic

structure of the maize genome reflects its complex evolutionary history. *PLoS Genet* 3: e123.

8.   Nguyen TD, Lawn RJ, Bielig LM (2012) Expression and inheritance of perenniality and

other qualitative traits in hybrids between mungbean cultivars and Australian wild

accessions. *Crop & Pasture Sci* 63:619-634.

9.   Ryder N, Dorn KM, Huitsing M, Adams M, Ploegstra J, DeHaan L, Larson S, Tintle N

(2018) Transcriptome assembly and annotation of johnsongrass (*Sorghum halepense*)

rhizomes identify candidate rhizome-specific genes. *Plant Direct* 2:e00065.

10. Emerson RA, Beadle GW (1930) A Fertile Tetraploid Hybrid Between *Euchlaena perennis*

and *Zea mays*. *Am Nat* 64:190-192.

11. Srinivasan G, Brewbaker, JL (1999) Genetic analysis of hybrids between maize and

perennial teosinte. I. Morphological traits. *Maydica* 44:353-369.

12. Galinat WC (1981a). Evergreen stalks as an indicator of perennialism. *MNL* 55:107.

(personal communication per the policy of MNL)

13. Mangelsdorf PC, Dunn ME (1984a) Various expressions and dominance relations of

perennialism. *MNL* 58:53. (personal communication per the policy of MNL)

14. Westerbergh A, Doebley J (2004) Quantitative trait loci controlling phenotypes related to the

perennial versus annual habit in wild relatives of maize. *Theor Appl Genet* 109:1544-1553.

15. Galinat WC (1981b) The inheritance and linkage of perennialism derived from diploperennis.

MNL 55:107. (personal communication per the policy of MNL)

16. Shaver DL (1967) Perennial maize. *J. Hered* 58:271-273.

17. Shaver DL (1964) Perennialism in *Zea*. *Genetics* 50:393-406.

18. Mangelsdorf PC, Dunn ME (1984b) Linkage relations of Pe*-d. *MNL* 58:54. (personal

communication per the policy of MNL)

19. Whipple CJ, Kebrom TH, Weber AL, Yang F, Hall D, Meeley R, Schmidt R, Doebley J,

Brutnell T, Jackson DP (2011) *grassy tillers1* promotes apical dominance in maize and

responds to shade signals in the grasses. *PNAS* 108:E506-E512.

20. Singleton WR (1946) Inheritance of indeterminate growth in maize. *J Hered* 37:61-64.

21. Colasanti J, Yuan Z, Sundaresan V (1998) The *indeterminate* gene encodes a zinc finger

protein and regulates a leaf-generated signal required for the transition to flowering in maize.

Cell 93:593-603.

22. Mangelsdorf PC, Roberts LM, Rogers S (1981) Crosses of *Zea diploperennis* with corn.

MNL 55:19-21. (personal communication per the policy of MNL)

23. Ting YC, Yu MK (1982) Further studies of F1 hybrids of maize x diploid perennial teosinte.

MNL 56:35-36. (personal communication per the policy of MNL)

24. Murry SC, Jessup RW. 2013. Breeding and genetics of perennial maize: Progress,

opportunities and challenges. In: *Perennial Crops for Food Security, Proceedings of the FAO*

Expert Workshop, August 28-30, 2013, Rome, Italy, pp103-111.

25. Camara-Hernandez J, Mangelsdorf PC (1981) Crosses of *Zea diploperennis* with corn. *MNL*

55:15-17. (personal communication per the policy of MNL)

26. Doebley J, Stec A, Hubbard L (1997) The evolution of apical dominance in maize. *Nature*

386:485-488.

27. Camara-Hernandez J, Mangelsdorf PC (1981) Crosses of *Zea diploperennis* with corn. *MNL*

55:15-17. (personal communication per the policy of MNL)

28. Brewbaker JL (2007) Grassy tiller and sweet corn. *MNL* 81:12. (personal communication per

the policy of MNL)

29. Washburn JD, Murray SC, Burson BL, Klein RR, Jessup R. 2013. Targeted mapping of

quantitative trait locus regions for rhizomatousness in chromosome SBI-01 and analysis of

overwintering in a *Sorghum bicolor* X *S. propinquum* population. *Mol Breed* 31:153-162.

30. Yun L, Larson SR, Mott IW, Jensen KR, Staub JE (2014) Genetic control of rhizomes and

genomic localization of a majoreffect growth habit QTL in perennial wildrye. *Mol Genet*

Genomics 289:383-397.

31. Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011) A

robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS*

ONE 6:e19379.

32. Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, et al. (2014) TASSEL-

GBS: A high capacity genotyping by sequencing analysis pipeline. PLoS ONE 9: e90346.

33. Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13-15.

34. Langmead B, Salzberg S (2012) Fast gapped-read alignment with Bowtie 2. *Nature Methods*

9:357-359.

35. Broman KW, Wu H, Sen Ś, Churchill GA (2003) R/qtl: QTL mapping in experimental

crosses. *Bioinformatics* 19:889-890.

Table legends:

Table 1. The regrowth (R) and the non-regrowth (NR) phenotypes and the marker genotypes of

Zea diploperennis (Zd), *Z. mays* B73 and their 134 F_2 plants*.

Table 2. Segregation of regrowth among the *Zea diploperennis-Z. mays* cv Rhee Flint F_2s and F_3s.

Table 3. Results of the χ^2 goodness-of-fit tests of three genetic models.

Table 4. The actions taken and the numbers of SNPs revealed in each filtration step of SNP

analysis of *Zea mays* B13 - *Z. diploperennis* F_2 plants.

Table 5. PCR primers used in this study.

Figure legends:

Figure 1. Photos of *Zea mays* and *Z. diploperennis* (Zd) F_1 plants. **A**: reciprocal Mo17-Zd (right)

and Zd-Mo17 (left) F_1 plants; **B**: reciprocal B73-Zd (right) and Zd-B73 (left) F_1 plants; **C**: RF-Zd

F_1 plant; **D**: regrowth of a Mo17-Zd F_1 plant; **E**: regrowth of a B73-Zd F_1 plant; and **F**: regrowth

of a RF-Zd F_1 plant. B73, Mo17 and RF represent, respectively, inbred lines B73 and Mo17 and

cultivar Rhee Flint of *Z. mays*.

Figure 2. Photos of abnormal F_1 plants of crosses of *Zea diploperennis* with *Z. mays* inbred lines

B73 (A & B) and Mo17 (C) or cv. Rhee Flint (D).

Figure 3. Photos of the ears produced from a *Zea mays* cv Rhee Flint x *Z. diploperennis* F_1 plant

in different seasons (the upper panel) and from F_2 in summer 2014 in greenhouse (the lower

panel).

Figure 4. Photos of *Zea mays* Mo17-*Z. diploperennis* F_2 plants, showing regrowth from the basal

node of a single-stalked plant (A) or non-regrowth from a multi-stalked plant (B).

Figure 5. Graphics showing LOD scores of the QTL mapping the B73-Zd F2 population without

(A) or with (B) chi-square imputation. The 95% threshold lines (the parallel red dash lines) were

calculated with 1,000 permutation. Significant QTL/loci are indicated by the location of the peak

SNPs of the loci.

Figure 6. Genetic map of 30 representing SNPs and genes *gt1*, *id1*, and *tb1*. Each SNP represents

a one- Mbp region except of SNP S2_27934739, which represent a SNP cluster.

Supplementary documents:

Supplementary Table S1. Segregation of regrowability among the B73-Zd F_3s.

Supplementary Table S2. Phenotypes and the *gt1*, *id1* and *tb1* haplotypes of 26 F_2 plants and

three F_3 populations of the *Zea mays* cv. Rhee Flint x *Z. diploperennis* cross.

Supplementary Table S3. Steps and codes in TASSEL pipeline including example command

lines and brief descriptions. Parameters are highlighted and described when first be used.

Supplementary Table S4. R codes used for candidate locus/QTL analyses.

Supplementary Figure S1. A photo showing the growth of *Zea diploperennis* and its F_1 with Z.

mays B73 or Mo17 in the field in the Summer 2017.

Supplementary Figure S2. An agarose gel image showing that two molecular markers confirmed

the heterozygosity of a *Z. diploperennis-Z. mays* cv. Rhee Flint F_1 plant over three life-cycles.

Supplementary Figure S3. A photo showing the regrowth of the B73 - *Z. diploperennis* F_4s in the

summer, 2017.

Table 1. Phenotypes and genotypes of the *Zea mays* B73-*Z. diploperennis* F2s

Plant*	PT	*gt1*	*tb1*	*id1*	Plant*	PT	*gt1*	*tb1*	*id1*	Plant*	PT	*gt1*	*tb1*	*id1*
Zea diploperennis	1	1	1	1	BZ2-006-9	1	1	1	2	BZ2-009-6	1	1	1	2
B73	2	2	2	2	BZ2-006-10	1	1	3	2	BZ2-009-7	2	1	-	2
BZ2-001-1	1	1	3	2	BZ2-006-11	2	3	2	2	BZ2-009-8	2	3	2	2
BZ2-001-2	1	3	3	1	BZ2-006-12	2	3	3	2	BZ2-009-9	1	3	1	2
BZ2-001-3	1	1	3	2	BZ2-006-13	2	3	2	1	BZ2-009-10	2	1	3	2
BZ2-001-4	1	1	3	2	BZ2-006-14	1	3	2	2	BZ2-009-11	1	1	3	2
BZ2-001-5	1	1	1	1	BZ2-007-1	2	3	2	2	BZ2-009-12	1	3	3	2
BZ2-002-1	2	1	3	2	BZ2-007-2	2	3	2	2	BZ2-010-1	1	3	2	2
BZ2-002-2	2	1	2	2	BZ2-007-3	1	1	2	2	BZ2-010-2	2	-	3	2
BZ2-002-3	1	3	3	2	BZ2-007-4	1	3	2	2	BZ2-010-3	1	3	1	2
BZ2-002-4	1	3	2	1	BZ2-007-5	1	3	3	2	BZ2-010-4	1	3	2	1
BZ2-002-5	2	3	3	2	BZ2-007-6	2	3	3	2	BZ2-010-5	2	3	1	2
BZ2-002-6	2	3	2	2	BZ2-007-7	1	1	3	2	BZ2-010-6	1	3	2	2
BZ2-002-7	2	3	2	2	BZ2-007-8	1	3	1	2	BZ2-010-7	2	3	1	2
BZ2-002-8	1	1	-	2	BZ2-007-9	1	3	1	2	BZ2-010-8	1	3	2	2
BZ2-002-9	1	3	2	2	BZ2-007-10	2	1	3	2	BZ2-010-9	1	1	3	2
BZ2-002-10	1	3	3	2	BZ2-007-11	2	3	1	2	BZ2-010-10	1	3	2	2
BZ2-002-11	2	1	2	1	BZ2-007-12	1	1	2	2	BZ2-010-11	1	2	3	2
BZ2-002-12	1	1	2	2	BZ2-007-13	1	3	1	2	BZ2-010-12	2	2	3	2
BZ2-002-13	1	3	3	2	BZ2-007-14	2	1	3	2	BZ2-010-13	1	3	3	2
BZ2-002-14	2	1	1	2	BZ2-007-15	2	3	1	2	BZ2-010-14	1	1	2	2
BZ2-002-15	1	1	3	1	BZ2-007-16	1	1	3	2	BZ2-010-15	2	3	3	2
BZ2-002-16	1	3	3	1	BZ2-007-17	2	1	-	1	BZ2-010-16	2	1	3	2
BZ2-002-17	2	3	2	2	BZ2-007-18	1	1	1	2	BZ2-010-17	1	1	3	2
BZ2-002-18	1	1	2	2	BZ2-007-19	1	3	2	2	BZ2-010-18	2	3	3	2
BZ2-002-19	1	1	2	2	BZ2-007-20	1	3	2	2	BZ2-010-19	1	2	2	1
BZ2-002-20	1	1	2	2	BZ2-007-21	2	3	2	2	BZ2-010-20	2	1	2	1
BZ2-002-21	1	1	2	1	BZ2-008-1	1	1	1	2	BZ2-010-21	2	3	3	2
BZ2-002-22	1	3	2	2	BZ2-008-2	1	3	1	2	BZ2-011-1	2	1	-	1
BZ2-002-23	1	3	2	1	BZ2-008-3	1	3	1	2	BZ2-011-2	1	3	2	2
BZ2-002-24	1	3	2	2	BZ2-008-4	2	3	2	2	BZ2-011-3	1	2	3	2
BZ2-002-25	2	1	2	2	BZ2-008-5	1	3	3	2	BZ2-011-4	1	3	1	2
BZ2-004-1	1	3	1	2	BZ2-008-6	2	3	2	2	BZ2-011-5	1	1	-	1
BZ2-004-2	1	1	3	2	BZ2-008-7	2	3	3	2	BZ2-011-6	2	3	2	2
BZ2-004-3	2	3	2	2	BZ2-008-8	1	3	2	2	BZ2-011-7	2	1	2	2
BZ2-004-4	1	1	3	2	BZ2-008-9	2	3	2	2	BZ2-011-8	2	2	1	1
BZ2-004-5	1	3	2	2	BZ2-008-10	1	1	1	2	BZ2-011-9	1	3	2	2
BZ2-004-6	1	3	1	2	BZ2-008-11	2	3	2	2	BZ2-011-10	1	3	1	2
BZ2-006-1	2	3	1	2	BZ2-008-12	1	3	3	2	BZ2-011-11	1	1	2	2
BZ2-006-2	2	1	1	2	BZ2-008-13	1	1	2	2	BZ2-011-12	1	1	3	2
BZ2-006-3	1	3	3	2	BZ2-008-14	2	1	3	2	BZ2-011-13	1	3	1	2
BZ2-006-4	1	3	2	2	BZ2-009-1	2	3	3	2	BZ2-011-14	2	3	1	2
BZ2-006-5	1	3	3	2	BZ2-009-2	1	1	3	1	BZ2-011-15	2	3	3	2
BZ2-006-6	2	3	3	2	BZ2-009-3	2	3	2	2	BZ2-011-16	2	1	2	2
BZ2-006-7	1	3	1	2	BZ2-009-4	1	1	3	2					
BZ2-006-8	1	3	1	2	BZ2-009-5	2	3	3	2					

* Bold: used for SNP calling in GBS; 1: homozygous Zd allele; 2:homozygous B73 allele; 3: heterozygous; -: missing data

Table 2. Segregation of regrowth among the *Zea mays* cv Rhee Flint-*Z. diploperennis* F2s and F3s.

Plant	PT	Plant	PT	Plant	PT	Plant	PT	Plant	PT	Plant	PT	Plant	PT
ZR2-001-1	1	ZR2-001-33	2	ZR2-001-69	0	ZR2-001-102	1	ZR2-001-132	2	ZR2-001-162	2	ZR3-005-7	1
ZR2-001-2	2	ZR2-001-34	1	ZR2-001-71	1	ZR2-001-103	1	ZR2-001-133	1	ZR2-001-163	1	ZR3-005-8	1
ZR2-001-3	1	ZR2-001-35	2	ZR2-001-72	0	ZR2-001-104	1	ZR2-001-134	1	ZR2-001-164	1	ZR3-005-9	1
ZR2-001-4	2	ZR2-001-36	2	ZR2-001-73	1	ZR2-001-105	1	ZR2-001-135	2	ZR2-001-165	2	ZR3-005-10	1
ZR2-001-5	1	ZR2-001-37	2	ZR2-001-74	1	ZR2-001-106	1	ZR2-001-136	2	ZR2-001-166	1	ZR3-005-11	2
ZR2-001-6	1	ZR2-001-38	2	ZR2-001-75	1	ZR2-001-107	2	ZR2-001-137	1	ZR2-001-167	2	ZR3-005-12	2
ZR2-001-7	2	ZR2-001-39	1	ZR2-001-77	0	ZR2-001-108	2	ZR2-001-138	1	ZR2-001-168	2	ZR3-005-13	1
ZR2-001-9	1	ZR2-001-40	2	ZR2-001-78	0	ZR2-001-109	2	ZR2-001-139	1	ZR2-001-169	2	ZR3-005-14	2
ZR2-001-10	2	ZR2-001-42	2	ZR2-001-79	1	ZR2-001-110	1	ZR2-001-140	2	ZR2-001-171	1	ZR3-005-15	2
ZR2-001-11	1	ZR2-001-43	1	ZR2-001-80	1	ZR2-001-111	2	ZR2-001-141	1	ZR3-003-1	1	ZR3-005-16	2
ZR2-001-12	1	ZR2-001-44	1	ZR2-001-81	1	ZR2-001-112	1	ZR2-001-142	1	ZR3-003-2	1	ZR3-009-1	1
ZR2-001-13	2	ZR2-001-45	1	ZR2-001-82	0	ZR2-001-113	2	ZR2-001-143	1	ZR3-003-3	1	ZR3-009-2	1
ZR2-001-14	2	ZR2-001-47	2	ZR2-001-83	0	ZR2-001-114	2	ZR2-001-144	2	ZR3-003-4	2	ZR3-009-3	1
ZR2-001-15	1	ZR2-001-48	2	ZR2-001-84	1	ZR2-001-115	2	ZR2-001-145	1	ZR3-003-6	1	ZR3-009-4	1
ZR2-001-16	2	ZR2-001-49	1	ZR2-001-85	1	ZR2-001-116	1	ZR2-001-146	1	ZR3-003-7	1	ZR3-009-5	1
ZR2-001-17	1	ZR2-001-51	2	ZR2-001-86	1	ZR2-001-117	1	ZR2-001-147	1	ZR3-003-8	1	ZR3-009-6	2
ZR2-001-18	2	ZR2-001-53	2	ZR2-001-87	0	ZR2-001-118	2	ZR2-001-148	1	ZR3-003-9	1	ZR3-009-7	1
ZR2-001-19	2	ZR2-001-54	1	ZR2-001-88	0	ZR2-001-119	1	ZR2-001-149	1	ZR3-003-10	1	ZR3-009-8	1
ZR2-001-20	1	ZR2-001-55	1	ZR2-001-89	0	ZR2-001-120	1	ZR2-001-150	1	ZR3-003-11	2	ZR3-009-9	1
ZR2-001-21	2	ZR2-001-56	1	ZR2-001-90	1	ZR2-001-121	2	ZR2-001-151	1	ZR3-003-12	1	ZR3-009-10	1
ZR2-001-22	2	ZR2-001-57	1	ZR2-001-91	1	ZR2-001-122	1	ZR2-001-152	2	ZR3-003-13	1	ZR3-009-11	1
ZR2-001-23	1	ZR2-001-58	2	ZR2-001-92	1	ZR2-001-123	2	ZR2-001-153	1	ZR3-003-14	1	ZR3-009-12	1
ZR2-001-24	1	ZR2-001-59	1	ZR2-001-93	1	ZR2-001-124	1	ZR2-001-154	1	ZR3-003-15	1	ZR3-009-13	1
ZR2-001-25	2	ZR2-001-60	1	ZR2-001-94	0	ZR2-001-125	1	ZR2-001-155	2	ZR3-003-16	2	ZR3-009-14	2
ZR2-001-26	1	ZR2-001-62	1	ZR2-001-95	0	ZR2-001-126	2	ZR2-001-156	1	ZR3-005-1	1	ZR3-009-15	2
ZR2-001-27	2	ZR2-001-63	1	ZR2-001-97	1	ZR2-001-127	2	ZR2-001-157	2	ZR3-005-2	1	ZR3-009-16	1
ZR2-001-28	2	ZR2-001-64	1	ZR2-001-98	1	ZR2-001-128	2	ZR2-001-158	2	ZR3-005-3	2		
ZR2-001-30	1	ZR2-001-65	1	ZR2-001-99	1	ZR2-001-129	1	ZR2-001-159	2	ZR3-005-4	2	PT: phenotype	
ZR2-001-31	1	ZR2-001-67	2	ZR2-001-100	1	ZR2-001-130	2	ZR2-001-160	1	ZR3-005-5	1	1: regrowth	
ZR20-01-32	1	ZR2-001-68	2	ZR2-001-101	1	ZR2-001-131	2	ZR2-001-161	1	ZR3-005-6	1	2: non-regrowth	

Table 3. Results of the χ2 goodness-of-fit tests of three simple genetic models

Generations	Observed			No. dominant genes (the expected R to NR ratio) and P(χ2)*		
	Total	**R**	**NR**	**1 (3:1)**	**2 (9:7)**	**3 (27:37)**
B73-Zd F_2	134	81	53	0.0001	**0.2964**	0.0001
B73-Zd F3	72	52	20	**0.5862**	0.0063	0.0001
Zd-RF F_2	160	92	68	0.0001	**0.7499**	0.0001
Zd-RF F_3-3	15	12	3	**0.6547**	0.0639	0.3000
Zd-RF F_3-5	16	9	7	0.0833	**1.0000**	0.2547
Zd-RF F_3-9	16	13	3	**0.5637**	0.0438	0.0016

*: the best fit models are in bold.

Table 4. Numbers of SNPs revealed in each chromosome of the B73-ZD F2 population after each filtering step

Chr	Raw SNP number	1st filter	2nd filter	3rd filter	4th filter
1	109,543	5,751	1,628	82	51
2	85,283	4,966	1,476	120	77
3	81,625	4,708	1,200	120	75
4	75,832	3,376	942	112	82
5	77,314	4,409	1,197	198	111
6	58,195	2,938	761	87	49
7	62,280	3,108	877	144	98
8	57,748	3,210	877	20	16
9	57,231	2,982	741	29	16
10	49,107	2,477	732	34	22
Total	**714,158**	**37,925**	**10,431**	**946**	**597**

Table 5. PCR primers used in this study

Primers	Sequences
tb1MF	5'–AGTAGGCCATAGTACGTAC–3'
tb1MR	5'–CTCTTTACCGAGCCCCTACA–3'
tb1ZF	5'–ACTCAACGGCAGCAGCTACCTA–3'
tb1ZR	5'–CGTGTGTGTGATCGAATGGT–3'
tga1cF:	5'–AATAAAATAGAGGAACGTCA–3'
tga1cR:	5'–TGCTGCAAAGGATTACTGAT–3'
mmc0381F	5'–GTGGCCCTGTTGATGAG–3'
mmc0381R	5'–CGACGAGTACCAGGCAT–3'
gt1-ZF:	5'–TCGCCTACATGACCGAGTAC–3'
gt1-ZR:	5'–ATACTCTCAGCTGCTACGCG–3'
gt1-MF:	5'–GAGACCGAGCTGCTGAAGAT–3'
gt1-MR:	5'–TGTAGCTGTTGTAGGCGTACT–3'

Supplementary Table S1. Segeagation of regrowability among the B73-Zd F3s

Plant	PT	Plant	PT	Plant	PT	Plant	PT
BZ3-010-1-1	1	BZ3-010-1-20	2	BZ3-010-1-46	2	BZ3-010-1-69	1
BZ3-010-1-2	1	BZ3-010-1-22	1	BZ3-010-1-47	1	BZ3-010-1-70	1
BZ3-010-1-3	1	BZ3-010-1-23	1	BZ3-010-1-48	1	BZ3-010-1-71	1
BZ3-010-1-4	1	BZ3-010-1-24	1	BZ3-010-1-50	1	BZ3-010-1-72	1
BZ3-010-1-5	1	BZ3-010-1-25	1	BZ3-010-1-51	2	BZ3-010-1-73	1
BZ3-010-1-6	1	BZ3-010-1-26	1	BZ3-010-1-52	1	BZ3-010-1-74	1
BZ3-010-1-7	1	BZ3-010-1-27	1	BZ3-010-1-53	1	BZ3-010-1-75	1
BZ3-010-1-8	2	BZ3-010-1-30	1	BZ3-010-1-54	1	BZ3-010-1-76	2
BZ3-010-1-9	2	BZ3-010-1-31	1	BZ3-010-1-56	1	BZ3-010-1-77	2
BZ3-010-1-11	2	BZ3-010-1-32	1	BZ3-010-1-57	1	BZ3-010-1-78	1
BZ3-010-1-12	1	BZ3-010-1-33	2	BZ3-010-1-68	2	BZ3-010-1-79	1
BZ3-010-1-13	2	BZ3-010-1-34	1	BZ3-010-1-59	1	BZ3-010-1-80	1
BZ3-010-1-14	2	BZ3-010-1-35	1	BZ3-010-1-61	2	BZ3-010-1-81	2
BZ3-010-1-15	1	BZ3-010-1-38	2	BZ3-010-1-62	1	BZ3-010-1-82	2
BZ3-010-1-16	1	BZ3-010-1-39	1	BZ3-010-1-63	1	BZ3-010-1-83	2
BZ3-010-1-17	1	BZ3-010-1-40	1	BZ3-010-1-64	2	BZ3-010-1-84	1
BZ3-010-1-18	1	BZ3-010-1-41	2	BZ3-010-1-67	1	BZ3-010-1-85	1
BZ3-010-1-19	1	BZ3-010-1-42	2	BZ3-010-1-68	1	BZ3-010-1-87	1

1: regrowth; 2: non-regrowth

Suplementary Table S2. Phenotypes and the *gt1* , *id1* and *tb1* haplotypes of 26 F2 plants and three F3 populations of the *Zea mays* cv Rhee Flint x *Z. diploperennis* cross*.

Line	PT	*tb1*	*id1*	*gt1*	Line	PT	*tb1*	*id1*	*gt1*	Line	PT	*tb1*	*id1*	*gt1*
F_2-1	1	3	1	3	F_3-3-2	1	1	2	1	F_3-5-13	1	1	2	1
F_2-2	2	2	2	2	F_3-3-3	1	1	2	1	F_3-5-14	2	1	2	1
F_2-3	1	1	2	3	F_3-3-4	2	1	2	3	F_3-5-15	2	1	2	1
F_2-4	2	2	2	2	F_3-3-6	1	1	2	3	F_3-5-16	2	1	2	1
F_2-5	1	1	2	1	F_3-3-7	1	1	2	1	F_3-9-1	1	1	2	1
F_2-6	1	3	1	1	F_3-3-8	1	1	2	3	F_3-9-2	1	1	2	2
F_2-7	2	2	2	2	F_3-3-9	1	1	2	3	F_3-9-3	1	1	2	2
F_2-9	1	1	2	3	F_3-3-10	1	1	2	3	F_3-9-4	1	1	2	3
F_2-10	2	3	1	3	F_3-3-11	2	1	2	3	F_3-9-5	1	1	2	1
F_2-11	1	2	2	3	F_3-3-12	1	1	2	1	F_3-9-6	2	1	2	2
F_2-12	1	3	1	1	F_3-3-13	1	1	2	3	F_3-9-7	1	1	2	3
F_2-13	2	3	1	3	F_3-3-14	1	1	2	1	F_3-9-8	1	1	2	1
F_2-14	2	2	2	3	F_3-3-15	1	1	2	1	F_3-9-9	1	1	2	3
F_2-15	1	3	1	1	F_3-3-16	2	1	2	2	F_3-9-10	1	1	2	3
F_2-16	2	2	2	1	F_3-5-1	1	1	2	1	F_3-9-11	1	1	2	1
F_2-17	1	3	1	3	F_3-5-2	1	1	2	1	F_3-9-12	1	1	2	3
F_2-18	2	3	1	2	F_3-5-3	2	1	2	1	F_3-9-13	1	1	2	1
F_2-19	2	3	1	2	F_3-5-4	2	1	2	1	F_3-9-14	2	1	2	3
F_2-20	1	3	1	3	F_3-5-5	1	1	2	1	F_3-9-15	2	1	2	3
F_2-21	2	2	2	3	F_3-5-6	1	1	2	1	F_3-9-16	1	1	2	1
F_2-22	2	2	2	3	F_3-5-7	1	1	2	1	F_3-12-1	1	1	2	1
F_2-23	1	2	2	3	F_3-5-8	1	1	2	1	F_3-12-2	1	1	2	1
F_2-24	1	1	2	3	F_3-5-9	1	1	2	1	F_3-12-3	1	1	2	1
F_2-25	1	2	1	1	F_3-5-10	1	1	2	1	F_3-12-4	1	3	1	1
F_2-26	1	3	2	1	F_3-5-11	2	1	2	1	F_3-12-5	1	3	1	1
F_3-3-1	1	1	2	3	F_3-5-12	2	1	2	1					

* For phenotype (PT), "1" is for regrowth and "2" is for non-regrowth; for genotypes, "1" is homozygous for the *Zea diploperennis* allele, "2" is homozygous for the *Z. mays* 's allele and "3" is for heterozygous.

Supplementary Table S3. Steps and codes in TASSEL pipeline including example command lines and brief descriptions. Parameters are highlighted and described when first be used.

	FastqToTagCountPlugin	
Step 1	Description	Derives a tagCount list for each FASTQ file. Keeps only good reads having a barcode and a cut site and no N's in the useful part of the sequence. Trims off the barcodes and truncates sequences that (1) have a second cut site, or (2) read into the common adapter.
	Code	*/home/tassel/run_pipeline.pl -fork1 -Xms8G -Xmx16G - FastqToTagCountPlugin -i fastq -k key.txt -e ApeKI -o tagcounts - endPlugin -runfork1*
	Parameters	-Xms8G, -Xmx16G — Set minimum and maximum usage of RAM.
	MergeMultipleTagCountPlugin	
Step 2	Description	Merges each tagCount file into a single "master" tagCount list.
	Code	*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 - MergeMultipleTagCountPlugin -i tagcounts -o mergedtagcounts/BCA965ANXX_4.cnt **-c 5** -endPlugin -runfork1*
	Parameters	-c 5 — Minimum number of times a tag must be present to be output, typically between 5 and 20. The lower value is, the more sequencing errors will be included analysis. Default: 1
	Note:	Sequencing errors can be tolerated and will be removed by following steps.
	TagCountToFastqPlugin	
Step 3	Description:	Converts a master tagCount file containing all the tags of interest from binary (.cnt) format into a FASTQ format file (.fq) that can then be used as input to bowtie2.
	Code	*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 - TagCountToFastqPlugin -i mergedtagcounts/BCA965ANXX_4.cnt -o BCA965ANXX_4.fq **-c 5** -endPlugin -runfork1*
	Bowtie2 Index	
Step 4	Description	Creates a series of support files needed to operate bowtie2.
	Code	*bowtie2-build chr1.fa,chr2.fa,chr3.fa,chr4.fa,chr5.fa,chr6.fa,chr7.fa,chr8.fa,chr9.fa,chr10.fa /home/gbs/b73ref/b73_ref.fa*
	Notes	Chromosome 1 to 10 from B73_Ref_V4 genome are used to create index files.
	Alignment with bowtie2	
Step 5	Description	Aligns the master set of GBS tags to the reference genome.
	Code	*bowtie2 **-p 8 -N 0 -L 20 -i S,1,0.50 -D 20 -R 3** -x /home/b73ref/b73_ref.fa - U /home/gbs/analysis/BCA965ANXX_4.fq -S /home/gbs/analysis/BCA965ANXX_4.sam*
	Parameters	-p 8 — The number of processors to be used.
		-N 0 — The number of mismatches to allowed in a seed alignment during multiseed alignment. Can be set to 0 or 1. Higher value makes alignment slower but increases sensitivity. Default: 0.
		-L 20 — The length of the seed substrings to align during multiseed alignment. Smaller values make alignment slower but more sensitive. Default: 20.
		-I S, 1, 0.50 — Sets a function governing the interval between seed substrings to use during multiseed alignment. Decide the seed frame length and interval which can influence the accuracy of alignment.

		-D 20	The times that consecutive seed extension attempts can "fail" before Bowtie 2 moves on. Default: 15.
		-R 3	the maximum number of times Bowtie 2 will "re-seed" reads with repetitive seeds. Default: 2.
	Note		Arguments and values used here are identical to –very-sensitive-local.
Step 6	**SAMConverterPlugin**		
	Description		Converts a SAM format alignment (.sam) file produced by bowtie2 into a binary tagsOnPhysicalMap (.topm) file that can be used by the TagsToSNPByAlignmentPlugin for calling SNPs.
	Code		*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 -SAMConverterPlugin -i BCA965ANXX_4.sam -o topm/BCA965ANXX_4.topm -endPlugin -runfork1*
Step 7	**FastqToTBTPlugin**		
	Description		Generates a TagsByTaxa file to parse FASTQ files containing raw GBS sequence data for good reads
	Code		*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 -FastqToTBTPlugin -i fastq -k key.txt -e ApeKI -o tbt -y -t mergedtagcounts/BCA965ANXX_4.cnt -endPlugin -runfork1*
	Parameters	-y	Output in TBTByte format (counts from 0-127) instead of TBTBit (0 or 1).
Step 8	**MergeTagsByTaxaFilesPlugin**		
	Description		Merges all .tbt.bin and/or (preferably) .tbt.byte files present in the input directory.
	Code		*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 -MergeTagsByTaxaFilesPlugin -i tbt -o mergedtbt/BCA965ANXX_4.tbt.byte -endPlugin -runfork1*
Step 9	**TagsToSNPByAlignmentPlugin**		
	Description		Aligns tags from the same physical location against one another, calls SNPs from each alignment, and then outputs the SNP genotypes to a HapMap format file (one file per chromosome).
	Code		*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 -TagsToSNPByAlignmentPlugin -i mergedtbt/BCA965ANXX_4.tbt.byte -y -m topm/BCA965ANXX_4.topm -o hapmap/raw/BCA965ANXX_4_chr+.hmp.txt **-mnF "-0.1"** -ref /home/b73ref/b73_chr.fastq **-sC 1 -eC 10** -endPlugin -runfork1*
	Parameters	-mnF "-0.1"	Minimum value of F. Samples that are not inbreed line should invoke this to be negative value.
		-sC 1	Start chromosome.
		-eC 10	End chromosome.
	Note		This command is only available in TASSEL 3. The DiscoverySNPCaller in TASSEL 5 replaces this command and remains the same SNP calling functions but generate a db file rather than hapmap file. Except listed parameters, all others were used as default.
Step 10	**MergeDuplicateSNPsPlugin**		
	Description		Finds duplicate SNPs in the input HapMap file, and merges them if they have the same pair of alleles
	Code		*/home/tassel/run_pipeline.pl -Xms8G -Xmx16G -fork1 -MergeDuplicateSNPsPlugin -hmp hapmap/raw/BCA965ANXX_4_chr +.hmp.txt -o hapmap/mergedSNPs/BCA965ANXX_4_chr+.hmp.txt **-misMat 0.1 -callHets** -sC 1 -eC 10 -endPlugin -runfork1*
	Parameters	-misMat 0.1	Threshold genotypic mismatch rate above which the duplicate SNPs won't be merged. Default: 0.05. For lines not fully inbred, value of 0.1 is recommended.
		-callHets	When two genotypes at a replicate SNP disagree for a taxon, call it a heterozygote. If the germplasm is not fully

		inbred and contains residual heterozygosity, then -callHets should be on.
	Note	The -callHets parameter is to guarantee heterouzygotes can be correctly output, rather than omitted as errors.
Step 11	colspan GBSHapMapFiltersPlugin	
	Description:	Filtering SNPs with specific requirements.
	Code:	*/home/tassel4/run_pipeline.pl -Xms10G -Xmx16G -fork1 -GBSHapMapFiltersPlugin -hmp hapmap/mergedSNPs/BCA965ANXX_4_chr+.hmp.txt -o hapmap/filt/BCA965ANXX_4_chr+.hmp.txt* **-mnTCov 0.01 -mnSCov 0.2 -mnMAF 0.01 -hLD -mnR2 0.2 -mnBonP 0.005** *-sC 1 -eC 10 -endPlugin -runfork1*
	Parameters:	
	-mnTCov 0.01	Minimum taxon coverage. The minimum SNP call rate for a taxon to be included in the output
	-mnSCov 0.2	Minimum site coverage. The minimum taxon call rate for a SNP to be included in the output
	-mnMAF 0.01	Minimum minor allele frequency
	-hLD True	filtered for those in statistically significant LD with at least one neighboring SNP

Supplementary Table S4. R codes used for candidate locus/QTL analyses

```
library(qtl)
all <- read.cross("csv", file="SNP.csv", genotypes = c("AA","AB", "BB"),
na.strings = "NA", alleles = c("A", "B"))
all <- calc.genoprob(all, step=1.0, off.end = 0.0, error.prob = 1.0e-
4,map.function = "haldane",stepwidth = "fixed")
all <- sim.geno(all, n.draws=32, step=1.0, off.end = 0.0, error.prob = 1.0e-
4,map.function = "haldane",stepwidth = "fixed")
all.scan1 <- scanone(all, pheno.col=2, model="binary", method = "em")
all.scan1.perm <- scanone(all, pheno.col = 2, model = "binary", method="em",
n.perm = 1000)
plot(all.scan1,main="LOD plot of regrowth",ylim = c(0,6))
threshold <- summary(all.scan1.perm, alpha=c(0.1, 0.05, 0.01))
abline(h=threshold[1], lty="dashed", lwd=1, col="blue")
abline(h=threshold[2], lty="dashed", lwd=1, col="yellow")
abline(h=threshold[3], lty="dashed", lwd=1, col="red")
summary(all.scan1, perm=all.scan1.perm, lodcolumn=1, alpha=0.1)
mkname1 <- find.marker(all, chr=2, pos=24.244290)
mkname2 <- find.marker(all, chr=7, pos=5.060739)
effectplot(all,pheno.col=2,mname1= mkname1), ylim=c(0,1))
effectplot(all,pheno.col=2,mname1= mkname2), ylim=c(0,1))
write.csv(all.scan1, "all.scan1.csv",row.names = TRUE)
```

Figures

Figure 1:

Figure 2:

Figure 3:

Figure 4:

Figure 5:

Chromosomes

Figure 6:

Figure S1:

Figure S2:

Figure S3: