Stimulus-induced Gamma Power Predicts the Amplitude of the Subsequent Visual Evoked Response

Mats W.J. van Es[1], Jan-Mathijs Schoffelen[1]

[1] Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen,

Kapittelweg 29, 6525 EN Nijmegen, Netherlands

Corresponding author: Mats W.J. van Es

Donders Institute for Brain, Cognition and Behaviour

Radboud University Nijmegen

P.O. Box 9101 6500 HB Nijmegen, Netherlands

Phone number: +31 24 36 68291

e-mail: m.vanes@donders.ru.nl

Manuscript information:

- 32 pages
- 6 figures
- 0 tables
- 243 words for Abstract
- 566 words for Introduction
- 1254 words for Discussion

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by The Netherlands Organisation for Scientific

Research (NWO Vidi grant to J.M. Schoffelen).

Abstract

The efficiency of neuronal information transfer in activated brain networks may affect behavioral performance. Gamma-band synchronization has been proposed to be a mechanism that facilitates neuronal processing of behaviorally relevant stimuli. In line with this, it has been shown that strong gamma-band activity in visual cortical areas leads to faster responses to a visual go cue. We investigated whether there are directly observable consequences of trial-by-trial fluctuations in non-invasively observed gamma-band activity on the neuronal response. Specifically, we hypothesized that the amplitude of the visual evoked response to a go cue can be predicted by gamma power in the visual system, in the window preceding the evoked response. Thirty-three human subjects (22 female) performed a visual speeded response task while their magnetoencephalogram (MEG) was recorded. The participants had to respond to a pattern reversal of a concentric moving grating. We estimated single trial stimulus-induced visual cortical gamma power, and correlated this with the estimated single trial amplitude of the most prominent event-related field (ERF) peak within the first 100 ms after the pattern reversal. In parieto-occipital cortical areas, the amplitude of the ERF correlated positively with gamma power, and correlated negatively with reaction times. No effects were observed for the alpha and beta frequency bands, despite clear stimulus onset induced modulation at those frequencies. These results support a mechanistic model, in which gamma-band synchronization enhances the neuronal gain to relevant visual input, thus leading to more efficient downstream processing and to faster responses.

Significance statement

Gamma-band activity has been associated with many cognitive functions and improved behavioral performance. For example, high amplitude gamma-band activity in visual cortical areas before a go cue leads to faster reaction times. However, it remains unclear through which neural mechanism(s) gamma-band activity eventually affects behavior. We tested whether the strength of induced gamma-band activity affects evoked activity elicited by a subsequent visual stimulus. We found enhanced amplitudes of early visual evoked activity, and faster responses with higher gamma power. This suggests that gamma-band activity affects the neuronal gain to new sensory input, and thus these results bridge the gap between gamma power and behavior, and support the putative role of gamma-band activity in the efficiency of cortical processing.

Introduction

Mesoscopic and macroscopic electrophysiological signals, as measured invasively as local field potentials (LFPs) or non-invasively as the magneto/electroencephalogram (MEG/EEG), can often be characterized by rhythmic activity patterns in a broad range of frequencies (Buzsáki and Draguhn, 2004). Experimentally, distinct frequency bands have been implicated in various cognitive processes. For instance, cortical gamma-band activity (30-90 Hz) has been associated with attention (Tiitinen et al., 1993; Fries et al., 2001; Taylor et al., 2005), memory (Jensen and Lisman, 1996; Carr et al., 2012) and perception (Gray and Singer, 1989; Llinas et al., 1994). Gamma rhythms result from a balanced interplay between neuronal excitation and inhibition. Fast-spiking interneurons bring about the inhibition of the excitatory drive within a population. Once the inhibition fades off, the excitatory drive activates pyramidal cells and in turn, excites the feedback loop of fast-spiking interneurons. This interaction synchronizes the IPSPs in pyramidal neurons and generates gamma rhythms at the population level (Buzsáki and Wang, 2012). Fries (2015) proposed that this mechanism functions to synchronize inputs down the processing hierarchy, thereby making communication between neuronal groups more effective.

Given its putative mechanistic role in affecting the outcome of cortical computations, gamma-band synchronization has become a popular neural substrate to quantify in relation to behavior during cognitive experiments. This has led to evidence for a relationship between gamma-band synchronization and behavior, both in humans and other mammals. Multiple studies have found a larger pre-stimulus gamma power for perceived versus unperceived stimuli (Makeig and Jung, 1996; Linkenkaer-Hansen, 2004; Hanslmayr et al., 2007; Wyart and Tallon-Baudry, 2008), and strong gamma power in visual areas leads to faster responses to a visual go cue (Womelsdorf et al., 2006; Koch et al., 2009; Hoogenboom et al., 2010). These results are in line with the idea that gamma-band synchronization facilitates stimulus processing, and more specifically, they suggest a behavioral relevance of the strength of gamma-band synchronization in task-relevant areas.

Although the relation between the amplitude of the gamma rhythm and behavior has been

established, relatively little is known about how gamma-band synchronization in sensory cortical

areas affects the chain of neuronal events, leading to an eventual effect on behavior. One way to

investigate this would be to relate trial-by-trial fluctuations of the gamma amplitude and/or phase,

estimated at the moment of task-relevant stimulus onset, with the transient event-related response

to this stimulus. Most studies investigating the mechanisms of gamma-band facilitation used invasive

recording techniques, and focused on the relevance of the gamma phase (Fries et al., 2001; Cardin et

al., 2009; Ni et al., 2016). Ni et al (2016) show, at the mesoscopic scale of LFPs and multiunit activity,

that gamma-band oscillations lead to rhythmic fluctuations in neuronal gain, such that inputs at

phases of high gain elicit stronger multiunit activity.

In the present research, we investigated the effect of trial-by-trial fluctuations in MEG-derived

gamma-band activity on stimulus-evoked activity. Using a visual stimulation paradigm that is known

to robustly induce gamma-band activity in early visual cortical areas, we instructed participants to

respond as fast as possible to an unpredictable salient change in a moving grating. We hypothesized

that intrinsic variability in gamma power reflects variability in the efficiency of information transfer in

the visual processing stream, which would manifest itself as correlated amplitude variability of the

early evoked responses. More salient activation in sensory areas would in turn lead to enhanced

processing in downstream areas, eventually causing a faster behavioral response.

Materials and Methods

Subjects

33 healthy volunteers, of which 22 females and 11 males, participated in the study. Their age range was 18-63 years (mean ± SD: 27 ± 10 years) and they all had normal or corrected-to-normal vision. All subjects gave written informed consent according to the Declaration of Helsinki. The study was approved by the local ethics committee (CMO region Arnhem/Nijmegen). One subject was excluded from analysis due to a technical error, which corrupted one of the data files.

Experimental Design

Stimuli

The experimental task was programmed in MATLAB (R2012b, Mathworks, RRID: SCR_001622) using Psychophysics Toolbox (Brainard and Vision, 1997, RRID: SCR_002881). All stimuli were presented against a gray background. A fixation dot was present throughout the experiment, the color of which indicated when the participant was allowed to blink with their eyes (green for blinking, red for not blinking). A concentric sinusoidal grating was presented at 100% black/white contrast and was tapered towards the edges with a Hanning mask, such that edge effects were excluded (see figure 1). The grating was present at the center of the screen, with a visual angle of 7.1°, 2 sinusoidal cycles per degree and a contraction speed of 2 cycles per second.

Experimental equipment

Stimuli were presented by back-projection onto a semi translucent screen (width 48 cm) by an PROPixx projector with a refresh rate of 120 Hz and a resolution of 1920 x 1080 pixels. Subjects were seated at a distance of 76 cm from the projection screen in a magnetically shielded room. MEG was

recorded throughout the experiment with a 275-channel axial gradiometer CTF MEG system at a

sampling rate of 1200 Hz. In addition, subject's gaze position was continuously recorded using an SR

Research Eyelink 1000 eye-tracking device (RRID: SCR_009602). Head position was monitored in real-

time during the experiment by using head-positioning coils at the nasion and left and right ear canals

of the subject (Stolk et al., 2013). When head position deviated more than 5 mm from the position at

the start of the experiment, subjects readjusted to the original position. Behavioral responses during

the MEG session were recorded using a fiber optic response pad (FORP).

In addition to the MEG recording, anatomical T1 scans of the brain were acquired with a 3T Siemens

MRI system (Siemens, Erlangen, Germany). In order for co-registration of the MEG and MRI datasets,

the scalp surface was mapped using a Polhemus 3D electromagnetic tracking device (Polhemus,

Colchester, Vermont, USA).

Procedure

Subjects were instructed to keep fixation at the fixation dot throughout the experiment (see figure

1). The fixation dot was colored red most of the time, but turned green during the eye-blink period.

After a 1.0 second eye-blink period and a 1.5-2.0 second baseline window, a contracting grating was

presented at the center of the screen. The grating contracted for 1.0-3.0 seconds, after which a

pattern reversal of the stimulus occurred. This functioned as a go cue. Participants had to respond as

fast and as accurately as possible to the go cue by pressing a button with the right index finger.

Responses had to be made within 700 ms. Ten percent of the trials were catch trials, in which no

stimulus change occurred. After the stimulus change the grating continued to contract for another

750 ms, until the end of the trial. There was no feedback of task performance, but participants were

trained before the experiment to make sure they understood the task. Participants completed a

maximum of thirteen blocks, each consisting of forty trials, or until one hour had passed. In between

blocks there was a self-paced break, if needed followed by repositioning of the subject to the original

position (see Experimental Equipment). In total, participants completed between 400 and 520 trials.

Data Analysis

MEG preprocessing

The MEG data was preprocessed offline in MATLAB (2015b, Mathworks, RRID: SCR_001622) using

FieldTrip toolbox (Oostenveld et al., 2011, RRID:SCR_004849) and custom written code. First,

excessively noisy channels and trials were removed from the data by visual inspection. Additionally,

trials with squid jumps or muscle artifacts were removed from the data. Eyetracker data was visually

inspected to discard trials with eye blinks within the latency of interest and trials where the eye

position exceeded 5 degrees from the fixation dot were removed likewise.

The data were demeaned, and high pass filtered at 1 Hz using a finite impulse response windowed

sinc (FIRWS; Widmann, 2006) filter. Power line interference (50 Hz) and its harmonics were removed

using a discrete Fourier transform (DFT) filter. Further, signals relating to cardiac activity or eye blinks

and eye movements were identified and removed from the data using independent component

analysis (ICA). Lastly, the trials of interest were defined as those where a stimulus change was

present and where a behavioral response was made within 700 ms of that event.

MRI processing

MRI data were co-registered to the MEG-based coordinate system using the head-positioning coils

and the digitized scalp surface. Using SPM8 (Penny et al., 2011) we created volume conduction

models of the head, and individual meshes of dipole positions, consisting of a cortically constrained

surface-based mesh with 15,784 vertex locations. These meshes were created using Freesurfer

(RRID: SCR_001847) and HCP workbench (RRID: SCR_008750). The dipole positions were used for the

identification of a virtual channel with the strongest gamma-band response or low frequency

response (see *Single-trial power*), and evoked responses were modeled on a parcellated version of

this mesh. The vertices were grouped into 374 parcels based on a refined version of the Conte69

atlas (Van Essen et al., 2012) in order to reduce the dimensionality of the data (similar to Schoffelen

et al., 2017). Forward models were computed using single-shell volume conductor models that were

derived from individual structural MR images (Nolte, 2003).

Time-frequency analysis

Time-resolved spectral power was estimated for low (2-30 Hz) and high (28-100 Hz) frequencies after

padding the data with zeros to six seconds. For low frequencies, a Hanning tapered 500 ms sliding

time window was used in steps of 50 ms, with 2 Hz resolution. High frequency power was estimated

using a DPSS multi-taper approach with a sliding time window of 250 ms and steps of 50 ms, 4 Hz

resolution and 8 Hz smoothing. Time-frequency activity was expressed relative to a baseline, defined

as [-1.0 -0.25] seconds, time locked to stimulus onset. For initial exploration, spectral decomposition

was performed on synthetic planar gradient data (Bastiaansen and Knösche, 2000), and combined

into a single spectrum per sensor. This way, power spectra could easily be averaged across subjects

for visualization purposes.

Peak frequency

Subject-specific gamma power was estimated on the individualized gamma peak frequency. In order

to estimate peak frequencies, the power spectrum after stimulus onset was contrasted with the pre-

stimulus baseline. First, trials were separated into baseline and stimulus presentation epochs, where

the first 400 ms of stimulus presentation were discarded in order to prevent spectral effects of

evoked activity. Next, trial epochs were cut into 500 ms snippets, with fifty percent overlap. Spectral

power was then estimated on these snippets in the 30-90 Hz range, after tapering with a Hanning

window. Finally, the gamma peak frequency was determined at the maximum power ratio of

stimulus presentation over baseline period, averaged over occipital MEG channels. A similar

approach was used for low frequencies, in the 2-30 Hz range, but here the negative peak (i.e.

showing the largest power reduction from baseline) was used.

Single-trial power

To get an optimal estimate of single-trial gamma power, we created subject-specific virtual channels,

using a DICS beamformer (Gross et al., 2001), scanning the cortically constrained mesh of dipole

positions. The 750 ms of data before the stimulus change were used to ensure the best signal-to-

noise ratio, while at the same time ensuring that the estimate of gamma-band activity was as little as

possible affected by evoked activity. These 750 ms epochs were padded with zeros to one second,

and a multi-taper Fast Fourier transform (FFT) with 8 Hz spectral smoothing was applied to these

data. The same was done for a 750 ms baseline window. Spatial filters were created for each of the

vertex locations, using the cross-spectral density estimated from the concatenated data, at the

subject-specific peak frequencies, and a regularization parameter of 5%. Next, the virtual channel

was selected as the vertex that showed the largest increase in gamma power, relative to baseline.

Next, the single-trial gamma power was estimated on these virtual channels, in the 200 ms just

before the ERF window (see *Single-trial event-related responses*; the window in which power was

estimated ended 20 ms before the start of the ERF window), using a spatial filter with fixed dipole

orientation, optimized for this time window.

To estimate single-trial power estimates for the alpha-beta band, a similar procedure was used. The

cross-spectral density matrix was estimated at individual peak-frequency with 2.5 Hz smoothing,

based on the 400 ms before the ERF window. Since the induced low frequency response was a power

decrease relative to baseline, the vertex location that showed the largest decrease was selected as

the virtual channel.

Single-trial event-related responses

The event-related response to the stimulus change was modelled using a Linearly Constrained

Minimum Variance (LCMV) beamformer on the cortically-constrained meshes of dipole positions,

followed by a parcellation based on an anatomical atlas (see *MRI preprocessing*). Data, time locked to

stimulus change, were selected and baseline corrected based on 100 ms prior to the change. For

each anatomical parcel, the source time courses of the dipoles belonging to this parcel were

concatenated, and subjected to a principal component analysis (PCA). The first spatial component

that explained most variance in the signal was used as a representation of single-trial activity for this

parcel. In order to account for the beamformer's depth bias, the data were normalized by an

estimate of the noise using the covariance matrix of the 200 ms prior to the go cue. The resulting

time courses were low-pass filtered at 30 Hz using a finite impulse response (FIR) windowed sync

function. The data were filtered from right to left, to avoid leakage of pre-change signal into the post-

change estimates. Next, the amplitudes of the single-trial visual evoked responses were estimated in

a time window showing the most prominent peak in the trial averaged ERF, in the first 100ms after

stimulus change, on a subjects-by-subject basis. These windows were manually defined by visual

inspection of the source-level activity time courses.

Correlation of single-trial power and ERF amplitude

Correlations between ERF amplitude, alpha-beta power and gamma power, and response speed, and

between gamma power, response speed and trial length were computed at the single-subject level

using Spearman's rank correlation coefficient. We also computed partial correlations between alpha-

beta/gamma power and response speed, each time accounting for trial length and power values in

the other frequency band.

The correlation between alpha-beta power and gamma power, response speed and trial length was

statistically evaluated using a parametric t-test (against 0) on the distribution of correlation

coefficients over subjects (alpha = 0.05).

Statistical significance of the correlation between alpha-beta/gamma power and ERF amplitude, and

between ERF amplitude and reaction times was assessed using non-parametric permutation tests

(based on 10000 permutations) combined with spatial clustering for family-wise error control (Maris

and Oostenveld, 2007). Under the null hypothesis of no systematic relationship across participants

between gamma power and the ERF amplitude, we created a reference distribution of the group-

level t-statistic of the correlation against zero, using sign swapping of the correlation for random

subsets of subjects. Spatially adjacent parcels with t-values corresponding to a nominal alpha

threshold of 0.05 (0.01 for the correlation between ERF amplitude and RT) were grouped into

clusters, and cluster-level statistics were computed as the sum of t-values within a cluster. The null-

hypothesis was rejected if the maximum cluster-level statistic in the observed data was in the

positive tail of the permutation distribution of cluster-level statistics for the correlation between ERF

amplitude and gamma power, and in the negative tail for the correlation between alpha-beta power

and ERF amplitude, and between ERF amplitude and reaction times, at a level of <0.05 one-sided.

Results

Out of the 400-520 completed trials per subject, on average fifty of them were catch trials, i.e. these trials did not require a response. Overall, the subjects performed with a mean accuracy of 94% (SD = 5.8%). Excluding catch trials and trials with artifacts or excessive eye movements, on average 339 trials per subject (SD = 61) were considered for further analysis. Of this pool, the mean performance rate was 94% (SD = 5.5%) and the mean reaction time over subjects was 371 ms (SD = 56 ms).

Stimulus protocol leads to reliable stimulus-induced changes in gamma power and visual event-related responses

The contracting grating stimuli used here are known to robustly induce gamma-band synchronization (Hoogenboom et al., 2006; Swettenham et al., 2009; Van Pelt and Fries, 2013). In order to verify the spectral characteristics of the stimulus-onset induced neuronal response, we conducted a time-frequency analysis at the channel-level. We contrasted spectral power after stimulus onset with the average spectrum in the baseline window. Figure 2 shows the average spectral power for all subjects in the gamma-frequency range (30-90 Hz). As expected, it was highest in occipital channels and it remained high throughout the whole stimulus presentation. Sources of the activity were localized to early visual areas (see figure 2e). In order to assess the gamma power increase quantitatively we estimated the power increase from baseline at the individual gamma peak frequency (figure 2d) and at the occipital channel that showed maximal increase. Over subjects, gamma power increased on average with 124% (mean, SD = 124%) during stimulation (figure 2d, right). Besides a gamma-band increase, a decrease in power was generally observed in the alpha/low beta band (8-20 Hz). This phenomenon presented itself mainly in occipital channels, and was also strongest in occipital source parcels (see figure 3).

In addition to the stimulus inducing a robust gamma-band response, the stimulus change caused an

event-related field (ERF). Figure 3a shows the ERF of an example subject over trials, together with the

topography of the trial average of the P100 response in figure 3c. The signal-to-noise ratio (SNR) of

the evoked response is relatively poor at the single-trial level. Also, the spatial topography was

variable over subjects (data not shown). In order to reduce the spatial variability over subjects and to

boost the SNR, all further analyses were conducted on the source level. Figure 3b shows a superior

SNR for single trials on the source level compared to the channel level, together with the source

topography in figure 3d. Despite the large variability of the response evoked by the stimulus change

and the percentage increase in induced gamma, all subjects did show the neuronal response that

was expected.

Pre-stimulus gamma power correlates with reaction times

In order to evaluate the effect of pre-stimulus gamma power on behavior, we calculated the

correlation between reaction times and the gamma power preceding the stimulus change. Gamma

power was estimated on a virtual channel in source space. There was a relatively weak but highly

significant, negative correlation of -0.072 (SD = 0.064, t(31) = -6.3, p = $5.1*10^{-7}$), which is in line with

previous research (Womelsdorf et al., 2006; Hoogenboom et al., 2010).

One potential factor that might explain the correlation between gamma power and reaction times is

stimulus jitter (i.e., the time between stimulus onset and the stimulus change). Stimulus jitter was

uniformly distributed, and by consequence the instantaneous probability of a reversal event (hazard

rate) increased over time. There could be a common dependence of gamma power and reaction

times on stimulus expectancy. In order to investigate this possibility, we computed partial

correlations between reaction time, gamma power and stimulus jitter, each time accounting for the

third variable. Since we also found a stimulus-induced power reduction in the alpha-beta band, there

is also a possibility that the correlation between gamma power and reaction times is actually caused

by a common dependence on power in this frequency band. Therefore, we also partialled out power

in the alpha-beta band. Reaction times correlated negatively with both gamma power (M = -0.068,

$t(31) = -6.1$, $p = 8.9 * 10^{-7}$, uncorrected) and stimulus jitter (M = -0.17, $t(31) = -7.0$, $p = 6.9 * 10^{-8}$,

uncorrected), but there was no correlation between gamma power and stimulus jitter ($t(31) = 0.85$, p

= 0.40, uncorrected). Additionally, no significant correlation was found between reaction times and

low frequency power when accounting for gamma power and stimulus jitter ($t(31) = 1.1$, p= 0.27,

uncorrected), nor between low frequency power and gamma power ($t(31) = 0.03$, p = 0.97,

uncorrected). These results indicate that the correlation between gamma power and reaction times

is not likely to be the result of a build-up in expectancy, nor a result of power correlations between

frequency bands. Further, there is no effect of low frequency power on reaction times, above and

beyond the effect of gamma.

Pre-stimulus gamma power predicts ERF amplitude

Considering the vast amount of variability in the evoked response over subjects at the channel level,

and the low signal-to-noise ratio of single trial event-related responses, we estimated the single trial

responses to the stimulus change at the source-level, before quantifying the relation between

gamma power and ERF amplitude. The time courses of the evoked response were projected into

source space with an LCMV beamformer and combined into parcels according to an anatomical brain

atlas (Van Essen et al., 2012). The relation between gamma power and the ERF was quantified as a

Spearman rank correlation at the single subject level and can be seen in figure 5. Group statistical

evaluation showed that the correlation differed significantly from zero (M = 0.027, p=0.01,

nonparametric permutation test, corrected). This difference was supported by a cluster of positive

correlation in source parcels in occipital and parietal areas (figure 5a), supporting the hypothesis that

increased gamma power leads to an increased amplitude of the stimulus-evoked transient.

Additionally, we correlated ERF amplitude and reaction times (figure 6b). This correlation was

significantly lower than zero (M = -0.027, p=0.037, nonparametric permutation test, corrected),

indicating that a higher amplitude of early evoked activity leads to a faster behavioral response. The

cluster that mostly contributed to this effect was found in source parcels belonging to the visual

cortex of the right hemisphere (figure 6a), conform our hypothesis. Although we did not find a

correlation between low frequency power and reaction times, low frequency power might still affect

ERF amplitude. In order to ensure that this was not the case, we tested whether low frequency

power before the ERF was predictive of ERF amplitude. Conform the stimulus-induced alpha-beta

power decrease, if any, a negative correlation was expected with ERF amplitude. This correlation was

not significant at the group level (p = 0.66, nonparametric permutation test, corrected).

Discussion

In this experiment, we investigated the neuronal consequences of trial-by-trial variability in induced gamma-band activity, using a visual stimulus change detection paradigm. We hypothesized that higher gamma power before a go cue would facilitate the efficiency of the processing of the response cue, leading to a more strongly synchronized response in early visual areas, as reflected by higher early latency ERF amplitudes. In turn, the increased processing efficiency would lead to a faster behavioral response.

We computed single trial estimates of gamma power in visual areas, in the time window just prior to the stimulus change, and replicated the finding that higher gamma amplitude leads to faster reaction times in response to the go cue (Hoogenboom et al., 2010). Moreover, we correlated the single-trial gamma power with the amplitude of early latency source-reconstructed event-related activity, and observed a significant group-level effect, where the early latency event-related response in parieto-occipital areas correlated positively with pre-stimulus gamma power. In turn, the amplitude of event-related response in visual areas correlated negatively with reaction times. These findings are consistent with the hypothesis that strong local gamma-band synchronization facilitates the neuronal response to a change in the stimulus, which eventually leads to improved behavioral performance.

Here, the effect of oscillatory activity on the subsequent neuronal response was specific to the gamma band. We also analyzed the effect of alpha/beta activity on the event-related response, since activity in these frequency bands was also prominently modulated by the onset of the visual stimulus. In contrast to the gamma band, we did not observe a significant association between trial-by-trial power fluctuations in these lower frequencies, and trial-by-trial fluctuations in the event-related response, and response speed.

Although Hoogenboom et al. (2010) did not specifically investigate the relation between gamma-band activity and the ERF, the authors did quantify the relation between ERF amplitude and reaction times, and found no significant effect. This latter null-finding is in contrast with our analysis of the

current data. Most likely this discrepancy is due to the fact that the aforementioned study used a

temporally ill-defined stimulus change (change in drift speed). This did not elicit prominent evoked

activity and thus prohibited the reliable estimation of evoked activity. In contrast, we used a pattern

reversal as stimulus change, precisely because this is known to elicit prominent evoked activity

(Nakamura et al., 1997; Di Russo et al., 2005; Barnikol et al., 2006; Perfetti et al., 2007). Because of

this, we were able to reliably estimate the amplitude of early visual components on single trials and

demonstrate a positive correlation between gamma power and ERF amplitude, and a negative

correlation between ERF amplitude and reaction times, in support of our hypothesis.

One possible concern that might confound a mechanistic interpretation of the relation between

gamma-band activity, response speed, and the event-related transients could be a latent variable

that correlates with these measures, causing spurious, indirect correlations. Specifically, the time

interval between a warning cue and an upcoming stimulus is well known to be a determinant of

response speed (Schoffelen et al., 2005; Beck et al., 2014), and temporally better predictable stimuli

are associated with higher amplitudes in early components of evoked activity (Doherty, 2005; Lange

et al., 2006; Dassanayake et al., 2016). Additionally, the hazard rate has been shown to correlate with

spectral characteristics in alpha/beta and gamma band, (Schoffelen et al., 2005; Rolke and Hofmann,

2007; Tsunoda and Kakei, 2008; Rohenkohl and Nobre, 2011) and low frequency spectral responses

have been shown to be anti-correlated with high frequency responses (Hoogenboom et al., 2006;

Womelsdorf et al., 2006; Scheeringa et al., 2011; Spaak et al., 2012) Therefore, variability in the low

frequency response, and/or stimulus expectancy might be a common determinant for gamma power,

and response speed. We checked for these possibilities by estimating the partial correlation between

gamma power and reaction time, controlling for differences in stimulus expectancy and power in low

frequencies. The partial correlations were still significant, and specifically there was no additional

effect of low frequency power on reaction times. This further corroborates the absence of a trial-by-

trial effect of low frequency alpha/beta activity on the ERF amplitude. These results highlight the

relevance and uniqueness of gamma power in behavior, and are in support of a model in which

gamma-band activity modulates neuronal processing in order to affect behavior.

Even though to our knowledge there is no further literature supporting a correlation between

reaction times and the amplitude of early visual evoked components, correlations have been found

with their peak latency (Kammer et al., 1999; Gerson et al., 2005). In contrast to the current

experiment, where the stimulus was constant in every trial and we made use of the natural

variations in the physiological and behavioral response, these studies manipulated either luminance

or natural scenes in order to do so. Disregarding the source of variation in the physiological and the

behavioral response, it is conceivable that pre-stimulus change gamma-band activity might also

affect the latency of the evoked response in addition to its amplitude, and the combination of both

ultimately affects behavior. This is beyond the scope of the current study, but would be an

interesting topic in future research.

Our main effect is in contrast to Privman et al. (2011). The authors used a repetition suppression

paradigm, and found a reduction in ERP power in higher order visual areas as a function of gamma

power in response to the second stimulus. The authors hypothesized that the gamma-band activity

caused by the first stimulus might be sustained after its offset and disrupts synchronization of the

neural population, selective for the second incoming stimulus. Thus, their findings might be specific

to the simulation protocol used, which is further supported by the finding that the repetition

suppression effect is largest when the stimuli are more similar, leading to larger overlapping

neuronal representations (Grill-Spector et al., 2006).

In this study, we used non-invasive MEG recordings in human participants. In contrast to invasive

recordings, MEG lacks the high spatial resolution and high signal-to-noise ratio to allow for a detailed

functional and spatial interpretation of our findings. In contrast to the present findings, recent work

using invasive data from macaques and cats (Ni et al., 2016), showed that the gain of the multiunit

response in primary visual cortex is dependent on the gamma phase of the local field potential.

However, the authors did not investigate the functional relevance of gamma amplitude, nor did we

study gamma phase. Still, their results and our results are not contradictive: the amount of

synchronization on the one hand, reflected by gamma power, and high excitability phases on the

other hand, might both contribute to enhanced neuronal gain.

In addition to the relatively limited spatial resolution, the high spatiotemporal variability in the

response across subjects did not allow for a consistent assignment of even the early ERF components

to a specific subregion in the visual system. The amplitude of the ERF was estimated as the most

prominent peak within the first 100 ms after the go cue, which in terms of latency is well beyond the

first geniculate input into primary visual cortex and might even reflect extrastriate activity, and thus

likely reflects a more widespread activation of several cortical areas. Despite this limitation, our

findings indicate that gamma-band activity increases the neuronal gain to new visual input. In

addition, the fact that this effect can be shown at the spatial scale at which MEG operates, provides

further justification to use gamma-band responses as a physiologically and mechanistically inspired

dependent variable in non-invasive human cognitive neuroscience experiments.

References

Barnikol UB, Amunts K, Dammers J, Mohlberg H, Fieseler T, Malikovic A, Zilles K, Niedeggen M, Tass PA (2006) Pattern reversal visual evoked responses of V1/V2 and V5/MT as revealed by MEG combined with probabilistic cytoarchitectonic maps. NeuroImage 31:86–108.

Bastiaansen MC., Knösche TR (2000) Tangential derivative mapping of axial MEG applied to event-related desynchronization research. Clin Neurophysiol 111:1300–1305.

Beck MR, Hong SL, van Lamsweerde AE, Ericson JM (2014) The Effects of Incidentally Learned Temporal and Spatial Predictability on Response Times and Visual Fixations during Target Detection and Discrimination Hamed SB, ed. PLoS ONE 9:e94539.

Brainard DH, Vision S (1997) The psychophysics toolbox. Spat Vis 10:433–436.

Buzsáki G, Draguhn A (2004) Neuronal oscillations in cortical networks. science 304:1926–1929.

Buzsáki G, Wang X-J (2012) Mechanisms of Gamma Oscillations. Annu Rev Neurosci 35:203–225.

Cardin JA, Carlén M, Meletis K, Knoblich U, Zhang F, Deisseroth K, Tsai L-H, Moore CI (2009) Driving fast-spiking cells induces gamma rhythm and controls sensory responses. Nature 459:663–667.

Carr MF, Karlsson MP, Frank LM (2012) Transient Slow Gamma Synchrony Underlies Hippocampal Memory Replay. Neuron 75:700–713.

Dassanayake TL, Michie PT, Fulham R (2016) Effect of temporal predictability on exogenous attentional modulation of feedforward processing in the striate cortex. Int J Psychophysiol 105:9–16.

Di Russo F, Pitzalis S, Spitoni G, Aprile T, Patria F, Spinelli D, Hillyard SA (2005) Identification of the neural sources of the pattern-reversal VEP. NeuroImage 24:874–886.

Doherty JR (2005) Synergistic Effect of Combined Temporal and Spatial Expectations on Visual Attention. J Neurosci 25:8259–8266.

Fries P, Neuenschwander S, Engel AK, Goebel R, Singer W (2001) Rapid feature selective neuronal synchronization through correlated latency shifting. Nat Neurosci 4:194–200.

Gerson AD, Parra LC, Sajda P (2005) Cortical origins of response time variability during rapid discrimination of visual objects. NeuroImage 28:342–353.

Gray CM, Singer W (1989) Stimulus-specific neuronal oscillations in orientation columns of cat visual cortex. Proc Natl Acad Sci 86:1698–1702.

Grill-Spector K, Henson R, Martin A (2006) Repetition and the brain: neural models of stimulus-specific effects. Trends Cogn Sci 10:14–23.

Gross J, Kujala J, Hamalainen M, Timmermann L, Schnitzler A, Salmelin R (2001) Dynamic imaging of coherent sources: Studying neural interactions in the human brain. Proc Natl Acad Sci U S A 98:694–699.

Hanslmayr S, Aslan A, Staudigl T, Klimesch W, Herrmann CS, Bäuml KH (2007) Prestimulus oscillations
predict visual perception performance between and within subjects. NeuroImage 37:1465–
1473.

Hoogenboom N, Schoffelen JM, Oostenveld R, Parkes LM, Fries P (2006) Localizing human visual
gamma-band activity in frequency, time and space. NeuroImage 29:764–773.

Hoogenboom N, Schoffelen JM, Oostenveld R, Fries P (2010) Visually induced gamma-band activity
predicts speed of change detection in humans. NeuroImage 51:1162–1167.

Jensen O, Lisman JE (1996) Hippocampal CA3 region predicts memory sequences: accounting for the
phase precession of place cells. Learn Mem 3:279–287.

Jensen O, Mazaheri A (2010) Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by
Inhibition. Front Hum Neurosci 4 Available at:
http://journal.frontiersin.org/article/10.3389/fnhum.2010.00186/abstract [Accessed August
2, 2018].

Kammer T, Lehr L, Kirschfeld K (1999) Cortical visual processing is temporally dispersed by luminance
in human subjects. Neurosci Lett 263:133–136.

Koch SP, Werner P, Steinbrink J, Fries P, Obrig H (2009) Stimulus-Induced and State-Dependent
Sustained Gamma Activity Is Tightly Coupled to the Hemodynamic Response in Humans. J
Neurosci 29:13962–13970.

Lange K, Krämer UM, Röder B (2006) Attending points in time and space. Exp Brain Res 173:130–140.

Linkenkaer-Hansen K (2004) Prestimulus Oscillations Enhance Psychophysical Performance in
Humans. J Neurosci 24:10186–10190.

Llinas R, Ribary U, Joliot M, Wang X-J (1994) Human oscillatory brain activity near 40 Hz coexists with
cognitive temporal binding. Proc Natl Acad Sci 19:11748–11751.

Makeig S, Jung TP (1996) Tonic, phasic, and transient EEG correlates of auditory awareness in
drowsiness. Cogn Brain Res 4:15–25.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J Neurosci
Methods 164:177–190.

Nakamura A, Kakigi R, Hoshiyama M, Koyama S, Kitamura Y, Shimojo M (1997) Visual evoked cortical
magnetic fields to pattern reversal stimulation. Cogn Brain Res 6:9–22.

Ni J, Wunderle T, Lewis CM, Desimone R, Diester I, Fries P (2016) Gamma-Rhythmic Gain Modulation.
Neuron 92:240–251.

Nolte G (2003) The magnetic lead field theorem in the quasi-static approximation and its use for
magnetoencephalography forward calculation in realistic volume conductors. Phys Med Biol
48:3637–3652.

Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open Source Software for Advanced
Analysis of MEG, EEG, and Invasive Electrophysiological Data. Comput Intell Neurosci 2011:1–
9.

Perfetti B, Franciotti R, Della Penna S, Ferretti A, Caulo M, Romani GL, Onofrj M (2007) Low- and
high-frequency evoked responses following pattern reversal stimuli: A MEG study supported
by fMRI constraint. NeuroImage 35:1152–1167.

Privman E, Fisch L, Neufeld MY, Kramer U, Kipervasser S, Andelman F, Yeshurun Y, Fried I, Malach R
(2011) Antagonistic Relationship between Gamma Power and Visual Evoked Potentials
Revealed in Human Visual Cortex. Cereb Cortex 21:616–624.

Rohenkohl G, Nobre AC (2011) Alpha Oscillations Related to Anticipatory Attention Follow Temporal
Expectations. J Neurosci 31:14076–14084.

Rolke B, Hofmann P (2007) Temporal uncertainty degrades perceptual processing. Psychon Bull Rev
14:522–526.

Scheeringa R, Fries P, Petersson K-M, Oostenveld R, Grothe I, Norris DG, Hagoort P, Bastiaansen
MCM (2011) Neuronal Dynamics Underlying High- and Low-Frequency EEG Oscillations
Contribute Independently to the Human BOLD Signal. Neuron 69:572–583.

Schoffelen JM, Hultén A, Lam N, Marquand AF, Uddén J, Hagoort P (2017) Frequency-specific
directed interactions in the human brain network for language. Proc Natl Acad Sci 114:8083–
8088.

Schoffelen JM, Oostenveld R, Fries P (2005) Neuronal Coherence as a Mechanism of Effective
Corticospinal Interaction. 308:111–114.

Spaak E, Bonnefond M, Maier A, Leopold DA, Jensen O (2012) Layer-Specific Entrainment of Gamma-
Band Neural Activity by the Alpha Rhythm in Monkey Visual Cortex. Curr Biol 22:2313–2318.

Stolk A, Todorovic A, Schoffelen J-M, Oostenveld R (2013) Online and offline tools for head
movement compensation in MEG. NeuroImage 68:39–48.

Swettenham JB, Muthukumaraswamy SD, Singh KD (2009) Spectral Properties of Induced and Evoked
Gamma Oscillations in Human Early Visual Cortex to Moving and Stationary Stimuli. J
Neurophysiol 102:1241–1253.

Taylor K, Mandon S, Freiwald WA, Kreiter AK (2005) Coherent oscillatory activity in monkey area v4
predicts successful allocation of attention. Cereb Cortex 15:1424–1437.

Tiitinen H, Sinkkonen J, Reinikalnen K, Alho K, Lavikainen J, Näätänen R (1993) Selective attention
enhances the auditory 40-Hz transient response in humans. Nature 364:59–60.

Tsunoda Y, Kakei S (2008) Reaction time changes with the hazard rate for a behaviorally relevant
event when monkeys perform a delayed wrist movement task. Neurosci Lett 433:152–157.

Van Essen DC, Glasser MF, Dierker DL, Harwell J, Coalson T (2012) Parcellations and Hemispheric
Asymmetries of Human Cerebral Cortex Analyzed on Surface-Based Atlases. Cereb Cortex
22:2241–2262.

Van Pelt S, Fries P (2013) Visual stimulus eccentricity affects human gamma peak frequency.
NeuroImage 78:439–447.

Widmann A (2006) Firfilt EEGLAB Plugin, Version 1.5. 1. Leipz Univ Leipz.

Womelsdorf T, Fries P, Mitra PP, Desimone R (2006) Gamma-band synchronization in visual cortex
predicts speed of change detection. Nature 439:733–736.

Wyart V, Tallon-Baudry C (2008) Neural Dissociation between Visual Awareness and Spatial
Attention. J Neurosci 28:2667–2679.

Legends

Figure 1. Task time-line. Each trials starts with a 1.0 second blink period, followed by a variable baseline period (1.5-2.0 s). A concentric drifting grating is presented for 1.0-3.0 seconds, after which a stimulus reversal occurs. The grating continues drifting for 0.75 seconds, until the end of the trial, during which a response has to be made.

Figure 2. Stimulus induces strong gamma synchronization. A) Group-average time-frequency spectrum. Gamma (50-70 Hz) power peaks right after stimulus onset and is sustained throughout stimulus presentation. B) Topography of the stimulus-induced gamma-band activity. Circles reflect selected channels, shown in A. C) Individual (gray line) and group-average (black line) power spectra on the channel level. D) Left: gamma peak frequencies. Generally, peak frequencies were in the 40-70 Hz range. Right: power changes (right) at individual gamma peak frequencies. Power changes were highly variable. E) Source level activity of induced gamma the group average t-value. Induced gamma activity was strongest in occipital regions.

Figure 3. Stimulus induces desynchronization in alpha-beta band (8-20 Hz). Similar to figure 2 but for low frequencies. Power in de alpha and low beta range decreased after stimulus onset, which was strongest in occipital parts of the cortex.

Figure 4. Signal-to-noise ratio of evoked activity is higher at the source level than at the channel level. A-B: single-trial time courses of evoked activity for the channel/parcel depicted in C/D, for a representative subject. C-D: topography of the P100 (depicted in A/B), evoked by the stimulus change.

Figure 5. Occipital gamma power correlates with ERF amplitude. A) Group statistic of the correlation

between gamma power and ERF amplitude. B) boxplot (left) of correlation values and individual

correlations (right). For each subject, correlations were averaged over those parcels belonging to the

cluster that contributed to the significant effect.

Figure 6. ERF amplitude in occipital cortex correlates with reaction times. A) Group statistic of the

correlation between ERF amplitude and reaction times. B) boxplot (left) of correlation values and

individual correlations (right). For each subject, correlations were averaged over those parcels

belonging to the cluster that contributed to the significant effect.

Illustrations and tables

Figure 1.

Figure 2.

Figure 3.

Figure 4.

A

B

Figure 5.

Figure 6.