**The C-terminal tail of the bacterial translocation ATPase SecA modulates its activity**

Running title: Regulation of SecA by its C-terminal tail

Mohammed Jamshad[1,2], Timothy J. Knowles[1,2], Scott A. White[2], Douglas G. Ward[3], Fiyaz

Mohammed[4], Kazi Rahman[2], Gareth W. Hughes[2], Günter Kramer[5], Bernd Bukau[5], and

Damon Huber[2,*].

[1]These authors contributed equally to this work

[2]Institute for Microbiology and Infection

School of Biosciences

University of Birmingham

Edgbaston, Birmingham, UK

[3]Institute of Cancer and Genomic Sciences

University of Birmingham

Edgbaston, Birmingham, UK

[4]Institute of Immunology and Immunotherapy

University of Birmingham

Edgbaston, Birmingham, UK

[5]Center for Molecular Biology of Heidelberg University (ZMBH); German Cancer Research

Center (DKFZ), ZMBH-DKFZ Alliance

Heidelberg

Germany

*To whom correspondence should be addressed:

School of Biosciences

University of Birmingham

Edgbaston

Birmingham B15 2TT

UK

d.huber@bham.ac.uk

**Abstract**

SecA is an evolutionarily conserved and essential ATPase that is required for the

translocation of a subset of proteins across the cytoplasmic membrane in bacteria. SecA can

recognise proteins that are destined for translocation as they are still being synthesised in

order to deliver them to the membrane-bound Sec machinery. However, the mechanism of

cotranslational substrate recognition is not well defined. In *E. coli*, SecA contains a relatively

long C-terminal tail (CTT), which consists of a small metal-binding domain (MBD) that is

attached to the C-terminus of the catalytic core by a flexible linker (FLD). In this study, we

investigated the role of the CTT in the interaction of SecA with the ribosome and nascent

polypeptides.  Previous work indicates that the CTT is required for interaction with the

molecular chaperone SecB. However, phylogenetic analysis suggested that the CTT (and the

MBD in particular) has an additional function. Binding experiments indicated that the CTT

interacts with 70S ribosomes, and disruption of the entire CTT moderately reduced the

affinity of SecA for ribosomes. However, disruption of the MBD alone significantly

increased the affinity of SecA for ribosomes and inhibited the interaction of SecA with

substrate protein, suggesting that the FLD affects the conformation of SecA.

Photocrosslinking and mass spectrometry indicated that the FLD is bound at the site where

SecA binds to substrate proteins. Structural analysis by x-ray crystallography and small-angle

x-ray scattering (SAXS) provided insight into how the CTT influences the structure of SecA

in solution. Finally, site-specific crosslinking experiments indicated that binding to nascent

substrate protein affects the conformation of SecA. Taken together, our results suggest that

the CTT regulates the ability of SecA to interact with substrate proteins.

## Introduction

In *Escherichia coli*, approximately a quarter of all newly synthesised proteins are

transported across the cytoplasmic membrane by the Sec machinery (Cranford Smith &

Huber, 2018; Van den Berg et al., 2004). Of these, the majority (~65%) require the assistance

of SecA, a evolutionarily conserved and essential motor ATPase that drives protein

translocation in bacteria (Cranford Smith & Huber, 2018). The catalytic core of SecA (amino

acids ~1-832 in *E. coli*) contains five domains (**supplemental figure S1**): nucleotide binding

domain-1 (NBD1; amino acids 9-220 & 378-411), nucleotide binding domain-2 (NBD2; 412-

620), the polypeptide crosslinking domain (PPXD; 221-377), the α-helical scaffold domain

(HSD; 621-672 & 756-832) and the α-helical wing domain (HWD; 673-755). Binding and

hydrolysis of ATP at the interface of NBD1 and NBD2 cause conformational changes in the

HSD and HWD, which drive translocation (Collinson, Corey, & Allen, 2015; Cranford Smith

& Huber, 2018). In addition, the PPXD undergoes a large conformational change, swinging

from a position near the HWD (the "closed" conformation) to a position near NBD2 (the

"open" conformation) (Chen, Bauer, Rapoport, & Gumbart, 2015; Zimmer, Nam, &

Rapoport, 2008; Zimmer & Rapoport, 2009). SecA binds to substrate protein in the groove

formed between the PPXD and the two NBDs, and the PPXD serves as a "clamp" that locks

unfolded substrate proteins into this groove when it is in the open conformation (Zimmer &

Rapoport, 2009).

SecA can recognise proteins that are destined for translocation across the cytoplasmic

membrane while they are still being synthesised (i.e. cotranslationally) (Huber et al., 2017;

Huber et al., 2011).  Previous work indicates that SecA binds to the ribosome and that

ribosome binding facilitates its interaction with nascent chains (Chun & Randall, 1994;

Huber et al., 2017; Huber et al., 2011). The binding site for SecA on the ribosome includes

ribosomal protein uL23, which is located adjacent to the opening of the polypeptide exit

channel (Huber et al., 2011). Binding is mediated by the N-terminal α-helix of SecA and the

N-terminal portion of the HSD (Huber et al., 2011; Singh et al., 2014). The structure of the

SecA-ribosome complex was recently determined at medium resolution (~11 Å) by cryo-

electron microscopy (Singh et al., 2014). However, the molecular mechanism governing the

recognition of nascent substrate proteins is not known.

In addition to the catalytic core, most SecA proteins contain a relatively long C-

terminal tail (CTT; also known as the C-terminal linker (Hunt et al., 2002)), whose function

is not well understood (**figure 1A**). In *E. coli*, the CTT (833-901) contains of a small metal

binding domain (MBD; 878-901) and a structurally flexible linker domain (FLD; 833-877).

Although it is not required for protein translocation (Fekkes, van der Does, & Driessen, 1997;

Or, Boyd, Gon, Beckwith, & Rapoport, 2005; Or, Navon, & Rapoport, 2002), *E. coli* strains

producing a C-terminally truncated SecA protein display modest translocation defects

(Grabowicz, Yeh, & Silhavy, 2013; Or et al., 2005). The MBD coordinates a $Zn^{2+}$ ion *via*

three cysteines and a histidine (or in some species, four cysteines) (Dempsey et al., 2004;

Fekkes, de Wit, Boorsma, Friesen, & Driessen, 1999), and in *E. coli* it is required for

interaction of SecA with SecB (Breukink et al., 1995; Fekkes et al., 1997; Kimsey, Dagarag,

& Kumamoto, 1995). SecB is a molecular chaperone that is required for the secretion of a

subset of Sec substrate proteins (Randall & Hardy, 2002). Although not all SecA proteins

contain an MBD, the MBD is conserved in many species that do not have a SecB homologue

(Dempsey et al., 2004).

In this study, we investigated the role of the CTT in the recognition of nascent

substrate proteins by SecA. Phylogenetic analysis of the CTT suggested that it could be

involved in binding of SecA to ribosomes, which we confirmed using ribosome

cosedimentation and chemical crosslinking approaches. Strikingly, disruption of the MBD

alone or the entire CTT had opposing effects on multiple different activities of SecA,

suggesting that the CTT affects conformation of the catalytic core. Mass spectrometry, x-ray

crystallography, and small-angle x-ray scattering experiments indicated that the FLD is

bound in the substrate binding groove and affects the conformation of the PPXD. Finally, site

specific chemical crosslinking suggested that binding of the MBD to the ribosome allows

full-length SecA to interact with nascent substrate proteins. Taken together, our results

provide insight into the molecular mechanism underlying nascent substrate recognition by

SecA.

**Results**

**Evolutionary distribution of the MBD of SecA.** To investigate the evolutionary

distribution of the CTT, we analysed the sequences of 156 SecA proteins from bacterial

species in 155 phylogenetic families using ClustalOmega (McWilliam et al., 2013). The

phylogenetic tree produced by this analysis generally placed SecA proteins from more closely

related species (*e.g.* those in the same phylogenetic class) into similar groups (**figure 1B**).

The majority of SecA proteins (143) contained a CTT (**figure 1B, red and black branches**).

Of these, 117 contained an MBD (**figure 1B, black branches**). A small minority (13) lacked

the CTT entirely (**figure 1B, yellow branches**). Of the SecA proteins in the 69 species that

contained a SecB homologue (**figure 1B, starred species**), only two lacked an MBD. The

strong conservation of the MBD in species that contain SecB suggests that there is strong

selective pressure to maintain the MBD in species containing SecB. However, a significant

number of species (52) have a SecA protein containing an MBD but lack a SecB homologue.

In addition, many of the residues implicated in SecB binding were strongly conserved even in

the MBDs of species that lacked SecB (**figure 1C, arrowheads**) (Zhou & Xu, 2003). These

results suggested that the MBD has an evolutionarily conserved function in addition to its

role in binding to SecB.

**The MBD binds to the ribosome near uL23.** Many of the most highly conserved

residues in the MBD consensus sequence are positively charged (**figure 1C**), suggesting that

the MBD could interact with the negatively charged surface of the ribosome. To determine

whether the CTT could bind to ribosomes, we fused the C-terminal 70 amino acids of *E. coli*

SecA to the small ubiquitin-like modifier from *Saccharomyces cerevisiae* (SUMO-CTT).

Sedimentation of SUMO-CTT through a 30% sucrose cushion during ultracentrifugation was

dependent on the presence of ribosomes in the binding reaction, indicating that it could bind

to ribosomes (**figure 1D, lanes 1 & 2**). The ribosome binding activity could be attributed to

the MBD since a shorter protein fusion containing only the MBD (SUMO-MBD) also

cosedimented with ribosomes (**supplemental figure S2**). The presence of purified SecA or

SecB in binding reactions inhibited cosedimentation (**figure 1D, lanes 3-5**), indicating that

binding to ribosomes and to SecB are mutually exclusive.

We also determined the site of interaction between the CTT and the ribosome using

chemical crosslinking. Full-length SecA crosslinks to uL23 in the presence of the non-

specific crosslinking agent EDC (Huber et al., 2011). Incubation of SUMO-CTT with

ribosomes in the presence of 5 mM and 25 mM EDC resulted in the appearance of several

crosslinking products, which crossreacted with antibodies against uL23 and SecA (**figure**

**1E**). These results suggested that the CTT binds in the vicinity of the opening of the

ribosomal exit channel.

**The C-terminal tail affects the affinity of SecA for ribosomes.** To determine the

contribution of the MBD to binding of SecA to the ribosome, we constructed two C-terminal

truncation variants of SecA and measured their affinity for the ribosome using fluorescence

anisotropy (Huber et al., 2011) (**figure 1A**). The dissociation constant ($K_D$) of the complex

between full-length SecA and ribosomes was ~ 640 nM, similar to previously published

figures (Huber et al., 2011). Truncation of the C-terminal 69 amino acids of SecA

(SecAΔCTT) caused a modest, but reproducible, increase in the $K_D$ of the complex (920 nM)

(**figure 2A & table 1**), consistent with the idea that the CTT contributes to ribosome binding.

However, truncation of the C-terminal 21 amino acids (SecAΔMBD) significantly increased

the affinity of SecA for the ribosome ($K_D$ = 160 nM) (**figure 2A & table 1**). These

differences in affinity were also evident from the amount of SecA that cosedimented with

ribosomes during ultracentrifugation (**figure 2B, lanes 1-6**).

**Affinity of truncated SecA proteins for substrate polypeptides.** We reasoned that

the truncation variants could occupy distinct conformations with differing affinities for the

ribosome and that the truncations might therefore affect other activities of SecA. First, we

examined binding of SecA to nascent SecM, a SecA substrate protein that has previously

been used as a model for binding to nascent chains (Huber et al., 2017; Huber et al., 2011).

Binding of SecA to non-translating ribosomes was sensitive to the presence of salt in the

binding buffer (**figure 2B, lanes 7-9**). Binding of SecA and SecAΔCTT to ribosome-nascent

chain complexes containing nascent SecM (SecM-RNCs) was stable in the presence of 250

mM potassium acetate (**figure 2B, lanes 10 & 12**), indicating that they interacted with

nascent SecM. However, nascent SecM did not stabilise binding of SecAΔMBD to RNCs

under the same conditions (**figure 2B, lane 11**).

We next examined the effect of the truncations on the affinity of SecA for free

substrate protein. To this end, we determined the affinity of SecA for a short peptide (VipB),

which was labelled with an environmentally sensitive fluorophore (IAANS-VipB). The

amino acid sequence for VipB does not resemble a Sec targeting signal (Pietrosiuk et al.,

2011), and binding to VipB was used as a model for binding of SecA to the mature portion of

Sec substrate proteins. The affinities of SecA and SecAΔCTT for  IAANS-VipB ($K_D$ = 0.9

μM and 1.7 μM, respectively) were consistent with the previously reported affinity of SecA

for unfolded substrate protein, but the affinity of SecAΔMBD for IAANS-VipB was

significantly lower ($K_D$ = 5.9 μM) (**figure 2C** & **table 1**) (Gouridis, Karamanou, Gelis,

Kalodimos, & Economou, 2009). Furthermore, alanine substitutions in two of the metal-

coordinating cysteines (SecA$^{C885A/C887A}$) greatly reduced the affinity of SecA for IAANS-

VipB (**table 1**), suggesting that disrupting the structure of the MBD was sufficient to cause

this decrease in affinity.

**Effect of alterations to the CTT on the ATPase activity of SecA.** To investigate the

effect of the truncations on the ATPase activity of SecA, we determined the basal ATPase

rates of SecA, SecAΔMBD, SecAΔCTT and SecA$^{C885A/C887A}$. The ATP turnover rate for full-

length SecA was 0.05 s$^{-1}$ (**table 1**), consistent with previously reported figures (Huber et al.,

2011). Deletion of the entire CTT caused a >10-fold increase in the basal ATPase activity

compared to the full-length protein (0.9 s$^{-1}$) (**table 1**), suggesting that the FLD inhibits the

ATPase activity of SecA. SecAΔMBD and SecA$^{C885A/C887A}$ did not hydrolyse ATP at a

detectable rate, suggesting that the MBD is required to relieve the FLD-mediated

autoinhibition.

**C-terminal truncations differentially affect the $T_M$ of folding of SecA.** To

investigate the effect of the C-terminal truncations on the folding of SecA, we examined the

secondary structure content and thermal stability of SecA, SecAΔCTT and SecAΔMBD

using circular dichroism (CD) spectroscopy. The CD spectra of the three proteins indicated

that they were fully folded (**supplemental figure S3A**). However, the denaturation midpoint

temperature ($T_M$) of SecAΔMBD (42°C) was ~1.5°C higher relative to that of the full-length

protein and ~2°C higher than that of SecAΔCTT (**table 1, supplemental figure S3B**),

indicating that SecAΔMBD was more stably folded than SecA or SecAΔCTT.

**SecA truncation variants have differing abilities to complement the growth**

**defect of a ΔsecA mutation.** To investigate the effect of the C-terminal truncations on the

function of SecA *in vivo*, we constructed strains in which the sole copy of the *secA* gene

produced a truncated version of the protein in an IPTG-inducible fashion. Because SecA is

required for viability, growth of these strains was a reflection of the activity of the SecA

variant *in vivo*. Genes producing all three proteins complemented the viability defect of

Δ*secA* mutant (**figure 2D**), indicating that they were functional *in vivo*. SecAΔCTT and

SecAΔMBD displayed cold-sensitive growth defects compared to the strain expressing wild-

type SecA, consistent with a defect in the ability of the proteins to interact with SecB

(Shimizu, Nishiyama, & Tokuda, 1997). However, in contrast to cells producing SecAΔCTT,

cells producing SecAΔMBD as the sole form of SecA displayed a severe slow growth defect

even at the permissive temperature at all IPTG concentrations tested. These results confirmed

that the function of SecAΔMBD was also defective *in vivo*.

219            **The CTT is bound in the substrate binding groove of full-length SecA.** In order to

affect such a range of activities of SecA, we reasoned that the CTT was likely interacting

with the catalytic core of the protein. To investigate this possibility, we incorporated the non-

natural amino acid benzophenylalanine (Bpa) into the CTT at positions 852, 893 and 898

using nonsense suppression (Chin, Martin, King, Wang, & Schultz, 2002). Bpa contains a

photoactivatable side chain that forms covalent crosslinks to nearby molecules containing C-

H bonds. In order to distinguish between early termination products and full-length SecA, we

used a variant of SecA containing a biotin-attachment peptide at its C-terminus (SecA-biotin)

(Huber et al., 2011; Tagwerker et al., 2006). To aid in purification, we fused SecA to the C-

terminus of SUMO, which contained an N-terminal hexahistidine tag. In all cases, the

purified protein cross-reacted with streptavidin (**figure 3A**), indicating that Bpa was

efficiently incorporated. However, SecA containing Bpa at position 852 (SecA[Bpa852]-biotin)

migrated more rapidly than the other proteins in SDS-PAGE (**figure 3A**). Production of this

faster migrating species was very efficient: virtually all of the purified protein migrated with

a lower apparent molecular weight even before photoactivation (**figure 3B**). In addition,

SecA[Bpa852]-biotin exhibited a very low affinity for substrate protein and displayed no

detectable ATPase activity (**table 1**). These results suggested that SecA[Bpa852]-biotin contained

an internal crosslink, which locked it into a conformation similar to that of SecAΔMBD. To

investigate the site of this internal crosslink, we determined the molecular weights of the

tryptic peptides of SecA[Bpa852]-biotin using mass spectrometry (MALDI-TOF). As expected,

the tryptic peptide containing position 852 (851-877) was absent from the mass spectrum of

SecA[Bpa852]-biotin. The only other peptide absent from SecA[Bpa852]-biotin but present in wild-

type SecA-biotin was the peptide corresponding to amino acids 361-382 (**figure 3C**),

suggesting that the crosslink is to one or more of the residues in this sequence. These amino

acids are located in the groove where SecA binds to substrate protein in one of the two

strands linking the PPXD to NBD1 (**figure 3D**) (Cranford Smith & Huber, 2018). The site of

this crosslink was also consistent with the location of the the extreme N-terminal residues of

the FLD in the crystal structure of SecA from *Bacillus subtilis* (1M6N) (Hunt et al., 2002).

**Structural analysis of the SecA truncation variants.** To investigate the effect of the

FLD on the conformation of the SecA catalytic core, we determined the crystal structure of

SecAΔMBD at 3.5Å resolution (6GOX; **figure 4A** and **supplemental table S1**). SecAΔMBD

crystallised as a symmetric dimer in a head-to-tail configuration (**figure 4A**), similar to that

reported for the *E. coli* SecA homodimer in complex with ATP (Papanikolau et al., 2007)

(PDB file 2FSG). Consistent with previous studies, the structure of the PPXD was less well

defined relative to the other domains of the catalytic core, supporting the idea that the PPXD

is structurally mobile (Gold, Whitehouse, Robson, & Collinson, 2013; Zimmer & Rapoport,

2009).  However, the FLD was not resolved, and its effect on the structure of the ribosome-

binding surface could not be determined.

In order to determine the effect of the truncations on the conformation of the PPXD

and the CTT, we investigated the structures of SecA, SecAΔCTT and SecAΔMBD in

solution using small-angle x-ray scattering (SAXS) (**supplemental table S2**). The SAXS

spectra for all three proteins were similar in the low-$q$ region, indicating that the overall

shapes of the three proteins were similar, and the radii of gyration suggested that they were

dimeric, consistent with previous studies (Woodbury, Hardy, & Randall, 2002)

(**supplemental figure S4**). However, the spectra of the three proteins diverged significantly

in the mid-$q$ region (**supplemental figure S4, arrow**), indicating that there were differences

in domain organisation. SecA has been crystallised in several distinct dimer configurations,

and the physiological configuration of the dimer and its relevance is an issue of on-going

dispute (see discussion in Cranford Smith and Huber (2018)). However, fitting of structural

models of the *E. coli* SecA dimer based on PDB files 2FSG (Papanikolau et al., 2007), 2IBM

(Zimmer, Li, & Rapoport, 2006), 1M6N (Hunt et al., 2002), 1NL3 (Sharma et al., 2003),

2IPC (Vassylyev et al., 2006) and 6GOX indicated that under the experimental conditions,

the conformation of the dimer was similar to that found in 2FSG ($\chi^2 = 3.66$) and 6GOX ($\chi^2 =$

5.25) (**supplemental table S3**). To gain insight into the structural differences between the

three proteins, we modelled the SAXS data by structural fitting. Initial fitting runs indicated

that the CTT was in close proximity to the catalytic core in models of full-length SecA and

SecAΔMBD, and further fitting runs were carried out by fixing the FLD to a position

consistent with the Bpa crosslinking results. These models suggested that the PPXD was

positioned much closer to NBD2 (i.e. more "open") in SecA and SecAΔMBD than in

SecAΔCTT ($p = 2.0 \times 10^{-5}$ and $1.1 \times 10^{-7}$, respectively) (**figure 4F**). In models of

SecAΔMBD and SecAΔCTT, the PPXDs in the two protomers of the dimers were positioned

asymmetrically ($p = 1.8 \times 10^{-8}$ and 0.0085, respectively) (**figure 4B-D, F**). Finally, in models

of full-length SecA, the MBD was positioned between NBD2 and the C-terminal portion of

the HSD (amino acids 756-832) in both protomers of the dimer (**figure 4E**). Localisation of

the MBD to this region would position it directly adjacent to the ribosome-binding surface on

the catalytic core (Huber et al., 2011; Singh et al., 2014).

**Site specific crosslinking of SecA to ribosomes.** To investigate the effect of the CTT

on binding of SecA to the ribosome, we incorporated Bpa at 12 positions located on the face

of SecA that binds to the ribosome, including positions 56, 260, 299, 399, 406, 625, 647, 665,

685, 695, 748 and 796 (**figure 5A & B**) (Huber et al., 2011; Singh et al., 2014). In the

presence of purified 70S ribosomes, SecA containing Bpa at positions 399 (SecA$^{Bpa399}$) and

406 (SecA$^{Bpa406}$) produced an additional high molecular weight band in SDS-PAGE, which

was recognised by α-SecA antiserum (**figure 5C**). Analysis of the high-molecular weight

band produced by SecA$^{Bpa399}$ by mass-spectrometry (LC-MS/MS) indicated that it was an

adduct between SecA and ribosomal protein uL29. uL29 is located adjacent to uL23 on the

ribosomal surface, and both F399 and K406 appear to contact uL29 in the structure of the

SecA-ribosome complex determined by cryo-electron microscopy (**figure 5A**) (Singh et al.,

2014). To investigate the effect of a nascent chain on crosslinking between SecA$^{Bpa399}$ and

uL29, we incubated SecA$^{Bpa399}$ with RNCs containing nascent SecM or maltose binding

protein (MBP) (**figure 5D** and **supplemental figure S5**). The presence of a nascent

polypeptide completely inhibited crosslinking of SecA$^{Bpa399}$ to uL29, suggesting that binding

to nascent polypeptide causes a conformational change in SecA that affects its interaction

with the ribosomal surface. However, SecAΔMBD$^{Bpa399}$ produced crosslinks to uL29 in both

the presence and the absence of nascent polypeptide (**figure 5D**), suggesting that

displacement of the FLD is required for interaction with nascent substrate proteins.

**Discussion**

Our results suggest that the CTT controls the conformation of SecA and regulates its

activity. Disruption of the MBD (i) increases the affinity of SecA for the ribosome, (ii)

decreases the affinity of SecA for substrate protein, (iii) inhibits the ATPase activity of SecA,

(iv) increases the thermal stability of SecA, (v) prevents SecA from undergoing a

conformational change upon binding to nascent substrate protein and (vi) causes a defect in

SecA function *in vivo*. However, SecAΔCTT behaves very similarly to full-length SecA,

indicating that the FLD mediates these effects. Chemical crosslinking and structural

modelling of the SAXS data for wild-type SecA suggest that the FLD causes these effects by

binding in the substrate protein binding groove and causing a conformational change in the

PPXD. Gold et al. (2013) have suggested that opening of the PPXD when SecA is bound to

substrate protein (*i.e.* enclosing the substrate protein in the binding groove by the PPXD

clamp) activates the ATPase activity of SecA. Our work suggests that enclosure of the FLD

by the PPXD has the opposite effect—that is, autoinhibiting SecA. We propose that the MBD

is the key for unlocking this autoinhibited conformation in the full-length protein. Previous

work suggests that interaction of the MBD with SecB increases the affinity of SecA for

polypeptides (Gelis et al., 2007). Our results indicate that the MBD also binds to the

ribosome, and modelling of the SAXS data reveal that the MBD is ideally positioned to

interact with the ribosomal surface (Singh et al., 2014).

Taken together, our results allow us to propose a putative mechanism for the

recognition of nascent substrate proteins by SecA (**figure 6**): (i) interaction of the MBD with

the ribosomal surface upon binding of SecA to the ribosome destabilises binding of FLD in

the substrate protein binding groove; (ii) destabilisation of the FLD allows SecA to sample

nascent polypeptides; (iii) the stable interaction of SecA with nascent substrate protein

displaces the FLD from the substrate binding groove; (iv) binding of SecA to nascent

substrate protein causes a conformational change in SecA that leads to release from the

ribosome.

While our results clearly indicate that the CTT is responsible for autoinhibition of

SecA, the physiological role of this autoinhibition is not clear. One possibility is that it

prevents the spurious interaction of SecA with non-substrate proteins in the cytoplasm.

Restricting the ability of SecA to interact with unfolded polypeptides to certain conditions

(*e.g.* in the presence of the ribosome or SecB) would effectively reduce the pool of potential

substrate proteins that SecA needs to screen. Indeed, artificially increasing the pool of SecA-

substrate proteins causes translocation defects *in vivo* (Muller, Reinert, & Malke, 1989;

Oliver & Beckwith, 1982; Wagner et al., 2007). In addition, the spurious translocation of

certain proteins (*e.g.* β-galactosidase) could be toxic (Emr, Schwartz, & Silhavy, 1978; van

Stelten, Silva, Belin, & Silhavy, 2009).

While the basic features of the catalytic core of SecA are highly conserved amongst

bacteria, the SecA proteins of different phylogenetic groups contain a diverse array of loops

and extensions. For example, our phylogenetic analysis indicated that a large number of

bacterial species contain alternative CTTs with structures that are significantly different from

that of *E. coli*. These differences could allow the activity of SecA to be regulated in response

to interaction with a different subset of interaction partners. Nonetheless, many of these

alternative CTTs are highly positively charged (*e.g.* those of many Actinobacteria),

suggesting that they may retain the interaction with the ribosome. Some phylogenetic groups

(*e.g.* the Cyanobacteria) lack a CTT entirely. However, most of these species contain large

loops in between the conserved elements of the catalytic core of the protein. Indeed, *E. coli*

SecA also contains a "variable" subdomain in NBD2 (amino acids ~519-547), which has

been proposed to regulate its activity (Das et al., 2012). It is therefore reasonable to speculate

that large loops in between the conserved features of the catalytic core might carry out

functions analogous to the CTT in *E. coli*.

**Methods**

**Chemicals and media.** All chemicals were purchased from Sigma-Aldrich unless

otherwise indicated. Rabbit anti-SecA antiserum was a laboratory stock. Sheep anti-uL23

antiserum was a kind gift from R. Brimacombe. IR700-labelled anti-rabbit and IR800-

labelled anti-sheep antibodies were purchased from Rockland (Philadelphia, PA).

Horseradish peroxidase (HRP)-labelled anti-rabbit antibody was purchased from GE

Healthcare. HRP-coupled streptactin was purchased from IBA Lifesciences (Goettingen,

Germany). Bpa was purchased from Bachem (Santa Cruz, CA). Strains were grown in

lysogeny broth (LB) containing kanamycin (30 μg/ml) or ampicillin (100 μg/ml) as required.

**Strains and plasmids.** Strains and plasmids were constructed using standard methods

(Miller, 1992; Sambrook & Russell, 2001). For protein-expression plasmids, the DNA

encoding full-length SecA or fragments of SecA were amplified by PCR and ligated into

plasmid pCA528 (His$_6$-SUMO) or pCA597 (Strep$_3$-SUMO) using the *BsaI* and *BamHI*

restriction sites (Andreasson, Fiaux, Rampelt, Mayer, & Bukau, 2008). UAG stop codons

were introduced into plasmids expressing His-SUMO-SecA or His-SUMO-SecA-biotin using

QuikChange (Agilent). Strains DRH1119, DRH1120 and DRH1121, which produce were

constructed as follows: *secA* genes producing full-length SecA, SecAΔMBD and SecAΔCTT

were cloned into pDSW204 (Weiss, Chen, Ghigo, Boyd, & Beckwith, 1999) and introduced

onto the chromosome of strain DRH663 (MC4100 Δ*secA*::Kan$^R$ + pTrcSpc-*secA*)(Huber et

al., 2011) using lambda InCh (Boyd, Weiss, Chen, & Beckwith, 2000). The SecA-producing

plasmid was then cured from these strains by plating on LB containing 1 mM isopropyl-

thiogalactoside (IPTG). All three strains required > 10μM IPTG for growth on LB.

**Phylogenetic analysis.** The sequences of SecA for the given UniProtKB entry names

(The UniProt, 2017) were analysed using ClustalOmega (McWilliam et al., 2013). The

unrooted phylogenetic tree was rendered using iTOL (Ciccarelli et al., 2006). The logo of the

consensus MBD sequence was generated using WebLogo

(https://weblogo.berkeley.edu/logo.cgi).

**Ribosome and protein purification.** Ribosomes and arrested RNCs were purified as

previously described (Huber et al., 2011; Rutkowska et al., 2008). For all SecA proteins,

BL21(DE3) (laboratory stock) was transformed with the appropriate plasmid and grown in

LB in the presence of kanamycin at 37°C to $OD_{600}$ ~1, induced using 1 mM IPTG and shifted

to 18°C overnight. Cells were then harvested by centrifugation and lysed by cell disruption in

buffer (50 mM K·HEPES, pH 7.5, 500 mM NaCl and 0.5 mM TCEP [tris(2-

carboxyethyl)phosphine]) containing cOmplete EDTA-free protease inhibitor cocktail

(Roche). Unlabelled His-tagged proteins were affinity purified by passing over a 5 ml Ni-

NTA HiTrap column (GE Healthcare), washed with buffer containing 50 mM imidazole and

eluted from the column in buffer containing 250 mM imidazole. The eluted protein was cut

with the SUMO-protease Ulp1 and the cut SUMO was removed by passing over a 5 ml Ni-

NTA HiTrap column. The partially purified protein was then concentrated (Centricon) and

purified by size exclusion chromatography using a sepharose S-200 column (GE Healthcare).

Bpa-labelled proteins were purified as described by Huber et al. (2017). For Strep-tagged

proteins, lysates from cells producing SUMO-CTT and SUMO-MBD were passed over

streptactin-coupled sepharose beads (IBA Lifesciences), washed extensively with buffer and

eluted using buffer containing 10 mM desthiobiotin.

**Western blotting.** Western blots were carried out as previously described (Sambrook

& Russell, 2001). Protein samples were resolved using "Any kD" SDS-PAGE gels (BioRad)

and transferred to nitrocellulose membranes. Membranes were probed using the indicated

primary and secondary antisera or with HRP-streptactin. For HRP-based detection,

membranes were developed using ECL (GE Healthcare) and visualised using a BioRad Gel-

Doc. For IR700- and IR800-based detection, membranes were visualised using a LI-COR

Odyssey scanner.

**Ribosome cosedimentation.** Ribosome cosedimentation experiments were carried

out as previously described (Huber et al., 2017). Binding reactions were incubated in buffer

containing 10 mM HEPES potassium salt, pH 7.5, 100 mM potassium acetate, 10 mM

magnesium acetate, 1 mM β-mercaptoethanol for >10 minutes. The reaction mixture was

then layered on top of a 30% sucrose cushion made with the same buffer and centrifuged at

>200,000 x *g* for 90 minutes. The supernatant was discarded. The concentration of ribosomes

in the pellet fractions were normalised using the absorbance at 260 nm.

**Fluorescence anisotropy.** Determination of the $K_D$ of the SecA-ribosome complex

by fluorescence anisotropy was determined as previously described (Huber et al., 2011).

SecA, SecAΔMBD and SecAΔMBD were labelled with $Ru(bpy)_2(dcbpy)$ and the

fluorescence anisotropy was determined on a Jasco FP-6500 fluorometer containing an

ADP303 attachment using an excitation wavelength of 426 nm (slit width 5 nm) and an

emission wavelength of 640 nm (slit width 10 nm).

**CD spectroscopy.** The CD spectra of 2μM solutions of full-length SecA,

SecAΔMBD, or SecAΔCTT in 10 mM potassium phosphate buffer (pH 7.5) were measured

at temperatures that promote folding (10°C) and denaturation (85°C) in a 0.1cm cuvette using

a Jasco J750 CD spectrometer. For thermal titrations, the temperature was raised 0.5 K/min

from 20 to 60°C and circular dichroism was measured at 222 nm.

**Peptide binding.** 600 nM VipB peptide labelled with IAANS (Pietrosiuk et al., 2011)

was incubated with increasing concentrations of SecA or the indicated SecA variant. The

increase in IAANS fluorescence upon binding of SecA was measured using a Jasco FP-6500

fluorometer or a BMG Labtech CLARIOStar.

**ATPase assays.** ATPase activities were determined by measuring the rate of NADH

oxidation in a coupled reaction (Kiianitsa, Solinger, & Heyer, 2003). 1µM SecA, or the

respective SecA variant, was added to a solution containing 250 mM NADH, 0.5 mM

phosphoenolpyruvate, 2 mM ATP, 20/ml lactate dehydrogenase, 100 U/ml pyruvate kinase

and incubated at 25°C, 50 mM K·HEPES, pH 7.5 and 500 mM NaCl. The decrease in

absorbance at 340 nm from the oxidation of NADH to $NAD^+$ was measured using an Anthos

Zenyth 340rt (Biochrom) absorbance photometer equipped with ADAP software. The rate of

ATP hydrolysis was determined from the rate of NADH oxidation by dividing the rate of

decrease in the absorbance by the extinction coefficient for NADH ($6220$ $M^{-1}$ $cm^{-1}$).

**Chemical crosslinking.** Photo-crosslinking of Bpa SecA and non-specific

crosslinking of SUMO-CTT to the ribosome using 1-ethyl-3-(3-dimethylaminopropyl)

carbodiimide (EDC) were carried out as described previously (Huber et al., 2017; Huber et

al., 2011).

**Mass spectrometry.** Auto-crosslinked samples were digested with sequencing grade

trypsin and the masses of the tryptic peptides were determined using MALDI-TOF mass

spectrometry. The identity of ribosomal crosslinking adducts was determined by excising the

protein band from a Coomassie-stained gel and analysing the tryptic peptide fragments using

liquid chromatography-tandem mass spectrometry (LC-MS/MS) identification (The

Advanced Mass Spectrometry Facility, School of Biosciences, University of Birmingham).

**X-ray crystallography.** SecAΔMBD was crystallised by mixing 2 µl of purified

protein (100 µM) with 2 µl of a 9:1 mixture of Peg Ion 2 # 39 buffer (0.04M Citric acid,

0.06M BIS-TRIS propane, pH6.4, 20% Polyethylene glycol 3,350) and Morpheus condition

47, box 2 (0.1M Amino acids, 0.1M Buffer system 3, pH 8.5, 50% v/v Precipitant Mix 3) in a

48-well MRC MAXI plate (Molecular Dimensions, Newmarket, UK). Crystals appeared

within six days and were fully matured by two weeks. Crystals were analysed at Diamond

light source, and the structure was solved by molecular replacement using PDB file 2FSG at

3.5 Å resolution (**Table S1**). The structure was deposited at RCSB under PDB file 6GOX.

**SAXS measurements.** Synchrotron radiation X-ray scattering data were collected on

the ESRF BM29 BioSAXS beamline (Grenoble) (**Table S2**). An in-line Superose 6 column

was used to ensure that the protein was free from aggregates and that it occupied a single

oligomeric state during data collection. The sample-to-detector distance was 3 m, covering a

range of momentum transfer s = 0.03-0.494 Å$^{-1}$ (s = (4π·sin θ)/ λ, where 2θ is the scattering

angle, and λ = 0.992 Å is the X-ray wavelength). Data from the detector were normalized to

the transmitted beam intensity, averaged, placed on absolute scale relative to water and the

scattering of buffer solutions subtracted. All data manipulations were performed using

PRIMUS*qt* and ATSAS (Petoukhov et al., 2012). The forward scattering *I(0)* and radius of

gyration, $R_g$ were determined by Guinier analysis. These parameters were also estimated from

the full scattering curves using the indirect Fourier transform method implemented in the

program GNOM, along with the distance distribution function *p(r)* and the maximum particle

dimensions $D_{max}$. Molecular masses of solutes were estimated from SAXS data by comparing

the extrapolated forward scattering with that of a reference solution of bovine serum albumin.

Computation of theoretical scattering intensities was performed using the program CRYSOL.

**Molecular modelling of SAXS data.** SAXS data has been deposited at the SASBDB

(www.sasbdb.org) with accession codes: SASDDY9 (full-length SecA), SASDDZ9

(SecAΔMBD) and SASDE22 (SecAΔCTT). For modelling based on SAXS data, multiple

runs were performed to verify the stability of the solution, and to establish the most typical

3D reconstructions using DAMAVER. Guinier analysis of the SAXS data indicated that the

protein was dimeric under the conditions used for SAXS. Structural models of the *E. coli*

SecA dimer were generated by aligning the structure of SecA in PDB file 2VDA to PDB files

2FSG, 2IBM, 2IPC, 1M6N, 1NL3 and 6GOX using PyMol v. 1.8.0.5 and refining using

GROMACS (Pronk et al., 2013). Because the CTT is not resolved in the structures used for

modelling, these models were fit to the SAXS spectrum of SecAΔCTT using FoXS

(Schneidman-Duhovny, Hammel, Tainer, & Sali, 2016) (**supplemental table S3**). The

structures of full-length SecA, SecAΔMBD and SecAΔCTT were modelled by fitting the

2FSG dimer to the respective SAXS data by multi-step rigid body refinement using CORAL

(Petoukhov et al., 2012). The positions of NBD1, NBD2, HSD and HWD were fixed in all

models. The regions corresponding to residues 1-8, 220-231, 367-375 were defined as linkers

and modelled as flexible. The PPXD was allowed rigid body movement in all three models.

The FLD (residues 829-832 in SecAΔCTT and 829-880 in SecAΔMBD and full-length

SecA) were modelled as flexible. For full length SecA, the MBD was modelled using PDB

file 1SX0 and allowed rigid-body movement. Because initial modelling indicated that the

FLD was in close contact with the catalytic core, and because photocrosslinking indicated the

FLD was bound in the substrate binding groove, residues 851-854 were modelled to form a

small β-sheet with residues 222-225 and 373-375 and allowed rigid body movement. All ten

independently generated fits for SecA and SecAΔMBD produced plausible structural models

($\chi^2 = 0.97 \pm 0.02$ and $1.87 \pm 0.09$, respectively). Ten of 15 of the fits of SecAΔCTT produced

plausible structural models ($\chi^2 = 1.82 \pm 0.18$). In the remaining five fits, the position of the

PPXD in one of the two protomers was inconsistent with previously published structures of

SecA and occupied a non-realisitic conformation, suggesting increased mobility of the PPXD

in SecAΔCTT.

**<u>Acknowledgements</u>**

We thank B. Zachmann-Brand, Jingli Yu and the functional genomic facility (School of

Biosciences, University of Birmingham) for technical assistance. We thank A. McNally and

members of the Bukau, Mayer, Henderson, Lund, Grainger, Cole and Rossiter groups for

helpful advice and discussions.

**Funding**

DH and MJ were funded by Biological and Biomedical Research Council (BBSRC) grant

BB/L019434/1. GK and BB were funded by the Deutsche Forschungsgemeinschaft (FOR

1805, SFB 638). FM is funded by Wellcome Trust grant 099266/Z/12/Z. TK is funded by

BBSRC grant BB/P009840/1.

**Competing interests**

The authors declare no competing interests.

**References**

Andreasson, C., Fiaux, J., Rampelt, H., Mayer, M. P., & Bukau, B. (2008). Hsp110 is a

nucleotide-activated exchange factor for Hsp70. *J Biol Chem, 283*(14), 8877-8884.

Boyd, D., Weiss, D. S., Chen, J. C., & Beckwith, J. (2000). Towards single-copy gene

expression systems making gene cloning physiologically relevant: lambda InCh, a

simple Escherichia coli plasmid-chromosome shuttle system. *J Bacteriol, 182*(3), 842-

847.

Breukink, E., Nouwen, N., van Raalte, A., Mizushima, S., Tommassen, J., & de Kruijff, B.

(1995). The C terminus of SecA is involved in both lipid binding and SecB binding. *J

Biol Chem, 270*(14), 7902-7907.

Chen, Y., Bauer, B. W., Rapoport, T. A., & Gumbart, J. C. (2015). Conformational Changes

of the Clamp of the Protein Translocation ATPase SecA. *J Mol Biol, 427*(14), 2348-

2359. doi:10.1016/j.jmb.2015.05.003

Chin, J. W., Martin, A. B., King, D. S., Wang, L., & Schultz, P. G. (2002). Addition of a

529             photocrosslinking amino acid to the genetic code of Escherichiacoli. *Proc Natl Acad*

530             *Sci U S A, 99*(17), 11020-11024.

531     Chun, S. Y., & Randall, L. L. (1994). In vivo studies of the role of SecA during protein

532             export in Escherichia coli. *J Bacteriol, 176*(14), 4197-4203.

533     Ciccarelli, F. D., Doerks, T., von Mering, C., Creevey, C. J., Snel, B., & Bork, P. (2006).

534             Toward automatic reconstruction of a highly resolved tree of life. *Science, 311*(5765),

535             1283-1287. doi:10.1126/science.1123061

536     Collinson, I., Corey, R. A., & Allen, W. J. (2015). Channel crossing: how are proteins

537             shipped across the bacterial plasma membrane? *Philos Trans R Soc Lond B Biol Sci,*

538             *370*(1679). doi:10.1098/rstb.2015.0025

539     Cranford Smith, T., & Huber, D. (2018). The way is the goal: how SecA transports proteins

540             across the cytoplasmic membrane in bacteria. *FEMS Microbiol Lett*.

541             doi:10.1093/femsle/fny093

542     Das, S., Grady, L. M., Michtavy, J., Zhou, Y., Cohan, F. M., Hingorani, M. M., & Oliver, D.

543             B. (2012). The variable subdomain of Escherichia coli SecA functions to regulate

544             SecA ATPase activity and ADP release. *J Bacteriol, 194*(9), 2205-2213.

545             doi:10.1128/JB.00039-12

546     Dempsey, B. R., Wrona, M., Moulin, J. M., Gloor, G. B., Jalilehvand, F., Lajoie, G., . . .

547             Shilton, B. H. (2004). Solution NMR structure and X-ray absorption analysis of the

548             C-terminal zinc-binding domain of the SecA ATPase. *Biochemistry, 43*(29), 9361-

549             9371. doi:10.1021/bi0493057

550     Emr, S. D., Schwartz, M., & Silhavy, T. J. (1978). Mutations altering the cellular localization

551             of the phage lambda receptor, an Escherichia coli outer membrane protein. *Proc Natl*

552             *Acad Sci U S A, 75*(12), 5802-5806.

Fekkes, P., de Wit, J. G., Boorsma, A., Friesen, R. H., & Driessen, A. J. (1999). Zinc

stabilizes the SecB binding site of SecA. *Biochemistry, 38*(16), 5111-5116.

Fekkes, P., van der Does, C., & Driessen, A. J. (1997). The molecular chaperone SecB is

released from the carboxy-terminus of SecA during initiation of precursor protein

translocation. *Embo J, 16*(20), 6105-6113.

Gelis, I., Bonvin, A. M., Keramisanou, D., Koukaki, M., Gouridis, G., Karamanou, S., . . .

Kalodimos, C. G. (2007). Structural basis for signal-sequence recognition by the

translocase motor SecA as determined by NMR. *Cell, 131*(4), 756-769.

Gold, V. A., Whitehouse, S., Robson, A., & Collinson, I. (2013). The dynamic action of

SecA during the initiation of protein translocation. *Biochem J, 449*(3), 695-705.

doi:10.1042/BJ20121314

Gouridis, G., Karamanou, S., Gelis, I., Kalodimos, C. G., & Economou, A. (2009). Signal

peptides are allosteric activators of the protein translocase. *Nature, 462*(7271), 363-

367.

Grabowicz, M., Yeh, J., & Silhavy, T. J. (2013). Dominant negative lptE mutation that

supports a role for LptE as a plug in the LptD barrel. *J Bacteriol, 195*(6), 1327-1334.

doi:10.1128/JB.02142-12

Huber, D., Jamshad, M., Hanmer, R., Schibich, D., Doring, K., Marcomini, I., . . . Bukau, B.

(2017). SecA Cotranslationally Interacts with Nascent Substrate Proteins In Vivo. *J

Bacteriol, 199*(2). doi:10.1128/JB.00622-16

Huber, D., Rajagopalan, N., Preissler, S., Rocco, M. A., Merz, F., Kramer, G., & Bukau, B.

(2011). SecA interacts with ribosomes in order to facilitate posttranslational

translocation in bacteria. *Mol Cell, 41*(3), 343-353.

doi:doi:10.1016/j.molcel.2010.12.028

Hunt, J. F., Weinkauf, S., Henry, L., Fak, J. J., McNicholas, P., Oliver, D. B., & Deisenhofer,

J. (2002). Nucleotide control of interdomain interactions in the conformational

reaction cycle of SecA. *Science, 297*(5589), 2018-2026. doi:10.1126/science.1074424

297/5589/2018 [pii]

Kiianitsa, K., Solinger, J. A., & Heyer, W. D. (2003). NADH-coupled microplate

photometric assay for kinetic studies of ATP-hydrolyzing enzymes with low and high

specific activities. *Anal Biochem, 321*(2), 266-271.

Kimsey, H. H., Dagarag, M. D., & Kumamoto, C. A. (1995). Diverse effects of mutation on

the activity of the Escherichia coli export chaperone SecB. *J Biol Chem, 270*(39),

22831-22835.

McWilliam, H., Li, W., Uludag, M., Squizzato, S., Park, Y. M., Buso, N., . . . Lopez, R.

(2013). Analysis Tool Web Services from the EMBL-EBI. *Nucleic Acids Res,*

*41*(Web Server issue), W597-600. doi:10.1093/nar/gkt376

Miller, J. H. (1992). *A Short Course in Bacterial Genetics*. Cold Spring Harbor, NY: Cold

Spring Harbor Press.

Muller, J., Reinert, H., & Malke, H. (1989). Streptokinase mutations relieving Escherichia

coli K-12 (prlA4) of detriments caused by the wild-type skc gene. *J Bacteriol, 171*(4),

2202-2208.

Oliver, D. B., & Beckwith, J. (1982). Regulation of a membrane component required for

protein secretion in Escherichia coli. *Cell, 30*(1), 311-319.

Or, E., Boyd, D., Gon, S., Beckwith, J., & Rapoport, T. (2005). The bacterial ATPase SecA

functions as a monomer in protein translocation. *J Biol Chem, 280*(10), 9097-9105.

Or, E., Navon, A., & Rapoport, T. (2002). Dissociation of the dimeric SecA ATPase during

protein translocation across the bacterial membrane. *Embo J, 21*(17), 4470-4479.

Papanikolau, Y., Papadovasilaki, M., Ravelli, R. B., McCarthy, A. A., Cusack, S.,

Economou, A., & Petratos, K. (2007). Structure of dimeric SecA, the Escherichia coli

preprotein translocase motor. *J Mol Biol, 366*(5), 1545-1557.

Petoukhov, M. V., Franke, D., Shkumatov, A. V., Tria, G., Kikhney, A. G., Gajda, M., . . .

Svergun, D. I. (2012). New developments in the ATSAS program package for small-

angle scattering data analysis. *J Appl Crystallogr, 45*(Pt 2), 342-350.

doi:10.1107/S0021889812007662

Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C.,

& Ferrin, T. E. (2004). UCSF Chimera--a visualization system for exploratory

research and analysis. *J Comput Chem, 25*(13), 1605-1612. doi:10.1002/jcc.20084

Pietrosiuk, A., Lenherr, E. D., Falk, S., Bonemann, G., Kopp, J., Zentgraf, H., . . . Mogk, A.

(2011). Molecular basis for the unique role of the AAA+ chaperone ClpV in type VI

protein secretion. *J Biol Chem, 286*(34), 30010-30021. doi:M111.253377 [pii]

10.1074/jbc.M111.253377

Pronk, S., Pall, S., Schulz, R., Larsson, P., Bjelkmar, P., Apostolov, R., . . . Lindahl, E.

(2013). GROMACS 4.5: a high-throughput and highly parallel open source molecular

simulation toolkit. *Bioinformatics, 29*(7), 845-854. doi:10.1093/bioinformatics/btt055

Randall, L. L., & Hardy, S. J. (2002). SecB, one small chaperone in the complex milieu of the

cell. *Cell Mol Life Sci, 59*(10), 1617-1623.

Rutkowska, A., Mayer, M. P., Hoffmann, A., Merz, F., Zachmann-Brand, B., Schaffitzel, C.,

. . . Bukau, B. (2008). Dynamics of trigger factor interaction with translating

ribosomes. *J Biol Chem, 283*(7), 4124-4132.

Sambrook, J., & Russell, D. W. (2001). *Molecular Cloning: A Laboratory Manual, 3rd ed.*

Cold Spring Harbor, NY: Cold Spring Harbor Press.

Schneidman-Duhovny, D., Hammel, M., Tainer, J. A., & Sali, A. (2016). FoXS, FoXSDock

and MultiFoXS: Single-state and multi-state structural modeling of proteins and their

complexes based on SAXS profiles. *Nucleic Acids Res, 44*(W1), W424-429.

doi:10.1093/nar/gkw389

Schuwirth, B. S., Borovinskaya, M. A., Hau, C. W., Zhang, W., Vila-Sanjurjo, A., Holton, J.

M., & Cate, J. H. (2005). Structures of the bacterial ribosome at 3.5 A resolution.

*Science, 310*(5749), 827-834.

Sharma, V., Arockiasamy, A., Ronning, D. R., Savva, C. G., Holzenburg, A., Braunstein, M.,

. . . Sacchettini, J. C. (2003). Crystal structure of Mycobacterium tuberculosis SecA, a

preprotein translocating ATPase. *Proc Natl Acad Sci U S A, 100*(5), 2243-2248.

doi:10.1073/pnas.0538077100

Shimizu, H., Nishiyama, K., & Tokuda, H. (1997). Expression of gpsA encoding biosynthetic

sn-glycerol 3-phosphate dehydrogenase suppresses both the LB- phenotype of a secB

null mutant and the cold-sensitive phenotype of a secG null mutant. *Mol Microbiol,*

*26*(5), 1013-1021.

Singh, R., Kraft, C., Jaiswal, R., Sejwal, K., Kasaragod, V. B., Kuper, J., . . . Bhushan, S.

(2014). Cryo-Electron Microscopic Structure of SecA Bound to the 70S Ribosome. *J*

*Biol Chem*. doi:M113.506634 [pii]

10.1074/jbc.M113.506634

Tagwerker, C., Flick, K., Cui, M., Guerrero, C., Dou, Y., Auer, B., . . . Kaiser, P. (2006). A

tandem affinity tag for two-step purification under fully denaturing conditions:

application in ubiquitin profiling and protein complex identification combined with in

vivocross-linking. *Mol Cell Proteomics, 5*(4), 737-748.

The UniProt, C. (2017). UniProt: the universal protein knowledgebase. *Nucleic Acids Res,*

*45*(D1), D158-D169. doi:10.1093/nar/gkw1099

Van den Berg, B., Clemons, W. M., Jr., Collinson, I., Modis, Y., Hartmann, E., Harrison, S.

C., & Rapoport, T. A. (2004). X-ray structure of a protein-conducting channel.

*Nature, 427*(6969), 36-44.

van Stelten, J., Silva, F., Belin, D., & Silhavy, T. J. (2009). Effects of antibiotics and a proto-

oncogene homolog on destruction of protein translocator SecY. *Science, 325*(5941),

753-756.

Vassylyev, D. G., Mori, H., Vassylyeva, M. N., Tsukazaki, T., Kimura, Y., Tahirov, T. H., &

Ito, K. (2006). Crystal structure of the translocation ATPase SecA from Thermus

thermophilus reveals a parallel, head-to-head dimer. *J Mol Biol, 364*(3), 248-258.

Wagner, S., Baars, L., Ytterberg, A. J., Klussmeier, A., Wagner, C. S., Nord, O., . . . de Gier,

J. W. (2007). Consequences of membrane protein overexpression in Escherichia coli.

*Mol Cell Proteomics, 6*(9), 1527-1550. doi:10.1074/mcp.M600431-MCP200

Weiss, D. S., Chen, J. C., Ghigo, J. M., Boyd, D., & Beckwith, J. (1999). Localization of FtsI

(PBP3) to the septal ring requires its membrane anchor, the Z ring, FtsA, FtsQ, and

FtsL. *J Bacteriol, 181*(2), 508-520.

Woodbury, R. L., Hardy, S. J., & Randall, L. L. (2002). Complex behavior in solution of

homodimeric SecA. *Protein Sci, 11*(4), 875-882.

Zhou, J., & Xu, Z. (2003). Structural determinants of SecB recognition by SecA in bacterial

protein translocation. *Nat Struct Biol, 10*(11), 942-947.

Zimmer, J., Li, W., & Rapoport, T. A. (2006). A novel dimer interface and conformational

changes revealed by an X-ray structure of B. subtilis SecA. *J Mol Biol, 364*(3), 259-

265.

Zimmer, J., Nam, Y., & Rapoport, T. A. (2008). Structure of a complex of the ATPase SecA

and the protein-translocation channel. *Nature, 455*(7215), 936-943.

Zimmer, J., & Rapoport, T. A. (2009). Conformational flexibility and peptide interaction of

the translocation ATPase SecA. *J Mol Biol, 394*(4), 606-612. doi:S0022-

2836(09)01264-9 [pii]

10.1016/j.jmb.2009.10.024

**Table 1. Biochemical properties of mutant SecA proteins**

| SecA variant | $K_D$ Ribosomes* | $K_D$ VipB | Basal ATPase activity | $T_M$ |
|---|---|---|---|---|
| Wild type | 640 nM | 0.94 μM | $0.053 \pm 0.02$ s$^{-1}$ | $40.7 \pm 0.09$ °C |
| SecAΔZnBD | 160 nM | 1.74 μM | $< 0.001$ s$^{-1}$ | $42.0 \pm 0.08$ °C |
| SecAΔCTT | 920 nM | 5.93 μM | $0.91 \pm 0.02$ s$^{-1}$ | $40.0 \pm 0.1$ °C |
| SecA$^{C885A/C887A}$ | ND* | >10 μM | $< 0.001$ s$^{-1}$ | ND* |
| SecA$^{Bpa852}$ | ND* | >10 μM | $< 0.001$ s$^{-1}$ | ND* |

681 *not determined

Figure 1.

**Figure 1. Phylogenetic analysis of the CTT and binding of *E. coli* CTT to the ribosome.**

(A) Schematic diagram of the primary structure of SecA, SecAΔMBD and SecAΔCTT.

Structures are oriented with the N-termini to the left, and the amino acid positions of the N-

and C-termini are given. Residues of the catalytic core and the CTT are indicated below.

Catalytic core, black; FLD, yellow; MBD, red. (B) Phylogenetic tree of the SecA proteins of

156 representative species from 155 different bacterial families. Species names are given as

the five-letter organism mnemonic in UniProtKB (The UniProt, 2017). Taxonimical classes

are colour-coded according to the legend. Leaves representing SecA proteins with an MBD

are coloured black. Those with CTTs lacking a MBD are coloured red, and those that lack a

CTT entirely are coloured yellow. Species that also contain a SecB protein are indicated with

a star (*). (C) Logo of the consensus sequence of the MBD generated from the 117 species

containing the MBD in the phylogenetic analysis. Positions of the metal-coordinating amino

acids are indicated above. Amino acids that make contact with SecB in the structure of the

MBD-SecB complex (Zhou & Xu, 2003) (1OZB) are indicated by arrowheads below. (D) 10

μM Strep-SUMO-CTT was incubated in the presence or absence of 1 μM ribosomes and

layered on a 30% sucrose cushion. Ribosomes were then sedimented through the cushion by

ultracentrifugation. Where indicated, 1 μM wild-type SecA, 1 μM containing alanine

substitutions at position 625 and 633 (SecA$^{KK}$) or 10 μM tetrameric SecB were included in

the binding reaction. Samples were resolved on SDS-PAGE and probed by western blotting

using HRP-coupled Streptactin against the Strep tag. (E) 10 μM SUMO-CTT containing an

N-terminal Strep(II)-tag was incubated with 1 μM purified ribosomes and treated with 5 mM

or 25 mM EDC, as indicated. Samples were resolved by SDS-PAGE and analysed by western

blotting by simultaneously probing against SecA (red) and ribosomal protein uL23 (green).

Running positions of SUMO-CTT, L23 and crosslinking adducts between them (*) are

indicated at left.

Figure 2.

**Figure 2. Effect of C-terminal truncations on SecA *in vitro* and *in vivo*.** (A) 900 nM

Ru(bpy)$_2$(dcbpy)-labelled SecA (Wild type; squares), SecAΔMBD (ΔMBD; triangles) or

SecAΔCTT (ΔCTT; circles) was incubated in the presence of increasing concentrations of

purified 70S ribosomes. The equilibrium dissociation constant (K$_D$) of the complex was

determined by fitting the increase in fluorescence anisotropy from the Ru(bpy)$_2$(dcbpy)

(lines; **Table 1**). (B) 0.5 μM SecA, SecAΔMBD or SecAΔCTT was incubated in the absence

(lanes 1-3) of ribosomes, in the presence of 0.5 μM vacant 70S ribosomes (lanes 4-9) or in

the presence of 0.5 μM RNCs containing nascent SecM peptide (lanes 10-12). Where

indicated, binding reactions were incubated in the presence of 100 mM (lanes 1-6) or 250

mM (lanes 7-12) potassium acetate (KOAc). Binding reactions were layered on a 30%

sucrose cushion and ribosomes were sedimented through the sucrose cushion by

ultracentrifugation. Ribosomal pellets were resolved by SDS-PAGE and stained by

Coomassie. (C) 600 nM IAANS-VipB peptide was incubated with increasing concentrations

of SecA (Wild type; circles), SecAΔMBD (ΔMBD; triangles) or SecAΔCTT (ΔCTT;

squares). The $K_D$ for the SecA-peptide complex was determined by fitting the increase in

IAANS fluorescence upon binding to SecA (lines; **table 1**). (D) Growth of strains producing

SecA (DRH1119; bottom left), SecAΔMBD (DRH1120; bottom right) and SecAΔCTT

(DRH1121; top) on LB plates containing 100 μM IPTG.

Figure 3.

**Figure 3. Auto-crosslinking of the CTT to the catalytic core of SecA.** (A & B) 1 μM

SUMO-tagged SecA-biotin containing Bpa at position 852, 893 or 898 in the CTT was

incubated in the absence (-) or presence (+) of light (UV) at 350 nm. The protein samples

were resolved using SDS-PAGE and visualised by (A) western blotting against the C-

terminal biotin tag or (B) Coomassie staining. The running positions of full-length SUMO-

SecA and the internal crosslinking adduct (*) are indicated. (C) Mass spectra of wild-type

SecA-biotin (above, blue) and SecA[Bpa852]-biotin (below, red) in the region of 2450-2750 Da

region. Wild-type SecA-biotin and SecA[Bpa852]-biotin were exposed to light at 350 nm and

subsequently digested with trypsin. The masses of the tryptic fragments were determined

using MALDI-TOF. (D) Structure of SecA (2VDA (Gelis et al., 2007)). The main body of

the catalytic core is coloured blue, the PPXD is coloured cyan and the tryptic peptide that

crosslinks to position 852 (amino acids 361-382) is highlighted in orange. The structural

model was rendered using Chimera v. 1.12 (Pettersen et al., 2004).

Figure 4.

A

X-ray crystal structure of SecAΔMBD

PPXD

Protomer 1          Protomer 2

Fits of SAXS data

B

PPXD
"closed"

PPXD
"open"

Full-length SecA

C

PPXD

MBD

D

PPXD
"closed"

PPXD
"open"

SecAΔMBD

E

PPXD
"closed"

PPXD
"open"

SecAΔCTT

F

Distance between α-carbons
of amino acids 314 and 596

"closed"

Protomer 2 (Å)

"open"

Protomer 1 (Å)

| Marker | SecA | $\chi^2$ diagonal | p-value |
|---|---|---|---|
| ● | FL | 8.39 | 0.50 |
| ▲ | ΔMBD | 54.10 | <0.01 |
| ■ | ΔCTT | 22.14 | <0.01 |

**Figure 4. SAXS analysis of SecA truncation variants.** (A) X-ray crystal structure of

SecAΔMBD at 3.5Å solved by molecular replacement. The asymmetric unit (Protomer 1) is

coloured orange with the PPXD highlighted in cyan. The crystallographic mate (Protomer 2)

interacts with promoter 1 using an interface similar to that found in 2FSG (Papanikolau et al.,

2007). (B-E) Overlay of 10 independent structural models of SecA (B, C), SecAΔMBD (D)

and SecAΔCTT (E) generated from fitting to the SAXS data using CORAL. The main body

of the catalytic core is coloured grey, and the flexible residues are not displayed. (B, D, E) To

facilitate visualization of the asymmetry in the in the dimeric models, both protomeric

partners of the dimer were overlaid and the PPXD was coloured (blue/magenta) according to

the protomer. The MBD is not displayed in panel B. (C) To facilitate visualization of the

position of the MBD in the full-length protein, both protomeric partners of the dimer were

overlaid and the MBD of the dimer pair that was located nearest to position 596 of the

depicted protomer (orange) was displayed. In panel C, the PPXDs of two protomers, which

occupy the same space as the MBDs, are not displayed. (F) Plot of the position of the PPXD

in partners of the SecA dimer predicted by structural modelling. The distance between the α-

carbon of amino acid 314, which is located near the centroid of the PPXD, and amino acid

596 in NBD2 was determined for each protomer and plotted against the distance in the

second protomer. SecA, black circles (FL); SecAΔMBD, orange triangles (ΔMBD);

SecAΔCTT, blue squares (ΔCTT). The grey diagonal line indicates the position of the

distances if the dimers were symmetric. $\chi^2$ values to the diagonal were calculated and used to

determine p-values to test whether the asymmetry in the dimer was statistically significant.

**Figure 5. Site specific crosslinking of SecA to purified ribosomes and ribosome-nascent**

**chain complexes.** (A & B) Sites of incorporation of Bpa in the structure of *E. coli* SecA. (A)

Fit of the high resolution structure of SecA (PDB code 2VDA (Gelis et al., 2007)) and the

70S ribosome (PDB code 4V4Q (Schuwirth et al., 2005)) to the cryoEM structure of the

SecA ribosome complex (EMD-2565 (Singh et al., 2014)). (B) View of SecA from the

ribosome-interaction surface. Amino acid positions where Bpa was incorporated are

represented in space fill (yellow). Positions that crosslink to ribosomal proteins are coloured

red. The locations of the N-terminal α-helix of SecA and of ribosomal proteins uL23 (dark

blue), uL29 (purple) and uL24 (cyan) are indicated. Structural models were rendered using

Chimera v. 1.12  (Pettersen et al., 2004). (C) Bpa-mediated photocrosslinking of SecA

variants to vacant 70S ribosomes. 1 μM purified ribosomes were incubated with 1 μM SecA

containing BpA at the indicated position and exposed to light at 350 nm (above) or incubated

in the dark. Crosslinking adducts consistent with the molecular weight of a covalent crosslink

to ribosomal proteins are indicated with red arrowheads. The positions of full-length SecA

782    and uncleaved SUMO-SecA protein are indicated to the right. (D) 1 μM SecA$^{Bpa399}$ or

783    SecAΔMBD$^{Bpa399}$ was incubated with 1 μM non-translating 70S ribosomes or 1 μM arrested

784    RNCs containing nascent SecM (SecM-RNCs) and exposed to light at 350 nm. The positions

785    of full-length SecA and the SecA-uL29 crosslinking adduct are indicated. In C & D, samples

786    were resolved using SDS-PAGE and probed by western blotting using anti-SecA antiserum.

**Figure 6. Diagram of the proposed mechanism for recognition of nascent substrate proteins by SecA.** (a) In solution, SecA occupies an autoinhibited conformation with the FLD bound stably in the substrate protein binding site and the PPXD in the open conformation. (b) Binding of both the catalytic core and the MBD to the ribosomal surface causes the PPXD to shift to the open conformation, which destabilises binding of the FLD and allows SecA to sample nascent polypeptides. (c) Binding to the nascent substrate protein displaces the FLD from the substrate protein binding site and the PPXD returns to the open conformation, stabilising this interaction. Binding to nascent substrate releases SecA from the ribosomal surface.

1
**Supplemental figure S1. Structural model of the catalytic core of SecA in the "closed"**

**conformation.** Structural model of *E. coli* SecA from PDB file 2VDA (Gelis et al., 2007) in

ribbon diagram. The model is coloured according to domains described in the main text.

NBD1, dark blue; NBD2 cyan; PPXD, light blue; HSD, red; HWD, orange. The side chains

of lysines 625 (K625) and 633 (K633), which were identified by Huber et al. (2011) to be

involved in ribosome binding, are depicted in space-fill. The N-terminal α-helix (aa1-38),

which was identified by Singh et al. (2014) to be involved in ribosome binding is coloured

green. The CTT is not resolvable in high-resolution structures of SecA and is therefore not

depicted.

1μM ribosomes:     -      +

5μM SUMO-MBD:     +      +

 SUMO-MBD

Streptactin-HRP

**Supplemental figure S2 (associated with figure 1). SUMO-MBD cosediments with**

**ribosomes.** The C-terminal 27 amino acids of SecA were fused to the C-terminus of Strep-

tagged SUMO (SUMO-MBD). 5 μM SUMO-MBD was incubated in the presence or absence

of 1 μM 70S ribosomes. Binding reactions were layered on a 30% sucrose cushion and

subjected to ultracentrifugation to sediment ribosomes. The pellet fractions were then

resolved by SDS-PAGE and analysed by western blotting against the Strep tag.

**Supplemental figure S3 (associated with table 1). CD spectra and thermal denaturation**

**plots of SecA, SecAΔMBD and SecAΔCTT.** (A) Far-UV circular dichroism (CD) spectra of

2 μM solutions of SecA, SecAΔMBD, and SecAΔCTT in 10mM potassium phosphate (pH

7.5). (B) Representative plot of the thermal denaturation of SecA as determined by CD

spectroscopy. The α-helical content of 2 μM solutions of SecA, SecAΔMBD, and

SecAΔCTT in 10mM potassium phosphate (pH 7.5) was determined by measuring molar

ellipticity at 222 nm while the temperature of the solution was raised from 30 to 50°C. The

$T_{MS}$ given in **Table 1** were determined by van't Hoff analysis.

**30** **Supplemental table S1. Data collection and refinement statistics for the crystal structure**

**31** **of SecAΔMBD.**

| | SecAΔMBD |
|---|---|
| **Wavelength** | 0.9795 |
| **Resolution range** | 47.5 - 3.5 (3.625 - 3.5)* |
| **Space group** | C 1 2 1 |
| **Unit cell** | 163.1 106.8 76.0 90 102.8 90 |
| **Total reflections** | 32115 (3197)* |
| **Unique reflections** | 16106 (1609)* |
| **Multiplicity** | 2.0 (2.0)* |
| **Completeness (%)** | 99.66 (99.81)* |
| **Mean I/sigma(I)** | 13.51 (1.51)* |
| **Wilson B-factor** | 132.20 |
| **R-merge** | 0.03651 (0.5532)* |
| **R-meas** | 0.05163 (0.7823)* |
| **R-pim** | 0.03651 (0.5532)* |
| **CC1/2** | 1 (0.672)* |
| **CC*** | 1 (0.897)* |
| **Reflections used in refinement** | 16106 (1609)* |
| **Reflections used for R-free** | 785 (89)* |
| **R-work** | 0.2316 (0.3407)* |
| **R-free** | 0.2927 (0.3891)* |
| **CC(work)** | 0.900 (0.645)* |
| **CC(free)** | 0.869 (0.405)* |

| | |
|---|---|
| **No. non-hydrogen atoms** | 6182 |
| **Protein residues** | 777 |
| **RMS(bonds/angles)** | 0.011/1.47 |
| **Ramachandran (%)** | 91.42/5.07/3.51 |
| favoured/allowed/outliers | |
| **Rotamer outliers (%)** | 3.64 |
| **Clashscore** | 14.32 |
| **Average B-factor** | 158.55 |
| **PDB** | 6GOX |

*Statistics for the highest-resolution shell are shown in parentheses.

**Supplemental figure S4 (associated with figure 4). SAXS analysis of the solution**

**structure of SecA, SecAΔMBD and SecAΔCTT.** X-ray scattering plots for SecA (black),

SecAΔMBD (red) and SecAΔCTT (green). The region of divergence between the three

SAXS traces in the mid-$q$ region is indicated (black arrow).

**Supplemental table S2. SAXS data collection and processing details for SecA,**

**SecAΔMBD and SecAΔCTT.**

| Sample Details | |
| --- | --- |

| **SecA** | |
| --- | --- |
| Organism | *Escherichia coli* |
| Source | *Escherichia coli* |
| UniProt sequence ID | P10408 (amino acids 1-901) |
| Extinction coefficient (A$_{280}$, 0.1% (w/v)) | 0.742 |
| M from chemical composition (Da) | 102022.99 (monomer) |
| Average C in merged data (mg/ml) | 0.5 |
| Solvent | 20mM HEPES, 100mM NaCl pH 8, 1mM TCEP |

| **SecAΔMBD** | |
| --- | --- |
| Organism | *Escherichia coli* |
| Source | *Escherichia coli* |
| UniProt sequence ID | P10408 (amino acids 1-880) |
| Extinction coefficient (A$_{280}$, 0.1% (w/v)) | 0.745 |
| M from chemical composition (Da) | 99665.28 (monomer) |
| Average C in merged data (mg/ml) | 0.5 |
| Solvent | 20mM HEPES, 100mM NaCl pH 8, 1mM TCEP |

**SecAΔCTT**

| | |
|---|---|
| Organism | *Escherichia coli* |
| Source | *Escherichia coli* |
| UniProt sequence ID | P10408 (amino acids 1-832) |
| Extinction coefficient ($A_{280}$, 0.1% (w/v)) | 0.787 |
| M from chemical composition (Da) | 94329.44 (monomer) |
| Average C in merged data (mg/ml) | 0.5 |
| Solvent | 20mM HEPES, 100mM NaCl pH 8, 1mM TCEP |

**SAXS Data collection parameters**

| | |
|---|---|
| **Instrument** | ESRF BM29 BioSAXS |
| **Wavelength (Å)** | 0.919 |
| **Beamsize (μm)** | 700 x 700 |
| **Detector distance (m)** | 2.867 |
| **Q measurement range ($\text{Å}^{-1}$)** | 0.0034-0.494 |
| **Absolute scaling method** | Comparison with scattering from 1mm pure $H_2O$ |
| **Monitoring for radiation damage** | Data frame-by-frame comparison |
| **Exposure time** | 1 s per frame |
| **Sample temperature (ºC)** | 20 |

**Software**

**SAXS data reduction**

| | |
|---|---|
| *I(q)* versus *q* | BsxCuBE |
| solvent subtraction, averaging and | PRIMUS (ATSAS 2.8.3) |

merging

| | |
|---|---|
| **Extinction coefficient estimate** | ProtParam |
| **Basic analyses: Guinier, P(r), $V_p$** | Primus (ATSAS 2.8.3), FoXS, ScÅtter |
| **Atomic structure modelling** | Gromacs, PyMOL 1.7 |
| **Three-dimensional graphic model representation** | PyMOL 1.7 |
| **SAXS Molecular weight** | SAXSMoW |

**Structural parameters**

**SecA**

Guinier analysis

| | |
|---|---|
| *I(0)* | $144.47 \pm 0.076$ |
| $R_g$ (nm) | $4.22 \pm 0.00$ |
| $q_{min}$ (nm$^{-1}$) | 0.0174 |
| $qR_g$ max ($q_{min}$ = nm$^{-1}$) | 1.29 |
| Coefficient of correlation, $R^2$ | 0.99 |
| M from *(0)* (ratio to predicted)* | 224.7  (10.1 % to dimer) |

**SecAΔMBD**

Guinier analysis

| | |
|---|---|
| *I(0)* | $160.86 \pm 0.045$ |
| $R_g$ (nm) | $4.15 \pm 0.045$ |
| $q_{min}$ (nm$^{-1}$) | 0.0129 |
| $qR_g$ max ($q_{min}$ = nm$^{-1}$) | 1.29 |
| Coefficient of correlation, $R^2$ | 0.92 |
| M from *(0)* (% discrepancy)* | 210.8 (5.73% to dimer) |

**SecAΔCTT**

Guinier analysis

| | |
|---|---|
| *I(0)* | $53.41 \pm 0.086$ |
| $R_g$ (nm) | $4.51 \pm 0.01$ |
| $q_{min}$ (nm$^{-1}$) | 0.0304 |
| $qR_g$ max ($q_{min}$ = nm$^{-1}$) | 1.28 |
| Coefficient of correlation, $R^2$ | 0.95 |
| M from *(0)* (ratio to predicted)* | 232.3 (23.1 % to dimer) |

44
45

46    **Supplemental table S3. Fitting parameters of models of the *E. coli* SecA dimer.**

| Model[a] | $\chi^2$ | $c_1$[b] | $c_2$[c] |
|:---:|:---:|:---:|:---:|
| 2FSG | 3.66 | 0.99 | 3.36 |
| 2IBM | 14.99 | 1.05 | 4.00* |
| 2IPC | 24.08 | 1.05 | 0.51 |
| 1M6N | 30.81 | 1.05 | 4.00* |
| 1NL3 | 25.01 | 1.05 | 1.73 |
| 6GOX | 5.25 | 1.03 | 1.38 |

[a]SecA template structures used to generate models of the *E. coli* SecA homodimer

[b]Variable for scaling of the atomic radius, which also controls the excluded volume of the

molecule (permitted range: 0.99 - 1.05)

[c]Variable used to adjust the difference between the densities of the hydration layer and the

bulk water (permitted range: -2.0 - 4.0)

*Fitting reached limit of permitted range, which suggests overfitting of the data.

**Supplemental figure S5 (associated with figure 5). Binding of SecA to the ribosome.** 1

μM SecA[Bpa399] was incubated with non-translating 70S ribosomes (vacant) or RNCs

containing arrested nascent SecM (SecM-RNCs) or maltose binding protein (MBP-RNCs).

Where indicated, samples were exposed to light at 350 nm (UV). Samples were then resolved

using SDS-PAGE and probed by western blotting against SecA. The running positions of

SecA and the crosslinking adduct between SecA and ribosomal protein uL29 are indicated.

65