A validation of the diathesis-stress model for depression in Generation Scotland.

Running title: **The diathesis-stress model for depression in Generation Scotland.**

Aleix Arnau-Soler[1], Mark J. Adams[2], Toni-Kim Clarke[2], Donald J. MacIntyre[2], Keith Milburn[3], Lauren Navrady[2], Generation Scotland[4], Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium[5], Caroline Hayward[6], Andrew McIntosh[2,7], Pippa A. Thomson[1,7]

[1] Medical Genetics Section, University of Edinburgh, Centre for Genomic and Experimental Medicine and MRC Institute of Genetics and Molecular Medicine, Edinburgh, UK.
[2] Division of Psychiatry, Deanery of Clinical Sciences, University of Edinburgh, Royal Edinburgh Hospital, Morningside Park, Edinburgh EH10 5HF, UK.
[3] Health Informatics Centre, University of Dundee, UK.
[4] A collaboration between the University Medical School and NHS in Aberdeen, Dundee, Edinburgh and Glasgow, Scotland, UK.
[5] For a full list of MDD working group of the PGC investigators, see the Supplementary Material
[6] Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK.
[7] Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK.

Correspondence should be addressed to: P. Thomson (Pippa.Thomson@ed.ac.uk) or A. Arnau-Soler (aleix.arnau.soler@igmm.ed.ac.uk)

Centre for Genomic and Experimental Medicine
Institute of Genetics and Molecular Medicine
University of Edinburgh
Western General Hospital
Crewe Road South
Edinburgh, EH4 2XU, UK
Phone: +44(0)1316518756

Abstract: 250
Main body (excl. abstract): 4 730
No. Tables: 1
No. Figures: 3
No. References: 73
Supplementary Information: 1 file
       Supplementary Tables: 1
       Supplementary Figures: 2

ABSTRACT

Depression has well-established influences from genetic and environmental risk

factors. This has led to the *diathesis-stress* theory, which assumes a multiplicative

gene-by-environment interaction (GxE) effect on risk. Recently, *Colodro-Conde et al.*

empirically tested this theory, using the polygenic risk score for major depressive

disorder (PRS, genes) and stressful life events (SLE, environment) effects on depressive

symptoms, identifying significant GxE effects with an additive contribution to liability.

We have tested the *diathesis-stress* theory on an independent sample of 4 919

individuals.

We identified nominally significant positive GxE effects in the full cohort (R^2 = 0.08%, *p*

= 0.049) and in women (R^2 = 0.19%, *p* = 0.017), but not in men (R^2 = 0.15%, *p* = 0.07).

GxE effects were nominally significant, but only in women, when SLE were split into

those in which the respondent plays an active or passive role (R^2 = 0.15%, *p* = 0.038; R^2

= 0.16%, *p* = 0.033, respectively). High PRS increased the risk of depression in

participants reporting high numbers of SLE (*p* = 2.86 x 10^{-4}). However, in those

participants who reported no recent SLE, a higher PRS appeared to increase the risk of

depressive symptoms in men (β = 0.082, *p* = 0.016) but had a protective effect in

women (β = -0.061, *p* = 0.037). This difference was nominally significant (*p* = 0.017).

Our study reinforces the evidence of additional risk in the aetiology of depression due

to GxE effects. However, larger sample sizes are required to robustly validate these

findings.

INTRODUCTION

Stressful life events (SLE) have been consistently recognized as a determinant of depressive symptoms, with many studies reporting significant associations between SLE and major depressive disorder (MDD)[1-7]. Some studies suggest that severe adversity is present before the onset of illness in over 50% of individuals with depression[8] and may characterize a subtype of cases[9]. However, some individuals facing severe stress never present symptoms of depression[10]. This has led to a suggestion that the interaction between stress and an individual's vulnerability, or *diathesis*, is a key element in the development of depressive symptoms. Such vulnerability can be conceived as a set of biological factors that predispose to illness. Several *diathesis-stress* models have been successfully applied across many psychopathologies[11-15].

The *diathesis-stress* model proposes that a latent *diathesis* may be activated by stress before psychopathological symptoms manifest. Some levels of *diathesis* to illness are present in everybody, with a threshold over which symptoms will appear. Exceeding such a threshold depends on the interaction between *diathesis* and the degree of adversity faced in SLE, which increases the liability to depression beyond the combined additive effects of the *diathesis* and stress alone[11]. Genetic risk factors can, therefore, be conceived as a genetic *diathesis*. Thus, this genetically driven effect produced by the *diathesis-stress* interaction can be seen as a gene-by-environment interaction (GxE).

MDD is characterized by a highly polygenic architecture, composed of common variants with small effect and/or rare variants[16]. Therefore, interactions in depression are also expected to be highly polygenic. In recent years, with the increasing success of

genome-wide association studies, GxE studies in depression have shifted towards

hypothesis-free genome-wide and polygenic approaches that capture liability to

depression using genetic data[17-25]. Recent advances in genomics and the massive effort

from national institutions to collect genetic, clinical and environmental data on large

population-based samples now provide an opportunity to empirically test the

diathesis-stress model for depression. The construction of polygenic risk scores (PRS)

offers a novel paradigm to quantify genetic *diathesis* into a single genetic measure,

allowing us to study GxE effects with more predictive power than any single variant[26-29].

PRS are genetic indicators of the aggregate number of risk alleles carried by an

individual weighted by their allelic effect estimated from genome-wide association

studies. This polygenic approach to assessing the *diathesis-stress* model for depression

has been tested using either childhood trauma[17,19,25] or adult SLE[18,23,25] as measures of

environmental adversity.

Recently, Colodro-Conde *et al.*[23] provided a direct test of the *diathesis-stress* model for

recent SLE and depressive symptoms. In this study, Colodro-Conde *et al.* used PRS

weighted by the most recent genome-wide meta-analysis conducted by the Psychiatric

Genetics Consortium (PGC; N = 159 601), and measures of three environmental

exposures: lack of social support, "personal" SLE, and "network" SLE. Colodro-Conde

et al. reported a significant additive risk on liability to depression due to a GxE effect in

individuals who combine a high genetic predisposition to MDD and a high number of

reported "personal" SLE, mainly driven by effects in women. A significant effect of

interaction was not detected in males. They found no significant interaction between

the genetic *diathesis* and "network" SLE or social support. They concluded that the

effect of stress on risk of depression was dependent on an individual's *diathesis,* thus

supporting the *diathesis-stress* theory. In addition, they suggested possible sex-specific

differences in the aetiology of depression. However, Colodro-Conde *et al.* findings

have not, to our knowledge, been independently validated.

In the present study we aim to test the *diathesis-stress* model in an independent

sample of 4 919 unrelated white British participants from a further longitudinal follow-

up from Generation Scotland and assess the differences between women and men,

using self-reported depressive symptoms and recent SLE.

MATERIALS AND METHODS

Sample description

Generation Scotland is a family-based population cohort recruited throughout

Scotland by a cross-disciplinary collaboration of Scottish medical schools and the

National Health Service (NHS) between 2006 and 2011[30]. At baseline, blood and

salivary DNA samples from Generation Scotland participants were collected, stored

and genotyped at the Wellcome Trust Clinical Research Facility, Edinburgh. Genome-

wide genotype data were generated using the Illumina HumanOmniExpressExome-8

v1.0 DNA Analysis BeadChip (San Diego, CA, USA) and Infinium chemistry[31]. The

procedures and further details for DNA extraction and genotyping have been

extensively described elsewhere[32,33]. In 2014, 21 525 participants from Generation

Scotland eligible for re-contact were sent self-reported questionnaires as part of a

further longitudinal assessment funded by a Wellcome Trust Strategic Award

"STratifying Resilience and Depression Longitudinally" (STRADL)[34] to collect new and

updated mental health questionnaires including psychiatric symptoms and SLE

measures. 9 618 re-contacted participants from Generation Scotland agreed to provide

new measures to the mental health follow-up[34] (44.7% response rate). Duplicate

samples, those showing sex discrepancies with phenotypic data, or that had more than

2% missing genotype data, were removed from the sample, as were samples identified

as population outliers in principal component analysis (mainly non-Caucasians and

Italian ancestry subgroup). In addition, individuals with diagnoses of bipolar disorder,

or with missing SLE data, were excluded from the analyses. SNPs with more than 2% of

genotypes missing, Hardy-Weinberg Equilibrium test $p < 1 \times 10^{-6}$, or a minor allele

frequency lower than 1%, were excluded. Individuals were then filtered by degree of

relatedness (pi-hat < 0.05) using PLINK v1.9[35], maximizing retention of those

participants reporting higher numbers of SLE (see phenotype assessment below). After

quality control, the final dataset comprised 4 919 unrelated individuals of European

ancestry and 560 351 SNPs (mean age at questionnaire: 57.2, s.d. = 12.2, range 22-95;

women: n = 2 990 - 60.8%, mean age 56.1, s.d. = 12.4; *men*: n = 1 929 - 39.2%, mean

age 58.7, s.d. = 11.8). Further details on the recruitment procedure and Generation

Scotland profile are described in detail elsewhere[30,32,36-38]. All participants provided

written consent. All components of Generation Scotland and STRADL obtained ethical

approval from the Tayside Committee on Medical Research Ethics on behalf of the

National Health Service (reference 05/s1401/89). Generation Scotland data is available

to researchers on application to the Generation Scotland Access Committee

(access@generationscotland.org).

Phenotype assessment

Participant self-reported current depressive symptoms through the 28-item scaled

version of The General Health Questionnaire[39,40]. The General Health Questionnaire is

a reliable and validated psychometric screening tool to detect common psychiatric and

non-psychotic conditions (General Health Questionnaire Cronbach alpha coefficient:

0.82 – 0.86)[41]. This consists of 28 items designed to identify whether an individual's

current mental state has changed over the last 2 weeks from their typical state. The

questionnaire captures core symptoms of depression through subscales for severe

depression, emotional (e.g. anxiety and social dysfunction) and somatic symptoms

linked to depression. These subscales are highly correlated[42] and suggest an overall

general factor of depression[43]. Participants rated the 28 items on a four-point Likert

scale from 0 to 3 to assess its degree or severity[41] (e.g., *Have you recently felt that life*

is entirely hopeless? "Not at all", "No more than usual", "Rather more than usual",

"Much more than usual"), resulting on an 84-point scale depression score. The Likert

scale, which provides a wider and smoother distribution[41], could be more sensitive to

detect changes in mental status in those participants with chronic conditions or

chronic stress who may feel their current symptoms as "usual"[44], and to detect

psychopathology changes as response to stress. The final depression score was log

transformed to reduce the effect of positive skew and provide a better approximation

to a normal distribution. In addition, participants completed the Composite

International Diagnostic Interview–Short Form, which diagnoses lifetime history of

MDD according to DSM-IV criteria[45]. The depression score predicted lifetime history of

MDD (odd ratio = 1.91, 95% confidence intervals 1.80-2.02, p = 1.55 x 10^{-102}, N = 8

994), with a 3.8-fold increased odd of having a lifetime history of MDD between

participants in the top and bottom deciles, thus supporting the usefulness of the

depression score in understanding MDD. For a better interpretation, we scaled the

depression score to a mean of 0 when required (Figure 3).

Data from a self-reported questionnaire based on the List of Threating Experiences[46]

was used to construct a measure of common SLE over the previous 6 months. The List

of Threatening Experiences is a reliable psychometric device to measure psychological

"stress"[47,48]. It consists of a 12-item questionnaire to assess SLE with considerable

long-term contextual effects (e.g., *Over last 6 months, did you have a serious problem*

with a close friend, neighbour or relatives?). A final score reflecting the total number of

SLE (TSLE) ranging from 0 to 12 was constructed by summing the "yes" responses.

Additionally, TSLE was split into two categories based on those items measuring SLE in

which the individual may play and active role exposure to SLE, and therefore in which

the SLE is influenced by genetic factors and thus subject to be "dependent" on an

individual's own behaviour or symptoms (DSLE; 6 items, e.g., *a serious problem with a*

close friend, neighbour or relatives may be subject to a respondent's own behaviour),

or SLE that are not influenced by genetic factors, likely to be "independent" on a

participant's own behaviour (ISLE; 5 items, e.g., *a serious illness, injury or assault*

happening to a close relative is potentially independent of a respondent's own

behaviour)[46,49]. The item "*Did you/your wife or partner give birth?*" was excluded from

this categorization. In addition, SLE reported were categorized to investigate the

diathesis effect at different levels of exposure, including a group to test the *diathesis*

effect when SLE is not reported. 3 levels of SLE reported were defined (0 SLE = "none",

1 or 2 SLE = "low", and 3 or more SLE = "high") to retain a large enough sample size for

each group to allow meaningful statistical comparison.

Polygenic profiling & statistical analysis

Polygenic risk scores (PRS) were generated by PRSice[50], whose functionality relies

mostly on PLINK v1.9[35], and were calculated using the genotype data of Generation

Scotland participants (i.e. target sample) and summary statistics for MDD from the

PGC-MDD2 GWAS release (July 2016, discovery sample) used by Colodro-Conde *et*

al.[23], with the added contribution from QIMR cohort and the exclusion of Generation

Scotland participants, resulting in summary statistics for MDD derived from a sample

of 50 455 cases and 105 411 controls.

Briefly, PRSice removed strand-ambiguous SNPs and clump-based pruned ($r^2 = 0.1$,

within a 10Mb window) our target sample to obtain the most significant independent

SNPs in approximate linkage equilibrium. Independent risk alleles were then weighted

by the allelic effect sizes estimated in the independent discovery sample and

aggregated into PRS. PRS were generated for eight p thresholds (p thresholds: $< 5 \times 10^{-8}$

, $< 1 \times 10^{-5}$, < 0.001, < 0.01, < 0.05, < 0.1, < 0.5, $<=1$) determined by the discovery

sample and standardized (See Supplementary Table 1 for summary of PRS).

A genetic relationship matrix (GRM) was calculated for each dataset (i.e. *full cohort*,

women and *men*) using GCTA1.26.0[51]. Mixed linear models using the GRM were used

to estimate the variance in depression score explained by PRS, SLEs and their

interaction; and stratified by sex. 20 principal components were calculated for the

datasets.

The mixed linear model used to assess the effects of PRS is as follows:

$$Depression = \beta_0 + \beta_1 PRS + GRM + Covariates$$

225   Mixed linear models used to assess the effect of the stressors are as follows:

$$Depression = \beta_0 + \beta_1 TSLE + GRM + Covariates$$

$$Depression = \beta_0 + \beta_1 DSLE + GRM + Covariates$$

$$Depression = \beta_0 + \beta_1 ISLE + GRM + Covariates$$

Following Colodro-Conde *et al.*[23], covariates (i.e. age, age^2, sex, age-by-sex and age^2-

by-sex interactions, and 20 principal components) were regressed from PRS (PRS') and

SLE scores (i.e. TSLE', DSLE' and ISLE'; SLEs') before fitting models in GCTA to guard

against confounding influences on the PRS-by-SLEs interactions[52]. PRS' and SLEs' were

standardized to a mean of 0 and a standard deviation of 1. The Mixed linear models

(i.e. the *diathesis-stress* model) used to assess GxE effects are as follows:

$$Depression = \beta_0 + \beta_1 PRS' + \beta_2 TSLE' + \beta_3 PRS'xTSLE' + GRM + Covariates$$

$$Depression = \beta_0 + \beta_1 PRS' + \beta_2 DSLE' + \beta_3 PRS'xDSLE' + GRM + Covariates$$

$$Depression = \beta_0 + \beta_1 PRS' + \beta_2 ISLE' + \beta_3 PRS' xISLE' + GRM + Covariates$$

Covariates fitted in the models above were age, age^2, sex, age-by-sex, age^2-by-sex and

20 principal components. Sex and its interactions (age-by-sex and age^2-by-sex) were

omitted from the covariates when stratifying by sex. All parameters from the models

were estimated using GCTA and the significance of the effect (β) from fixed effects

assessed using a Wald test. The significance of main effects (PRS and SLEs) allowed for

nominally testing the significance of interactions at p-threshold = 0.05. To account for

multiple testing correction, a Bonferroni's adjustment correcting for 8 PRS and 3

measures of SLE tested (24 tests) was used to establish a robust threshold for

significance at $p = 2.08 \times 10^{-3}$.

The PRS effect on depression score at different levels of exposure was further

examined for the detected nominally significant interactions by categorizing

participants on three groups based on the number of SLE reported (i.e. "none", "low"

or "high"). Using linear regression, we applied a least squares approach to assess PRS'

effects on the depression score in each SLE category. Further conservative Bonferroni

correction to adjust for the 3 SLE categories tested established a threshold for

significance of $p = 6.94 \times 10^{-4}$.

Differences on the estimated size of GxE effect between women and men were

assessed by comparing a z-score to the standard normal distribution (α = 0.05, one-

tailed). Z-scores were derived from GxE estimates (β) and standard errors (SE)

detected in women and men as follows:

$$Z - score = \frac{\beta_{women} - \beta_{men}}{\sqrt{SE(\beta_{women)}^2 + SE(\beta_{men})^2}}$$

RESULTS

PRS for MDD significantly predicted the depression score across the whole sample (β =

0.080, s.e. = 0.014, p = 7.53 x 10^{-9}) explaining 0.64% of the variance at its best p-

threshold (p-threshold = 0.1; Figure 1a). Stratifying by sex, PRS significantly predicted

the depression score in both sexes, explaining 0.59% in men and 0.67% in women

(*men*: p-threshold = 0.1, β = 0.077, s.e. = 0.022, p = 2.09 x 10^{-4}; *women*: p-threshold =

0.1, β = 0.082, s.e. = 0.018, p = 4.93 x 10^{-6}; Figure 1a). Self-reported SLE over the last 6

months (TSLE, mean = 1.3 SLE, s.d. = 1.5) also significantly predicted depression score

for the whole sample and stratified by sex (*full cohort*: variance explained = 4.91%, β =

0.222, s.e. = 0.014, p = 9.98 x 10^{-59}; *men*: 4.19%, β = 0.205, s.e. = 0.021, p = 2.23 x 10^{-22};

women: 5.33%, β = 0.231, s.e. = 0.018, p = 7.48 x 10^{-38}; Figure 1b). Overall, significant

additive contributions from genetics and SLE in depression score were detected in all

participants and across sexes. There was no significant difference in the direct effect of

TSLE between women and men (p = 0.17). However, the variance in depression score

explained by the TSLE appeared to be lower than the variance explained by the

measure of personal SLE (PSLE) used in Colodro-Conde *et al.*[23] (12.9%). This may, in

part, be explained by different contributions of dependent and independent SLE items

screened in Colodro-Conde *et al.* compared to our study. Although questions about

dependent SLE (DSLE, mean = 0.4 SLE) represented over 28% of the TSLE-items

reported in our study, the main effect of DSLE explained approximately 93% of the

amount of variance explained by TSLE (*full cohort*: variance explained = 4.56%, β =

0.212, s.e. = 0.014, p = 1.73 x 10^{-54}; *men*: 3.74%, β = 0.193, s.e. = 0.021, p = 9.66 x 10^{-21};

women: 5.07%, β = 0.225, s.e. = 0.018, p = 8.09 x 10^{-35}; Figure 1b). Independent SLE

(ISLE, mean = 0.85 SLE), which represented over 69% of TSLE-items, explained

approximately 57% of the amount of variance explained by TSLE (*full cohort*: variance

explained = 2.80%, β = 0.167, s.e. = 0.014, *p* = 1.32 x 10^{-33}; *men*: 2.44%, β = 0.156, s.e. =

0.022, *p* = 2.88 x 10^{-13}; *women*: 3.02%, β = 0.174, s.e. = 0.018, *p* = 5.20 x 10^{-22}; Figure

1b). To explore the contribution from each measure, we combined DSLE and ISLE

together in a single model. DSLE explained 3.34% of the variance of depressive score

compared to 1.45% of the variance being explained by ISLE, suggesting that DSLE have

a greater effect on liability to depressive symptoms than ISLE.

A *diathesis-stress* model for depression was tested to assess GxE effects. We detected

significant, albeit weak, GxE effects on depression score (Figure 2). The PRS interaction

with TSLE was nominally significant in the full cohort (β = 0.028, s.e. = 0.014, R^2 =

0.08%, *p* = 0.049) and slightly stronger in women (β = 0.044, s.e. = 0.018, R^2 = 0.19%, *p*

= 0.017; Figure 2a), compared to men in which the effect was not significant (β =

0.039, s.e. = 0.022, R^2 = 0.15%, *p* = 0.07). However, these results did not survive

correction for multiple testing (*p* > 2.08 x 10^{-3}).

The best-fit threshold was much lower in women (*p*-threshold = 1 x 10^{-5}) compared to

the full sample (*p*-threshold = 0.01). The size of GxE across sexes at *p*-threshold = 1 x

10^{-5} were significantly different (GxE*sex *p* = 0.017), but not at the best p-threshold in

the full cohort (*p*-threshold = 0.01, GxE*sex *p* = 0.32; Figure 2a). In women, GxE effect

with DSLE predicted depression score (*p*-threshold = 1 x 10^{-5}; β = 0.039, s.e. = 0.019, R^2

= 0.15%, *p* = 0.038; Figure 2b and Supplementary Figure 2a), as did the GxE effect with

ISLE (*p*-threshold = 1 x 10^{-5}; β = 0.040, s.e. = 0.019, R^2 = 0.16%, *p* = 0.033; Figure 2c and

Supplementary Figure 2b). No significant interaction was detected in men (best-fit *p*-

threshold = 0.1) with either TSLE (β = 0.039, s.e. = 0.022, R^2 = 0.15%, *p* = 0.072; Figure

300 2a), DSLE (β = 0.024, s.e. = 0.022, R^2 = 0.06%, p = 0.28; Figure 2b) or ISLE (β = 0.043, s.e.

301 = 0.022, R^2 = 0.18%, p = 0.055; Figure 2c).

302 To examine these results further and investigate the *diathesis* effect at different levels

303 of stress, nominally significant GxE were plotted between PRS and categories of SLE

304 (i.e, "none", "low" and "high" SLE reported; Figure 3). Examining the interaction found

305 at the full cohort (PRS at PGC-MDD GWAS p-threshold = 0.01), we detected a

306 significant direct *diathesis* effect on the risk of depressive symptoms in those

307 participants reporting SLE, with a higher risk when greater numbers of SLE were

308 reported ("low" number of SLE reported: PRS' β = 0.043, s.e. = 0.021, p = 0.039; "high"

309 number of SLE reported: PRS' β = 0.142, s.e. = 0.039, p = 2.86 x 10^{-4}; see Table 1 and

310 Figure 3a). Whereas, in participants who reported no SLE over the preceding 6 months,

311 the risk of depressive symptoms was the same regardless of their *diathesis* risk

312 ("none" SLE reported: PRS' β = 0.021, s.e. = 0.022, p = 0.339). Stratifying these results

313 by sex, we found the same pattern as in the full cohort in women ("none": p = 0.687;

314 "low": p = 0.023; "high": p = 2 x 10^{-3}), but not in men ("none": p = 0.307; "low": p =

315 728; "high": p = 0.053; see Table 1 and Figure 3a). However, the lack of significant

316 *diathesis* effect in men may be due to their lower sample size and its corresponding

317 reduced power.

318 Examining the interaction with PRS (at PGC-MDD GWAS p-threshold = 1 x 10^{-5}) with

319 which a significant interaction was detected in women, we only detected a significant

320 *diathesis* effect on depression score when stratifying by sex in those participants who

321 did not reported SLE over the last 6 months (see Table 1). The *diathesis* effect was

322 positive in men (PRS' β = 0.082, s.e. = 0.034, p = 0.016, R^2 = 0.7%; Figure 3b), consistent

323 with the contribution of risk alleles. Conversely, the *diathesis* effect was negative in

women (PRS' β = -0.061, s.e. = 0.029, p = 0.037, R^2 = 0.4%; Figure 3b), suggesting a

protective effect of increasing PRS in those women reporting no SLE, and consistent

with the contribution of alleles to individual sensitivity to both positive and negative

environmental effects (i.e. "plasticity alleles" rather than "risk alleles")[53,54]. This PRS

accounted for the effect of just 34 SNPs, and the size of its GxE across sexes were

significantly different (GxE*sex p = 0.017; Figure 2a), supporting possible differences in

the underlying stress-response mechanisms between women and men.

DISCUSSION

The findings reported in this study support those from Colodro-Conde *et al.*[23], in an independent sample of similar sample size and study design, and also supports possible sex-specific differences in the effect of genetic risk of MDD in response to SLE. Both Colodro-Conde *et al.* and our study suggest that individuals with an inherent genetic predisposition to MDD, reporting high number of recent SLE, are at additional risk of depressive symptoms due to GxE effects, thus validating the *diathesis-stress* theory. We identified nominally significant GxE effects in liability to depression at the population level ($p = 0.049$) and in women ($p = 0.017$), but not in men ($p = 0.072$). However, these interactions did not survive multiple testing correction ($p > 2.08 \times 10^{-3}$) and the power of both studies to draw robust conclusions remains limited[55]. With increased power these studies could determine more accurately both the presence and magnitude of a GxE effect in depression. To better understand the effect of PRS at different levels of exposure to stress, we examined the nominally significant interactions detected in the full sample by categorizing participants on three groups based on the number of SLE reported (i.e. "none", "low" or "high"). We detected a significant *diathesis* effect on risk of depression only in those participants reporting SLE, but not in those participants that reported no SLE over the last 6 month. Furthermore, the *diathesis* effect was stronger on those participants reporting a "high" number of SLE ($\beta = 0.142$, $p = 2.86 \times 10^{-4}$) compared to those participants reporting a "low" number of SLE ($\beta = 0.043$, $p = 0.039$). The former effect was robustly significant and survived a conservative Bonferroni correction to adjust for multiple testing ($p < 6.94 \times 10^{-4}$). This finding corroborates the *diathesis-stress* model for depression and supports Colodro-Conde *et al.* results using an independent sample.

To investigate the relative contribution of the GxE to the variance of depression, we

examined in the full cohort the total variance of depression score explained by the PRS

main effect and the significant GxE effect jointly. Together, they explained 0.34% of

the variance, of which 0.07% of the variance of the depression score was attributed to

the GxE effect (p-threshold = 0.01; PRS p = 1.19 x 10^{-4}, GxE p = 0.049; both derived

from the full diathesis-model with TSLE). This is lower than the proportion of variance

attributed to common SNPs (8.9%) in the full PGC-MDD analysis[16]. As Colodro-Conde *et*

al. noted, this result aligns with estimates from experimental organisms suggesting

that around 20% of the heritability may be typically attributed to the effects of GxE[56],

although it is inconsistent with the majority of human traits with the potential

exception of depression[57].

Consistent with PRS predicting "personal" SLE in Colodro-Conde *et al.,* PRS for MDD

predicted SLE in our study (see Supplementary Figure 1), although not at the *p*-

threshold at which significant GxE effects were detected. Genetic factors predisposing

to MDD may contribute to individuals exposing themselves to, or showing an increased

reporting of, SLE via behavioural or personality traits[58,59]. Such genetic mediation of

the association between depression and SLE would disclose a gene-environment

correlation (i.e. genetic effects on the probability of undergoing a SLE) that hinders to

interpret our findings as pure GxE effects[60,61]. To address this limitation and assess this

aspect, following Colodro-Conde *et al.*, we split the 12-items TSLE measure into SLE

that are either potentially "dependent" on a participant's own behaviour (DSLE;

therefore, potentially driven by genetic factors) or not ("independent" SLE; ISLE)[46,49].

DSLE are reported to be more heritable and have stronger associations with MDD than

ISLE[49,62,63]. In our sample, reporting DSLE is significantly heritable (h^2_{SNP} = 0.131, s.e. =

0.071, p = 0.029), supporting a genetic mediation of the association, whereas reporting

ISLE is not significantly heritable (h^2_{SNP} = 0.000, s.e. = 0.072, p = 0.5). Nominally

significant GxE effects were seen in women for both DSLE and ISLE, suggesting that

both GxE and gene-environment correlation co-occur. Colodro-Conde *et al.* did not

identify significant GxE using independent SLE as the exposure.

Between-sex differences on stress response could help to explain previous differences

seen between sexes in depression such as those in associated risk (i.e. approximately

1.5 - 2-fold higher in women), symptoms reported and/or coping strategies (e.g.,

whereas women tend to cope through verbal and emotional strategies, men tend to

cope by doing sport and consuming alcohol)[64-68]. This also aligns with an increased risk

associated with a lack of social support seen in women compared to men[23].

Furthermore, although we do not know whether participants experienced recent

events with positive effects, we saw a protective effect in those women who did not

experienced recent SLE (p = 0.037), suggesting that some genetic variants associated

with MDD may operate as "plasticity alleles" and not just as "risk alleles"[53,54]. This

effect was neutralized in the full cohort due to an opposite effect in men (p = 0.016),

but it is supported by previous protective effects reported when using a serotoninergic

multilocus profile score and absence of SLE in young women[69]. These findings would

be consistent with a differential-susceptibility model[70,71] of depression, also suggested

by the interaction effects seen between the serotonin transporter linked promoter

region gene (5-HTTLPR) locus and family support and liability to adolescent depression

in boys[72]. However, our results and the examples given are only nominally significant

and will require replication in larger samples. Robustly identified sex-specific

differences in genetic stress-response could improve personalized treatments and

therapies such as better coping strategies.

There are notable differences between our study and Colodro-Conde *et al.* to consider

before accepting our findings as a replication of Colodro-Conde *et al.* results. First,

differences in PRS profiling may have affected replication power. We used the same

equivalent PGC-MDD2 GWAS as discovery sample. However, whereas Colodro-Conde

et al. generated PRS in their target sample containing over 9.5M imputed SNP, in this

study we generated PRS in a target sample of over 560K genotyped SNPs (see

Supplementary table 1 for comparison). This potentially results in a less informative

PRS in our study, with less predictive power, although the variance explained by our

PRS was slightly larger (0.64% vs. 0.46%). The size of the discovery sample is key to

constructing an accurate predictive PRS, but to exploit the most of the variants

available may be an asset[55]. Secondly, different screening tools were used to measure

both current depression and recent environmental stressors across the two studies.

Both studies transformed their data, using item response theory or by log-

transformation, to improve the data distribution. However, neither study used

depression scores that were normally distributed. The scale of the instruments used

and their corresponding parameterization to test an interaction could have a direct

effect on the size and significance of their interaction[56,73]; so findings from GxE must

be taken with caution. Furthermore, although both screening methods have been

validated and applied to detect depressive symptoms, different questions may cover

and emphasise different features of the illness, which may result in different outputs.

The same applies to the measurement of environmental stressors in the two studies.

Both covering of a longer time-period and upweighting by "dependent" SLE items may

explain the increased explanatory power of "personal" SLE (12.9%) in Colodro-Conde

et al. to predict depression score compared to our "total" SLE measure (4.91%). Finally,

the unmeasured aspects of the exposure to SLE or its impact may also contribute to

lack of stronger replication and positive findings.

In conclusion, despite differences in the measures used across studies, we saw

concordance and similar patterns between our results and those of Colodro-Conde *et*

al.[23]. Our findings are consistent with Colodro-Conde *et al.* and, therefore, add validity

to the *diathesis-stress* theory for depression. Empirically demonstrating the *diathesis-*

stress theory for depression would validate recent[20-22,24] and future studies using a

genome-wide approach to identify genetic mechanisms and interactive pathways

involved in GxE underpinning the causative effect of "stress" in the development of

depressive symptoms and mental illness in general. This study adds to our

understanding of gene-by-environment interactions, although larger samples will be

required to confirm differences in *diathesis-stress* effects between women and men.

440 **ACKNOWLEDGMENTS**

441 The 1[st] author AAS is funded by the University of Edinburgh (www.ed.ac.uk) and MRC

442 for his PhD study at the University of Edinburgh Institute of Genetics and Molecular

443 Medicine (www.ed.ac.uk/igmm). DJM acknowledges the financial support of NHS

444 Research Scotland (NRS) through NHS Lothian. MA is supported by STRADL through a

445 Wellcome Trust Strategic Award (reference 104036/Z/14/Z). Generation Scotland

446 received core support from the Chief Scientist Office of the Scottish Government

447 Health Directorates [CZD/16/6] and the Scottish Funding Council [HR03006]. The

448 genotyping of the GS:SFHS samples was carried out by the Genetics Core Laboratory at

449 the Wellcome Trust Clinical Research Facility, Edinburgh, Scotland and was funded by

450 the Medical Research Council UK and the Wellcome Trust (Wellcome Trust Strategic

451 Award "STratifying Resilience and Depression Longitudinally" (STRADL) Reference

452 104036/Z/14/Z). The Major Depressive Disorder Working Group of the Psychiatric

453 Genomics Consortium depends on the contributions of many parties.

454

455 **FINANCIAL DISCLOSURE**

456 The authors declare no conflict of interest.

REFERENCES

1.   Hammen, C. Stress and depression. *Annu Rev Clin Psychol* **1**, 293-319
(2005).
2.   Kessler, R.C. The effects of stressful life events on depression. *Annu Rev*
Psychol **48**, 191-214 (1997).
3.   Kendler, K.S., Karkowski, L.M. & Prescott, C.A. Causal relationship between
stressful life events and the onset of major depression. *Am J Psychiatry* **156**,
837-41 (1999).
4.   Paykel, E.S. Life events and affective disorders. *Acta Psychiatr Scand Suppl*,
61-6 (2003).
5.   Stroud, C.B., Davila, J. & Moyer, A. The relationship between stress and
depression in first onsets versus recurrences: a meta-analytic review. *J*
Abnorm Psychol **117**, 206-13 (2008).
6.   Ensel, W.M., Peek, M.K., Lin, N. & Lai, G. Stress in the life course: a life history
approach. *J Aging Health* **8**, 389-416 (1996).
7.   Kendler, K.S., Karkowski, L.M. & Prescott, C.A. Stressful life events and major
depression: risk period, long-term contextual threat, and diagnostic
specificity. *J Nerv Ment Dis* **186**, 661-9 (1998).
8.   Mazure, C.M. Life Stressors as Risk Factors in Depression. *Clinical*
Psychology: Science and Practice **5**, 291-313 (1998).
9.   Lichtenberg, P. & Belmaker, R.H. Subtyping major depressive disorder.
Psychother Psychosom **79**, 131-5 (2010).
10.   Elisei, S., Sciarma, T., Verdolini, N. & Anastasi, S. Resilience and depressive
disorders. *Psychiatr Danub* **25 Suppl 2**, S263-7 (2013).
11.   Monroe, S.M. & Simons, A.D. Diathesis-stress theories in the context of life
stress research: implications for the depressive disorders. *Psychol Bull* **110**,
406-25 (1991).
12.   Vogel, F. Schizophrenia genesis: The origins of madness. *American Journal of*
Human Genetics **48**, 1218-1218 (1991).
13.   Mann, J.J., Waternaux, C., Haas, G.L. & Malone, K.M. Toward a clinical model
of suicidal behavior in psychiatric patients. *Am J Psychiatry* **156**, 181-9
(1999).
14.   Riemann, D. *et al.* The hyperarousal model of insomnia: a review of the
concept and its evidence. *Sleep Med Rev* **14**, 19-31 (2010).
15.   Bolt, M.A., Helming, L.M. & Tintle, N.L. The Associations between Self-
Reported Exposure to the Chernobyl Nuclear Disaster Zone and Mental
Health Disorders in Ukraine. *Front Psychiatry* **9**, 32 (2018).
16.   Wray, N.R. & Sullivan, P.F. Genome-wide association analyses identify 44
risk variants and refine the genetic architecture of major depression.
bioRxiv (2017).
17.   Peyrot, W.J. *et al.* Effect of polygenic risk scores on depression in childhood
trauma. *Br J Psychiatry* **205**, 113-9 (2014).
18.   Musliner, K.L. *et al.* Polygenic risk, stressful life events and depressive
symptoms in older adults: a polygenic score analysis. *Psychol Med* **45**, 1709-
20 (2015).
19.   Peyrot, W.J. *et al.* Does Childhood Trauma Moderate Polygenic Risk for
Depression? A Meta-analysis of 5765 Subjects From the Psychiatric
Genomics Consortium. *Biol Psychiatry* (2017).

20. Dunn, E.C. *et al.* Genome-Wide Association Study (Gwas) and Genome-Wide
by Environment Interaction Study (Gweis) of Depressive Symptoms in
African American and Hispanic/Latina Women. *Depress Anxiety* **33**, 265-80
(2016).
21. Otowa, T. *et al.* The First Pilot Genome-Wide Gene-Environment Study of
Depression in the Japanese Population. *PLoS One* **11**, e0160823 (2016).
22. Ikeda, M. *et al.* Genome-wide environment interaction between depressive
state and stressful life events. *J Clin Psychiatry* **77**, e29-30 (2016).
23. Colodro-Conde, L. *et al.* A direct test of the diathesis-stress model for
depression. *Mol Psychiatry* (2017).
24. Coleman, J.R.I., Eley, T.C. & Breen, G. Genome-wide gene-environment
analyses of depression and reported lifetime traumatic experiences in UK
Biobank. *bioRxiv* (2018).
25. Mullins, N. *et al.* Polygenic interactions with environmental adversity in the
aetiology of major depressive disorder. *Psychol Med* **46**, 759-70 (2016).
26. Iyegbe, C., Campbell, D., Butler, A., Ajnakina, O. & Sham, P. The emerging
molecular architecture of schizophrenia, polygenic risk scores and the
clinical implications for GxE research. *Soc Psychiatry Psychiatr Epidemiol* **49**,
169-82 (2014).
27. McGrath, J.J., Mortensen, P.B., Visscher, P.M. & Wray, N.R. Where GWAS and
epidemiology meet: opportunities for the simultaneous study of genetic and
environmental risk factors in schizophrenia. *Schizophr Bull* **39**, 955-9
(2013).
28. Plomin, R. Commentary: missing heritability, polygenic scores, and gene-
environment correlation. *J Child Psychol Psychiatry* **54**, 1147-9 (2013).
29. Wray, N.R. *et al.* Research review: Polygenic methods and their application
to psychiatric traits. *J Child Psychol Psychiatry* **55**, 1068-87 (2014).
30. Smith, B.H. *et al.* Cohort Profile: Generation Scotland: Scottish Family Health
Study (GS:SFHS). The study, its participants and their potential for genetic
research on health and illness. *Int J Epidemiol* **42**, 689-700 (2013).
31. Gunderson, K.L. Whole-Genome Genotyping on Bead Arrays. in *DNA*
Microarrays for Biomedical Research: Methods and Protocols (ed. Dufva, M.)
197-213 (Humana Press, Totowa, NJ, 2009).
32. Kerr, S.M. *et al.* Pedigree and genotyping quality analyses of over 10,000
DNA samples from the Generation Scotland: Scottish Family Health Study.
BMC Med Genet **14**, 38 (2013).
33. Nagy, R. *et al.* Exploration of haplotype research consortium imputation for
genome-wide association studies in 20,032 Generation Scotland
participants. *Genome Med* **9**, 23 (2017).
34. Navrady, L.B. *et al.* Cohort Profile: Stratifying Resilience and Depression
Longitudinally (STRADL): a questionnaire follow-up of Generation Scotland:
Scottish Family Health Study (GS:SFHS). *Int J Epidemiol* (2017).
35. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and
population-based linkage analyses. *Am J Hum Genet* **81**, 559-75 (2007).
36. Smith, B.H. *et al.* Generation Scotland: the Scottish Family Health Study; a
new resource for researching genes and heritability. *BMC Med Genet* **7**, 74
(2006).
37. Fernandez-Pujals, A.M. *et al.* Epidemiology and Heritability of Major
Depressive Disorder, Stratified by Age of Onset, Sex, and Illness Course in

Generation Scotland: Scottish Family Health Study (GS:SFHS). *PLoS One* **10**,
e0142197 (2015).

38. Amador, C. *et al.* Recent genomic heritage in Scotland. *BMC Genomics* **16**,
437 (2015).

39. Goldberg, D.P. & Hillier, V.F. A scaled version of the General Health
Questionnaire. *Psychol Med* **9**, 139-45 (1979).

40. Sterling, M. General Health Questionnaire - 28 (GHQ-28). *J Physiother* **57**,
259 (2011).

41. Goldberg, D.P. *et al.* The validity of two versions of the GHQ in the WHO
study of mental illness in general health care. *Psychol Med* **27**, 191-7 (1997).

42. Banks, M.H. Validation of the General Health Questionnaire in a young
community sample. *Psychol Med* **13**, 349-53 (1983).

43. Marks, A.D.G., Horrocks, K.A. & Schutte, N.S. Emotional intelligence mediates
the relationship between insecure attachment and subjective health
outcomes. *Personality and Individual Differences* **98**, 188-192 (2016).

44. O'Rourke, S., MacHale, S., Signorini, D. & Dennis, M. Detecting psychiatric
morbidity after stroke: comparison of the GHQ and the HAD Scale. *Stroke*
29, 980-5 (1998).

45. Kessler, R.C., Andrews, G., Mroczek, D., Ustun, B. & Wittchen, H.-U. The
World Health Organization Composite International Diagnostic Interview
short-form (CIDI-SF). *International Journal of Methods in Psychiatric*
Research **7**, 171-185 (1998).

46. Brugha, T., Bebbington, P., Tennant, C. & Hurry, J. The List of Threatening
Experiences: a subset of 12 life event categories with considerable long-
term contextual threat. *Psychol Med* **15**, 189-94 (1985).

47. Brugha, T.S. & Cragg, D. The List of Threatening Experiences: the reliability
and validity of a brief life events questionnaire. *Acta Psychiatr Scand* **82**, 77-
81 (1990).

48. Motrico, E. *et al.* Psychometric properties of the List of Threatening
Experiences--LTE and its association with psychosocial factors and mental
disorders according to different scoring methods. *J Affect Disord* **150**, 931-
40 (2013).

49. Kendler, K.S., Karkowski, L.M. & Prescott, C.A. The assessment of
dependence in the study of stressful life events: validation using a twin
design. *Psychol Med* **29**, 1455-60 (1999).

50. Euesden, J., Lewis, C.M. & O'Reilly, P.F. PRSice: Polygenic Risk Score
software. *Bioinformatics* **31**, 1466-8 (2015).

51. Yang, J., Lee, S.H., Goddard, M.E. & Visscher, P.M. GCTA: a tool for genome-
wide complex trait analysis. *Am J Hum Genet* **88**, 76-82 (2011).

52. Keller, M.C. Gene x environment interaction studies have not properly
controlled for potential confounders: the problem and the (simple) solution.
Biol Psychiatry **75**, 18-24 (2014).

53. Belsky, J. & Beaver, K.M. Cumulative-genetic plasticity, parenting and
adolescent self-regulation. *J Child Psychol Psychiatry* **52**, 619-26 (2011).

54. Belsky, J. *et al.* Vulnerability genes or plasticity genes? *Mol Psychiatry* **14**,
746-54 (2009).

55. Dudbridge, F. Power and predictive accuracy of polygenic risk scores. *PLoS*
Genet **9**, e1003348 (2013).

56. Eaves, L.J., Last, K., Martin, N.G. & Jinks, J.L. A progressive approach to non-
additivity and genotype-environmental covariance in the analysis of human
differences. *British Journal of Mathematical and Statistical Psychology* **30**, 1-
42 (1977).
57. Polderman, T.J. *et al.* Meta-analysis of the heritability of human traits based
on fifty years of twin studies. *Nat Genet* **47**, 702-9 (2015).
58. Clarke, T. *et al.* Genetic and environmental determinants of stressful life
events and their overlap with depression and neuroticism [version 1;
referees: 3 approved with reservations]. *Wellcome Open Research* **3**(2018).
59. Kendler, K.S., Kuhn, J. & Prescott, C.A. The interrelationship of neuroticism,
sex, and stressful life events in the prediction of episodes of major
depression. *Am J Psychiatry* **161**, 631-6 (2004).
60. Kendler, K.S. & Eaves, L.J. Models for the joint effect of genotype and
environment on liability to psychiatric illness. *Am J Psychiatry* **143**, 279-89
(1986).
61. Plomin, R., DeFries, J.C. & Loehlin, J.C. Genotype-environment interaction
and correlation in the analysis of human behavior. *Psychol Bull* **84**, 309-22
(1977).
62. Plomin, R., Lichtenstein, P., Pedersen, N.L., McClearn, G.E. & Nesselroade, J.R.
Genetic influence on life events during the last half of the life span. *Psychol*
Aging **5**, 25-30 (1990).
63. Clarke, T.K. *et al.* Genetic and environmental determinants of stressful life
events and their overlap with depression and neuroticism [version 1;
referees: awaiting peer review]. in *Wellcome Open Research* Vol. 3 %M
10.12688/wellcomeopenres.13893.1 (2018).
64. Weissman, M.M. *et al.* Sex differences in rates of depression: cross-national
perspectives. *J Affect Disord* **29**, 77-84 (1993).
65. Van de Velde, S., Bracke, P. & Levecque, K. Gender differences in depression
in 23 European countries. Cross-national variation in the gender gap in
depression. *Soc Sci Med* **71**, 305-13 (2010).
66. Labonte, B. *et al.* Sex-specific transcriptional signatures in human
depression. *Nat Med* (2017).
67. Angst, J. *et al.* Gender differences in depression. Epidemiological findings
from the European DEPRES I and II studies. *Eur Arch Psychiatry Clin*
Neurosci **252**, 201-9 (2002).
68. Piccinelli, M. & Wilkinson, G. Gender differences in depression. Critical
review. *Br J Psychiatry* **177**, 486-92 (2000).
69. Vrshek-Schallhorn, S. *et al.* Additive genetic risk from five serotonin system
polymorphisms interacts with interpersonal stress to predict depression. *J*
Abnorm Psychol **124**, 776-90 (2015).
70. Belsky, J. & Pluess, M. Beyond diathesis stress: differential susceptibility to
environmental influences. *Psychol Bull* **135**, 885-908 (2009).
71. Belsky, J., Bakermans-Kranenburg, M.J. & van Ijzendoorn, M.H. For Better
and for Worse: Differential Susceptibility to Environmental Influences.
Current Directions in Psychological Science **16**, 300-304 (2007).
72. Li, J.J., Berk, M.S. & Lee, S.S. Differential susceptibility in longitudinal models
of gene-environment interaction for adolescent depression. *Dev*
Psychopathol **25**, 991-1003 (2013).

73.   Kang, S.-M. & G. Waller, N. *Moderated Multiple Regression, Spurious*
Interaction Effects, and IRT, 87-105 (2005).
652

FIGURE LEGENDS

Figure 1. a) Association between polygenic risk scores (PRS) and depression score

(main effects, one-sided tests). PRS were generated at 8 *p*-threshold levels using

summary statistics from the Psychiatric Genetic Consortium MDD GWAS (released July

2016) with the exclusion of Generation Scotland participants. The depression score

was derived from The General Health Questionnaire. The Y-axis represents the % of

variance of depression score explained by PRS main effects. The full cohort (yellow)

was split into men (blue) and women (red). In Colodro-Conde *et al.,* PRS for MDD

significantly explained up to 0.46% of depression score in their sample (~0.39% in

women and ~0.70% in men). **b)** Association between reported number of SLE and

depression score (main effect, one-sided tests, results expressed in % of depression

score explained). SLE were self-reported through a brief life-events questionnaire

based on the List of Threating Experiences and categorized into: total number of SLE

reported (TSLE), "dependent" SLE (DSLE) or "independent" SLE (ISLE). The full cohort

(yellow) was split into men (blue) and women (red). In Colodro-Conde *et al.,*

"personal" SLE significantly explained up to 12.9% of depression score variance in their

sample (~11.5% in women and ~16% in men).

Figure 2. Association between GxE effect and depression score. The results represent

percentage of depression score explained by the interaction term (two-sided tests)

fitted in linear mixed models to empirically test the *diathesis-stress* model. Red

numbers show significant interactions *p*-values. *Shows significance of difference

between sexes when comparing the size of the estimated GxE effects. The full cohort

(yellow) was split into men (blue) and women (red). PRS were generated at 8 *p*-

threshold levels using summary statistics from the Psychiatric Genetic Consortium

MDD GWAS (released July 2016) with the exclusion of Generation Scotland

participants. The interaction effect was tested with **a)** the number of SLE reported

(TSLE), **b)** "dependent" SLE (DSLE) and **c)** "independent" SLE (ISLE). In Colodro-Conde *et*

al., the variance of depression score explained in their sample by GxE was 0.12% (p = 7

x 10^{-3}). GxE were also significant in women (p = 2 x 10^{-3}) explaining up to 0.25% of

depression score variation, but not in men (p = 0.059; R^2 = 0.17%; negative/protective

effect on depression score).

Figure 3. Scatterplot of *diathesis-stress* interactions on depression score. Interactions

with PRS at which nominally significant GxE effects were detected in **a)** full cohort (p-

threshold = 0.01) and **b)** in women (p-threshold = 1 x 10^{-5}) are shown. At bottom, the

remaining samples (i.e., *full cohort*, *women* or *men*) at same p-threshold are shown for

comparison. The X-axis represents the direct effect of PRS (standard deviation from

the mean) based on **a)** p-threshold = 0.01 and **b)** p-threshold = 1 x 10^{-5}, using the total

number of SLE reported by each participant (dot) as environmental exposures at three

SLE levels represented by colours. Blue: 0 SLE, "no stress", n = 1 833/1 041/792; green:

1 or 2 SLE, "low stress", n = 2 311/1 459/852; red: 3 or more SLE, "high stress", n =

775/490/285; in the full cohort, women and men, respectively. Y-axis reflects the

depression score standardized to mean of 0 and standard deviation of 1. Lines

represent the increment in risk of depression under a certain degree of "stress"

dependent on a genetic predisposition (= *diathesis*).

Table 1. *Diathesis* effect on depression score under SLE categories. Reported values at p-thresholds where nominally significant GxE effects were detected.

	PRS at p value threshold = 0.01								
Sample	***FULL COHORT**			**WOMEN**			**MEN**		
SLE category	none	low	high	none	low	high	none	low	high
N	1833	2311	**775**	1041	1459	490	792	852	285
Effect	0.021	0.043	**0.142**	0.0118	0.0617	0.1538	0.0346	0.0113	0.1227
s.e.	0.022	0.021	**0.039**	0.029	0.027	0.049	0.034	0.032	0.063
t	0.957	2.07	**3.644**	0.403	2.274	3.112	1.021	0.348	1.947
***p* value**	0.339	0.039	**2.86×10^{-4}**	0.687	0.023	0.002	0.307	0.728	0.053
CI (95%)	-0.022, 0.065	0.002, 0.084	**0.065, 0.218**	-0.046, 0.069	0.008, 0.115	0.057, 0.251	-0.032, 0.101	-0.052, 0.075	-0.001,0.247

	PRS at p value threshold = 1×10^{-5}								
Sample	**FULL COHORT**			***WOMEN**			**MEN**		
SLE category	none	low	high	none	low	high	none	low	high
N	1833	2311	775	1041	1459	490	792	852	285
Effect	-0.0022	0.0032	0.0705	-0.061	0.014	0.078	0.082	-0.0176	0.0548
s.e.	0.022	0.021	0.04	0.029	0.027	0.049	0.034	0.033	0.07
t	-0.098	0.153	1.76	-2.086	0.541	1.609	2.416	-0.537	0.778
***p* value**	0.922	0.878	0.079	0.037	0.589	0.108	0.016	0.592	0.437
CI (95%)	-0.046, 0.041	-0.037, 0.044	-0.008, 0.149	-0.119, -0.004	-0.038, 0.066	-0.017, 0.174	0.015, 0.149	-0.082,0.047	-0.084, 0.193

*Sample where nominally significant GxE was detected. SLE categories (number of SLE reported): 0 SLE = "none", 1 or 2 SLE = "low", and 3 or more SLE = "high". In red, nominally significant effects. In **bold red**, robustly significant effect after conservative Bonferroni correction ($p < 6.94 \times 10^{-4}$).

Figure 1.

Figure 2.

Figure 3.