Hyperspectral Reflectance-Derived Relationship Matrices for Genomic Prediction of

Grain Yield in Wheat

Margaret R. Krause[*], Lorena González-Pérez[†], José Crossa[†], Paulino Pérez-Rodríguez[‡],

Osval Montesinos-López[§], Ravi P. Singh[†], Susanne Dreisigacker[†], Jesse Poland[**], Jessica

Rutkoski[††], Mark Sorrells[*], Michael A. Gore[*], Suchismita Mondal[†]

[*]Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell

University, Ithaca, New York, 14853, USA

[†]Global Wheat Program, International Maize and Wheat Improvement Center

(CIMMYT), Ciudad de México, 06600, México

[‡]Colegio de Postgraduados, CP 56230, Montecillos, Edo. de México

[§]Facultad de Telemática, Universidad de Colima, Colima, Colima, 28040, México

[**] Department of Plant Pathology, Kansas State University, Manhattan, Kansas, 66506,

USA

[††] International Rice Research Institute (IRRI), DAPO Box 7777, Metro Manila, 1301,

Philippines

Corresponding Author:

Suchismita Mondal

Km45 Carretera Mex-Veracruz

El Batán, Texcoco

México. C.P 56237

Ph. +52 55-58042004 ext 1135

S.Mondal@cgiar.org

1   ABSTRACT

Hyperspectral reflectance phenotyping and genomic selection are two emerging

technologies that have the potential to increase plant breeding efficiency by improving

prediction accuracy for grain yield. Hyperspectral cameras quantify canopy reflectance

across a wide range of wavelengths that are associated with numerous biophysical and

biochemical processes in plants. Genomic selection models utilize genome-wide marker

or pedigree information to predict the genetic values of breeding lines. In this study, we

propose a multi-kernel GBLUP approach to genomic selection that uses genomic marker-

, pedigree-, and hyperspectral reflectance-derived relationship matrices to model the

genetic main effects and genotype $\times$ environment ($G \times E$) interactions across

environments within a bread wheat (*Triticum aestivum* L.) breeding program. We utilized

an airplane equipped with a hyperspectral camera to phenotype five differentially

managed treatments of the yield trials conducted by the Bread Wheat Improvement

Program, International Maize and Wheat Improvement Center (CIMMYT) at Ciudad

Obregón, México over four breeding cycles. We observed that single-kernel models

using hyperspectral reflectance-derived relationship matrices performed similarly or

superior to marker- and pedigree-based genomic selection models when predicting within

and across environments. Multi-kernel models combining marker/pedigree information

with hyperspectral reflectance phentoypes had the highest prediction accuracies;

however, improvements in accuracy over marker- and pedigree-based models were

marginal when correcting for days to heading. Our results demonstrates the potential of

hyperspectral imaging in predicting grain yield within a multi-environment context, it

also supports further studies on integration of hyperspectral reflectance phenotyping in

breeding programs.

3   INTRODUCTION

The aim of plant breeding programs is to develop and deliver new, high-yielding crop

varieties that are adapted to a range of environmental conditions. Two major challenges

of breeding for multiple environments are: (1) being able to account for the presence of

genotype-by-environment interaction ($G \times E$), and (2) the high costs of evaluating trials

at multiple locations. Statistical, genomics, and phenomics tools that enable the accurate

prediction and selection of candidate lines appropriate for each target environment may

serve to increase the rate of genetic gain while reducing costs associated with large-scale

multi-environment field trials.

In recent years, genomic selection (GS) and high-throughput phenotyping (HTP)

have emerged as potential technologies for improving breeding efficiency (Furbank and

Tester, 2011; Cabrera-Bosquet *et al.* 2012; Crossa *et al.* 2017). In GS, genome-wide

marker effects are estimated for a "training set" of lines that has been phenotyped and

genotyped (Meuwissen *et al.* 2001). Those estimates are then applied to selection

candidates prior to phenotyping to predict trait values, which may reduce the time and

cost of testing breeding lines. While GS was initially developed to predict within

individual environments, recent studies have extended the genomic best linear unbiased

prediction (GBLUP) model to accommodate $G \times E$ interactions. Burgueño *et al.* (2012),

Heslot *et al.* (2013), Jarquín *et al.* (2014), López-Cruz *et al.* (2015), and Pérez-Rodríguez

et al. (2017) reported increases in prediction accuracies with extended models relative to

single-environment analyses.

HTP is based on the remote and proximal sensing of a large number of crop plants

to collect relevant phenotypes while reducing labor time and cost (White *et al.* 2012;

Araus and Cairns, 2014; Pauli *et al.* 2016). When deployed across different

developmental growth stages and in multiple environments, HTP can drastically increase

the phenotypic information available to breeding programs, which may help to improve

selection accuracy. Although HTP traits represent indirect estimations and may not be

able to provide the level of precision of direct measurement, they may be of particular use

at the early generation stage in breeding programs when seed is limited. In wheat

breeding, large numbers of lines are sown in small unreplicated plots for the purpose of

visual selection and seed increase prior to grain yield testing in large replicated plots.

Measurements of grain yield from small plots are not meaningful, nor are they feasible to

collect when thousands of lines are being screened. Therefore, acquiring accurate

predictions of grain yield at the early generation stage using HTP may serve to improve

selection accuracy.

A range of ground-based and aerial HTP platforms have been recently developed

to improve the accuracy, efficiency, and scope of phenotypic data collection (Andrade-

Sanchez *et al.* 2014; Crain *et al.* 2016; Haghighattalab *et al.* 2016). A major advantage of

aerial platforms is their ability to phenotype large areas of field trials in minimal time.

This enhanced efficiency increases the spatial and temporal resolution of the phenotypic

data and may be critical when assessing breeding trials at multiple locations. A number of

recent studies have integrated traits collected with HTP into GS to increase prediction

accuracy for grain yield in wheat (Rutkoski *et al.* 2014; Sun *et al.* 2017; Crain *et al.*

2018).

To date, many of the applications of aerial HTP within plant breeding have

focused on measuring the spectral reflectance of the crop canopy. Plant cells, tissues, and

pigments have wavelength-specific light absorption, reflectance, and transmittance

patterns that may, for example, differentiate between healthy and stressed plants (Li *et al.*

2014). Vegetation indices (VIs) provide a convenient way of summarizing spectral

reflectance information into scores that may be predictive of economically important

traits (Govaerts *et al.* 1999). VIs such as the normalized difference vegetation index

(NDVI) have been shown to be predictive of grain yield in wheat (Aparicio *et al.* 1999;

Labus *et al.* 2002). However, since VIs are calculated from only a few wavelengths, they

cannot leverage the high density of information captured by hyperspectral cameras,

which record reflectance at a large number of narrowband wavelengths in the visible and

near infrared regions of the light spectrum (Viña *et al.* 2011). While hyperspectral data

may have a greater capacity than VIs to detect phenotypic differences between

individuals, the high dimensionality of the data may complicate the interpretation. For

these data to become meaningful for plant breeding, methods that can derive useful

information on traits relevant to breeding are needed.

To address the high dimensionality of hyperspectral data, Aguate *et al.* (2017)

found that integrating information from all hyperspectral wavelengths using ordinary

least squares, partial least squares, and Bayesian shrinkage resulted in higher prediction

accuracy than what could be achieved using individual VIs in maize. Montesinos-López

et al. (2017a) proposed a Bayesian functional regression analysis using hyperspectral

wavelengths that likewise resulted in higher accuracies for predicting grain yield in wheat

when compared to a range of VIs. Montesinos-López *et al.* (2017b) further extended this

method to incorporate genomic and pedigree information, in addition to accommodating

$G \times E$ by modeling hyperspectral band-by-environment ($B \times E$) interactions. Their study

found that models that included the $B \times E$ term had higher prediction accuracies than

those that did not, suggesting that hyperspectral reflectance may be a useful phenotype

for modeling $G \times E$ interactions.

When collecting hyperspectral data within a multi-environment context, the

number of predictors increases in proportion to the number of environments and

phenotyping time-points observed, which may come at a computational cost depending

on the type of prediction model used (Montesinos-López *et al.* 2017b). One possible

approach that may minimize computation time would be to use the hyperspectral bands

as a high dimensional predictor set, similar to the case of prediction with genomic

markers in GBLUP, that is, to create a relationship matrix between individuals using the

hyperspectral bands. This way, the number of bands could be very large without

increasing the complexity of the GBLUP prediction model. Separate genomic

marker/pedigree and hyperspectral reflectance kernels could be integrated to model the

main genetic effects and $G \times E$ interactions, respectively.

Multi-environment field trials are often unbalanced, which can complicate their

use in prediction across environments. When deploying HTP in large breeding programs,

it can be difficult to ensure that all locations are phenotyped at the same stage and the

same number of times throughout the season. Weather conditions, technical difficulties

with the cameras or sensors, and scheduling with contracted pilots/airports may prohibit

the use of aerial HTP on certain days, and lines grown at different sites may develop at

faster or slower rates depending on weather and management conditions. As a result, sites

may have different number of observed HTP time-points, as in Sun *et al.* (2017). It is

well documented that canopy spectral reflectance varies according to crop phenology

(Viña *et al.* 2003; Zhang *et al.* 2003). One strategy for comparing HTP time-points across

varying sites may be to classify them according to the predominant developmental

growth stage at the time of phenotyping.

To test these approaches, we deployed HTP on the CIMMYT Bread Wheat

Improvement Program's multi-environment yield evaluations of advanced germplasm to

phenotype a range of differentially managed treatments with a hyperspectral camera at

multiple time-points throughout the growing season. The main objectives of this research

were to: (1) propose a multi-kernel, multi-environment GBLUP model that involves

modeling genetic main effects using genomic markers or pedigrees and modeling $G \times E$

interactions using relationship matrices derived from hyperspectral reflectance data, (2)

compare the prediction accuracies of models using genomic marker/pedigree main effects

kernels and hyperspectral $G \times E$ interaction kernels separately and in combination, and

(3) determine the optimal developmental growth stages for hyperspectral phenotyping

based on grain yield prediction accuracy.

18   MATERIALS AND METHODS

Experimental Data

The dataset included a total of 3,771 bread wheat lines evaluated at the Campo

Experimental Norman E. Borlaug in Ciudad Obregón, México over the course of four

breeding cycles. In each cycle, lines were sown under five differentially managed

treatments: Optimal Bed, Optimal Flat, Moderate Drought, Severe Drought, and Heat.

Descriptions of the managed treatments are given in Table 1. The 20 managed treatment-

breeding cycle combinations are herein referred to as site-years. Within each of the five

managed treatments, 1092 lines were arranged into 39 trials in a α-lattice design with

three replicates and six incomplete blocks per replicate. Each replicate contained two

repeated checks. 'Kachu #1' was sown as a check in the Optimal Bed, Optimal Flat, and

Moderate Drought managed treatments. 'Baj #1' was used as a check in the Severe

Drought and Heat managed treatments. 'Borlaug100 F2014' was sown as the other check

in all managed treatments. All lines within a breeding cycle were evaluated in all five

managed treatments, while no lines other than checks overlapped between cycles.

Records for some lines were removed for the analysis due to unavailability of genotypic

or agronomic data, resulting in the final dataset: 588 lines in 2013-14, 1,033 lines in

2014-15, 1,063 lines in 2015-16, and 1,087 lines in 2016-17, for a total of 56,565 plots

phenotyped. Each breeding cycle contained full-sib families with an average of two full-

sibs per family.

15   Grain yield (GY) in t ha^{-1} and lodging evaluated on an ordinal scale (0: no

lodging; 5: completely lodged) were assessed in all three replicates in each site-year.

Heading date was recorded as the date of 50% spike emergence within the plot. Maturity

date was recorded as the date of senescence of the peduncle for 50% of stems in the plot.

Days to heading (DTHD) and days to maturity (DTMT) were measured as the number of

days to reach heading and maturity, starting from the date of the first irrigation if sowing

was on dry soil or from the sowing date if sowing was in pre-irrigated fields. DTHD and

DTMT were assessed for the first replicate only in each site-year due to the high

heritability of the traits.

Hyperspectral Phenotyping

Hyperspectral reflectance data were collected with a hyperspectral camera (A-series,

Micro-Hyperspec VNIR, Headwall photonics Fitchburg, MA, USA) as part of the Alava

Remote Sensing Spectral Solution (ARS3, Alava Ingenieros, Madrid, Spain) mounted in

a Piper PA-16 Clipper aircraft. The camera's sensor had a 12-bit radiometric resolution,

covering the light spectrum in the 400–850nm region with a 7.5nm full width at half

maximum and was set with an integration time of 18ms. The flights were scheduled

around noon (GMT-7) and aligned to the solar azimuth angle at a height of 300m,

resulting in 30cm resolution. The aerial imagery acquisition during the growing cycle

was spaced approximately at seven- to ten-day intervals on mostly clear days.

Radiometric calibration of the sensor was done using coefficients derived from a

calibrated uniform light source and an integrating sphere (CSTMUSS2000C Uniform

Source System, LabSphere, North Sutton, NH, USA). Dark frame correction was

performed for each flight dataset. Atmospheric calibration was performed using

irradiance measurements acquired at the beginning and end of each flight using a Jaz

spectrometer with a CC-3 Cosine Corrector (Ocean Optics Inc, FL, USA) for the 2016-17

breeding cycle. For remaining three breeding cycles, irradiance was modeled using

aerosol optical depth from sun-photometer measurements (Microtops II, Solar Light

Company, Glenside, PA, USA) based on the SMARTS simulation model (Gueymard,

1995, 2001).

Ortho-rectification and georeferencing of the imagery were performed using

PARGE (ReSe Applications Schläpfer, Wil, Switzerland) based on data from the intertial

navigation system (INS) attached to the camera (IG-500N model, SGB systems S. A. S.,

Carrières-sur-Seine, France). Hyperspectral reflectance data were extracted from the

aerial images using the mean value of the pixels inside the central area of each observed

plot, avoiding 0.5m from the plot border.

Each hyperspectral phenotyping time-point within each site-year was assigned a

developmental growth stage classification according to the predominant growth stage of

the lines within the site-year at the time of phenotyping (Table 2). The vegetative (VEG)

stage was defined as the period between germination and 50% of plots at heading. The

heading stage (HEAD) was defined as the period between 50% of plots at heading and

100% of plots at heading. The grain fill stage (GF) was defined as the period between

100% of plots at HD and 100% of plots at maturity.

Genotypic Data

All lines were genotyped using genotyping-by-sequencing (Elshire *et al.* 2011) according

to the pipeline described in Poland *et al.* (2012). From the initial set of 34,900 single

nucleotide polymorphisms (SNPs), 8,519 SNPs remained after excluding all markers with

more than 70% of missing data or minor allele frequency less than 0.05. For each marker,

missing data were imputed using the sample mean of observed values (Poland *et al.*,

2012).

Basic Statistical Models

Within each site-year, best linear unbiased estimates (BLUEs) were calculated for the

agronomic traits and for each hyperspectral band at each phenotyping time-point using

the following model:

23
$$y_{ijkl} = \mu + g_i + t_j + r_{k(j)} + b_{l(jk)} + \varepsilon_{ijkl}, \qquad (1)$$

where y_{ijkl} is the trait value for genotype i within trial j, replicate k, and block l; μ is the

overall mean; g_i is the fixed effect for genotype i; t_j is the random effect for trial j, which

are assumed to be independently and identically distributed according to a normal

distribution with mean zero and variance σ_t^2, that is, $t_j \sim$ iid $N(0, \sigma_t^2)$; $r_{k(j)} \sim$ iid $N(0, \sigma_r^2)$ is

the random effect for replicate k within trial j; $b_{l(jk)} \sim$ iid $N(0, \sigma_b^2)$ is the random effect for

block l within replicate k and trial j; and $\varepsilon_{ijkl} \sim iid\ N(0, \sigma_\varepsilon^2)$ is the residual effect. For

DTHD and DTMT, which were recorded for only one replicate, the replicate and block

effects were excluded from the model. For validation of prediction models, best linear

unbiased predictions (BLUPs) for GY within each site-year were calculated by treating

the effect for genotype g_i in model (1) as random with $g_i \sim iid\ N(0, \sigma_g^2)$ and by including

a covariate for lodging as a fixed effect, fit only when lodging was observed within a site-

year. To correct for the influence of phenology, BLUPs for GY were calculated again

using DTHD included as a fixed effect in model (1).

BLUEs were also calculated for each hyperspectral band on a development growth

stage basis according to the growth stage classifications detailed in Table 2. BLUEs for

each hyperspectral band for the VEG, HEAD, and GF stages were estimated by fitting the

following model:

$$y_{ijklm} = \mu + g_i + d_j + t_{k(j)} + r_{l(jk)} + b_{m(jkl)} + \varepsilon_{ijklm}, \qquad (2)$$

where y_{ijkl} is the hyperspectral reflectance trait value for genotype i on time-point j and

within trial k, replicate l, and block m; μ is the overall mean, g_i is the fixed genetic effect

for genotype i; $d_j \sim iid\ N(0, \sigma_d^2)$ is a random effect for time-point j; $t_{k(j)} \sim iid\ N(0, \sigma_d^2)$ is

random effect for trial k nested within time-point j; $r_{l(jk)} \sim iid\ N(0, \sigma_r^2)$ is the random

effect for replicate l nested within time-point j and trial k; $b_{m(jkl)} \sim iid\ N(0, \sigma_b^2)$ is the

random effect for block m nested within time-point j, trial k, and rep l; and $\varepsilon_{ijklm} \sim iid\ N(0,$

$\sigma_\varepsilon^2)$ is the residual effect. BLUEs were also calculated for each hyperspectral band across

all available phenotyping time-points (ALL) using model (2).

Broad-sense heritability within site-years was calculated for GY (Bernardo 2010) and

each hyperspectral band as:

10
$$H^2 = \frac{\sigma_g^2}{\sigma_g^2 + \frac{\sigma_\varepsilon^2}{nreps}} \qquad (3)$$

where $\sigma_g{}^2$ is the genetic variance, σ_ε^2 is the error variance, and *nreps* is the number of

replicates (*nreps* = 3). For each breeding cycle, Pearson's correlations for GY between

managed treatments were calculated using the BLUEs derived from model (1). Pearson's

correlations were also calculated between GY BLUEs and the BLUEs for each

hyperspectral band calculated in model (2).

Relationship Matrices

Genetic relationships between individuals were modeled using genomic markers,

pedigrees, and hyperspectral reflectance phenotypes. The genomic relationship matrix

(**G**) was calculated according to Endelman and Jannink (2012). The additive relationship

matrix (**A**) was derived from pedigrees and calculated as twice the coefficient of

parentage.

Hyperspectral reflectance-based relationship matrices (H) were calculated within

each site-year using the hyperspectral BLUEs calculated for 1) the individual time-points

from model (1), 2) the developmental growth stages from model (2), and 3) all time-

points from model (2). The matrices were calculated as $H = \frac{SS'}{nbands}$, where S is a matrix

of the centered and standardized BLUEs of the hyperspectral bands and *nbands* is the

total number of hyperspectral bands observed.

Prediction Models

Genetic Main Effects: To compare the utility of hyperspectral reflectance-based models

to standard genomic prediction models, the following single-kernel genetic main effects

model was fitted using marker- and pedigree-derived relationship matrices:

13                              $$y_{ij} = \mu + E_i + g_j + \varepsilon_{ij} \tag{4}$$

where y_{ij} is the BLUE of GY for genotype j in site-year i, μ is the overall mean, E_i is the

fixed effect for site-year ($i=1,...,I$), g_j is the random effect for genotype j ($j=1,...,J$), and

ε_{ij} is the residual effect. We assume that the joint distribution of genotype effects is

distributed according to a multivariate normal distribution with mean $\mathbf{0}$ and variance-

covariance matrix $\sigma_g^2 K$, that is, $\mathbf{g} = (g_1, ..., g_J)^T \sim MVN(\mathbf{0}, \sigma_g^2 K)$, where σ_g^2 denotes the

genomic variance and K represent the genomic relationship matrix between individuals G

($K=G$) or the numerical relationship matrix A ($K=A$).

Hyperspectral Reflectance Main Effects: To mimic a situation in which genomic

marker and pedigree information are not available, the following single-kernel model was

fitted using the hyperspectral reflectance-derived relationship matrices only:

$$y_{ij} = \mu + E_i + Eh_{ij} + \varepsilon_{ij} \tag{5}$$

where y_{ij}, μ, and E_i were as defined in model (4). Eh_{ij} is the random effect of the

hyperspectral bands for genotype j in site-year i with the joint distribution of the

hyperspectral bands as $\boldsymbol{hE} = (hE_{11}, \dots, hE_{IJ})^T \sim MVN(\boldsymbol{0}, \sigma_h^2 \boldsymbol{H})$, where σ_h^2 is the

hyperspectral band variance and $\boldsymbol{H}$ is the hyperspectral reflectance-derived relationship

matrix. Unlike matrices $\boldsymbol{G}$ and $\boldsymbol{A}$, which have only one row and column for each unique

genotype evaluated, the matrix $\boldsymbol{H}$ has a row and column for each unique site-year-

genotype combination where $I \times J$ results in the total number of site-year-genotype

combinations.

Genetic Main Effects + Genetic G × E: As a basis for comparison to assess the

advantage of $G \times E$ models that integrate both markers or pedigrees with hyperspectral

reflectance phenotypes, model (4) was extended to accommodate $G \times E$ interactions

using marker or pedigree information. The following multi-kernel model was fit:

$$y_{ij} = \mu + E_i + g_j + Eg_{ij} + \varepsilon_{ij} \tag{6}$$

where y_{ij}, μ, E_i, and g_j were as defined in model (4). The gE_{ij} term was assumed to have

multivariate normal distribution $\boldsymbol{gE} = (gE_{11}, ..., gE_{IJ})^T \sim MVN(\boldsymbol{0},$

$(\mathbf{Z_g}K\mathbf{Z_g}^T)\#(\mathbf{Z_E}\mathbf{Z_E}^T)\sigma_{gE}^2)$ where $\mathbf{Z_g}$ and $\mathbf{Z_E}$ are incidence matrices for genotypes and site-

years, K represents the genomic relationship matrix $(\boldsymbol{K=G})$ or the numerical relationship

matrix A $(\boldsymbol{K=A})$, and σ_{gE}^2 is the variance component for gE_{ij} (Jarquín *et al.* 2014).

Genetic Main Effects + Hyperspectral Reflectance G × E: Finally, a multi-kernel

model using marker or pedigree information to estimate the genetic main effects and

hyperspectral reflectance phenotypes to model the $G \times E$ interactions was fitted:

11                           $y_{ij} = \mu + E_i + g_j + hE_{ij} + \varepsilon_{ij}$ (7)

Here, y_{ij}, μ, E_i, and g_j were defined as above in model (4). The term hE_{ij} was assumed

to have multivariate normal distribution $\boldsymbol{hE} = (hE_{11}, ..., hE_{IJ})^T \sim MVN(\boldsymbol{0},$

$(\mathbf{Z_g}H\mathbf{Z_g}^T)\#(\mathbf{Z_E}\mathbf{Z_E}^T)\sigma_{hE}^2)$ where $\mathbf{Z_g}$ and $\mathbf{Z_E}$ are incidence matrices for genotypes and site-

years and σ_{hE}^2 is the variance component for hE_{ij}. The $(\mathbf{Z_g}H\mathbf{Z_g}^T)\#(\mathbf{Z_E}\mathbf{Z_E}^T)$ term was

obtained with the block diagonal matrix $BDiag(\boldsymbol{H}_{11},...,\boldsymbol{H}_{II})$ where $\boldsymbol{H}_{ii}$ represent the

hyperspectral relationship matrices for genotypes in site-year $i=1,...,I$.

Software

Processing of the hyperspectral images was performed using the ARS3 hyproQ software

(Álava Ingenieros, Madrid, Spain). Plot polygons for tabular data extraction were

generated using ArcGIS (ESRI, Redlands, California, US). Images were aligned

manually in ArcGIS if they did not overlay the plot polygons due to INS inaccuracy.

All models were fit using the R statistical programming language (R Core Team,

2018). Basic models (1-3) were fit with the package "ASReml-R" (Gilmour *et al.*, 2014)

for R, while the prediction models (4-7) were fit using the "BGLR" package (de los

Campos and Pérez-Rodríguez, 2014) for R. The marker-based genetic relationship

matrices were calculated using the A.mat() function within the "rrBLUP" package

(Endelman 2011) for R. The coefficients of parentage were estimated using the "Browse"

application within the International Crop Information System software package

(McLaren *et al.* 2000).

Assessing Model Prediction Accuracy

The above models were fit to assess model prediction accuracy for three prediction

strategies representing different testing and evaluation problems relevant to plant

breeding programs: 1) **within site-year**, 2) **within breeding cycle/across managed**

treatments, and 3) **across breeding cycles/within managed treatment**.

A train-test (TRN-TST) validation scheme (Daetwyler *et al.*, 2013) of 20 random

partitions was used to assess model prediction accuracy for all prediction strategies. For

within site-year prediction, a random 80% of records within a given site-year were

assigned to the TRN set, and the remaining 20% were used as the TST set for prediction.

This reflects a scenario in which prediction of the trait values for a set of unobserved

lines is performed within a managed treatment and breeding cycle of interest. Models (4)

and (5) were fitted without the E_i term for site-year and models (6) and (7) were not

tested for the within site-year prediction scheme because multiple site-years were not

considered.

For prediction within breeding cycle/across managed treatments, predictions were

carried out across the five managed treatments within a single breeding cycle. The TRN

set consisted of all records from four out of the five managed treatments within a

breeding cycle plus 20% of records from the fifth managed treatment. The remaining

80% of records from the fifth managed treatment were assigned to the TST set for

prediction. In this situation, lines have been previously characterized in all managed

treatments except one, where only 20% of records were available.

For prediction across breeding cycles/within managed treatment, predictions were

performed across the four breeding cycles but within a single managed treatment. The

TRN set contained all records from three out of the four breeding cycles for a given

managed treatment plus 20% of records from the fourth breeding cycle. The remaining

80% of records from the fourth breeding cycle were assigned to the TST set for

prediction. This scenario predicts performance of unobserved lines that are 1-3 breeding

cycles removed from lines in the training set.

Table S1 illustrates examples of TRN-TST partitioning for the three prediction

schemes. For all prediction strategies, model prediction accuracy was assessed as the

Pearson's correlation between predictions for GY and GY BLUPs with and without

correction for DTHD calculated in model (1). Reported values are the mean and standard

deviation of the 20 random TRN-TST partitions implemented. The same partitioning

"splits" were used to assess all models to ensure fair comparisons.

Data Availability

All phenotypic and genotypic data required to confirm the results presented in this study

are available on CIMMYT Dataverse link: hdl:11529/10548109. The "GID" column

denotes the unique identifiers for the genotypes. Supplementary file sets, tables and

figures are available at FigShare.

6   RESULTS

Descriptive Statistics

To evaluate the potential of integrating aerial hyperspectral reflectance phenotypes into

GS to improve prediction accuracy for GY in wheat, we deployed an airplane equipped

with a hyperspectral camera to phenotype wheat breeding trials in 20 site-years at

multiple time-points throughout the growing season. Across the site-years, the Optimal

Bed and Optimal Flat managed treatments had higher GY than the stressed Moderate

Drought, Severe Drought, and Heat managed treatments (Figure 1). The standard

deviations for GY remained relatively stable across site-years, ranging from 0.30 to 0.68 t

ha^{-1} (Table S2). Broad-sense heritability for GY was high across site-years, ranging from

0.58 to 0.94 (Table S2). Overall, correlations for GY between managed treatments were

moderately positive with the managed treatments that received similar levels of irrigation

(e.g., Optimal Bed with Optimal Flat; Moderate Drought with Severe Drought) showing

the strongest correlations (Figure S1).

Between 3 and 11 hyperspectral phenotyping time-points were collected within

each site year (Table 2). In most site-years, at least one hyperspectral phenotyping time-

point was collected during each of the three developmental growth stages. While the

managed treatments within a breeding cycle were often phenotyped on the same date, the

growth stage at the time of phenotyping often differed among the managed treatments

(Figure 2). During the 2015-16 cycle, technological issues with the camera prevented

early season phenotyping. As a result, no hyperspectral data were collected during the

VEG stage. The Optimal Bed, Optimal Flat, and Heat managed treatments had

hyperspectral phenotypes for the HEAD and GF stages only, while the Moderate Drought

and Severe Drought managed treatments had hyperspectral phenotypes for the GF stage

only due to early maturation of the crop under stress.

Broad-sense heritability estimates for the hyperspectral bands were generally

between 0.5 and 0.8 for most phenotyping time-points (Figure 3). Heritabilities were

relatively homogeneous across individual time-points and developmental growth stages.

In all site-years, the lowest heritabilities were observed for the hyperspectral bands

between 398 and 425 nm. Heritabilities in the 2013-14 Optimal Flat environment were

notably lower than in other site-years, falling between 0 and 0.25, though the GF time-

points were more heritable with values around 0.4. Low heritabilities (<0.2) were also

observed for the 9 March time-point in the 2015-16 Moderate Drought environment.

Correlations between the individual hyperspectral bands and GY ranged from -

0.68 to 0.64 across the 20 site-years and were most frequently the strongest above 500

nm (Figure 4). While there were some observable similarities in correlation patterns

among time-points taken during the same growth stage within the same site-year, patterns

were not uniform across site-years, breeding cycles, or managed treatments. In some site-

years, there were clear differences in correlation patterns between developmental growth

stages. For example, in the 2014-15 Optimal Bed site-year, correlations between GY and

hyperspectral reflectance in the 398-700 nm range were around 0 during the VEG stage

but become progressively more negative during the HEAD and GF stages, reaching

around -0.4. For the same managed treatment during the 2016-17 breeding cycle, distinct

differences in correlation patterns between the three developmental growth stages were

also observed; however, the patterns do not reflect those observed in 2014-15. Despite

irregularities in correlation patterns, the correlations between hyperspectral bands from

the GF stages and GY were, on average, stronger by 0.10 than for hyperspectral

phenotypes taken during the VEG and HEAD stages.

Model Prediction Accuracy

Within Site-Year: Five types of relationship matrices were used within models (4) and

(5) for within site-year prediction: genomic (G), pedigree (A), individual hyperspectral

phenotyping time-points (e.g. H_{10Jan}, H_{23Mar}, etc.), hyperspectral phenotype BLUEs

calculated from multiple time-points for each developmental growth stage (H_{VEG}, H_{HEAD},

H_{GF}), and hyperspectral phenotype BLUEs calculated using all available time-points

(H_{ALL}). Model (4) was used to assess accuracies using the G and A matrices for

prediction. Model (5) was fit to assess accuracies using the hyperspectral relationship

matrices.

Prediction accuracies of models using the individual hyperspectral time-point-

derived relationship matrices ranged from 0.00 to 0.75 with a mean of 0.42 (Figure 5A).

These results are similar to the prediction accuracies observed for G and A, which

recorded means of 0.41 and 0.42 and ranges of 0.19-0.60 and 0.19-0.67, respectively. In

18 and 17 of the 20 site-years, at least one time-point showed accuracies that were

equivalent or superior to G and A, respectively. In considering the most highly predictive

time-point in each site-year, 11 were recorded during the GF stage, followed by 6 and 3

in the HEAD and VEG stages, respectively, although the site-years had different numbers

of hyperspectral phenotyping time-points from each growth stage and some site-years

had no observations recorded during the VEG and HEAD stages.

The level of prediction accuracy for the individual hyperspectral time-points was

highly correlated with the strength of the relationship between hyperspectral reflectance

and GY (Figure 6). For a given time-point within a site-year, the correlation between

hyperspectral reflectance and GY was taken for each of the 62 wavelengths. The average

of the absolute values of those correlations, or their relative magnitudes, was then

compared to the level of prediction accuracy for that time-point. For the 71 time-

point/site-year combinations, the average magnitude of the correlation between

hyperspectral reflectance and GY explained 51 percent of the variation in prediction

accuracy, though the two characters exhibited a non-linear relationship (Figure 6A).

Correlations beyond 0.25 between hyperspectral reflectance and GY provided little

marginal increase in prediction accuracy. When GY was corrected for DTHD, this

relationship was linear and the correlation between reflectance and GY explained 26

percent of the variation in prediction accuracy (Figure 6B).

In considering accuracies using the hyperspectral phenotype BLUEs for each

growth stage, H_{GF} had the highest prediction accuracies in 11 of the 20 site-years,

followed by H_{HEAD} in 8 site-years and H_{VEG} in 1 site-year. Combining multiple time-

points did not necessarily increase accuracy. The accuracies of H_{VEG}, H_{HEAD}, and H_{GF}

were comparable to the mean of the accuracies of the individual time-points. Likewise,

the accuracy of H_{ALL}, which combined all available time-points, was on average slightly

1. higher than the mean of the accuracies of the individual time-points but lower than the

2. most predictive time-point.

3.   While correcting for DTHD had negligible impacts on prediction accuracy with

4. respect to the G and A models, the accuracies of the hyperspectral reflectance models

5. decreased by 0.10 on average (Figure 5B). The greatest reductions were observed in the

6. Severe Drought and Moderate Drought treatments, which showed average decreases in

7. accuracy of 0.20 and 0.18, respectively. After the DTHD correction, at least one-time

8. point recorded prediction accuracies that were equivalent or superior to the accuracies of

9. G and A in 13 out of the 20 site-years,

10.

11. **Within Breeding Cycle/Across Managed Treatments:** For prediction across managed

12. treatments within a breeding cycle, the genomic, pedigree, and hyperspectral reflectance

13. matrices were tested individually in single-kernel models and in combination in multi-

14. kernel models (Figure 7). Single-kernel models included the following: genetic main

15. effects (G and A) from model (4) and hyperspectral reflectance main effects (H_{VEG},

16. H_{HEAD}, H_{GF}, and H_{ALL}) from model (5). Multi-kernel models were built by combining a

17. main effects kernel with a $G \times E$ interaction kernel in models (6) and (7). In model (6),

18. the $G \times E$ interaction kernel was also fit using G or A. These models are herein referred

19. to as G $G \times E$ and A $G \times E$. In model (7), the hyperspectral reflectance matrices were used

20. to estimate the $G \times E$ interaction kernel. These models are referred to as $G + H_{VEG}$, $A +$

21. H_{HEAD}, etc. according to the respective relationship matrices used to estimate the main

22. and $G \times E$ interaction effects. The H_{VEG}, $G + H_{VEG}$, and $A + H_{VEG}$ models were assessed

23. for 14 out of the 20 site-years due to the unavailability of hyperspectral data at the

vegetative stage in the remaining 6 site-years. Likewise, the H_{HEAD}, $G + H_{HEAD}$, and $A +$

H_{HEAD} models were assessed for 18 out of the 20 site-years. The remaining models with

H_{GF} and H_{ALL} were tested for all site-years.

When considering the single-kernel hyperspectral main effect models, 2013-14

Optimal Flat had prediction accuracies close to zero. The 2013-14 Optimal Flat site-year

had high levels of lodging. This may have affected the reflectance signatures of the crop

canopy. The hyperspectral reflectance phenotypes for 2013-14 Optimal Flat had lower

heritabilities than all other site-years. To summarize trends for the remaining 19 site-

years, the results from the single-kernel hyperspectral main effects models for 2013-14

Optimal Flat were removed as outliers. The accuracies of H_{VEG} averaged 0.37 across the

site-years where it was tested, which was similar to the accuracies of G (0.35) and A

(0.34) (Figure 7A). The H_{HEAD}, H_{GF}, and H_{ALL} showed slightly higher averages

accuracies of 0.44, 0.44, and 0.46, respectively. The greatest differences in prediction

accuracy between the hyperspectral reflectance-based models and the G and A models

were observed in the 2015-16 and 2016-17 breeding cycles. In these cycles, the H_{HEAD},

H_{GF}, and H_{ALL} models showed accuracies that were on average 0.22 greater than the G

and A models. When correcting all models for DTHD, mean accuracies of the H_{VEG}

(0.29), H_{HEAD} (0.34), H_{GF} (0.30), and H_{ALL} (0.34) models were more similar to the G

(0.33) and A (0.32) models (Figure 7B).

Expanding the G and A models to account for the $G \times E$ interactions in the G

$G{\times}E$ and A $G{\times}E$ improved prediction accuracies to a level of 0.43 on average, an

increase of 0.08 and 0.09 over the G and A models, respectively. These improvements in

accuracy were more pronounced during the 2016-17 breeding cycle, recording gains in

accuracy of 0.16 and 0.19 for **G G×E** and **A G×E**, respectively. Likewise, slightly

greater improvements were observed for the Severe Drought (gains of 0.13 and 0.16 for

G G×E and **A G×E**, respectively) managed treatment. Similar trends were observed

when correcting for DTHD, though accuracies were 0.04 lower on average.

The multi-kernel models that estimated the main effects using **G** or **A** and the

$G \times E$ interactions using hyperspectral reflectance gave the highest accuracies overall.

Hyperspectral reflectance matrices $\boldsymbol{H_{VEG}}$, $\boldsymbol{H_{HEAD}}$, $\boldsymbol{H_{GF}}$, and $\boldsymbol{H_{ALL}}$ had average accuracies

of 0.54, 0.56, 0.56, and 0.58, respectively, when integrated with **G** and average accuracies

of 0.53, 0.55, 0.55, and 0.57, respectively, when integrated with **A**. In most site-years, no

clear "best model" could be identified among the multi-kernel models within a site-year.

Accuracies were similar regardless of during which growth stage the hyperspectral data

were recorded or whether genomic markers or pedigrees were used to model the main

effects.

When compared to the **G G×E** and **A G×E** models, the use of hyperspectral

reflectance to estimate the $G \times E$ interactions increased prediction accuracies by an

average of 0.10 and 0.12, respectively,   and by an average of 0.06 and 0.08, respectively

after correcting for DTHD. Compared to the corresponding single-kernel hyperspectral

reflectance models, the integration of markers or pedigree with hyperspectral reflectance

in multi-kernel models increased prediction accuracies on an average by 0.11 and 0.17

respectively, and 0.14 and 0.19, respectively when correcting for DTHD.

Across Breeding Cycles/Within Managed Treatment: The models tested for prediction

across breeding cycles/within managed treatment were the same as those tested for

prediction within breeding cycle/across managed treatments (Figure 8). On average,

accuracies were 0.08 lower in this prediction scheme than for prediction within breeding

a cycle and across managed treatments.

For the single-kernel hyperspectral main effects models, H_{GF} in 2013-14 Heat and

all single-kernel H models for 2013-14 Optimal Flat showed prediction accuracies close

to zero. The 2013-14 Heat site-year was phenotyped once during the GF late in the

growing season (21 May) which was close to maturity. Figure 4 shows that the

correlation between hyperspectral reflectance and GY for that time-point is close to zero

for each of the 62 wavelengths. To consider average trends, these were removed as

outliers. Mean accuracies of the H_{VEG}, H_{HEAD}, H_{GF}, and H_{ALL} models were 0.37, 0.45,

0.45, and 0.46, respectively (Figure 8A). The accuracies of the G and A models were

slightly lower, recording averages of 0.25 and 0.28, respectively. When correcting for

DTHD, the mean accuracies of H_{VEG}, H_{HEAD}, H_{GF}, and H_{ALL} models were 0.28, 0.34,

0.32, and 0.34, respectively (Figure 8B). While closer in performance to G and A, the

single-kernel hyperspectral main effects models still conferred an advantage in some site-

years.

Expanding the G and A models to account for the $G \times E$ interactions marginally

increased accuracies to 0.32 and 0.31 for the G $G\times E$ and A $G\times E$ models, respectively.

As with prediction within breeding cycles/across managed treatments, the highest

accuracies were observed for the multi-kernel models that estimated the main effects

using markers or pedigrees and the $G \times E$ interactions using hyperspectral reflectance.

Hyperspectral reflectance matrices H_{VEG} had an average accuracy of 0.39 when

integrated with G and 0.41 when integrated with A. H_{HEAD}, H_{GF}, and H_{ALL} had average

accuracies of 0.46, 0.46, and 0.48, respectively, when integrated with G or A. As before,

for most site-years, these multi-kernel models were observed to have similar levels of

accuracy within each site-year, irrespective of which hyperspectral reflectance matrix

was used to model the $G \times E$ interactions or whether genomic markers or pedigrees were

used to model the main effects.

When compared to the G $G×E$ and A $G×E$ models, estimating the $G \times E$

interactions using hyperspectral reflectance increased prediction accuracy by an average

of 0.12 and 0.14, respectively, and by an average of 0.07 and 0.08, respectively, when

correcting for DTHD. Average improvements over the corresponding single-kernel

hyperspectral reflectance main effects models ranged from 0.0.04 to 0.06 and from 0.06

to 0.08 after correcting for DTHD.

Mean accuracies and standard deviations of all models for the three prediction

schemes can be found in Files S1, S2, and S3 respectively.

15   DISCUSSION

We proposed a multi-kernel, across-environment GBLUP model that uses

relationship matrices derived from genomic markers, pedigrees, and aerial hyperspectral

reflectance phenotypes to estimate both genetic main effects and $G \times E$ interactions in the

context of a wheat breeding program. Our study found that deriving a relationship matrix

from high-dimensional hyperspectral reflectance phenotypes - as if they were genetic

markers - for use in GBLUP can be an effective approach for predicting GY in wheat and

in many situations resulted in predictions accuracies that were equivalent or superior to

the use of genomic markers or pedigrees. This is consistent with similar studies that used

linear and non-linear modeling approaches including ordinary least squares, partial least

squares, Bayesian shrinkage, and functional regression methods to predict grain in maize

and wheat with hyperspectral reflectance data (Aguate *et al.* 2017; Montesinos-Lopéz *et*

al. 2017a).

The heritabilities of the hyperspectral wavelengths and their correlations with GY

were not homogeneous across time-points, growth stages, breeding cycles or managed

treatments. Wavelengths between 398 and 425 nm were observed to have the lowest

heritabilities while those below 500 nm tended to have the weakest correlations with GY.

These results are consistent with a similar study of canopy reflectance in wheat (Hansen

et al. 2003) that found low signal-to-noise between 400 and 438 nm. According to

Mahlein *et al.* (2015), the hyperspectral imaging systems most commonly used in

agriculture have poor sensitivity to reflectance in the blue region of the light spectrum

(400-500 nm). Despite this, Montesinos-Lopéz *et al.* (2017a) showed that removing

wavelengths with low heritability did not improve prediction accuracies when using

hyperspectral reflectance to predict GY in wheat. Based on this result, we used all

hyperspectral wavelengths to build relationship matrices for prediction. However, the

blue region of the light spectrum contains important information on the optical properties

of plants, including the absorbance maxima of chlorophyll a, chlorophyll b, and

carotenoids (Lichenthaler, 1987; Horton *et al*. 1996). Further advancements in the ability

of hyperspectral imaging to accurately record reflectance in the blue region may

potentially improve prediction accuracies for grain yield.

When predicting line performance within a site-year of interest, most site-years

recorded at least one individual hyperspectral time-point with prediction accuracy

equivalent or superior to prediction with markers or pedigrees, suggesting that

hyperspectral reflectance phenotypes may be a useful alternative for generating

predictions when markers and pedigree are not available. We also found that combining

multiple time-points into hyperspectral BLUEs for a given growth stage generally did not

improve accuracy beyond the average accuracy across the individual time-points. In

practice, however, GY data would not be available to inform the identification of the

most predictive time-points. Therefore, phenotyping at multiple time-points throughout

the growing season to develop multi-time-point BLUEs may be an effective approach to

buffer against time-points with low prediction accuracy.

It is also possible that some information pertaining to plant growth and

development is lost when integrating multiple time-points using means-based approaches.

An alternative that may more fully capture temporal variations could be to model a

growth curve based on reflectance data for each line in each site-year, as in Verhulst *et al*.

(2011). A greater number of time-points than were measured in this study may be

required to accurately develop growth curves. In addition, statistical methods that can

integrate multiple curves from a range of hyperspectral bands may be needed. Further

research in this area should be performed to compare prediction accuracies of

hyperspectral reflectance-based growth curves versus those achieved in this study.

Correcting results for DTHD reduced accuracies when hyperspectral reflectance

was used, but not for markers or pedigrees. These results suggest that the hyperspectral

reflectance measurements are also capturing physiological parameters associated with

relative maturity. The greatest reductions in accuracy were observed in the Drought and

Severe Drought managed treatments, where GY is typically associated with earliness.

This is consistent with Rutkoski *et al.* (2014), which found that correcting for DTHD in

wheat GY reduced the prediction accuracies of multivariate GS models integrating NDVI

and canopy temperature measurements with markers and pedigrees. The CIMMYT bread

wheat breeding program maintains a high level of diversity for DTHD in germoplasm

development due to the wide target of geographic regions.. For breeding programs with

high levels of variation for DTHD, it may be advisable to perform a correction for DTHD

when predicting GY using hyperspectral reflectance so as to avoid indirect selection on

relative maturity.

   To test our proposed prediction approaches in a multi-environment context, we

developed two prediction schemes. In the first, predictions were performed within a

breeding cycle for genotypes that have been evaluated under some treatments but not

others. In the second, predictions were performed across breeding cycles onto genotypes

that were not previously evaluated. Overall, prediction accuracies in the second scheme

were lower than for the corresponding models in the first, which is consistent with

previous studies showing that predicting the performance of newly developed lines is

more challenging than the prediction of lines that have been evaluated in correlated

environments (Burgueño *et al.* 2012; Crossa *et al.* 2014).

   If markers or pedigrees are not available, our results showed that predicting GY

with hyperspectral reflectance alone in single-kernel models could provide similar results

in terms of accuracy. It should be noted that this study was conducted at the yield trial

stage of the breeding program when families contain fewer full-sibs and the variance due

to Mendelian sampling is low. As pedigrees do not account for Mendelian sampling, it is

possible that prediction with hyperspectral reflectance may be more advantageous than

pedigree-based GS at earlier stages of the breeding program in which families contain

greater numbers of full-sibs, though further study is needed to assess the ability of

hyperspectral reflectance to distinguish within-family variation.

The optimal prediction accuracies were achieved by building combined models

that used markers or pedigrees to model the genetic main effects and hyperspectral

reflectance to model the $G \times E$ interactions. However, in prediction across breeding

cycles within a managed treatment, the improvements in accuracy with the addition of

markers or pedigrees were marginal when compared to the corresponding single-kernel

hyperspectral reflectance models. This result was similar to Montesinos-Lopéz *et al.*

(2017b), which found that the addition of markers or pedigrees did not significantly

improve accuracy. The modest increases observed when predicting across breeding

cycles within a managed treatment may not justify the added investment of marker

genotyping.

When considering the optimal developmental growth stages during which to

collect hyperspectral reflectance data, there were some differences in accuracy among

growth stages when predicting with single hyperspectral phenotyping time-points within

a site-year. Overall, the time-points from the HEAD and GF stages showed slightly

higher accuracies than those collected during the VEG stage. However, when predicting

across site-years, the advantage of phenotyping later in the season was reduced,

particularly for the multi-kernel models that combined markers or pedigrees with

hyperspectral reflectance, where most often there was no significant difference in

accuracy among the different growth stages. Montesinos-Lopéz *et al.* (2017a) observed

clear optimal time-points for hyperspectral phenotyping in a multi-environment context.

However, their study represented a balanced scenario in which all lines in all

environments were phenotyped an equal number of times, and individual phenotyping

time-points could be used to predict across environments. While classifying time-points

according to the predominant growth stage at the time of phenotyping represents a simple

and efficient method for predicting across site-years, it is somewhat difficult to compare

results between growth stages. The numbers of time-points observed for each growth

stage were not consistent within and across site-years. In addition, the growth stage

classifications used were dependent on the predominant growth stage of the site-year and

did not reflect the variation in phenology within the site-year at the time of phenotyping.

These challenges may warrant further investigation into multi-environment prediction

when HTP datasets are unbalanced in the number of time-points observed and at which

stage of crop development those observations were recorded.

While our results suggest that hyperspectral reflectance data have the potential to

add value to a breeding program by providing accurate GY predictions, the data used in

this study were collected on large plot sizes that are suitable for measuring GY *per se*.

While our approach may provide breeding programs with GY predictions earlier in the

growing season, enabling more efficient allocation of resources at harvest, a potentially

greater benefit could come from utilizing this method for smaller plot sizes where there is

limited seed available to replicate and reliably assess yield. To address this, we are

currently evaluating the use of aerial HTP at the early generation stage when measuring

GY is not feasible.

The integration of highly dimensional data, such as hyperspectral reflectance, into

relationship matrices for use in GBLUP has several advantages. First, there is no

additional development in statistical software required to implement the proposed

models. Many options for fitting GBLUP such as "rrBLUP", "GAPIT", and "BGLR" are

currently available and are being increasingly used. The models proposed here can be

readily implemented in these or other existing software without requiring additional user

training. In addition, with the development of improved hyperspectral sensors that

register reflectance at wavelengths ranging from 400 to 2500 nm, it is likely that the

number of data points available for prediction will continue to increase. By using the

relationship matrix-based approach to GBLUP that we have proposed, these increases in

data dimensionality will neither influence the complexity of the GBLUP model, nor will

they increase the computation time of the model itself. This approach could also be useful

for integrating different types of highly dimensional phenotypes for prediction, such as

ionomics and metabolomics data (Riedelsheimer *et al.* 2012). Rincent *et al.* (2018)

recently showed that relationship matrices derived from near infrared spectroscopy

absorbance of winter wheat grains and powdered leaf tissue between 400 and 2500 nm

provided yield prediction accuracies that were superior to marker-based GBLUP

approaches. Further research should be performed to assess the potential for this

relationship matrix-based approach to be applied to other forms of biological data.

19   CONCLUSION

In this study, we have proposed a multi-kernel GBLUP model that uses genomic

marker-, pedigree-, and hyperspectral reflectance-derived relationships matrices to model

the genetic main effects and $G \times E$ interactions across environments within a bread wheat

breeding program. We have shown that deriving relationship matrices from hyperspectral

reflectance phenotypes can effectively predict GY in wheat within and across managed

treatments and breeding cycles. Accuracies when testing single-kernel models using

hyperspectral reflectance data alone are similar to those achieved with markers or

pedigrees. Our results also show that combining markers/pedigrees with hyperspectral

reflectance data in multi-kernel models can increase accuracies over single-kernel

approaches, but in some prediction scenarios, these increases were modest. We also

suggested a method for addressing the issue of unbalanced HTP datasets involving the

classification of time-points according to the predominant developmental growth stage

observed at the time of phenotyping. Accuracies of multi-kernel models were roughly

equivalent irrespective of the growth stage in which hyperspectral phenotyping was

performed. Further research on how best to leverage multi-temporal phenotypes when the

amount of data differs across site-years is needed. The methods we have proposed

provide a simple and computationally efficient approach for integrating highly

dimensional aerial HTP information into genomic selection and should be tested on other

forms of high dimensional biological data.

17   ACKNOWLEDGEMENTS

We are thankful to the National Science Graduate Research Fellowship under (Grant No.

DGE-1650441) and the U.S. Borlaug Fellowship in Global Food Security for supporting

the graduate studies of Margaret Krause. The work presented here was supported in part

by the Delivering Genetic Gain in Wheat project, supported by aid from the U.K.

Government's Department of International Development (DFID) and the Bill & Melinda

Gates Foundation (OPP113319). Partial support was also provided by the Agriculture and

22   REFERENCES

Aguate, F.M., S. Trachsel, L. González-Pérez, J. Burgueño, J. Crossa, *et al.*, 2017 Use of

hyperspectral image data outperforms vegetation indices in prediction of maize

yield. Crop Sci. 57: 2517-2524.

Andrade-Sanchez, P., M. A., Gore, J. T. Heun, K. R. Thorp, A. E. Carmo-Silva, *et al.*,

2014 Development and evaluation of a field-based high-throughput phenotyping

platform. Funct. Plant Biol. 41: 68-79.

Aparicio, N., D. Villegas, J. Casadesus, J. L. Araus, and C. Royo, 1999 Spectral

vegetation indices as nondestructive tools for determining durum wheat yield.

Agron. J. 92: 83-91.

Araus, J. L., and J. E. Cairns, 2014 Field high-throughput phenotyping: the new crop

breeding frontier. Trends Plant Sci.19: 52-61.

Bernardo, R., 2010 *Breeding for quantitative traits in plants*. Stemma Press, Woodbury,

Minnesota.

Burgueño, J., G. de los Campos, K. Weigel, and J. Crossa, 2012 Genomic prediction of

breeding values when modeling genotype $\times$ environment interaction using

pedigree and dense molecular markers. Crop Sci. 52: 707–719.

Cabrera-Bosquet, L., J. Crossa, J. von Zitzewitz, M. D. Serret, and J. L. Araus, 2012

High-throughput phenotyping and genomic selection: the frontiers of crop

breeding coverage. J. Integr. Plant Biol. 54: 312-320.

Crain, J., Y. Wei, J. Barker, S. M. Thompson, P. D. Alderman, *et al.*, 2016 Development

and deployment of a portable field phenotyping platform. Crop Sci. 56: 965-975.

Crain, J., S. Mondal, J. E. Rutkoski, R. Singh, J. A. and Poland, 2018 Combining high-

throughput phenotyping and genomic information to increase prediction and

selection accuracy in wheat breeding. Plant Genome 11: 170043.

Crossa, J., P. Pérez, J. Hickey, J. Burgueño, L. Ornella, *et al.*, 2014 Genomic prediction

in CIMMYT maize and wheat breeding programs. Heredity 112: 48-60.

Crossa, J., P. Pérez-Rodríguez, J. Cuevas, O. Montesinos-López, D. Jarquín, *et al.*, 2017

Genomic selection in plant breeding: methods, models, and perspectives. Trends

Plant Sci. 22: 961-975.

Daetwyler, H.D., Calus, M.P.L., Pong-Wong, R., de los Campos, G. and J.M. Hickey,

2013 Genomic prediction in animals and plants: simulation of data, validation,

reporting, and benchmarking. Genetics 193:347-365.

de los Campos, G., and P. Pérez-Rodríguez, 2014 Bayesian generalized linear regression.

R package version 1.0.5. URL https://CRAN.R-project.org/package=BGLR.

Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto, *et al.*, 2011 A robust,

simple genotyping-by-sequencing (GBS) approach for high diversity species.

PLoS One 6: e19379.

Endelman, J. B., 2011 Ridge regression and other kernels for genomic selection with R

package rrBLUP. Plant Genome 4: 250-255.

Endelman, J. B., and J. L. Jannink, 2012 Shrinkage estimation of the realized relationship

matrix. G3 Genes Genomes Genet 2: 1405-1413.

Furbank, R. T., and M. Tester, 2011 Phenomics – technologies to relieve the phenotyping

bottleneck. Trends Plant Sci. 16: 635-644.

Gilmour, A. R., B. J. Gogel, B. R. Cullis, S. J. Welham, and R. Thompson, 2014 ASReml

user guide release 4.1 structural specification. VSN International, Hemel

Hempstead, UK.

Govaerts, Y. M., M. M. Verstraete, B. Pinty, and N. Gobron, 1999 Designing optimal

spectral indices: a feasibility and proof of concept study. Int. J. Remote Sens.

1999: 1853-1873.

Haghighattalab, A., L. González-Pérez, S. Mondal, D. Singh, D. Schinstock, *et al.*, 2016

Application of unmanned aerial systems for high throughput phenotyping of large

wheat breeding nurseries. Plant Methods: 12, 35.

Hansen, P.M., and J.K. Schjoerring, 2003 Reflectance measurement of canopy biomass

and nitrogen status in wheat crops using normalized difference vegetation indices

and partial least squares regression. Remote Sens. Environ. 86:542-553.

Heslot, N., J. Jannink, and M. E. Sorrells, 2013 Using genomic prediction to characterize

environments and optimize prediction accuracy in applied breeding data. Crop

Sci. 53: 921-933.

Horton, P., Ruban, A.V., and R.G. Walters, 1996 Regulation of light harvesting in green

plants. Annu. Rev. Plant Physiol. Plant Mol. Biol. 47:655-684.

Jarquín, D., J. Crossa, X. Lacaze, P. Du Cheyron, J. Daucourt, *et al.*, 2014 A reaction

norm model for genomic selection using high-dimensional genomic and

environmental data. Theor. Appl. Genet. 127: 595-607.

Labus, M., G. Nielsen, R. Lawrence, R. Engel, and D. Long, 2002 Wheat yield estimates

using multi-temporal NDVI satellite imagery. Int. J. Remote Sens. 23: 4169-4180.

Li, L., Q. Zhang, and D. Huang, 2014 A review of imaging techniques for plant
     phenotyping. Sensors 14: 20078-20111.

Lichtenthaler, H.K., 1987 Chlorophylls and carotenoids: pigments of photosynthetic
     biomembranes. Methods Enzymol. 148:350-382.

López-Cruz, M., J. Crossa, D. Bonnett, S. Dreisigacker, J. Poland, *et al.*, 2015 Increased
     prediction accuracy in wheat breeding trials using markers × environment
     interaction genomic selection model. G3 Genes Genomes Genet. 5: 569-582.

Mahlein, A.K., Hammersley, S., Oerke, E.C., Dehne, H.W., Goldbach, H., *et al.*, 2015
     Supplemental blue LED lighting array to improve signal quality in hyperspectral
     imaging of plants. Sensors (Basel) 15:12834-12840.

McLaren, C. G., L. Ramos, C. Lopez, and W. Eusebio, 2000 Applications of the
     geneaology manegment system, pp. 5.8-5.13 in *International Crop Information*
     System. Technical Development Manual, version VI, edited by C. G. McLaren,
     J.W. White and P.N. Fox. CIMMyT, México: CIMMyT and IRRI.

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard, 2001 Prediction of total genetic
     value using genome-wide dense marker maps. Genetics 157: 1819-1829.

Montesinos-López, O. A., A. Montesinos-López, J. Crossa, G. de los Campos, G.
     Alvarado, *et al.*, 2017a Predicting grain yield using hyperspectral reflectance in
     wheat breeding data. Plant Methods 13: 4.

Montesinos-López, A., O. A. Montesinos-López, J. Cuevas, W. A. Mata-López, J.
     Burgueño, *et al.*, 2017b Genomic Bayesian functional regression models with
     interactions for predicting wheat grain yield using hyperspectral reflectance. Plant
     Methods 13: 62.

Pauli, D., S. C. Chapman, R. Bart, C. Topp, C. J. Lawrence-Dill, *et al.,* 2016 The quest for understanding phenotypic variation via integrated approaches in the field environment. Plant Physiol. 172: 622-634.

Pérez-Rodríguez, P., J. Crossa, J. Rutkoski, J. Poland, R. Singh, *et al.*, (2017). Single-step genomic and pedigree genotype × environment interaction models for predicting wheat lines in international envrionments. Plant Genome 10. doi:10.3835/plantgenome2016.09.0089.

Poland, J., J. Endelman, J. Dawson, J. Rutkoski, S. Wu, *et al.*, 2012 Genomic selection in wheat breeding using genotyping-by-sequencing. Plant Genome 5: 103-113.

R Core Team, 2018 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Riedelsheimer, C., A. Czedik-Eysenberg, C. Grieder, J. Lisec, F. Technow, *et al.* 2012 Genomic and metabolic prediction of complex heterotic traits in hybrid maize. Nat. Genet. 44: 217-220.

Rincent, R., Charpentier, J.P., Faivre-Rampant, P., Paux, E., Le Gouis, J., et al., 2018 Phenomic selection is a low-cost and high-throughput method based on indirect predictions: proof of concept on wheat and poplar. G3 Genes Genomes Genet. 8 doi: 10.1534/g3.118.200760.

Rutkoski, J. E., J. A. Poland, S. Mondal, E. Autrique, L. González-Pérez, L., et al., 2016 Canopy temperature and vegetation indices from high-throughput phenotyping improve accuracy of pedigree and genomic selection for grain yield in wheat. G3 Genes Genomes Genet. 6: 2799-2808.

Sun, J., J. E. Rutkoski, J. A. Poland, J. Crossa, J. Jannink, *et al*., 2017 Multitrait, random

regression, or simple repeatability model in high-throughput phenotyping data

improve genomic prediction for wheat grain yield. Plant Genome 10: 1-12.

Verhulst, N., B. Govaerts, V. Nelissen, K. D. Sayre, J. Crossa, *et al.*, 2011 The effect of

tillage, crop rotation and residue management on maize and wheat growth and

development evaluated with an optical sensor. Field Crops Res. 120: 58-67.

Viña, A., A. A. Gitelson, D. C. Rundquist, G. Keydan, B. Leavitt, *et al.*, 2003 Monitoring

maize (*Zea mays* L.) phenology with remote sensing. Agron. J. 96: 1139-1147.

Viña, A., A. A. Gitelson, A. L. Nguy-Robertson, and Y. Peng, 2011 Comparison of

different vegetation indices for the remote assessments of green leaf area index of

crops. Remote Sens. Environ. 115: 3468-3478.

White, J.W., P. Andrade-Sanchez, M. A. Gore, K. F. Bronson, T. A. Coffelt, *et al.* 2012

Field-based phenomics for plant genetics research. Field Crops Res. 133: 101-

112.

Zhang, X., M. A. Friedl, C. B. Schaaf, A. H. Strahler, J. C. F. Hodges, *et al*., 2003

Monitoring vegetation phenology using MODIS. Remote Sens. Environ. 84: 471-

475.

1   **TABLE 1** Description of the field management conditions in each of the five managed treatments

2   sown at CIMMYT Research Station in Ciudad Obregon, México

MANAGED TREATMENT	PLANTING DATE	PLOT TYPE	PLOT DIMENSIONS	IRRIGATION METHODS
Optimal Bed	Late November/ Early December	Two beds with 3 rows per bed	2.8m × 0.8m	Five furrow irrigations
Optimal Flat	Late November/ Early December	Flat sown plot with 6 rows	4.0m × 1.3m	Five flood irrigations
Moderate Drought	Late November/ Early December	Two beds with 3 rows per bed	2.8m × 0.8m	Two furrow irrigations
Severe Drought	Late November/ Early December	Flat sown plot with 6 rows	4.0m × 1.3m	Three minimal irrigations through drip
Heat	Late February	Two beds with 3 rows per bed	2.8m × 0.8m	Five furrow irrigations

1 **TABLE 2** Classification of hyperspectral phenotyping time-points as vegetative (VEG), heading

2 (HEAD), and grain fill (GF) phenological stages

BREEDING CYCLE	PHENOTYPING DATE	OPTIMAL BED	OPTIMAL FLAT	MODERATE DROUGHT	SEVERE DROUGHT	HEAT
2013-14	10 Jan	-	VEG	VEG	-	-
	17 Jan	-	VEG	VEG	VEG	-
	30 Jan	VEG	VEG	VEG	VEG	-
	07 Feb	VEG	VEG	VEG	VEG	-
	14 Feb	HEAD	VEG	HEAD	HEAD	-
	19 Feb	HEAD	HEAD	HEAD	HEAD	-
	27 Feb	GF	HEAD	GF	GF	-
	11 Mar	GF	GF	GF	GF	-
	17 Mar	GF	GF	GF	-	-
	25 Apr	-	-	-	-	VEG
	02 May	-	-	-	-	HEAD
	07 May	-	-	-	-	HEAD
	21 May	-	-	-	-	GF
2014-15	10 Jan	VEG	VEG	VEG	VEG	-
	19 Jan	VEG	VEG	VEG	VEG	-
	04 Feb	VEG	VEG	VEG	VEG	-
	09 Feb	VEG	VEG	HEAD	HEAD	-

	25 Feb	HEAD	HEAD	GF	GF	-
	10 Mar	GF	GF	GF	GF	-
	15 Mar	GF	GF	GF	GF	-
	23 Mar	GF	GF	GF	-	-
	26 Mar	GF	GF	-	-	-
	07 Apr	-	GF	-	-	-
	14 Apr	-	-	-	-	VEG
	24 Apr	-	-	-	-	HEAD
	28 Apr	-	-	-	-	HEAD
	06 May	-	-	-	-	GF
2015-16	26 Feb	HEAD	HEAD	GF	GF	-
	03 Mar	HEAD	HEAD	GF	GF	-
	09 Mar	GF	GF	GF	GF	-
	15 Mar	GF	GF	GF	GF	-
	22 Mar	GF	GF	GF	GF	-
	02 May	-	-	-	-	HEAD
	09 May	-	-	-	-	GF
	14 May	-	-	-	-	GF
2016-17	10 Jan	VEG	VEG	VEG	VEG	-
	17 Jan	VEG	VEG	VEG	VEG	-

	23 Jan	VEG	VEG	VEG	VEG	-
	02 Feb	VEG	VEG	VEG	HEAD	-
	10 Feb	VEG	VEG	HEAD	HEAD	-
	16 Feb	HEAD	VEG	HEAD	GF	-
	22 Feb	HEAD	HEAD	GF	GF	-
	15 Mar	GF	GF	-	GF	-
	29 Mar	GF	-	-	-	-
	05 Apr	GF	-	-	-	-
	11 Apr	GF	-	-	-	-
	02 May	-	-	-	-	HEAD
	22 May	-	-	-	-	GF
	31 May	-	-	-	-	GF

2 **FIGURE 1** Boxplot of BLUEs for wheat grain yield (t ha^{-1}) in each of the 20 observed site-years

FIGURE 2 A graphical representation of the unbalanced nature of the hyperspectral reflectance

phenotypic data

Four site-years are represented: 2014-15 Optimal Flat, 2014-15 Severe Drought, 2016-17

Optimal Flat and 2016-17 Severe Drought. The histograms represent heading dates. Each

dashed line corresponds to a hyperspectral phenotyping date colored according to the

predominant growth stage of the lines at the time of phenotyping.

2 **FIGURE 3** Broad-sense heritabilities of the hyperspectral wavelengths for each phenotyping time-

3 point within each site-year

5 Each boxplot represents the distribution of broad-sense heritability values for the 62

6 hyperspectral wavelengths observed. The colors correspond to the developmental growth stage

7 of the site-year at the time of hyperspectral phenotyping.

2 **FIGURE 4** Empirical correlations between grain yield and hyperspectral reflectance BLUEs within

3 each site-year

5 Each line represents a phenotyping time-point and the correlation between grain yield and the 62

6 hyperspectral wavelengths observed. Lines are colored according to the predominant

7 developmental growth stage of the site-year at the time of hyperspectral phenotyping.

1   **A. WITHOUT DTHD CORRECTION:**

4   **B. WITH DTHD CORRECTION:**

7   **FIGURE 5** Within site-year prediction accuracies, with and without correction for DTHD

9   Accuracy is expressed as the average Pearson's correlation between predictions and observed

10   BLUPs for GY across 20 random TRN-TST partitions. Results shown according to the type of

1 relationship matrix tested: Genomic (**G)**, pedigree (**A**), individual hyperspectral time-points (e.g.

2 **H10Jan**, **H.23Mar**, etc.), hyperspectral BLUEs for each developmental growth stage (**H.VEG**,

3 **H.HEAD**, **H.GF**), and hyperspectral BLUEs across all time-points (**H.ALL**). The color corresponds

4 to the predominant developmental growth stage of the site-year at the time of phenotyping. Error

5 bars are the standard deviation of prediction accuracy for the 20 random partitions

A. WITH DTHD CORRECTION:

B. WITH DTHD CORRECTION:

FIGURE 6 The relationship between individual hyperspectral time-point prediction accuracy and

the average magnitude of the correlations between hyperspectral bands and GY

The absolute values of the correlations between hyperspectral bands and GY were calculated

and then averaged across the 62 bands for each time-point within each site-year. Plotted on the

x-axis, this represents the average strength of the relationship between hyperspectral reflectance

1   and GY for each time-point within each site-year. The y-axis shows the prediction accuracy for

2   each individual hyperspectral time-point in within site-year prediction.

1   **A. WITHOUT DTHD CORRECTION:**

3   B. **WITH DTHD CORRECTION:**

5   **FIGURE 7** Within breeding cycle/across managed treatments prediction accuracies, with and

6   without correction for DTHD

Accuracy is expressed as the average Pearson's correlation between predictions and observed

BLUPs for GY across 20 random TRN-TST partitions. In each partition, the TRN set consisted of

all records from four out of the five managed treatments within the breeding cycle plus 20% of

records from the fifth managed treatment. Single-kernel models tested were genetic main effects

only (**G** or **A**) and hyperspectral reflectance main effects plus GxE (**H.VEG, H.HEAD, H.GF,**

H.ALL). Multi-kernel models assessed were genetic main effects plus genetic GxE (**G GxE**, **A**

GxE) and genetic main effects plus hyperspectral reflectance GxE (e.g., **G + H.VEG**, **A + H.ALL**,

etc.). The color corresponds to the developmental growth stage of the site-year at the time of

phenotyping. Error bars are the standard deviation of prediction accuracy for the 20 random

partitions

1   **A. WITHOUT DTHD CORRECTION:**

2

3   **B. WITH DTHD CORRECTION:**

4

5   **FIGURE 8** Across breeding cycles/within managed treatment prediction accuracies, with and

6   without correction for DTHD

Accuracy is expressed as the average Pearson's correlation between predictions and observed

BLUPs for GY across 20 random TRN-TST partitions. Single-kernel models tested were genetic

main effects only (**G** or **A**) and hyperspectral reflectance main effects plus GxE (**H.VEG, H.HEAD,**

H.GF, H.ALL). Multi-kernel models assessed were genetic main effects plus genetic GxE (**G**

GxE, **A GxE**) and genetic main effects plus hyperspectral reflectance GxE (e.g., **G + H.VEG**, **A +**

H.ALL, etc.). The color corresponds to the developmental growth stage of the site-year at the

time of phenotyping. Error bars are the standard deviation of prediction accuracy for the 20

random partitions.