CRISPR/Cas9-mediated gene deletion of the *ompA gene* in symbiotic

Enterobacter* impairs biofilm formation and reduces gut colonization of *Aedes

***aegypti* mosquitoes**

Shivanand Hegde[1,4], Pornjarim Nilyanimit[2], Elena Kozlova[1], Hema P. Narra[1], Sanjeev

K. Sahni[1], Grant L. Hughes[3,4,#]

[1] Department of Pathology, University of Texas Medical Branch, Galveston, TX, USA.

[2] Center of Excellence in Clinical Virology, Chulalongkorn University, Bangkok,

Thailand.

[3] Department of Pathology, Institute for Human Infections and Immunity, Center for

Tropical Diseases, Center for Biodefense and Emerging Infectious Disease. University

of Texas Medical Branch, Galveston, TX, USA.

[4] Current address: Department of Vector Biology and Department of Parasitology,

Liverpool School of Tropical Medicine, Pembroke Place, Liverpool, UK.

[#]Corresponding author. Email: grant.hughes@lstmed.ac.uk

Running head: CRISPR/Cas9 engineering of a gut bacterium.

Abstract

Background: Symbiotic bacteria are pervasive in mosquitoes and their presence can influence many host phenotypes that affect vectoral capacity. While it is evident that environmental and host genetic factors contribute in shaping the microbiome of mosquitoes, we have a poor understanding regarding how bacterial genetics affects colonization of the mosquito gut. The CRISPR/Cas9 gene editing system is a powerful tool to alter bacterial genomes facilitating investigations into host-microbe interactions but has yet to be applied to insect symbionts.

Methodology/Principal Findings: To investigate the role of bacterial factors in mosquito biology and in colonization of mosquitoes we used CRISPR/Cas9 gene editing system to mutate the outer membrane protein A (*ompA*) gene of an *Enterobacter* symbiont isolated from *Aedes* mosquitoes. The *ompA* mutant had an impaired ability to form biofilms and poorly infected *Ae. aegypti* when reared in a mono-association under gnotobiotic conditions. In adults the mutant had a significantly reduced infection prevalence compared to the wild type or complement strains, while no differences in prevalence were seen in larvae, suggesting bacterial genetic factors are particularly important for adult gut colonization. We also used the CRISPR/Cas9 system to integrate genes (antibiotic resistance and fluorescent markers) into these symbionts genome and demonstrated that these genes were functional in vitro and in vivo.

Conclusions/Significance: Our results shed insights onto the role of *ompA* gene in host-microbe interactions in *Ae. aegypti* and confirm that CRISPR/Cas9 gene editing can be employed for genetic manipulation of non-model gut microbes. The ability to use this technology for site-specific integration of genes into the symbiont will facilitate the

development of paratransgenic control strategies to interfere with arboviral pathogens

such Chikungunya, dengue, Zika and Yellow fever viruses transmitted by *Aedes*

mosquitoes.

Author summary

Microbiota profoundly affect their host but few studies have investigated the role of

bacterial genetics in host-microbe interactions in mosquitoes. Here we applied the

CRISPR/Cas9 gene editing system to knock out a membrane protein in *Enterobacter*,

which is a dominant member of the mosquito microbiome. The mutant strain lacked the

capacity to form biofilms, infected larvae and adults at lower titers, and had a reduced

prevalence in adults. The lower prevalence in adults, but not larvae, likely reflects the

difference in the modes of bacterial acquisition from the larval water of these two life

stages. Importantly from an applied perspective, we also demonstrated that this editing

technology can be harnessed for site-specific integration of genes into the bacterial

chromosome. In proof-of-principle studies we integrated either a fluorescent protein or

gene conferring antibiotic resistance into the bacterial genome and showed these

transgenes were functional in mosquitoes. The specificity, flexibility, and simplicity of

this editing approach in non-model bacteria will be useful for developing novel symbiotic

control strategies to control arthropod-borne disease.

Keywords: CRISPR/Cas9, Gut microbe, Gut colonization, Biofilm,

Paratransgenesis, Paratransgenic

Introduction.

Mosquitoes harbor a community of microbes within their guts. In general, the gut-associated microbiome of mosquitoes tends to have low species richness but can differ greatly between individuals and habitats [1-8]. Importantly, these microbes can modulate many host phenotypes, several of which can influence vectorial capacity [9-11]. As such, it is imperative that we understand how the microbiome is acquired and maintained within mosquito vectors. While environmental factors unquestionably influences the mosquito microbiome composition and abundance [2-4, 8], studies are elucidating the role of microbial interactions[5, 7, 12, 13] and host genetic factors [14-18] in shaping the microbiome. However, we have a poor understanding of bacterial factors that influence colonization of the mosquito gut and this is likely an underappreciated force influencing host-microbe interactions in mosquitoes.

In other invertebrates, several bacterial genes have been implicated in gut colonization. For example, a genome wide screen exploiting transposon-sequencing found a suite of genes from the bacterium *Snodgrasselia* involved in colonization of the honey bee gut [19]. These bacterial genes were classified into the broad categories of extracellular interactions, metabolism and stress response [19]. Knock out of a purine biosynthesis gene in *Burkholderia* impaired biofilm formation and reduced bacterial colonization rates in a bean bug [20]. Biofilm formation was also shown to play a role in virulence of pathogenic *Pseudomonas* in artificial infections of *Drosophila,* with strains that lacked the capacity to form biofilms being more virulence to the host, while a hyperbiofilm strain was less virulent than the WT strain [21]. In other blood feeding invertebrates, bacterial

genetics also appears critical for host colonization. Knockout of the type II secretion

system in *Aeromonas veronii* reduced infection in *Hirudo verbena* leeches [22]. In

Tsetse flies, the outer-membrane protein A (*ompA*) gene of *Sodalis glossinidius* is

essential for symbiotic interactions [23]. *Sodalis* mutants lacking the *ompA* gene poorly

colonized the fly gut compared to the wild type (WT) symbionts [23] and the mutant

strain also had a reduced capacity to form biofilms [24]. Heterologous expression of the

ompA gene from pathogenic *Escherichia coli* in *Sodalis* mutants induced mortality in the

fly implicating this gene as a virulence factor in pathogenic bacteria [23]. Taken

together, these studies suggest that bacterial genetic factors are critical for host

colonization of invertebrates and that biofilm formation facilitates symbiotic associations

in insects.

In mosquitoes, few studies have investigated how bacterial genetics affect gut

colonization. However, evidence from experimental evolution studies suggests bacterial

genetics plays a critical role. In two separate studies, *Enterobacter* was selected for

increased persistence in the gut of *Anopheles gambiae* mosquitoes, the major malaria

vector in sub-Saharan Africa, by repeatedly infecting mosquitoes with strains that

persisted in the gut for longer periods of time [25, 26]. Transcriptomics comparisons of

effective and ineffective colonizers in liquid media identified 41 genes that were

differentially expressed between these two strains [26], further implicating the

importance of bacterial genetics in mosquito infection, however the role of these genes

in colonization of the mosquito gut has not been resolved. In a separate study, *in vitro*

screening of a transposon mutant library of *Enterobacter* identified a *waaL* gene mutant

that was insensitive to oxidative stress [27]. The *waaL* gene encodes an O antigen

ligase which is needed for attachment of the O antigen to lipopolysaccharide and the

mutant was found to have lower rates of colonization of the midguts of *Anopheles*

mosquitoes [27].

Gene knockouts approaches in bacteria provide compelling evidence of the role of

bacterial genes in host-microbe interactions [22-24, 27-29]. In general, most studies use

transposon mutagenesis for gene knockout, which requires screening of the mutant

library. A targeted gene knockout approach is highly desirable to investigate the

functionality of bacterial genes in host-microbe interactions. In the past few years, the

CRISPR/Cas9 gene editing system has been employed to modify bacterial genomes

[30-32] . While much of the work has been done in model bacterial species [31-37],

editing approaches have expanded into non-model bacterial systems [38-43]. Despite

this expansion, the approach has been used less frequently used for host-associated

microbes [39, 44], and rarely for arthropod symbionts. In the vector biology field, gene

knockout approaches can be used to interrogate the role of bacterial genes responsible

for host-microbe interactions, while the ability to integrate genes into the bacterial

symbiont genome has great potential for applied paratransgenic control strategies [10,

45-47]. Previously, manipulation of non-model symbionts that associate with insect

vectors have has been accomplished by plasmid transformation [48-55] or stable

transformation of the genome using transposons or integrative plasmids [56-63], but the

use of CRISPR/Cas9 gene editing in insect gut symbionts has yet to be accomplished.

For paratransgenic strategies, stable site-specific integration of transgenes into the

symbiont genome is critical, and as such, the application of CRISPR/Cas9 gene editing

technology to non-model bacteria that associate with insect vectors will stimulate

research in this field.

We therefore undertook studies to develop CRISPR/Cas9 genome editing approaches

in an *Enterobacter* species isolated from *Aedes* mosquitoes. We used the <u>S</u>carless

<u>C</u>as9 <u>A</u>ssisted <u>R</u>ecombineering (no-SCAR) method to disrupt the *ompA* gene of the

non-model *Enterobacter* species [35]. After characterization of the mutant *in vitro*, we

examined the role of the *ompA* gene in host-microbe interactions by re-infecting

bacteria into mosquito in a mono-association. To demonstrate that the CRISPR/Cas9

gene-editing system could be useful for applied symbiotic control approaches we

inserted genes conferring antibiotic resistance or a fluorescent protein into the bacterial

genome and re-infected the altered strains back into mosquitoes. Our result sheds

insights into the role of the *ompA* gene in host-microbe interactions in *Ae. aegypti* and

confirm that CRISPR/Cas9 gene editing can be a powerful tool for genetic manipulation

of native gut-associated microbes of mosquitoes.

Results

***Enterobacter* biofilm formation in *Ae. aegypti* guts**

Over the course of conducting mono-axenic infections in *Ae. aegypti* mosquitoes with an *Enterobacter* symbiont, we repeatedly observed a conglomeration of bacterial cells in the gut that was indicative of a biofilm (Figure 1, Figure S1 A-C). This formation of bacteria has a similar appearance to biofilms observed in the guts of other insects [21, 24]. No bacteria were observed in the intestinal track of *Ae. aegypti* when infected with *E. coli* (Figure 1, S1 D-F), although as seen previously, infection with this bacterium enabled mosquito development [64]. We therefore examined the role of bacterial genetics in biofilm formation and host colonization of this gut-associated bacterium of *Aedes* mosquitoes. While several genes have been implicated in biofilm formation [21, 24], we chose to knockout the *ompA* gene of *Enterobacter* given that this gene has been demonstrated to influence biofilm formation and gut colonization of *Sodalis* [23, 24]*,* an *Enterobacteriaceae* symbiont of Tsetse flies. We used the CRISRP/Cas9 genome editing system to mutate the symbionts genome.

Genome editing in *Enterobacter* bacteria isolated from mosquitoes.

To edit the *Enterobacter* isolate that resides within the gut of *Aedes* mosquitoes, we employed the no-SCAR gene editing approach that had been developed in *E. coli* [35]. To optimize the approach in our hands, we performed initial experiments in *E. coli* to delete a ~1 kb region of the *ompA* gene (Figure 2A). As the no-SCAR approach exploits the λ-Red recombineering system to repair double stranded breaks, we transformed bacteria with a double stranded DNA template that had regions of homology flanking the

gRNA site (250 bp for each arm). Using this approach, we successfully deleted a 1001

bp fragment of the *ompA* gene. From the colonies we screened, we saw an editing at a

frequency of 6.25% (N = 48) (Figure 2A). For *Enterobacter*, we altered our editing

procedure to delete a 598 bp fragment from the *ompA* gene. This was done to enhance

the efficiency of obtaining mutants [35] and accommodate the PAM site which was at a

different location in the *ompA* gene in *Enterobacter*. Using a donor template designed

for the *Enterobacter ompA* gene that had similar length flanking homology arms as the

previous experiment done in *E. coli*, we obtained mutant knockouts at a rate of 32% (N

= 50) (Figure 2B). For both bacterial species, Sanger sequencing across the integration

site indicated the deletion occurred at the expected loci in the bacterial genome (Figure

2C; S1 Appendix).

Characterization of the *Enterobacter ompA* mutant.

We quantified the growth rates of the ∆*ompA* mutant in comparison to the WT

Enterobacter and the ∆*ompA/ompA* complement in liquid LB media. We saw minimal

differences in the growth between the WT, the ∆*ompA* mutant or the ∆*ompA/ompA*

complement (Figure 3A). To examine the stability of the deletion, we subcultured the

∆*ompA* mutant on LB media for 10 generations and performed PCR to amplify across

the deletion. At alternative generations PCR analysis indicated the deletion was present

indicating genomic stability at this site (Figure 3B).

Previously, *ompA* has been shown to be important in biofilm formation as *Sodalis*

deletion mutants were unable to form biofilms [24]. As such we characterized *in vitro*

biofilm formation using the crystal violet (CV) biofilm assay. After visual inspection, it

was clear the Δ*ompA* mutant had distinctly less biofilm deposition compared to either

the WT or the Δ*ompA/ompA* complement (Figure 3C), and after quantification and

normalization to account for any difference in growth between the strains, biofilm

formation was confirmed to be significantly different between the Δ*ompA* mutant and the

WT (Figure 3D; Tukey's multiple comparisons test, P < 0.0001) or Δ*ompA/ompA*

complement (Tukey's multiple comparisons test, P < 0.0001), while there was no

significant differences between the WT and the Δ*ompA/ompA* complement (Tukey's

multiple comparisons test P = 0.2).

The role of *ompA* gene in mosquito infection.

To examine the importance of the *ompA* gene on bacterial colonization of mosquitoes,

we infected *Ae. aegypti* mosquitoes in a mono-association under gnotobiotic conditions

[64]. This infection method was used to avoid other gut-associated microbes influencing

host colonization rates [7] and it also enable straightforward quantification of introduced

bacteria by measuring colony forming units (CFUs). In larvae we saw a significant

reduction in bacterial titer in the mutant compared to both the WT (Kruskal-Wallis test; P

< 0.01) and the Δ*ompA/ompA* complement (Kruskal-Wallis test; P < 0.05) (Figure 4A).

Similarly, in adults, there was a significant reduction in bacterial infection in the Δ*ompA*

mutant compare to either the WT or Δ*ompA/ompA* complement (Kruskal-Wallis test; P <

0.001) (Figure 4B). While no significant changes were seen in the prevalence of

infection (number of mosquitoes infected) in the larval stage (Figure 4C, Fisher's exact

test; WT compared to Δ*ompA* P = 0.24 and Δ*ompA* compared to Δ*ompA/ompA* P =

0.24), in adults, the prevalence of infection was significantly different (Figure 4D,

Fisher's exact test; WT compared to Δ*ompA* P < 0.0001 and Δ*ompA* compared to

Δ*ompA/ompA* P < 0.0001), with only 45% of adults infected by the Δ*ompA* mutant

compared to 95% and 88% by the WT and Δ*ompA/ompA* complement, respectively. We

also examined the growth rates of mosquitoes administered with the WT, Δ*ompA*

mutant and Δ*ompA/ompA* complement. No significant differences were seen in the time

to pupation (Figure 5A) or percentage of first instar larvae that reached adulthood

(Figure 5B) between any of the strains.

Integration of genes into the *Enterobacter* chromosome.

We undertook experiments to demonstrate the CRISPR/Cas9 gene-editing approaches

can be used to integrate genes into the chromosome of non-model bacteria that

associate with mosquitoes. We created two independent transgenic strains that had

either, a gene encoding mCherry fluorescence or a gene encoding resistance to the

antibiotic gentamicin, inserted into the bacterial chromosome. These genes were

integrated into the genome using the same gRNA that was used for deletional

mutagenesis (Table S1), and as such, these insertions also disrupted the *ompA* gene.

Sequencing across the integration site indicated the insertion of these genes occurred

within the *ompA* gene and thereby disrupted its function (Figure 6A and 6D). Continual

subculturing was undertaken for both strains and molecular analysis indicated the

stability of these lines for ten generations (Figure 6B and 6E). To demonstrate the

integrated genes were functional, we observed expression of mCherry fluorescence and

successfully cultured the strain containing gentamicin resistance on plates containing

the antibiotic (Figure 6C and 6F). Finally, we infected these transgenic strains into

mosquitoes to demonstrate that these strains were able to colonize the mosquito gut

and functionality of the integrated gene was confirmed by observing fluorescence or by

rearing the *Enterobacter ompA*::gentamicin strain in mosquitoes administered sugar

supplemented with gentamicin. Fluorescent bacteria were observed in the gut of

mosquitoes while no signal was seen in controls (WT *Enterobacter* infected mosquitoes)

(Figure 5G). The *Enterobacter ompA*::gentamicin was successfully rescued from

mosquitoes reared on gentamicin and was seen to stably infect mosquitoes over time at

a density of 1×10^{4} CFUs/mosquito. Consistent with our previous finding (Figure 4B), the

WT bacteria initially infected mosquitoes at higher titers (T test; day 0 $P < 0.001$).

However, at 4 days post infection (dpi), the total bacterial load of culturable microbes in

mosquitoes supplemented with WT *Enterobacter* was significantly reduced when reared

on sugar supplemented with antibiotic (T test; day 4 $P < 0.05$), and no CFUs were

recovered after at 6 dpi (T test; day 6 $P < 0.001$) (Figure 6H).

Discussion.

We harnessed the CRISPR/Cas9 gene editing system to create knockout mutants in an *Enterobacter* gut symbiont of *Ae. aegypti* mosquitoes enabling us to examine the role of bacterial genetics, specifically the *ompA* gene, in biofilm formation and gut colonization. A deletion of the *ompA* gene of *Enterobacter* decreased bacterial colonization of the mosquito host at both the larval and adult stages after infection in a mono-association. Strikingly, we found this effect was most pronounced in adult mosquitoes with more than half of the mosquitoes not possessing any culturable mutants, while there was no difference in prevalence of infection between the mutant and WT bacteria in larvae. The reduced prevalence of mutant bacteria in adults likely reflects differences in microbial colonization of each mosquito life stage. Larvae are continually subjected to bacteria in the larval water habitat while adults only have a short time frame to acquire bacteria from the aquatic environment immediately after eclosion, when they are thought to imbibe a small amount of larval water which seeds the gut with microbiota [65]. Our data shows greater variation in colonization of the adult stage between the mutant and WT strains, indicating that the *ompA* gene, and potentially bacterial factors in general, may be critical for colonization of the adult gut. These findings are also consistent with other sequence-based studies, that indicate adult stages have greater variability in species composition of their microbiota, while the microbiome of immature stages is similar to the microbiota in larval water habitat [2-5, 8, 66].

Overall, our findings are similar to studies done in Tsetse flies whereby an *ompA* mutant of *Sodalis*, an *Enterobacteriaceae* symbiont, has impaired biofilm formation and

reduced colonization rates [23, 24]. These studies, in conjunction with our work,

suggests that the *ompA* gene is imperative for symbiotic associations within dipterans. It

also suggests that biofilm formation may be a strategy employed by bacteria to colonize

the gut of insects. In pathogenic infections in mammals, biofilms enable bacteria to

colonize new niches, promote infection and are associated with virulence [67] . Although

less is known regarding the importance of biofilm formation in insects, in an artificial

Pseudomonas-Drosophila infection model, biofilm formation was associated with

virulence and host survival [21]. In a natural symbiotic association between bean bugs

and *Burkholderia*, disruption of a purine biosynthesis gene in the bacterium also reduce

biofilm formation and colonization of the insect [20] In mosquitoes, gut biofilm formation

could also have implications for vector competence as *Chromobacterium*, which was

isolated from *Aedes* mosquitoes, produced molecules that inhibited dengue virus only

when grown *in vitro* as a biofilm but not when grown in a planktonic state [68]. Despite

this, it was unknown if biofilm formation occurred *in vivo* in the mosquito [68]. Our data

provide evidence that biofilms occur within the gut of mosquitoes and facilitate host

colonization.

While we have shown that the *ompA* gene of *Enterobacter* is important for host

colonization, we see no evidence that deletion of this gene alters mosquito development

or growth rates. This is in contrast to the *Riptortus-Burkholderia* symbiosis whereby

mutation of the *purT* gene in *Burkholderia* resulted in reduced growth rates and

reduction in body weight of the host compared to insects that were infected with the WT

bacterium [20]. The difference in our study to the findings in the *Riptortus-Burkholderia*

symbiosis could be related to different requirements of the bean bug compared to the

mosquito host as well as the different genes mutated in the symbionts. Our findings are

consistent with a previous study in *Ae. aegypti* whereby an *ompA* mutant of *E. coli* did

not influence growth reared in a mono-association [69]. Using a similar gnotobiotic

system that exploits the ability to sterilize mosquito eggs and rescue development by

nutritional supplementation, several recent reports describe approaches to create

bacteria-free mosquitoes [69, 70]. Here, we reared mosquitoes in a mono-association

where they were only subjected to *Enterobacter.* However*,* more than half the adult

mosquitoes inoculated with the $\Delta ompA$ mutant were not infected by bacteria, as

evidenced by the inability to culture bacteria from these insects. Nevertheless, these

mosquitoes had similar development and growth rates compared to mosquito

possessing WT bacteria. The use of mutant bacteria that rescue development but have

an impaired ability to colonize mosquitoes may provide a simple means to create axenic

adult mosquitoes.

CRISPR/Cas9 gene editing has revolutionized genetic approaches in model and non-

model bacteria [31-43]. However, there has been limited use of this technology in

symbiotic microbes of arthropods. Here we demonstrate that editing approaches

functional in *E. coli* can be easily applied with minimal adaptation to phylogenetically

related symbiotic bacteria that associate within the guts of mosquitoes. The application

of CRISPR/Cas9 genome editing to gut-associated bacteria of mosquitoes has

significant applied potential. Paratransgenesis strategies are being evaluated in a range

of medical and agricultural systems to mitigate pathogen transmission from insect

vectors, however, most approaches engineer symbionts by plasmid transformation [49-

55, 71] and where genome integration has been accomplished in symbionts [58-61], it

has often been done with technologies that did not allow for site specific integration.

Paratransgenic approaches suitable for use in the field will need to stably integrate

genes into the bacterial genome in a manner that does not compromise bacterial

fitness. Exploiting the flexibility and specificity of the CRISPR/Cas9 to integrate genes in

intergenic regions of the bacterial chromosome will undoubtedly be beneficial for these

applied approaches.

In summary, we have demonstrated that the CRISPR/Cas9 gene editing system can be

applied to symbiotic bacteria that associate with eukaryotic hosts to interrogate the role

of bacterial genes in host-microbe associations. We created knockout and knockin

mutants by deleting and disrupting the *ompA* gene of *Enterobacter*. The knockout

mutant displayed a reduced ability to form biofilms and colonize the gut of *Ae. aegypti*

mosquitoes in a mono-association, demonstrating bacterial genetic factors are

important determinants that influence colonization of mosquito guts. *Aedes* mosquitoes

are becoming powerful systems to investigate the genetics of host-microbe interactions

given the scientific community has simple and efficient approaches to alter both the

microbes (this study) and mosquito host genome [72, 73] at their disposal, as well as

methods to create mono-associated mosquito lines[7, 64]. Finally, rapid, efficient, and

site specific gene editing approaches for gut bacteria that associate with mosquitoes will

facilitate the development of novel paratransgenic approaches to control arthropod-

borne disease [57].

Experimental procedures

Bacterial and mosquito strains. *E. coli* BL21(DE3) (NEB) and an *Enterobacter* strain previous isolated from a lab-reared colony of *Ae. albopictus* (Galveston) mosquitoes [7] were used in this study. Cultures were grown in liquid LB media at 37°C with the appropriate antibiotic unless stated otherwise. Mosquitoes were reared in the UTMB insectary under conventional conditions or in mono-associations (described below).

CRISPR gene editing. Editing the *ompA* gene of *E. coli* and *Enterobacter* was complete as described in Reisch and Prather [35]. The protospacer sequence for the *ompA* gene was designed using the CHOPCHOP [74, 75], and cloned into pKDsgRNA-ack plasmid [35] directly upstream of gRNA scaffold using REPLACR mutagenesis protocol [76]. Two protospacer sequences were designed for each gene and the one which had lower escape rate after plating with or without aTC (S1 Table). The plasmids were acquired from Addgene (S2 Table; Addgene plasmid 62655 and 62654). The resulting plasmids pKDsgRNA-Ec-*ompA* and pKDsgRNA-Ent-*ompA* were Sanger sequenced to confirm insertion of protospacer sequence. These plasmids were then transformed into either *E. coli* or *Enterobacter* containing the pCas9-CR4 plasmid. Transformants were selected at 30°C on LB agar plate containing spectinomycin (50 µg/mL), chloramphenicol (34 µg/mL), and with or without anhydrotetracycline (aTC) at 100ng/mL. Colonies from the –aTC plate were grown overnight in LB broth with the appropriate antibiotic at 30°C. A 1:100 diluted overnight culture was (grown until 0.4 OD_{600}) supplemented with 1.2% arabinose to induce the expression of λ-Red recombinase. Cells were then transformed with 1-1.5 µg of double stranded donor DNA

that flanked the PAM site for homologous recombination. Donor DNA was created by

either PCR amplification or by gene synthesis (Genewiz). Regardless of the method of

construction, each donor had flanking regions of 250 bp homologous to the target DNA.

The resulting colonies were screened for mutations by colony PCR with primers flanking

the integration site and positive clones were Sanger sequenced (S3 Table). Positive

colonies were grown in LB broth and genomic DNA was isolated. For further validation,

the flanking regions of deletion or insertions were amplified and the PCR product

Sanger sequenced.

Stability of insertion. The stability of the knockout Δ*ompA* mutant and the knockin

ompA::gentamicin and *ompA*::mCherry strains was assessed in LB medium. The

*omp*A::mCherry and knockout Δ*ompA* mutant cultures were grown for 10 passages in

LB broth. At each passage 40 µl of culture was transferred into 4ml fresh LB medium.

The *ompA*::gentamicin strain was grown with or without gentamicin (50 µg/mL).

Genomic DNA was isolated from the 0, 2, 4, 6, 8 and 10th subculture and PCR that

amplified across the integration site was performed.

Complementation of *ompA* mutant. Functional rescue of the *omp*A mutation was

achieved by complementing the mutant with the WT gene. The WT *omp*A gene was

amplified from *Enterobacter* genomic DNA and cloned into the pRAM-mCherry vector[7]

and thereby creating pRAM-mCherry-*Ent-OmpA.* The Sanger sequence-verified

plasmid was transformed into the Δ*ompA* mutant, thereby generating the Δ*ompA/ompA*

complement strain. Colonies that acquired the plasmid were selected on LB plates

containing kanamycin (50 µg/mL).

***In vitro* characterization of *Enterobacter* strains.** To assess the impact of the gene

deletion on bacterial growth the WT, $\Delta ompA$ mutant and $\Delta ompA/ompA$ complement

were grown in LB broth and the density of bacteria (OD_{600}) was quantified by

spectrophotometer. A 1:100 dilution of an overnight culture was inoculated into a 5 ml

LB broth in 50 ml tube and incubated at 37°C for 24 hrs. At 2, 4, 6, 8, 10, 12 and 24

hours growth was recorded at OD_{600}. The biofilm assay was performed as described

previously [77, 78]. Briefly, biofilm formation by *Enterobacter* strains was quantified on

polystyrene microtiter plates after 72 h of incubation at 37°C by CV staining. Three

independent experiments were performed, and the data were represented as CV OD_{570}

after normalizing by CFUs.

Mosquito infections. Mono-association in *Ae. aegypti* mosquitoes were done using

gnotobiotic infection procedure [7, 64],+ with slight modifications. Briefly, mosquito eggs

were sterilized for 5 min in 70% ethanol, 3 min 3% bleach+0.01% Coverage Plus NPD

(Steris Corp.), 5 min in 70% ethanol then rinsed three times in sterile water. Eggs were

vacuumed hatched for 30-45 min and left overnight at room temperature to hatch any

remaining eggs. Exactly twenty L1 larvae were transferred to T175 flask containing 60

ml of sterile water and fed on alternative days with 60 µl of fish food (1 µg/µl). Larvae

were inoculated with 1×10^{7}/ml of either the WT *Enterobacter*, the $\Delta ompA$ mutant or the

$\Delta ompA/ompA$ complement. The WT and $\Delta ompA$ strains were transformed with the

pRAM-mCherry plasmid that conferred resistance to kanamycin (but did not possess a

functional *ompA* gene). L4 larvae were collected, washed three times with 1X PBS, and

then homogenized in 500 µl of 1X PBS and 50 µl of homogenate was plated on LB agar

containing 50 µg/mL kanamycin. Similarly, adult mosquitoes were collected 3-4 days

post emergence and bacterial infection was quantified in the same manner as larvae. In

order to assess the growth of the mosquitoes, time to pupation and growth rate were

observed. Time to pupation was determined by quantifying the number of pupae each

day post hatching, while survival to adulthood was calculated by quantifying the number

of L1 larvae that reached adulthood. The experiment was repeated three times.

Knock-in mutants were administered to adult *Ae. aegypti* in a sugar meal. Three to four

day old mosquitoes were fed with 1×10^{7} of WT or the $\Delta ompA$::gentamicin strain for

three days in 10% sucrose solution. After three days, mosquitoes were either

administered sugar supplemented with gentamicin (50 µg/mL) or sugar without

antibiotic. CFUs were determined at days 0, 2, 4, and 6 dpi by plating homogenized

mosquitoes (N=10) on LB agar. Similarly, the $\Delta ompA$::mCherry and WT *Enterobacter*

were fed to mosquitoes and midguts were dissected to assess the colonization of

bacteria in the tissue. For visualization of bacteria, midguts were fixed in 1%

paraformaldehyde (PFA) in 1X PBS for 30 minutes and permeabilized with 0.01% Triton

X-100 in 1X PBS for 20 min. The tissues were stained with 1:250 diluted Phalloidin

(Sigma) for 20 minutes and samples were washed twice with 1X PBS for 10 minutes.

Finally, midguts were then stained with 1:500 diluted DAPI (Invitrogen) for 10 min.

Samples were transferred to slides and mounted with ProLong™ Gold Antifade

(Invitrogen). The slides were observed under Revolve FL (ECHOLAB).

Acknowledgements.

We would like to thank the UTMB insectary core for providing mosquitoes. GLH was

supported by NIH grants (R21AI138074, R21AI124452 and R21AI129507), a University

of Texas Rising Star award, the Western Gulf Center of Excellence for Vector-borne

Diseases (CDC grant CK17-005), the Robert J. and Helen Kleberg Foundation and the

Gulf Coast Consortia. This work was also supported by a James W. McLaughlin

postdoctoral fellowship at the University of Texas Medical Branch to SH and a Dutsadi

Piphat Scholarship and Thailand Research Chair Grant from NSTDA (p-15-50004) to

PN. We also wish to thank Dr Ulrike Munderloh (University of Minnesota) for the kind

gift of the pRAM plasmid.

Competing interests.

The authors declare no competing interests.

Figure legends.

Figure 1. Midgut infection of *Enterobacter* and *E. coli* in mono-associations of *Aedes* mosquitoes. *Enterobacter* forms a biofilm in the gut of *Aedes aegypti* mosquitoes (left) while no bacteria were observed in the gut of mosquitoes reared with *E. coli* under gnotobiotic conditions (right). Bacteria possessed a plasmid expressing mCherry. Blue – host nuclei. Green – host actin cytoskeleton stained with phalloidin. The scale bar is 70 μm.

Figure 2. CRISPR/Cas9 genome editing in bacteria. A schematic of the editing approach and screening of putative mutants in *E. coli* (A) and *Enterobacter* (B). A ~1kb fragment of *E. coli* BL21(DE3) was deleted using no-SCAR protocol. The 250 bp of left arm (LA) and right arm (RA) was assembled to generate 500 bp donor DNA. The transformants were screened via colony PCR with primers binding in regions flanking the deletion. Similar to strategy employed in *E. coli*, knockout of the *ompA* gene from *Enterobacter* isolated from the mosquito gut was created by deleting the 598 bp fragment. The grayed area indicates the PAM site in the *ompA* gene and arrow shows cleavage site in the genome. (C) The sequence of the *ompA* mutation in *E. coli* and *Enterobacter* was confirmed by Sanger sequencing. The sequence above the gene within the dotted line has been deleted. The chromatogram shows the 10 bp flanking the deletion.

Figure 3. In vitro characterization of the *ompA* mutation. The *Enterobacter* ΔompA mutant had a similar growth rate compared to both the WT and the Δ*ompA/ompA* complement in liquid LB media (A). The stability of mutant was evaluated *in vitro* by continuous subculturing in LB medium (B). Genomic DNA of alternative subcultures was used as template for PCR using gene specific primers that amplified across the deletion. Two separate gel images were merged to create the figure 2B. Passage 8 was run on a separate gel to passages 0 – 6. Biofilm formation was assessed using the CV biofilm assay for the WT, Δ*ompA* mutant and the Δ*ompA/ompA* complement (C). Quantification of the relative biofilm formation normalized by the number of bacteria per well (D).

Figure 4. The Δ*ompA* mutant poorly infected mosquitoes. Infection of *Enterobacter* strains (WT, Δ*ompA* mutant and Δ*ompA/ompA* complement) reared in a mono-association using a gnotobiotic rearing approach for larvae (A) and adults (B). L4 and 3-4 days post emergence adults were screened for bacterial load by plating on LB media to quantify the bacteria. The prevalence of infection (number of mosquitoes infected) between the treatments was calculated comparing number of infected to uninfected larvae (C) or adults (D).

Figure 5. The Δ*ompA* mutant does not affect growth rates or development of mosquitoes. The growth rate (time to pupation) (A) and development (percentage of L1 larvae to reach adulthood) (B) was observed in mosquitoes infected with *Enterobacter*

strains (WT, Δ*ompA* mutant and Δ*ompA/ompA* complement) reared in a mono-

association.

Figure 6. Integration of mCherry and gentamicin into the *Enterobacter* genome.

Sanger sequence across the integration site, stability of the inserted gene and *in vitro*

expression of the inserted gene for the Δ*ompA*::mCherry (A-C) and the Δ*ompA*::

gentamicin (D-F) strains. The chromatogram shows the sequence spanning the inserted

sites. Strains were continually subcultured for 10 passages and PCR was done to

examine the stability of the insert (B; Δ*ompA*::mCherry plus WT; E Δ*ompA*::gentamicin

passaged with (ab+) or without (ab-) gentamicin plus WT). mCherry fluorescence or

ability to grow on selective media containing gentamicin confirmed the expression of the

transgene *in vitro*. Mosquitoes were inoculated with the *Enterobacter* strains to confirm

expression of the transgene *in vivo*. Dissected midgut infected with Δ*ompA*::mCherry

(left) or negative control (right; WT bacteria without expression plasmid) (G). Midguts

were stained with phalloidin (green) and DAPI (blue). The scale bar is 30 μM. The WT

and Δ*ompA*::gentamicin *Enterobacter* strains were fed to adult mosquitoes for 3 days in

a sugar meal before gentamicin was administered to mosquitoes in a sugar meal (H).

Mosquitoes were collected every second day and CFUs assessed. Pairwise

comparisons were conducted at each time point using a T test (* - $P < 0.05$, *** $P <$

0.001, **** $P < 0.0001$).

S1. Figure. Midgut infection of *Enterobacter* and *E. coli* in mono-associations of

***Aedes* mosquitoes.** Different gut tissue locations showing the conglomeration of

bacterial cells when infected in mono-association in *Aedes* mosquitoes with

Enterobacter (A-C). However, infection of *E. coli* did not show any infection in the gut of

the mosquitoes (D-F). The pictures were taken from different field of view of the gut

dissected from different mosquitoes.

References.

1.   Osei-Poku J, Mbogo CM, Palmer WJ, Jiggins FM. Deep sequencing reveals extensive variation in the gut microbiota of wild mosquitoes from Kenya. Molecular Ecology. 2012;21(20):5138-50. doi: 10.1111/j.1365-294X.2012.05759.x.

2.   Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, et al. Midgut microbiota of the malaria mosquito vector *Anopheles gambiae* and interactions with *Plasmodium falciparum* infection. PLoS Pathogens. 2012;8(5):e1002742. doi: 10.1371/journal.ppat.1002742.

3.   David MR, Santos LMBD, Vicente ACP, Maciel-de-Freitas R. Effects of environment, dietary regime and ageing on the dengue vector microbiota: evidence of a core microbiota throughout *Aedes aegypti* lifespan. Memórias do Instituto Oswaldo Cruz. 2016;111(9):577-87. doi: 10.1590/0074-02760160238.

4.   Muturi EJ, Kim C-H, Bara J, Bach EM, Siddappaji MH. *Culex pipiens* and *Culex restuans* mosquitoes harbor distinct microbiota dominated by few bacterial taxa. Parasites & Vectors. 2016;9(1):18. doi: 10.1186/s13071-016-1299-6.

5.   Hughes GL, Dodson BL, Johnson RM, Murdock CC, Tsujimoto H, Suzuki Y, et al. Native microbiome impedes vertical transmission of *Wolbachia* in *Anopheles* mosquitoes. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(34):12498-503. doi: 10.1073/pnas.1408888111.

6.   Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are essential for development but vary greatly between local habitats. Molecular Ecology. 2016;25(22):5806-26. doi: 10.1111/mec.13877.

7.   Hegde S, Khanipov K, Albayrak L, Golovko G, Pimenova M, Saldaña MA, et al.

Microbiome interaction networks and community structure from laboratory-reared and

field-collected *Aedes aegypti*, *Aedes albopictus*, and *Culex quinquefasciatus* mosquito

vectors. Frontiers in Microbiology. 2018;9:715. doi: 10.3389/fmicb.2018.02160.

8.   Wang Y, Gilbreath TM, Kukutla P, Yan G, Xu J. Dynamic gut microbiome across

life history of the malaria mosquito *Anopheles gambiae* in Kenya. PLoS ONE.

2011;6(9):e24767. doi: 10.1371/journal.pone.0024767.

9.   Weiss BL, Aksoy S. Microbiome influences on insect host vector competence.

Trends in Parasitology. 2011;27(11):514-22. doi: 10.1016/j.pt.2011.05.001.

10.   Saldaña MA, Hegde S, Hughes GL. Microbial control of arthropod-borne disease.

Memórias do Instituto Oswaldo Cruz. 2017;112(2):81-93. doi: 10.1590/0074-

02760160373.

11.   Hegde S, Rasgon JL, Hughes GL. The microbiome modulates arbovirus

transmissionin mosquitoes. Current Opinion in Virology. 2015;15:97-102. doi:

10.1016/j.coviro.2015.08.011.

12.   Audsley MD, Ye YH, McGraw EA. The microbiome composition of *Aedes aegypti*

is not critical for *Wolbachia*-mediated inhibition of dengue virus. PLoS Neglected

Tropical Diseases. 2017;11(3):e0005426. doi: 10.1371/journal.pntd.0005426.

13.   Muturi EJ, Bara JJ, Rooney AP, Hansen AK. Midgut fungal and bacterial

microbiota of *Aedes triseriatus* and *Aedes japonicus* shift in response to La~Crosse

virus infection. Molecular Ecology. 2016;25(16):4075-90. doi: 10.1111/mec.13741.

14.   Kumar S, Molina-Cruz A, Gupta L, Rodrigues J, Barillas-Mury C. A

peroxidase/dual oxidase system modulates midgut epithelial immunity in *Anopheles*

gambiae. Science. 2010;327(5973):1644-8. doi: 10.1126/science.1184008.

15.   Pang X, Xiao X, Liu Y, Zhang R, Liu J, Liu Q, et al. Mosquito C-type lectins

maintain gut microbiome homeostasis. Nature Microbiology. 2016;1(5):16023. doi:

10.1038/nmicrobiol.2016.23.

16.   Short SM, Mongodin EF, MacLeod HJ, Talyuli OAC, Dimopoulos G. Amino acid

metabolic signaling influences *Aedes aegypti* midgut microbiome variability. PLoS

Neglected Tropical Diseases. 2017;11(7):e0005677-29. doi:

10.1371/journal.pntd.0005677.

17.   Stathopoulos S, Neafsey DE, Lawniczak MKN, Muskavitch MAT, Christophides

GK. Genetic dissection of *Anopheles gambiae* gut epithelial responses to *Serratia*

marcescens. PLOS Pathogens. 2014;10(3):e1003897. doi:

10.1371/journal.ppat.1003897.s015.

18.   Xiao X, Yang L, Pang X, Zhang R, Zhu Y, Wang P, et al. A Mesh-Duox pathway

regulates homeostasis in the insect gut. Nature Microbiology. 2017;2(5):17020. doi:

10.1038/nmicrobiol.2017.20.

19.   Powell JE, Leonard SP, Kwong WK, Engel P, Moran NA. Genome-wide screen

identifies host colonization determinants in a bacterial gut symbiont. Proceedings of the

National Academy of Sciences of the United States of America. 2016:201610856. doi:

10.1073/pnas.1610856113.

20.   Kim JK, Kwon JY, Kim S-K, Han SH, Won YJ, Lee J-H, et al. Purine

biosynthesis, biofilm formation, and persistence of an snsect-microbe gut symbiosis.

Appl Environ Microbiol. 2014;80(14):4374-82. doi: 10.1128/AEM.00739-14.

21.   Mulcahy H, Sibley CD, Surette MG, Lewenza S. *Drosophila melanogaste*r as an

animal model for the study of *Pseudomonas aeruginosa* biofilm infections in vivo. PLOS

Pathogens. 2011;7(10):e1002299-14. doi: 10.1371/journal.ppat.1002299.

22.   Maltz M, Graf J. The type II secretion system is essential for erythrocyte lysis and

gut colonization by the leech digestive tract symbiont *Aeromonas veronii*. Applied and

Environmental Microbiology. 2011;77(2):597-603. doi: 10.1128/AEM.01621-10.

23.   Weiss BL, Wu Y, Schwank JJ, Tolwinski NS, Aksoy S. An insect symbiosis is

influenced by bacterium-specific polymorphisms in outer-membrane protein A.

Proceedings of the National Academy of Sciences of the United States of America.

2008;105(39):15088-93. doi: 10.1073/pnas.0805666105.

24.   Maltz MA, Weiss BL, O apos Neill M, Wu Y, Aksoy S. OmpA-mediated biofilm

formation is essential for the commensal bacterium *Sodalis glossinidius* to colonize the

tsetse fly gut. Applied and Environmental Microbiology. 2012;78(21):7760-8. doi:

10.1128/AEM.01858-12.

25.   Riehle MA, Jacobs-Lorena M. Using bacteria to express and display anti-parasite

molecules in mosquitoes: current and future strategies. Insect Biochemistry and

Molecular Biology. 2005;35(7):699-707. doi: 10.1016/j.ibmb.2005.02.008.

26.   Dennison NJ, Saraiva RG, Cirimotich CM, Mlambo G, Mongodin EF, Dimopoulos

G. Functional genomic analyses of *Enterobacter*, *Anopheles* and *Plasmodium* reciprocal

interactions that impact vector competence. Malaria Journal. 2016;15(1):425. Epub 3.

doi: 10.1186/s12936-016-1468-2.

27.   Pei D, Jiang J, Yu W, Kukutla P, Uentillie A, Xu J. The *waaL* gene mutation

compromised the inhabitation of *Enterobacter sp. Ag1* in the mosquito gut environment.

Parasites & Vectors. 2015:1-10. doi: 10.1186/s13071-015-1049-1.

28.   Enomoto S, Chari A, Clayton AL, Dale C. Quorum sensing attenuates virulence

in *Sodalis praecaptivus*. Cell Host & Microbe. 2017;21(5):629-36.e5. doi:

10.1016/j.chom.2017.04.003.

29.   Dale C, Young SA, Haydon DT, Welburn SC. The insect endosymbiont *Sodalis*

glossinidius utilizes a type III secretion system for cell invasion. Proceedings of the

National Academy of Sciences. 2001;98(4):1883-8. doi: 10.1073/pnas.021450998.

30.   Selle K, microbiology RBTi, 2015. Harnessing CRISPR–Cas systems for

bacterial genome editing. Elsevier. doi: 10.1016/j.tim.2015.01.008.

31.   Sander JD, Joung JK. CRISPR-Cas systems for editing, regulating and targeting

genomes. Nature Biotechnology. 2014;32(4):347-55. doi: 10.1038/nbt.2842.

32.   Barrangou R, van Pijkeren JP. Exploiting CRISPR-Cas immune systems for

genome editing in bacteria. Current Opinion in Biotechnology. 2016;37:61-8. doi:

10.1016/j.copbio.2015.10.003.

33.   Jiang Y, Chen B, Duan C, Sun B, Yang J, Yang S. Multigene editing in the

Escherichia coli genome via the CRISPR-Cas9 system. Applied and Environmental

Microbiology. 2015;81(7):2506-14. doi: 10.1128/AEM.04023-14.

34.   Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA. RNA-guided editing of

bacterial genomes using CRISPR-Cas systems. Nature Biotechnology. 2013;31(3):233-

9. doi: 10.1038/nbt.2508.

35.   Reisch CR, Prather KLJ. The no-SCAR (Scarless Cas9 Assisted

Recombineering) system for genome editing in *Escherichia coli.* Scientific Reports.

2015;5:15096. doi: 10.1038/srep15096.

36.   Li Y, Lin Z, Huang C, Zhang Y, Wang Z, Tang Y-j, et al. Metabolic engineering of

Escherichia coli using CRISPR–Cas9 meditated genome editing. Metabolic

Engineering. 2015;31:13-21. doi: 10.1016/j.ymben.2015.06.006.

37.   Ronda C, Pedersen LE, Sommer MOA, Nielsen AT. CRMAGE: CRISPR

Optimized MAGE Recombineering. Scientific Reports. 2016;6(1):1200. doi:

10.1038/srep19452.

38.   Tong Y, Robertsen HL, Blin K, Weber T, Lee SY. CRISPR-Cas9 Toolkit for

Actinomycete Genome Editing. Methods in molecular biology 2018;1671(1):163-84. doi:

10.1007/978-1-4939-7295-1_11.

39.   Oh J-H, van Pijkeren JP. CRISPR-Cas9-assisted recombineering in *Lactobacillus*

reuteri. Nucleic Acids Research. 2014;42(17):e131-e. doi: 10.1093/nar/gku623.

40.   Mougiakos I, Bosma EF, Weenink K, Vossen E, Goijvaerts K, van der Oost J, et

al. Efficient genome editing of a facultative thermophile using mesophilic spCas9. ACS

Synthetic Biology. 2017;6(5):849-61. doi: 10.1021/acssynbio.6b00339.

41.   Li K, Cai D, Wang Z, He Z, Chen S. Development of an efficient genome editing

tool in *Bacillus licheniformis* ssing CRISPR-Cas9 nickase. Applied and Environmental

Microbiology. 2018:AEM.02608-17. doi: 10.1128/AEM.02608-17.

42.   Jiang Y, Qian F, Yang J, Liu Y, Dong F, Xu C, et al. CRISPR-Cpf1 assisted

genome editing of *Corynebacterium glutamicum*. Nat Commun. 2017;8:15179. Epub

2017/05/05. doi: 10.1038/ncomms15179.

43.   Cobb RE, Wang Y, Zhao H. High-efficiency multiplex genome editing of

Streptomyces species using an engineered CRISPR/Cas system. ACS Synth Biol.

2015;4(6):723-8. Epub 2014/12/03. doi: 10.1021/sb500351f.

44.   Waller MC, Bober JR, Nair NU, Beisel CL. Toward a genetic tool development

pipeline for host-associated bacteria. Current Opinion in Microbiology. 2017;38:156-64.

doi: 10.1016/j.mib.2017.05.006.

45.   Wilke ABB, Marrelli MT. Paratransgenesis: a promising new strategy for

mosquito vector control. Parasites & Vectors. 2015;8(1):391-19. doi: 10.1186/s13071-

015-0959-2.

46.   Arora AK, Douglas AE. Hype or opportunity? Using microbial symbionts in novel

strategies for insect pest control. Journal of insect physiology. 2017;103:10-7. doi:

10.1016/j.jinsphys.2017.09.011.

47.   Ricci I, Damiani C, Capone A, DeFreece C, Rossi P, Favia G.

Mosquito/microbiota interactions: from complex relationships to biotechnological

perspectives. Current Opinion in Microbiology. 2012;15(3):278-84. doi:

10.1016/j.mib.2012.03.004.

48.   Beard CB, Mason PW, Aksoy S, Tesh RB, Richards FF. Transformation of an

insect symbiont and expression of a foreign gene in the Chagas' disease vector

Rhodnius prolixus. American Journal of Tropical Medicine and Hygiene.

1992;46(2):195-200.

49.   Bisi DC, Lampe DJ. Secretion of anti-Plasmodium effector proteins from a natural

Pantoea agglomerans isolate by using PelB and HlyA secretion signals. Applied and

Environmental Microbiology. 2011;77(13):4669-75. doi: 10.1128/AEM.00514-11.

50.   Hughes GL, Allsopp PG, Webb RI, Yamada R, Iturbe-Ormaetxe I, Brumbley SM,

et al. Identification of yeast associated with the planthopper, *Perkinsiella saccharicida*:

potential applications for Fiji leaf gall control. Current Microbiology. 2011;63(4):392-401.

doi: 10.1007/s00284-011-9990-5.

51.   Medina F, Li H, Vinson SB, Coates CJ. Genetic transformation of midgut bacteria

from the red imported fire ant (*Solenopsis invicta*). Current Microbiology.

2009;58(5):478-82. doi: 10.1007/s00284-008-9350-2.

52.   Bextine B, Lauzon C, Potter S, Lampe D, Miller TA. Delivery of a genetically

marked *Alcaligenes sp.* to the glassy-winged sharpshooter for use in a paratransgenic

control strategy. Current Microbiology. 2004;48(5):327-31. doi: 10.1007/s00284-003-

4178-2.

53.   Hurwitz I, Hillesland H, Fieck A, Das P, Durvasula R. The paratransgenic sand

fly: A platform for control of *Leishmania* transmission. Parasites & Vectors.

2011;4(1):82. doi: 10.1186/1756-3305-4-82.

54.   Durvasula RV, Gumbs A, Panackal A, Kruglov O, Taneja J, Kang AS, et al.

Expression of a functional antibody fragment in the gut of *Rhodnius prolixus* via

transgenic bacterial symbiont *Rhodococcus rhodnii*. Medical and Veterinary

Entomology. 1999;13(2):115-9.

55.   Wang S, Ghosh AK, Bongio N, Stebbings KA, Lampe DJ, Jacobs-Lorena M.

Fighting malaria with engineered symbiotic bacteria from vector mosquitoes.

Proceedings of the National Academy of Sciences of the United States of America.

2012;109(31):12734-9. doi: 10.1073/pnas.1204158109.

56.   Wu SC-Y, Maragathavally KJ, Coates CJ, Kaminski JM. Steps toward targeted

insertional mutagenesis with class II transposable elements. Methods in molecular

biology 2008;435:139-51. doi: 10.1007/978-1-59745-232-8_10.

57.   Tikhe CV, Martin TM, Howells A, Delatte J, Husseneder C. Assessment of

genetically engineered *Trabulsiella odontotermitis* as a 'Trojan Horse' for

paratransgenesis in termites. BMC Microbiology. 2016;16(1):355. doi: 10.1186/s12866-

016-0822-4.

58.   Pittman GW, Brumbley SM, Allsopp PG, O apos Neill SL. Assessment of gut

bacteria for a paratransgenic approach to control *Dermolepida albohirtum* larvae.

Applied and Environmental Microbiology. 2008;74(13):4036-43. doi:

10.1128/AEM.02609-07.

59.   Pontes MH, Dale C. Lambda red-mediated genetic modification of the insect

endosymbiont *Sodalis glossinidius*. Applied and Environmental Microbiology.

2011;77(5):1918-20. doi: 10.1128/AEM.02166-10.

60.   Wang S, Dos-Santos ALA, Huang W, Liu KC, Oshaghi MA, Wei G, et al. Driving

mosquito refractoriness to *Plasmodium falciparum* with engineered symbiotic bacteria.

Science (New York, NY). 2017;357(6358):1399-402. doi: 10.1126/science.aan5478.

61.   Dotson EM, Plikaytis B, Shinnick TM, Durvasula RV, Beard CB. Transformation

of *Rhodococcus rhodni*i, a symbiont of the Chagas disease vector *Rhodnius prolixus*,

with integrative elements of the L1 mycobacteriophage. Infection, genetics and

evolution. 2003;3(2):103-9.

62.   Bextine B, Lampe D, Lauzon C, Jackson B, Miller TA. Establishment of a

genetically marked insect-derived symbiont in multiple host plants. Curr Microbiol.

2005;50(1):1-7. doi: 10.1007/s00284-004-4390-8.

63.   Wu P, Sun P, Nie K, Zhu Y, Shi M, Xiao C, et al. A gut commensal bacterium

promotes mosquito permissiveness to arboviruses. Cell Host & Microbe.

2019;25(1):101-12.e5. doi: 10.1016/j.chom.2018.11.004.

64.   Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut

microbiota for development. Molecular Ecology. 2014;23(11):2727-39. doi:

10.1111/mec.12771.

65.   Lindh JM, Borg-Karlson AK, Faye I. Transstadial and horizontal transfer of

bacteria within a colony of *Anopheles gambiae* (Diptera: Culicidae) and oviposition

response to bacteria-containing water. Acta Tropica. 2008;107(3):242-50. doi:

10.1016/j.actatropica.2008.06.008.

66.   Coon KL, Brown MR, Strand MR. Gut bacteria differentially affect egg production

in the anautogenous mosquito *Aedes aegypti* and facultatively autogenous mosquito

Aedes atropalpus (Diptera: Culicidae). Parasites & Vectors. 2016;9(1):375. doi:

10.1186/s13071-016-1660-9.

67.   Ribet D, Cossart P. How bacterial pathogens colonize their hosts and invade

deeper tissues. Microbes and infection. 2015;17(3):173-83. doi:

10.1016/j.micinf.2015.01.004.

68.   Ramirez JL, Short SM, Bahia AC, Saraiva RG, Dong Y, Kang S, et al.

Chromobacterium Csp_P reduces malaria and dengue infection in vector mosquitoes

and has entomopathogenic and in vitro anti-pathogen activities. PLoS Pathogens.

2014;10(10):e1004398. doi: 10.1371/journal.ppat.1004398.

69.   Valzania L, Martinson VG, Harrison RE, Boyd BM, Coon KL, Brown MR, et al.

Both living bacteria and eukaryotes in the mosquito gut promote growth of larvae. PLoS

Neglected Tropical Diseases. 2018;12(7):e0006638. doi: 10.1371/journal.pntd.0006638.

70.   Correa MA, Matusovsky B, Brackney DE, Steven B. Generation of axenic *Aedes*

aegypti demonstrate live bacteria are not required for mosquito development. Nature

Communications. 2018;9(1):R37. doi: 10.1038/s41467-018-07014-2.

71.   Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, et al. Bacteria of

the genus *Asaia* stably associate with *Anopheles stephensi,* an Asian malarial mosquito

vector. Proceedings of the National Academy of Sciences. 2007;104(21):9047-51. doi:

10.1073/pnas.0610451104.

72.   Chaverra-Rodriguez D, Macias VM, Hughes GL, Pujhari S, Suzuki Y, Peterson

DR, et al. Targeted delivery of CRISPR-Cas9 ribonucleoprotein into arthropod ovaries

for heritable germline gene editing. Nature Communications. 2018;9(1):245. doi:

10.1038/s41467-018-05425-9.

73.   Li M, Bui M, Yang T, Bowman CS, White BJ, Akbari OS. Germline Cas9

expression yields highly efficient genome engineering in a major worldwide disease

vector,*Aedes aegypti*. Proceedings of the National Academy of Sciences of the United

States of America. 2017;114(49):E10540-E9. doi: 10.1073/pnas.1711538114.

74.   Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E. CHOPCHOP: a

CRISPR/Cas9 and TALEN web tool for genome editing. Nucleic Acids Research.

2014;42(W1):W401-W7. doi: 10.1093/nar/gku410.

75.   Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E. CHOPCHOP v2: a

web tool for the next generation of CRISPR genome engineering. Nucleic Acids

Research. 2016;44(W1):W272-W6. doi: 10.1093/nar/gkw398.

76.   Trehan A, Kiełbus M, Czapinski J, Stepulak A, Huhtaniemi I, Rivero-Müller A.

REPLACR-mutagenesis, a one- step method for site-directed mutagenesis by

recombineering. Scientific Reports. 2015:1-9. doi: 10.1038/srep19121.

77.   Kozlova EV, Khajanchi BK, Popov VL, Wen J, Chopra AK. Impact of QseBC

system in c-di-GMP-dependent quorum sensing regulatory network in a clinical isolate

SSU of *Aeromonas hydrophila*. Microbial pathogenesis. 2012;53(3-4):115-24. doi:

10.1016/j.micpath.2012.05.008.

78.   O' Toole GA, Kolter R. Flagellar and twitching motility are necessary for

Pseudomonas aeruginosa biofilm development. Molecular Microbiology.

1998;30(2):295-304. doi: 10.1046/j.1365-2958.1998.01062.x.

S1 Table. gRNA sequences tested in this study. Capital letters indicate the protospacer

sequence.

gRNA	Sequence (5'-3')
Ec.gRNA1	CTACCCGTCTGGAATACCAGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtcggtgc
Ec.gRNA2	GGTTCTGACGCTTACAACCAgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtcggtgc
Ent. gRNA1	CTACCCGTCTGGAATACCAGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtcggtgc
Ent. gRNA2	GCTGCAGAAGCAGCTCTCGGgttttagagctagaaatagcaagttaaaataaggctagtccgttatcaacttgaaaaagtggcaccgagtcggtgc

792
793

S2 Table. Plasmids and bacterial strains used in this study.

Plasmid and strains	Relevent charecteristics	Referece
E. coli BL21 (DE3)	*fhuA2 [lon] ompT gal (λ DE3) [dcm] ∆hsdS, λ DE3 = λ sBamHIo ∆EcoRI-B int::(lacI::PlacUV5::T7 gene1) i21 ∆nin5*	NEB
Ec. Δ*ompA*	*E. coli* BL21 (DE3) Δ*ompA*	This study
Enterobacter symbiont		
Δ*ompA*	Enterobacter Δ*ompA*	This study
Ent::Δ*ompA*	Enterobacter ΔompA pRAM_mCherry_*ompA*; Kan^r	This study
ompA::mChery	Enterobacter ΔompA::AmTr::mCherry	This study
ompA::gentamicin	Enterobacter ΔompA::AmTr::gentamicin; Gent^r	This study
pKDsgRNA-xxx	gRNA amplification plasmid with scaffold sequence under P$_{TET}$ promoter & λ-Red system under control of the arabinose inducible promoter P$_{araB}$;Spec^r	36
pcas9cr4	expressed under control of the PTET promoter and tetR constitutively express	36
pRAM_mCherry	mCherry expressing plasmid under the *ompA* promoter	This study

796
797

810 **S3 Table**. Primers used in this study.

Primer	Sequence (5'-3')	Description
InF_BL21ompAgRNA1_F	CTA CCC GTC TGG AAT ACC AGG TTT TAG AGC TAG AAA TAG C	Amplification of *E. coli ompA* gRNA1 and cloning
InF_BL21ompAgRNA1_R	CTG GTA TTC CAG ACG GGT AGG TGC TCA GTA TCT CTA TCA C	Amplification of *E. coli ompA* gRNA1 and cloning
InF_BL21ompAgRNA2_F	GGT TCT GAC GCT TAC AAC CAG TTT TAG AGC TAG AAA TAG C	Amplification of *E. coli ompA* gRNA2 and cloning
InF_BL21ompAgRNA2_R	TGG TTG TAA GCG TCA GAA CCG TGC TCA GTA TCT CTA TCA C	Amplification of *E. coli ompA* gRNA2 and cloning
BL21ompA1_LAFw	CTT TTT TTT CAT ATG CCT GAC GG	Amplification of left arm of donor dsDNA *E.coli*
BL21ompA1_LARv	AAG CCT GCG GCT GAG TTA GCG ATA GCT GTC TTT TTC AT	Amplification of left arm of donor dsDNA *E. coli*
BL21ompA1_RAFw	TAA CTC AGC CGC AGG CTT AAG TTC TCG TCT GGT AGA A	Amplification of right arm ofdonor dsDNA *E. coli*
BL21ompA1_RARv	GTC GCG GTA AAA CGC TTT CTG	Amplification of right arm ofdonor dsDNA *E. coli*
OmpA1_COLPCR_F	CTACCCGTCTGGAATACCAG	*ompA* Colony PCR primer
OmpA2_COLPCR_F	GGTTCTGACGCTTACAACCA	*ompA* Colony PCR primer
pRAM_CompV_F	TAACTCAGCCTGCGGCATAAGGCTAATAAGCAACCGTTCT	Amplification of complement vector
pRAM_CompV_R	GCGATAGCTGTCTTTTTCATATGTAAAACCTTAATCAAAA	Amplification of complement vector
ompA_F	TTTTGATTAAGGTTTTACATATGAAAAAGACAGCTATCGC	Amplification of *Enterobacter ompA* gene
ompA_R	AGAACGGTTGCTTATTAGCCTTATGCCGCAGGCTGAGTTA	Amplification of *Enterobacter ompA* gene
ompAprom_COLPCR_F	CTATGTTATTAATTATAATAG	colony PCR primer
EntompALA_F	ATGAAAAAGACAGCTATCGC	Amplification of left arm of dsDNA for ompA deletion
EntompALA_R	CGTTTTTCAGACAGTTTCCAAAACCAACGTACGGGTTAAC	Amplification of left arm of dsDNA for ompA deletion
EntompARA_F	GAAACTGTCTGAAAAACGTGCACAGAGCGTTGTTGAT	Amplification of right arm of dsDNA for ompA deletion
EntompARA_R	TTATGCCGCAGGCTGAGTTAC	Amplification of right arm of dsDNA for ompA deletion
COLPCR_R	GACGCTTCTCAAAACTACATT	colony pcr primer
Ent. ompA _LAF	ATGGGTTACGACTGGCTTGGC	Amplification of left arm of donor dsDNA for insertion
Ent. ompA _LAR	GTTCTAGGAGGAGCCCGATGTCACGGGTAACTGCCCAC	Amplification of left arm of donor dsDNA for insertion
AmTr prom_F	GTTACCCGTGACATCGGGCTCCTCCTAGAACGATCGCCGC	Amplification of AmTr promoter for dsDNA for insertion
AmTr prom.gent_R	TGCTGCTGCGTAACATTATAATATCCCTTATGTTACTC	Amplification of AmTr promoter for dsDNA for gentamicin insertion
Gent. F	CATAAGGGATATTATAATGTTACGCAGCAGCAACGATG	Amplification of Gentamicin for dsDNA
Gent. R	CGATGTTGTTAACCCATTAGGTGGCGGTACTTGGGTCG	Amplification of Gentamicin for dsDNA
Ent. ompA _RAF1-gent	CAAGTACCGCCACCTAATGGGTTAACAACATCGGTGATGC	Amplification of right arm of donor dsDNA for gentamicin insertion
Ent.ompA_RAR	TCAGCTGGGTGTACAGCTGATC	Amplification of right arm of donor dsDNA for insertion
AmTr mChery_R	CTCCCTTGCTGACCATTATAATATCCCTTATGTTACTC	Amplification of AmTr promoter for dsDNA for mCherry insertion
mCherry. F	CATAAGGGATATTATAATGGTCAGCAAGGGAGAGGAAG	Amplification of mCherry for dsDNA
mCherry. R	CGATGTTGTTAACCCACTATTTGTATAATTCGTCCATTC	Amplification of mCherry for dsDNA
pkdsgRNA_COLPCR_F	GTC TGA CGC TCA GTG GAA CG	Colony PCR primer for pKDsgRNA plamsid
pkdsgRNA_COLPCR_R	CTG GCA GTT CCC TAC TCT CG	Colony PCR primer for pKDsgRNA plamsid
InF_EntinvgRNA1_F	AAGCGACGCATGGCATCCGG gttttagagctagaaatagc	Amplification of *Enterobacter ompA* gRNA1 and cloning
InF_EntinvgRNA1_R	CCGGATGCCATGCGTCGCTT gtgctcagtatctctatcac	Amplification of *Enterobacter ompA* gRNA1 and cloning
InF_EntinvgRNA2_F	GCTGCAGAAGCAGCTCTCGG gttttagagctagaaatagc	Amplification of *Enterobacter ompA* gRNA2 and cloning
InF_EntinvgRNA2_R	CCGAGAGCTGCTTCTGCAGC gtgctcagtatctctatcac	Amplification of *Enterobacter ompA* gRNA2 and cloning

811
812

 S1. Appendix. Multiple sequence alignment of WT and mutant *ompA* sequences of

814 *Enterobacter* and *E. coli.*

```
815   Multiple sequence alignment E. coli WT ompA and mutant ompA
816
817
818   WT        ATGAAAAAGACAGCTATCGCGATTGCAGTGGCACTGGCTGGTTTCGCTACCGTAGCGCAG
819   •ompA     ATGAAAAAGACAGCTATCGC----------------------------------------
820             *******************
821
822   WT        GCCGCTCCGAAAGATAACACCTGGTACACTGGTGCTAAACTGGGCTGGTCCCAGTACCAT
823   •ompA     ------------------------------------------------------------
824
825
826   WT        GACACTGGTTTCATCAACAACAATGGCCCGACCCATGAAAACCAACTGGGCGCTGGTGCT
827   •ompA     ------------------------------------------------------------
828
829
830   WT        TTTGGTGGTTACCAGGTTAACCCGTATGTTGGCTTTGAAATGGGTTACGACTGGTTAGGT
831   •ompA     ------------------------------------------------------------
832
833
834   WT        CGTATGCCGTACAAAGGCAGCGTTGAAAACGGTGCATACAAAGCTCAGGGCGTTCAACTG
835   •ompA     ------------------------------------------------------------
836
837
838   WT        ACCGCTAAACTGGGTTACCCAATCACTGACGACCTGGACATCTACACTCGTCTGGGTGGT
839   •ompA     ------------------------------------------------------------
840
841
842   WT        ATGGTATGGCGTGCAGACACTAAATCCAACGTTTATGGTAAAAACCACGACACCGGCGTT
843   •ompA     ------------------------------------------------------------
844
845
846   WT        TCTCCGGTCTTCGCTGGCGGTGTTGAGTACGCGATCACTCCTGAAATCGCTACCCGTCTG
847   •ompA     ------------------------------------------------------------
848
849
850   WT        GAATACCAGTGGACCAACAACATCGGTGACGCACACACCATCGGCACTCGTCCGGACAAC
851   •ompA     ------------------------------------------------------------
852
853
854   WT        GGCATGCTGAGCCTGGGTGTTTCCTACCGTTTCGGTCAGGGCGAAGCAGCTCCAGTAGTT
855   •ompA     ------------------------------------------------------------
856
857
858   WT        GCTCCGGCTCCAGCTCCGGCACCGGAAGTACAGACCAAGCACTTCACTCTGAAGTCTGAC
859   •ompA     ------------------------------------------------------------
860
861
862   WT        GTTCTGTTCAACTTCAACAAAGCAACCCTGAAACCGGAAGGTCAGGCTGCTCTGGATCAG
863   •ompA     ------------------------------------------------------------
864
865
866   WT        CTGTACAGCCAGCTGAGCAACCTGGATCCGAAAGACGGTTCCGTAGTTGTTCTGGGTTAC
867   •ompA     ------------------------------------------------------------
868
869
870   WT        ACCGACCGCATCGGTTCTGACGCTTACAACCAGGGTCTGTCCGAGCGCCGTGCTCAGTCT
871   •ompA     ------------------------------------------------------------
872
873
874   WT        GTTGTTGATTACCTGATCTCCAAAGGTATCCCGGCAGACAAGATCTCCGCACGTGGTATG
875   •ompA     ------------------------------------------------------------
876
877
878   WT        GGCGAATCCAACCCGGGTTACTGGCAACACCTGTGACAACGTGAAACAGCGTGCTGCACTG
879   •ompA     ------------------------------------------------------------
```

```
880
881
882   WT        ATCGACTGCCTGGCTCCGGATCGTCGCGTAGAGATCGAAGTTAAAGGTATCAAAGACGTT
883   •ompA     ------------------------------------------------------------
884
885
886   WT        GTAACTCAGCCGCAGGCTTAA
887   •ompA     -TAACTCAGCCGCAGGCTTAA
888             *******************
894   Alignment of Enterobacter WT ompA and mutant ompA sequence
895
896
897   WT        atgaaaaagacagctatcgcgattgcagtggcactggctggcttcgctaccgtagcgcag 60
898   •ompA     atgaaaaagacagctatcgcgattgcagtggcactggctggcttcgctaccgtagcgcag 60
899             ************************************************************
900
901   WT        gccgcaccgaaagataacacctggtatgcaggtggtaaactgggctggtctcagttccac 120
902   •ompA     gccgcaccgaaagataacacctggtatgcaggtggtaaactgggctggtctcagttccac 120
903             ************************************************************
904
905   WT        gataccggctggtataacaaggacctgcaaaacaacggtaacacccacgaaagccagctg 180
906   •ompA     gataccggctggtataacaaggacctgcaaaacaacggtaacacccacgaaagccagctg 180
907             ************************************************************
908
909   WT        ggcgctggtgccttcggtggttatcaggttaacccgtacgttggttttgaaatgggttac 240
910   •ompA     ggcgctggtgccttcggtggttatcaggttaacccgtacgttggttttgaaa-------- 232
911             ****************************************************
912
913   WT        gactggcttggccgtatgccttacaaaggcagcgtaactagcggtgccttcaaagctcag 300
914   •ompA     ------------------------------------------------------------ 232
915
916
917   WT        ggcgtacagctgaccactaaactgggttacccaatcactgacgacctggacatctacacc 360
918   •ompA     ------------------------------------------------------------ 232
919
920
921   WT        cgtctgggcggcatggtatggcgtgcagactctacaaacaccatcgctggtaaaaaccac 420
922   •ompA     ------------------------------------------------------------ 232
923
924
925   WT        gacaccggtgtttcccagtattcgctggtggtgttgagtgggcagttacccgtgacatc 480
926   •ompA     ------------------------------------------------------------ 232
927
928
929   WT        gctacccgtctggaataccagtgggttaacaacatcggtgatgcacagaccgttggcgtt 540
930   •ompA     ------------------------------------------------------------ 232
931
932
933   WT        cgtcctgacaacggcatgctgagcgtaggtgtttcctaccgctttggccagcaggaagaa 600
934   •ompA     ------------------------------------------------------------ 232
935
936
937   WT        gctgcaccagttgtagctccggctccagctccagctcctgaagtacagaccaaacacttc 660
938   •ompA     ------------------------------------------------------------ 232
939
940
941   WT        actctgaagtctgacgtcctgttcaacttcaacaaagcaaccctgaaaccagaaggtcag 720
942   •ompA     ------------------------------------------------------------ 232
```

```
943
944
945   WT      caggcactggatcagctgtacacccagctgagcaacctggatcctaaagatggttccgtt 780
946   •ompA   ------------------------------------------------------------ 232
947
948
949   WT      gttgttctgggctacaccgaccgtatcggttccgagcagtacaacctgaaactgtctgaa 840
950   •ompA   ---------------------------------------------actgtctgaa 242
951                                                         *********
952
953   WT      aaacgtgcacagagcgttgttgattacctgatctctaaaggtatcccagcgaacaaaatc 900
954   •ompA   aaacgtgcacagagcgttgttgattacctgatctctaaaggtatcccagcgaacaaaatc 302
955           ************************************************************
956
957   WT      tctccacgtggcatgggcaaagctaacccagttaccggcaacacctgtgacaaagtagcg 960
958   •ompA   tctccacgtggcatgggcaaagctaacccagttaccggcaacacctgtgacaaagtagcg 362
959           ************************************************************
960
961   WT      cctaaagctaaactgatcgactgcctggctccagatcgtcgcgttgagatcgaagttaaa 1020
962   •ompA   cctaaagctaaactgatcgactgcctggctccagatcgtcgcgttgagatcgaagttaaa 422
963           ************************************************************
964
965   WT      ggtatcaaagaagttgtaactcagcctgcggcataa 1056
966   •ompA   ggtatcaaagaagttgtaactcagcctgcggcataa 458
967           ************************************

968
```

A

Chr — *ompA* — *AmTr prom* — *mCherry* — *ompA*

CGTGACATCG GGCTCCTCCT

TGTTAACCCA CTATTTGTAT

B

P0 P2 P4 P6 P8 P10 *NTC* M
a b a b a b a b a b a b

2 kb →
1kb →

a- WT, b-*ompA*::mCherry

C

WT

ΔompA::mCherry

D

Chr — *ompA* — *AmTr prom* — *gentamicin* — *ompA*

CGTGACATCG GGCTCCTCCT

CGCCACCTAA TGGGTTAACA

E

P0 P2 P4 P6 P8 P10 NTC M
a b a b c a b c a b c a b c a b c

1.8 kb →
1kb →

a- WT, b-*ompA*::gentamicin(ab-), c-*ompA*::gentamicin(ab+)

F

WT

ΔompA::gentamicin

G

H

$P < 0.0001$ $P = 0.9705$ $P < 0.05$ $P < 0.001$

CFU/adult

Time in days

WT   ●   *ompA*::gentamicin   ■