Whole-genome sequencing of three native cattle breeds originating from the northernmost cattle farming regions

Melak Weldenegodguad[1,2], Ruslan Popov[3], Kisun Pokharel[1], Innokentyi Ammosov[4], Ming

Yao[5], Zoya Ivanova[6] and Juha Kantanen[1*]

[1]Production Systems Department, Natural Resources Institute Finland (Luke), Jokioinen, Finland

[2]Department of Environmental and Biological Sciences, University of Eastern Finland, Kuopio,
Finland

[3]Yakutian Research Institute of Agriculture (FGBNU Yakutskij NIISH), Yakutsk, Russia

[4]Board of Agricultural Office of Eveno-Bytantaj Region, Yakutia, Russia

[5]BGI-Shenzhen, Shenzhen, China

[6] Autumnwood Ct SE 22344, Yelm WA 98597, USA

* Correspondence:

Juha Kantanen

juha.kantanen@luke.fi

Keywords: adaptation, demographic history, Finncattle, indels, selective sweeps, SNPs, WGS,

Yakutian cattle

Abstract

Northern Fennoscandia and the Sakha Republic in the Russian Federation represent the northernmost

regions on Earth where cattle farming has been traditionally practiced. In this study, we performed

whole-genome resequencing to genetically characterize three rare native breeds Eastern Finncattle,

Western Finncattle and Yakutian cattle adapted to these northern Eurasian regions. We examined the

demographic history, genetic diversity and unfolded loci under natural or artificial selection. On

average, we achieved 13.01-fold genome coverage after mapping the sequencing reads on the bovine

reference genome (UMD 3.1) and detected a total of 17.45 million single nucleotide polymorphisms

(SNPs) and 1.95 million insertions-deletions (indels). We observed that the ancestral species (*Bos*

primigenius) of Eurasian taurine cattle experienced two notable prehistorical declines in effective

population size associated with dramatic climate changes. The modern Yakutian cattle exhibited a

higher level of within-population variation in terms of number of SNPs and nucleotide diversity than

the contemporary European taurine breeds. This result is in contrast to the results of marker-based

cattle breed diversity studies, indicating assortment bias in previous analyses. Our results suggest that

the effective population size of the ancestral Asiatic taurine cattle may have been higher than that of

the European cattle. Alternatively, our findings could indicate the hybrid origins of the Yakutian

cattle ancestries and possibly the lack of intensive artificial selection. We identified a number of

genomic regions under selection that may have contributed to the adaptation to the northern and

subarctic environments, including genes involved in disease resistance, sensory perception, cold

adaptation and growth. By characterizing the native breeds, we were able to obtain new information

on cattle genomes and on the value of the adapted breeds for the conservation of cattle genetic

resources.

Introduction

During their 8,000-10,000 years of domestication, taurine cattle (*Bos taurus*) have adapted to a wide

variety of biogeographic zones and sociocultural environments as a result of natural and human-

derived selection (Felius, 1995). Fennoscandia along with northwestern Russia and the region of

Sakha (Yakutia) in eastern Siberia, are the northernmost territories where cattle farming has had a

relatively long tradition as the livelihood of local people (Kopoteva and Partanen, 2009; Bläuer and

Kantanen, 2013; Cramp et al., 2014; Egorov et al., 2015). In prehistoric and historic times, animal

husbandry faced several challenges in these northern climatic conditions, such as short summers and

limited vegetation resources for feeding during the long winters, and this practice required well-

adapted animals that were suited to the available environmental resources and socioeconomic and

cultural conditions (Kantanen et al., 2009a; Bläuer and Kantanen, 2013; Egorov et al., 2015).

Cattle breeds such as Eastern Finncattle, Icelandic cattle, Swedish Mountain cattle, Yakutian cattle

and other northern native cattle breeds are assumed to have their origins in the near-eastern

domesticated taurine cattle that once spread to these northern regions (Kantanen et al., 2000, 2009a;

Li et al., 2007). Herd books, pedigree registers and breeding associations were established in the late

19th and early 20th centuries. Early native breeds had a pivotal socioeconomic role in dairy and beef

production in the northern Eurasian regions but have been almost exclusively replaced by

commercial international cattle populations bred for high-input, high-output farming systems.

Exceptions to this trend are Yakutian cattle in Siberia and Icelandic cattle, which continue to have

high regional importance in food production (Kantanen et al., 2000, 2009a). The conservation of the

genetic resources of native, typically low-profit breeds is often motivated by the fact that these breeds

may possess valuable genetic variations for future animal breeding and to address the challenges that

animal production will face during adaptation to future conditions, brought about by factors such as

climate change (Odegård et al., 2009; Boettcher et al., 2010; Kantanen et al., 2015). In addition,

breeds such as Yakutian cattle exhibit adaptation in demanding environments and may be extremely

useful for enabling animal production in marginal regions (Kantanen et al., 2015).

Previous studies on the characterization of cattle genetic resources in northern Eurasian breeds have

used various methods to study within-breed genetic diversity, population structure, demographic

factors and interbreed relationships, e.g., autosomal and Y-chromosomal microsatellites,

mitochondrial D-loop and whole-genome SNP-marker scans (Li et al., 2007; Kantanen et al., 2009b;

Iso-Touru et al., 2016). These studies have indicated, for example, the genetic distinctiveness of the

native northern European cattle breeds (e.g., the Finnish native breeds and Yakutian cattle) from

modern commercial dairy breeds (such as the Finnish Ayrshire and Holstein breeds). In addition, a

whole-genome SNP genotyping analysis detected genomic regions targeted by selection, which, for

example, contain immune-related genes (Iso-Touru et al., 2016). Whole-genome sequencing (WGS)-

based approaches provide additional possibilities for investigation of the genetic diversity of

livestock breeds adapted to various biogeographic regions and production environments. Moreover,

recent advancements in bioinformatics and statistical tools have enhanced our understanding of the

demographic evolution of domestic animal species, the possible role of genomic structural variations

in the adaptation of livestock breeds in the course of domestication and selection and the biological

functions of these genomic variations (Gutenkunst et al., 2009; Li and Durbin, 2011; Alachiotis et al.,

2012; Pavlidis et al., 2013; Wang et al., 2014b; Librado et al., 2015).

To expand our knowledge of genomic variations in northern Eurasian taurine cattle, we performed

whole-genome resequencing of five animals from each of three northern native breeds, namely,

Eastern Finncattle, Western Finncattle and Yakutian cattle (Figure 1). We examined the genetic

diversity and population structures of the breeds and identified chromosomal regions and genes under

selection pressure. We also studied the demographic history of the northern Eurasian taurine cattle by

using the whole-genome sequence data.

MATERIALS AND METHODS

Ethics statement

Blood samples of animals for DNA extraction were collected by using a protocol approved by the

Animal Experiment Board of MTT Agrifood Research Finland (currently the Natural Resources

Institute Finland, Luke) and the Board of Agricultural Office of Eveno-Bytantaj Region, Sakkyryr,

Sakha, Russia.

DNA sample preparation and sequencing

DNA extracted from blood samples was available for the two Finnish cattle breeds (Eastern

Finncattle and Western Finncattle) and one Siberian breed (Yakutian cattle) from a previous study

(Li et al., 2007). Five unrelated individuals from each breed (14 females and one Yakutian cattle bull)

were examined. Genomic DNA was extracted using a standard phenol/chloroform-based protocol

(Malke, 1990). For sequencing library preparation following the manufacturer's specifications, the

genomic DNA of each individual was fragmented randomly. After electrophoresis, DNA fragments

of desired length were gel purified. One type of library was constructed for each sample (500 bp

insert size); 15 paired-end DNA libraries were constructed for the 15 samples. Adapter ligation and

DNA cluster preparation were performed, and the DNA was subjected to Illumina HiSeq 2000

sequencing using the 2×100 bp mode at Beijing Genomics Institute (BGI). Finally, paired-end

sequence data were generated. To ensure quality, the raw data was modified by the following 2 steps:

first, the contaminating adapter sequences from the reads were deleted, and then, the reads that

contained more than 50% low-quality bases (quality value≤5) were removed.

Short read alignment and mapping

For short read alignment, the bovine reference genome (UMD 3.1), including regions that were not

assembled into chromosomes (Zimin et al., 2009), were downloaded from the Ensembl database

release 71 (Flicek et al., 2013) and indexed using SAMtools v0.1.19 (Li et al., 2009). Paired-end 100-

bp short reads from each individual sample were mapped against the bovine reference genome

assembly UMD 3.1 using BWA v0.7.5a with the default parameters. After mapping, for downstream

SNP and insertion-deletion (indel) detection, the SAM files that were generated from BWA were

converted to the corresponding binary equivalent BAM files and sorted simultaneously using

SortSam.jar in Picard tools v1.102 (http://picard.sourceforge.net/). We used Picard tools to remove

PCR duplicates from the aligned reads and then used the uniquely mapped reads for variant calling.

SNP and indel detection

We used the Genome Analysis Toolkit (GATK) v2.6-4 according to the GATK best practices

pipeline (McKenna et al., 2010; DePristo et al., 2011; Van der Auwera A. et al., 2013) for

downstream SNP and indel calling. We used RealignerTargetCreater to identify poorly mapped

regions (nearby indels) from the alignments and realigned these regions using IndelRealigner. Next,

the UnifiedGenotyper was used to call SNPs and indels with a Phred scale quality greater than 30.

After SNP calling, we used VariantFiltration to discard sequencing and alignment artifacts from the

SNPs with the parameters "MQ0 $\geq$ 4 && ((MQ0 / (1.0 * DP)) > 0.1)", "SB $\geq$ -1.0, QUAL < 10", and

"QUAL < 30.0 || QD < 5.0 || HRun > 5 || SB > -0.10" and from the indels with the parameters "QD <

2.0 ", "FS > 200.0" and "ReadPosRankSum < -20.0". All the variants that passed the above filtering

criteria were used in the downstream analysis and compared to the cattle dbSNP148 (Van der

Auwera A. et al., 2013) to identify novel variants.

SNP and indel annotation and gene ontology analysis

ANNOVAR (Wang et al., 2010) was used to annotate the functions of the variants (exonic, intronic,

5′ and 3′ UTRs, splicing, intergenic) using Ensembl release 71. SNPs that were identified in the

exonic regions were classified as synonymous or nonsynonymous SNPs. In recent studies, numerous

phenotypes have been associated with the genes containing the highest number of nonsynonymous

SNPs (nsSNPs) (Kawahara-Miki et al., 2011; Li et al., 2014). We performed gene ontology (GO)

analysis for genes containing nsSNPs and indels using the GO Analysis Toolkit and Database for

Agricultural Community (AgriGO) (Du et al., 2010). In this analysis, we selected genes containing

>5 nsSNPs for each breed. The significantly enriched GO terms were assessed by Fisher's exact test

with the Bonferroni correction using default parameters (P-value, 0.05; at least 5 mapping entries).

Out of four indel classes (frameshift, nonframeshift, stopgain and stoploss), we annotated frameshift

indels in exonic regions using default parameters in ANNOVAR. Frameshift indels may change

amino acid sequences and thereby affect protein function.

Identification and annotation of selective sweeps

We investigated the signatures of selection using site frequency spectrum (SFS)-based α statistics in

SweeD (Pavlidis et al., 2013) with default parameters, except setting the grid as the only parameter.

SweeD detects the signature of selection based on the composite likelihood ratio test (CLR) using

SFS-based statistics. SweeD was run separately for each chromosome by setting the grid parameter at

5-kb equidistant positions across the chromosome (size of the chromosome/5 kb). We used BEAGLE

program ver.4 (Browning and Browning, 2007) to impute missing alleles and infer the haplotype

phase for all individual Western Finncattle, Yakutian cattle and Eastern Finncattle simultaneously

(among the Eastern Finncattle, we excluded one inbred animal; see Results). The BEAGLE program

infers the haplotype information of each chromosome, which is required for α statistics. Following

the approaches described in previous studies (Wang et al., 2014b; McManus et al., 2015), we selected

the outliers falling within the top 0.5% of the CLR distribution. The cutoff value for α statistics was

taken as the 99.5 percentile of the empirical distribution of the 5-kb equidistant positions across the

genome for each chromosome. Annotation of the candidate sites that exhibited a signal of selection

was performed using Ensembl BioMart (Kinsella et al., 2011) by considering a 150-kb sliding

window on the outlier sites. Candidate genes exhibiting signatures of selection were subjected to GO

analysis with same parameters applied in the variant annotation using AgriGO.

Population genetics analysis

The average pairwise nucleotide diversity within a population (π) and the proportion of polymorphic

sites (Watterson's θ) were computed using the Bio::PopGen::Statistics package in BioPerl (v1.6.924)

(Stajich et al., 2002). Principal component analysis (PCA) was conducted using smartpca in

EIGENSOFT3.0 software (Patterson et al., 2006) on biallelic autosomal SNPs that were genotyped in

all individuals. Significant eigenvectors were determined using Tracy-Widom statistics with the

twstats program implemented in the same EIGENSOFT package.

Demographic history inference

We used the pairwise sequentially Markovian coalescent (PSMC) model (Li and Durbin, 2011) to

construct the demographic history of the three breeds. For the analysis, one individual per breed with

highest sequence depth was selected to explore changes in local density of heterozygous sites across

the cattle genome. The following default PSMC parameters were set: –N25, –t15, –r5 and –p

'4+25*2+4+6'. To scale the PSMC output to real time, we assumed a neutral mutation rate of $1.1 \times$

10^{-8} per generation and an average generation time of 5 years (Kumar and Subramanian, 2002;

Murray et al., 2010; MacLeod et al., 2013). As the power of the PSMC approach to reconstruct recent

demographic history is not reliable (Li and Durbin, 2011; MacLeod et al., 2013; Zhao et al., 2013),

we reconstructed a more recent demographic history of the Finnish and Yakutian populations using

the diffusion approximation for demographic inference ($\partial a \partial i$) program (dadi-1.6.3) (Gutenkunst et

al., 2009). We used the intergenic sites from the identified SNPs in the 15 individuals to compute the

folded SFS. We merged the results for the Eastern and Western Finncattle breeds, as these breeds

exhibited similar genetic diversity measures (Figure S4). Since we had 10 Finncattle and 5 Yakutian

samples, we downscaled the Finncattle sample size to be equal to that of the Yakutian cattle. We ran

the $\partial a \partial i$ algorithm multiple times to ensure convergence and selected the optimal parameters with the

highest likelihood as the final result. As $\partial a \partial i$ requires ancestral population size (Na), we calculated

Na using the formula $NA = \theta / 4 \mu L$, where θ was the observed number of segregating sites divided by

the sum of the expected SFS using the best-fit parameters of our model, L was the effective sequence

length, and μ was the mutation rate per generation per site. We used a mutation rate of $1.0 \times 10\text{-}8$

mutations per generation assuming that one generation was equal to 5 years (Kumar and

Subramanian, 2002), and the effective sequence length (intergenic regions) was 10,836,904. We

calculated population size and divergence time between the Finnish and Yakutian populations based

on NA. Finally, using the parameters described previously, we generated the demographic model

using $\partial a \partial i$ as shown in Figure S5. The optimal model identified the change from the ancestral

population size (NA) to the effective population size (nua) from the time Ta to the time Td. Ta was

the time period when the change in NA started and Td was the time when the divergence between the

Finnish and Yakutian cattle occurred. nu1F and nu2Y were the effective population sizes during the

split. To calculate the statistical confidence in the estimated parameter values, we estimated the

parameter uncertainties using the Hessian method (a.k.a. the Fisher information matrix).

RESULTS

Sequence data

A total of 521 gigabases (Gb) of paired-end DNA sequence data was obtained after removing adapter

sequences and low-quality reads (Table 1, Table S1). On average, each sample had 347.4 million (M)

reads, 98.45% of which were successfully mapped to the bovine reference genome UMD3.1 (Table

1, Table S1), representing 12.38-fold coverage.

Identification and annotation of variants

A total of 17.45 M SNPs were detected in the mapped reads across all 15 samples, with Yakutian

cattle exhibiting the highest number of SNPs (Table 2, Figure 2a, Table S2). The average number of

SNPs detected per individual within the breeds was 5.73 M, 6.03 M and 7.12 M in Eastern Finncattle,

Western Finncattle and Yakutian cattle, respectively (Table S2). A total of 6.3 M (36.1%) SNPs were

shared by the three breeds, and as expected, the Finnish breeds shared the highest number (n=8.06 M,

46.2%) of SNPs (Figure 1a). Moreover, we found that 1.85 M SNPs (16.83%) in Eastern Finncattle,

1.60 M (15.15%) in Western Finncattle and 3.96 M (32.33%) in Yakutian cattle were private SNPs in

our data (Figure 2a). The transition-to-transversion (TS/TV) ratios were 2.20 and 2.23 in the

Finncattle and Yakutian cattle, respectively (Table S2). The observed Ts/Tv ratios were consistent

with those observed in previous studies in mammalian systems (Lachance et al., 2012; Choi et al.,

2013, 2014), indicating the quality of our SNP data.

Of the SNPs identified in our analysis, 1.28 M (6.9%) SNPs were found to be novel when compared

to NCBI dbSNP bovine build 148. At the breed level, 3.1%, 2.8% and 5.3% of the total SNPs in the

Eastern Finncattle, Western Finncattle and Yakutian cattle, respectively, were novel. Furthermore,

out of the novel SNPs identified for each breed, 278,399 (82.57%), 235,741(80.72%) and 618,717

(94.85%) were breed-specific SNPs in Eastern Finncattle, Western Finncattle and Yakutian cattle

(Figure 2b), respectively. A summary of the homozygous and heterozygous SNPs is given in Tables

S2 and S3. One Eastern Finncattle cow (sample_3 in Table S3) exhibited exceptionally low diversity,

with only 1.66 M (32.58%) heterozygous and 3.44 M (67.42%) homozygous SNPs. This animal

originated from an isolated, inbred herd and represented one relict Eastern Finncattle line (herd) that

passed through the breed's demographic bottleneck (Kantanen et al., 2000). After excluding this

sample, the average number of SNPs detected per Eastern Finncattle individual was 5.88 M, and the

Eastern Finncattle animals exhibited 2.63 M (44.83%) homozygous and 3.24 M (55.17%)

heterozygous SNPs, with a ratio of 1:1.23 (homozygous:heterozygous). Apparently, the number of

homozygous SNPs in the Eastern Finncattle was higher than that in the other two breeds.

In total, we detected 2.12 M indels, 79.8% of which were found in the dbSNP build 148, with 20.2%

being novel (Figure 2C, Table S2). At the breed level, 13.0%, 11.7% and 16.1% of the total indels in

the Eastern Finncattle, Western Finncattle and Yakutian cattle, respectively, were novel.

In our data, on average, 0.65% of the SNPs were detected in exonic regions, 25.1% in intronic

regions, 72.6% in intergenic regions, and 1.65% in UTRs and in regions upstream and downstream of

genes (Table 2 and Table S4). In general, all the three breeds exhibited similar distributions of SNPs

in various functional categories. A total of 76,810, 71,256 and 84,927 exonic SNPs were identified in

the Eastern Finncattle, Western Finncattle and Yakutian cattle, respectively. Of the exonic SNPs in

the Eastern Finncattle, Western Finncattle and Yakutian cattle, 31,299, 29,035 and 33,111,

respectively, were nonsynonymous SNPs (nsSNPs) (Table 2) and were found in 10,309, 9,864 and

10,429 genes, respectively.

The functional categories of the indel mutations are presented in Table 2 and Table S4. In total,

1,045, 927 and 1,148 of the indels were frameshift indels that were associated with 808, 770 and 895

genes in Eastern Finncattle, Western Finncattle and Yakutian cattle, respectively (Supplementary

Data 1, 2 and 3).

GO analysis of the SNPs and indels

GO enrichment analysis of 1,331, 1,170 and 1,442 genes containing >5 nsSNPs (Supplementary Data

4, 5 and 6), identified 111, 113 and 95 significantly enriched GO terms in Eastern Finncattle,

Western Finncattle and Yakutian cattle, respectively (Supplementary Data 7, 8 and 9). A total of 38,

43 and 38 GO terms were associated with biological processes in Eastern Finncattle, Western

Finncattle and Yakutian cattle, respectively (Supplementary Data 7, 8, 9).

A detailed comparison of the biological processes associated with genes with >5 nsSNPs with the

bovine Ensembl gene set (n=25,160) is shown in Figure S1. The GO enrichment analysis revealed

that a majority of the significantly enriched GO terms were shared by the three cattle breeds.

"Response to stimulus, GO:00050896" was associated with approximately 50% of the genes in

Eastern Finncattle (n=611), Western Finncattle (n =544) and Yakutian cattle (n=629) (see Figure S1).

In addition, this analysis showed that in each breed, a large number of genes were associated with

immune functions, such as "Immune response, GO:0006955", "Defense response, GO:0006952",

"Antigen processing and presentation, GO:0019882", and "Immune system process, GO:0002376".

Among the three breeds, the Yakutian cattle had more enriched genes associated with immune

functions than the two Finncattle breeds. On the other hand, in the Finncattle breeds, a large number

of genes were associated with sensory perception functions, such as "Sensory perception,

GO:0007600", "Sensory perception of smell, GO:0007608" and "Detection of chemical stimulus

involved in sensory perception, GO:0050907". In Yakutian cattle, none of the GO terms associated

with sensory perception were enriched. However, 55 genes associated with "Developmental growth,

GO: 0048589" were enriched in only Yakutian cattle.

We further identified the top genes, namely, *TTN, PKHD1, GPR98* and *ASPM*, that had at least 40

nsSNPs in all the breeds. These genes have large sizes; *TTN* is 274 kb in size, *PKHD1* if 455 kb,

GPR98 is 188 kb and ASPM is 64 kb. Among the genes with nsSNPs, *TTN* contained the highest

number of nsSNPs: 68, 63 and 87 nsSNPs in Eastern Finncattle, Western Finncattle and Yakutian

cattle, respectively. The *TTN* gene is present on chromosome 2 and is associated with meat quality

(Sasaki et al., 2006; Watanabe et al., 2011).

A total of 709, 675 and 772 genes associated with frameshift indels in these breeds were linked to at

least one GO term (Figure S2, Supplementary Data 10, 11 and 12). The results indicated that a

majority of the significantly enriched GO terms were shared by the breeds. The GO terms "Defense

response, GO:0006952" and "Female pregnancy, GO:0007565" were enriched exclusively in

Yakutian cattle. In total, 96 genes were enriched in "Defense response, GO:0006952".

Selection signatures

We identified 2,528 sites exhibiting signatures of selection in each breed, of which 58%, 61% and

53% mapped to gene regions in Eastern Finncattle, Western Finncattle and Yakutian cattle,

respectively (Figure S3). Information regarding the SNPs found in selective sweep regions in each

breed is shown in Table S5.

Chromosome 1 exhibited the highest (n=159) number of selection signals and chromosome 25 the

lowest (n=43). Considering a 150-kb window centered on the candidate site, Western Finncattle

exhibited the highest number (n=371) of candidate genes with selection signatures, followed by

Eastern Finncattle (n=331), while Yakutian cattle exhibited the lowest number (n=249)

(Supplementary Data 13, 14 and 15). Apparently, 36 (Eastern Finncattle), 35 (Western Finncattle)

and 20 (Yakutian cattle) candidate gene IDs lacked gene descriptions (Supplementary Data 16, 17

and 18). Seven genes with greater than 5 nsSNPs in Eastern Finncattle (*CCSAP, CEP72, GBP5,*

LOC100297846, GBP2, LOC613867 and ENSBTAG00000045571), Western Finncattle (*CDH23,*

PCDHB4, PCDHB6, PCDHB7, SIRPB1, LOC783488 and ENSBTAG00000012326) and Yakutian

cattle (*FER1L6, GBP5*, ENSBTAG00000015464, ENSBTAG00000025621, *GBP2*,

ENSBTAG00000039016 and *LOC101902869*) exhibited the strongest signatures of selection. Of the

genes with the strongest signatures of selection, one gene each from Eastern

(ENSBTAG00000045571) and Western Finncattle (ENSBTAG00000012326) and three genes from

Yakutian cattle (ENSBTAG00000015464, ENSBTAG00000025621, ENSBTAG00000039016)

lacked gene descriptions (Table S6).

A total of 28, 67 and 13 GO terms were significantly enriched in Eastern Finncattle, Western

Finncattle and Yakutian cattle, respectively (Supplementary Data 19, 20 and 21). We found only one

significantly enriched GO term ("GMP binding, GO:0019002") that was shared by the three cattle

breeds. The GO terms "Homophilic cell adhesion, GO:0007156", "Calcium-dependent cell-cell

adhesion,GO:0016339" and "Multicellular organism reproduction, GO:0032504" were shared by the

Finncattle breeds. Most of the significantly enriched GO terms (23, 62 and 12 in Eastern Finncattle,

Western Finncattle and Yakutian cattle, respectively) were 'breed-specific' in our data. In addition,

we examined the significantly enriched GO terms that were potentially involved in cold adaptation

by assuming that in extremely cold environments, energy requirement is high and fat and lipids are

the main sources of energy (Liu et al., 2014). The levels of fatty acids, lipids and phospholipids

typically increase with decreasing temperatures (Purać et al., 2011). The significantly enriched GO

terms associated with Western Finncattle included "Lipid localization, GO:0010876", "Lipid

digestion, GO:0044241", "Unsaturated fatty acid biosynthetic process, GO:0006636" and

"Unsaturated fatty acid metabolic process, GO:0033559 ". However, no significantly enriched GO

terms associated with fatty acid and lipid metabolism and biosynthesis were identified in Eastern

Finncattle and Yakutian cattle.

We examined the candidate selective sweep genes in each breed. A number of genes potentially

associated with cold adaptation (Cardona et al., 2014) were present in Eastern Finncattle (*DNAJC28,*

HSP90B1, AGTRAP, TAF7, TRIP13, NPPA and NPPB), Western Finncattle (*CD14, COBL,*

JMJD1C, KCNMA1, PLA2G4, SERPINF2, SRA1 and *TAF7*) and Yakutian cattle (*DNAJC9, SOCS3,*

TRPC7, SLC8A1 GLP1R, PKLR and *TCF7L2*).

Among the selective sweep genes, there were several genes that have been previously shown to be

associated with domestication-related changes, such as changes in disease resistance, neuronal and

brain development, growth, meat quality, pigmentation, sensory perception and milk production

(Gutiérrez et al., 2015). For example, the chromosomal regions exhibiting selective sweeps in

Eastern Finncattle included genes associated with disease resistance (*IFNAR1, IFNAR2, IL10RB* and

NOD2), neuronal and brain development (*OLIG1*), growth (*ACTA1*) and meat quality (*IGFBP5,*

NRAP, PC and *S1PR1*) (Supplementary Data 13). In Western Finncattle, selective sweeps were

detected in genes associated with pigmentation (*ULBP3*), sensory perception (*LOC521946,*

LOC783558 and *LOC783323*), meat quality (*COX5B, KAT2B* and *ITGB3*) and disease resistance

(*CD96, CD14, GZMB* and *IL17A*) (Supplementary Data 14). Similarly, selective sweep-influenced

genes in Yakutian cattle were associated with disease resistance (*PFKM, ADAM17* and *SIRPA*),

sensory perception (*OR13C8, LOC100336881, LOC101902265, LOC512488, LOC617388,*

LOC783884, LOC788031 and *LOC789957*), meat quality (*ALDH1B1, CAPNS1, COX7A1, PFKM,*

SLC8A1, SOCS3 and *THBS3*) and milk production (*MUC1*) (Supplementary Data 15).

Population genetics analysis

The overall genome-wide genetic diversity, as measured by Watterson's θ and pairwise nucleotide

diversity (π), were higher in the Yakutian cattle (0.001588 and 1.728×10^{-3}, respectively) than in

Eastern Finncattle (0.001445 and 1.559×10^{-3}, respectively) and Western Finncattle (0.001398 and

1.512×10^{-3}, respectively), and these results were inconsistent with those of previous studies based

on autosomal microsatellite and SNP data sets, which showed that Finncattle were more diverse than

the Yakutian cattle (Li and Kantanen, 2010).

We also applied PCA to examine the genetic relationships among the three cattle breeds. In the PCA

plot, the Finncattle and Yakutian cattle were grouped in the first eigenvectors, indicating clear

genetic differentiation (Figure S4). The inbred Eastern Finncattle animal grouped separately from the

other Finncattle animals.

Demographic population size history

The PSMC profiles of the contemporary Finnish and Siberian native cattle were used to construct the

demographic prehistory and evolution of ancestral populations of northern Eurasian cattle. As shown

in Figure 3, the temporal PSMC profiles of the three cattle genomes followed a similar pattern. The

ancestral species of northern Eurasian taurine cattle, the near-eastern aurochs (*Bos primigenius*)

(Kantanen et al., 2009a), experienced two population peaks starting at ~1 Mya and ~40 kya and two

bottlenecks at ~250 kya and ~12 kya (Figure 3). After the first population expansion, the population

size declined gradually. The second population expansion of the ancestral wild species began around

~80 kya and started to decline around ~30 kya, leading to a second bottleneck.

We also used the $\partial a \partial i$ program to reconstruct the recent northern European cattle demographic

history (from 418 kya to the present). The parameters Ta, Td, nua, nu1F and nu2Y in the

demographic model are shown and explained in Figure S5 and Table S7. Based on this model, we

estimated that the reference ancestral population size (NA) was 43,116. The optimal model fit for

each parameter and confidence interval (CI) are shown in Table S7 by fixing NA at 43,116 and

generation time at 5 years. Our best-fit model indicated that the ancestral population underwent a size

change to 51,883 (CI, 51,658-52,108) at 418 kya (95% CI, 413.96-409.47 kya) (Table S7). This

result is consistent with the PSMC profile (Figure 3). In addition, our model suggested that the

divergence of North European native cattle and East Siberian turano-mongolicus type of cattle

occurred 8,822 years ago (CI, 8,775-8,869 years ago).

DISCUSSION

To our knowledge, this is the first whole-genome sequence-based report on the genetic diversity of

Eurasian native cattle (*B. taurus*) breeds that have adapted to the northernmost cattle farming regions,

even subarctic regions. The contemporary genetic resources of the Eastern Finncattle, Western

Finncattle and Yakutian cattle breeds studied are the result of a complex process of genetic and

demographic events that occurred during the domestication and selection and even the evolution of

the ancestral species of northern Eurasian taurine cattle, namely, the near-eastern aurochs (*B.*

primigenius).

Demographic evolution of *Bos primigenius*

As shown in Figure 3, the auroch species (*B. primigenius*) experienced two notable prehistorical

population expansions, after which the population size declined gradually. The first marked decline

in the effective population size (Ne) occurred during the Middle Pleistocene period starting after ~1

Mya, which may have been associated with reduction in global temperatures and even with negative

actions of humans on the auroch population (Barnosky et al., 2004; Hughes et al., 2007). The second

marked decline in Ne prior to domestication was obviously caused by dramatic climate changes

during the last glacial maximum (Yokoyama et al., 2000). Although the sequencing depth attained in

this study was not ideal for PSMC analysis (typically >20×), our observations regarding the temporal

changes in the Ne of the aurochs during the Pleistocene period (Mei et al., 2018) followed the pattern

observed for ancestral populations of several other domestic mammalian species, such as pig (*Sus*

scrofa; (Groenen et al., 2012)), horse (*Equus caballus*; (Librado et al., 2016)) and sheep (*Ovis aries*;

(Yang et al., 2016)). The $\partial a \partial i$ results confirmed the past fluctuations in the prehistorical Ne of *B.*

primigenius (Table S7), and the comparison of the current SNP-based estimated Ne of the present

cattle breeds (~100; (Iso-Touru et al., 2016)) to the Ne of the corresponding early domesticated

ancestral populations showed that there was a dramatic decline in the Ne during domestication and

breed formation. In addition, our demographic analysis (Figure S5) provided new knowledge of the

prehistory of northern Eurasian native cattle. As suggested by a previous study (Kantanen et al.,

2009b), both the Finnish and Yakutian native cattle descended from the near-eastern aurochs

domesticated 8,000-10,000 years ago. Here, our results have shown that the two northern Eurasian

native cattle lineages may have already diverged in the early stage of taurine cattle domestication,

more than 8,000 years ago.

High genetic variability in the Yakutian cattle

The total number of sequence variants identified on average in Eastern Finncattle and Western

Finncattle animals (e.g., 5.88 M and 6.03 M SNPs, respectively, exhibiting a minor allele frequency

> 0.05) corresponded well to numbers found typically in European taurine animals. In contrast, we

found that the Yakutian cattle exhibited a higher number of SNPs on average per individual (7.12 M

SNPs) than the number of SNPs detected in European and Asiatic humpless cattle to date (Tsuda et

al., 2013; Choi et al., 2014; Szyda et al., 2015). According to (Szyda et al., 2015) and studies cited

therein, a European taurine animal may exhibit on average 2.06-6.12, 5.89-6.37, 5.85-6.40 and 5.93

M SNPs, while (Choi et al., 2014) detected 5.81M SNPs in a Korean Holstein cattle individual, a

breed that originated from western Europe and North America. Typically, it may be possible to detect

additional SNPs by increasing the sequencing depth (Szyda et al., 2015). In addition to the average

number of SNPs per individual, total number of SNPs and number of indels, the Yakutian cattle

exhibited the highest number of exonic SNPs and nsSNPs among the three northern native breeds

studied. However, although the Yakutian cattle had the highest number of nsSNPs and genes with >5

nsSNPs, the functional annotation of the exonic SNPs by GO analysis indicated that the lowest

number of significantly enriched GO terms was obtained for the Yakutian cattle.

Our estimates for the population-level diversity for the Eastern Finncattle, Western Finncattle and

Yakutian cattle (the nucleotide diversity (π) values were 1.559×10^{-3}, 1.512×10^{-3} and 1.728×10^{-3},

respectively, and the proportions of polymorphic sites (θ) were 0.001445, 0.001398 and 0.001588,

respectively) exceed those typically found in European taurine cattle breeds (Kim et al., 2017; Chen

et al., 2018; Mei et al., 2018). We observed that Yakutian cattle such as the Asiatic taurine cattle

breeds exhibit high levels of genomic diversity in terms of π and θ estimates. The typical nucleotide

diversity values for the European taurine cattle are $>1.0 \times 10e-3$, while those for the Asiatic taurine

breeds are closer to $\sim 2.0 \times 10e-3$ than to $1.0 \times 10e-3$ (Kim et al., 2017; Chen et al., 2018; Mei et al.,

2018). We observed higher within-population diversity for the Yakutian cattle than that observed for

several other taurine cattle breeds, which differs from previous estimates based on autosomal

microsatellites and whole-genome SNP data (Li et al., 2007; Iso-Touru et al., 2016), where lower

levels of variation were observed in Yakutian cattle, indicating that the genetic variation in Yakutian

cattle has been underestimated. The set of autosomal microsatellites recommended by FAO (the Food

and Agricultural Organization of the United Nations) for biodiversity analysis of cattle breeds and the

design of commercial SNP BeadChips used in cattle whole-genome genotyping were derived mainly

from the genetic data of western breeds, causing a bias in the diversity estimates of clearly

genetically distinct cattle breeds, such as Yakutian cattle (Li et al., 2007; Iso-Touru et al., 2016).

There could have been differences in the past effective population sizes of the European and Asiatic

taurine cattle, and the present elevated genomic diversity of the Asiatic taurine cattle breeds may

reflect the higher "ancient" effective sizes of the ancestral populations of the Asiatic taurine breeds

(Chen et al., 2018). However, the prehistory of domesticated cattle in East Asia appears to be more

complex than previously thought (Zhang et al., 2013; Gao et al., 2017; Chen et al., 2018), and an

additional speculative explanation for the elevated genomic diversity in the Yakutian cattle and

several other Asiatic taurine cattle breeds (or their ancestral populations) could be ancient

introgression with the East Asian aurochs (*B. primigenius*) that lived in the East Asian region during

the arrival of near-eastern taurine cattle (Chen et al., 2018). The previous mtDNA and Y-

chromosomal diversity study indicated the near-eastern origins of the ancestral population of the

Yakutian cattle (Kantanen et al., 2009b). The possible hybrid origins of the Yakutian cattle ancestries

may have increased the genetic variation in the ancestral population of Yakutian cattle seen even in

the current population and may have played a pivotal role in the process of adaptation of the

Yakutian cattle to the subarctic environment in the Sakha Republic, eastern Siberia.

The high number of SNPs and high genomic diversity found in the Yakutian cattle may be due partly

to the breed's selection history: the artificial selection by humans has not been intensive (Kantanen et

al., 2009b). The Yakutian cattle breed is an aboriginal taurine population, the gene pool of which has

been shaped by natural and artificial selection. However, the centuries-old "folk selection" methods

and traditional knowledge for the selection of the most suitable animals for the challenging subarctic

environment followed the methods used by local people rather than the breeding implemented by

organizations or institutions (Kantanen et al., 2009a). When compared with the Western Finncattle

and Eastern Finncattle in the present study, the Yakutian cattle exhibited distinctly low numbers of

candidate genes that exhibited selection signatures (n=371, n=331 and n=249, respectively). Among

these three breeds, Western Finncattle have been subjected to the most intensive artificial selection

for milk production characteristics, while the production selection program of Eastern Finncattle was

stopped in the 1960s, when the census population size of this native breed declined rapidly.

Currently, *in vivo* and *in vitro* conservation activities are being implemented for Eastern Finncattle

(and for Western Finncattle and Yakutian cattle). In addition, although Yakutian cattle had the

highest number of genes containing SNPs (also nsSNPs) among the three breeds, the GO analysis

indicated that this breed had the lowest number of significantly enriched GO terms (Eastern

Finncattle, 111; Western Finncattle, 113; and Yakutian cattle, 95). This difference between the native

Finnish cattle and Yakutian cattle can be due to the differences in the selection histories of these

breeds.

Genomic characteristics of the northern Eurasian taurine cattle breeds

The GO enrichment analysis of genes harboring >5 nsSNPs indicated that genes related, e.g., to

immunity and "response to stimulus" are overrepresented in the set of genes identified in the northern

Eurasian native cattle breeds in this study. "Response to stimulus" refers to a change in the state or

activity of a cell or an organism as a result of the detection of a stimulus, e.g., a change in enzyme

production or gene expression (Gene Ontology Browser). This observation was consistent with

previous cattle resequencing analyses (Choi et al., 2014; Stafuzza et al., 2017; Mei et al., 2018) and

suggests that these genes were under positive selection during the course of cattle evolution and

provided survival benefits, e.g., during environmental changes (Nielsen et al., 2007). Interestingly,

genes related to the GO term "Sensory perception" were enriched in Eastern Finncattle and Western

Finncattle but not Yakutian cattle. We performed a manual search for genes associated with "Sensory

perception" genes. We found that 47 of these genes exhibited >5 nsSNPs in Eastern Finncattle and

Western Finncattle, most of which were olfactory receptor genes. We determined the number of

SNPs and nucleotide diversity of this set of genes and found that the Yakutian cattle exhibited less

variation than the two Finnish native breeds (the number of SNPs and π-estimates for Eastern

Finncattle, Western Finncattle and Yakutian cattle were 2,298 and $1.864 \times 10e\text{-}3$; 2,091 and $1.792 \times$

$10e\text{-}3$; 1,478 and $1.113 \times 10e\text{-}3$, respectively), which is in contrast to the number of SNPs and π-

estimates obtained for the entire genomes of the breeds. Great variations in the number of olfactory

receptor genes and structural variations in these genes among mammalian species and even

individuals within species (e.g., in humans) have been interpreted as reflecting the effects of

environmental factors on the genetic diversity of this multigene family and demonstrate the

importance of these genes from the evolutionary point of view (Niimura, 2011; Niimura et al., 2014).

Therefore, we hypothesize that the reduced genetic diversity in the evolutionarily important genes in

Yakutian cattle could be associated with gradual adaptation to the challenging subarctic environment

along with human movements from the southern Siberian regions to more northern environment

(Librado et al., 2015). Cattle (and horses) may have been introduced to the Yakutian region after the

9th century, perhaps as late as the 13th century (Kopoteva and Partanen, 2009). Compared to

European taurine cattle, this is a relatively short time period in terms of intervals between cattle

generations. In our study, genes related to the GO term "Developmental growth" were enriched in

only Yakutian cattle. (Stothard et al., 2011) suggested that genes associated with the GO term

"Growth" may be related to the increase in the mass of intensively selected Black Angus (beef breed)

and Holstein (dairy breed) cattle. However, Yakutian cattle have not been selected for increased body

size as that would be less desirable characteristic in Yakutian conditions. Instead, we hypothesize that

the enrichment of these growth-related genes in Yakutian cattle may be a signature of adaptation to

the harsh environment. The Yakutian cattle exhibit unique morphoanatomical adaptations to the

subarctic climate and are characterized by their small live weights (adult cows typically weigh 350-

400 kilograms, with heights of 110-112 centimeters); deep but relatively narrow chests; and short,

firm legs (Kantanen et al., 2009a). The Yakutian cattle are unique remnants of the Siberian Turano-

Mongolian type of taurine cattle (Kantanen et al., 2009a) and can be distinguished from the European

humpless cattle by these anatomical characteristics.

We performed genome-wide selection-mapping scans for the three northern cattle breeds and found a

great majority of SNPs exhibiting selection signatures in noncoding genomic regions. This finding

indicates that selection occurs specifically via the regulatory elements of genomes (see also (Librado

et al., 2015)). We found that the studied breeds exhibited 'private' (breed-specific) selection

signature patterns, indicating distinctiveness in their selection histories. We further investigated the

proportions of genes exhibiting selection signatures among the breeds and found that only 5 genes

from this set of genes were shared by the three breeds. Only 37 genes were shared by the two Finnish

native cattle breeds, while 13 'selection signature' genes were shared by Western Finncattle and

Yakutian cattle and 11 by Eastern Finncattle and Yakutian cattle. In addition, the breeds did not share

any of the genes exhibiting the strongest selection signatures and harboring >5 nsSNPs, and the GO

term enrichment analysis of this set of genes indicated that only one GO term ("GMP binding") was

significantly enriched in all three breeds.

We identified several positively selected candidate genes underlying adaptation, appearance and

production of Eastern Finncattle, Western Finncattle and Yakutian cattle. For example, in Eastern

Finncattle, selection signatures were detected in *NRAP* and *IGFBP5*, both of which have been

previously identified as candidate genes for muscle development and meat quality in cattle (Williams

et al., 2009), and in *NOD2*, which is a candidate gene for dairy production (Ogorevc et al., 2009). In

Western Finncattle, we detected selection signatures in, e.g., candidate genes for beef production,

such as *COX5B* and *ITGB3* (Williams *et al.*, 2009), and dairy production, such as *CD14* (Ogorevc et

al., 2009). In Yakutian cattle, several genes exhibiting selection signatures were candidate genes for

muscle development and meat quality, such as *COX7A1*, *THBS3*, *PFKM*, and *SOCS3* (Williams *et*

al., 2009) but also for color pattern (*ADAM17*; (Gutiérrez-Gil et al., 2015)) and milk production traits

(*MUC1*; (Ogorevc et al., 2009)). We were particularly interested in the genomic adaptation to North

Eurasian environments. (Cardona et al., 2014) listed in the supplementary materials of their

publication several potential candidate genes associated with biological processes and pathways

hypothesized to be involved in cold adaptation in indigenous Siberian human populations in terms of

response to temperature, blood pressure, basal metabolic rate, smooth muscle contraction and energy

metabolism. Several of these genes also exhibited significant selection signatures in our cattle

sequence data, as exemplified in the Results section of this paper. *SLC8A1* (sodium/calcium

exchanger 1), influencing the oxidative stress response, is an example of the genes with significant

selection signatures in Yakutian cattle, Siberian human populations (Cardona et al., 2014) and native

Yakutian horses (Librado et al., 2015). This example of selection signatures and associated genes

found in the Yakutian cattle and Siberian human populations (Cardona et al., 2014) indicates

convergent evolution between the mammalian populations adapted to subarctic environments.

Convergent evolution between mammalian species in adaptation to harsh environments has also

occurred, e.g., on the Tibetan plateau, as indicated by (Wang et al., 2014a, 2015; Yang et al., 2016).

CONCLUSIONS

We have investigated by whole-genome sequencing for the first time the genetic diversity of native

cattle breeds originating from the northernmost region of cattle farming in the world. We found novel

SNPs and indels and genes that have not yet been annotated. Our observations suggest that accurate

reference genome assemblies are needed for genetically diverse native cattle breeds showing genetic

distinctiveness, such as Yakutian cattle, in order to better understand the genetic diversity of the

breeds and the effects of natural and artificial selection and adaptation. We identified a number of

genes and chromosomal regions important for the adaptation and production traits of the breeds.

Moreover, GO terms such as defense response, growth, sensory perception and immune response

were enriched in the genes associated with selective sweeps. To improve our knowledge of the value

of native breeds as genetic resources for future cattle breeding and the power of selection signature

analyses, a greater number of animals of these breeds should be investigated in a wider breed

diversity context.

ABBREVIATIONS

nsSNPs: nonsynonymous SNPs; GO: gene ontology; CLR: composite likelihood ratio; SFS: site

frequency spectrum; PCA: principal component analysis; PSMC: pairwise sequentially Markovian

coalescent; $\partial a \partial i$: diffusion approximation for demographic inference; Gb: gigabases

DATA AVAILABILITY

The raw sequence reads (Fastq Files) for this study can be found in European Nucleotide Archive

(ENA) under the accession number PRJEB28185 (please see Table S1 for sample specific

accessions).

AUTHOR CONTRIBUTIONS

JK designed the study, and revised the manuscript. MW performed the bioinformatics and statistical

analyses and drafted the manuscript. JK, RP, IA and ZI collected the samples. RP, KP, IA, MY and

ZI participated in the experimental design and paper revision. All authors read and approved the final

manuscript.

FUNDING

This work was supported by the Academy of Finland (the Arctic Ark—project no. 286040 in the

ARKTIKO research program of the Academy of Finland), the Ministry of Agriculture and Forestry

in Finland, and the Finnish Cultural Foundation. MW was partly supported by grants from the Betty

Väänänen Foundation and Ella and Georg Ehrnrooth foundation, Doctoral School of University of

Eastern Finland in Environmental Physics, Health and Biology.

ACKNOWLEDGMENTS

We thank Tuula-Marjatta Hamama for DNA extraction and BGI for sequencing the samples. The

authors wish to acknowledge the CSC-IT Center for Science, Finland, for computational resources.

The owners of the animals included in the study are acknowledged for providing samples for this

study.

REFERENCES

Alachiotis, N., Stamatakis, A., and Pavlidis, P. (2012). OmegaPlus: a scalable tool for rapid detection

of selective sweeps in whole-genome datasets. *Bioinformatics* 28, 2274–2275.

doi:10.1093/bioinformatics/bts419.

Barnosky, A. D., Koch, P. L., Feranec, R. S., Wing, S. L., and Shabel, A. B. (2004). Assessing the

Causes of Late Pleistocene Extinctions on the Continents. *Science (80-.).* 306, 70. Available at:

http://science.sciencemag.org/content/306/5693/70.abstract.

Bläuer, A., and Kantanen, J. (2013). Transition from hunting to animal husbandry in Southern,

Western and Eastern Finland: new dated osteological evidence. *J. Archaeol. Sci.* 40, 1646–1666.

doi:http://dx.doi.org/10.1016/j.jas.2012.10.033.

Boettcher, P. J., Tixier-Boichard, M., Toro, M. A., Simianer, H., Eding, H., Gandini, G., et al. (2010).

Objectives, criteria and methods for using molecular genetic data in priority setting for

conservation of animal genetic resources. *Anim. Genet.* 41, 64–77. doi:10.1111/j.1365-

2052.2010.02050.x.

Browning, S. R., and Browning, B. L. (2007). Rapid and Accurate Haplotype Phasing and Missing-

Data Inference for Whole-Genome Association Studies By Use of Localized Haplotype

Clustering. *Am. J. Hum. Genet.* 81, 1084–1097. Available at:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2265661/.

Cardona, A., Pagani, L., Antao, T., Lawson, D. J., Eichstaedt, C. A., Yngvadottir, B., et al. (2014).

Genome-Wide Analysis of Cold Adaptation in Indigenous Siberian Populations. *PLoS One* 9,

e98076. doi:10.1371/journal.pone.0098076.

Chen, N., Cai, Y., Chen, Q., Li, R., Wang, K., Huang, Y., et al. (2018). Whole-genome resequencing

reveals world-wide ancestry and adaptive introgression events of domesticated cattle in East

Asia. *Nat. Commun.* 9, 2337. doi:10.1038/s41467-018-04737-0.

Choi, J.-W., Liao, X., Park, S., Jeon, H.-J., Chung, W.-H., Stothard, P., et al. (2013). Massively

Parallel Sequencing of Chikso (Korean Brindle Cattle) to Discover Genome-Wide SNPs and

InDels. *Mol. Cells* 36, 203–211. doi:10.1007/s10059-013-2347-0.

Choi, J.-W., Liao, X., Stothard, P., Chung, W.-H., Jeon, H.-J., Miller, S. P., et al. (2014). Whole-

Genome Analyses of Korean Native and Holstein Cattle Breeds by Massively Parallel

Sequencing. *PLoS One* 9, e101127. Available at:

http://dx.doi.org/10.1371%2Fjournal.pone.0101127.

Cramp, L. J. E., Jones, J., Sheridan, A., Smyth, J., Whelton, H., Mulville, J., et al. (2014). Immediate

replacement of fishing with dairying by the earliest farmers of the northeast Atlantic

archipelagos. *Proc. R. Soc. B Biol. Sci.* 281, 20132372. doi:10.1098/rspb.2013.2372.

DePristo, M. A., Banks, E., Poplin, R. E., Garimella, K. V, Maguire, J. R., Hartl, C., et al. (2011). A

framework for variation discovery and genotyping using next-generation DNA sequencing data.

Nat. Genet. 43, 491–498. doi:10.1038/ng.806.

Du, Z., Zhou, X., Ling, Y., Zhang, Z., and Su, Z. (2010). agriGO: a GO analysis toolkit for the

agricultural community. *Nucleic Acids Res.* 38, W64–W70. doi:10.1093/nar/gkq310.

Egorov, E. G., Nikiforov, M. M., and Danilov, Y. G. (2015). Unique experience and achievements of

sakha people in development of agriculture in the north. 9.

Felius, M. (1995). *Cattle Breeds, an Encyclopedia; Misset Uitgeverij: Doetinchem, The Netherlands*.

Flicek, P., Ahmed, I., Amode, M. R., Barrell, D., Beal, K., Brent, S., et al. (2013). Ensembl 2013.

Nucleic Acids Res. 41, D48–D55. doi:10.1093/nar/gks1236.

Gao, Y., Gautier, M., Ding, X., Zhang, H., Wang, Y., Wang, X., et al. (2017). Species composition

and environmental adaptation of indigenous Chinese cattle. *Sci. Rep.* 7, 16196.

doi:10.1038/s41598-017-16438-7.

Groenen, M. A. M., Archibald, A. L., Uenishi, H., Tuggle, C. K., Takeuchi, Y., Rothschild, M. F., et

al. (2012). Analyses of pig genomes provide insight into porcine demography and evolution.

Nature 491, 393–398. doi:10.1038/nature11622.

Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., and Bustamante, C. D. (2009). Inferring the

Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency

Data. *PLoS Genet* 5, e1000695. Available at:

http://dx.doi.org/10.1371%2Fjournal.pgen.1000695.

Gutiérrez-Gil, B., Arranz, J. J., and Wiener, P. (2015). An interpretive review of selective sweep

studies in Bos taurus cattle populations: identification of unique and shared selection signals

across breeds. *Front. Genet.* 6, 167. Available at:

https://www.frontiersin.org/article/10.3389/fgene.2015.00167.

Hughes, P. D., Woodward, J. C., and Gibbard, P. L. (2007). Middle Pleistocene cold stage climates in

the Mediterranean: New evidence from the glacial record. *Earth Planet. Sci. Lett.* 253, 50–56.

doi:https://doi.org/10.1016/j.epsl.2006.10.019.

Iso-Touru, T., Tapio, M., Vilkki, J., Kiseleva, T., Ammosov, I., Ivanova, Z., et al. (2016). Genetic

diversity and genomic signatures of selection among cattle breeds from Siberia, eastern and

northern Europe. *Anim. Genet.* 47, 647–657. doi:10.1111/age.12473.

Kantanen, J., Ammosov, I., Li, M., Osva, A., and Popov, R. (2009a). "A cow of the permafrost," in

Sakha Ynaga□: cattle of the Yakuts (Helsinki: Suomalaisen tiedeakatemian toimituksia), 19–44.

Kantanen, J., Edwards, C. J., Bradley, D. G., Viinalass, H., Thessler, S., Ivanova, Z., et al. (2009b).

Maternal and paternal genealogy of Eurasian taurine cattle (Bos taurus). *Heredity (Edinb).* 103,

404–415. Available at:

http://www.nature.com/hdy/journal/v103/n5/suppinfo/hdy200968s1.html.

Kantanen, J., Låvendahl, P., Strandberg, E., Eythorsdottir, E., Li, M.-H., Kettunen-Præbel, A., et al.

(2015). Utilization of farm animal genetic resources in a changing agro-ecological environment

in the Nordic countries. *Front. Genet.* 6, 52. doi:10.3389/fgene.2015.00052.

Kantanen, J., Olsaker, I., Holm, L.-E., Lien, S., Vilkki, J., Brusgaard, K., et al. (2000). Genetic

diversity and population structure of 20 north European cattle breeds. *J. Hered.* 91, 446–457.

doi:10.1093/jhered/91.6.446.

Kawahara-Miki, R., Tsuda, K., Shiwa, Y., Arai-Kichise, Y., Matsumoto, T., Kanesaki, Y., et al.

(2011). Whole-genome resequencing shows numerous genes with nonsynonymous SNPs in the

Japanese native cattle Kuchinoshima-Ushi. *BMC Genomics* 12, 103. Available at:

http://www.biomedcentral.com/1471-2164/12/103.

Kim, J., Hanotte, O., Mwai, O. A., Dessie, T., Bashir, S., Diallo, B., et al. (2017). The genome

landscape of indigenous African cattle. *Genome Biol.* 18, 34. doi:10.1186/s13059-017-1153-y.

Kinsella, R. J., Kähäri, A., Haider, S., Zamora, J., Proctor, G., Spudich, G., et al. (2011). Ensembl

BioMarts: a hub for data retrieval across taxonomic space. *Database* 2011, bar030-bar030.

Available at: http://dx.doi.org/10.1093/database/bar030.

Kopoteva, I., and Partanen, U. (2009). "Sakha Ynaga□: Cattle of the Yakuts.," in *A historical*

excursion to Northern Sakha., eds. J. Granberg, K. Soini, and J. Kantanen (Helsinki:

Suomalainen Tiedeakatemia), 75–116.

Kumar, S., and Subramanian, S. (2002). Mutation rates in mammalian genomes. *Proc. Natl. Acad.*

Sci. 99, 803–808. doi:10.1073/pnas.022629899.

Lachance, J., Vernot, B., Elbers, C. C., Ferwerda, B., Froment, A., Bodo, J.-M., et al. (2012).

Evolutionary history and adaptation from high-coverage whole-genome sequences of diverse

African hunter-gatherers. *Cell* 150, 457–469. doi:10.1016/j.cell.2012.07.009.

Li, H., and Durbin, R. (2011). Inference of human population history from individual whole-genome

sequences. *Nature* 475, 493–496. Available at:

http://www.nature.com/nature/journal/v475/n7357/abs/nature10231.html#supplementary-

information.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The Sequence

Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.

doi:10.1093/bioinformatics/btp352.

Li, M.-H., and Kantanen, J. (2010). Genetic structure of Eurasian cattle (*Bos taurus*) based on

microsatellites: clarification for their breed classification. *Anim. Genet.* 41, 150–158.

doi:10.1111/j.1365-2052.2009.01980.x.

Li, M., Tapio, I., Vilkki, J., Ivanova, Z., Kiselyova, T., Marzanov, N., et al. (2007). The genetic

structure of cattle populations (Bos taurus) in northern Eurasia and the neighbouring Near

Eastern regions: implications for breeding strategies and conservation. *Mol. Ecol.* 16, 3839–

3853. doi:10.1111/j.1365-294X.2007.03437.x.

Li, M., Tian, S., Yeung, C. K. L., Meng, X., Tang, Q., Niu, L., et al. (2014). Whole-genome

sequencing of Berkshire (European native pig) provides insights into its origin and

domestication. *Sci.Rep.* 4. Available at: http://10.0.4.14/srep04678.

Librado, P., Der Sarkissian, C., Ermini, L., Schubert, M., Jónsson, H., Albrechtsen, A., et al. (2015).

Tracking the origins of Yakutian horses and the genetic basis for their fast adaptation to

subarctic environments. *Proc. Natl. Acad. Sci.* 112, E6889–E6897.

doi:10.1073/pnas.1513696112.

Librado, P., Fages, A., Gaunitz, C., Leonardi, M., Wagner, S., Khan, N., et al. (2016). The

Evolutionary Origin and Genetic Makeup of Domestic Horses. *Genetics* 204, 423–434.

doi:10.1534/genetics.116.194860.

Liu, S., Lorenzen, E. D., Fumagalli, M., Li, B., Harris, K., Xiong, Z., et al. (2014). Population

Genomics Reveal Recent Speciation and Rapid Evolutionary Adaptation in Polar Bears. *Cell*

157, 785–794. doi:10.1016/j.cell.2014.03.054.

MacLeod, I. M., Larkin, D. M., Lewin, H. A., Hayes, B. J., and Goddard, M. E. (2013). Inferring

Demography from Runs of Homozygosity in Whole-Genome Sequence, with Correction for

Sequence Errors. *Mol. Biol. Evol.* 30, 2209–2223. doi:10.1093/molbev/mst125.

Malke, H. (1990). J. SAMBROCK, E. F. FRITSCH and T. MANIATIS, Molecular Cloning, A

Laboratory Manual (Second Edition), Volumes 1, 2 and 3. 1625 S., zahlreiche Abb. und Tab.

Cold Spring Harbor 1989. Cold Spring Harbor Laboratory Press. $ 115.00. ISBN: 0-87969-309-

6. *J. Basic Microbiol.* 30, 623. doi:10.1002/jobm.3620300824.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., et al. (2010). The

Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Res.* 20, 1297–1303. doi:10.1101/gr.107524.110.

McManus, K. F., Kelley, J. L., Song, S., Veeramah, K. R., Woerner, A. E., Stevison, L. S., et al.

(2015). Inference of Gorilla Demographic and Selective History from Whole-Genome Sequence

Data. *Mol. Biol. Evol.* 32, 600–612. doi:10.1093/molbev/msu394.

Mei, C., Wang, H., Liao, Q., Wang, L., Cheng, G., Wang, H., et al. (2018). Genetic Architecture and

Selection of Chinese Cattle Revealed by Whole Genome Resequencing. *Mol. Biol. Evol.* 35,

688–699. doi:10.1093/molbev/msx322.

Murray, C., Huerta-Sanchez, E., Casey, F., and Bradley, D. G. (2010). Cattle demographic history

modelled from autosomal sequence variation. *Philos. Trans. R. Soc. London B Biol. Sci.* 365,

2531–2539. Available at: http://rstb.royalsocietypublishing.org/content/365/1552/2531.abstract.

Nielsen, R., Hellmann, I., Hubisz, M., Bustamante, C., and Clark, A. G. (2007). Recent and ongoing

selection in the human genome. *Nat. Rev. Genet.* 8, 857–868. doi:10.1038/nrg2187.

Niimura, Y. (2011). Olfactory Receptor Multigene Family in Vertebrates: From the Viewpoint of

Evolutionary Genomics. *Curr. Genomics* 13, 103–114. doi:10.2174/138920212799860706.

Niimura, Y., Matsui, A., and Touhara, K. (2014). Extreme expansion of the olfactory receptor gene

repertoire in African elephants and evolutionary dynamics of orthologous gene groups in 13

placental mammals. *Genome Res.* 24, 1485–1496. doi:10.1101/gr.169532.113.

Odegård, J., Yazdi, M. H., Sonesson, A. K., and Meuwissen, T. H. E. (2009). Incorporating desirable

genetic characteristics from an inferior into a superior population using genomic selection.

Genetics 181, 737–45. doi:10.1534/genetics.108.098160.

Ogorevc, J., Kunej, T., Razpet, A., and Dovc, P. (2009). Database of cattle candidate genes and

genetic markers for milk production and mastitis. *Anim. Genet.* 40, 832–851.

doi:10.1111/j.1365-2052.2009.01921.x.

Patterson, N., Price, A. L., and Reich, D. (2006). Population Structure and Eigenanalysis. *PLoS*

Genet 2, e190. Available at: http://dx.plos.org/10.1371%2Fjournal.pgen.0020190.

Pavlidis, P., Živkovic, D., Stamatakis, A., and Alachiotis, N. (2013). SweeD: Likelihood-Based

Detection of Selective Sweeps in Thousands of Genomes. *Mol. Biol. Evol.* 30, 2224–2234.

doi:10.1093/molbev/mst112.

Purać, J., Pond, D. W., Grubor-Lajšić, G., Kojić, D., Blagojević, D. P., Worland, M. R., et al. (2011).

Cold hardening induces transfer of fatty acids between polar and nonpolar lipid pools in the

Arctic collembollan Megaphorura arctica. *Physiol. Entomol.* 36, 135–140. doi:10.1111/j.1365-

3032.2010.00772.x.

Sasaki, Y., Nagai, K., Nagata, Y., Doronbekov, K., Nishimura, S., Yoshioka, S., et al. (2006).

Exploration of genes showing intramuscular fat deposition-associated expression changes in

musculus longissimus muscle. *Anim. Genet.* 37, 40–46. doi:10.1111/j.1365-2052.2005.01380.x.

Stafuzza, N. B., Zerlotini, A., Lobo, F. P., Yamagishi, M. E. B., Chud, T. C. S., Caetano, A. R., et al.

(2017). Single nucleotide variants and InDels identified from whole-genome re-sequencing of

Guzerat, Gyr, Girolando and Holstein cattle breeds. *PLoS One* 12, e0173954. Available at:

https://doi.org/10.1371/journal.pone.0173954.

Stajich, J. E., Block, D., Boulez, K., Brenner, S. E., Chervitz, S. A., Dagdigian, C., et al. (2002). The

Bioperl Toolkit: Perl Modules for the Life Sciences. *Genome Res.* 12, 1611–1618.

doi:10.1101/gr.361602.

Stothard, P., Choi, J.-W., Basu, U., Sumner-Thomson, J. M., Meng, Y., Liao, X., et al. (2011). Whole

genome resequencing of black Angus and Holstein cattle for SNP and CNV discovery. *BMC*

Genomics 12, 559. doi:10.1186/1471-2164-12-559.

Szyda, J., FrÄ…szczak, M., Mielczarek, M., Giannico, R., Minozzi, G., Nicolazzi, E. L., et al.

(2015). The assessment of inter-individual variation of whole-genome DNA sequence in 32

cows. *Mamm. Genome* 26, 658–665. doi:10.1007/s00335-015-9606-7.

Tsuda, K., Kawahara-Miki, R., Sano, S., Imai, M., Noguchi, T., Inayoshi, Y., et al. (2013). Abundant

sequence divergence in the native Japanese cattle Mishima-Ushi (Bos taurus) detected using

whole-genome sequencing. *Genomics* 102, 372–378. doi:10.1016/j.ygeno.2013.08.002.

Van der Auwera A., G., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A., et

al. (2013). From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best

practices pipeline. *Curr. Protoc. Bioinforma. / Ed. board, Andreas D.Baxevanis ...[et al.]* 11,

11.10.1-11.10.33. doi:10.1002/0471250953.bi1110s43.

Wang, G.-D., Fan, R.-X., Zhai, W., Liu, F., Wang, L., Zhong, L., et al. (2014a). Genetic

Convergence in the Adaptation of Dogs and Humans to the High-Altitude Environment of the

Tibetan Plateau. *Genome Biol. Evol.* 6, 2122–2128. Available at:

http://dx.doi.org/10.1093/gbe/evu162.

Wang, K., Li, M., and Hakonarson, H. (2010). ANNOVAR: functional annotation of genetic variants

from high-throughput sequencing data. *Nucleic Acids Res.* 38, e164–e164.

doi:10.1093/nar/gkq603.

Wang, M.-S., Li, Y., Peng, M.-S., Zhong, L., Wang, Z.-J., Li, Q.-Y., et al. (2015). Genomic Analyses

Reveal Potential Independent Adaptation to High Altitude in Tibetan Chickens. *Mol. Biol. Evol.*

32, 1880–1889. Available at: http://dx.doi.org/10.1093/molbev/msv071.

Wang, M., Yu, Y., Haberer, G., Marri, P. R., Fan, C., Goicoechea, J. L., et al. (2014b). The genome

sequence of African rice (Oryza glaberrima) and evidence for independent domestication. *Nat.*

Genet. 46, 982–988. Available at: http://10.0.4.14/ng.3044.

Watanabe, N., Satoh, Y., Fujita, T., Ohta, T., Kose, H., Muramatsu, Y., et al. (2011). Distribution of

allele frequencies at TTN g.231054C > T, RPL27A g.3109537C > T and AKIRIN2 c.*188G >

A between Japanese Black and four other cattle breeds with differing historical selection for

marbling. *BMC Res. Notes* 4, 10. Available at: http://www.biomedcentral.com/1756-0500/4/10.

Williams, J. L., Dunner, S., Valentini, A., Mazza, R., Amarger, V., Checa, M. L., et al. (2009).

Discovery, characterization and validation of single nucleotide polymorphisms within 206

bovine genes that may be considered as candidate genes for beef production and quality. *Anim.*

Genet. 40, 486–491. doi:10.1111/j.1365-2052.2009.01874.x.

Yang, J., Li, W.-R., Lv, F.-H., He, S.-G., Tian, S.-L., Peng, W.-F., et al. (2016). Whole-Genome

Sequencing of Native Sheep Provides Insights into Rapid Adaptations to Extreme

Environments. *Mol. Biol. Evol.* 33, 2576–2592. doi:10.1093/molbev/msw129.

Yokoyama, Y., Lambeck, K., De Deckker, P., Johnston, P., and Fifield, L. K. (2000). Timing of the

Last Glacial Maximum from observed sea-level minima. *Nature* 406, 713. Available at:

http://10.0.4.14/35021035.

Zhang, H., Paijmans, J. L. A., Chang, F., Wu, X., Chen, G., Lei, C., et al. (2013). Morphological and

genetic evidence for early Holocene cattle management in northeastern China. *Nat. Commun.* 4,

2755. Available at: http://10.0.4.14/ncomms3755.

Zhao, S., Zheng, P., Dong, S., Zhan, X., Wu, Q., Guo, X., et al. (2013). Whole-genome sequencing

of giant pandas provides insights into demographic history and local adaptation. *Nat. Genet.* 45,

67–71. Available at:

http://www.nature.com/ng/journal/v45/n1/abs/ng.2494.html#supplementary-information.

Zimin, A., Delcher, A., Florea, L., Kelley, D., Schatz, M., Puiu, D., et al. (2009). A whole-genome

assembly of the domestic cow, Bos taurus. *Genome Biol.* 10, R42. Available at:

http://genomebiology.com/2009/10/4/R42.

Figures

Figure 1. Three North Eurasian native cattle breeds are included in this study. **(A)** Eastern Finncattle

are typically red-sided and polled. Cattle breeding in Finland was started with this breed, and the

breed's herd book was established in 1898. The breed was threatened with extinction in the 1970s

and 1980s. The current census size is 1,600 cows, and the annual milk yield on average 4,000 Kgs.

(B) Western Finncattle are solid light or dark brown and polled. The breed is one of the most

productive native cattle breeds: the average annual milk yield is about 7,000 Kgs.**(C)** The Yakutian

cattle are characterized by being purebred aboriginal native cattle from Sakha. Adult Yakutian cows

weigh typically 350-400 Kgs and their height at the withers is 110-112cm on average. The animals

are well adapted to Siberian harsh conditions where the temperature falls below -50°C in long

winters. The average annual milk yield is 1,000 Kgs. Please do not copy, use or upload the

photographs without permission of the copyright holders.

Figure 2. Venn diagram showing overlapping and unique SNPs/indels between the three breeds

(Eastern Finncattle, Western Finncattle and Yakutian). The numbers in parentheses outside the

circles are the total number of detected SNPs from each breed. The numbers in the circle components

show specific SNPs for each breed or overlapping SNPs/indels between any two breeds or among

three breeds. **(A)** The identified shared and specific SNPs for each breed, **(B)** the identified shared

and specific novel SNPs for each breed, and **(C)** the identified shared and specific indels for each

breed.

Figure 3 Demographic history of the northernmost cattle breed reconstructed from three cattle

genomes, one from each breed, by using PSMC. The X axis shows the time in thousand years

(Kyr), and the Y axis shows the effective population size.

Tables

Table 1. Summary of sequencing and short read alignment results

	Eastern Finncattle	Western Finncattle	Yakutian cattle	Overall sample
Number of individuals	5	5	5	15
Paired-end length (bp)	100	100	100	100

Average reads per individual	352.73 M	347.14 M	342.33 M	347.4 M
Average sequence depth per individual [a]	13.21X	13.00X	12.82X	13.01X
Average map reads per individual	348.42 M	340.12 M	337.50 M	342,02 M
Average unique map reads per individual	316.89 M	312.58 M	309.19 M	312.88 M
Average read mapping rate	98.78%	97.97%	98.59%	98.45%
Average coverage rate	98.42%	98.22%	98.46%	98.37%

822   [a] Average sequence depth per individual was computed by dividing the clean reads by the reference

823   genome size.

825   **Table 2.** Functional annotation of the detected SNPs and indels

			Eastern Finncattle	Western Finncattle	Yakutian cattle
S N P	**Total number of SNPs**		11,017,215	10,543,290	12,242,166
	Intergenic		7,998,914	7,662,604	8,845,911
	Intronic		2,764,951	2,643,821	3,114,622
	Exonic[a]				
		Nonsynonymous	30,982	28,733	32,782
		Stop gain	294	284	310
		Stop loss	23	18	19
		Synonymous	41,111	38,137	46,950
	Upstream		74,552	69,369	82,170
	Downstream		74,198	70,815	83,549
	Upstream; downstream[b]		1,636	1,534	2,036
	UTR[c]		25,545	23,342	28,273

	Splicing		417	388	459
	ncRNA		4,593	4,256	5,086
	Total number of indels		1,275,128	1,188,892	1,374,577
	Intergenic		942,143	878,007	1,012,733
	Intronic		332,504	310,089	363,783
	Exonic[a]				
I		Nonframeshift	397	327	427
n		Stop gain	24	22	27
		Stop loss	1	1	0
d		Frameshift	1,045	972	1,148
	Upstream		9,269	8,286	9,861
el	**Downstream**		10,611	9,609	11,377
	Upstream; downstream[b]		250	218	282
	UTR[c]		3,380	3,101	3,767
	Splicing		248	233	268
	ncRNA		406	371	437

826 [a] Exonic = "exonic" and "exonic; splicing" as annotated by ANNOVAR

827 [b] Upstream; downstream = variant located in downstream and upstream regions

828 [c] UTR = "UTR3" and "UTR5" as annotated by ANNOVAR