Sex differences in fear regulation and reward seeking behaviors in a fear-safety-reward discrimination task

Abbreviated title: Sex differences in conditioned safety

Eliza M. Greiner[1], Makenzie R. Norris[1], Iris Müller [1,2], Ka H. Ng[1,2], Susan Sangha[1,2]*

1. Department of Psychological Sciences
2. Purdue Institute for Integrative Neuroscience
Purdue University, West Lafayette, IN, USA 47907

*Correspondence to:
Susan Sangha
Department of Psychological Sciences
Purdue University
703 Third St., West Lafayette, IN 47907
(t) 765.494.0648
(e) sangha@purdue.edu

Number of figures: 4
Number of tables: 1

Abstract

Reward availability and the potential for danger or safety potently regulates emotion. Despite women being more likely than men to develop emotion dysregulation disorders, there are comparatively few studies investigating fear, safety and reward regulation in females. Here, we show that female Long Evans rats do not suppress conditioned freezing in the presence of a safety cue, nor do they extinguish their freezing response, whereas males do both. Females were also more reward responsive during the reward cue until the first footshock exposure, at which point there were no sex differences in reward seeking to the reward cue. Darting analyses indicate females might be able to regulate this behavior in response to the safety cue, suggesting they might be able to discriminate between fear and safety cues but do not demonstrate this with conditioned suppression of the freezing behavior. However, levels of darting in this study were too low to make any clear conclusions. In summary, females showed a significantly different behavioral profile than males in a task that tests the ability to discriminate among fear, safety and reward cues. This paradigm offers a great opportunity to test for mechanisms that are generating these behavioral sex differences in learned safety and reward seeking.

Keywords: sex differences; fear; safety; reward

1. Introduction

Clinical disorders arising from maladaptive emotion regulation present a large burden on society worldwide. Many of these disorders show comorbidity, for example, addiction with anxiety disorders (Grant et al., 2016). Cues predicting something aversive elicit avoidance and fear behaviors whereas cues predicting reward elicit approach and reward-seeking behaviors. Cues signifying safety have the power to modulate fear and reward-seeking behaviors by informing the organism whether or not the environment is safe (Walasek, Wesierska, & Zieliński, 1995). Thus, safety, fear and reward behaviors, and the circuitries governing these behaviors, are intertwined. The majority of studies on reward and fear processing have been conducted in parallel, investigating the circuitries separately in primarily male subjects. If we hope to understand and treat comorbid disorders resulting from maladaptive emotion regulation, increased efforts in investigating how these circuitries integrate their functions to influence behavior is needed in both male and female subjects.

Our laboratory has designed and validated a behavioral task in which fear, safety and reward cues are learned within the same session allowing us to assess the animal's ability to discriminate among these cues (Ng et al., 2018; Sangha et al., 2013; Sangha, Greba et al., 2014; Sangha, Robinson et al., 2014). Rats are exposed to cues associated with safety, fear (fear cue paired with footshock), and reward (reward cue paired with sucrose). Male rats consistently learn to discriminate among safety, fear and reward cues to 1) suppress conditioned freezing in the presence of a safety cue (fear+safety cue), and 2) increase reward seeking when reward is available (reward cue) (Ng et al., 2018; Sangha et al., 2013; Sangha, Greba, et al., 2014; Sangha, Robinson, et al., 2014). This paradigm also allows us to investigate how safety cues can regulate both fear and reward behaviors. Evidence suggests that reward learning mechanisms overlap with safety learning (Leknes et al., 2011; Pollak et al., 2008; Rescorla, 1969; Rogan et al., 2005; Sangha et al., 2013; Tanimoto et al., 2004; Walasek et al., 1995). For example, learned safety can act as a behavioral antidepressant in mice (Pollak et al., 2008), and animals will perform certain behaviors in order to turn on a safety signal (Rescorla, 1969; Rogan et al., 2005). Within the amygdala we have shown a subpopulation of neurons responding with the same level of excitation or inhibition during both the reward and safety cues (Sangha et al., 2013). Additionally, we have shown inactivation of the prelimbic or infralimbic cortices of the ventromedial prefrontal cortex have differential effects on reward and safety discrimination, respectively (Sangha, Robinson, et al., 2014). Thus, in male rats, our prior work

has already shown a critical involvement of the corticoamygdalar circuit in learning this fear-
safety-reward cue discrimination.

Much of the research investigating emotion regulation mechanisms have exclusively used male
subjects. In a study using male Vietnam veterans, PTSD patients show impairments in
suppressing their fear response in the presence of a safety cue (Jovanovic et al., 2009). But,
women are more than twice as likely to develop Post-Traumatic Stress Disorder (PTSD) than
men, with females having a lifetime prevalence of 8.5% in contrast to 3.4% in males (Mclean et
al., 2011). In fear studies that have included female rats, it has been shown that females exhibit
lower levels of freezing behavior than male rats after repeated fear cue presentations (Daviu et
al., 2014). These findings have been thought to indicate a difficulty in fear conditioning learning
in female rats. A more recent experiment has identified that approximately 40% of female rats
tested exhibit an alternate fear behavior in the form of fast paced movements called 'darting';
this was only seen in approximately 10% of male rats tested (Gruene et al., 2015). There is also
evidence of sex differences in the seeking of natural rewards, where it has been reported that
female rats consume more sucrose pellets than males and are willing to work harder for them
(Tapia, Lee, Weise, Tamasi, & Will, 2019). Dopamine signaling during reward tasks has also
been demonstrated to be different between sexes. For example, Conway et al (2019) showed
females continue to perform intracranial self-stimulation for brain stimulation reward while under
the influence of a kappa-opioid receptor agonist, which suppresses dopamine release, whereas
males decrease this behavior. Their data suggest that female rats may have an increased
capacity to produce and release dopamine compared to males, under these conditions. Our
prior work has shown, in males, that dopamine signaling in the basolateral amygdala contributes
to effective discrimination among fear, safety and reward cues (Ng et al., 2018).

Taken together, we hypothesized there would be sex differences in the ability to express clear
discrimination among fear, safety and reward cues. The inability of male PTSD patients to learn
safety signaling has been labeled a biomarker of the disorder (Jovanovic et al., 2012). Due to
sex-related differences in human diagnosis of PTSD, with women diagnosed at rates twice that
of men (Glover et al., 2015), any differences female rats have in the learning or retention of
safety signals could steer towards further research on the neurological processes underlying
these variations.

2. Materials and Methods

2.1 Subjects

A total of 24 adult male (215-375g) and 28 adult age-matched female (198-230g) Long Evans rats (Blue Spruce; Envigo, Indianapolis), were single-housed and handled for 1 week prior to testing. All procedures were performed during the light cycle and approved by the Purdue Animal Care and Use Committee. Rats had *ad libitum* access to food and water prior to the start of the experiment. After experiment onset, they were maintained on a food restricted diet (20g per day for males; 16g per day for females) until the last day of the experiment.

2.2 Apparatus

The rats were trained in operant conditioning chambers consisting of Plexiglas boxes (32cm length x 25cm width x 30cm height) encased in sound-attenuating chambers (Med Associates, ST Albans, VT). 10% liquid sucrose was delivered through a recessed port 2cm above the floor in the center of one wall. Two lights (28V, 100mA) were located 10.5cm from floor on either side of the port. A light (28V, 100mA) 27cm above the floor on the wall was on throughout the entire session. Auditory cues were delivered via a speaker (ENV-224BM) located 24cm from the floor on the same wall as the port. Footshocks were delivered through a grid floor via a constant current aversive stimulator (ENV-414S). An overhead video camera and side video camera recorded the sessions for subsequent offline video scoring.

2.3 Behavioral Procedures

Reward pre-training (5 sessions): An auditory cue is paired with 10% sucrose solution delivery (100µl) and serves as the reward cue (25 trials; ITI, 90-130s).

Habituation (1 session): Rats continue to receive 25 reward cue-sucrose pairings (ITI, 90-130s) in addition to 5 unreinforced presentations each of the future fear and safety cues in order to habituate the rats to their presentation, thereby reducing any baseline freezing to these novel cues.

Discriminative conditioning (DC) (4 sessions): Reward cue-sucrose pairings continue (15 trials). Another auditory cue is paired with a mild 0.5mA, 0.5s footshock and serves as the fear cue (4 trials). In separate trials the 20s fear cue is presented at the same time as a 20s safety light cue resulting in no footshock ('fear+safety', 15 trials). Trials in which the safety cue is presented alone without any footshock are also included to assess whether freezing develops to the safety cue as well as providing the animal with additional trials that contains a safety cue-no shock contingency (10 trials). Trials are presented pseudorandomly (ITI, 100-140 s). Cues were

counterbalanced across subjects with the caveat that the fear+safety compound cue was

composed of one auditory cue and one light cue. Eight of the male rats and 12 of the female

rats underwent DC training in which the reward cue was a continuous auditory cue (3 kHz, 20s

cue; 70dB), the fear cue a pulsing auditory cue (11 kHz, 20s; 70dB), and the safety cue was the

presentation of two lights (28V, 100mA located on both sides of the port). The remaining eight

male rats and eight female rats underwent training in which the fear and safety cue stimuli were

counterbalanced: the light served as the fear cue and the pulsing auditory cue served as the

safety cue.

Extinction Training (1 session): One day after the last DC session, both the reward cue and fear

cue were presented 20 times each in a pseudorandomized order without sucrose or footshock.

Extinction Test (1 session): One day after extinction training, rats were presented with the

reward (10 trials), fear (10 trials), fear+safety (5 trials) and safety (5 trials) cues in a

pseudorandomized order (ITI, 60-120s). None of the cues were presented with sucrose or

footshock.

To exclude possible sex differences in pain sensitivity and foot shock perception a separate

group of male (n=8) and age-matched female (n=8) rats was presented with a series of

unsignalled foot shocks of increasing intensities (0.3 mA, 0.35 mA, 0.4 mA, 0.45 mA, 0.5 mA,

0.55 mA, 0.6 mA, 0.7 mA, 0.8 mA, 0.9, 1.0 mA) with an inter-stimulus interval of 2 min. The

session was flanked with 5 min intervals in which no stimulus occurred.

2.4 Data analyses

Our experimental groups to directly compare males and females on discrimination behavior

consisted of 16-20 rats. Cohorts of 4 or 8 female rats were trained alongside cohorts of 4 male

rats for a total of 4 replications. Fear behavior was assessed manually offline from videos by

measuring freezing, defined as complete immobility with the exception of respiratory

movements, which is an innate defensive behavior (Blanchard & Blanchard, 1969; Fendt &

Fanselow, 1999). The total time spent freezing during each 20s cue was quantified and

expressed as a percentage. Measuring the total time the animal spent inside the reward port

and at the entrance of the port with nose positioned at port entrance during each cue assessed

reward seeking behavior and was expressed as a percentage. Darting behavior was detected

and quantified offline from videos recorded from overhead cameras via a custom MatLab

program, with movements of a velocity of 23.5cm/s or faster qualifying as a single dart (Gruene

et al., 2015); these were also confirmed manually. Darting was expressed as the averaged # of

darts per cue (sum of darts/ # trials) or trial (sum of darts). Since there were different number of

trials per reward, fear, fear+safety and safety cue in each DC session and test for extinction,

this was expressed as the sum of darts across trials divided by the number of trials for each cue

(sum of darts/ # trials). During extinction training, data for each individual trial is shown and the

dart rate is expressed the averaged sum of darts for each individual trial (sum of darts). Three

individuals performed manual offline behavioral scoring. Pearson's correlations of behavioral

values between scorers were greater than $r = 0.80$. Behavioral data were analyzed with one-

way or two-way repeated measures ANOVAs, with sex as the independent factor and condition

as the repeated factor, followed by *post hoc* Sidak's, Tukey's or Dunnett's multiple comparisons

tests with GraphPad Prism 8. P values were adjusted for multiple comparisons.

For shock sensitivity testing, freezing duration in the 2-min intervals between shock

presentations was scored manually as an indicator of fear, as well as darting and jumping as an

immediate shock response. For the freezing durations, a two-way repeated measures ANOVA

was carried out via GraphPad Prism 7, with sex as the independent factor and shock intensity

as the repeated factor. Darting and jumping were assessed as dichotomous variables with

darting/no darting and jumping/no jumping, respectively. For both, a Cochran test was

performed.

3. Results

3.1 Female rats spent more time reward seeking during reward pre-training

All rats first underwent 5 reward pre-training sessions in which the reward cue was paired with

sucrose delivery. The percent time spent at or in the reward port during each reward cue across

each reward session was quantified (Figure 1b). Two-way repeated-measures ANOVAs showed

main effects of session ($F_{(4,136)}=5.395$, p=0.0005) and sex ($F_{(1,34)}=10.83$, p=.0023), but no

significant interaction ($F_{(4,136)}=0.9031$, p=0.4641). *Post hoc* Sidak's multiple comparisons test

showed females spent significantly more time reward seeking during the reward cue than males

for sessions R2 (p=0.0274), R3 (p=0.0151) and R5 (p=0.0041). The latency, in seconds, to

enter the port post-cue onset was also calculated for each reward cue presentation across all

sessions (Figure 1c). Two-way repeated-measures ANOVAs showed a main effect of sex

($F_{(1,34)}=20.37$, p<.0001), but no significant interaction ($F_{(4,136)}=1.684$, p=0.1571) or main

effect of session ($F_{(4,136)}=0.7755$, p=0.5429). *Post hoc* Sidak's multiple comparisons test

showed females were significantly faster to enter the port than males during the last 3 reward

sessions (R3, p=0.001; R4, p=0.0391; R5, p=0.0014). Taken together, female rats consistently

spent more time than males in the reward port during the reward cue in reward pre-training

sessions.

3.2 Female rats did not show conditioned inhibition of freezing

After reward pre-training, rats were then exposed to sessions also consisting of reward, fear and

safety cues. The reward cue and sucrose reward were the same as the reward pre-training

sessions. The fear cue was paired with a 0.5mA footshock, and both the safety cue and

fear+safety cue did not result in footshock or sucrose.

The percent time spent at or in the reward port during each cue across session was quantified

for each DC session (Figure 2b). Two-way repeated-measures ANOVAs showed a significant

cue by sex effect, as well as main effects of cue and sex for DC1 (Table 1). *Post hoc* Sidak's

multiple comparisons test showed that, during DC1, females spent significantly more time

reward seeking during the reward cue compared to males (p<0.001), consistent to what was

seen in reward pre-training. For the remaining DC2-4 sessions, a main effect of cue was

observed (Table 1) and *post hoc* Sidak's multiple comparisons test showed that both male and

female rats spent significantly more time reward seeking during the reward cue compared to all

other cues (p<0.0001), with no significant differences between the males and females. Thus,

the noticeable increase in reward seeking in the females, that was seen during reward pre-

training, dissipated by the 2^{nd} DC session.

The percent time freezing during each cue across session was quantified for each DC session

(Figure 2c). Two-way repeated-measures ANOVAs showed a significant cue by sex effect for

sessions DC2-4, as well as main effects of cue and sex for every session (Table 1). *Post hoc*

Sidak's multiple comparisons tests showed that, for every session, females displayed

significantly more freezing to the fear+safety cue compared to males (DC1, p=0.0313; DC2,

p=0.007; DC3, p=0.0007; DC4, p<0.0001). Females also showed significantly higher freezing

levels to the fear cue compared to males during DC2 (p=0.0111). Males showed a significant

reduction in freezing levels to the fear+safety cue compared to the fear cue during sessions

DC3 (p=0.0156) and DC4 (p<0.0001), thus showing significant conditioned inhibition of freezing.

Females did not show a significant inhibition of freezing during any session.

The number of darts during each cue was also quantified for each DC session and expressed

as a dart rate (Figure 2d; sum of darts/ # trials). Darting behavior during cue presentation was

largely absent until DC3 and DC4. Two-way repeated-measures ANOVAs showed a significant

cue by sex effect for DC4, as well as main effects of cue, for DC2-4, and sex, for DC1 and DC4

(Table 1). *Post hoc* Sidak's multiple comparisons test showed that, during DC4, females

expressed more darting behavior compared to males during both the fear cue (p=0.0017) and

the fear+safety cue (p=0.0186). The females reduced their darting rate from 0.51 during the fear

cue to 0.32 during the fear+safety cue, but this was not statistically significant.

3.3 Female rats did not show significant extinction of freezing

The day after the last DC session all rats underwent fear and reward extinction within the same

session. That is, both the fear and reward cues were presented within the same training

session, without footshocks or sucrose presentations.

During extinction of reward, there was no main effect of reward trial (F(19,646)=1.526,

p=0.0704) or sex (F(1,34)=1.31, p=0.2603) and no interaction (F(19,646)=0.8927, p=0.5924);

there was also no significant difference between male and female groups for any trial (Figure

3Bi). One day later when rats were re-tested for extinction memory (Figure 3Bii), there was a

main effect of cue (2-way RM ANOVA; F(3,102)=134.7, p<0.0001) and sex (F(1,34)=6.217,

p=0.0177). *Post hoc* Sidak's multiple comparisons test showed females had significantly more

port activity than males just during the safety cue (p=0.0452), although this difference did not

reflect a large increase in port activity as females spent 6.38% +/- 0.86 of the safety cue in the

port compared to 2.66% +/- 0.86 in males. Overall, there appeared to be no differences in the

ability of males and females to extinguish their reward seeking responses.

To assess fear extinction the averaged percent time freezing during each trial of fear extinction

training was calculated (Figure 3Ci). There was a main effect of fear trial (2-way RM ANOVA;

F(19, 646)=7.69, p<0.0001) and sex (2-way RM ANOVA; F(1, 34)=4.607, p=0.0391), but no

significant interaction (F(19, 646)=1.566, p=0.059). Compared to trial 1, males showed

significantly reduced freezing in extinction trials 8, 9 and 11-20 (*post hoc* Tukey's multiple

comparisons test, p<0.05), demonstrating good fear extinction beginning around the 8[th] trial. In

contrast, females only showed a significant reduction in freezing during trial 19 compared to the

first trial (*post hoc* Tukey's multiple comparisons test, p=0.0394), demonstrating relatively

absent fear extinction. One day later when rats were retested for extinction memory (Figure

3Cii), there was a main effect of cue (F(3,102)=134.7, p<0.0001) and sex (F(1, 34)=6.217,

p=0.0177), as well as a significant interaction of cue X sex (F(3, 102)=3.481, p=0.0187). *Post*

hoc Sidak's multiple comparisons test showed that females froze significantly more than males
to the fear (p=0.0146) and fear+safety (p=0.0091) cues. This indicates the continued absence of
any extinction of freezing in females.
284
In response to each fear cue presentation across extinction, we also assessed darting levels
(Figure 3Di). There was a main effect of sex ($F_{(1,34)}=4.816$, p=0.0351), but no effect of trial
($F_{(6.957, 236.6)}=0.6941$, p=0.6762) and no significant interaction ($F_{(19, 646)}=1.083$, p=0.3640).
Post hoc Sidak's multiple comparisons test showed no significant differences between males
and females for any trial. For the extinction memory test one day later (Figure 3Dii), there was a
significant cue X sex interaction ($F_{(3,102)}=4.447$, p=0.0056), as well as a main effect of both
cue ($F_{(2.013, 68.44)}=4.248$, p=0.018) and sex ($F_{1, 34}=4.834$, p=0.0348). Females showed a
significantly higher dart rate than males during the fear cue (*post hoc* Sidak's multiple
comparisons test, p<0.05), which was also significantly higher than the dart rate to the reward
and safety cues in the females (*post hoc* Sidak's multiple comparisons test, p<0.05). However,
even though statistically significant, the amount of darting during the fear cue in females was
very low (0.05-0.4 in extinction training and 0.065 in the extinction memory test), and therefore
no clear conclusions can be made regarding darting and extinction in this study.
298
3.4 Shock reactivity in males versus females
To exclude possible sex differences in pain sensitivity and footshock perception, a separate
cohort of 8 male and 8 age-matched female rats received 11 unsignaled footshocks of
increasing intensities (0.3 mA, 0.35 mA, 0.4 mA, 0.45 mA, 0.5 mA, 0.55 mA, 0.6 mA, 0.7 mA,
0.8 mA, 0.9, 1.0 mA) with an inter-stimulus interval of 2 min. Freezing increased as a function of
shock intensities (Figure 4A; 2-way RM ANOVA; $F_{(11,121)}=25.9$, p<0.0001). No main effects of
sex ($F_{(1,121)}=0.2871$, p=0.6027) or sex by shock ($F_{(11,121)}=1.413$, p=0.1754) were observed.
Our experiments utilized a shock intensity of 0.5mA throughout this study. For this particular
intensity, we also noted the number of rats that jumped or darted in response to a 0.5mA shock
(Figure 4B,C). No sex differences in the number of rats jumping in response to the 0.5mA
footshock were observed ($\chi2$: p>0.9). The number of female rats darting after the 0.5mA
footshock was higher than males, but not significantly ($\chi2$: p =0.0769), with five of the eight
female rats tested exhibiting the behavior. A higher number of females darting in response to
the footshock in this test would still not explain the lack of conditioned inhibition of freezing in
the females, as freezing levels at 0.5mA was slightly lower than the males (Figure 4A). Our

results do not definitively show, but do suggest, that females may be more likely to respond to a

footshock with a darting response.

4. Discussion

In this study, we show females exhibit a significantly different behavioral profile than males in a

task that tests for reward, fear and safety cue discrimination, as well as conditioned inhibition

and extinction. Female Long Evans rats showed more reward seeking early in training and

persistently high freezing levels to the fear cue when in the presence of a safety cue or after

fear extinction. Darting behavior in the females late in training showed conditioned inhibition of

this behavior in the presence of a safety cue, suggesting the females are able to discriminate

between the fear and safety cues but do not suppress their freezing response. This data adds to

the growing body of evidence of sex differences in fear regulation and highlights the advantages

of using more complex learning paradigms with additional behavioral measurements.

Even though studies including female subjects have been proportionally low, several studies

have reported clear sex differences in fear regulation. Most of these are consistent with our

findings of reduced discrimination between fear and safety signals. For instance, female mice

show more generalization of fear to novel and safe contexts compared to males, and with this

generalization there is a concurrent increase in basal amygdala activity (Keiser et al., 2017).

Male and female rats also respond differently to the controllability of a stressor. Males display

reduced fear during escapable stress versus inescapable stress whereas females exhibit no

beneficial effects of perceiving a stressor as escapable and controllable (Baratta et al., 2018).

The buffering effects seen in these males were linked to prelimbic cortical neurons projecting to

the dorsal raphe nucleus, which do not appear to be engaged in females. Females displaying a

similar fear response to both inescapable and escapable stress is similar to our findings of

females showing equivalent freezing levels to the fear cue in the presence or absence of a

safety cue, in that there were no buffering effects seen by the safety cue. It appears that

females do not downregulate their fear response in situations cued as safe.

Our data showing an increase in darting behavior in female rats as the number of fear cue-

footshock trials increase is consistent with another report using female rats in a fear conditioning

and extinction paradigm (Gruene et al., 2015). Like us, Gruene et al (2015) also show darting

levels increase as learning about the fear cue advances. Compared to us, Gruene et al (2015)

report notably higher darting frequencies, which is most likely due to the differences in shock

intensities and number of trials; our study used 4 trials of 0.5mA per day for 4 days compared to
their study using 7 trials of 0.7mA on one day. Our study also includes reinforced reward trials
within the same sessions as the fear cue-footshock trials, which could alter the contextual
expectations of the training session and reduce overall darting levels. It would be interesting in
future studies to identify what leads a female to become a 'darter' versus 'non-darter'. As darting
is a more active response compared to freezing, the circuits engaged during potential threats
would likely be different in these two populations.
355
Our findings showing a lack of conditioned inhibition of freezing in females appear to be
inconsistent with a recent study demonstrating a lack of sex differences in conditioned inhibition
of freezing (Foilb et al., 2018). This is likely due to differences in our respective protocols. First,
their footshock intensity was 1.2mA, resulting in freezing levels >90% during the fear cue. As
footshock intensity and number of trials are consistently inconsistent across studies, it would be
interesting to assess if freezing and darting levels in females follow a linear trend with increasing
training intensity, or if there is instead a possibly U-shaped relationship. Foilb et al (2018) also
used separate presentations of the fear cue and safety cue throughout training and employed
the fear+safety cue summation test during recall, whereas we include fear+safety trials as part
of the training. In contrast, another study has shown females discriminate equally to males early
in training but then generalize their fear response to the safety cue with continued training (Day
et al., 2016). While the females in our study clearly showed equivalent freezing levels to both
the fear and fear+safety cues at all time points throughout training, they did not increase their
freezing levels to the safety cue when presented alone. And, lastly, our paradigm, unlike others,
includes reinforced reward trials during the training of fear and safety cues, which would change
the context from a 'threat-no threat' situation to a 'threat-no threat-reward' situation, inducing
approach behaviors on top of defensive behaviors.
373
Altogether, the data paints a consistent picture of females showing heightened fear responses
to cues signaling safety, mimicking the clinical picture in women (Gamwell et al., 2015; Lonsdorf
et al., 2015). The presentation of a safety signal not only decreases fear, but also stimulates
opposing neuronal activity. Field potential recordings in the striatum during safety signal
presentation has shown that brain regions dealing with approach and reward become activated
(Rogan et al., 2005). These findings have also been translated to using safety signals to
overcome anhedonia in rats (Pollak et al., 2008), showing that safety signals may also be
regulating emotion in addition to conditioned behavior (Foilb & Christianson, 2018).

In our study, females consistently showed elevated reward seeking behavior during the reward

cue compared to males beginning in the second reward pre-training session. This data appears

consistent with reward studies showing significant sex differences in response to sucrose, with

females willing to work more for sucrose in a progressive ratio paradigm (Tapia et al., 2019),

and in response to drugs of abuse, with female rats consistently self-administering drugs more

rapidly than males (Becker & Koob, 2016). The increased reward seeking in females seen in our

study remained until the end of the first DC session at which point they were equivalent to the

males. Interestingly, DC1 is the first time the animals are exposed to footshock. Taking into

account the lack of conditioned inhibition of freezing in the females, the females may no longer

be as motivated to seek rewards in the face of adverse footshocks. This would be consistent

with the report that female rats sacrifice their metabolic needs in order to avoid shocks more

than males (Pellmanet al., 2017).

Numerous sex differences have been reported in the functioning of the stress neuropeptide,

corticotropin-releasing factor (CRF), with differences in receptor expression, distribution,

trafficking and signaling (reviewed in (Bangasser & Wiersielis, 2018)). The majority of these

differences lead to enhanced CRF efficacy in females which may lead to heightened sensitivity

to stressors in females. Recently, the gene for CRH receptor 1 (*CRHR1*) has been identified as

a possible candidate gene for mood and anxiety disorders. Weber et al. (2016) have shown that

carrying the *CRHR1* minor rs17689918 allele increases the risk for panic disorders in women.

Patients carrying this risk allele also demonstrate more generalization of fear to a safety cue,

increased amygdala activation during the safety cue and decreased frontal cortex activation with

discriminative fear conditioning. Thus, aberrant CRF signaling can lead to sustained fear under

conditions cued as safe and can be manifested by changes in neural activity in the amygdala

and frontal cortex.

Neural activity in the amygdala and prefrontal cortex has been shown by our lab to also play a

critical role in effective discriminative conditioning in male rats. We have previously identified

neurons in the basolateral amygdala (BLA) that discriminate among safety, fear and reward

cues in male rats (Sangha et al., 2013); our future experiments will test if females show the

same discriminative neurons. Using reversible pharmacological inactivations in male rats, we

have also demonstrated that the infralimbic prefrontal cortex (IL) is necessary for suppression of

conditioned fear during a safety cue and the prelimbic prefrontal cortex (PL) is necessary for

fear expression and discriminatory reward seeking (Sangha, Robinson, et al., 2014). These

results indicate that activating the IL in the females may improve conditioned inhibition to the

combined fear and safety cues. Our results with male rats also show that manipulating D1-

receptor mediated dopamine activity in the BLA disrupts suppression of conditioned fear (Ng et

al., 2018), implicating dopaminergic ventral tegmental area (VTA) neurons projecting to the BLA

in safety-fear-reward discrimination.

Our findings are consistent with human studies where females show less discrimination

between the fear and safety signals than males (Gamwell et al., 2015; Lonsdorf et al., 2015),

which may reflect underlying mechanisms of increased prevalence for anxiety and stress-

related disorders in women. For example, a deficiency in effective safety signal processing has

been linked to Post-traumatic Stress Disorder (Jovanovic et al., 2009, 2010), panic disorder

(Gorka et al., 2014), and anxiety (Lissek et al., 2005), all disorders with a higher incidence in

women than men (Mclean et al., 2011). In our paradigm, females show a significantly different

behavioral profile than males that is consistent with the clinical picture, thus making it a great

tool to test for the neurobiological mechanisms underlying these sex differences.

5. Acknowledgements:

We thank Yolanda Jonker, Signe Hobaugh, Jessica Kerns, Emma Speckelson, Emily Willis, and

Mackenzie McIntosh for excellent animal care. IM was supported by the Alexander von

Humboldt foundation. This work was supported by the National Institutes of Health [NIMH

R01MH110425 to SS].

6. References

Bangasser, D. A., & Wiersielis, K. R. (2018). Sex differences in stress responses : a critical role for corticotropin-releasing factor. *Hormones*, *17*(1), 5–13.

Baratta, M. V, Leslie, N. R., Fallon, I. P., Dolzani, S. D., Chun, L. E., Tamalunas, A. M., … Maier, S. F. (2018). Behavioural and neural sequelae of stressor exposure are not modulated by controllability in females. *European Journal of Neuroscience*, *47*, 959–967. http://doi.org/10.1111/ejn.13833

Becker, J. B., & Koob, G. F. (2016). Sex Differences in Animal Models : Focus on Addiction. *Pharmacological Reviews*, *68*, 242–263.

Blanchard, R. J., & Blanchard, D. C. (1969). Crouching as an index of fear. *Journal of Comparative and Physiological Psychology*, *67*(3), 370–5.

Christianson, J. P., Fernando, a. B. P., Kazama, a. M., Jovanovic, T., Ostroff, L. E., & Sangha, S. (2012). Inhibition of Fear by Learned Safety Signals: A Mini-Symposium Review. *Journal of Neuroscience*, *32*(41), 14118–14124. http://doi.org/10.1523/JNEUROSCI.3340-12.2012

Conway, S. M., Puttick, D., Russell, S., Potter, D., Roitman, M. F., & Charto, E. H. (2019). Neuropharmacology Females are less sensitive than males to the motivational- and dopamine- suppressing e ff ects of kappa opioid receptor activation, *146*(November 2018), 231–241. http://doi.org/10.1016/j.neuropharm.2018.12.002

Daviu, N., Andero, R., Armario, A., & Nadal, R. (2014). Sex differences in the behavioural and hypothalamic-pituitary-adrenal response to contextual fear conditioning in rats. *Hormones and Behavior*, *66*(5), 713–723. http://doi.org/10.1016/j.yhbeh.2014.09.015

Day, H. L. L., Reed, M. M., & Stevenson, C. W. (2016). Sex differences in discriminating between cues predicting threat and safety. *Neurobiology of Learning and Memory*, *133*, 196–203. http://doi.org/10.1016/j.nlm.2016.07.014

Fendt, M., & Fanselow, M. S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. *Neuroscience & Biobehavioral Reviews*, *23*(5), 743–760. http://doi.org/10.1016/S0149-7634(99)00016-0

Foilb, A. R., Bals, J., Sarlitto, M. C., & Christianson, J. P. (2018). Sex differences in fear discrimination do not manifest as differences in conditioned inhibition. *Learning and Memory*, *25*(1), 49–53. http://doi.org/10.1101/lm.045500.117

Foilb, A. R., & Christianson, J. P. (2018). Chapter 11 Brain Mechanisms for Learning and Using Safety Signals. In S. Sangha & D. Foti (Eds.), *Neurobiology of Abnormal Emotion and Motivated Behaviors: Integrating Animal and Human Research* (1st editio, pp. 205–224). Cambridge, Massachusetts: Academic Press.

Gamwell, K., Nylocks, M., Cross, D., Bradley, B., Norrholm, S. D., & Jovanovic, T. (2015). Fear
Conditioned Responses and PTSD Symptoms in Children : Sex Differences in Fear-
Related Symptoms. *Developmental Psychobiology*, *57*, 799–808.
http://doi.org/10.1002/dev.21313

Glover, E. M., Jovanovic, T., & Norrholm, S. D. (2015). Estrogen and extinction of fear
memories: Implications for posttraumatic stress disorder treatment. *Biological Psychiatry*,
78(3), 178–185. http://doi.org/10.1016/j.biopsych.2015.02.007

Gorka, S. M., Lieberman, L., Nelson, B. D., Sarapas, C., & Shankman, S. A. (2014). Aversive
responding to safety signals in panic disorder: The moderating role of intolerance of
uncertainty. *Journal of Anxiety Disorders*, *28*(7), 731–736.
http://doi.org/10.1016/j.janxdis.2014.06.004

Grant, B. F., Saha, T. D., June Ruan, W., Goldstein, R. B., Patricia Chou, S., Jung, J., … Hasin,
D. S. (2016). Epidemiology of DSM-5 drug use disorder results from the national
epidemiologic survey on alcohol and related conditions-III. *JAMA Psychiatry*, *73*(1), 39–47.
http://doi.org/10.1001/jamapsychiatry.2015.2132

Gruene, T. M., Flick, K., Stefano, A., Shea, S. D., & Shansky, R. M. (2015). Sexually divergent
expression of active and passive conditioned fear responses in rats. *ELife*, *4*, 1–9.
http://doi.org/10.7554/eLife.11352

Hernandez-Avila, C. A., Rounsaville, B. J., & Kranzler, H. R. (2004). Opioid-, cannabis- and
alcohol-dependent women show more rapid progression to substance abuse treatment.
Drug and Alcohol Dependence, *74*(3), 265–272.
http://doi.org/10.1016/j.drugalcdep.2004.02.001

Jovanovic, T., Kazama, A., Bachevalier, J., & Davis, M. (2012). Impaired safety signal learning
may be a biomarker of PTSD. *Neuropharmacology*, *62*(2), 695–704.
http://doi.org/10.1016/j.neuropharm.2011.02.023

Jovanovic, T., Norrholm, S. D., Blanding, N. Q., Davis, M., Duncan, E., Bradley, B., & Ressler,
K. J. (2010). Impaired fear inhibition is a biomarker of PTSD but not depression.
Depression and Anxiety, *27*(3), 244–251. http://doi.org/10.1002/da.20663

Jovanovic, T., Norrholm, S. D., Fennell, J. E., Keyes, M., Fiallos, A. M., Myers, K. M., …
Duncan, E. J. (2009). Posttraumatic stress disorder may be associated with impaired fear
inhibition: Relation to symptom severity. *Psychiatry Research*, *167*(1–2), 151–160.
http://doi.org/10.1016/j.psychres.2007.12.014

Keiser, A. A., Turnbull, L. M., Darian, M. A., Feldman, D. E., Song, I., & Tronson, N. C. (2017).
Sex Differences in Context Fear Generalization and Recruitment of Hippocampus and

Amygdala during Retrieval. *Neuropsychopharmacology*, *42*(2), 397–407.

http://doi.org/10.1038/npp.2016.174

Keyes, K. M., Grant, B. F., & Hasin, D. S. (2008). Evidence for a closing gender gap in alcohol

use, abuse, and dependence in the United States population. *Drug and Alcohol*

Dependence, *93*(1–2), 21–29. http://doi.org/10.1016/j.drugalcdep.2007.08.017

Leknes, S., Lee, M., Berna, C., Andersson, J., & Tracey, I. (2011). Relief as a reward: Hedonic

and neural responses to safety from pain. *PLoS ONE*, *6*(4), e17870.

http://doi.org/10.1371/journal.pone.0017870

Lissek, S., Powers, A. S., McClure, E. B., Phelps, E. A., Woldehawariat, G., Grillon, C., & Pine,

D. S. (2005). Classical fear conditioning in the anxiety disorders: A meta-analysis.

Behaviour Research and Therapy, *43*(11), 1391–1424.

http://doi.org/10.1016/j.brat.2004.10.007

Lonsdorf, T. . B., Haaker, J., Schümann, D., Sommer, T., Brassen, S., Bunzek, N., … Kalisch,

R. (2015). Sex differences in CS-discrimination during context-dependent fear learning and

its retrieval - the role of biological sex, contraceptives and menstrual cycle phases. *Journal*

of Psychiatry and Neuroscience, *40*(6), 368–375. http://doi.org/10.1503/jpn.140336

Mclean, C., Asnaani, A., Litz, B., & Hofmann, S. (2011). Gender differences in anxiety disorders:

Prevalence, course of illness, comorbidity and burden of illness. *Journal of Psychiatric*

Research, *45*(8), 1027–1035. http://doi.org/10.1016/j.jpsychires.2011.03.006.Gender

Ng, K. H., Pollock, M. W., Urbanczyk, P. J., & Sangha, S. (2018). Altering D1 receptor activity in

the basolateral amygdala impairs fear suppression during a safety cue. *Neurobiology of*

Learning and Memory, *147*, 26–34. http://doi.org/10.1016/j.nlm.2017.11.011

Pellman, B. A., Schuessler, B. P., Tellakat, M., & Kim, J. J. (2017). Sexually Dimorphic Risk

Mitigation Strategies in Rats. *Eneuro*, *4*(1), e.0288-16.2017.

http://doi.org/10.1523/ENEURO.0288-16.2017

Pollak, D. D., Monje, F. J., Zuckerman, L., Denny, C. a., Drew, M. R., & Kandel, E. R. (2008).

An Animal Model of a Behavioral Intervention for Depression. *Neuron*, *60*(1), 149–161.

http://doi.org/10.1016/j.neuron.2008.07.041

Rescorla, R. A. (1969). Conditioned inhibition of fear resulting from negative CS-US

contingencies. *Journal of Comparative and Physiological Psychology*, *67*(4), 504–509.

http://doi.org/10.1037/h0027313

Rogan, M. T., Leon, K. S., Perez, D. L., & Kandel, E. R. (2005). Distinct neural signatures for

safety and danger in the amygdala and striatum of the mouse. *Neuron*, *46*(2), 309–20.

http://doi.org/10.1016/j.neuron.2005.02.017

Sangha, S., Chadick, J. Z., & Janak, P. H. (2013). Safety encoding in the basal amygdala. *The*
Journal of Neuroscience, *33*(9), 3744–51. http://doi.org/10.1523/JNEUROSCI.3302-
12.2013

Sangha, S., Greba, Q., Robinson, P. D., Ballendine, S. A., & Howland, J. G. (2014). Heightened
fear in response to a safety cue and extinguished fear cue in a rat model of maternal
immune activation. *Frontiers in Behavioral Neuroscience*, *8*, 168.
http://doi.org/10.3389/fnbeh.2014.00168

Sangha, S., Robinson, P. D., Greba, Q., Davies, D. A., & Howland, J. G. (2014). Alterations in
reward, fear and safety cue discrimination after inactivation of the rat prelimbic and
infralimbic cortices. *Neuropsychopharmacology*, *39*(10), 2405–13.
http://doi.org/10.1038/npp.2014.89

Tanimoto, H., Heisenberg, M., & Gerber, B. (2004). Event timing turns punishment to reward.
Nature, *430*, 983. http://doi.org/10.1038/nature02816

Tapia, M. A., Lee, J. R., Weise, V. N., Tamasi, A. M., & Will, M. J. (2019). Sex di ff erences in
hedonic and homeostatic aspects of palatable food motivation. *Behavioural Brain*
Research, *359*(November 2018), 396–400. http://doi.org/10.1016/j.bbr.2018.11.023

Walasek, G., Wesierska, M., & Zieliński, K. (1995). Conditioning of fear and conditioning of
safety in rats. *Acta Neurobiologiae Experimentalis*, *55*(2), 121–132.

Weber, H., Richter, J., Straube, B., Lueken, U., Domschke, K., Schartner, C., … Reif, A. (2016).
Allelic variation in CRHR1 predisposes to panic disorder: Evidence for biased fear
processing. *Molecular Psychiatry*, *21*(6), 813–822. http://doi.org/10.1038/mp.2015.125

 7. Figure Legends

Figure 1. **Females show increased reward seeking in response to the reward cue**. **A)**

Schematic depicting experimental outline. During reward pre-training, rats (16 males, 20

females) received 25 cue-sucrose pairings across 5 separate sessions. **B)** Averaged percent

time spent in the reward port during the five reward pre-training sessions (R1-5). Females spent

significantly more time in the port compared to males during R2, R3 and R5. **C)** Averaged

latency to enter the port after cue onset (in seconds). Females entered the port significantly

sooner than males during R3-5. Means +/- SEM. *p<0.05, **p<0.01.

Figure 2. Females do not show inhibition of conditioned freezing in the presence of the

safety cue. A) Schematic depicting experimental outline. During the 4 DC sessions, rats (16

males, 20 females) were presented with four types of cued trials: reward cue-sucrose, fear cue-

shock, fear+safety cue with no footshock and the safety cue presented alone without footshock.

B) Averaged percent time spent in the port during each cue across the 4 DC sessions. Both

males and females showed significantly higher reward seeking during the reward cue compared

to all other cues during every DC session. During DC1, females showed significantly higher

reward seeking to the reward cue compared to males. **C)** Averaged percent time spent freezing

during each cue across the 4 DC sessions. During DC3 and DC4, males showed significantly

lower freezing to the fear+safety cue (and reward and safety cues) when compared to the fear

cue. Females did not show significant inhibition of conditioned freezing to the fear+safety cue

compared to the fear cue during any DC session. Females also showed significantly higher

freezing to the fear+safety cue compared to males during every session. **D)** Darting behavior

during each cue across the 4 DC sessions. During DC4 females showed significantly more darts

than males during the fear and fear+safety cues. Means +/- SEM. # p<0.05, ####p<0.0001

within sex, between cue comparison; * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 within cue,

between sex comparison.

Figure 3. Females do not show significant extinction of fear. A) Schematic depicting

experimental outline. During extinction training both the reward and fear cues are presented in

the same session without sucrose or footshock. During the test for extinction memory 1 day

later all cues are presented without sucrose or footshock. **Bi)** Averaged percent time spent in

the port during each reward cue presentation during extinction training. No significant

differences were found between males and females during extinction training. **Bii)** Averaged

percent time spent in the port during each cue 1 day after extinction training. Females spent

significantly more time in the port than males during the safety cue. **Ci)** Averaged percent time

spent freezing during each fear cue presentation during extinction training. Compared to the first

trial of extinction, males showed significantly reduced freezing during trials 8, 9 and 11-20.

Freezing levels for females did not significantly decrease at any point in extinction training, with

the exception of trial 19. #$p<0.05$, compared to trial 1. **Cii)** Averaged percent time spent freezing

during each cue 1 day after extinction training. Males showed evidence of fear cue extinction

retention. Females froze significantly more than males during the fear and fear+safety cues. **Di)**

Averaged darting during each fear cue presentation during extinction training. No significant

post hoc differences found between males and females during extinction training. **Dii)** Averaged

darting during each cue 1 day after extinction training. Females had a significantly higher dart

rate than males during the fear cue, which was also significantly higher than the reward and

safety cues in females. Means +/- SEM. #$p<0.05$, ####$p<0.0001$ within sex, between cue/trial

comparisons. *$p<0.05$, **$p<001$, ****$p<0.0001$ within cue, between sex comparisons.

Figure 4*.* No significant differences in shock reactivity between age-matched male and

female rats. A) Male and female rats (n=8 each) were subjected to increasing footshock

intensities from 0.3mA to 1.0mA. No significant differences in freezing levels (means +/- SEM)

were detected between males and females after each shock presentation. The box around the

data at 0.5mA indicates the intensity used for the experiments in this study. There were no

significant differences in the number of males or females who jumped **(B)** or darted **(C)** in

response to the 0.5mA shock.

Reward seeking			
Session	*Cue x Sex effects*	*Main effect of cue*	*Main effect of sex*
DC1	$F(3,102) = 3.472$, **p=0.0189**	$F(3,102) = 95.16$, **p<0.0001**	$F(1,34) = 9.827$, **p=0.0035**
DC2	$F(3,102) = 0.7742$, p=0.5110	$F(3,102) = 227.9$, **p<0.0001**	$F(1,34) = 4.69$, **p=0.0374**
DC3*	$F(3,90) = 0.6512$, p=0.5843	$F(3,90) = 117$, **p<0.0001**	$F(1,30)=1.041$, p=0.3157
DC4	$F(3,102) = 2.255$, p=0.0864	$F(3,102) = 181.2$, **p<0.0001**	$F(1,34) = 2.453$, p=0.1266
Freezing			
Session	*Cue x Sex effects*	*Main effect of cue*	*Main effect of sex*
DC1	$F(3,102) = 2.245$, p=0.0876	$F(3,102) = 31.82$, **p<0.0001**	$F(1,34) = 5.045$, **p=0.0313**
DC2	$F(3,102) = 4.075$, **p=0.0089**	$F(3,102) = 103.4$, **p<0.0001**	$F(1,34) = 6.621$, **p=0.0146**
DC3*	$F(3,90) = 2.9$, **p=0.0393**	$F(3,90) = 151.3$, **p<0.0001**	$F(1,30)=9.719$, **p=0.0040**
DC4	$F(3,102) = 4.889$, **p=0.0032**	$F(3,102) = 198.9$, **p<0.0001**	$F(1,34) = 8.294$, **p=0.0068**
Darting			
Session	*Cue x Sex effects*	*Main effect of cue*	*Main effect of sex*
DC1	$F(3,102) = 1.98$, p=0.1216	$F(2.022,68.73) = 2.388$, p=0.0989	$F(1,34) = 4.146$, **p=0.0496**
DC2	$F(3,102) = 1.134$, p=0.3390	$F(1.923, 65.38) = 9.377$, **p=0.0003**	$F(1,34) = 3.667$, p=0.0640
DC3	$F(3,102) = 0.9158$ p=0.4361	$F(1.586,53.93) = 18.96$, **p<0.0001**	$F(1,34)=0.9579$, p=0.3346
DC4	$F(3,102) = 10.65$, **p<0.0001**	$F(1.684, 57.26) = 15.65$, **p<0.0001**	$F(1,34) = 13.34$, **p=0.0009**

*video files for 4 females were corrupted for this session (n=16 females, 16 males)

Table 1. Summary of two-way repeated-measures ANOVA analyses for reward seeking, freezing and darting behaviors during the four discriminative conditioning (DC) sessions.

A

Reward Pre-Training	Habituation	Discriminative Conditioning (DC)	Extinction Training	Extinction Test
5 sessions	*1 session*	*4 sessions*	*1 session*	*1 session*
reward cue→ sucrose	reward cue→ sucrose fear cue→ ⊘ safety cue → ⊘	reward cue→ sucrose fear cue→ shock fear+safety cue→⊘ safety cue→ ⊘	reward cue→⊘ fear cue→ ⊘	reward cue→⊘ fear cue→ ⊘ fear+safety cue→⊘ safety cue→ ⊘

B

C