Complex genetic and epigenetic regulation deviates gene expression from a unifying global transcriptional program

Mónica Chagoyen[1] and Juan F Poyatos[2,3]

[1] Computational Systems Biology Group (CNB-CSIC), Darwin 3, 28049, Madrid, Spain
[2] Logic of Genomic Systems Laboratory (CNB-CSIC), Darwin 3, 28049, Madrid, Spain
[3] Center for Genomics and Systems Biology, Department of Biology, New York University, New York, New York, United States of America

Email: *monica.chagoyen@cnb.csic.es* (MC), *jpoyatos@cnb.csic.es* (JFP)

Abstract

Environmental or genetic perturbations lead to gene expression changes. While most analyses of these changes emphasize the presence of qualitative differences on just a few genes, we now know that changes are widespread. This large-scale variation has been linked to the exclusive influence of a global transcriptional program determined by the new physiological state of the cell. However, given the sophistication of eukaryotic regulation, we expect to have a complex architecture of specific control affecting this program. Here, we examine this architecture. Using data of *Saccharomyces cerevisiae* expression in different nutrient conditions, we first propose a five-sector genome partition, which integrates earlier models of resource allocation, as a framework to examine the deviations from the global control. In this scheme, we recognize invariant genes, whose regulation is dominated by physiology, specific genes, which substantially depart from it, and two additional classes that contain the frequently assumed growth-dependent genes. Whereas the invariant class shows a considerable absence of specific regulation, the rest is enriched by regulation at the level of transcription factors (TFs) and epigenetic modulators. We nevertheless find markedly different strategies in how these classes deviate. On the one hand, there are TFs that act in a unique way between partition constituents, and on the other, the action of chromatin modifiers is significantly diverse. The balance between regulatory strategies ultimately modulates the action of the general transcription machinery and therefore limits the possibility of establishing a unifying program of expression change at a genomic scale.

Introduction

The limited availability of the components of the cell expression machinery, for example, free RNA polymerases, cofactors, ribosomes, etc., creates a resource allocation problem that affects their activities. This differential allocation eventually represents a global program of regulation (1)(2)(3)(4), with some genes expressed at the cost of others. As the dosage of these components can be modulated by the growth rate at balanced growth, the influence of the global program has typically been studied by quantifying how the expression of genes responds to growth.

This led to the identification of laws that predict the expression of different cellular elements (3)(4)(5). Two broad models have been considered. In a first model (model 1, Fig. 1A, top), the impact of the global program is recognized by partitioning the genome in three minimal sectors (4)(5). One of them contains ribosomal genes whose expression increases with growth rate verifying its role in driving cell growth (6). Two other sectors comprise genes whose expression decreases or remains invariant with growth. The three-sector model describes therefore fundamental aspects of cellular economics (7) already advanced in the early work of bacterial physiologists (8), and acts as a basic constituent to investigate many subjects. For instance, modifications to this framework were advanced to explain the cost of unnecessary gene expression (4), the dependence of cellular composition with antibiotics (9), or the rationale behind some seemingly wasteful carbon utilization strategies (10).

In a second model, the emphasis is not so much in the trade-offs between ribosomal genes and the rest, but between genes that follow a common pattern of expression and a minimal subset that diverges (model 2 containing two sectors: nonspecific and specific, Fig. 1A, bottom) (11). Here, the common pattern of expression is explained by a single scaling factor, which incorporates the resources of the global program that are

not involved in the activation of the specific genes. This implies that most changes in

expression result from a passive rather than active regulation what produces a unifying

pattern that "simplifies" the expected complexities of genome-wide expression

changes.

In this manuscript, we integrate both models (model 3, Fig. 1B). The new framework

discriminates a fine-structure within the nonspecific sector of model 2 that corresponds

to the three-sector partition representative of model 1. While the integration of both

views is significant *per se*, models 1 and 2 might appear unrelated, we also show that

this scheme is fundamental to help us expose how specific regulatory strategies

modulate the impact of the global transcriptional program at the genome-wide level.

To this aim, we first reexamined the original data that led to model 2, i.e., promoter

activity (PA) measurements of *Saccharomyces cerevisiae*'s genes obtained in different

growing conditions (11), to present our new scheme. However, this data only includes

a subgroup of ~900 yeast genes, and thus the quantification of the resource allocation

can only be approximated. To obtain a more precise classification we considered

genome-wide expression data of yeasts growing in chemostats (12) and developed a

method to define the new model at a large scale. We validate our partition through

functional analysis of the corresponding components.

Armed with this new framework, we focused on understanding how the global program

becomes modulated by specific regulatory mechanisms, a question which is only

beginning to be addressed. Indeed, within the framework of model 1 results are mixed.

For instance, several recent reports in bacterial systems demonstrated the prevalence

of the global expression program, while they have lowered the importance of

transcription factors (TFs) controlling the assumed deviations from it. TFs seem only to

fine-tune the action of the global regulation (13)(14) in combination with a few

metabolites (15). More recent work in yeast argued, in contrast, about the relevance of

epigenetic factors (promoter nucleosomal stability) as modulators of the global program

(16).

The function of the specific regulation might appear more straightforward *a priori* in the

scheme of model 2. Specific genes are indeed enriched by specific regulation,

generally coupled to the particular nutrient condition, whereas the behavior of those

genes within the nonspecific sector does not seem connected to a particular

transcription regulation strategy (11). We thus examine within our new framework to

what extent we could observe differential genetic and epigenetic regulation acting on

the genes constituting each sector. We discovered well-defined regulatory patterns.

More broadly, our results put forward an integrated view of the resource allocation

connected to genome-wide expression and emphasize how the global program is

eventually modified by the specific regulation. Active strategies of control are certainly

at work in genes within the fine-structure of the "nonspecific" response and can

eventually associate the metabolic status of the cell with gene expression. Overall, this

complex hierarchy of regulation in eukaryotes enhances the adjustment of genome-

wide expression patterns beyond the one that could be achieved through a passive

unifying program of global transcriptional control.

Results

Integration of two preceding resource allocation models. We first examined how

PA changes with the growth rate for a subset of *Saccharomyces cerevisiae* genes.

Keren *et al*. (11) presented a binary partition to describe these changes, recognizing a

common response in the absolute PA values of most genes and a specific one in a

much smaller subgroup. To focus on resource reallocation, we studied here fractional

PA activities instead of absolute values, i.e., the fraction of PA of each gene in a given

nutrient condition out of the summed activity of all genes in the set (3), and quantified

their change for each pair of conditions (from low to high growth rate).

Figure 1C shows a descriptive case (glycerol to glucose conditions). The most extreme

deviations of the global program correspond to the specific genes (11). Note also that

the stronger allocation of "expression resources" to these genes the fewer resources to

biosynthesis (reduction of the nonspecific sector) affecting growth rate (Fig. 1A,

bottom). We revised the two-sector partition by further separating specific genes as

specifically activated genes (fractional PA becomes much larger between conditions),

and specifically repressed genes (fractional PA becomes much smaller), and also

delimiting three components within the nonspecific sector: genes whose fractional PA

remains approximately invariant (diagonal in Fig. 1C), positive genes, whose fractional

PA moderately increases between conditions, and negative genes (fractional PA

decreases to only a limited extent, see Methods for details). These three sectors relate

model 2 to model 1.

We validated this new partition in two ways. First, that a set of genes follows a precise

proportional response between conditions was suggested by (11) as an evidence that

they share a common functionality (e.g., being part of the same pathway, etc.). Here,

we find that invariant, positive and negative genes also follow a precise proportional

response (Fig.S1) suggesting that our analysis identifies a biologically relevant fine-

grained structure. Second, we similarly expect that this fine-grained structure would

parallel that proposed in model 1 (4) in terms of functional categories. This is the case.

The invariant class is enriched by transcription regulation and ribosomal proteins; the

latter being more extensively observed in the positive class. Indeed, positive genes are

enriched by ribosomal genes (~65% of genes code for small or large subunits of the

ribosome), while negative genes are enriched in ATP metabolic processes, e.g.,

oxidative phosphorylation or the TCA cycle (Table S1, Methods).

The five-sector resource allocation model on a genomic scale. To substantiate the

previous five-sector model based on ~900 genes, we examined a genome-wide DNA

microarray dataset of yeast cells exhibiting the same range of growth rates for six

limiting nutrients (12). We studied again changes of relative expression (Methods) and

applied singular value decomposition (SVD) to each nutrient separately. The first and

second SVD components (Fig. 2A) explain >90% of the variance in each condition (the

components exhibited an analogous trend in all nutrients, Fig. S2). As a result, the

fractional expression response to growth of each gene can be approximated by the

linear combination of these two components (Fig. 2B).

Furthermore, we can interpret the first element of the linear combination ($\vec{v}_1$) as the

baseline fractional expression of the gene, which does not change with growth rate,

and the second element ($\vec{v}_2$) as its monotonic response to growth (Fig. 2B). This

reading enables us to generalize the previous partition framework obtained with PA

data. More specifically, a gene that maintains the same baseline expression (similar

loading of $\vec{v}_1$ that we denoted a_i) between two conditions involves a nonspecific

response. When this response in observed in >8 pairwise changes then we reason that

the gene is nonspecific (recall that there are 6 different nutrients and consequently a

total of 15 pairwise nutrient changes). Genes are considered specific otherwise (Fig.

2C).

Beyond the classification above, the second component ($\vec{v}_2$) provides a quantitative

score (the second loading, b_i) to classify genes as invariant, positive or negative (b_i~0,

b_i>0, b_i<0, respectively; Fig. 2C. Methods). We labelled as invariant those nonspecific

genes which exhibit this behavior in >3 nutrient conditions (total of 6). Nonspecific and

not invariant genes appearing more times as positive than as negative are categorized

as positive, and likewise for negative. Finally, specific genes which appear more times

as positive than as negative (again in all 6 conditions) are categorized as activated,

and analogously for repressed (Fig. 2C).

Finally, the functional analysis of genes within each sector agrees with that obtained

with the PA data and previous reports, what substantiates the biological significance of

the partition (Table S2; see also Table S1 showing how this classification maps onto

the subset of genes with PA data). In this way, we have generalized at a large scale

the new resource allocation model, a genome-wide classification that we use

throughout the next sections.

TFs are typically nonspecific genes. We inspect next the influence of the most direct

elements related to specific regulation, i.e., TFs. But before examining their impact on

the activity of the global transcription program, we asked first how TFs themselves are

framed in the previous partition.

We observed that most constituent TFs of the transcriptional regulatory network (122 of

a total of 133 composing the network, Methods) are nonspecific genes, i.e., they

present similar basal fractional expression (a_i loadings) across all pairwise condition

changes. Within this set, 31% presents a dominant invariant response ($b_i \sim 0$ in >3

nutrient conditions, of a total of 6), with five genes acting as invariant in all six

conditions (*rsc1*, *mbp1*, *pho2*, *rgr1*, and *swi6*). Two of these (*mbp1*, *swi6*) are at the top

of the network hierarchy (being involved in the mitotic cell cycle), and two are elements

of relevant complexes that interact with RNA polymerase II (*rsc1* of the RSC chromatin

complex, and *rgr1*/*med14* of the mediator complex); they can be considered as

elements of a general transcriptional machinery, for which maintaining its concentration

invariant across conditions could be essential. Moreover, 32% of nonspecific TFs are

dominantly negative, and only 4% positive. Of note, some of the TFs whose expression

decreases with growth ($b_i < 0$) are positive regulators of transcription in response to

stress (e.g., *bur6*, *gcn4*, *rpn4*) what justifies their overexpression at low growth rates.

Genes within each partition sector present different degree of transcriptional

regulation. Is the extent of regulation of target genes dependent on which sector they

belong to? We computed the mean number of regulators acting on genes within each

sector. Nonspecific genes are less regulated, on average, by TFs than specific ones as

expected [by 3.09 TFs *vs.* 5.06 TFs, p = $1.20\ 10^{-4}$, two-sample Kolmogorov-Smirnov

(KS) test]. Within nonspecific genes, invariant genes are less regulated than

nonspecific and *not* invariant ones (by 2.56 TFs *vs.* 3.3 TFs, p = $8.16\ 10^{-13}$, two-sample

KS test). Finally, nonspecific and positive genes are slightly more regulated than

nonspecific and negative genes (by 3.33 TFs *vs.* 3. 27 TFs, p = 0.0018, two-sample KS

test). Overall, specific genes are subjected to more regulation (larger number of TFs)

(Fig. 3A), while nonspecific and invariant ones show the least.

TFs and their regulated genes frequently exhibit the same response to growth.

Although Fig. 3A shows how the number of transcriptional interactions is reflected

differentially in the sectors of the partition it does not assure us whether these

interactions are functioning, e.g., regulation by TFs have been shown to be "inactive" in

bacteria (13). For this, we examined several features.

We initially inspected if target genes presenting a particular growth response are

enriched by TFs showing the very same response, as the similarity of the responses

could indicate that these TFs do influence the expression of the target. We thus

computed –for each target gene– the fraction of its regulators that behave as negative,

invariant, or positive with the growth rate in a given condition (TF_{neg}, TF_{inv}, TF_{pos},

respectively). Figure 3B shows the mean of these fractions for target genes which

themselves show a negative, invariant, or positive response, respectively. Negative

TFs are more likely to be found acting on target genes that are also negative (higher

mean TF_{neg} on negative genes), while invariant (TF_{inv}) and positive (TF_{pos}) TFs regulate

more often invariant and positive target genes, respectively (the latter signal is weaker

and depends on the particular condition, Fig. S3). Thus, TFs that exhibit the same

behavior as their cognate target gene tend to appear, on average, dominant on its

regulation; this indicates that part of the regulatory structure is functional.

Specific TFs regulate genes fitting to each partition sector. To further estimate the

active regulatory character of TFs, we measured the correlation of the response to

growth rate between any particular gene and all its cognate TFs ("regulatory

coherence", Methods). Specific genes showed stronger regulatory coherence than

nonspecific ones (Fig. S4A), and remain coherent in more nutrient conditions (Fig.

S4B), both results implying an existing contribution of TFs to deviate gene expression

from the global program.

Moreover, Fig. 3C shows those TFs whose operation is particularly coherent per

partition sector (Methods). One can identify here that TFs controlling specific genes are

normally not acting on nonspecific ones [this is supported by earlier reports[17]].

Moreover, some of the TFs that work coherently on positive genes (gray color, Fig. 3C)

are particularly absent in negative ones (pink color, Fig. 3C) and *vice versa*. In addition,

those that control nonspecific genes are higher up in the network hierarchy (Methods).

We also noted that some of these (significantly coherent) TFs are involved in chromatin

remodeling (Cyc8, Ume6, Spt6, Msn4, Abf1, Msn2, Nhp6A, acting on nonspecific

ones), or chromatin organization (Spt3, Spt2, Pho4, FKh2, Sin3, Spt20, Wtm2, Wtm1,

Hif1, acting on specific genes). We examine epigenetic aspects next.

The resource allocation model also reveals distinctive epigenetic regulation. To

inspect the function of epigenetic control mechanisms, we first quantified the proportion

of general transcription factors (GTFs) found within the set of TFs acting on a given

gene (18). GTFs (Rap1, Abf1, Reb1, Cbf1, and Mcm1) usually have little intrinsic

regulatory activity and together with the presence of chromatin remodelers (in

particular, RSC –Remodels Structure of Chromatin) control an alleged general

machinery of expression. We observed that GTFs constitute a larger and significant

fraction in the regulation of positive genes, while the opposite is observed for negative

ones (Fig. 4A).

GTFs are also associated with particularly fragile nucleosome promoter architectures

(19), a connection recently examined (16). Using this data, we computed the

nucleosome landscape for the different gene classes (Methods). Promoters of positive

genes are indeed enriched in fragile nucleosomes (Fig 4.B) while both negative and

invariant genes lack these structures. This indicates that positive genes are more

sensitive to adjust the global program of expression by means of chromatin modulation.

Enrichment of other promoter features contribute to this observation (Fig 4.C), like the

absence of TATA boxes, the action of TFIID over SAGA [but this precise grouping has

been recently reexamined (20)], the presence of nucleosomal free regions closer to the

transcriptional starting site (shNFRs) (partially associated to the previous score of

fragile nucleosomes), and the dominant effect of *trans* variability (Fig. S5) (21)**.**

Chromatin modifiers further modulate the global program. Finally**,** we examined

the effects of mutating different types of *trans*-acting chromatin regulators on the genes

constituting the sectors using a previously assembled compendium (22) (Methods). We

considered first the magnitude of change of expression (i.e., the *absolute* values of

expression) before/after the mutation of several types of modifiers. With the exception

of histone acetyltransferases (HATs) and TATA-binding protein related factors (TAFs),

the influence of most chromatin modifiers is dominant in specific genes as compared to

nonspecific ones (Fig. S6, Methods).

Within nonspecific genes, we also quantified the type of expression change

(experienced after mutation of the modifiers) and found three broad relationships (Fig.

5): 1) Epigenetic regulators acting as part of a general machinery (HATs –including

SAGA–, TAFs and methyltransferases) whose mutation causes a general decrease in

expression, very particularly in invariant and positive classes. Indeed, work by (20) and

(23) demonstrated that SAGA and TFIID are recruited to pol II promoters genome-wide

and that each complex is generally required for pol II transcription, i.e., its mutation

would lead to a genome-wide decrease of gene expression. 2) Regulators (e.g.,

histones, etc.) acting in a dual manner: increasing the expression of negative genes

after mutation (remodeler as a repressor) or reducing their expression in positives

(remodeler as an activator). This underlines the enrichment of negative and positive

classes by stress and ribosomal genes, respectively, which are largely regulated in an

opposite manner (24); a dual role of remodelers as activators and repressors have

been previously reported (25)(26). And 3) regulators as broad repressors (mutation

increases expression), e.g., those that represent regulation by gene silencing.

310   **Discussion**

Recent studies emphasized how most genome-wide expression changes might be the

consequence of a unifying global program of transcription control. This program is

determined by the availability of the general components of the expression machinery

and inevitably refers to a problem of resource reallocation, in which different parts of

the genome are to use the limited expression resources.

We followed here this interpretation to present a new partition of the yeast genome

through the integration of two earlier models, not connected until now (6)(11). Essential

to the framework is the faculty to discriminate what it was contemplated before as a

common nonspecific proportional response –subjected to the same global

regulation(11)– into three subclasses: invariant genes, that best follow the global

program, and positive and negative genes, which were broadly defined in other studies

as growth-related genes (12)(27)(28). The analysis of relative expression values is

important here as it allows us to appreciate expression reallocation among partition

sectors (3)(4).

We then focused on unravelling how global and specific regulation combine within this

new scheme. This is largely an open question, with only a few preliminary studies in

minimal bacterial systems suggesting a (somehow unexpected) minor role of TFs in the

control of expression (13)(14)(15). We studied here the problem at a genomic scale

using all available regulatory data to first identify invariant genes as the ones that are

least regulated by TFs. The amount of regulation increases among the rest of

nonspecific classes, and between these and the specific ones. Moreover, specific

genes also show a stronger regulatory coherence than nonspecific genes. Coherence

quantifies the similarity of expression response (to growth rate) between genes and the

TFs acting on them and a strong signal suggests an operative role of the latter. In

addition, among those TFs whose action is particularly coherent, we identify groups

that almost exclusively regulate specific or nonspecific genes, or positive and negative

genes, i.e., the functioning of the TF network is somehow segregated.

Beyond TF regulation, we discriminated two general promoter architectures. Those that

are TATA enriched and shNFR/TFIID depleted (nonspecific negative and specific

genes), and those that are TATA depleted and shNFR/TFIID enriched (nonspecific

invariant and positive genes); features that are similarly observed in metazoans

promoters (29) (Type I promoters, genes expressed in a tissue-specific manner, and

Type II promoters, ubiquitously expressed genes, respectively).

That (nonspecific) positive genes are moderately controlled by TFs (like negative) but

depleted in TATA box (unlike negative) could suggest certain expression features (e.g.,

high level of transcription, Fig. S7) and alternative modes of epigenetic regulation.

Indeed, positive genes are enriched in fragile nucleosomes, which highlights the

regulatory role of nucleosomal stability (16). This is supported by the particular action

of GTFs on these genes [as GTFs fine-tune nucleosomal stability (19)]. In addition, we

find the expression of nonspecific genes being adjusted in distinctive manner by

epigenetic modifiers, with three main configurations: 1/ general activators of invariant

and positive genes, 2/ remodelers working in a dual manner; repressors of negative

genes and activators of positive ones, and 3/ elements acting as general repressors of

negative genes.

Global and specific regulation in a broader context.

Some of the previous features match those observations related to environmental

stress response genes (ESR) (30), so it is interesting to examine how this set fits into

our partition. ESR genes included two complementary subsets, which are enriched in

nonspecific negative and repressed genes (induced ESR genes), or nonspecific

positive genes (repressed ESR genes, Methods). This confirms the suggestion of

previous studies that stress response genes were not responding directly to stress but

rather to the associated decrease in growth rate [but see (31)].

More generally, two models to coordinate gene expression to the available nutrients

can be imagined: a *feed-forward* regulation by signaling pathways that predict growth

rate in a certain environmental condition, or a *feedback* mechanism, which senses

growth rate, or other related internal cell variable, and then modifies expression (32). In

this context, a passive resource allocation model could explain that the global program

is responding always to the environment, although indirectly (as it can only use those

resources that were not consumed in the mounting of the specific response). This

validates, for instance, that ribosomal genes follow the feed-forward model (33). The

fine-grained structure of the nonspecific class (invariant/positive/negative) could

nevertheless monitor growth rate, at least partially, with the feedback being mediated

by epigenetic mechanisms (see below).

If, as suggested by Hansen and O'Shea (34), TFs can mostly transmit qualitative

(presence/absence of a particular nutrient) rather than quantitative (amount of nutrient)

information, how can we then explain the monotonic variation of fractional expression

(with nutrient dilution in the chemostats) of the genes in the negative and positive

partition components? One way is that metabolism, which is highly sensitive to the

limiting nutrient (35) acts as a regulator of the epigenetic factors discussed above.

Indeed, several metabolites (e.g., GlcNAc, NAD+, acetyl-CoA, alpha KG, ATP) are

known to regulate transcription through interactions with enzymes involved in

epigenetic modifications (36). For example, acetyl-CoA induces cell growth and

proliferation by promoting the acetylation of histones at growth genes (37) (histone

acetylation affects rather similarly specific and nonspecific genes, Fig. S6, what

supports its potential role as a widespread mechanism). Another explanation is that the

monotonic variation observed is the result of cell population shifts with growth rate,

instead of changes in single-cell resource allocations. Note that these shifts cannot be

attributed to the fact that slow-growing cells enlarged their G1 cell cycle phase as

neither (12) nor us observed a bias in positive/negative genes with any particular phase

of cell cycled genes.

In this work we have studied changes in fractional expression but not in mRNA

abundances. It is known that the global program dictates that the faster a population of

cells growths, the higher the promoter activity (rate of RNA synthesis) (11) or total

mRNA abundance (rate of RNA synthesis and degradation) (38). We expect most (if

not all) gene products to follow this (absolute) global program, with potential additional

layers of regulation (which are nutrient and gene dependent) that increment or

decrement mRNA levels. Expression of the genes within the invariant group best

follows the absolute global program, with positive genes being slightly above -and

negative genes slightly below- this response (but all of them incrementing mRNA levels

or promoter activities) (e.g., Fig. S1B). On the other hand, it would be interesting to

quantify the degree to which single cells can present a distribution of resources that is

separated from the model here discussed (39), as well as to understand the

mechanisms that lead to such divergence.

In summary, although one could argue that cellular physiology can indeed determine a

global transcriptional program of gene expression control, our work highlights that this

program is adjusted by the integration of effective genetic and epigenetic modes of

regulation. This modulation limits the prospect of "simplifying" our understanding of

genome-wide expression change.

Materials and Methods

Promoter activity (PA) data. Keren *et al.* (11) measured the activities of ~900 *S.*

cerevisiae promoters in 10 different growing conditions using a library of fluorescent

reporters. For each strain in every growth condition, PA was obtained as the YFP

production rate per OD per second in the window of maximal growth.

Five-sector partition based on PA data. We computed fractional PA (f_{PA}) for each

growth condition as the ratio of the PA of each gene to the summed PA of all promoters

(for a gene *i*, $f_{PA,i} = PA_i / \Sigma PA_i$). Ratios of f_{PA}s for each pair of conditions (with

increasing growth rate) were also calculated. We then computed the absolute distance

of these ratios to ratio 1 (i.e., same f_{PA} in both conditions), and defined as invariant

genes the top 350 genes (distance closest to 0) and as activated (repressed) the

bottom 50 with ratio >0 (< 0). The rest of genes with ratio >0 (<0), and both f_{PA}s > 10^{-4},

were designated as nonspecific positive (negative). We used the "typical" class of a

gene (the most frequently occurring category that a gene presents in all pairwise

growth rate changes) to functionally characterize the sectors (Table S1) what

confirmed their biological significance. Minor modifications on the threshold values

defining these sectors did not alter the conclusions.

Microarray data. Brauer *et al.* (12) grew yeast cultures in chemostats under different

continuous culture conditions (six different limiting nutrients each at six dilution rates)

and measured mRNA abundance with two-color microarrays. Since the original

reference channel for all samples corresponded to a particular glucose condition, which

mixes the response of different nutrients, we reanalyzed the data without this reference

by considering the red processed signal as independent channel (40), and normalizing

by the corresponding sum for each case to obtain a fractional score; more specifically,

the fractional expression value of gene *i* is given by $f_i = \log_{10}[10^6(g_i/\Sigma g_i)]$, with g_i being

the corresponding red-processed signal. SVDs were computed on this processed data.

Five-sector partition based on microarray data. We defined as nonspecific genes

those whose difference on the loadings of the 1st component (a_i's) between two

conditions is less, or equal, than three standard deviations of all gene differences (in

absolute values). Genes are otherwise considered specific (activated or repressed if

the difference of a_i's is positive or negative, respectively). Moreover, absolute values of

the loadings of the 2nd component (b_i's) were sorted to define those with smallest

values (top 2500) as invariant genes, with the rest being positive or negative

(determined by the sign of b_i). To define the partition, we classify as nonspecific those

genes that act as nonspecific in >8 pairwise conditions (out of 15). Nonspecific genes

acting as invariant in > 3 conditions (recall that the total number is 6) are labeled as

invariant. Nonspecific and not invariant genes appearing more times as positive than

as negative (in all 6 conditions) are categorized as positive, and likewise for negative.

Specific genes which appear more times as positive than as negative (again in all 6

conditions) are categorized as activated, and analogously for repressed. This partition

is the one used for the all the regulatory analysis (Table S2 shows the connection

between these sectors and those obtained with PA data). Finally, note that minor

modifications on the threshold values defining these sectors did not alter the

conclusions, and that the biological significance of the sectors are validated by the

regulatory and functional signals observed.

Regulatory network. We obtained regulatory data from

http://yeastmine.yeastgenome.org. No microarray data is considered for the TF info;

only data from different manuscripts using chromatin immunoprecipitation, chromatin

immunoprecipitation-chip, chromatin immunoprecipitation-seq, combinatorial evidence,

and computational combinatorial evidence for a total of 20,673 interactions with 133

TFs involved (12). We also calculated the hierarchical organization of the network (41).

Bas1, Mbp1, Med6, Spt7, and Swi6 appear at the top of the hierarchy.

Regulatory coherence. We identified the set of TFs regulating each gene and

quantified the Pearson's correlation coefficient between the expression vector (as a

function of growth rate) of each TF within the set and the target gene to then take the

mean. This is the (mean) regulatory coherence in a given nutrient condition.

Randomizing expression vectors for each gene, 1000 times, we obtained a score of

significance for each gene's regulatory coherence. With this, we identified a list of

genes displaying significant regulatory coherence. Identification of TFs acting more

significantly on each partition component is computed by first measuring how often it

acts on significantly coherent genes, within the five-sector partition, and then estimating

a null value by randomization of the partition classes.

Fragile nucleosome data. Nucleosome occupancy and position have been measured

by analysis of MNase-digested chromatin. Recent work noted that certain nucleosomes

were extremely sensitive to this digestion, and thus obtained a quantitative score of

nucleosome fragility that we used for our analysis, Table S6 in (19).

Chromatin compendium. This set includes 170 gene expression profiles for

chromatin-regulation related mutations (expressed in $\log_2$ ratios) taken from 26 different

publications (22). It covers more than 60 potential interacting chromatin modifiers such

as histone acetyltransferases (HATs; the NuA4, HAT1 and SAGA complexes), histone

deacetylases (HDACs; the RPD3, HDA1 and SET3 complexes), histone

methyltransferases (the COMPASS complex), ATP-dependent chromatin remodelers

(the SWI/SNF, SWR1, INO80, ISWI and RSC complexes), and other chromatin-

affecting genes and cofactors such as Spt10, Sir proteins and the TATA-binding protein

(TBP). We normalized each dataset to unit variance (21). For Fig. S6, we took absolute

values to estimate the strength of the chromatin regulator effect. Note here that growth

rate reduction can be connected to the impact associated to many of these deletions,

so we controlled for the possible contribution of cell cycle population shifts as described

(next section). This enables us to better identify expression changes due to regulation

(42).

Removal of the slow growth signature. We took the full data in (43) to obtain the

slow growth profile and remove the slow growth signature in the epigenetic data

following (42). In brief, the slow growth profile is obtained as the first-mode

approximation of the data after SVD decomposition. To compare with the epigenetic

compendium data, we chose the column of this approximation with the largest norm as

the slow growth signature. The correlation with the slow growth signature is removed

by transforming the epigenetic data in Gram-Schmidt fashion by subtracting from their

projection onto the basis vector, given by the normalized slow growth profile.

Statistical analysis. Null models associated to most results are obtained by randomly

assigning each gene to one of the sectors, to then compute the precise statistic (e.g.,

mean number or regulators in Fig. 3A). We typically considered 10000 randomizations

unless it is stated otherwise and show the mean and +/- 2 std deviations of the

corresponding statistic (e.g., gray shading in Fig. 3A). Moreover, all P values shown in

enrichment analyses (Table S1-2) are calculated using the Hypergeometric distribution

with Holm-Bonferroni correction for multiple testing.

ESR genes. There are 281 stress-induced and 585 stress-repressed genes –as

defined in (30)– within the set of genes delineating the five-sector partition. A subset of

nonspecific negative genes and specific repressed genes corresponds to stress-

induced (232 out of 2053, and 10 out of 70, respectively), while a subset of nonspecific

positive genes corresponds to stress-repressed (485 out of 1914). Note that most of

the features discussed in the main text associated with the five-sector partition remain

when controlling for ESR genes.

Acknowledgments

523 This work was supported by grant FIS2016-78781-R and the Salvador de Madariaga

524 program (grant PRX18/00439) from the Spanish Ministerio de Economía y

525 Competitividad (J.F.P.). We thank John C. Matese for his help with the analysis of the

526 microarray data.

References

1. Liang S-T, et al. (1999) Activities of constitutive promoters in Escherichia coli.
Journal of Molecular Biology 292(1):19–37.

2. Dennis PP, Ehrenberg M, Bremer H (2004) Control of rRNA Synthesis in
Escherichia coli: a Systems Biology Approach. *Microbiology and Molecular Biology*
Reviews 68(4):639–668.

3. Zaslaver A, et al. (2009) Invariant Distribution of Promoter Activities in Escherichia
coli. *PLoS Computational Biology* 5(10):e1000545.

4. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T (2010) Interdependence
of Cell Growth and Gene Expression: Origins and Consequences. *Science*
330(6007):1099–1102.

5. Maaløe O (1979) Regulation of the Protein-Synthesizing Machinery—Ribosomes,
tRNA, Factors, and So On. *Biological Regulation and Development*, ed Goldberger
RF (Springer US, Boston, MA), pp 487–542.

6. Li G-W, Burkhardt D, Gross C, Weissman JS (2014) Quantifying Absolute Protein
Synthesis Rates Reveals Principles Underlying Allocation of Cellular Resources.
Cell 157(3):624–635.

7. Bosdriesz E, Molenaar D, Teusink B, Bruggeman FJ (2015) How fast-growing
bacteria robustly tune their ribosome concentration to approximate growth-rate
maximization. *FEBS Journal* 282(10):2029–2044.

8. Schaechter M, MaalOe O, Kjeldgaard NO (1958) Dependency on Medium and
Temperature of Cell Size and Chemical Composition during Balanced Growth of
Salmonella typhimurium. *Journal of General Microbiology* 19(3):592–606.

9. Scott M, Klumpp S, Mateescu EM, Hwa T (2014) Emergence of robust growth laws
from optimal regulation of ribosome synthesis. *Molecular Systems Biology*
10(8):747–747.

10. Basan M, et al. (2015) Overflow metabolism in Escherichia coli results from
efficient proteome allocation. *Nature* 528(7580):99–104.

11. Keren L, et al. (2014) Promoters maintain their relative activity levels under
different growth conditions. *Molecular Systems Biology* 9(1):701–701.

12. Brauer MJ, et al. (2008) Coordination of Growth Rate, Cell Cycle, Stress Response,
and Metabolic Activity in Yeast. *Molecular Biology of the Cell* 19(1):352–367.

13. Berthoumieux S, et al. (2014) Shared control of gene expression in bacteria by
transcription factors and global physiology of the cell. *Molecular Systems Biology*
9(1):634–634.

14. Gerosa L, Kochanowski K, Heinemann M, Sauer U (2014) Dissecting specific and
global transcriptional regulation of bacterial gene expression. *Molecular Systems*
Biology 9(1):658–658.

15. Kochanowski K, et al. (2017) Few regulatory metabolites coordinate expression of
central metabolic genes in *Escherichia coli*. *Molecular Systems Biology* 13(1):903.

16. Kubik S, et al. (2015) Nucleosome Stability Distinguishes Two Different Promoter
Types at All Protein-Coding Genes in Yeast. *Molecular Cell* 60(3):422–434.

17. Fazio A, et al. (2008) Transcription factor control of growth rate dependent genes
in Saccharomyces cerevisiae: A three factor design. *BMC Genomics* 9(1):341.

18. Fourel G, Miyake T, Defossez P-A, Li R, Gilson É (2002) General Regulatory Factors
(GRFs) as Genome Partitioners. *Journal of Biological Chemistry* 277(44):41736–
41743.

19. Xi Y, Yao J, Chen R, Li W, He X (2011) Nucleosome fragility reveals novel functional
states of chromatin and poises genes for activation. *Genome Research* 21(5):718–
724.

20.   Baptista T, et al. (2017) SAGA Is a General Cofactor for RNA Polymerase II
579   Transcription. *Molecular Cell* 68(1):130-143.e5.

580   21.   Choi JK, Kim Y-J (2008) Epigenetic regulation and the variability of gene
581   expression. *Nature Genetics* 40(2):141–147.

582   22.   Steinfeld I, Shamir R, Kupiec M (2007) A genome-wide analysis in Saccharomyces
583   cerevisiae demonstrates the influence of chromatin modifiers on transcription.
584   *Nature Genetics* 39(3):303–309.

585   23.   Warfield L, et al. (2017) Transcription of Nearly All Yeast RNA Polymerase II-
586   Transcribed Genes Is Dependent on Transcription Factor TFIID. *Molecular Cell*
587   68(1):118-129.e5.

588   24.   Bajić D, Poyatos JF (2012) Balancing noise and plasticity in eukaryotic gene
589   expression. *BMC Genomics* 13(1):343.

590   25.   Sudarsanam P, Iyer VR, Brown PO, Winston F (2000) Whole-genome expression
591   analysis of snf/swi mutants of Saccharomyces cerevisiae. *Proceedings of the*
592   *National Academy of Sciences* 97(7):3364–3369.

593   26.   Holstege FC., et al. (1998) Dissecting the Regulatory Circuitry of a Eukaryotic
594   Genome. *Cell* 95(5):717–728.

595   27.   Regenberg B, et al. (2006) Growth-rate regulated genes have profound impact on
596   interpretation of transcriptome profiling in Saccharomyces cerevisiae. *Genome*
597   *Biology*:13.

598   28.   Castrillo JI, et al. (2007) Growth control of the eukaryote cell: a systems biology
599   study in yeast. *Journal of Biology* 6(2):4.

600   29.   Lenhard B, Sandelin A, Carninci P (2012) Metazoan promoters: emerging
601   characteristics and insights into transcriptional regulation. *Nature Reviews*
602   *Genetics* 13(4):233–245.

603   30.   Gasch AP, et al. (2000) Genomic Expression Programs in the Response of Yeast
604   Cells to Environmental Changes□D. *Molecular Biology of the Cell* 11:17.

605   31.   Ho Y-H, Shishkova E, Hose J, Coon JJ, Gasch AP (2018) Decoupling Yeast Cell
606   Division and Stress Defense Implicates mRNA Repression in Translational
607   Reallocation during Stress. *Current Biology* 28(16):2673-2680.e4.

608   32.   Levy S, Barkai N (2009) Coordination of gene expression with growth rate: A
609   feedback or a feed-forward strategy? *FEBS Letters* 583(24):3974–3978.

610   33.   Levy S, et al. (2007) Strategy of Transcription Regulation in the Budding Yeast.
611   *PLoS ONE* 2(2):e250.

34. Hansen AS, O'Shea EK (2015) Limits on information transduction through
amplitude and frequency regulation of transcription factor activity. *eLife* 4.
doi:10.7554/eLife.06559.

35. Boer VM, Crutchfield CA, Bradley PH, Botstein D, Rabinowitz JD (2010) Growth-
limiting Intracellular Metabolites in Yeast Growing under Diverse Nutrient
Limitations. *Molecular Biology of the Cell* 21(1):198–211.

36. Lu C, Thompson CB (2012) Metabolic Regulation of Epigenetics. *Cell Metabolism*
16(1):9–17.

37. Cai L, Sutter BM, Li B, Tu BP (2011) Acetyl-CoA Induces Cell Growth and
Proliferation by Promoting the Acetylation of Histones at Growth Genes.
Molecular Cell 42(4):426–437.

38. Athanasiadou N, et al. (2016) Growth Rate-Dependent Global Amplification of
Gene Expression. doi:10.1101/044735.

39. Gasch AP, et al. (2017) Single-cell RNA sequencing reveals intrinsic and extrinsic
regulatory heterogeneity in yeast responding to stress. *PLOS Biology*
15(12):e2004050.

40. 't Hoen PAC (2004) Intensity-based analysis of two-colour microarrays enables
efficient and flexible hybridization designs. *Nucleic Acids Research* 32(4):41e – 41.

41. Jothi R, et al. (2009) Genomic analysis reveals a tight link between transcription
factor dynamics and regulatory network architecture. *Molecular Systems Biology*
5. doi:10.1038/msb.2009.52.

42. O'Duibhir E, et al. (2014) Cell cycle population effects in perturbation studies.
Molecular Systems Biology 10(6):732–732.

43. Kemmeren P, et al. (2014) Large-Scale Genetic Perturbations Reveal Regulatory
Networks and an Abundance of Gene-Specific Repressors. *Cell* 157(3):740–752.

Figures

Fig. 1

Figure 1. A new resource allocation model partitions the genome in five expression
sectors. A) Top. Resource allocation model (Model 1) based on three sectors in which the
expression of their constituent genes increases (positive genes, blue), decreases (negative
genes, red), or remains constant (invariant genes, brown) with increasing growth rate.
Bottom. Resource allocation model (Model 2) based on a specific and nonspecific sectors
in which the expression of their constituent genes increases (nonspecific, white) or
decreases (specific, grey) with increasing growth rate. **B)** Partition of genome expression
into five sectors that combine the two previous models. A model-1-like partition appears as
a fine structure within the nonspecific sector. We labelled this scheme as Model 3. **C)**
Fractional promoter activity (fPA) for a transition between two example conditions
increasing growth (glycerol to glucose). Promoters can be classified into the five sectors of
model 3 depending on how their fPA changes (yellow dots indicate those with very low
activity in both conditions). Repressed and activated promoters constitute the specific
sector. Nonspecific promoters are constituted by one invariant type and two other
subclasses whose fPA depends on the growing condition.

Fig. 2

Figure 2. Genome-wide partition of gene expression in five sectors based on SVD. A)
SVD components $\vec{v}_1$ and $\vec{v}_2$ describe baseline fractional expression and dependence with
growth, respectively, and together explain most of the expression variance. **B)** The
fractional expression of every gene $\vec{g}_i$ as a function of growth rate can be approximated by
a linear combination of these two components, with loadings a_i and b_i. We show two
examples (purple circles denote the expression vector, while the black dots correspond to
the two-component approximation; lines added to help visualization) with the same
baseline (dashed line; same a_i) but whose expression increases ($b_i>0$) or decreases ($b_i<0$)
with growth. Data in A) and B) corresponds to growth in limiting glucose conditions. **C)** The
baseline expression of a gene between two conditions can change (top left) or not change
(top right). A gene is nonspecific if its baseline expression remains the same (similar a_i
loadings) in >8 pairwise change of conditions (total of 15), being specific otherwise. Within
nonspecific genes the second loadings b_i's on each nutrient condition (total of 6) enable us
to determine if the gene is positive ($b_i>0$), invariant ($b_i\sim0$) or negative ($b_i<0$). Specific genes
can similarly be divided between repressed or activated. In this way, we can partition the
genome into five sectors what generalizes the scheme discussed with PA data in a subset
of yeast genes (Figure 1). See main text for details.

 Fig. 3

Figure 3. Differential integration of transcriptional regulation and the five-sector resource allocation model. A) Mean number of regulators acting on genes as a function of their response to growth (blue dots; dashed line to help visualization). Grey shading denotes the average null values +/- 2 standard deviations obtained by randomization. **B)** Mean ratio of the fraction of TFs of a given class with respect to growth (e.g., TF_{neg} denotes TFs which are negative genes) for each group of target genes (also for a given class; here we do not distinguish between nonspecific and specific). Histogram obtained in glucose conditions, see also Fig. S5 for other nutrients (** $p < 0.001$, *´ $p < 0.01$, * $p < 0.05$, two-sided KS test). **C)** Regulators that act dominantly, or secondarily, in genes showing a significant regulatory coherence. Color denotes z-score with respect to a null obtained by randomization (positive values denoting enrichment; negative values denoting lack of it). Properties of some regulators are also included (arrows). See main text for details.

700 **Fig. 4**

701

702

703 **Figure 4. Differential integration of epigenetic regulation and the five-sector resource**
704 **allocation model. A)** Mean fraction of global transcriptional regulators (Rap1, Reb1, Cbf1
705 and Mcm1) within the full set of regulators acting on each gene. Grey shading denotes the
706 average null values +/- 2 standard deviations obtained by randomization. Dashed lines to
707 help visualization. **B)** Mean nucleosomal fragility. Shading/lines as before. **C)** Enrichment
708 of nucleosomal free regions (shNFR, blue), presence of TATA boxes (purple), or action of
709 TFIID global factor (green) as function of response class (measured as z-score with respect
710 to a null by randomization; dashed line indicates z-score = +/- 2).

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

Fig. 5

Figure 5. Chromatin modifiers act differentially on nonspecific genes. Mutations in
chromatin modifiers reveal their diverse regulatory role. For each modifier, we plotted its
mean effect, after mutation, on the genes constituting the negative (red circle), invariant
(brown circle), or positive (blue circle) sectors. A kernel density plot corresponding to a null
distribution obtained by permutation is also shown (Methods). We observed three broad
categories: 1/ modifiers acting as activators (mutation decreases significantly the
expression of invariant/positive genes), 2/ dual activator/repressor (mutation
decreasing/increasing significantly expression of positive/invariant or negative genes,
respectively), and 3/ repressors (mutation increasing significantly expression of negative
genes). TAF: TATA-binding protein related factors; HATs: histone acetyltransferases;
HDACs: histone deacetylases.

Supplementary Information

A)

B)

Figure S1. Scaling factors between pairs of conditions. A) Genes that follow a single proportional scaling may serve a definite cellular function according to *(11).* We find a single scaling that describes the change of promoter activity (PA) for different subsets of promoters according to the five-sector partition. The three classes within the nonspecific promoters (invariant, negative, positive) clearly show singular scaling. Shown also a null that corresponds to the ratio of growth rates between conditions (cyan curve). **B)** PA response of a typical invariant, positive and negative gene that corresponds to the *mrs11*, *rps6A* and *atp5*, respectively (conditions sorted by increasing growth rate; this is absolute PA not fractional PA). A null model of the dependence of PA with growth rate is given by the ratio of growth rates (empty circles). Gene categories within the global group clearly separate from the null.

Figure S2. SVD components. First and second SVD components exhibited an analogous trend in all conditions what underlines a core response. As a result, expression of each gene can be approximated by the linear combination of these two components on each nutrient.

778
Figure S3. Association between class of TF and class of cognate target gene for all
nutrients. Fraction of TF class (negative/invariant/positive) acting on target genes divided
also with respect to growth response (negative/invariant/positive; nonspecific and specific
genes were included that we denoted as allneg, etc.). Mean values of each grouping are
shown in bars, while the orange curves show the distribution of each class of TF on each
condition.

785
786
787

788
789

Figure S4. Regulatory coherence and five-sector resource allocation model. A)
Percentage of genes within each class that exhibit a *significant* regulatory coherence with
respect to a null in which classes were assigned randomly (1000 randomizations;
significance implies z-scores > 2). Note that invariant genes show minimal coherence. **B)**
Distribution of genes exhibiting *significant* regulatory coherence in 1 to 6 different nutrient
conditions. Specific genes show more cases of genes significantly coherent in more
different conditions, while invariant genes showed the opposite.

Figure S5. *Cis* and *trans* variability with respect to the five-sector resource allocation model. A cross between a standard laboratory yeast strain and a wild isolate allowed the computation of *cis* and *trans* effects on transcriptional variance *(21)*. For each partition, we quantified the mean of these measures and showed the associated z-score with respect to a null by randomization; dashed line indicates z-score = +/- 2. Positive genes show dominant effects associated with *trans* variability (trv and civ denote *trans* and *cis* variability, respectively).

Figure S6. Chromatin modifiers act differentially on specific and nonspecific genes. The chromatin regulatory effect (CRE; *x* axis) quantifies change in gene expression (absolute value) due to mutations in chromatin modifiers. CRE values larger than expected by a null (z-scores > 2, obtained by randomization) are observed for most modifiers on specific genes. Each circle type corresponds to a class of epigenetic modifier (TAF: TATA-binding protein related factors; HATs: histone acetyltransferases; HDACs: histone deacetylases); *y* axis denotes z-scores and dashed lines emphasizes z-scores within +/-2 values.

Figure S7. Expression abundance and noise with respect to the five-sector resource
allocation model. Mean expression and protein abundance (A) and protein noise (B) with
respect to the five-sector partition as compared to a null in which classes were assigned
randomly (10000 randomizations; y-axis is plotting the associated z-score, shading
corresponds to z-score values within a range of -/+ 2; SD/YEPD denote poor/rich growing
conditions). Nonspecific and positive genes showed high expression and low noise, a
signal that was associated to the presence of fragile nucleosomes in the promoter and the
action of general transcription factors [both enriched in nonspecific positive genes, see
main text and *(24)* for details on data].