Complex genetic and epigenetic regulation deviates gene expression from a unifying global transcriptional program

Mónica Chagoyen[1] and Juan F Poyatos[2]

[1] Computational Systems Biology Group (CNB-CSIC), Darwin 3, 28049, Madrid, Spain
[2] Logic of Genomic Systems Laboratory (CNB-CSIC), Darwin 3, 28049, Madrid, Spain

Email: *monica.chagoyen@cnb.csic.es* (MC), *jpoyatos@cnb.csic.es* (JFP)

Abstract

Environmental or genetic perturbations lead to gene expression changes. While most analyses of these changes emphasize the presence of qualitative differences on just a few genes, we now know that changes are widespread. This large-scale variation has been linked to the exclusive influence of a global transcriptional program determined by the new physiological state of the cell. However, given the sophistication of eukaryotic regulation, we expect to have a complex structure of deviations from the global program. Here, we examine the regulatory landscape that contributes to these deviations. Using data of *Saccharomyces cerevisiae* expression in different nutrient conditions, we first propose a five-component genome partition as a framework to understand expression variation. In this framework, we recognize invariant genes, whose regulation is dominated by the global program, specific genes, which substantially depart from it, and two additional classes that respond to intermediate regulatory schemes. Whereas the invariant class shows a considerable absence of specific regulation, the rest is enriched by regulation at the level of transcription factors (TFs) and epigenetic modulators. We nevertheless find markedly different strategies in how these classes deviate. On the one hand, there are TFs that act in an exclusive way between partition constituents, and on the other, the action of chromatin modifiers is significantly diverse. The balance between regulatory strategies ultimately modulates the action of the general transcription machinery and therefore limits the possibility of establishing a unifying program of expression change at a genomic scale.

Introduction

A global program of regulation impacts the expression of most genes (Liang et al., 1999, Dennis et al., 2004, Zaslaver et al., 2009, Scott et al., 2010). This program depends on the availability of individual components of the cellular expression machinery, e.g., free RNA polymerases, co-factors, ribosomes, etc., which affects the rates of transcription and/or translation and thus determines the overall state of the cell.

The study of this program goes back to the research of the early school of bacterial physiologists who introduced the notion of steady state of cellular growth, in which growth rate acted as a valid proxy for cell physiology and consequently for the global program (Schaechter et al., 1958, Neidhardt and Magasanik, 1960, Maaløe, 1979). Indeed, these initial studies documented that the macromolecular composition of cells is a function of the growth rate. With such experimental approach, the general question of how physiology influences gene expression is confronted by studying how expression depended on the growth rate.

More recent work resumed this research (Zaslaver et al., 2009, Scott et al., 2010, Klumpp and Hwa, 2014, Bosdriesz et al., 2015), emphasizing a framework of distribution of limited resources associated with gene expression, i.e., several cellular parameters manifest as resource trade-offs. Changes in expression as a function of growth rate can be understood by means of coarse genomic partitions that gain differential access to these resources according to separate functional categories; the minimal partition being that between ribosomal and metabolic genes (Zaslaver et al., 2009, Scott et al., 2010, Hui et al., 2015). Likewise, a recent model proposed a broader picture of resource allocation in which only a small group of genes, specific genes, deviate from the global expression program that is constraint by the remaining resources not involved in the activation of the specific response (Keren et al., 2014).

Understanding what precise control mechanisms decouple gene expression from such

global transcriptional program is therefore of great interest. For instance, results in

bacteria have demonstrated the prevalence of the global expression program, while

they have lowered the importance of transcription factors (TFs) controlling the assumed

deviations from it. TFs seem only to complement the action of the global regulation

(Berthoumieux et al., 2014, Gerosa et al., 2014), in combination with a few metabolites

(Kochanowski et al., 2017). Results in eukaryotes are however lacking (Keren et al.,

2014, Metzl-Raz et al., 2017), partly because one has to study a more complex basal

regulatory machinery.

In this work, we wanted to characterize which molecular elements of control influence

the deviation from the global transcriptional program in eukaryotes. For this, we have

outlined in detail the groups of genes that can *a priori* be more sensitive to the global

program. We do this by introducing a partition of the yeast genome into five

components, which consolidates previous models (Zaslaver et al., 2009, Scott et al.,

2010, Keren et al., 2014). In this framework, we recognize invariant genes, whose

regulation is dominated by the global program, specific genes, which are those that

seemingly deviate the most from it, and two intermediate classes of genes that

experience both global and specific regulation. We then study which regulatory factors

act on the components of each element of the partition. We focus both on the influence

of TFs and chromatin modifiers. Our results allow us to fully appreciate the integration

between the specific and global mechanisms controlling genome-wide expression in

eukaryotes, as well as the genetic and epigenetic factors that contribute to this

integration. More broadly, our study emphasizes the limitations of achieving a unique

genome-wide program of expression control.

Results

A five-component partition captures large-scale changes of gene expression. We first examined the changes with the growth rate in the promoter activity (PA) of a subset of ~900 *Saccharomyces cerevisiae* genes in 10 growing conditions (Keren et al., 2014). Studying absolute PA values Keren et al. presented a binary partition that helped explain these changes, recognizing a global proportional response in most genes and a specific one in a much smaller set. To focus on resource reallocation, we considered here instead fractional activities, i.e., the fraction of PA of each gene in a given nutrient condition [out of the summed activity of all genes in the dataset (Maaløe, 1979, Zaslaver et al., 2009)], and quantified their change for each pair of conditions (from low to high growth rate).

We interpreted these changes by delimiting a five-component partition of all genes (Methods). Figure 1A shows a descriptive case (glycerol to glucose growing conditions). Genes whose fractional PA remains approximately invariant (diagonal in Fig. 1A) constitute the first partition element. One can broadly introduce other four components: 1/positive genes (whose fractional PA moderately increases between conditions), 2/negative genes (fractional PA decreases to only a limited extent), 3/specifically activated genes (fractional PA becomes much larger), and 4/ specifically repressed genes (fractional PA becomes much smaller). Note that by reducing the partition to two components, specific and global, the stronger allocation of "expression resources" to specific genes in glycerol as compared to glucose becomes manifest (Fig. 1.B, white/grey pie charts). This leaves fewer resources to biosynthesis (reduction of the global component) affecting growth rate (Keren et al., 2014).

Our analysis identifies a fine-grained structure within global genes (invariant, positive and negative genes), each class following a precise proportional response between

conditions (Fig.S1). Within this structure, we also observed how the resources taken by

positive genes in a condition of high growth rate limit, in turn, those available for the

expression of negative genes (Fig. 1B, brown/red/blue pie charts). The resource

reallocation is also revealed by the (absolute) PA response to growth that exhibits this

type of genes, PA that cannot be simply explained in terms of the associated changes

in growth rate (Maaløe, 1979) (Keren et al., 2014) (Fig. 1C).

With respect to the type of genes that comprise each component, the invariant class is

enriched by transcription regulation and ribosomal proteins; the latter being more

extensively observed in the positive class. Indeed, positive genes are enriched by

ribosomal genes (~65% of genes code for small or large subunits of the ribosome),

while negative genes are enriched in ATP metabolic processes, e.g., oxidative

phosphorylation or the TCA cycle. Lastly, activated and repressed genes indicated

specific expression programs related to the particular carbon source, and the pathways

of the central metabolism that facilitate the transition between fermentation or

respiration routes (Table S1, Methods).

The five-component partition can be delineated on a genomic scale. To generalize

the preceding analysis to a genome-wide scale we analyzed a DNA microarray dataset

of yeast cells exhibiting a range of growth rates for several limiting nutrients (Brauer et

al., 2008). We first applied singular value decomposition (SVD) to the fractional gene

expression quantified on each nutrient separately (Methods). Notably, the first and

second SVD components (Fig. 2A) explain >90% of the variance in each condition. As

a result, fractional expression of each gene can be approximated by the linear

combination of these two components (Fig. 2B), which exhibited an analogous trend in

all nutrients (Fig. S2).

Moreover, while the first component ($\vec{v}_1$) does not change with growth rate, we

observed that the second component ($\vec{v}_2$) exhibits a monotonic behavior (Fig, 2A for

glucose, and Fig. S2 for other nutrients). We can thus interpret the first element of the

linear combination as the baseline fractional expression of the gene, and the second

element as its monotonic behavior with growth rate (Fig. 2B). This interpretation

enables us to generalize the partition framework previously introduced with PA data.

Therefore, a change in the loading of $\vec{v}_1$ (a_i) in two different nutrients implies that the

corresponding gene is specific, as it changes between conditions, and global otherwise

(Fig. 2C) (note that $\vec{v}_1$ is quantitatively similar in all nutrients, Fig. S2). This supports

the framework in (Keren et al., 2014) (Fig. 1B). Comparison of these gene loadings in

the six nutrients revealed that they are fairly similar (minimal correlations found of

~0.96), i.e., much of the gene response is global.

In contrast, the second component ($\vec{v}_2$) provides a quantitative score (the second

loading, b_i) to classify genes as invariant, positive or negative, as before (Fig. 2C,

Methods). Some genes have the same classification in two different nutrients, but this

does not have to be necessarily the case. With the use of the second component, we

can also evaluate how the response to growth rate depends on the exact nutrient

(Brauer et al., 2008). We found that nutrient condition particularly matters in the range

of slow growth and that some nutrients trigger a more similar response to growth than

others (Gutteridge et al., 2010). Finally, the functional analysis of genes within each

class agrees with PA data and previous reports (Table S2).

The partition composition presents different transcriptional regulation. TFs are

the most direct elements that can deviate from the global transcription program the

expression of genes. Before examining this effect, we asked how TFs themselves are

framed in the previous partition. After assembling a transcriptional regulatory network

with existing data (Methods), we observed that most constituent TFs (122 of a total of

133 composing the network) exhibit similar basal fractional expression (a_i loadings)

across all pairwise condition changes, i.e., they are global genes. Within this set, 31%

presents a dominant invariant response ($b_i \sim 0$ in >3 nutrients, of a total of 6), with five

genes acting as invariant in all six conditions (*rsc1*, *mbp1*, *pho2*, *rgr1*, and *swi6*). Two

of these (*mbp1*, *swi6*) are at the top of the network hierarchy (being involved in the

mitotic cell cycle), and two are elements of relevant complexes that interact with RNA

polymerase II (*rsc1* of the RSC chromatin complex, and *rgr1/med14* of the mediator

complex); they can be considered as elements of a general transcriptional machinery,

for which maintaining its concentration invariant across conditions could be essential.

Moreover, 32% of global TFs are dominantly negative, and only 4% positive. Of note,

some of the TFs whose expression decreases with growth ($b_i<0$) are positive regulators

of transcription in response to stress (e.g., *bur6*, *gcn4*, *rpn4*) what justifies their

overexpression at low growth rates.

To what extent is the regulation of target genes dependent on which component of the

partition they belong to? We labeled target genes as global if they showed a global

response (similar a_i loadings) in >8 pairwise change of conditions (total of 15). Genes

are considered specific otherwise. Global genes are less regulated on average than

specific ones [by 3.09 TFs *vs.* 5.06 TFs, p = $1.20 \ 10^{-4}$, two-sample Kolmogorov-

Smirnov (KS) test]. Within global genes, we described as invariant –following again the

second loading score, b_i – those which exhibit this pattern in >3 conditions. Global and

invariant genes are less regulated on average than global and *not* invariant genes (by

2.56 TFs *vs.* 3.3 TFs, p = $8.16 \ 10^{-13}$, two-sample KS test). Finally, global and positive

genes are slightly more regulated than global and negative genes (by 3.33 TFs *vs.* 3.

27 TFs, p = 0.0018, two-sample KS test). Overall, specific genes are subjected to more

regulation (larger number of TFs) (Fig. 3A), while global and invariant genes are the

ones that show the least.

Specific TFs regulate genes belonging to each partition sector. Although Fig. 3A

shows how the structure of transcriptional interactions is reflected differentially in the

components of the partition it does not assure us when these interactions are active,

e.g., (Berthoumieux et al., 2014). For this, we examined several features.

We first inspected if target genes presenting a particular growth response are enriched

by TFs showing the very same response, as the similarity of the responses could imply

that part of the regulatory structure is active. We thus computed –for each target gene–

the fraction of its regulators that behave as negative, invariant, or positive (TF_{neg}, TF_{inv},

TF_{pos}, respectively) with growth rate in a given condition. Figure 3B shows the mean of

the fractions for target genes whose response is negative, invariant or positive.

Negative TFs are more likely to be found acting on target genes that are also negative

(higher mean TF_{neg} on negative genes), while invariant (TF_{inv}) and positive (TF_{pos}) TFs

regulate more often invariant and positive target genes, respectively (the latter signal is

weaker and depends on the particular condition, Fig. S3). Thus, TFs that exhibit the

same behavior as their cognate target gene tend to appear, on average, dominant on

its regulation; part of the regulatory structure seems then functional.

To further test the active effect of TFs, we measured the correlation of the response to

growth rate between any particular gene and all its cognate TFs ("regulatory

coherence", Methods). Specific genes showed stronger regulatory coherence than

global ones (Fig. S4A), and remain coherent in more nutrient conditions (Fig. S4B),

both results implying an active contribution of TFs to deviate gene expression from the

global program. Moreover, Fig. 3C shows those TFs whose action is particularly

coherent *per* partition component (Methods). One can identify here two broad groups,

which control global or specific genes [this is supported by earlier reports (Fazio et al.,

2008)]. In this way, within the regulatory network, we find TFs that act more

significantly on different types of genes. Notably, those that work on global genes are

higher up in the network hierarchy (Methods). We also noted that some these

(significantly coherent) TFs are involved in chromatin remodeling (Cyc8, Ume6, Spt6,

Msn4, Abf1, Msn2, Nhp6A, acting on global ones), or chromatin organization (Spt3,

Spt2, Pho4, FKh2, Sin3, Spt20, Wtm2, Wtm1, Hif1, acting on specific genes). We

examine epigenetic aspects next.

The partition composition also reveals distinctive epigenetic regulation. To

inspect the function of epigenetic control mechanisms, we first quantified the proportion

of general transcription factors (GTFs) found within the set of TFs acting on a given

gene (Fourel et al., 2002). GTFs (Rap1, Abf1, Reb1, Cbf1, and Mcm1) usually have

little intrinsic regulatory activity and together with the presence of chromatin remodelers

(in particular, RSC –Remodels Structure of Chromatin) control an alleged general

machinery of expression. We observed that GTFs constitute a larger and significant

fraction in the regulation of positive genes, while the opposite is observed for negative

ones (Fig. 4A).

GTFs are also connected to particularly fragile nucleosome promoter architectures (Xi

et al., 2011), a connection recently examined (Kubik et al., 2015). Using this data, we

computed the nucleosome landscape for the different gene classes (Methods).

Promoters of positive genes are indeed enriched in fragile nucleosomes (Fig 4.B) while

both negative and invariant genes typically lack these structures. This suggests that

positive genes are more sensitive to deviate from the global program of expression

(implemented by a general transcriptional machinery) by means of chromatin

modulation. Enrichment of other promoter features contribute to this model (Fig 3.C),

like the absence of TATA boxes, the action of TFIID over SAGA [but this precise

grouping has been recently revaluated (Baptista et al., 2017)], the presence of

nucleosomal free regions closer to the transcriptional starting site (shNFRs) (partially

associated to the previous score of fragile nucleosomes), and the dominant effect of

trans variability (Fig. S5) (Choi and Kim, 2008)**.**

We examined additional chromatin modifiers using a previously assembled

compendium (Steinfeld et al., 2007) (Methods). Figure 5 shows the effects of mutating

different types of *trans*-acting chromatin regulators on the genes constituting the

partitions. Note here that growth rate reduction can be connected to many of these

deletions, so we controlled for the possible contribution of cell cycle population shifts as

described (Methods). This enables us to better identify expression changes due to

regulation (O'Duibhir et al., 2014). With the exception of histone acetyltransferases

(HATs) and TATA-binding protein related factors (TAFs), the effect of most chromatin

modifiers is dominant in specific genes (Fig. 5A, Methods). Within global genes we

found three main configurations (Fig. 5B): 1) Epigenetic regulators acting as part of a

general machinery (HATs –including SAGA–, TAFs and methyltransferases) whose

mutation causes a general decrease in expression, very particularly in invariant and

positive classes. Indeed, work by (Baptista et al., 2017) and (Warfield et al., 2017)

demonstrated that SAGA and TFIID are recruited to pol II promoters genome-wide and

that each complex is generally required for pol II transcription, i.e., its mutation would

lead to a genome-wide decrease of gene expression. 2) Regulators (histones, and

chromatin remodelers) acting in a dual manner: increasing the expression of negative

genes (remodeler as a repressor) or reducing their expression in positives (remodeler

as an activator). This underlines the enrichment of negative and positive classes by

stress and ribosomal genes, respectively, which are largely regulated in opposite

manner (Bajić and Poyatos, 2012); a dual role of remodelers as activators and

repressors have been previously reported (Sudarsanam et al., 2000, Holstege et al.,

1998). And 3) regulators as broad repressors that represent regulation by gene

silencing.

Discussion

Could one interpret genome-wide expression changes as deviations from a global

program of transcriptional control? In this work, we propose to answer this question by

capturing these deviations in a five-component partition of the yeast genome. The

analysis of relative expression values is necessary here as it helps us to appreciate

expression reallocation among partition constituents. Therefore, this integrates an

earlier model (Keren et al., 2014) that reduced variation between conditions to a

proportional response shared by most genes and induced by a global transcriptional

program (with only a limited number of specifically expressed genes), while also

discriminating three subclasses within such response. Invariant genes, that best follow

the global program, and positive and negative genes, which were broadly defined in

other studies as growth-related genes (Regenberg et al., 2006, Castrillo et al., 2007,

Brauer et al., 2008). The biological significance of our framework is reinforced by its

differential promoter and regulatory architecture ranging from a model of almost

passive control to one exhibiting complex combinatorial regulation.

More explicitly, invariant genes are those subjected to less regulation by TFs,

regulation that increases among the rest of global classes, and between these and the

specific ones. Specific genes also show a stronger regulatory coherence than global

genes (similarity of expression response to that of the TFs acting on them). In addition,

among TFs whose action is particularly coherent, we identify two groups that almost

exclusively regulate global or specific genes: the action of the TF network is somehow

segregated. Beyond TF regulation, we can discriminate between two broad promoter

architectures. Those that are TATA enriched and shNFR/TFIID depleted (global

negative and specific genes), and those that are TATA depleted and shNFR/TFIID

enriched (global invariant and positive genes). Notably, these features are similarly

observed in metazoans promoters (Lenhard et al., 2012) (Type I promoters, genes

expressed in a tissue-specific manner, and Type II promoters, ubiquitously expressed

genes, respectively).

That (global) positive genes are moderately controlled by TFs (like negative) but

depleted in TATA box (unlike negative) could suggest certain expression features (e.g.,

high level of transcription, Fig. S6) and alternative modes of regulation. Indeed, positive

genes are enriched in fragile nucleosomes, which highlights the regulatory role of

nucleosomal stability. This is supported by the particular action of GTFs on these

genes [as GTFs fine-tune nucleosomal stability (Xi et al., 2011, Kubik et al., 2015)]. In

addition, we find the expression of global genes being adjusted in distinctive manner by

epigenetic modifiers, with three main configurations: 1/ HATs, TAFs and

methyltransferases working as general activators of invariant and positive genes, 2/

histone and chromatin remodelers working in a dual manner; repressors of negative

genes and activators of positive ones, and 3/ gene silencing elements acting as

general repressors of negative genes.

The five-component partition in a broader context.

Some of the previous features discussed in our framework of the five-component

partition match those observations related to environmental stress response genes

(ESR) (Gasch et al., 2000), so it is interesting to examine how this set fits into our

partition. ESR genes included two complementary subsets, which are enriched in our

global negative and repressed genes (induced ESR genes), or global positive genes

(repressed ESR genes, Methods). This confirms the suggestion of previous studies

that stress response genes were not responding directly to stress but rather to the

associated decrease in growth rate. More generally, two models to coordinate gene

expression to the available nutrients can be imagined: a feed-forward regulation by

signaling pathways that predict growth rate in a certain environmental condition, or a

feedback mechanism, which senses growth rate, or other related internal cell variable,

and then modifies expression (Levy and Barkai, 2009). In this context, a passive

resource allocation model could explain that the global program is responding always

to the environment, although indirectly (as it can only use those resources that were

not consumed in the mounting of the specific response). This validates, for instance,

that ribosomal genes follow the feed-forward model (Levy et al., 2007). The fine-

grained structure of the global class (invariant/positive/negative) could nevertheless

monitor growth rate, at least partially, with the feedback being mediated by epigenetic

mechanisms (see below).

If, as suggested by (Hansen and O'Shea, 2015), TFs can mostly transmit qualitative

(presence/absence of a particular nutrient) rather than quantitative (amount of nutrient)

information, how can we then explain the monotonic variation of fractional expression

with nutrient dilution in the chemostat of the genes in the negative and positive partition

components? One way is that metabolism, which is highly sensitive to the limiting

nutrient (Boer et al., 2010) acts as a regulator of the epigenetic factors discussed

above. Indeed, several metabolites (e.g., GlcNAc, NAD+, acetyl-CoA, alpha KG, ATP)

are known to regulate transcription through interactions with enzymes involved in

epigenetic modifications (Lu and Thompson, 2012). For example, acetyl-CoA induces

cell growth and proliferation by promoting the acetylation of histones at growth genes

(Cai et al., 2011) (histone acetylation affects rather similarly specific and global genes,

Fig. 5A, what supports its potential role as a widespread mechanism). Another

explanation is that the monotonic variation observed is the result of cell population

shifts with growth rate, instead of changes in single-cell resource allocations. Note that

these shifts cannot be attributed to the fact that slow-growing cells enlarged their G1

cell cycle phase as neither (Brauer et al. 2008) nor us observed a bias in

positive/negative genes with any particular phase of cell cycled genes.

In this work we have studied changes in fractional expression but not in mRNA

abundances. It is known that the global program dictates that the faster a population of

cells growths, the higher the promoter activity (rate of RNA synthesis) (Keren et al.,

2014) or total mRNA abundance (rate of RNA synthesis and degradation)

(Athanasiadou et al., 2016). We expect most (if not all) gene products to follow this

(absolute) global program, with potential additional layers of regulation (which are

nutrient and gene dependent) that increment or decrement mRNA levels. The invariant

group best describes the absolute global program, while positive genes are slightly

above and negative genes slightly below this program (but all of them incrementing

mRNA levels or promoter activities) (e.g., Fig. 1C). On the other hand, it would be

interesting to quantify the degree to which single cells can present a distribution of

resources that is separated from the model here discussed (Gasch et al., 2017), as

well as to understand the mechanisms that lead to such divergence.

In summary, although one could argue that cellular physiology can indeed determine a

global transcriptional program of gene expression control, our work highlights that this

program is mediated by the integration of genetic and epigenetic modes of regulation,

what limits the prospect of "simplifying" our understanding of genome-wide expression

change.

Materials and Methods

Promoter activity (PA) data. (Keren et al., 2014) measured the activities of ~900 *S.*

cerevisiae promoters in 10 different growing conditions using a library of fluorescent

reporters. For each strain in every growth condition, promoter activity was obtained as

the YFP production rate per OD per second in the window of maximal growth.

Genome partition based on PA data. Fractional promoter activity (f_{PA}) for each

growth condition and ratios of f_{PA}s for each pair of conditions (with increasing growth

rate) were calculated. We then computed the absolute distance of these ratios to ratio

1 (i.e., same f_{PA} in both conditions), and defined as invariant genes the top 350 genes

(distance closest to 0) and as activated (repressed) the bottom 50 with ratio >0 (< 0).

The rest of genes with ratio >0 (<0), and both f_{PA}s > 10^{-4}, were designated as global

positive (negative). We used the "typical" class of a gene (the most frequently occurring

category that a gene presents in all pairwise growth rate changes) to select the

examples of Fig. 1C, and to characterize the partition in functional terms (Table S1).

Microarray data. (Brauer et al., 2008) grew yeast cultures in chemostats under

different continuous culture conditions (six different limiting nutrients each at six dilution

rates) and measured mRNA abundance with two-color microarrays. Since the original

reference channel for all samples corresponded to a particular glucose condition, which

mixes the response of different nutrients, we reanalyzed the data without this reference

by considering the red processed signal as independent channel ('t Hoen, 2004), and

normalizing by the corresponding sum for each case (to obtain a fractional score).

SVDs were computed on this processed data.

Genome partition based on microarray data. Global genes are those whose

difference on the loadings of the 1st component (a_i's) between two conditions is less, or

equal, than three standard deviations of all gene differences (in absolute values).

Genes are otherwise considered specific (activated or repressed if the difference of a_i's

is positive or negative, respectively). Moreover, absolute values of the loadings of the

2nd component (b_i's) were sorted to define those with smallest values (top 2500) as

invariant genes, with the rest being positive or negative (determined by the sign of b_i).

To define the partition, we classify as global those genes that act as global in >8

pairwise conditions (out of 15). Global genes acting as invariant in > 3 conditions (recall

that the total number is 6) are labeled as invariant. Global and not invariant genes

appearing more times as positive than as negative (in all 6 conditions) are categorized

as positive, and likewise for negative. Specific genes which appear more times as

positive than as negative (again in all 6 conditions) are categorized as activated, and

analogously for repressed.

Regulatory network. We obtained regulatory data from

http://yeastmine.yeastgenome.org. No microarray data is considered for the TF info;

only data from different manuscripts using chromatin immunoprecipitation, chromatin

immunoprecipitation-chip, chromatin immunoprecipitation-seq, combinatorial evidence,

and computational combinatorial evidence for a total of 20,673 interactions with 133

TFs involved (Brauer et al., 2008). We also calculated the hierarchical organization of

the network (Jothi et al., 2009). Bas1, Mbp1, Med6, Spt7, and Swi6 appear at the top of

the hierarchy.

Fragile nucleosome data. Nucleosome occupancy and position have been measured

by analysis of MNase-digested chromatin. Recent work noted that certain nucleosomes

were extremely sensitive to this digestion, and thus obtained a quantitative score of

nucleosome fragility that we used for our analysis, Table S6 in (Kubik et al., 2015).

Chromatin compendium. This set includes 170 gene expression profiles for

chromatin-regulation related mutations (expressed in $\log_2$ ratios) taken from 26 different

publications (Steinfeld et al., 2007). It covers more than 60 potential interacting

chromatin modifiers such as histone acetyltransferases (HATs; the NuA4, HAT1 and

SAGA complexes), histone deacetylases (HDACs; the RPD3, HDA1 and SET3

complexes), histone methyltransferases (the COMPASS complex), ATP-dependent

chromatin remodelers (the SWI/SNF, SWR1, INO80, ISWI and RSC complexes), and

other chromatin-affecting genes and cofactors such as Spt10, Sir proteins and the

TATA-binding protein (TBP). We normalized each dataset to unit variance (Choi and

Kim, 2008). For Fig. 5A, we took absolute values to estimate the strength of the

chromatin regulator effect.

Regulatory coherence. We identified the set of TFs regulating each gene and

quantified the Pearson's correlation coefficient between the expression vector (as a

function of growth rate) of each TF within the set and the target gene to then take the

mean. This is the (mean) regulatory coherence in a given nutrient condition.

Randomizing expression vectors for each gene, 1000 times, we obtained a score of

significance for each gene's regulatory coherence. With this, we identified a list of

genes displaying significant regulatory coherence. Identification of TFs acting more

significantly on each partition component is computed by first measuring how often it

acts on significantly coherent genes, within the five-component partition grouping, and

then estimating a null value by randomization of the partition classes.

Removal of the slow growth signature. We took the full data in (Kemmeren et al.,

2014) to obtain the slow growth profile and remove the slow growth signature in the

epigenetic data following (O'Duibhir et al., 2014). In brief, the slow growth profile is

obtained as the first-mode approximation of the data after SVD decomposition. To

compare with the epigenetic compendium data, we chose the column of this

approximation with the largest norm as the slow growth signature. The correlation with

the slow growth signature is removed by transforming the epigenetic data in Gram-

Schmidt fashion by subtracting from their projection onto the basis vector, given by the

normalized slow growth profile.

ESR genes. There are 281 stress-induced and 585 stress-repressed genes –as

defined in (Gasch et al., 2000)– within the set of genes delineating the five-component

partition. A subset of global negative genes and specific repressed genes corresponds

to stress-induced (232 out of 2053, and 10 out of 70, respectively), while a subset of

global positive genes corresponds to stress-repressed (485 out of 1914). Note that

most of the features discussed in the main text associated with the five-component

partition remain when controlling for ESR genes.

Acknowledgments

This work was supported by grant FIS2016-78781-R from the Spanish Ministerio de

Economía y Competitividad (J.F.P.). We thank John C. Matese for his help with the

analysis of the microarray data. **Authors contributions:** MC and JFP conceived the

project, performed the analysis, discussed the results, and wrote the paper.

Competing interests: The authors declare that no competing interests exist in relation

to this manuscript.

References

Athanasiadou, N., Neymotin, B., Brandt, N., Miller, D., Tranchina, D., Gresham, D.,
2016. Growth Rate-Dependent Global Amplification of Gene Expression.
bioRxiv https://doi.org/10.1101/044735.
Bajić, D., Poyatos, J.F., 2012. Balancing noise and plasticity in eukaryotic gene
expression. BMC Genomics 13, 343.
Baptista, T., Grünberg, S., Minoungou, N., Koster, M.J.E., Timmers, H.T.M., Hahn, S.,
Devys, D., Tora, L., 2017. SAGA Is a General Cofactor for RNA Polymerase II
Transcription. Molecular Cell 68, 130-143.e5.
Berthoumieux, S., de Jong, H., Baptist, G., Pinel, C., Ranquet, C., Ropers, D.,
Geiselmann, J., 2014. Shared control of gene expression in bacteria by
transcription factors and global physiology of the cell. Molecular Systems
Biology 9, 634–634.
Boer, V.M., Crutchfield, C.A., Bradley, P.H., Botstein, D., Rabinowitz, J.D., 2010.
Growth-limiting Intracellular Metabolites in Yeast Growing under Diverse
Nutrient Limitations. Molecular Biology of the Cell 21, 198–211.
Bosdriesz, E., Molenaar, D., Teusink, B., Bruggeman, F.J., 2015. How fast-growing
bacteria robustly tune their ribosome concentration to approximate growth-rate
maximization. FEBS Journal 282, 2029–2044.
Brauer, M.J., Huttenhower, C., Airoldi, E.M., Rosenstein, R., Matese, J.C., Gresham,
D., Boer, V.M., Troyanskaya, O.G., Botstein, D., 2008. Coordination of Growth
Rate, Cell Cycle, Stress Response, and Metabolic Activity in Yeast. Molecular
Biology of the Cell 19, 352–367.
Cai, L., Sutter, B.M., Li, B., Tu, B.P., 2011. Acetyl-CoA Induces Cell Growth and
Proliferation by Promoting the Acetylation of Histones at Growth Genes.
Molecular Cell 42, 426–437.
Castrillo, J.I., Zeef, L.A., Hoyle, D.C., Zhang, N., Hayes, A., Gardner, D.C., Cornell,
M.J., Petty, J., Hakes, L., Wardleworth, L., Rash, B., Brown, M., Dunn, W.B.,
Broadhurst, D., O'Donoghue, K., Hester, S.S., Dunkley, T.P., Hart, S.R.,
Swainston, N., Li, P., Gaskell, S.J., Paton, N.W., Lilley, K.S., Kell, D.B., Oliver,
S.G., 2007. Growth control of the eukaryote cell: a systems biology study in
yeast. Journal of Biology 6, 4.
Choi, J.K., Kim, Y.-J., 2008. Epigenetic regulation and the variability of gene
expression. Nature Genetics 40, 141–147.
Dennis, P.P., Ehrenberg, M., Bremer, H., 2004. Control of rRNA Synthesis in
Escherichia coli: a Systems Biology Approach. Microbiology and Molecular
Biology Reviews 68, 639–668.

Fazio, A., Jewett, M.C., Daran-Lapujade, P., Mustacchi, R., Usaite, R., Pronk, J.T., Workman, C.T., Nielsen, J., 2008. Transcription factor control of growth rate dependent genes in Saccharomyces cerevisiae: A three factor design. BMC Genomics 9, 341.

Fourel, G., Miyake, T., Defossez, P.-A., Li, R., Gilson, É., 2002. General Regulatory Factors (GRFs) as Genome Partitioners. Journal of Biological Chemistry 277, 41736–41743.

Gasch, A.P., Spellman, P.T., Kao, C.M., Carmel-Harel, O., Eisen, M.B., Storz, G., Botstein, D., Brown, P.O., 2000. Genomic Expression Programs in the Response of Yeast Cells to Environmental Changes□D. Molecular Biology of the Cell 11, 17.

Gasch, A.P., Yu, F.B., Hose, J., Escalante, L.E., Place, M., Bacher, R., Kanbar, J., Ciobanu, D., Sandor, L., Grigoriev, I.V., Kendziorski, C., Quake, S.R., McClean, M.N., 2017. Single-cell RNA sequencing reveals intrinsic and extrinsic regulatory heterogeneity in yeast responding to stress. PLOS Biology 15, e2004050.

Gerosa, L., Kochanowski, K., Heinemann, M., Sauer, U., 2014. Dissecting specific and global transcriptional regulation of bacterial gene expression. Molecular Systems Biology 9, 658–658.

Gutteridge, A., Pir, P., Castrillo, J.I., Charles, P.D., Lilley, K.S., Oliver, S.G., 2010. Nutrient control of eukaryote cell growth: a systems biology study in yeast. BMC Biology 8, 68.

Hansen, A.S., O'Shea, E.K., 2015. Limits on information transduction through amplitude and frequency regulation of transcription factor activity. eLife 4.

Holstege, F.C.., Jennings, E.G., Wyrick, J.J., Lee, T.I., Hengartner, C.J., Green, M.R., Golub, T.R., Lander, E.S., Young, R.A., 1998. Dissecting the Regulatory Circuitry of a Eukaryotic Genome. Cell 95, 717–728.

Hui, S., Silverman, J.M., Chen, S.S., Erickson, D.W., Basan, M., Wang, J., Hwa, T., Williamson, J.R., 2015. Quantitative proteomic analysis reveals a simple strategy of global resource allocation in bacteria. Molecular Systems Biology 11, e784–e784.

Jothi, R., Balaji, S., Wuster, A., Grochow, J.A., Gsponer, J., Przytycka, T.M., Aravind, L., Babu, M.M., 2009. Genomic analysis reveals a tight link between transcription factor dynamics and regulatory network architecture. Molecular Systems Biology 5.

Kemmeren, P., Sameith, K., van de Pasch, L.A.L., Benschop, J.J., Lenstra, T.L., Margaritis, T., O'Duibhir, E., Apweiler, E., van Wageningen, S., Ko, C.W., van Heesch, S., Kashani, M.M., Ampatziadis-Michailidis, G., Brok, M.O., Brabers, N.A.C.H., Miles, A.J., Bouwmeester, D., van Hooff, S.R., van Bakel, H., Sluiters, E., Bakker, L.V., Snel, B., Lijnzaad, P., van Leenen, D., Groot Koerkamp, M.J.A., Holstege, F.C.P., 2014. Large-Scale Genetic Perturbations Reveal Regulatory Networks and an Abundance of Gene-Specific Repressors. Cell 157, 740–752.

Keren, L., Zackay, O., Lotan-Pompan, M., Barenholz, U., Dekel, E., Sasson, V., Aidelberg, G., Bren, A., Zeevi, D., Weinberger, A., Alon, U., Milo, R., Segal, E., 2014. Promoters maintain their relative activity levels under different growth conditions. Molecular Systems Biology 9, 701–701.

Klumpp, S., Hwa, T., 2014. Bacterial growth: global effects on gene expression, growth feedback and proteome partition. Current Opinion in Biotechnology 28, 96–102.

Kochanowski, K., Gerosa, L., Brunner, S.F., Christodoulou, D., Nikolaev, Y.V., Sauer, U., 2017. Few regulatory metabolites coordinate expression of central metabolic genes in *Escherichia coli*. Molecular Systems Biology 13, 903.

Kubik, S., Bruzzone, M.J., Jacquet, P., Falcone, J.-L., Rougemont, J., Shore, D., 2015. Nucleosome Stability Distinguishes Two Different Promoter Types at All Protein-Coding Genes in Yeast. Molecular Cell 60, 422–434.

Lenhard, B., Sandelin, A., Carninci, P., 2012. Metazoan promoters: emerging
characteristics and insights into transcriptional regulation. Nature Reviews
Genetics 13, 233–245.
Levy, S., Barkai, N., 2009. Coordination of gene expression with growth rate: A
feedback or a feed-forward strategy? FEBS Letters 583, 3974–3978.
Levy, S., Ihmels, J., Carmi, M., Weinberger, A., Friedlander, G., Barkai, N., 2007.
Strategy of Transcription Regulation in the Budding Yeast. PLoS ONE 2, e250.
Liang, S.-T., Bipatnath, M., Xu, Y.-C., Chen, S.-L., Dennis, P., Ehrenberg, M., Bremer,
H., 1999. Activities of constitutive promoters in Escherichia coli. Journal of
Molecular Biology 292, 19–37.
Lu, C., Thompson, C.B., 2012. Metabolic Regulation of Epigenetics. Cell Metabolism
16, 9–17.
Maaløe, O., 1979. Regulation of the Protein-Synthesizing Machinery—Ribosomes,
tRNA, Factors, and So On, in: Goldberger, R.F. (Ed.), Biological Regulation and
Development. Springer US, Boston, MA, pp. 487–542.
Metzl-Raz, E., Kafri, M., Yaakov, G., Soifer, I., Gurvich, Y., Barkai, N., 2017. Principles
of cellular resource allocation revealed by condition-dependent proteome
profiling. eLife 6, e28034.
Neidhardt, F.C., Magasanik, B., 1960. Studies on the role of ribonucleic acid in the
growth of bacteria. Biochimica et Biophysica Acta 42, 99–116.
O'Duibhir, E., Lijnzaad, P., Benschop, J.J., Lenstra, T.L., van Leenen, D., Groot
Koerkamp, M.J., Margaritis, T., Brok, M.O., Kemmeren, P., Holstege, F.C.,
2014. Cell cycle population effects in perturbation studies. Molecular Systems
Biology 10, 732–732.
Regenberg, B., Grotkj, T., Åkesson, M., Bro, C., Hansen, L.K., Brunak, S., Nielsen, J.,
2006. Growth-rate regulated genes have profound impact on interpretation of
transcriptome profiling in Saccharomyces cerevisiae. Genome Biology 13.
Schaechter, M., MaalOe, O., Kjeldgaard, N.O., 1958. Dependency on Medium and
Temperature of Cell Size and Chemical Composition during Balanced Growth
of Salmonella typhimurium. Journal of General Microbiology 19, 592–606.
Scott, M., Gunderson, C.W., Mateescu, E.M., Zhang, Z., Hwa, T., 2010.
Interdependence of Cell Growth and Gene Expression: Origins and
Consequences. Science 330, 1099–1102.
Steinfeld, I., Shamir, R., Kupiec, M., 2007. A genome-wide analysis in Saccharomyces
cerevisiae demonstrates the influence of chromatin modifiers on transcription.
Nature Genetics 39, 303–309.
Sudarsanam, P., Iyer, V.R., Brown, P.O., Winston, F., 2000. Whole-genome
expression analysis of snf/swi mutants of Saccharomyces cerevisiae.
Proceedings of the National Academy of Sciences 97, 3364–3369.
't Hoen, P.A.C., 2004. Intensity-based analysis of two-colour microarrays enables
efficient and flexible hybridization designs. Nucleic Acids Research 32, 41e –
41.
Warfield, L., Ramachandran, S., Baptista, T., Devys, D., Tora, L., Hahn, S., 2017.
Transcription of Nearly All Yeast RNA Polymerase II-Transcribed Genes Is
Dependent on Transcription Factor TFIID. Molecular Cell 68, 118-129.e5.
Xi, Y., Yao, J., Chen, R., Li, W., He, X., 2011. Nucleosome fragility reveals novel
functional states of chromatin and poises genes for activation. Genome
Research 21, 718–724.
Zaslaver, A., Kaplan, S., Bren, A., Jinich, A., Mayo, A., Dekel, E., Alon, U., Itzkovitz, S.,
2009. Invariant Distribution of Promoter Activities in Escherichia coli. PLoS
Computational Biology 5, e1000545.

Figures

Fig. 1

Figure 1. Partition of gene expression in five sectors. A) Fractional promoter activity
(fPA) between two example conditions. Promoters can be classified into five categories
(inset) depending on how their fPA changes (yellow dots indicate those with very low
activity in both conditions). **B)** Repressed and activated promoters constitute the specific
response, whose fPA is greater the lower the growth rate (at the cost of the promoters
changing in a global manner). Global promoters are constituted by one invariant type and
two other subclasses whose fPA depends on the growing condition. Note how the portion
of expression, within global genes, of positive genes increases with growth rate, while it
decreases for negative ones (colors as in A). **C)** Absolute promoter activity (PA) response
of a typical invariant, positive and negative gene that corresponds to the *mrs11*, *rps6A* and
atp5, respectively (conditions sorted by increasing growth rate). A null model of the
dependence of PA with growth rate is given by the ratio of growth rates (empty circles).
Gene categories within the global group clearly separate from the null (see main text).

647 **Fig. 2**

648
649

Figure 2. Genome-wide partition of gene expression in five sectors based on SVD. A)
SVD components $\vec{v}_1$ and $\vec{v}_2$ describe baseline fractional expression and dependence with
growth, respectively, and together explain most of the expression variance. **B)** The
fractional expression of every gene $\vec{g}_i$ as a function of growth rate can be approximated by
a linear combination of these two components, with loadings a_i and b_i. We show two
examples (purple circles denote the expression vector, while the black dots correspond to
the two-component approximation; lines added to help visualization) with the same
baseline (dashed line; same a_i) but whose expression increases ($b_i > 0$) or decreases ($b_i < 0$)
with growth. Data in A) and B) corresponds to growth in limiting glucose conditions. **C)** A
given gene can be considered specific or global when its baseline fractional expression
(bars) changes (left) or does not change (right) with different nutrients (different or
comparable a_i, respectively). Beyond the baseline value on each condition, expression can
increase, decrease or remain constant with growth-rate (μ) change. We thus compare a_i's
between pairwise conditions (total of 15) to define a gene as specific or global, whereas the
second loadings b_i's on each condition (total of 6) enable us to determine if the gene is
positive, invariant or negative.

666
667
668

Fig. 3

Figure 3. Genes within each partition sector show differential transcriptional
regulation. **A)** Mean number of regulators acting on genes as a function of their response
to growth (blue dots; dashed line to help visualization). Grey shading denotes the average
null values +/- 2 standard deviations obtained by randomization. **B)** Mean ratio of the
fraction of TFs of a given class with respect to growth (e.g., TF_{neg} denotes TFs which are
negative genes) for each group of target genes (also for a given class; here we do not
distinguish between global and specific). Histogram obtained in glucose conditions, see
also Fig. S5 for other nutrients (** $p < 0.001$, *´ $p < 0.01$, * $p < 0.05$, two-sided KS test). **C)**
Regulators that act dominantly in genes showing a significant regulatory coherence. Colour
denotes z-score with respect to a null obtained by randomization (positive values denoting
enrichment). Properties of some regulators are also included (arrows). See main text for
details.

Fig. 4

Figure 4. Genes within each partition sector show differential epigenetic regulation.
A) Mean fraction of global transcriptional regulators (Rap1, Reb1, Cbf1 and Mcm1) within
the full set of regulators acting on each gene. Grey shading denotes the average null
values +/- 2 standard deviations obtained by randomization. Dashed lines to help
visualization. **B)** Mean nucleosomal fragility. Shading/lines as before. **C)** Enrichment of
nucleosomal free regions (shNFR, blue), presence of TATA boxes (purple), or action of
TFIID global factor (green) as function of response class (measured as z-score with respect
to a null by randomization; dashed line indicates z-score = +/- 2).

Fig. 5

Figure 5. Chromatin modifiers act differentially on genes within each partition. A) The
chromatin regulatory effect quantifies change in gene expression (absolute value) due to
mutations in chromatin modifiers. Most modifiers show a significant effect on specific genes
as compared to a null (obtained by randomization; grey shading indicates non-significant
values, and dashed lines denote z-scores of +/-2). **B)** Mutations in chromatin modifiers
reveal their diverse regulatory role when acting on negative, invariant, or positive genes.
We observed three main categories: 1/ modifiers acting as activators (mutation decreases
significantly the expression of invariant/positive genes), 2/ dual activator/repressor
(mutation decreasing/increasing significantly expression of positive/invariant or negative
genes, respectively), and 3/ repressors (mutation increasing significantly expression of
negative genes). Significance and shading as before (note that a significant increase or
decrease in expression correspond to positive or negative z-scores, respectively). Different
type of modifiers corresponds to filled/empty circles of different size, whose colour denotes
the precise target gene class considered. TAF: TATA-binding protein related factors; HATs:
histone acetyltransferases; HDACs: histone deacetylases.

 Supplementary Information

Figure S1. Scaling factors between pairs of conditions. A single proportionality
(scaling) factor describes the change of promoter activity for different subsets of promoters
according to the five-component partition. The three classes within the global promoters
(invariant, negative, positive) clearly show different scaling. Shown also a null that
corresponds to the ratio of growth rates between conditions.

Figure S2. SVD components. First and second SVD components exhibited an analogous trend in all conditions what underlines a core response. As a result, expression of each gene can be approximated by the linear combination of these two components on each nutrient.

730
Figure S3. Association between class of TF and class of cognate target gene for all
nutrients. Fraction of TF class (negative/invariant/positive) acting on target genes divided
also with respect to growth response (negative/invariant/positive; global and specific genes
were included that we denoted as allneg, etc.). Mean values of each grouping are shown
in bars, while the orange curves show the distribution of each class of TF on each
condition.

737
738
739

740
741

Figure S4. Regulatory coherence and five-component partition. A) Percentage of
genes within each class that exhibit a significant regulatory coherence with respect to a null
in which classes were assigned randomly (1000 randomizations; significance implies z-
scores > 2). Note that invariant genes show minimal coherence. B) Distribution of genes
exhibiting significant regulatory coherence in 1 to 6 different nutrient conditions. Specific
genes show more cases of genes significantly coherent in more different conditions, while
invariant genes showed the opposite.

750
Figure S5. *Cis* and *trans* variability with respect to the five-component partition. A
cross between a standard laboratory yeast strain and a wild isolate allowed the
computation of *cis* and *trans* effects on transcriptional variance (Choi and Kim, 2008). For
each partition, we quantified the mean of these measures and showed the associated z-
score with respect to a null by randomization; dashed line indicates z-score = +/- 2. Positive
genes show dominant effects associated with *trans* variability (trv and civ denote *trans* and
cis variability, respectively).

759
760
Figure S6. Expression abundance and noise with respect to the five-component
partition. Mean expression and protein abundance (A) and protein noise (B) with respect
to the five-component partition as compared to a null in which classes were assigned
randomly (10000 randomizations; y-axis is plotting the associated z-score, shading
corresponds to z-score values within a range of -/+ 2; SD/YEPD denote poor/rich growing
conditions). Global and positive genes showed high expression and low noise, a signal that
was associated to the presence of fragile nucleosomes in the promoter and the action of
general transcription factors [both enriched in global positive genes, see main text and
(Bajić and Poyatos, 2012*)* for details on data].

28