1  Pleotropic effects of mutations in the effector domain of influenza A virus NS1 protein
2

3  Carina F. Pereira[1‡], Helen M. Wise[1,2†], Dominic Kurian[2], Rute M. Pinto[2], Maria J. Amorim[1,3],
4  Andrew C. Gill[2, 4] and Paul Digard[1,2#]
5
[1]Division of Virology, Department of Pathology, University of Cambridge, Tennis Court Road,
Cambridge, CB2 1QP, UK.
[2]The Roslin Institute, University of Edinburgh, Easter Bush, Midlothian EH25 9RG, UK.
[3]Cell Biology of Viral Infection, Instituto Gulbenkian de Ciência, *Rua da Quinta Grande 6,* P-
2780-156 Oeiras, Portugal.
[4]School of Chemistry, Joseph Banks Laboratories, University of Lincoln, Green Lane, Lincoln,
Lincolnshire LN6 7DL
13
14  [‡]Present address: European Commission, Place Rogier 16, B-1210, Brussels, Belgium
15  [†]Present address: Department of Engineering and Physical Sciences, Heriot Watt University,
16  Edinburgh, UK.
23  [#]Corresponding author: Paul Digard
24  Tel:           +44 131 651 9240
25  Fax:           +44 131 651 9105
26  Email          paul.digard@roslin.ed.ac.uk
27
28  Other author email addresses:
29
CFP: carina.pereira@ec.europa.eu
HMW: h.wise@hw.ac.uk
RMP: s1564909@sms.ed.ac.uk
DK: Dominic.Kurian@roslin.ed.ac.uk
MJA: mjamorim@igc.gulbenkian.pt
ACG: angill@lincoln.ac.uk

## Abstract

Objective: The multifunctional NS1 protein of influenza A virus has roles in antagonising

cellular innate immune responses and promoting viral gene expression.  To better

understand the interplay between these functions, we tested the effects of NS1 effector

domain mutations known to affect homo-dimerisation or interactions with cellular PI3

kinase or Trim25 on NS1 ability to promote nuclear export of viral mRNAs.

Results: The NS1 dimerisation mutant W187R retained the functions of binding cellular

NXF1 as well as stabilising NXF1 interaction with viral segment 7 mRNAs and promoting

their nuclear export. Two PI3K-binding mutants, NS1 Y89F and Y89A still bound NXF1 but no

longer promoted NXF1 interactions with segment 7 mRNA or its nuclear export. The Trim25-

binding mutant NS1 E96A/E97A bound NXF1 and supported NXF1 interactions with segment

7 mRNA but no longer supported mRNA nuclear export. Analysis of WT and mutant NS1

interaction partners identified hsp70 as specifically binding to NS1 E96A/E97A. Whilst these

data suggest the possibility of functional links between NS1's effects on intracellular

signalling and its role in viral mRNA nuclear export, they also indicate potential pleiotropic

effects of the NS1 mutations; in the case of E96A/E97A possibly via disrupted protein folding

leading to chaperone recruitment.

Keywords: influenza A virus, NS1, PI3K, Trim25, RIG-I, nuclear export

## Introduction

Influenza A virus (IAV) infects both man and animals, including species important for food security [1]. A better understanding of its replication cycle is important to aid control measures. Like all viruses, IAV co-opts multiple cellular systems for replication and also has to counteract innate immune defence mechanisms. A key virally-encoded factor in both processes is the NS1 protein. As a nuclear-replicating virus, IAV utilises the NXF1 cellular mRNA export pathway to direct several of its mRNAs to the cytoplasm [2] and we and others have recently found that NS1 acts as an adaptor between segment 7 mRNAs (encoding the M1 or M2 proteins) and cellular mRNA processing/export pathways [3, 4]. NS1 is also the primary viral antagonist of innate immune responses, interfering with the function of multiple cellular polypeptides including PKR, 2'-OAS, CPSF30 and Trim25 [5]. NS1 also activates the phosphoinositide 3-kinase (PI3K) complex, possibly to inhibit apoptosis [6]. Structurally, NS1 consists of an N-terminal RNA-binding domain and a C-terminal "effector" domain followed by a disordered tail, with no identified enzymatic activities. Instead, it functions by binding to multiple different proteins and RNAs (viral and cellular) and altering their activities [5]. Various NS1 point mutations have been identified that inhibit these interactions and comparative analyses of the behaviour of wild type (WT) and mutant IAV strains with these alterations have provided important information on the biological significance of each particular function. Here, we tested whether mutations identified as affecting NS1 interactions with the p85ß subunit of PI3K (Y89A and Y89F), Trim25 (E96A/E97A) and NS1 effector domain-mediated oligomerisation (W187R) [7-11] also affected NS1's ability to promote nuclear export of segment 7 viral mRNAs.

**Methods.**

All IAV plasmid constructs and viruses were based on the A/PR/8/34 (PR8) strain described

previously [12]. NS1 mutations used in this study were introduced by standard PCR

mutagenesis reactions; primer sequences are available on request. Methods for virus

reverse genetics, infection, minireplicon assays, fluorescent *in situ* hybridization (FISH),

immunofluorescence and confocal microscopy, GFP-trap pulldowns, western blots and

reverse transcriptase-radioactive primer extension are described elsewhere [4]. LC-MS

analysis was performed using a RSLCnano system (Thermo Fisher Scientific) coupled to a

micrOTOF QII mass spectrometer (Bruker) on in-gel trypsin digested protein bands excised

from Coomassie Blue-stained SDS-PAGE gels [13]. Raw spectral data were processed to peak

lists and searched using Mascot 2.4 server (Matrix Science) against the Uniprot Human

sequence database containing 93,786 entries.

**Results.**

Our previous work led to the hypothesis of NS1 acting as an adaptor protein to feed

viral segment 7 mRNA(s) into the cellular NXF1 export pathway, by forming a ternary

complex between NS1, NXF1 and the mRNA that requires both the NS1 effector domain and

a functional RNA-binding domain [4]. To further test the role of the NS1 effector domain, a

set of point mutations in this region of the protein were tested (Table 1), firstly in a subviral

"minireplicon" system. 293T cells were transfected with plasmids to produce segment 7

ribonucleoproteins (RNPs) as well as with either WT or mutant NS1s as EGFP-fusion

proteins. Cells were fixed 24 h post transfection and segment 7 mRNA localisation detected

**Table 1. Genetic and phenotypic summary of the NS1 polypeptides used in this study.**

| NS1 protein | mRNA export | NXF1 binding | M1 mRNA binding | Functional alteration | Reference |
|---|---|---|---|---|---|
| WT | + | + | + | n/a | [12] |
| N81[a,b] | - | - | - | Effector domain deletion | [14] |
| R38A/K41A [b] | - | - | - | dsRNA-binding | [15] |
| F103S/M106I [b] | + | + | + | Gain of CPSF inhibition | [16] |
| Y89A | - | + | - | PI3K-binding | [17] |
| Y89F | - | + | - | PI3K-binding | [8] |
| E96A/97A | - | + | + | Trim25-binding | [9] |
| W187R | + | + | + | Homo-dimerisation | [11] |

[a]Stop codon added at position 82. [b]See [4] for experimental data.

by FISH (**Figure 1A**). NS1 expression was confirmed by detection of GFP signal as well as by

western blot, although the E96A/E97A mutant accumulated to lower levels than the other

NS1s (**Figure 1B**). WT NS1-GFP and most mutant proteins localised largely to the cytoplasm,

but the W187R mutant was found both in nucleus and cytoplasm (**Figure 1A**). No mRNA

staining was detected in the negative control lacking a complete RNP. As expected [4],

when the minimal components for RNP reconstitution were present, segment 7 mRNA was

detected primarily in cell nuclei, whilst on addition of NS1-GFP, it translocated to the

cytoplasm. However, in the presence of PI3K-binding mutants NS1-Y89A or NS1-Y89F or the

Trim25-binding mutant NS1-E96A/E97A, segment 7 mRNA was clearly still retained in the

nucleus. In contrast, the dimerization mutant NS1-W187R caused segment 7 mRNA to

localise to the cytoplasm. To quantify these effects, cells were visually scored according to

the predominant cellular localisation of segment 7 mRNA (nuclear, cytoplasmic or both). In

the absence of NS1, segment 7 was mainly nuclear in ~80% of cells scored, while the

remaining cells presented mRNA staining both in the nucleus and the cytoplasm (**Figure 1C**).

When WT NS1 or NS1-W187R were added, a statistically significant shift of segment 7 mRNA

localisation was confirmed, with ~ 75% of cells showing mRNA in the cytoplasm. However,

addition of NS1-E96A/E97A, NS1-Y89A or NS1-Y89F did not produce this shift, with a clear

majority of cells displaying nuclear segment 7 mRNA, similarly to samples with no NS1-GFP.

To further test the mRNA export activity of the NS1 effector domain mutants, their

phenotype was examined in the context of virus infection. 293T cells were mock infected or

infected with either WT PR8 or the NS1 mutant viruses and at 6 h post infection (p.i.), fixed

and processed for segment 7 FISH analysis as well as immunofluorescently stained for NS1

(**Figure 1D**). All viruses generally gave cytoplasmic NS1 signal above the uninfected cell

background, although staining of cells infected with NS1-Y89 and NS1-E96A/E97A mutants

was noticeably fainter than the WT and W187R viruses. No viral mRNA signal was detected

in the mock sample while segment 7 mRNA was cytoplasmic in WT and W187R virus-

infected cells. In cells infected with NS1-Y89A, NS1-Y89F and NS1-E96A/E97A, segment 7

mRNA was predominantly nuclear. Thus, in both minireplicon and viral settings, the NS1

dimerization mutant W187R supported normal segment 7 mRNA nuclear export, but the

PI3K-binding and Trim25-binding mutants showed loss-of-function.

NS1 both binds NXF1 and promotes the ability of NXF1 to bind segment 7 mRNA [4,

18-20]. To test whether the effector domain mutants retained NXF1-binding activity, 293T

cells were transfected with either GFP or GFP-NXF1 and 48 h later mock infected or infected

with the panel of viruses. At 6 h p.i., cells were lysed and GFP-Trap precipitations were

performed on the supernatants. Western blot analyses of total and bound fractions showed

that GFP and GFP-NXF1 were both expressed and collected in comparable amounts (**Figure**

**2A**). Blotting for NS1 confirmed successful infection, although the NS1-E96A/E97A

polypeptide again accumulated to lower levels than the other NS1s. Nevertheless, all NS1

polypeptides co-precipitated with GFP-NXF, indicating that the mutations did not abrogate

the interaction.

To test whether the effector domain mutations affected the ability of NS1 to

promote a stable interaction between NXF1 and segment 7 mRNA, 293T cells were

transfected with GFP or GFP-NXF1 and infected with WT or mutant viruses as above. At 6 h

p.i., cells were lysed and total RNA was extracted before or after the lysates had been

subjected to GFP-Trap pulldown. Segment 7 RNA species were then detected by primer

extension. Viral mRNAs and genomic vRNA were detected in the total fraction of every

infected sample (**Figure 2B**, lanes 1-2 and 4-7). Analysis of the bound fraction from WT virus

infected cells showed that both segment 7 mRNA species but not genomic vRNA co-

precipitated with GFP-NXF1 (lane 9) while no viral RNAs bound to GFP alone (lane 8),

consistent with previous work [4, 18]. In contrast, viral mRNAs were not detected in GFP-

NXF1 pulldowns from cells infected with either NS1-Y89A or NS1-Y89F viruses (lanes 11 and

12). The NS1-W187R and -E96A/E97A mutants also supported NXF1 interactions with the

viral mRNAs (lane 13, 14). Since the E96A/E97A mutation blocked successful nuclear export

of segment 7 mRNA, this finding indicated that, like the interaction between NS1 and NXF1,

the interaction between NXF1 and segment 7 is necessary, but not sufficient for the

successful nuclear export of segment 7 mRNA.

The behaviour of the NS1 E96A/E97A mutant – unable to mediate nuclear export of

segment 7 mRNA despite binding to NXF1 and promoting its interaction with the mRNA –

suggested the involvement of another essential factor in the hypothesised ternary export

complex. Accordingly, we examined cellular interaction partners of WT NS1 and the

mutants, including a complete deletion of the effector domain (NS1-N81), an RNA-binding

domain mutant (R38A/K41A) and a gain-of-function mutant that restores CPSF30-binding

activity (F103S/M106I), of which only the latter supported mRNA nuclear export (Table 1;

[4]). 293T cells were transfected with GFP or GFP-tagged NS1 mutant plasmids. Two days

later, cells were collected, lysed and subjected to GFP-Trap. Bound fractions were loaded

onto SDS-PAGE gels and stained with Coomassie Blue. No obvious contamination from

cellular GFP-binding proteins was seen from cells transfected with the GFP control plasmid

(**Figure 2C**, lane 1). Major polypeptide species of the expected size were detected in all

other samples, including the truncated form NS1-N81, confirming successful expression

(lanes 2-9). Various other minor polypeptide species were also visible that were mostly

common to all NS1-containing samples. However, one clear difference was observed; a

distinct novel polypeptide in the E96A/E97A preparation (lane 8) that was absent from

other samples. LC-MS analysis of the excised band unambiguously identified this

polypeptide as heat shock protein 70 (hsp70) through multiple peptide matches (**Table S1**).

To confirm the interaction between NS1-E96A/E97A and hsp70, repeat GFP-trap

pulldowns were analysed by western blotting for hsp70 as well as for the GFP-tagged "bait"

proteins. Transfection was confirmed by the detection of GFP and GFP-tagged NS1 mutant

proteins, while equivalent amounts of the cellular proteins hsp70, Nup62 and UAP56 in the

total fractions of all samples confirmed that comparable amounts of cell lysates had been

generated (**Figure 2D**, lanes 1-9). However, in the bound fraction, hsp70 was detected only

in the NS1-E96A/E97A sample (lane 17), confirming the specific interaction with hsp70 for

NS1 E96A/E97A.

**Discussion**

The data reported here extend our previous investigation [4] into the role of NS1 in

facilitating nuclear export of viral mRNA. Firstly, we show that higher order multimerization

of NS1 via its effector domain is not needed for viral mRNA nuclear export. Secondly, the

finding that point mutations in the NS1 effector domain, and not just its complete deletion

along with the C-terminal unstructured region [4], can block viral mRNA nuclear export,

strengthens the argument for the functional involvement of this domain in the process.

Thirdly, identification of an NS1 mutant (E96A/E97A) that still binds NXF1 and enhances its

interaction with segment 7 mRNA without upregulating mRNA nuclear export suggests that

at least one other factor is needed for export. Finally, the effect of mutations previously

considered to specifically inhibit PI3K activation or inhibition of RIG-I signalling on M1 mRNA

localisation could suggest the involvement of one or both of these pathways in viral mRNA

nuclear export.

Of broader significance than the mechanism of IAV mRNA nuclear export, our data

add to the evidence suggesting pleiotropic effects from NS1 mutations commonly used as

tools to interrogate viral interactions with the PI3K and RIG-I pathways (*e.g.* [10, 17, 21-

28]). Potential pleiotropic effects of the E96A/E97A mutation have been noted previously,

including poor expression of the polypeptide as seen here [25] and an intriguing effect on

PI3K activation [29]. Indeed a recent report detailing the crystal structure of NS1-Trim25

complexes concluded that NS1 residues E96 and 97 do not directly contact Trim25 but

instead support a network of contacts within the local region of the effector domain in

which L95 and S99 actually make contact with Trim25 [30]. Our finding that the E96A/E97A

mutation also induces binding of hsp70 further suggests effects on protein folding and that

interpreting results obtained with this mutation may be complicated.

**Limitations**

Attempts to probe the role of PI3K activation in IAV mRNA export using small molecule

inhibitors gave inconsistent results (data not shown) and could not be further pursued

because of time and funding constraints. It would also have been interesting to test the

effects of hsp70 inhibitors [31] had time allowed. The effect of NS1 mutations L95A/S99A on

mRNA export should also be tested.

**Declarations**

**Ethics approval and consent to participate:** not applicable.

**Availability of data and material:** available from the corresponding author on reasonable

request.

**Funding**: PD, ACG, DK and HW were supported by grants BBS/E/D/20241864 and

BBS/E/D/20002173 from the British Biotechnology and Biological Sciences Research Council.

CFP was supported by PhD studentship no. SFRH/BD/60299/2009 from the Portuguese

Fundação para a Ciência e a Tecnologia. MJA was supported by grant no. IF/00899/2013

from the Portuguese Fundação para a Ciência e a Tecnologia and the Fundação Calouste

Gulbenkian. RMP acknowledges PhD studentship support from the Roslin Institute. None of

the funding bodies had any role in directing or interpreting the research.

**Acknowledgements:** we thank Professor Adrian Whitehouse for the gift of pGFP-NXF1.

**Consent for publication:** not applicable.

**Competing interests:** none to declare.

**Author's contributions:** CFP carried out the majority of the experimental work with input

from MJA, HW and RMP. DK and ACG performed the LC-MS analyses. CFP and PD drafted

the manuscript and all authors read, corrected and approved the final manuscript.

256
257
258
259

**Figures and Legends**

262
263
**Figure 1. Effect of mutations in the NS1 effector domain on segment 7 mRNA localisation.**
A-C) 293T cells were transfected with plasmids to recreate segment 7 RNPs or, as a negative

control with a combination that excluded a plasmid expressing PB2 (2PNP), together with
the indicated NS1-GFP polypeptides. (A) Cells were processed for FISH analysis at 24 h post
transfection (p.t.) using Cy3-labelled RNA probes for detection of positive sense segment 7
viral RNAs (red). GFP fluorescence was also detected (green). Images were captured using a
Leica-TCS confocal microscope and Leica TCS analysis software. (B) Duplicate cell samples
were lysed and examined by SDS-PAGE and western blotting for GFP and tubulin. (C)
Individual cells were scored according to the predominant cellular localisation of segment 7
mRNA considering three phenotypes: nuclear, cytoplasmic or mixed. Values are the mean ±
range from two independent experiments. A 2-way ANOVA was used to test for statistically
significant differences between the values for each category from those obtained for WT-
NS1 (NS, not significant; **** $P < 0.0001$).  (D) 293T cells were mock infected or infected
with the indicated viruses at an M.O.I. of 5. At 6 h post infection (p.i.), cells were processed
for FISH as above (red) and stained with anti-NS1 serum (green) and DAPI (for DNA; blue).
Scale-bars indicate 10 µm. Lanes 1 and 2 in panel (B) and a portion of the corresponding
numerical data in panel (C) are taken with permission from Figure 4C and 4B respectively of
[4].
282

283
284
**Figure 2. Interactions between NS1, cellular proteins and segment 7 mRNAs.** (A) 293T cells
were transfected with plasmids encoding GFP or GFP-NXF1 and 48 h later either mock
infected or infected with the indicated NS1 mutant viruses at an M.O.I. of 5. Cells were
harvested at 6 h p.i. and cell lysates examined by western blotting for the indicated proteins
before (Total) or after (Bound) being subjected to GFP-Trap pulldown assays. (B) 293T cells
were transfected with GFP-NXF1 (GFP-NXF1 +) or with GFP alone (GFP-NXF1 -) and 48 h later
either mock infected or infected with the indicated viruses at an M.O.I. of 10, in duplicate
sets. At 6 h p.i., cells were harvested and total cellular RNA was extracted from one set
while the second set was subjected to GFP-Trap pulldown assays prior to total RNA
extraction. RNA was analysed by reverse transcription with radiolabelled primers specific for
segment 7 RNAs followed by urea-PAGE and autoradiography. (C, D) 293T cells were
transfected with either GFP alone or with GFP-tagged NS1 mutant protein expressing
plasmids as indicated and 48 h later cells were harvested and subjected to GFP-Trap pull
down assays. Eluted samples were loaded onto SDS-PAGE gels and (C) stained with
Coomassie Blue dye or (D) western blotted for GFP or hsp70 before (Totals) or after (Bound)
GFP-trap fractionation. Lanes 1-3 and 8-10 from panel (A) are reproduced with permission
from Figure 7 of [4].
302

## References

304 **References**
305
306
1. Taubenberger JK, Kash JC: **Influenza virus evolution, host adaptation, and pandemic**
**formation.** *Cell Host Microbe* 2010, **7**:440–451.

2. York A, Fodor E: **Biogenesis, assembly, and export of viral messenger ribonucleoproteins**
**in the influenza A virus infected cell.** *RNA Biol* 2013, **10**:1274–1282.

3. Mor A, White A, Zhang K, Thompson M, Esparza M, Muñoz-Moreno R, Koide K, Lynch KW,
García-Sastre A, Fontoura BMA: **Influenza virus mRNA trafficking through host nuclear**
**speckles**. *Nature Microbiology* 2016, **1**:1–12.

4. Pereira CF, Read EKC, Wise HM, Amorim MJ, Digard P: **The influenza A virus NS1 protein**
**promotes efficient nuclear export of unspliced viral M1 mRNA.** *J Virol* 2017:JVI.00528–17.

5. Hale BG, Randall RE, Ortín J, Jackson D: **The multifunctional NS1 protein of influenza A**
**viruses.** *J Gen Virol* 2008, **89**:2359–2376.

6. Ayllon J, García-Sastre A, Hale BG: **Influenza A viruses and PI3K: Are there time, place**
**and manner restrictions?** *Virulence* 2012, **3**:411–414.

7. Aramini JM, Ma L-C, Zhou L, Schauder CM, Hamilton K, Amer BR, Mack TR, Lee H-W,
Ciccosanti CT, Zhao L, Xiao R, Krug RM, Montelione GT: **Dimer interface of the effector**
**domain of non-structural protein 1 from influenza A virus: an interface with multiple**
**functions.** *J Biol Chem* 2011, **286**:26050–26060.

8. Hale BG, Jackson D, Chen YH, Lamb RA, Randall RE: **Influenza A virus NS1 protein binds**
**p85β and activates phosphatidylinositol-3-kinase signaling.** *Proc Natl Acad Sci USA* 2006,
**103**:14194–14199.

9. Gack MU, Albrecht RA, Urano T, Inn K-S, Huang I-C, Carnero E, Farzan M, Inoue S, Jung JU,
García-Sastre A: **Influenza A virus NS1 targets the ubiquitin ligase TRIM25 to evade**
**recognition by the host viral RNA sensor RIG-I.** *Cell Host Microbe* 2009, **5**:439–449.

10. Hrincius ER, Hennecke A-K, Gensler L, Nordhoff C, Anhlan D, Vogel P, McCullers JA,
Ludwig S, Ehrhardt C: **A single point mutation (Y89F) within the non-structural protein 1 of**
**influenza A viruses limits epithelial cell tropism and virulence in mice.** 2012, **180**:2361–
2374.

11. Hale BG, Barclay WS, Randall RE, Russell RJ: **Structure of an avian influenza A virus NS1**
**protein effector domain.** *Virology* 2008, **378**:1–5.

12. de Wit E, Spronken MIJ, Bestebroer TM, Rimmelzwaan GF, Osterhaus ADME, Fouchier
RAM: **Efficient generation and growth of influenza virus A/PR/8/34 from eight cDNA**
**fragments.** *Virus Res* 2004, **103**:155–161.

13. Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M: **In-gel digestion for mass**
**spectrometric characterization of proteins and proteomes.** *Nat Protoc* 2006, **1**:2856–2860.

14. Zürcher T, Marión RM, Ortín J: **Protein synthesis shut-off induced by influenza virus**
**infection is independent of PKR activity.** *Journal of Virology* 2000, **74**:8781–8784.

15. Wang W, Riedel K, Lynch P, Chien CY, Montelione GT, Krug RM: **RNA binding by the**
**novel helical domain of the influenza virus NS1 protein requires its dimer structure and a**
**small number of specific basic amino acids.** *RNA* 1999, **5**:195–205.

16. Kochs G, García-Sastre A, Martínez-Sobrido L: **Multiple anti-interferon actions of the**
**influenza A virus NS1 protein.** 2007, **81**:7011–7021.

17. Lopes AM, Domingues P, Zell R, Hale BG: **Structure-Guided Functional Annotation of the**
**Influenza A Virus NS1 Protein Reveals Dynamic Evolution of the p85β-Binding Site during**
**Circulation in Humans.** *J Virol* 2017, **91**:e01081–17.

18. Wang W, Cui Z-Q, Han H, Zhang Z-P, Wei H-P, Zhou Y-F, Chen Z, Zhang X-E: **Imaging and**
**characterizing influenza A virus mRNA transport in living cells.** *Nucleic Acids Research*
2008, **36**:4913–4928.

19. Larsen S, Bui S, Perez V, Mohammad A, Medina-Ramirez H, Newcomb LL: **Influenza**
**polymerase encoding mRNAs utilize atypical mRNA nuclear export.** *Virol J* 2014, **11**:154.

20. Satterly N, Tsai P-L, van Deursen J, Nussenzveig DR, Wang Y, Faria PA, Levay A, Levy DE,
Fontoura BMA: **Influenza virus targets the mRNA export machinery and the nuclear pore**
**complex.** *PNAS* 2007, **104**:1853–1858.

21. Jackson D, Killip MJ, Galloway CS, Russell RJ, Randall RE: **Loss of function of the**
**influenza A virus NS1 protein promotes apoptosis but this is not due to a failure to**
**activate phosphatidylinositol 3-kinase (PI3K).** *Virology* 2010, **396**:94–105.

22. Lu X, Masic A, Li Y, Shin Y, Liu Q, Zhou Y: **The PI3K/Akt pathway inhibits influenza A**
**virus-induced Bax-mediated apoptosis by negatively regulating the JNK pathway via ASK1.**
*Journal of General Virology* 2010, **91**:1439–1449.

23. Ayllon J, Hale BG, Garcia-Sastre A: **Strain-Specific Contribution of NS1-Activated**
**Phosphoinositide 3-Kinase Signaling to Influenza A Virus Replication and Virulence.** *Journal*
*of Virology* 2012, **86**:5366–5370.

24. Rajsbaum R, Albrecht RA, Wang MK, Maharaj NP, Versteeg GA, Nistal-Villán E, García-
Sastre A, Gack MU: **Species-specific inhibition of RIG-I ubiquitination and IFN induction by**
**the influenza A virus NS1 protein.** *PLoS Pathog* 2012, **8**:e1003059.

25. Khaperskyy DA, Hatchette TF, McCormick C: **Influenza A virus inhibits cytoplasmic stress**
**granule formation.** *FASEB J* 2012, **26**:1629–1639.

26. Liedmann S, Hrincius ER, Guy C, Anhlan D, Dierkes R, Carter R, Wu G, Staeheli P, Green
DR, Wolff T, McCullers JA, Ludwig S, Ehrhardt C: **Viral suppressors of the RIG-I-mediated**
**interferon response are pre-packaged in influenza virions.** *Nat Commun* 2014, **5**:5645.

27. Moriyama M, Chen I-Y, Kawaguchi A, Koshiba T, Nagata K, Takeyama H, Hasegawa H,
Ichinohe T: **The RNA- and TRIM25-Binding Domains of Influenza Virus NS1 Protein Are**

**Essential for Suppression of NLRP3 Inflammasome-Mediated Interleukin-1β Secretion.** *J*
*Virol* 2016, **90**:4105–4114.

28. Tang Q, Wang X, Gao G: **The Short Form of the Zinc Finger Antiviral Protein Inhibits**
**Influenza A Virus Protein Expression and Is Antagonized by the Virus-Encoded NS1**. *Journal*
*of Virology* 2017, **91**:e01909–16.

29. Hale BG, Kerry PS, Jackson D, Precious BL, Gray A, Killip MJ, Randall RE, Russell RJ:
**Structural insights into phosphoinositide 3-kinase activation by the influenza A virus NS1**
**protein.** *Proc Natl Acad Sci USA* 2010, **107**:1954–1959.

30. Koliopoulos MG, Lethier M, Veen AG, Haubrich K, Hennig J, Kowalinski E, Stevens RV,
Martin SR, Sousa CRE, Cusack S, Rittinger K: **Molecular mechanism of influenza A NS1-**
**mediated TRIM25 recognition and inhibition**. *Nat Commun* 2018, **9**:1820.

31. Chatterjee S, Burns TF: **Targeting Heat Shock Proteins in Cancer: A Promising**
**Therapeutic Approach.** *Int J Mol Sci* 2017, **18**:1978.