The Metabotropic Glutamate Receptor 5 Negative Allosteric Modulator Fenobam: Pharmacokinetics, Side Effects, and Analgesic Effects in Healthy Human Subjects.

Laura F. Cavallone[1], Michael C. Montana[1], Karen Frey[1], Dorina Kallogjeri[2], James M. Wages[3], Thomas. Rodebaugh[4], Tina Doshi[5], Evan D. Kharasch[6] & Robert W. Gereau, IV[1,7]

1. Department of Anesthesiology, Washington University in Saint Louis School of Medicine, Saint Louis, MO

2. Department of Otolaryngology-Head and Neck Surgery, Washington University School of Medicine in St Louis, St Louis, MO

3. Pain Center, Precision Spine Care, Texarkana, TX

4. Department of Psychological and Brain Sciences, Washington University in Saint Louis, Saint Louis, MO

5. Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, Baltimore, MD

6. Department of Anesthesiology, Duke University School of Medicine
Durham, NC

7. Washington University Pain Center, St Louis, MO

Corresponding Author: Laura F. Cavallone, MD, MSCI
Washington University School of Medicine
Anesthesiology Department
660 South Euclid Ave
Campus Box 8054
St. Louis, MO 63110
Ph. (314) 362-2358
Email: cavallonel@wustl.edu

Abstract

Metabotropic glutamate receptor 5 (mGlu5) has been shown to modulate nociception in animals, but no mGlu5 antagonists have been developed commercially as analgesics. The mGlu5 antagonist fenobam [N-(3-chlorophenyl)-N'-(4,5-dihydro-1-methyl-4-oxo-1H-imidazole-2-yl)urea] was originally evaluated for development as a non-benzodiazepine anxiolytic. Fenobam is analgesic in numerous mouse pain models, acting exclusively via mGlu5 blockade. Furthermore, fenobam showed no signs of analgesic tolerance with up to two weeks of daily dosing in mice. Analgesic effects of fenobam in humans have not been reported. The purpose of this investigation was to evaluate fenobam pharmacokinetics and analgesic effects in humans. We first evaluated single-dose oral fenobam disposition in a parallel-group dose-escalation study in healthy volunteers. A second investigation tested the analgesic effects of fenobam in an established experimental human pain model of cutaneous sensitization utilizing capsaicin cream and heat, in a double-blind placebo-controlled crossover study. The primary outcome measure was the area of hyperalgesia and allodynia around the area applied with heat/capsaicin. Secondary outcome measures included nociception, measured as pain rating on a visual analog scale, heat-pain detection threshold, and effects on cognition and mood. Fenobam plasma exposures showed considerable interindividual variability, and were not linear with dose. Fenobam reduced sensitization vs placebo at a single time-point (peak plasma concentration); we found no other difference between fenobam and placebo. Our results suggest highly variable fenobam disposition, and minimal analgesic effects at the dose tested. We suggest that future studies testing analgesic effects of mGlu5 blockade are warranted, but such studies should employ molecules with improved pharmacokinetic profiles.

Introduction

Metabotropic glutamate receptor 5 (mGlu5) has emerged as a potential candidate for the development of a new class of analgesic drugs. Despite demonstrated anti-nociceptive effects of mGlu5 antagonists in a broad range of animal pain models[6; 15] , so far, no mGlu5 antagonists have been developed commercially as analgesics.

The investigational drug fenobam [N-(3-chlorophenyl)-N'-(4,5-dihydro-1-methyl-4-oxo-1H-imidazole-2-yl)urea] was originally evaluated for development by Ortho-McNeil (McN-3377) in the late 1970s - early 1980s as a non-benzodiazepine anxiolytic, with a then unknown molecular target[13; 16; 23; 24]. While the commercial development of fenobam was not pursued, in 2005, Porter et al. characterized fenobam as a selective, non-competitive mGlu5 antagonist[26]. In agreement with earlier reports of the role of mGlu5 in nociceptive pathways[29] and the selectivity of fenobam for mGlu5, the analgesic effect of fenobam has been demonstrated in multiple mouse models of inflammatory, neuropathic, and visceral pain[7; 14; 17; 21], with no analgesic effect in mGlu5 knockout mice[21]. Additionally, two-week daily dosing did not result in tolerance to fenobam analgesia in mice[22]. Pre-clinical studies demonstrating the analgesic potential of fenobam, and the safety profile of fenobam observed in small clinical studies [3; 4; 13; 16; 23; 24; 32] encouraged us to utilize fenobam to test the hypothesis that mGlu5 modulates nociceptive sensitization in humans.

In prior human studies, fenobam was administered to a limited number of subjects, and the disposition was not thoroughly investigated. We sought to evaluate the pharmacokinetics of a single dose of fenobam, and to then test the analgesic effects of fenobam in a human experimental pain model. The first study was a dose escalation protocol in adult healthy volunteers. The highest dose administered (150 mg) was the highest single dose reportedlywell tolerated in previous studies [3; 13; 24]. The second

study evaluated 150 mg oral fenobam analgesia in the validated heath/capsaicin model of cutaneous sensitization[9; 25].   Our hypothesis was that the area of cutaneous sensitization[25] that would develop under the effect of fenobam would be reduced in size compared to the baseline area (measured on the training day) across time-points, while, with placebo, the area would not differ from baseline. Therefore, we expected the areas of sensitization to be significantly different under the two conditions (fenobam/placebo). The primary outcome measure was suppression of the development of cutaneous hyperalgesia and allodynia around the area treated with heat/capsaicin. The outcome definition was size of the area of hypersensitivity around the area treated with heat/capsaicin quantified by cutaneous stimulation with foam brush strokes and von Frey filament. Heat/capsaicin application was timed for maximal sensitization to occur at the time of fenobam C_{max} , as determined in the first investigation. Secondary outcomes of interest were: Heat Pain Detection Thresholds (HPDT); pain scores after thermal stimulation of normal (untreated) skin, and side effects.

Given the reported anxiolytic properties of fenobam [13; 16; 23; 24; 26], we considered the possible implications of anxiolysis on pain modulation; therefore subject mood and affect changes were evaluated by a short version [19] of the brief Positive and Negative Affect Scale (PANAS)[30] and Brief State Anxiety Measure (BSAM)[2]. Moreover, because pre-clinical studies reported effects of mGlu5 modulation on working memory and cognitive performance[1; 14; 20; 27; 28],   subjects underwent the Letter and Number Sequencing Assessment (LNS), a subtest of the Wechsler Adult Intelligence Scale-Fourth Edition (WAIS-IV)[31] to evaluate working memory as a measure of cognitive function before and after administration of fenobam or placebo.

Methods

All studies were carried out in accordance with ethical principles of Good Clinical Practice and the Declaration of Helsinki and its guidelines, and approved by Washington University Institutional Review Board, following submission of an Investigational New Drug (IND) application to the FDA (IND#117,989). Subjects provided written informed consent. The investigations were registered on ClinicalTrials.gov NCT01806415 and NCT01981395. Fenobam was manufactured under GMP guidelines by Scynexis, Inc. (Durham, NC) as powder and then compounded in gelatin capsules with lactose monohydrate at Washington University in St. Louis investigational Pharmacy under strict adherence to USP 795.

Study Setting

Both studies ("PK Study" and "Hyperalgesia Study") were conducted in the Washington University Clinical Research Unit (CRU) and/or the Anesthesiology Human Studies Lab at Washington University.

PK Study. Pharmacokinetics and Side Effects of Oral Fenobam in Adult Healthy Volunteers

This was a randomized, double blind, single dose, parallel group, placebo controlled study to evaluate the pharmacokinetics and side effects of fenobam (Flow Chart - Fig.1A). The primary objective of this study was to obtain pharmacokinetic data after oral administration of 50, 100 and 150 mg of fenobam in groups of healthy individuals. A secondary objective was to compare the side effects of a single dose of 50 mg, 100 mg or 150 mg of fenobam to placebo.

Screening Session/Pre-study Period

Potential candidates were screened by the PI (LFC) and by the Senior Research Coordinator (KF) according to the inclusion and exclusion criteria. Inclusion criteria were age 18-50 yr, good general health with no remarkable medical conditions (e.g., history of liver, kidney, heart, or lung disease), and BMI < 33. Exclusion criteria were medication use (prescription or non-prescription medications, vitamins, herbals, dietary and mineral supplements and grapefruit products during or within 14 days prior to study participation; excluding contraceptives), history of addiction to drugs or alcohol (prior or present addiction or treatment for addiction), pregnancy or nursing, history of lactose intolerance, and smoking. Study participants were asked to abstain from drinking alcohol for 24 hours before the study, and to abstain from eating and drinking after midnight on the night before the study. An initial visit included patient health self-assessment, medical history and physical examination. Vital signs were then recorded (heart rate, respiratory rate, blood pressure, temperature and oxygen saturation). Each subject who qualified for entry into the study on the basis of inclusion/exclusion criteria, agreement with informed consent and pre-study evaluation was assigned the next available patient number. Thirty-two subjects were enrolled, with 8 subjects assigned to each group. In this Phase I, single oral dose pharmacokinetic and side effects study, a formal sample size calculation was not performed and sample size justification was based on prior experience with clinical pharmacokinetic investigations.

Randomization

After successful completion of the screening session, a computer generated randomization schedule prepared by the Washington University in St. Louis investigational Pharmacy assigned subjects to administration of either 50 mg, 100 mg, 150 mg of

fenobam ([1-(3-chlorophenyl)-3-(1-methyl-4-oxo-2-imidazolidinylidine) urea hydrate]) or placebo (lactose monohydrate 150 mg in gelatin capsules). Therefore each subject received a single dose of fenobam or placebo. Subjects and study personnel were blinded to drug/placebo by administration in identical gelatin capsules provided by the Investigational Pharmacy.

Study Period

Vital signs were recorded and a peripheral IV catheter was inserted in an arm for blood sampling. A baseline blood sample was used for a complete blood cell count and comprehensive metabolic panel. Fenobam or placebo was administered with a sip of water and venous blood samples were drawn before and after fenobam/placebo administration (at 0.5,1,2,3,4,5,6,10 and 24 hours). Blood samples were centrifuged at 3000 rpm for 10 minutes at room temperature. Plasma was then removed and frozen at -20^{o}C for later analysis. Vital signs were simultaneously recorded and subjects were queried for side effects (confusion, visual changes/blurred vision, dizziness, light-headedness, weakness, speech difficulties, abnormal cutaneous sensations, tingling, or numbness, nausea/vomiting, headache, metallic taste, hot flashes and any abnormal feelings). An additional blood sample was obtained on day 2 for a complete blood cell count and comprehensive metabolic panel.

One week after the study completion subjects were asked again to answer a questionnaire regarding any abnormal/unusual feeling they might be experiencing. The 1 week interview was conducted by phone.

Laboratory Analysis

Plasma fenobam was quantified using tandem mass spectrometry, as previously described by our laboratory[21]. The calibration range was 6 to 16,000 ng/ml.

Data Analysis

Descriptive statistics were generated for reported measures, and difference in distribution of baseline characteristics among groups administered fenobam 50mg, 100mg, 150mg or placebo were analyzed with the Kruskal Wallis H test using SPSS statistical software. The correlation between fenobam dose and number of side effects was investigated with Kendall's tau coefficient of rank correlation (tau_b). Proportions of patients with side effects in the different groups were compared using Fisher's exact test. For all analyses, careful attention was given to whether the data satisfied the distributional and model-specific assumptions of the procedures used.

C_{max}, t_{max} and AUC were determined based on the measurements collected and using the statistical software package Sigma Plot 13.0 (Systat Software, Inc., San Jose, USA); Spearman's correlation coefficient (r_s) was calculated to measure the association between dose of fenobam administered and C_{max} and AUC.

Hyperalgesia Study: Anti-Hyperalgesic Effect of a Single Dose of Fenobam on Heat/Capsaicin-Induced Cutaneous Hyperalgesia in Adult Healthy Volunteers

This study was designed as a randomized, double-blinded, placebo-controlled, two-way cross-over trial with 32 healthy volunteers who received either 150 mg fenobam or placebo (lactose monohydrate) and were then tested for cutaneous hypersensitivity using the heat/capsaicin model of cutaneous sensitization (Flow chart - Fig 1.B).

The study consisted of an initial screening/training session to evaluate subjects, establish eligibility and explain the study procedures, and two subsequent study sessions (placebo or fenobam, randomized order) that were conducted one week apart. During the two study sessions, blood samples were collected hourly for 7 hours after administration of fenobam, and fenobam plasma concentrations were then determined by tandem mass

spectrometry. Measures of hyperalgesia and hypersensitivity were taken at regular time points during each study session, and included the area of cutaneous sensitivity to calibrated von Frey filament and to brush stroke stimulation, and Heat Pain Detection Thresholds (HPDTs). Assessments of mood/affect and cognitive function also occurred at the same time points. A complete overview of the study timeline is shown in Fig 2.

Screening Session/Pre-study Period

Subjects who were potential candidates for the study were evaluated for study eligibility, according to the inclusion and exclusion criteria. These were the same as for the pharmacokinetic study, with the additional exclusions of anatomical malformation of an upper extremity, recent trauma or chronic lesions on either forearm, history of allergy or intolerance to capsaicin, and history of multiple drug allergies. An initial visit included a health self-assessment form, collection of medical history and physical examination. Vital signs were then recorded (heart rate, respiratory rate, blood pressure, temperature and oxygen saturation). Each subject who qualified for entry into the study on the basis of inclusion/exclusion criteria, agreement with informed consent and pre-study evaluation was assigned the next available patient number.

Study Period

Subjects participated in three sessions: one screening/training session and two study sessions, each scheduled approximately one week apart. The *training session* was held on the same day as the above mentioned screening session. In this session the enrolled subjects were given a demonstration of the actual study procedures to be conducted during the study sessions (see below). The duration of the subjects' participation was approximately four weeks, with the training session on day 1, the first study session within 14 days of the training session, and Session #2 approximately one week following Session

#1. During the training session, subjects experienced the heat/capsaicin sensitization procedure and measurements of cutaneous sensitization as described in detail below. Subjects were withdrawn from the study if their heat pain detection threshold was greater than 47°C at baseline or they failed to develop an area of measurable cutaneous sensitization following heat and capsaicin stimulation. The training session was intended to familiarize subjects with the experimental procedures so that communication between experimenter and subject about the subjective experiences of painful stimulation could be as precise as possible. No drug was administered during the training session.

During the study sessions subjects underwent a complete set of sensitization procedures; mapping of the area of sensitization; measurement of pain from thermal stimulation and heat pain detection thresholds at baseline, and then multiple times following rekindling of the sensitization via heat application.

Sample Size Estimates

The primary outcome measure was the size of the area of hyperalgesia and allodynia around the area sensitized by heat/capsaicin as quantified by cutaneous stimulation with foam brush strokes and a von Frey filament requiring 26 g of bending force. Preliminary data on the development of cutaneous sensitization with the heat/capsaicin model obtained from our group[5] enabled us to estimate the variability of the primary outcome measure, so that we could calculate the sample size sufficient to detect a 20% difference in the area of hyperalgesia in the two treatment conditions.

In the population of 15 subjects in the study referenced above[5], reproducing areas of skin sensitization to von Frey filament stimulation with the heat/capsaicin model in the same subjects 1 week apart, the within-day standard deviation of the area ranged from 26 (within day standard deviation of initial and final areas for the same subject on Session 2)

to 30 (within day standard deviation of initial and final areas for the same subject on Session 1). The standard deviation of the difference between the areas for the same subject between days (Session 1 vs Session 2) was 13.

Based on these standard deviations, we calculated the sample size needed to detect a 20% difference in the area of sensitization measured in the two treatment conditions (80% power, alpha = 0.05, two-sided). A total of 32 subjects were estimated to be sufficient for this two-treatment crossover study. To detect a 30% difference in size of area of hyperalgesia between treatments, 15 subjects would be sufficient.

Randomization

A computer-generated randomization schedule prepared by the Washington University in St. Louis investigational Pharmacy assigned subjects to fenobam 150 mg or placebo administration, and they were then crossed over to the other drug for the second session one week later.

Study Procedures and Measurements

Administration Protocol

The heat/capsaicin hyperalgesia model combines heat stimulation (heat ramps from 32°C to 45°C at a rate of 1°C per second and hold at 45°C for 5 min) applied to a 9 cm^2 area of skin on the forearm followed by topical low dose capsaicin (0.1% Capzacin-HP Cream) applied to the same area. The sensitization is then rekindled with subsequent applications of heat (40^o for 5 min) at 35-45 minutes intervals. This procedure generates temporary pain and the sensory changes associated with peripheral and central sensitization for up to 4 hours[25]. Thermal stimulations were applied in a precise and controlled manner using a Medoc Advanced Thermal Stimulator (Medoc, Israel and North Carolina, USA)

driving a 9 cm^2 thermode. The thermode was a computer-controlled Peltier device that warms the skin from 32°C to a safety cutoff of 52°C in 1 C/sec increments.

Measurement of Pain at Thermal Stimulation, Pain Thresholds and Areas of Hypersensitivity

The following methods were used to induce and quantify pain and sensitization: 1) the pain intensity and area (cm^2) of secondary hyperalgesia and allodynia induced by the heat/capsaicin model 2) heat pain detection threshold; 3) pain intensity produced by 1 minute 45°C thermal stimulation.

1) *Heat/Capsaicin Sensitization Procedure and Mapping an Area of Secondary Hypersensitivity*: Sensitization was established by heating a 9 cm^2 surface of the dominant forearm, and then applying 14 ml of 0.1% capsaicin cream as described above.   Subjects were asked to rate their pain on a Visual Analog Scale (VAS) at the start of the 30 min period and then for every 5 min until the cream was wiped off. At the end of the 30 min capsaicin application, measurements were performed to determine the areas of hypersensitivity and allodynia on the forearm. The borders of secondary mechanical allodynia and hyperalgesia were mapped using a 1 inch foam brush and a von Frey filament (26g bending force). Subjects were asked to close their eyes during these procedures. First, the brush was applied along four linear paths between the thermode outline and 1) the antecubital fossa, 2) the wrist joint, 3) the lateral aspect of the forearm in anatomical position, and 4) the medial aspect of the forearm. Stimulation started distant from the heated area and worked closer in 5.0 mm steps at 1sec intervals. Subjects were asked to say when the stimulation first became painful, and that location was marked. This procedure was then repeated with the von Frey filament. The area of hypersensitivity was calculated as the distance between the farthest points on the rostral/caudal axis multiplied

by the distance between the farthest points on the medial/lateral axis (in cm^2), then subtracting the heated area of the thermode (Fig. 3).

2) *Heat Pain Detection Threshold (HPDT):* Thresholds for heat pain detection were determined by using a thermal ramp protocol on a marked location on the volar surface of the forearm. The temperature applied through the Medoc thermode was increased from 32°C to the 52°C safety cutoff at 1°C/s. Subjects were requested to turn off the heated thermode by pressing a button at "the lowest temperature that they perceive as painful." Four thermal ramps were performed 10 seconds apart and the median value perceived as painful will be calculated. To avoid testing individuals whose pain threshold approached the safety cutoff, subjects with HPDTs greater than 47°C were excluded from the study.

3) *Pain During Thermal Stimulation (PTS*): Acute pain was induced by a 1 min 45°C heat stimulus on a marked location on the upper non-dominant arm (deltoid). Subjects were asked to rate their pain intensity during the 1 min heat stimulus continuously using an electronic Visual Analog Scale (VAS) ranging from 0 to 100 where 0 indicates "no pain sensation" and 100 indicates "the most intense pain imaginable."

Rekindling Procedures

On drug study days (Session1 and 2), hypersensitivity was maintained by rekindling the site of heat/capsaicin application. This was accomplished by re-stimulating the previously treated skin four times at approximately 40-45 min intervals, with the thermode increasing from 32°C to 40°C) at a rate of 1° per second and held at 40°C for 5 min. Subjects rated their pain on a continuous visual analogue scale (VAS) during rekindling.

Measurement/Evaluation of Mood and Affect Changes

Using a combination of Brief Positive and Negative Affect Scale (PANAS) and Brief State Anxiety Measure (BSAM)[19] we aimed to quantify changes from baseline in the subjects' mood and affect following administration of the drug and after sensitization.

Assessment of Cognitive Function

Changes in working memory (attention, concentration and mental control) were assessed with the Letter-Number Sequencing (LNS) test [31]. Changes in performance from baseline following administration of the drug and after sensitization were recorded.

Data analysis

Our primary outcome measure was the size of von Frey and brush areas of cutaneous sensitization to heat and capsaicin.

A mixed within–between subject linear model approach using SAS Proc Mixed procedure was used to analyze the data. The mixed model used restricted maximum likelihood estimation for linear models with degrees of freedom adjusted using Kenward-Roger procedure. This model allowed controlling for potential confounders. Type III tests of fixed effects were used to evaluate the main effects of treatment group, time and interaction of treatment group with time.

The possibility of a carryover effect was explored by comparing baseline values between the two treatment Sessions in the same group (Wilcoxon signed ranks test), as well as by testing the sequence of randomization in the mixed model analysis. Since we found that there was a significant carryover effect, we analyzed separately data from Session 1 of the study. Independent samples t-test or non-parametric equivalent Wilcoxon rank sum test were used to explore differences in distribution of continuous level

characteristics and baseline measures between subjects randomized to fenobam or

Placebo in Session 1. In essence, the mixed model approach allowed us to explore the

difference in size of von Frey area of sensitization through different time points and

compare area sizes between the two treatment groups. The same analytical approach was

also used for brush area and HPDT measures.

Results

1. Pharmacokinetic Study. Pharmacokinetics and Side Effects of the mGlu5 Negative Allosteric Modulator Fenobam in Adult Healthy Volunteers

A total of 32 adult healthy volunteers (Table 1) were randomized to four groups to receive oral administration of placebo or fenobam (50, 100 or 150 mg). Fenobam plasma concentration curves after 50,100, and 150 mg oral doses are shown in Fig.4. C_{max}, t_{max} and AUC median, minimum and maximum values for the three groups receiving fenobam are summarized in Table 2.

After oral administration of 50 mg fenobam, t_{max} was between 2 and 4 hours. C_{max} was extremely variable, between 0 and 48.4 ng/ml. T_{max} for the 100mg and 150mg oral doses was between 2 and 6 hours; for the 100 mg dose C_{max} were between 0.5 and 3.7 ng/ml, and for the 150 mg dose C_{max} were between 0.1 and 32.2 ng/ml. There was no significant association between fenobam dose administered and C_{max} (r_s = 0.236) or AUC. (r_s = 0.295).

Safety and Tolerability

Fenobam was well tolerated up to the highest oral dose of 150 mg. Adverse events included headache, nausea, metallic or weird taste, and fatigue. All adverse events were described as "mild" by the subjects. Adverse events associated with either fenobam or placebo administration are presented in Table 3. No serious adverse events occurred in any subject. We found a weak negative correlation between fenobam dose and presence of side effects (Kendall's tau_b -0.48; p=0.05) and the number of subjects with side effects was not significantly different between fenobam and placebo (Fisher Exact test, p= 0.82).

2. Hyperalgesia Study. Effects of 150 mg Orally Administered Fenobam on the Development of Cutaneous Sensitization in the Heat Capsaicin Test.

Demographic information and baseline characteristics of subjects who received the heat/capsaicin sensitization procedures are presented in Table 4. Disposition of oral fenobam 150 mg in this cohort is shown in Fig 5.

Anti-hyperalgesic Effects

The primary outcome measure was size of areas of cutaneous sensitization to brush and von Frey filament. Taking into account the crossover design of the study, the presence of a carryover effect was explored by comparing baseline (M-test) area measures – i.e. immediately following sensitization with heat-capsaicin - between the 2 treatment sessions, as well as by testing the sequence of randomization in the mixed model analysis (SAS Proc Mixed procedure). We found a significant (p=0.001 for von Frey areas and p<0.001 for brush areas) and consistent (present in 26/32 subjects) reduction in von Frey and brush area measures at the first measurement in Session 2 compared to the same time-point (M-test) in Session 1, independent of treatment. Both von Frey and brush areas were reduced by 20% in Session 2 (fenobam) compared to Session 1 in the group that had received placebo in Session 1. In the group that had received fenobam in Session 1, von Frey areas in Session 2 (placebo) were reduced by 25%, and brush areas were reduced by approximately 50%. In light of these findings that we interpreted as the presence of a carryover effect of the pain testing paradigm, we focused our analysis on Session 1, with 17 subjects randomized to receive fenobam and 15 subjects randomized to receive placebo.

In Session 1, the overall (unadjusted) effect of fenobam on von Frey area compared to placebo was an average reduction in area size by 18.1 cm^2 (95% C.I. -4.6 – 40.8). The

overall effect on brush area was an average reduction by 13.9 cm^2 (-2.4-30.3). HPDTs (unadjusted results) were on average 0.9 ^{o}C higher in patients receiving fenobam compared to placebo (C.I. -0.3-2.0).

After controlling for area of sensitization to von Frey filament on the training day and alcohol use, which were found to be significant confounders of the drug effect, the change of von Frey area through different time points (M-test = Measurements taken immediately post sensitization, through M4= measurements taken after the 4th rekindling procedure; time-points illustrated in Fig.2) was significant, with the lowest values noted at time M4. These area reductions did not differ significantly between the placebo and fenobam groups. The average (all subjects) reduction of von Frey area at M4 compared to the area measured at M-test was 44.0 cm^2 (95% CI: 28.2 to 59.8) in the placebo group, and 35.4 cm^2 (95% CI: 20.4-50.4) in the fenobam group. Overall, subjects treated with fenobam had a von Frey area 0.76 cm^2 smaller than subjects treated with placebo; however this difference was not statistically significant (95% CI:-16.03 to 17.56).

A similar pattern was also observed in brush area measurements. After controlling for area of sensitization on training day and forearm circumference (confounding variables of the drug effect in this model), the mean reduction of brush area at M4 compared to M-test was 46.9 cm2 (95% CI: 34.3 to 59.4) in the placebo group, and 44 cm2 (95% CI 32.0-55.9) in the fenobam group. Overall, subjects treated with fenobam had a brush area 7.5 cm^2 smaller than placebo group with no statistical significance (95% CI:-7.1 to 22.1).

When von Frey and the brush areas measured immediately post-sensitization, at the single time point closest to the C$_{max}$ of fenobam (= M-test on Study Day, Fig.2) were compared with areas obtained at M-test on the training day within the same group of subjects (but without fenobam or placebo administration), both areas were significantly

reduced in size (median von Frey areas 74.6cm^2 vs 90.1cm^2; brush areas 54.7cm^2 vs 72.65 cm2; p=0.025 and 0.028 respectively), while the areas measured at M-test of Session 1 in the placebo group were not different from training day (median von Frey areas 112.4cm^2 vs 115.8cm^2; brush areas 66.8 cm^2 vs 77.9;p = 0.069 and p = 0.1 (Fig.6A and 6B).

After controlling for HPDT at baseline (Session 1), over all time points, mean HPDTs in the fenobam group were 0.42 ^{o}C higher than the placebo group; however this difference was not significant (95% CI -0.38 - 1.23)). At the M-test time-point there was no difference in HPDTs between the fenobam and placebo group.

Anti-nociceptive Effect

Overall after controlling for PTS at S1 baseline, there was no significant difference in maximum VAS (PTS MAX) scores recorded between the 2 groups and across all time-points (data not shown).

PNAS/BSAM and LNS

In a multi-level model to determine whether the mood of the subjects and their anxiety level varied with fenobam plasma concentration over time we could not observe any statistically significant effect of fenobam compared to placebo (not reported). Measures of cognitive function (attention, concentration and mental control) were also not significantly different between groups (Fig.7).

Discussion

Our study of the pharmacokinetic properties and side effect profile of fenobam after oral administration in human subjects showed highly variable disposition and t_{max} between 2 and 6 hours with marked inter-individual differences. These observations are similar to prior findings on the variability of fenobam disposition: in a pilot study on a limited number of subjects with Fragile-X syndrome (FXS), Berry-Kravis et al.[3] found that fenobam plasma concentrations were dose-dependent but variable, with mean C_{max} of 39.7 ±18.4 ng/ml at 180 min following oral administration of fenobam150 mg. In the Berry-Kravis study, three healthy volunteers (control group) had a mean peak plasma concentration of fenobam that did not differ from that seen in the FXS patients. The timing to mean peak plasma concentrations was similar in FXS patients and in healthy controls. Neither fenobam dose nor concentrations after 60 or 120 min correlated with improvement of the clinical outcome of interest (behavioral response).

In contrast to results obtained in pre-clinical studies by our group[21; 22], we observed a transient reduction in the area of hypersensitivity at the time of peak fenobam plasma concentration, but did not observe any persistent anti-hyperalgesic or anti-nociceptive effect of fenobam compared to placebo in this cutaneous sensitization experimental pain model. The relatively unimpressive effect of fenobam in this human study compared to the rodent studies may be due to multiple factors, including the unknown relationship between plasma levels and receptor occupancy, and accordingly the potential for insufficient fenobam dosing in the present study, or of course simply, reduced drug effect in humans relative to rodents.

In an attempt to reproduce in human the same conditions in which we had observed an analgesic effect of fenobam in rodents, we utilized an experimental model that constituted the direct translation of rodent experimental pain models that we had used [21;

22]. The single dose of fenobam that we administered in our human study was approximately equivalent to the dose that we administered in mice with analgesic effect (allometric scaling suggests that 30 mg/kg in mice corresponds to 2.4 mg/kg in human, with a total dose of 150 mg being the calculated dose for an ideal body weight of 60 kg) (**ref:** "Guidance for Industry" : "Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers" Published by the U.S. Department of Health and Human Services; Food and Drug Administration Center for Drug Evaluation and Research (CDER); July 2005).

We observed a reduction of von Frey area through different time points (M-test = immediately post sensitization, through M4= after the 4th rekindling procedure), with the lowest values noted at time M4. These area reductions did not differ significantly between the placebo and fenobam groups. This finding was consistent with previously reported progressive fading of areas of hyperalgesia over time[5]. The presence of a strong carryover effect of the sensitization model from Session 1 to Session 2, and possibly unexplained interactions between the carryover effects of both the sensitization model and of the drug, prevented us from taking advantage of the original cross-over design.

Before starting the study, we were aware of potential issues with the heat/capsaicin model, its reproducibility and presence of a carryover effect causing areas of sensitization in Session 2 to be smaller than Session 1 in a crossover design[5]. However, based on our pre-clinical data, we had anticipated that the magnitude of the analgesic effect of fenobam would be sufficiently large to detect despite the carryover effect of the model. A complete unknown here is the relative receptor occupancy by fenobam in the rodent and human studies. Additionally, despite the purported[9; 10; 25] adequacy of the washout period for fenobam demonstrated by PK analysis, we cannot exclude the possibility that an effect of fenobam on cutaneous sensitization in Session 1 could have affected the sensitization

process in Session 2, one week later, when the same subjects were receiving placebo. Our inability to exclude such an effect is based on the observation that areas in Sessions 2 have been consistently found to be smaller than Session 1. A possible explanation of this phenomenon is that cutaneous sensitization one week before may affect the development of a "re-sensitization" the following week. If this is true, any effect of fenobam on the sensitization process in Session 1 may have an indirect, unpredictable influence on the following week's process. A parallel comparison of two groups receiving fenobam in the two consecutive sessions or placebo in both could have provided more information on the effect of the drug versus the effect of the repeated exposure to heat and capsaicin on skin sensitization. Unfortunately we are not able to administer repeated doses of fenobam to the same subjects due to regulatory constraints.

Once a strong carryover effect was detected, the decision was made to examine data from Session 1 separately, an approach that has precedent [8; 18]. However this approach decreases the statistical power of the study and has been deemed "at risk of bias"[8; 11; 12]. We recognize this as a significant limitation of our study, and we focused the presentation and interpretation of our findings on 95% confidence intervals.

With the limitations noted above, in the heat/capsaicin cutaneous sensitization model, contrary to our hypothesis that the size of the area of hypersensitivity would remain different in the two treatment groups across subsequent time-points as fenobam concentration was decreasing, we found no difference between fenobam and placebo in the overall analgesic effect. However, our study design also took into account the possibility that there might be only a narrow window around the time of C_{max} to detect an effect of fenobam in our constrained experimental conditions, and that this effect would not last throughout the sensitization procedures. To maximize the sensitivity of the clinical model, we performed the preliminary PK study and then planned the skin sensitization

procedures so that initial sensitization would coincide approximately with peak plasma fenobam concentration.

Plasma fenobam analysis confirmed that initial heat and capsaicin sensitization did coincide with peak plasma fenobam concentrations.   By analyzing sensitization data from the single time-point closest to C_{max} of fenobam (M-test) we found that both von Frey and brush areas were significantly reduced in size compared to areas obtained at baseline on the training day in the same subjects, when no drug was administered, while areas measured at M-test in the placebo group were not statistically different from training day. While this single time-point finding was observed both in the von Frey and the brush areas, possible explanation for the absence of a persistent anti-hyperalgesic effect of fenobam might be the rapid elimination of fenobam compared to the duration of cutaneous sensitization.

Availability of validated models to reliably create experimental hyperalgesia in healthy volunteers is limited, but - from a safety standpoint -we considered necessary to test fenobam in human healthy volunteers first, before conducting a prospective randomized clinical trial in patients affected by a clinical pain condition. With these considerations in mind, we relied on the prior extensive use of the model, and our direct knowledge and experience with the heat/capsaicin sensitization procedures. However, our limited and constrained experimental conditions might have impaired our ability to detect a persistent effect on the areas of sensitization over time. Obviously, it is also possible that, despite the encouraging pre-clinical data, fenobam, and possibly mGlu5 antagonists in general, may not have robust analgesic effects in human. In our opinion, a clear answer on this issue requires additional studies where compounds with improved pharmacokinetic properties, and known receptor engagement, are utilized.

In conclusion, in our limited experimental conditions with human healthy volunteers, we did not observe any clinically or statistically significant persistent analgesic (anti-hyperalgesic) effect of fenobam compared to placebo over the study time course. Fenobam, administered orally to human healthy volunteers at a single dose of 50, 100 or 150 mg, showed highly variable disposition and t_{max} between 2 and 6 hours with marked inter-individual differences.

Given the significant limitations introduced by the highly variable plasma exposure of fenobam, we feel that attempts to assess the potential utility of mGlu5 modulation for pain necessitate the use of a compound with improved pharmacokinetics and known target engagement. Prospective randomized clinical trials with such an improved molecule are needed to clarify the role of mGlu5 modulation in the development and maintenance of acute and chronic pain conditions in human.

Acknowledgements

The authors wish to thank Dr. Eric Lenze for his contribution as a consultant on the evaluation and measurement of mood and affect changes and measures of cognitive function and Alicia Neiner for the quantification of fenobam in plasma.

None of the authors has reported any conflict of interest, other than the financial support as reported below.

Financial Support

This work was supported by the National Institutes of Health - National Institute of Neurological Disorders and Stroke (NS48602 to R.W.G.) and by a Grant of the Barnes Jewish Hospital Foundation and Washington University Institute of Clinical and Translational Sciences (ID# CTSA310 to R.W.G and L.F.C.).

References

[1] Ballard TM, Woolley ML, Prinssen E, Huwyler J, Porter R, Spooren W. The effect of the mGlu5 receptor antagonist MPEP in rodent tests of anxiety and cognition: a comparison. Psychopharmacology (Berl) 2005;179(1):218-229.

[2] Berg CZ, Shapiro N, Chambless DL, Ahrens AH. Are emotions frightening? II: An analogue study of fear of emotion, interpersonal conflict, and panic onset. Behav Res Ther 1998;36(1):3-15.

[3] Berry-Kravis E, Hessl D, Coffey S, Hervey C, Schneider A, Yuhas J, Hutchison J, Snape M, Tranfaglia M, Nguyen DV, Hagerman R. A pilot open label, single dose trial of fenobam in adults with fragile X syndrome. J Med Genet 2009;46(4):266-271.

[4] Bhave G, Karim F, Carlton SM, Gereau RWt. Peripheral group I metabotropic glutamate receptors modulate nociception in mice. Nat Neurosci 2001;4(4):417-423.

[5] Cavallone LF, Frey K, Montana MC, Joyal J, Regina KJ, Petersen KL, Gereau RWt. Reproducibility of the heat/capsaicin skin sensitization model in healthy volunteers. J Pain Res 2013;6:771-784.

[6] Chiechio S. Modulation of Chronic Pain by Metabotropic Glutamate Receptors. Advances in pharmacology (San Diego, Calif) 2016;75:63-89.

[7] Crock LW, Kolber BJ, Morgan CD, Sadler KE, Vogt SK, Bruchas MR, Gereau RWt. Central amygdala metabotropic glutamate receptor 5 in the modulation of visceral pain. The Journal of neuroscience : the official journal of the Society for Neuroscience 2012;32(41):14217-14226.

[8] Curtin F, Elbourne D, Altman DG. Meta-analysis combining parallel and cross-over clinical trials. III: The issue of carry-over. Stat Med 2002;21(15):2161-2173.

[9] Dirks J, Petersen KL, Dahl JB. The heat/capsaicin sensitization model: a methodologic study. The journal of pain : official journal of the American Pain Society 2003;4(3):122-128.

[10] Dirks J, Petersen KL, Rowbotham MC, Dahl JB. Gabapentin suppresses cutaneous hyperalgesia following heat-capsaicin sensitization. Anesthesiology 2002;97(1):102-107.

[11] Freeman PR. The performance of the two-stage analysis of two-treatment, two-period crossover trials. Stat Med 1989;8(12):1421-1432.

[12] Higgins JPT, Green, S. . Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. , 2011.

[13] Itil T.M. SBA, Huque M, Mukhopadhyay S, Blasucci D, Nq K.T., Ciccone P.E. The clinical and quantitative EEG effects and plasma levels of fenbam (McN-3377) in subjects with anxiety: an open rising dose tolerance and efficacy study. Curr Ther Res 1978 24:708–724.

[14] Jacob W, Gravius A, Pietraszek M, Nagel J, Belozertseva I, Shekunova E, Malyshkin A, Greco S, Barberi C, Danysz W. The anxiolytic and analgesic properties of fenobam, a potent mGlu5 receptor antagonist, in relation to the impairment of learning. Neuropharmacology 2009;57(2):97-108.

[15] Kolber BJ. mGluRs head to toe in pain. Progress in molecular biology and translational science 2015;131:281-324.

[16] Lapierre YD, Oyewumi,L.K. Fenobam: another anxiolytic? Curr Ther Res 1982(31):95-101.

[17] Lax NC, George DC, Ignatz C, Kolber BJ. The mGluR5 antagonist fenobam induces analgesic conditioned place preference in mice with spared nerve injury. PloS one 2014;9(7):e103524.

[18] Lehmacher W. Analysis of the crossover design in the presence of residual effects. Stat Med 1991;10(6):891-899.

[19] Mackinnon A, Jorm, A. F., Christensen, H., Korten, A. E., Jacomb, P. A. and Rodgers, B. A short form of the Positive and Negative Affect Schedule: Evaluation of factorial validity and invariance across demographic variables in a community sample. Personality and Individual Differences 1999;27(3):405-416.

[20] Mikulecka A, Mares P. Effects of mGluR5 and mGluR1 antagonists on anxiety-like behavior and learning in developing rats. Behav Brain Res 2009;204(1):133-139.

[21] Montana MC, Cavallone LF, Stubbert KK, Stefanescu AD, Kharasch ED, Gereau RWt. The metabotropic glutamate receptor subtype 5 antagonist fenobam is analgesic and has improved in vivo selectivity

compared with the prototypical antagonist 2-methyl-6-(phenylethynyl)-pyridine. J Pharmacol Exp Ther 2009;330(3):834-843.

[22] Montana MC, Conrardy BA, Cavallone LF, Kolber BJ, Rao LK, Greco SC, Gereau RWt. Metabotropic glutamate receptor 5 antagonism with fenobam: examination of analgesic tolerance and side effect profile in mice. Anesthesiology 2011;115(6):1239-1250.

[23] Pecknold J.C. MDJ, and Appeltauer L. Fenobam in anxious outpatients. Curr Ther Res 1980(27):119-123.

[24] Pecknold J.C. MDJ, Appeltauer L., Wrzesinski L., and Allan T. J. Treatment of anxiety using fenobam (a nonbenzodiazepine) in a double-blind standard (diazepam) placebo-controlled study. Clin Psychopharmacol 1982(2):129–133.

[25] Petersen KL, Rowbotham MC. A new human experimental pain model: the heat/capsaicin sensitization model. Neuroreport 1999;10(7):1511-1516.

[26] Porter RH, Jaeschke G, Spooren W, Ballard TM, Buttelmann B, Kolczewski S, Peters JU, Prinssen E, Wichmann J, Vieira E, Muhlemann A, Gatti S, Mutel V, Malherbe P. Fenobam: a clinically validated nonbenzodiazepine anxiolytic is a potent, selective, and noncompetitive mGlu5 receptor antagonist with inverse agonist activity. J Pharmacol Exp Ther 2005;315(2):711-721.

[27] Quarta D, Naylor CG, Morris HV, Patel S, Genn RF, Stolerman IP. Different effects of ionotropic and metabotropic glutamate receptor antagonists on attention and the attentional properties of nicotine. Neuropharmacology 2007;53(3):421-430.

[28] Semenova S, Markou A. The effects of the mGluR5 antagonist MPEP and the mGluR2/3 antagonist LY341495 on rats' performance in the 5-choice serial reaction time task. Neuropharmacology 2007;52(3):863-872.

[29] Walker K, Reeve A, Bowes M, Winter J, Wotherspoon G, Davis A, Schmid P, Gasparini F, Kuhn R, Urban L. mGlu5 receptors and nociceptive function II. mGlu5 receptors functionally expressed on peripheral sensory neurones mediate inflammatory hyperalgesia. Neuropharmacology 2001;40(1):10-19.

[30] Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. J Pers Soc Psychol 1988;54(6):1063-1070.

[31] Wechsler DA. Wechsler Adult Intelligence Scale (4th ed.). SanAntonio, TX: SanAntonio, TX: Psychological Corporation., 2008.

[32] Zhu CZ, Hsieh G, Ei-Kouhen O, Wilson SG, Mikusa JP, Hollingsworth PR, Chang R, Moreland RB, Brioni J, Decker MW, Honore P. Role of central and peripheral mGluR5 receptors in post-operative pain in rats. Pain 2005;114(1-2):195-202.

A   **CONSORT 2010 Flow Diagram: PK Study**

B   **CONSORT 2010 Flow Diagram: Hyperalgesia Study**

Fig. 1 CONSORT Flow Diagrams: **A**. Study of the pharmacokinetics of fenobam ("PK study"); **B.** Study of the analgesic effects of fenobam on the heat/capsaicin-induced cutaneous hyperalgesia in adult healthy volunteers ("Hyperalgesia study").

A

Timeline of Screening/Training Day Procedures

B

Timeline of Study Day Procedures

Fig. 2 Timeline of the procedures on Screening/Trinaing Day (A) and Study Day (B). In each timeline, time points (min) are in relation to the "time 0" of start of the heat/capsaicin sensitization procedure; negative time points mark the pre-sensitization procedures and positive time points delineate the post-sensitization phase. **(A)** M = measurements; vF = von Frey; HPDTs= heat pain detection thresholds; PTS = pain at thermal stimulation. **(B)** PANAS = positive and negative affect scale; BSAM = Brief State Anxiety Measure; LNS = Letter andNumber Sequencing; RK =rekindling of sensitization.

Fig. 3 Measurement of cutaneous hypersensitivity after heat and capsaicin sensitization. (A)
Thermode outline on a subject's forearm and markings indicating points of change in sensation on the skin
from "non-painful" to "painful" at foam brush (X) and von Frey filament (II) stimulation. The square marked
with a broken line (- -) inside the thermode's outline represents the heated surface of the thermode which
measured 9 cm2. **(B)** Stimulation with brush strokes is shown as performed along a linear path between the
wrist joint and the thermode outline. **(C)** Measurement of an area of hypersensitivity (brush area shown
here): [(ab) x (cd)] – 9 cm^2.

Fig. 4 PK Study: Fenobam plasma concentration curves after 50 mg,100 mg, and 150 mg oral doses.
Curves of individual subjects are shown in **A, C, E**: each symobl/line combination represents a different subject for each group. Median concentrations for each dosage are shown in **B, D, F**.

Fig. 5 Hyperalgesia Study: fenobam plasma concentration following 150 mg oral dose. Median values (Min-Max) at different time points in relation to the start of the skin sensitization procedures (time 0) are shown.

Fig. 6 von Frey (A) and brush (B) areas of cutaneous hypersensitivity measured immediately after sensitization (M-test time-point) on Session 1 (S1) compared to areas on Training Day. Median; minimum; maximum values of areas; lower and upper quartiles are shown. The difference between median **Training Day** areas (no treatment) and median areas on **Session 1** is significant when fenobam is administered [*p = 0.025 (von Frey, panel A); *p = 0.028 (brush, panel B)] but not when placebo is administered.

Fig. 7 Mean Letter and Number Sequencing Assessment scores of subjects receiving fenobam or placebo in Session1. In this test subjects are instructed to sort and repeat progressively longer sequences of letters and numbers. Both groups show a mild improvement in scores over time that could be explained by a learning effect with the repetition of letter and number sequences. No significant difference is detected between the two groups.

	All subjects N=32	Placebo (n=8) Median (Min-Max)	Fenobam 50 mg (n=8) Median (Min-Max)	Fenobam 100 mg (n=8) Median (Min-Max)	Fenobam 150 mg (n=8) Median (Min-Max)
Sex					
Female	18 (56%)	7 (87.5%)	6 (75%)	2 (25%)	3 (37.5%)
Male	14 (44%)	1 (12.5%)	2 (25%)	6 (75%)	5 (62.5%)
Age	28 (20-49)	29 (20-49)	30 (21-35)	24 (21-46)	29 (21-40)
Height (cm)	170 (157-198)	170 (157-175)	169 (163-180)	176.5 (165-185)	174 (160-198)
Weight	77 (53-105)	82 (66-95)	71 (54-90)	77 (54-98)	78 (53-105)
BMI	26 (19-33)	28 (22-33)	25 (20-29)	25 (19-30)	25 (20-30)

Table 1. PK Study: demographic information and baseline characteristics.

Fenobam Dose	C_{max} (ng/ml)	t_{max} (hrs)	AUC ng*h/ml
50 mg	1.4 (0;48.4)	2 (2;4)	67.8 (0; 2405)
100 mg	2.9 (0.5;3.7)	3 (2;5)	167.4 (76.8;344.4)
150 mg	3.6 (0.1;32.2)	4 (2;6)	183.6 (10.8; 3862.8)

Table 2. C_{max}, t_{max} and AUC value medians (min; max) for the three groups of subject receiving 50 mg, 100 mg or 150 mg of fenobam.

Reported Side Effect	Placebo (n=40) (%)	Fenobam 50 mg (n=8) (%)	Fenobam 100 mg (n=8) (%)	Fenobam 150 mg (n=40) (%)
Headache	4 (10)	4 (50)	2 (25)	4 (10)
Nausea	3 (7.5)	1 (12.5)	1 (12.5)	0
Metallic/weird taste	1 (2.5)	1 (12.5)	1 (12.5)	0
Fatigue	1 (2.5)	1 (12.5)	0	0
Drowsiness	11 (27.5)	0	0	10 (25)
Lack of Concentration	2 (5)	0	0	0
Tingling	0	0	0	1 (2.5)
Dry mouth	3 (7.5)	0	0	1 (2.5)

Table 3. Side Effects in subjects receiving placebo or different doses of fenobam.
Rate of side effects in the groups **"Placebo"** and **"Fenobam 150 mg"** refer to subjects from both the PK study (8 subjects in each group) and hyperalgesia study (32 subjects receiving fenobam or placebo).

Baseline characteristics (on training day)	Placebo (n=15) Median (Min-Max)	Fenobam (n=17) Median (Min-Max)
Age (y)	28 (22-34)	29(19-45)
Height(cm)	173(157-191)	170(157-188)
Weight (kg)	71(59-105)	79(65-90)
BMI	26 (20-32)	27(21-33)
Coffee cups	2(1-4)	1(0.5-3)
Forearm circumf. (cm)	26(23-33)	26(23-30)
HPDT(^{o}C)pre-sensitization	43(38-47)	43(37-46)
HPDT(^{o}C)post-sensitization	37(35-38)	37(35-39)
Brush Area (cm^2)	87(54-152)	82(46-109)
vF area (cm^2)	125(85-215)	99(90-133)
	n (%)	n (%)
Sex		
Female	8(53%)	11(65%)
Male	7(47%)	6(35%)
Daily Caffeine intake		
Yes	10(67%)	11(65%)
No	5(33%)	6(35%)
Alcohol in last 6 months		
Yes	14(93%)	15(88%)
No	1(7%)	2(12%)
Dominant Arm		
Left	4(27%)	2(12%)
Right	11(73%)	15(88%)

Table 4. Hyperlagesia Study: demographic information and baseline characteristics of subjects receiving placebo or fenobam in Session 1.
HPDT = Heat Pain Detection Threshold.

A　　　　　　　　CONSORT 2010 Flow Diagram: PK Study

B　　　　　　　　CONSORT 2010 Flow Diagram: Hyperalgesia Study

Fig. 1

A

Timeline of Screening/Training Day Procedures

B

Timeline of Study Day Procedures

Fig. 2

Fig. 3

Fig. 4

Fig. 5

A   **von Frey**

B   **Brush**

Fig. 6

Fig. 7

Figure Legends

Fig. 1 CONSORT Flow Diagrams: **A**. Study of the pharmacokinetics of fenobam ("PK study"); **B.** Study of the analgesic effects of fenobam on the heat/capsaicin-induced cutaneous hyperalgesia in adult healthy volunteers ("Hyperalgesia study").

Fig. 2 Timeline of the procedures on Screening/Trinaing Day (A) and Study Day (B). In each timeline, time points (min) are in relation to the "time 0" of start of the heat/capsaicin sensitization procedure; negative time points mark the pre-sensitization procedures and positive time points delineate the post-sensitization phase. **(A)** M = measurements; vF = von Frey; HPDTs= heat pain detection thresholds; PTS = pain at thermal stimulation. **(B)** PANAS = positive and negative affect scale; BSAM = Brief State Anxiety Measure; LNS = Letter andNumber Sequencing; RK =rekindling of sensitization.

Fig. 3 Measurement of cutaneous hypersensitivity after heat and capsaicin sensitization. (A) Thermode outline on a subject's forearm and markings indicating points of change in sensation on the skin from "non-painful" to "painful" at foam brush (X) and von Frey filament (II) stimulation. The square marked with a broken line (- -) inside the thermode's outline represents the heated surface of the thermode which measured 9 cm2. **(B)** Stimulation with brush strokes is shown as performed along a linear path between the wrist joint and the thermode outline. **(C)** Measurement of an area of hypersensitivity (brush area shown here): $[(ab) \times (cd)] - 9 \text{ cm}^2$.

Fig. 4 PK Study: Fenobam plasma concentration curves after 50 mg,100 mg, and 150 mg oral doses. Curves of individual subjects are shown in **A, C, E**: each symobl/line combination represents a different subject for each group. Median concentrations for each dosage are shown in **B, D, F**.

Fig. 5 Hyperalgesia Study: fenobam plasma concentration following 150 mg oral dose. Median values (Min-Max) at different time points in relation to the start of the skin sensitization procedures (time 0) are shown.

Fig. 6 von Frey (A) and brush (B) areas of cutaneous hypersensitivity   measured immediately after sensitization (M-test time-point) on Session 1 (S1) compared to areas on Training Day. Median; minimum; maximum values of areas; lower and upper quartiles are shown. The difference between median **Training Day** areas (no treatment) and median areas on **Session 1** is significant when fenobam is administered [*p = 0.025 (von Frey, panel A); *p = 0.028 (brush, panel B)] but not when placebo is administered.

Fig. 7 Mean Letter and Number Sequencing Assessment scores of subjects receiving fenobam or placebo in Session1. In this test subjects are instructed to sort and repeat progressively longer sequences of letters and numbers. Both groups show a mild improvement in scores over time that could be explained by a learning effect with the repetition of letter and number sequences. No significant difference is detected between the two groups.

	All subjects N=32	Placebo (n=8) Median (Min-Max)	Fenobam 50 mg (n=8) Median (Min-Max)	Fenobam 100 mg (n=8) Median (Min-Max)	Fenobam 150 mg (n=8) Median (Min-Max)
Sex					
Female	18 (56%)	7 (87.5%)	6 (75%)	2 (25%)	3 (37.5%)
Male	14 (44%)	1 (12.5%)	2 (25%)	6 (75%)	5 (62.5%)
Age	28 (20-49)	29 (20-49)	30 (21-35)	24 (21-46)	29 (21-40)
Height (cm)	170 (157-198)	170 (157-175)	169 (163-180)	176.5 (165-185)	174 (160-198)
Weight	77 (53-105)	82 (66-95)	71 (54-90)	77 (54-98)	78 (53-105)
BMI	26 (19-33)	28 (22-33)	25 (20-29)	25 (19-30)	25 (20-30)

Table 1. PK Study: demographic information and baseline characteristics.

Fenobam Dose	C_{max} (ng/ml)	t_{max} (hrs)	AUC ng*h/ml
50 mg	1.4 (0;48.4)	2 (2;4)	67.8 (0; 2405)
100 mg	2.9 (0.5;3.7)	3 (2;5)	167.4 (76.8;344.4)
150 mg	3.6 (0.1;32.2)	4 (2;6)	183.6 (10.8; 3862.8)

Table 2. C_{max}, t_{max} and AUC value medians (min; max) for the three groups of subject receiving 50 mg, 100 mg or 150 mg of fenobam.

Reported Side Effect	Placebo (n=40) (%)	Fenobam 50 mg (n=8) (%)	Fenobam 100 mg (n=8) (%)	Fenobam 150 mg (n=40) (%)
Headache	4 (10)	4 (50)	2 (25)	4 (10)
Nausea	3 (7.5)	1 (12.5)	1 (12.5)	0
Metallic/weird taste	1 (2.5)	1 (12.5)	1 (12.5)	0
Fatigue	1 (2.5)	1 (12.5)	0	0
Drowsiness	11 (27.5)	0	0	10 (25)
Lack of Concentration	2 (5)	0	0	0
Tingling	0	0	0	1 (2.5)
Dry mouth	3 (7.5)	0	0	1 (2.5)

Table 3. Side Effects in subjects receiving placebo or different doses of fenobam. Rate of side effects in the groups **"Placebo"** and **"Fenobam 150 mg"** refer to subjects from both the PK study (8 subjects in each group) and hyperalgesia study (32 subjects receiving fenobam or placebo).

Baseline characteristics (on training day)	Placebo (n=15) Median (Min-Max)	Fenobam (n=17) Median (Min-Max)
Age (y)	28 (22-34)	29(19-45)
Height(cm)	173(157-191)	170(157-188)
Weight (kg)	71(59-105)	79(65-90)
BMI	26 (20-32)	27(21-33)
Coffee cups	2(1-4)	1(0.5-3)
Forearm circumf. (cm)	26(23-33)	26(23-30)
HPDT(^{o}C)pre-sensitization	43(38-47)	43(37-46)
HPDT(^{o}C)post-sensitization	37(35-38)	37(35-39)
Brush Area (cm^2)	87(54-152)	82(46-109)
vF area (cm^2)	125(85-215)	99(90-133)
	n (%)	n (%)
Sex		
Female	8(53%)	11(65%)
Male	7(47%)	6(35%)
Daily Caffeine intake		
Yes	10(67%)	11(65%)
No	5(33%)	6(35%)
Alcohol in last 6 months		
Yes	14(93%)	15(88%)
No	1(7%)	2(12%)
Dominant Arm		
Left	4(27%)	2(12%)
Right	11(73%)	15(88%)

Table 4. Hyperlagesia Study: demographic information and baseline characteristics of subjects receiving placebo or fenobam in Session 1.

HPDT = Heat Pain Detection Threshold.