Transcriptomic response in symptomless roots of clubroot infected kohlrabi (*Brassica oleracea* var. *gongylodes*) mirrors resistant plants

Stefan Ciaghi[1,$], Arne Schwelm[1,2,$], Sigrid Neuhauser[1,*]

[1] *University of Innsbruck, Institute of Microbiology, Technikerstraße 25, 6020 Innsbruck, Austria*
[2] *Swedish University of Agricultural Sciences, Department of Plant Biology, Uppsala BioCenter, Linnean Centre for Plant Biology, P.O. Box 7080, SE-75007 Uppsala, Sweden*

[*] Correspondence: Sigrid.Neuhauser@uibk.ac.at
[$] Contributed equally to this work

Abstract

Background: Clubroot disease caused by *Plasmodiophora brassicae* (Phytomyxea, Rhizaria) is one of the economically most important diseases of *Brassica* crops. The formation of hypertrophied roots accompanied by altered metabolism and hormone homeostasis is typical for infected plants. Not all roots of infected plants show the same phenotypic changes. While some roots remain uninfected, others develop galls of diverse size. The aim of this study was to analyse and compare the intra-plant heterogeneity of *P. brassicae* root galls and symptomless roots of the same host plants (*Brassica oleracea* var. *gongylodes*) collected from a commercial field in Austria using transcriptome analyses.

Results: Transcriptomes were markedly different between symptomless roots and gall tissue. Symptomless roots showed transcriptomic traits previously described for resistant plants. Genes involved in host cell wall synthesis and reinforcement were up-regulated in symptomless roots indicating elevated tolerance against *P. brassicae*. By contrast, genes involved in cell wall degradation and modification processes like expansion were up-regulated in root galls. Hormone metabolism differed between symptomless roots and galls. Brassinosteroid-synthesis was down-regulated in root galls, whereas jasmonic acid synthesis was down-regulated in symptomless roots. Cytokinin metabolism and signalling were up-regulated in symptomless roots with the exception of one CKX6 homolog, which was strongly down-regulated. Salicylic acid (SA) mediated defence response was up-regulated in symptomless roots, compared with root gall tissue. This is probably caused by a secreted benzoic acid salicylic acid methyl transferase from the pathogen (PbBSMT), which was one of the highest expressed pathogen genes in gall tissue. The PbBSMT derived Methyl-SA potentially leads to increased pathogen tolerance in uninfected roots.

Conclusions: Infected and uninfected roots of clubroot infected plants showed transcriptomic differences similar to those previously described between clubroot resistant and susceptible hosts. The here described intra-plant heterogeneity suggests, that for a better understanding of clubroot disease targeted, spatial analyses of clubroot infected plants will be vital in understanding this economically important disease.

Keywords: Clubroot, Host-pathogen interaction, *Plasmodiophora brassicae*, *Brassica oleracea*, root transcriptome, protist

List of abbreviations

ABA: Abscisic acid; BG: Brown spindle gall; BR: Brassinosteroids; C: Control; CDS: Coding sequence; CK: Cytokinin; COG: Clusters of Orthologous Groups; DEG: Differentially expressed gene; dpi: days post inoculation; ET: Ethylene; FDR: False discovery rate; FPKM: Fragments per kilobase per million; HR: hypersensitive response; IsoPct: Isoform percentage; JA: Jasmonic acid; MeSA: Methyl-salicylate; NCBI: National Center for Biotechnology Information; ORF: Open reading frame; PbraAT: Austrian *P. brassicae* field population; PR: Pathogenesis related; RNA: Ribonucleic acid; SA: Salicylic acid; SAM: S-adenosylmethionine; SL: Symptomless root; TAIR: The Arabidopsis Information Resource; WG: White spindle gall.

Background

Clubroot disease is one of the most important diseases of *Brassica* crops worldwide accounting for approximately 10% loss in *Brassica* vegetable, fodder, and oilseed crops (Dixon, 2009). Clubroot is caused by *Plasmodiophora brassicae*, an obligate biotrophic protist, taxonomically belonging to Phytomyxea within the eukaryotic super-group Rhizaria (Bulman and Braselton, 2014; Neuhauser *et al.*, 2014). This soil borne pathogen has a complex life cycle: zoospores infect root hairs where primary plasmodia form. These plasmodia develop into secondary zoospores, which are released into the soil and re-infect the root cortex where secondary plasmodia develop (Kageyama and Asano, 2009). The secondary plasmodia mature into resting spores, which are released into the soil. In infected host tissue division and elongation of cells is triggered upon infection, which leads to hypertrophies of infected roots resulting in the typical root galls or clubroots (Fig. 1).

Figure 1: Sampling site and plant material. a: Field where plants were sampled (Ranggen, Tyrol; Austria). b: Normally developed kohlrabi plant. Roots were analysed as negative control (C). c: Clubroot infected kohlrabi plants with three different root

79   phenotypes: symptomless roots (SL), white spindle galls (WG), and brownish spindle
80   galls (BG). Scale bar: 1 cm.
81
P. brassicae can only be grown and studied in co-culture with its host. This has
hampered both, targeted and large scale studies on the molecular basis of *P. brassicae*
and the interactions with its host (Schwelm *et al.*, 2018). Because of the economic
importance of clubroot disease, numerous studies analysed specific aspects of the
biology, physiology and molecular biology of the plant-pathogen interaction to better
understand and control the disease. The first of these experimental studies were based
on the *Arabidopsis/Plasmodiophora* pathosystem (e.g. Devos *et al.*, 2006; Siemens *et
al.*, 2006). During the last years an increasing number of *Brassica* (host) genomes
became available (Wang *et al.*, 2011; Chalhoub *et al.*, 2014; Liu *et al.*, 2014; Cheng *et
al.*, 2016), as did several *P. brassicae* genomes (Schwelm *et al.*, 2015; Bi *et al.*, 2016b;
Rolfe *et al.*, 2016; Daval *et al.*, 2018) permitting new research approaches, including
targeted transcriptome studies. There are analyses of (plant) transcriptomes of roots of
clubroot infected plants compared with uninfected plants (Agarwal *et al.*, 2011; Bi *et
al.*, 2016a; Malinowski *et al.*, 2016; Zhao *et al.*, 2017), of host varieties susceptible and
tolerant to clubroot (Jubault *et al.*, 2008; Bi *et al.*, 2016a), or the host response to
different *P. brassicae* isolates (Siemens *et al.*, 2006; Agarwal *et al.*, 2011; Lovelock *et
al.*, 2013; Zhang *et al.*, 2016).
99
Plants infected with *P. brassicae* show marked physiological changes including cell
wall biosynthesis, plant hormone metabolism and plant defence related processes.
Expansin genes, involved in plant cell expansion and elongation (Cosgrove, 2005),
were up-regulated in *P. brassicae* infected roots (Siemens *et al.*, 2006; Agarwal *et al.*,
2011; Irani *et al.*, 2018). In *P. brassicae* infected roots enzymatic activity of
Xyloglucan endo Transglucosylase/Hydrolases (XTHs) increases (Devos *et al.*, 2005),
while an early response to *P. brassicae* infection was up-regulation of the
phenylpropanoid pathway that provides lignin precursors (Zhao *et al.*, 2017). With
progression of clubroot development, the lignification of clubroot tissue was reduced
(Deora *et al.*, 2013) and genes involved in lignification processes were down-regulated
(Cao *et al.*, 2008). On the other hand cell wall thickening and lignification was
suggested to limit the spread of the pathogen in tolerant *B. oleracea* (Donald *et al.*,
2008) and *B. rapa* (Takahashi *et al.*, 2001).
113
The development of clubroot symptoms is accompanied by changes of plant hormone
homeostasis (Siemens *et al.*, 2006; Ludwig-Müller *et al.*, 2009; Malinowski *et al.*,
2016). During clubroot development, auxins mediate host cell divisions and elongation.
Auxins increase over time during clubroot development and accumulate in *P. brassicae*
infected tissues in a sink like manner (Ludwig-Müller *et al.*, 2009). In addition, genes
belonging to the auxin conjugating GH3 protein family are regulated differentially
during clubroot development (Jahn *et al.*, 2013) with one GH3 protein gene (PbGH3;
CEP01995.1) identified in the *P. brassicae* genome (Schwelm *et al.*, 2015). Cytokinins
(CKs) increase initially, but decrease again with the onset of gall formation
(Malinowski *et al.*, 2016). At the same time *P. brassicae* plasmodia produce minute
amounts of CKs (Müller and Hilgenberg, 1986). Therefore, CKs play a crucial role in
disease development not only through their regulation of cell division but also through
their interference in the sugar metabolism and invertase production, which might be
crucial for the nutrition of *P. brassicae* (Siemens *et al.*, 2011; Walerowski *et al.*, 2018).
128

Stress- and defence related phytohormones like salicylic acid (SA), jasmonic acid (JA), brassinosteroids (BR), and ethylene (ET) and their regulatory pathways also change in response to pathogen infection (Kazan and Lyons, 2014). The accumulation of SA plays a key role in plant defence against biotrophic pathogens, often resulting in a localized hypersensitive response and induction of pathogenesis-related (PR) genes. Systemic acquired resistance (SAR) is a form of induced resistance that is activated by SA throughout a plant after being exposed to elicitors from microbes or chemical stimuli (Klessig *et al.*, 2018). High endogenous levels of SA and exogenous SA reduced the infection of the host by *P. brassicae* (Lovelock *et al.*, 2013; Lovelock *et al.*, 2016). In tolerant hosts SA related genes are induced upon infection (Siemens *et al.*, 2009; Agarwal *et al.*, 2011; Zhang *et al.*, 2016). The SAR-deficient *npr1-1* and SA-deficient isochorismate synthase 1 (ICS1) *sid2 Arabidopsis* mutants showed an increased susceptibility to *P. brassicae*, whereas the *bik-1* mutant, with elevated SA levels, was more resistant (Chen *et al.*, 2016b). Pathogenesis-related defence proteins were induced by SAR and expressed more highly in resistant than in susceptible *B. rapa* and *Arabidopsis* species (Jubault *et al.*, 2013; Bi *et al.*, 2016a; Jia *et al.*, 2017). *P. brassicae* might counteract the plant SA-mediated defence via a secreted methyltransferase (PbBSMT; AFK13134.1). This SABATH-like methyltransferase has been shown to convert SA to methyl-salicylate (MeSA) *in vitro* (Ludwig-Müller *et al.*, 2015). The proposed function *in planta* is the removal of SA in local infected tissue as MeSA is volatile. *Arabidopsis* mutants expressing the PbBMST gene showed a higher susceptibility towards *P. brassicae* (Bulman *et al.*, 2018).

The *P. brassicae* PbGH3 was also able to conjugate JA with amino acids *in vitro* (Schwelm *et al.*, 2015). In general, JA is associated with resistance against necrotrophic microbes (Pieterse *et al.*, 2012; Fu and Dong, 2013). In *A. thaliana* Col-0 several JA-responsive genes were induced in infected root tissues and JA accumulates in galls (Siemens *et al.*, 2006; Gravot *et al.*, 2012). *Jasmonate resistant 1* (*jar1*) mutants, impaired in JA-Ile accumulation, showed a higher susceptibility to *P. brassicae* (Agarwal *et al.*, 2011; Gravot *et al.*, 2012). Thus, JA responses contributed to a basal resistance against some strains of *P. brassicae* in *A. thaliana* Col-0 (Gravot *et al.*, 2012). But in partially resistant *Arabidopsis* Bur-0 only weak JA responses compared with the susceptible Col-0 were found (Lemarié *et al.*, 2015). Generally, clubroot susceptible hosts show a high level of JA response, whereas it is reduced in resistant hosts (Jubault *et al.*, 2008; Bi *et al.*, 2016a). Those differences might be due to occur if aliphatic or aromatic glucosinolate production is induced by JA in the particular host (Xu *et al.*, 2018).

The aim of this study was to generate the first data set of root tissue specific transcriptomic response of individual plants during clubroot development. Usually, clubroot infected plants do not develop symptoms uniformly on all root parts with some roots showing strong symptoms and others not showing symptoms at all (Fig. 1). We collected samples of kohlrabi (*Brassica oleracea* var. *gongylodes*) infected with *P. brassicae* (PbraAT) from a field in Austria. We compared morphologically different clubroots and symptomless roots of the same infected plants and a control plant. We analysed similarities and differences of their transcriptomic profile focussing on cell wall metabolism, hormone metabolism, and defence response.

Methods

Sampling

Kohlrabi "purple Vienna" plants with clubroots and without visible root infections were collected from a *P. brassicae* infested field in Ranggen (Tyrol, Austria, 47°15'27"N 11°12'37"E) in August 2016 with the consent of the farmer. No other permissions were necessary to collect these samples. Root samples were classified by visible properties into symptomless roots (SL), smaller white spindle galls with waxy appearance (WG) and larger brownish spindle galls (BG) (Fig. 1). Samples were taken in triplicates from three individual clubroot infected plants. Only one plant without apparent infection could be identified. All other plants collected from the vegetable plot without easily visible symptoms where infected when examined closer in the lab. Nevertheless we additionally took samples from the one uninfected control plant (C) (Fig. 1). Galls and roots were thoroughly washed with tap water before samples were taken (categories C, SL, WG, and BG) and transferred to RNA*later*™ (Ambion, Austin, TX, USA) until RNA extraction.

RNA extraction and sequencing

The outer layer of the root galls was trimmed-off and the trimmed galls and symptomless roots were snap-frozen in liquid nitrogen and transferred to 1.5 mL tubes containing RNase free zirconia beads (0.5 mm and 2 mm in diameter). Samples were homogenized using a FastPrep (MP Biomedicals, Santa Ana, CA, USA) for 40 s at 6 m s^{-1} followed by manual grinding with pestles after repeated snap-freezing. Total RNA was extracted using the Qiagen RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions, but with an additional 80% ethanol column wash prior elution. RNA quantity and quality were determined using Agilent Bioanalyzer 2100 (Agilent Technologies, Palo Alto, CA, USA). Additional RNA quality assessment, polyA selection (SENSE mRNA-Seq Library Prep Kit; Lexogen, Vienna, Austria), library construction (10 libraries; 1x C, 3x SL, 3x WG, and 3x BG) and sequencing was performed at the VBCF NGS Unit (Vienna, Austria). Sequencing was performed on the Illumina HiSeq 2500 platform (Illumina, San Diego, CA, USA) with a strand specific paired end library (2x 125 bp) using v4 chemistry.

Bioinformatics

Raw reads were quality checked using FastQC (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Illumina adapters were removed and good quality reads were kept using Trimmomatic v0.36 (sliding window 5 bp; average quality score > 20) (Bolger *et al.*, 2014). Only reads with a minimum length of 75 bp were processed further after a repeated FastQC check to confirm quality improvement. From the uninfected control library, 75% of the reads were randomly picked three times to generate pseudo-triplicates. Transcripts were *de novo* assembled using Trinity v2.2 (Grabherr *et al.*, 2011) with strand specific library type (RF) and jaccard clip options. Expression estimation was performed using Trinity embedded RSEM (Li and Dewey, 2011) keeping only transcripts with more than at least one fragment per kilobase per million (FPKM > 1) and an isoform percentage (IsoPct) > 1%.

The assembled transcripts were blasted using BlastN (Altschul *et al.*, 1990) against the coding sequences (CDS) of *B. oleracea* (Liu *et al.*, 2014; http://brassicadb.org/brad/datasets/pub/Genomes/Brassica_oleracea/V1.1/) and a

custom database containing the CDS of the *P. brassicae* isolates e3 (Schwelm *et al.*,
2015) and PT3 (Rolfe *et al.*, 2016) to identify, if the transcript derived from the
pathogen or host (E-value $< 10^{-5}$). Transcripts with blast hits in both reference databases
were analysed manually to identify their origin according to sequence identity and
E-value. Transcripts with no hit in any reference were blasted (BlastP) against National
Center for Biotechnology Information (NCBI) non redundant protein database and
manually assigned to the corresponding species or discarded for further analysis.
Transcripts with a best hit to a Brassicaceae reference sequence were assigned to the
host transcriptome. Transcripts with hits to *P. brassicae* sequences were assigned to the
pathogen transcriptome. Open reading frames (ORFs) were predicted using
TransDecoder v3.0.1 (https://github.com/TransDecoder/). Only the longest ORF per
transcript was used for further analysis. Translated peptide sequences were annotated
using the KEGG (Kyoto Encyclopedia of Genes and Genomes) Automatic Annotation
Server (KAAS) (Moriya *et al.*, 2007) and eggNOG-mapper v0.99.3 (Huerta-Cepas *et
al.*, 2016). Kohlrabi genes were additionally annotated using Mercator (Lohse *et al.*,
2014) with default settings. Mercator categories were used for MapMan v3.6.0RC1
(Usadel *et al.*, 2009) to bin predicted genes into groups. Putative secreted proteins of
P. brassicae were predicted with Phobius v1.01 (Käll *et al.*, 2004) and SignalP v4.1
(Nielsen, 2017) in combination with TMHMM v2.0 (Krogh *et al.*, 2001). Carbohydrate
active enzymes were predicted using dbCAN (Yin *et al.*, 2012).
248
Log$_2$-fold changes of differentially expressed genes (DEGs) were calculated using
edgeR with default settings (Robinson *et al.*, 2010). All DEGs with false discovery rate
(FDR) < 0.05 were treated as DEGs. Heatmaps for selected Mercator/MapMan
categories were created using R v3.3.2 (R Core Team, 2016) with the package
'pheatmap' v1.0.8 (https://cran.r-project.org/web/packages/pheatmap/index.html)
applying UPGMA clustering. Labelling of the predicted *B. oleracea* genes was done
according to their homologous *A. thaliana* genes from TAIR (The Arabidopsis
Information Resource) and adapted if necessary. Abundance of DEGs was visualized
using the R package 'ggplot2' v2.2.1 (Wickham, 2009).
258
259

Results
Transcriptome analyses
In total 162 million good quality reads with an average length of 125 bp were obtained
from all libraries (Additional file 2: Table S1). A total of 10940 genes including
isoforms were predicted for *P. brassicae* and 42712 for *B. oleracea*. About 50% of the
P. brassicae and 85% of the kohlrabi transcripts could be functionally annotated using
eggNOG-mapper. Only 0.0005% of the reads of the control plant (C) and the
symptomless root samples (SL) matched *P. brassicae* transcripts, which indicates that
these roots were not infected by *P. brassicae*. In the white spindle galls (WG) and
brownish spindle galls (BG) libraries 23% and 33% of the reads matched *P. brassicae*
(Additional file 1: Figure S1).
271
The transcriptomes of infected plants (SL, WG, BG) contained a total of 5204
differentially expressed genes (DEGs). Compared with SL, in WG 1619 DEGs were
up-regulated and 2280 were down-regulated (Fig. 2, Additional file 2: Table S2), while
in BG 942 DEGs were up- and 2571 were down-regulated. 790 plant DEGs were
assigned to the COG (Cluster of Orthologous Groups) category "Information and

Storage Processing", 1401 to "Metabolism", 1245 to "Cellular Process and Signalling"
and 1768 to "Poorly Characterized" by eggNOG-mapper (Fig. 2, Additional file 2:
Table S3). Only 19 plant genes were differentially expressed between BG and WG
(Additional file 2: Tables S4). Between the control plant and the three root tissue types
of the infected plants (C vs SL/WG/BG) 19230 DEGs were found (Additional file 2:
Tables S5, S6).

Figure 2: Numbers of differentially expressed genes (DEGs) in clubroot infected
kohlrabi roots per COG category. DEGs were split into up- and down-regulated
genes. Total number of DEGs in each panel is given.

Plant cell wall metabolism
In *B. oleracea* 161 of the 5204 DEGs within infected plants (SL vs WG vs BG) were
involved in cell wall synthesis, modification, degradation, or phenylpropanoid
metabolism. Cellulose, hemicellulose, pectin, and lignin synthesis genes involved in
cell wall reinforcement were up-regulated in SL compared with gall tissues (Fig. 3),
whereas genes involved in cell wall modification and degradation were down-regulated
(Fig. 3). The changes in expression of cell wall genes were more prominent between
SL and BG than between SL and WG. In SL a UDP-D-glucuronate 4-epimerase 6
(GAE6) homolog responsible for the synthesis of UDP-D-glucuronic acid, the main
building block for pectins (Harholt *et al.*, 2010) was up-regulated compared with gall
tissue. Predicted expansin (EXP) and expansin-like (EXL) genes were mainly down-
regulated in SL compared with WG and BG (Fig. 3). Genes coding for XTHs were
among the strongest down-regulated DEGs in SL, with XTH24 being the strongest
down-regulated transcript of all DEGs. The phenylpropanoid pathway was up-
regulated in SL compared with WG and BG (Fig. 3). This includes the phenylalanine
ammonia-lyase 1 (PAL1) homolog, a key enzyme in lignin biosynthesis. Lignin is also
part of the xylem and xylogenesis genes which were up-regulated in SL compared with
BG. Flavonoid metabolism was also induced in SL (Additional file 1: Figure S2).
Compared with an uninfected control plant cell wall synthesis genes were up-regulated
in SL (Additional file 1: Figure S3).

Figure 3: Kohlrabi cell wall metabolism. Clustered heatmaps of $\log_2$ fold change values of DEGs. Genes involved in anabolic processes of cell wall components were generally up-regulated in SL compared with WG and BG, whereas catabolic and modifying genes were mainly down-regulated. No DEGs were present comparing WG with BG. Up-regulated genes are shaded in purple and down-regulated genes in green. *Arabidopsis* homologs are given. Genes are annotated and categorized according to MapMan/Mercator. NA: not assigned by MapMan. Supplementary information for the genes including KEGG, eggNog, and TAIR annotation are given in Additional file 3.

Plant hormones
The CK and auxin metabolism was altered in WG and BG compared with SL (Fig. 4).
So was a homolog of CKX6 (cytokinin oxidase/dehydrogenase 6) down-regulated in
SL compared with WG. Homologs of CKX5, CK receptors, and a CK-regulated UDP-
glucosyltransferase were up-regulated in SL. The CK synthesis genes CYP735A2
(cytochrome P450) and LOG1 (lonely guy 1) were up-regulated in SL compared with
root galls (Fig. 4). Down-stream in CK-signalling we found DEGs within ARR
(Arabidopsis response regulator; CK-signalling target) genes (Fig. 4). Type-B ARRs
were differentially expressed in WG and BG whereas differentially expressed Type-A
ARRs were exclusively found in BG. A putative AHK4 (Arabidopsis histidine kinase 4;
CK-receptor) gene was up-regulated in SL compared with BG where it was not detected
although expression values in SL were low. No DEGs of AHPs (Arabidopsis histidine
phosphotransfer protein) were found in SL compared with root galls. Most auxin related
DEGs, auxin response factors (ARFs and IAAs) and IAA amino acid conjugate
synthetases (GH3) were up-regulated in SL compared with gall tissue (Additional
file 1: Figure S4). However, an IAA7 (repressor of auxin inducible genes) and a
homolog of GH3.2 were down-regulated. Expression of PIN-FORMED 1 (PIN1) genes
was reduced in SL (Fig. 4), whereas SAUR (small auxin up-regulated RNA) and AIR12
(auxin-induced in root cultures protein 12-like) genes, were found in up- and down-
regulated DEGs (Fig. 4). Myrosinases and nitrilases were up-regulated in SL compared
with galls (Additional file 1: Figure S5). Compared with the unifected control both CK
and auxin metabolism were up-regulated in SL (Additional file 1: Figure S3). In BG
two transcripts related to auxin synthesis and regulation were down-regulated
compared with WG (Additional file 2: Table S4).
342
343 Early sterol biosynthesis genes, such as the steroid reductase DET2, were up-regulated
344 in SL compared with galls (Fig. 4). However, BR biosynthesis genes were generally
345 down-regulated in SL (Fig. 4), including key genes like DWF1 (dwarf 1) or BRI1 (BR
346 receptor brassinosteroid insensitive 1).
347
348 Abscisic acid (ABA) signal transduction related genes like ABI1 (ABA insensitive 1)
349 and HSI2 (high-level expression of sugar-inducible gene 2) were up-regulated in SL
350 compared with WG and BG (Fig. 4) whereas ABA related transcription factors
351 WRKY18 and HVA22A/C homologs were down-regulated in SL (Fig. 4, Additional
352 file 1: Figure S6).
353

Figure 4: Kohlrabi phytohormone metabolism. Heatmaps of $\log_2$ fold change values of DEGs. Genes involved in cytokinin, jasmonic acid, salicylic acid, and abscisic acid metabolism were up-regulated in SL. Genes coding for brassinosteroids clustered into two groups: later stages in BR biosynthesis (down-regulated) and early sterol biosynthesis (up-regulated). Genes involved in auxin metabolism were found within the up- and down-regulated DEGs in SL compared with root galls. One DEGs (JAR1) was found between WG and BG. Up-regulated genes are shaded in purple and down-regulated genes in green. *Arabidopsis* homologs are given. Genes are annotated and categorized according to MapMan/Mercator, except for cytokinin metabolism for which genes were categorized as described by (Malinowski *et al.*, 2016). NA: not assigned by MapMan. Supplementary information for the genes including KEGG, eggNog, and TAIR annotation are given in Additional file 4.

Plant defence

Generally, genes for disease resistance proteins were up-regulated in SL compared with WG and BG (Fig. 5, Additional file 1: Figure S7). From pathogen recognition genes via signalling proteins and transcription factors to pathogenesis related (PR) proteins, the whole signal cascade of pathogen defence was affected. Of the predicted defence related DEGs within infected plants (SL vs WG vs BG) 60 were assigned as TIR-NBS-LRR (Toll/interleukin-1 receptor nucleotide-binding site leucine-rich repeat) class proteins. Within R-genes only PP2-A (phloem protein 2-A) genes were up-regulated in SL compared with root galls (Fig 5). Expressed genes for disease resistance protein RSP4 were not altered between the tissue types (Additional file 5). Signalling genes encoding for MLO (mildew resistance locus o) and MKS1 (MAP kinase substrate 1) were up-regulated in SL (Fig. 5), while no differences in the expression of CHORD or SRFR1 (suppressor of RSP4-RDL1) genes was observed (Additional file 5). Two transcription factors involved in the biotic stress signalling cascade (MapMan bin 20.1.5) were up-regulated in SL compared with the root galls (Fig 5). MAP3K (mitogen activated kinase kinase kinase) and RSH (RELA/SPOT homolog) genes were not differentially expressed within infected plants (Additional file 5).

The BIK1 (botrytis-induced kinase 1), which interacts with BRI1 and BAK1 (BRI1-associated receptor kinase) to induce defence responses was up-regulated in SL compared with galls (Additional file 1: Figure S8) and up-regulated in SL compared with C (Additional file 1: Figure S3).

In the SL samples JA-related genes such as LOX2 (lipoxygenase 2), AOC (allene oxide cyclase), and HPL (hyperoxide lyase) were down-regulated, while other LOX genes and the JA amido synthetase genes JAR1 were up-regulated in SL compared with galls (Fig. 4). One down-regulated isoform of JAR1 was found between BG and WG. We found no glucosinolate biosynthesis genes in the DEGs in our study.

The ICS1 gene and the ICS1 activating transcription factor WRKY28 (van Verk *et al.*, 2011) were up-regulated in SL compared with the control plant (data not shown). Genes for SA modification, like the SA methylating SABATH methyl transferase genes (PXMT1, GAMT1) and a SA-glucosidase (UGT74F2) were up-regulated in SL compared with galls (Fig. 4). The SA induced PR1 gene was induced in SL (Fig. 5, Additional file 1: Figure S7). The PR-gene expression regulator NPR1 (non-expressor of PR1) was not differentially expressed in our samples. Of genes that regulate PR1 expression via NPR1, WRKY70 was down-regulated in SL whereas NPR3 and TGA3 were up-regulated (Fig. 5, Additional file 1: Figure S7). Compared with the uninfected plant, the TGA3 gene expressions appeared to be induced in SL. Genes for the TAO1 disease resistant protein, which induces PR1 expression were up-regulated in SL, as well as the NDR1 (non-race specific disease resistance 1) gene, required for the establishment of hypersensitive response and SAR. Genes for the disease resistant protein RPS2, activated by NDR1, were also up-regulated in SL (Fig. 5, Additional file 1: Figure S7). Additionally, other defence related genes coding for protease inhibitor genes, R-genes or some chitinases were down-regulated in SL (Fig. 5). Comparing biotic stress response genes of SL with the control plant (C), we observed an up-regulation of 164 of the total 190 DEGs in SL (Additional file 1: Figure S3).

Figure 5: Biotic stress response of kohlrabi. Heatmaps of log$_2$ fold change values of
DEGs. Almost all DEGs were up-regulated in SL compared with root galls. Two down-
regulated DEGs, a chitinase (CHI) and a gene of unknown function, were found
between WG and BG. Up-regulated genes are shaded in purple and down-regulated
genes in green. *Arabidopsis* homologs are given. Genes are annotated and categorized
according to MapMan/Mercator. NA: not assigned by MapMman. Supplementary
information for the genes including KEGG, eggNog, and TAIR annotation are given in
Additional file 5.

P. brassicae gene expression

The *P. brassicae* genes with the highest FPKM values belonged to growth and cellular
process related COG categories such as translation, transcription, and signal
transduction, but also in energy conversion and carbohydrate and lipid metabolism
(Additional file 1: Figure S9). The *P. brassicae* PbBSMT gene was amongst the highest
expressed pathogen genes (Additional file 2: Tables S7, S8). Other highly expressed
genes were HSPs (heat shock proteins), a glutathione-S-transferase, an ankyrin repeat
domain-containing protein, ribosomal genes, and genes of unknown function
(Additional file 2: Tables S7, S8). The PbGH3 gene was not expressed in our samples.
The *P. brassicae* protease gene PRO1, proposed to be involved in spore germination
(Feng *et al.*, 2010), was expressed in both WG and BG.

Between WG and BG samples only five *P. brassicae* DEGs were identified, coding for
a HSP, a chromosomal maintenance protein, a DNA-directed RNA-polymerase, a
retrotransposon and a Scl Tal1 interrupting locus protein (Additional file 2: Table S9).
Cumulating all FPKM values revealed that most sequenced reads from *P. brassicae*
RNA extracted from root galls (WG and BG) mapped to the COG categories "Post-

445 translational modification, protein turnover, chaperon functions" and "Translation"
446 (Additional file 1: Figure S10). Very few *P. brassicae* reads were found in the data
447 obtained from the control plant and SL (Additional file 1: Figure S11), those were most
448 likely from attached spores or contamination via soil particles.
449
450

Discussion

Symptomless roots of clubroot infected plants show transcriptomic traits of clubroot resistant/tolerant plants

We found that symptomless roots and clubroots originating from the same plant showed
differences in their transcriptomic profile similar to previously described differences of
roots between resistant and susceptible plants (Chen *et al.*, 2016a; Jia *et al.*, 2017). Gene
expression patterns of symptomless roots were similar to the patterns described for
resistant hosts, while in clubroot tissue patterns were similar to those observed in
susceptible plants.
460
Reinforcement of cell walls has previously been reported to hamper the development
of *P. brassicae* in resistant *B. oleracea* (Donald *et al.*, 2008) and *B. rapa* callus cultures
(Takahashi *et al.*, 2001). Lignin biosynthesis genes were up-regulated in SL tissue
compared with root gall tissue (Fig. 3). Induced lignification processes were observed
in shoots of infected plants (Irani *et al.*, 2018) and between resistant and susceptible
B. oleracea cultivars (Zhang *et al.*, 2016). PAL1, a key enzyme in lignin, SA (discussed
below) and flavonoid biosynthesis (Chu *et al.*, 2014; Song *et al.*, 2016; Lahlali *et al.*,
2017) was up-regulated in symptomless roots compared with clubroot tissue (Fig. 3).
Increased lignin biosynthesis and up-regulation of PAL1 has been described for a
clubroot resistant oilseed *B. rapa* line carrying the resistance gene Rcr1 (Lahlali *et al.*,
2017), while callus cultures overexpressing PAL1 were resistant to infection by
P. brassicae (Takahashi *et al.*, 2001). Root reinforcement, therefore, seems to be a part
of the tolerance mechanisms of plants against *P. brassicae,* which has only a limited
arsenal of plant cell wall degrading enzymes in its genome (Schwelm *et al.*, 2015; Rolfe
et al., 2016) and infects its hosts via mechanical force with a specialised extrusosome
called "Stachel and Rohr" (Kageyama and Asano, 2009). Once *P. brassicae*
successfully infected its host, movement and spread within root tissues has been
discussed to happen via the plasmodesmata (Donald *et al.*, 2008; Riascos *et al.*, 2011;
Badstoeber *et al.*, 2018). Increased stability of the cell wall, as indicated by gene
expression patterns in symptomless roots (Fig. 3) or described for resistant plants
(Takahashi *et al.*, 2001; Donald *et al.*, 2008), requires higher mechanical force for
successful (primary) infection of yet uninfected roots or movement in between host
cells. Therefore, it is likely that cell wall reinforcement of the host cells is a considerable
obstacle for *P. brassicae*. The result of increased lignification in SL compared with
galls could also be the result of repressed xylem development in *P. brassicae* infected
hosts (Malinowski *et al.*, 2012).
487
In symptomless roots of infected plants, defence related pathways regulated by
hormones show patterns usually linked to induced plant defence (Fig. 4). Clubroot
tissue on the other hand showed suppression of SA-defence related processes. Genes of
SA biosynthesis were up-regulated in SL, but were down-regulated in galls. Salicylic
acid can be synthesised via isochorismate (ICS pathway) or from phenylalanine (via
PAL pathway) (Pieterse *et al.*, 2012; Lovelock *et al.*, 2016). The up-regulation of the

PAL1 gene in symptomless roots could therefore also be linked to the PAL-dependent
synthesis of SA, but the majority of SA in clubroot is produced via the ICS pathway
(Lovelock *et al.*, 2016). Based on the higher expression of ICS1 and WRKY28 in SL
compared with the control, the synthesis of SA was likely induced in SL. In clubroot
resistant hosts SA-defence is usually induced (Jubault *et al.*, 2013; Bi *et al.*, 2016a;
Lovelock *et al.*, 2016), whereas the SA deficient *sid2* (ICS1) mutant of *Arabidopsis*
was more susceptible to *P. brassicae* infection (Chen *et al.*, 2016b).

501
High SA levels in plant tissues helped to reduce new *P. brassicae* infections (Lovelock
et al., 2013), but SA alone is not sufficient to induce resistance against *P. brassicae*
(Lovelock *et al.*, 2016). Because SA levels increase in clubroot tissue *P. brassicae* is
thought to secret a SABATH-type methyltransferase (PbBSMT; Ludwig-Müller *et al.*,
2015; Bulman *et al.*, 2018), which was one of the highest expressed genes of
P. brassicae in this study (Additional file 2: Tables S7, S8). PbBSMT has been shown
to methylate salicylic acid, contributing to a local reduction of SA in the galls (Ludwig-
Müller *et al.*, 2015). MeSA is the major transport form of SA in the plant and has a key
role in inducing SAR (Park *et al.*, 2007; Vlot *et al.*, 2009). So based on our data we
hypothesise that *P brassicae* reduces SA concentrations in the galls via PbBMST
mediated methylation (Ludwig-Müller *et al.*, 2015). The produced MeSA could trigger
SA-related defences in distant plant parts, which become resilient towards new
pathogen infection.

515
Processes downstream from SA are mediated via NPR1 (not differentially expressed in
our dataset). NPR1 induces PR-gene expression when bound to TGA3 (Saleh *et al.*,
2015). In its interaction with WRKY70, NPR1 serves as a negative regulator of the SA
biosynthesis gene ICS1 (Wang *et al.*, 2006). Thus the observed up-regulation of TGA3
in SL (Additional file 1: Figure S7) would lead to an induction of expression of PR-
genes in SL. The reduced expression of WRKY70 in SL compared with the galls would
lead to a higher SA production in the uninfected root tissue. Additionally, we found an
up-regulation of NPR3 genes, coding for repressors of SA-defence genes (Ding *et al.*,
2018), in SL compared with the galls (Fig. 5). When bound to SA, NPR3 would lose
its function of repressing SA-defence genes (Ding *et al.*, 2018). The higher expression
of NPR3 in the SL might be necessary to compensate for negative effects of the SA
synthesis in SL roots.

528
Jasmonic acid contributed to a basal resistance against some strains of *P. brassicae*.
A. thaliana Col-0 and *A. thaliana* mutants impaired in JA-Ile accumulation showed a
higher susceptibility to *P. brassicae* (Agarwal *et al.*, 2011; Gravot *et al.*, 2012). In
A. thaliana Col-0 several JA-responsive genes were induced in infected root tissues and
JA accumulates in galls (Siemens *et al.*, 2006; Gravot *et al.*, 2012). But in partially
resistant *Arabidopsis* Bur-0 only weak JA responses compared with the susceptible
Col-0 were found (Lemarié *et al.*, 2015). Those differences might be due to if aliphatic
or aromatic glucosinolate production is induced by JA in the particular host (Xu *et al.*,
2018). Generally, clubroot susceptible hosts show a high level of JA response, whereas
it is reduced in resistant hosts (Jubault *et al.*, 2008; Bi *et al.*, 2016a). In our samples,
JAZ genes were up-regulated in SL compared with the gall tissue (Fig. 4). JAZ proteins
act as JA co-receptors and transcriptional repressors in JA signalling (Kazan and
Manners, 2012) reducing the JA synthesis in SL. This was previously observed in
B. oleracea plants, where JAZ expression was up-regulated in resistant plants and JA
synthesis was highly induced in susceptible plants (Zhang *et al.*, 2016). Among JA-

metabolism genes HPL, and LOX2 expression were up-regulated in the galls. LOX2 is
essential for the formation of oxylipin volatiles (Mochizuki *et al.*, 2016). The HPL
protein competes with substrates essential for JA-synthesis, producing volatile and non-
volatile oxylipins (Wasternack and Hause, 2013). The higher expression of HPL and
LOX2 in the galls might lead to the production of volatile aldehydes rather than a JA
accumulation in the galls.
550
In SL BR-synthesis and BR-signalling genes were down-regulated (Fig. 4). BRs are
necessary for the development of clubroot tissue (Schuller *et al.*, 2014), hence a
reduction in BRs impairs *P. brassicae* growth and development in SL. The receptor-
like cytoplasmic kinase BIK1, a negative regulator of BR-synthesis (Lin *et al.*, 2013),
was induced in SL (Additional file 1: Figure S8). *Arabidopsis bik1* knockout mutants
have an increased tolerance to *P. brassicae* lacking the typical pathogen phenotype (Bi
et al., 2016a; Chen *et al.*, 2016c). For BIK1 the expression of the SL roots differed to
resistant plants, as *Arabidopsis bik1* knockout mutants have an increased tolerance to
P. brassicae (Bi *et al.*, 2016a; Chen *et al.*, 2016c). In *Arabidopsis bik1* clubroot
resistance was likely not due to the regulatory function of BIK1 for BR, but to increased
PR1 expression in this mutant (Chen *et al.*, 2016c).
562

Transcriptomes of symptomless roots and clubroots of the same plant are markedly different

Clubroots and symptomless roots of the same *P. brassicae* infected plants showed
markedly different gene expression patterns. The morphological changes of the
clubroots go in hand with expression of genes that reduce cell wall stability and growth
related processes. Genes for cell wall-loosening processes such as expansins and XTHs
(Downes *et al.*, 2001; Sun *et al.*, 2005), were up-regulated in gall tissue (Fig. 3). Up-
regulation of expansins was reported from *Arabidopsis* clubroot tissue (Siemens *et al.*,
2006; Irani *et al.*, 2018) while XTH activity was reported in *B. rapa* clubroots (Devos
et al., 2005). Suppression of xyloglucan, xylan, hemicellulose, pectin and lignin
synthesis in gall tissues implies additional reduction of the cell wall stability and
rigidity in gall tissues similar to previous findings (Schuller *et al.*, 2014; Zhang *et al.*,
2016). GAE6 expression was reduced in the galls supporting clubroot development:
Arabidopsis gae1, *gae6*, and *gae1/gae6* mutants contained lower levels of pectin in
their leaf cell walls making them more susceptible to *Pseudomonas syringae* and
Botrytis cinerea (Bethke *et al.*, 2016). A similar mechanism might benefit *P. brassicae*.
Induction of the lignin pathway was an early response (48h) in *Arabidopsis* (Zhao *et
al.*, 2017). In the brownish kohlrabi galls investigated in our study, lignification genes
were down-regulated suggesting *P. brassicae* suppresses host lignification in tissues
where it is already established. In *B. napus* reduced lignification was also implied by
the down-regulation of CCoAOMT (caffeoyl-CoA O-methyltransferase) (Cao *et al.*,
2008). Here, although not statistically significant, CCoAOMT was also lower
expressed in root galls compared with SL, implying a decreased lignin biosynthesis in
clubroots.
587
The marked hypertrophies of the plant roots go hand in hand with changes in the
homeostasis of the growth hormones CK and auxins which appear to be host and time
dependent (Ludwig-Müller *et al.*, 2009; Jia *et al.*, 2017). With the exception of CKX6,
cytokinin related genes were up-regulated in SL (Fig. 4). Fine tuning of the hormone
balance appears to be essential in clubroot disease development (Ludwig-Müller *et al.*,
2009). Elevated CK levels are important for the onset of disease development via

increasing cell divisions. However, at the onset of gall formation, CK metabolism genes
including CK synthesizing and degrading enzymes are repressed (Siemens *et al.*, 2006;
Malinowski *et al.*, 2016). The more active CK metabolism in the SL might interfere
with clubroot development as CKX overexpressing *Arabidopsis* mutant showed
reduced gall formation (Siemens *et al.*, 2006). In our kohlrabi samples, CKX6 was
strongly down-regulated in SL compared with WG but not compared with BG (Fig. 4).
The high expression of CKX6 in WG indicates the presence of plasmodia, as the gene
was found to be strongly up-regulated only in cells containing *P. brassicae* plasmodia
(Schuller *et al.*, 2014). Although AHK4 was described to be highly induced in infected
A. thalinana plants 10 dpi (Siemens *et al.*, 2006), in a different study expression of
AHK and AHP genes did not differ between infected and uninfected *Arabidopsis* at
later time points (Malinowski *et al.*, 2016). Similar to these findings, in our much longer
infected kohlrabi plants AHK and AHP genes were not differential expressed between
SL and root galls. Expression levels were also very low indicating no major role of
those genes in the analysed roots. The ARR5 gene was down-regulated in infected roots
and hypocotyl tissues 16 and 26 dpi in *A. thaliana* (Malinowski *et al.*, 2016) but in a
different study ARR5 was up-regulated 10 dpi in infected roots (Siemens *et al.*, 2006).
For *Brassica* hosts, the expression of ARR5 in clubroot infected roots has not been
described. In our samples ARR5 levels increase in BG, but no difference is seen
between SL and WG. The ARR5 gene does not appearto have a prominent role at the
stage of the here investigated infections. Also. evidence that *P. brassicae* interferes
with the CK balance via the PbCKX of *P. brassicae* in gall tissues has not been seen in
this study as the PbCKX gene was not expressed.
617
Myrosinases and nitrilases that can synthesize auxins from secondary metabolites or
aromatic amino acids were up-regulated in SL compared with galls (Additional file 1:
Figures S6), but were previously reported to be induced in *Arabidopsis* galls (Grsic-
Rausch *et al.*, 2000; Siemens *et al.*, 2006). The auxin-induced GH3 gene family, which
conjugates IAA to several amino acids, is involved in various responses of plants to
abiotic and biotic stresses. The GH3.2 gene was shown to be specifically expressed in
clubroots of *Arabidopsis* (Jahn *et al.*, 2013), and was also up-regulated in the kohlrabi
galls. Expression of the *P. brassicae* PbGH3 gene was not detected in this study and
appears not to play a role in the auxin (or JA) homeostasis at the stage of our gall
samples.
628
Besides the described defence responses in SL that are similar to defence responses of
resistance plants, we observed up-regulation of other defence related genes in the SL
(Fig. 5, Additional file 1: Figure S7). Homologues of the *Arabidopsis* Toll-IL-1
receptor disease resistance proteins TAO1, NDR1 and RPS2 genes were up-regulated
in SL. Those genes confer resistance to biotrophic bacterial pathogens in *Arabidopsis*
when recognizing effectors (Coppinger *et al.*, 2004; Eitas *et al.*, 2008). In roots of
beans, NDR1 also suppressed nematode parasitism by activating defence responses
(McNeece *et al.*, 2017). Thus, we found indications that those proteins might also be
involved in a defence response towards *P. brassicae*.
638
639 As a result of gall formation and a reduced number of fine roots, clubroot infected plants
640 face abiotic stress like lower water and nutrient supply. The differential expression of
641 ABA related genes (Fig. 4) are therefore likely a response to the abiotic stress in the
642 galls. Lower water supply could also be a consequence of reduced xylem production
643 (Malinowski *et al.*, 2012).

Conclusions

Clubroots and symptomless roots of the same *P. brassicae* infected plants showed very different gene expression patterns. The differences in the plant hormone metabolism might be responsible for the different outcomes in gall tissue and in symptomless roots as is the increased cell wall stability in symptomless roots. These results highlight, that interpreting clubroot transcriptomes or any other data originating from whole root systems might result in a dilution of biologically relevant signatures. This clearly calls for further studies analysing intra- and inter-tissue specific pattern of clubroot infected plants. As genes involved in resistance responses to *P. brassicae* were up-regulated in symptomless roots, this might aid the identification of novel traits for resistance breeding.

Acknowledgements

We thank Srilakshmy Harikrishnan for discussion of bioinformatic analyses. M. H. Borham (Agriculture and Agri-Food Canada, 107 Science Place, Saskatoon, SK S7N0X2, Canada) provided the PT3 data. Illumina sequencing was performed at the VBCF NGS Unit (www.vbcf.ac.at).

Funding

S.C and S.N. were funded by the Austrian Science Fund (grant Y0801-B16) and A.S. by Formas, the Swedish Research Council (grant 2015-1317), financial support only.

Availability of data and materials

The datasets generated and analysed during the current study are available in the European Nucleotide Archive (ENA; https://www.ebi.ac.uk/ena) repository under the project PRJEB26435 (Accessions ERR2567399-ERR2567408) or are available from the corresponding author on request.

Authors' contributions

Experimental concept and design: SN. Wet lab work: SC. Bioinformatic and statistic analysis: SC. Analysis of Results: SC, AS, SN. Manuscript writing: SC, AS, SN. Figures and tables: SC. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Additional files

Additional file 1: Figure S1 - Figure S11
Additional file 2: Table S1 - Table S9
Additional file 3: Additional data for Figure 3
Additional file 4: Additional data for Figure 4
Additional file 5: Additional data for Figure 5

References

Agarwal A, Kaul V, Faggian R, Rookes JE, Ludwig-Muller J, Cahill DM (2011). Analysis of global host gene expression during the primary phase of the arabidopsis thaliana-plasmodiophora brassicae interaction. *Functional Plant Biology* **38:** 462-478.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990). Basic local alignment search tool. *J Mol Biol* **215:** 403-410.

Badstoeber J, Gachon CMM, Sandbichler AM, Neuhauser S (2018). Biotrophic interactions disentangled: In-situ localisation of mrnas to decipher plant and algal pathogen - host interactions at single cell level. *bioRxiv***:** 378794.

Bethke G, Thao A, Xiong G, Li B, Soltis NE, Hatsugai N *et al* (2016). Pectin biosynthesis is critical for cell wall integrity and immunity in arabidopsis thaliana. *Plant Cell* **28:** 537-556.

Bi K, He Z, Gao Z, Zhao Y, Fu Y, Cheng J *et al* (2016a). Integrated omics study of lipid droplets from plasmodiophora brassicae. *Scientific reports* **6:** 36965.

Bi K, He ZC, Gao ZX, Zhao Y, Fu YP, Cheng JS *et al* (2016b). Integrated omics study of lipid droplets from plasmodiophora brassicae. *Scientific reports* **6**.

Bolger AM, Lohse M, Usadel B (2014). Trimmomatic: A flexible trimmer for illumina sequence data. *Bioinformatics* **30:** 2114-2120.

Bulman S, Braselton JP (2014). 4 rhizaria: Phytomyxea. In: McLaughlin DJ, Spatafora JW (eds). *Systematics and evolution* Springer Berlin Heidelberg. pp 99-112.

Bulman S, Richter F, Marschollek S, Benade F, Jülke S, Ludwig-Müller J (2018). *Arabidopsis thaliana* expressing pbbsmt, a gene encoding a sabath-type methyltransferase from the plant pathogenic protist *plasmodiophora brassicae*, show leaf chlorosis and altered host susceptibility. *Plant Biology*.

Cao T, Srivastava S, Rahman MH, Kav NNV, Hotte N, Deyholos MK *et al* (2008). Proteome-level changes in the roots of brassica napus as a result of plasmodiophora brassicae infection. *Plant Science* **174:** 97-115.

Chalhoub B, Denoeud F, Liu S, Parkin IA, Tang H, Wang X *et al* (2014). Plant genetics. Early allopolyploid evolution in the post-neolithic brassica napus oilseed genome. *Science* **345:** 950-953.

739 Chen J, Pang W, Chen B, Zhang C, Piao Z (2016a). Transcriptome analysis of brassica
740 rapa near-isogenic lines carrying clubroot-resistant and -susceptible alleles in response
741 to plasmodiophora brassicae during early infection. *Front Plant Sci* **6:** 1183.
742
743 Chen T, Bi K, He Z, Gao Z, Zhao Y, Fu Y *et al* (2016b). Arabidopsis mutant bik1
744 exhibits strong resistance to plasmodiophora brassicae. *Front Physiol* **7:** 402.
745
746 Chen T, Bi K, He ZC, Gao ZX, Zhao Y, Fu YP *et al* (2016c). Arabidopsis mutant bik1
747 exhibits strong resistance to plasmodiophora brassicae. *Front Physiol* **7**.
748
749 Cheng F, Sun R, Hou X, Zheng H, Zhang F, Zhang Y *et al* (2016). Subgenome parallel
750 selection is associated with morphotype diversification and convergent crop
751 domestication in brassica rapa and brassica oleracea. *Nat Genet* **48:** 1218-1224.
752
753 Chu MG, Song T, Falk KC, Zhang XG, Liu XJ, Chang A *et al* (2014). Fine mapping
754 of rcr1 and analyses of its effect on transcriptome patterns during infection by
755 plasmodiophora brassicae. *Bmc Genomics* **15**.
756
757 Coppinger P, Repetti PP, Day B, Dahlbeck D, Mehlert A, Staskawicz BJ (2004).
758 Overexpression of the plasma membrane-localized ndr1 protein results in enhanced
759 bacterial disease resistance in arabidopsis thaliana. *Plant J* **40:** 225-237.
760
761 Cosgrove DJ (2005). Growth of the plant cell wall. *Nat Rev Mol Cell Bio* **6:** 850-861.
762
763 Daval S, Belcour A, Gazengel K, Legrand L, Gouzy J, Cottret L *et al* (2018).
764 Computational analysis of the plasmodiophora brassicae genome: Mitochondrial
765 sequence description and metabolic pathway database design. *Genomics* **In Press**.
766
767 Deora A, Gossen BD, McDonald MR (2013). Cytology of infection, development and
768 expression of resistance to plasmodiophora brassicae in canola. *Ann Appl Biol* **163:** 56-
769 71.
770
771 Devos S, Vissenberg K, Verbelen JP, Prinsen E (2005). Infection of chinese cabbage
772 by plasmodiophora brassicae leads to a stimulation of plant growth: Impacts on cell
773 wall metabolism and hormone balance. *New Phytol* **166:** 241-250.
774
775 Devos S, Laukens K, Deckers P, Van Der Straeten D, Beeckman T, Inze D *et al* (2006).
776 A hormone and proteome approach to picturing the initial metabolic events during
777 plasmodiophora brassicae infection on arabidopsis. *Mol Plant Microbe Interact* **19:**
778 1431-1443.
779
780 Ding Y, Sun T, Ao K, Peng Y, Zhang Y, Li X *et al* (2018). Opposite roles of salicylic
781 acid receptors npr1 and npr3/npr4 in transcriptional regulation of plant immunity. *Cell*
782 **173:** 1454-1467 e1415.
783
784 Dixon GR (2009). The occurrence and economic impact of plasmodiophora brassicae
785 and clubroot disease. *J Plant Growth Regul* **28:** 194-202.
786

787 Donald EC, Jaudzems G, Porter IJ (2008). Pathology of cortical invasion by
788 plasmodiophora brassicae in clubroot resistant and susceptible brassica oleracea hosts.
789 *Plant Pathology* **57:** 201-209.
790
791 Downes BP, Steinbaker CR, Crowell DN (2001). Expression and processing of a
792 hormonally regulated β-expansin from soybean. *Plant Physiology* **126:** 244-252.
793
794 Eitas TK, Nimchuk ZL, Dangl JL (2008). Arabidopsis tao1 is a tir-nb-lrr protein that
795 contributes to disease resistance induced by the pseudomonas syringae effector avrb.
796 *Proc Natl Acad Sci U S A* **105:** 6475-6480.
797
798 Feng J, Hwang R, Hwang SF, Strelkov SE, Gossen BD, Zhou QX *et al* (2010).
799 Molecular characterization of a serine protease pro1 from plasmodiophora brassicae
800 that stimulates resting spore germination. *Mol Plant Pathol* **11:** 503-512.
801
802 Fu ZQ, Dong X (2013). Systemic acquired resistance: Turning local infection into
803 global defense. *Annu Rev Plant Biol* **64:** 839-863.
804
805 Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I *et al* (2011). Full-
806 length transcriptome assembly from rna-seq data without a reference genome. *Nat
807 Biotechnol* **29:** 644-652.
808
809 Gravot A, Deleu C, Wagner G, Lariagon C, Lugan R, Todd C *et al* (2012). Arginase
810 induction represses gall development during clubroot infection in arabidopsis. *Plant
811 Cell Physiol* **53:** 901-911.
812
813 Grsic-Rausch S, Kobelt P, Siemens JM, Bischoff M, Ludwig-Muller J (2000).
814 Expression and localization of nitrilase during symptom development of the clubroot
815 disease in arabidopsis. *Plant Physiol* **122:** 369-378.
816
817 Harholt J, Suttangkakul A, Vibe Scheller H (2010). Biosynthesis of pectin. *Plant
818 Physiol* **153:** 384-395.
819
820 Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC *et al* (2016).
821 Eggnog 4.5: A hierarchical orthology framework with improved functional annotations
822 for eukaryotic, prokaryotic and viral sequences. *Nucleic Acids Res* **44:** D286-293.
823
824 Irani S, Trost B, Waldner M, Nayidu N, Tu J, Kusalik AJ *et al* (2018). Transcriptome
825 analysis of response to plasmodiophora brassicae infection in the arabidopsis shoot and
826 root. *BMC Genomics* **19:** 23.
827
828 Jahn L, Mucha S, Bergmann S, Horn C, Staswick P, Steffens B *et al* (2013). The
829 clubroot pathogen (plasmodiophora brassicae) influences auxin signaling to regulate
830 auxin homeostasis in arabidopsis. *Plants* **2:** 726-749.
831
832 Jia H, Wei X, Yang Y, Yuan Y, Wei F, Zhao Y *et al* (2017). Root rna-seq analysis
833 reveals a distinct transcriptome landscape between clubroot-susceptible and clubroot-
834 resistant chinese cabbage lines after plasmodiophora brassicae infection. *Plant and Soil*
835 **421:** 93-105.
836

837 Jubault M, Hamon C, Gravot A, Lariagon C, Delourme R, Bouchereau A *et al* (2008).
838 Differential regulation of root arginine catabolism and polyamine metabolism in
839 clubroot-susceptible and partially resistant arabidopsis genotypes. *Plant Physiology*
840 **146:** 2008-2019.
841
842 Jubault M, Lariagon C, Taconnat L, Renou JP, Gravot A, Delourme R *et al* (2013).
843 Partial resistance to clubroot in arabidopsis is based on changes in the host primary
844 metabolism and targeted cell division and expansion capacity. *Funct Integr Genomics*
845 **13:** 191-205.
846
847 Kageyama K, Asano T (2009). Life cycle of plasmodiophora brassicae. *J Plant Growth*
848 *Regul* **28:** 203-211.
849
850 Käll L, Krogh A, Sonnhammer ELL (2004). A combined transmembrane topology and
851 signal peptide prediction method. *J Mol Biol* **338:** 1027-1036.
852
853 Kazan K, Manners JM (2012). Jaz repressors and the orchestration of phytohormone
854 crosstalk. *Trends Plant Sci* **17:** 22-31.
855
856 Kazan K, Lyons R (2014). Intervention of phytohormone pathways by pathogen
857 effectors. *Plant Cell* **26:** 2285-2309.
858
859 Klessig DF, Choi HW, Dempsey DA (2018). Systemic acquired resistance and salicylic
860 acid: Past, present and future. *Mol Plant Microbe Interact*.
861
862 Krogh A, Larsson B, von Heijne G, Sonnhammer EL (2001). Predicting transmembrane
863 protein topology with a hidden markov model: Application to complete genomes. *J Mol*
864 *Biol* **305:** 567-580.
865
866 Lahlali R, Song T, Chu M, Yu F, Kumar S, Karunakaran C *et al* (2017). Evaluating
867 changes in cell-wall components associated with clubroot resistance using fourier
868 transform infrared spectroscopy and rt-pcr. *International Journal of Molecular*
869 *Sciences* **18:** 2058.
870
871 Lemarié S, Robert-Seilaniantz A, Lariagon C, Lemoine J, Marnet N, Jubault M *et al*
872 (2015). Both the jasmonic acid and the salicylic acid pathways contribute to resistance
873 to the biotrophic clubroot agent plasmodiophora brassicae in arabidopsis. *Plant Cell*
874 *Physiol* **56**.
875
876 Li B, Dewey CN (2011). Rsem: Accurate transcript quantification from rna-seq data
877 with or without a reference genome. *BMC Bioinformatics* **12:** 323.
878
879 Lin WW, Lu DP, Gao XQ, Jiang S, Ma XY, Wang ZH *et al* (2013). Inverse modulation
880 of plant immune and brassinosteroid signaling pathways by the receptor-like
881 cytoplasmic kinase bik1. *P Natl Acad Sci USA* **110:** 12114-12119.
882
883 Liu S, Liu Y, Yang X, Tong C, Edwards D, Parkin IA *et al* (2014). The brassica oleracea
884 genome reveals the asymmetrical evolution of polyploid genomes. *Nat Commun* **5:**
885 3930.
886

887 Lohse M, Nagel A, Herter T, May P, Schroda M, Zrenner R *et al* (2014). Mercator: A
888 fast and simple web server for genome scale functional annotation of plant sequence
889 data. *Plant Cell Environ* **37:** 1250-1258.
890
891 Lovelock DA, Donald CE, Conlan XA, Cahill DM (2013). Salicylic acid suppression
892 of clubroot in broccoli (brassicae oleracea var. Italica) caused by the obligate biotroph
893 plasmodiophora brassicae. *Australas Plant Path* **42:** 141-153.
894
895 Lovelock DA, Sola I, Marschollek S, Donald CE, Rusak G, van Pee KH *et al* (2016).
896 Analysis of salicylic acid-dependent pathways in arabidopsis thaliana following
897 infection with plasmodiophora brassicae and the influence of salicylic acid on disease.
898 *Mol Plant Pathol* **17:** 1237-1251.
899
900 Ludwig-Müller J, Prinsen E, Rolfe SA, Scholes JD (2009). Metabolism and plant
901 hormone action during clubroot disease. *J Plant Growth Regul* **28:** 229-244.
902
903 Ludwig-Müller J, Jülke S, Geiß K, Richter F, Mithöfer A, Šola I *et al* (2015). A novel
904 methyltransferase from the intracellular pathogen plasmodiophora brassicae methylates
905 salicylic acid. *Mol Plant Pathol* **16**.
906
907 Malinowski R, Smith JA, Fleming AJ, Scholes JD, Rolfe SA (2012). Gall formation in
908 clubroot-infected arabidopsis results from an increase in existing meristematic
909 activities of the host but is not essential for the completion of the pathogen life cycle.
910 *Plant J* **71:** 226-238.
911
912 Malinowski R, Novák O, Borhan MH, Spíchal L, Strnad M, Rolfe SA (2016). The role
913 of cytokinins in clubroot disease. *European Journal of Plant Pathology* **145:** 543-557.
914
915 McNeece BT, Pant SR, Sharma K, Niruala P, Lawrence GW, Klink VP (2017). A
916 glycine max homolog of non-race specific disease resistance 1 (ndr1) alters defense
917 gene expression while functioning during a resistance response to different root
918 pathogens in different genetic backgrounds. *Plant Physiol Biochem* **114:** 60-71.
919
920 Mochizuki S, Sugimoto K, Koeduka T, Matsui K (2016). Arabidopsis lipoxygenase 2
921 is essential for formation of green leaf volatiles and five-carbon volatiles. *Febs Lett*
922 **590:** 1017-1027.
923
924 Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M (2007). Kaas: An automatic
925 genome annotation and pathway reconstruction server. *Nucleic Acids Res* **35:** W182-
926 185.
927
928 Müller P, Hilgenberg W (1986). Isomers of zeatin and zeatin riboside in clubroot tissue:
929 Evidence for trans-zeatin biosynthesis by plasmodiophora brassicae. *Physiologia*
930 *Plantarum* **66:** 245-250.
931
932 Neuhauser S, Kirchmair M, Bulman S, Bass D (2014). Cross-kingdom host shifts of
933 phytomyxid parasites. *BMC evolutionary biology* **14:** 33.
934

Nielsen H (2017). Predicting secretory proteins with signalp. In: Kihara D (ed). *Protein*
function prediction: Methods and protocols. Springer New York: New York, NY. pp
59-73.

Park SW, Kaimoyo E, Kumar D, Mosher S, Klessig DF (2007). Methyl salicylate is a
critical mobile signal for plant systemic acquired resistance. *Science* **318:** 113-116.

Pieterse CM, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SC (2012).
Hormonal modulation of plant immunity. *Annu Rev Cell Dev Biol* **28:** 489-521.

R Core Team (2016). R: A language and environment for statistical computing. R
Foundation for Statistical Computing: Vienna, Austria.

Riascos D, Ortiz E, Quintero D, Montoya L, Hoyos-Carvajal L (2011).
Histopathological and morphological alterations caused by plasmodiophora brassicae
in brassica oleracea l. *Agronomía Colombiana* **29:** 57-67.

Robinson MD, McCarthy DJ, Smyth GK (2010). Edger: A bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26:** 139-
140.

Rolfe SA, Strelkov SE, Links MG, Clarke WE, Robinson SJ, Djavaheri M *et al* (2016).
The compact genome of the plant pathogen plasmodiophora brassicae is adapted to
intracellular interactions with host brassica spp. *BMC Genomics* **17:** 1-15.

Saleh A, Withers J, Mohan R, Marques J, Gu Y, Yan S *et al* (2015). Posttranslational
modifications of the master transcriptional regulator npr1 enable dynamic but tight
control of plant immune responses. *Cell Host Microbe* **18:** 169-182.

Schuller A, Kehr J, Ludwig-Muller J (2014). Laser microdissection coupled to
transcriptional profiling of arabidopsis roots inoculated by plasmodiophora brassicae
indicates a role for brassinosteroids in clubroot formation. *Plant Cell Physiol* **55:** 392-
411.

Schwelm A, Fogelqvist J, Knaust A, Julke S, Lilja T, Bonilla-Rosso G *et al* (2015). The
plasmodiophora brassicae genome reveals insights in its life cycle and ancestry of chitin
synthases. *Scientific reports* **5:** 11153.

Schwelm A, Badstober J, Bulman S, Desoignies N, Etemadi M, Falloon RE *et al*
(2018). Not in your usual top 10: Protists that infect plants and algae. *Mol Plant Pathol*
19: 1029-1044.

Siemens J, Keller I, Sarx J, Kunz S, Schuller A, Nagel W *et al* (2006). Transcriptome
analysis of arabidopsis clubroots indicate a key role for cytokinins in disease
development. *Mol Plant Microbe Interact* **19:** 480-494.

Siemens J, Bulman S, Rehn F, Sundelin T (2009). Molecular biology of
plasmodiophora brassicae. *J Plant Growth Regul* **28:** 245-251.

Siemens J, Gonzalez MC, Wolf S, Hofmann C, Greiner S, Du Y *et al* (2011).
Extracellular invertase is involved in the regulation of clubroot disease in arabidopsis
thaliana. *Mol Plant Pathol* **12:** 247-262.

Song T, Chu M, Lahlali R, Yu F, Peng G (2016). Shotgun label-free proteomic analysis
of clubroot (plasmodiophora brassicae) resistance conferred by the gene rcr1 in brassica
rapa. *Front Plant Sci* **7:** 1013.

Sun Y, Veerabomma S, Abdel-Mageed HA, Fokar M, Asami T, Yoshida S *et al* (2005).
Brassinosteroid regulates fiber development on cultured cotton ovules. *Plant Cell*
Physiol **46:** 1384-1391.

Takahashi H, Muraoka S, Ito K, Mitsui T, Hori H, Kiso A (2001). Resting spore of
plasmodiophora brassicae proliferates only in the callus of clubroot disease-susceptible
turnip but increases the pal activity in the callus of clubroot disease-resistant turnip.
Plant Biotechnology **18:** 267-274.

Usadel B, Poree F, Nagel A, Lohse M, Czedik-Eysenberg A, Stitt M (2009). A guide
to using mapman to visualize and compare omics data in plants: A case study in the
crop species, maize. *Plant Cell Environ* **32:** 1211-1229.

van Verk MC, Bol JF, Linthorst HJ (2011). Wrky transcription factors involved in
activation of sa biosynthesis genes. *BMC Plant Biol* **11:** 89.

Vlot AC, Dempsey DA, Klessig DF (2009). Salicylic acid, a multifaceted hormone to
combat disease. *Annu Rev Phytopathol* **47:** 177-206.

Walerowski P, Gundel A, Yahaya N, Truman W, Sobczak M, Olszak M *et al* (2018).
Clubroot disease stimulates early steps of phloem differentiation and recruits sweet
sucrose transporters within developing galls. *Plant Cell* **30:** 3058-3073.

Wang D, Amornsiripanitch N, Dong X (2006). A genomic approach to identify
regulatory nodes in the transcriptional network of systemic acquired resistance in
plants. *PLoS Pathog* **2:** e123.

Wang X, Wang H, Wang J, Sun R, Wu J, Liu S *et al* (2011). The genome of the
mesopolyploid crop species brassica rapa. *Nat Genet* **43:** 1035-1039.

Wasternack C, Hause B (2013). Jasmonates: Biosynthesis, perception, signal
transduction and action in plant stress response, growth and development. An update to
the 2007 review in annals of botany. *Ann Bot* **111:** 1021-1058.

Wickham H (2009). Ggplot2: Elegant graphics for data analysis. Springer-Verlag: New
York.

Xu L, Yang H, Ren L, Chen W, Liu L, Liu F *et al* (2018). Jasmonic acid-mediated
aliphatic glucosinolate metabolism is involved in clubroot disease development in
brassica napus l. *Front Plant Sci* **9:** 750.
1032

1033 Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y (2012). Dbcan: A web resource for
1034 automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* **40:** W445-451.
1035

1036 Zhang X, Liu Y, Fang Z, Li Z, Yang L, Zhuang M *et al* (2016). Comparative
1037 transcriptome analysis between broccoli (brassica oleracea var. Italica) and wild
1038 cabbage (brassica macrocarpa guss.) in response to plasmodiophora brassicae during
1039 different infection stages. *Frontiers in Plant Science* **7**.
1040

1041 Zhao Y, Bi K, Gao Z, Chen T, Liu H, Xie J *et al* (2017). Transcriptome analysis of
1042 arabidopsis thaliana in response to plasmodiophora brassicae during early infection.
1043 *Front Microbiol* **8:** 673.
1044

1045