1

Fitness Effects of Somatic Mutations Accumulating during Vegetative Growth

Mitchell B. Cruzan[1, 2], Matthew A. Streisfeld[3], and Jaime A. Schwoch[1]

[1]Department of Biology, Portland State University, Portland, OR 97207

[3]Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403

[2]Author for correspondence: cruzan@pdx.edu

Abstract

The unique life form of plants promotes the accumulation of somatic mutations that can be passed to offspring in the next generation, because the same meristem cells responsible for vegetative growth also generate gametes for sexual reproduction. However, little is known about the consequences of somatic mutation accumulation for offspring fitness. We evaluate the fitness effects of somatic mutations in *Mimulus guttatus* by comparing progeny from self-pollinations made within the same flower (autogamy) to progeny from self-pollinations made between stems on the same plant (geitonogamy). The effects of somatic mutations are evident from this comparison, as autogamy leads to homozygosity of a proportion of somatic mutations, but progeny from geitonogamy remain heterozygous for mutations unique to each stem. In two different experiments, we find consistent fitness effects of somatic mutations from individual stems. Surprisingly, several progeny groups from autogamous crosses displayed increases in fitness compared to progeny from geitonogamy crosses, indicating that beneficial somatic mutations were prevalent in some stems. These results support the hypothesis that somatic mutations accumulate during vegetative growth, but they are filtered by different forms of selection that occur throughout development, resulting in the culling of expressed deleterious mutations and the retention of beneficial mutations.

Key words: Acquired mutations, autogamy, autogamy depression, cell lineage selection, dominance, *Erythranthe guttata*, geitonogamy, *Mimulus guttatus*, somatic mutations.

Introduction

Mutation is the source of variation for evolution and adaptation, but organisms differ in whether mutations originating during gamete formation (meiosis) or somatic growth (mitosis) contribute to heritable variation. For the vast majority of organisms, including viruses, unicellular microbes, and some multicellular eukaryotes, sexual reproduction is rare or absent. In these organisms, mutations can occur during mitotic cell replication, and the primary mechanism for adaptation and diversification occurs via selection on cell lineages without recombination. By contrast, acquired mutations occurring during somatic growth of animals are not heritable. This is because in metazoans (but possibly excepting corals; Schweinsberg et al. 2014; Barfield et al. 2016), the germline is determined early in development and relatively few cell divisions occur before the formation of gametes (the Weismann Barrier; Buss 1983). Consequently, heritable mutations only occur in animals during the development of gonads and gametes.

Plants differ from animals and microbes, because mutations contributing to heritable variation can arise both during gamete formation and somatic growth (Antolin and Strobeck 1985; Klekowski and Godfrey 1989). This is due to the fact that plants lack a separate germline; the same germ cells within apical meristems subtend both vegetative growth and gamete development. Consequently, plants have the capacity to accumulate somatic mutations during vegetative growth that can be passed to the next generation (McKnight et al. 2002; Klekowski 2003; Schultz and Scofield 2009; Bobiwash et al. 2013; Dubrovina and Kiselev 2016; Watson et al. 2016; Schmid-Siegert et al. 2017; Yu et al. 2020). This aspect of plant biology is well known (Monro and Poore 2009; Ally et al. 2010; Reusch and Bostrom 2011), and somatic mutation accumulation has been important in agriculture, where the origin of many clonally-derived varieties of fruits, including citrus, apples, and wine grapes, have been cultivated by grafting from genetically differentiated bud tips (Aradhya et al. 2003; McKey et al. 2010; Miller and Gross 2011; Vezzulli et al. 2012; Jarni et al. 2015; Pelsy et al. 2015). However, there is disagreement over the extent and evolutionary importance of somatic mutation accumulation in plants (Schultz and Scofield 2009; Burian et al. 2016; Watson et al. 2016; Kuhlemeier 2017; Schmid-Siegert et al. 2017; Plomion et al. 2018; Hanlon et al. 2019).

In general, it is difficult to assess the effects of somatic mutations accumulating during growth because clonal cell lineages lack recombination, and the fitness effects of beneficial mutations can be muted by the co-occurrence of detrimental genetic variants (Hill-Robertson Effect; Selective Interference; Hill and Robertson 1966; Fogle et al. 2008). Although sexual reproduction does allow mutations to segregate among progeny, it is difficult to detect the fitness effects of somatic mutations in

eukaryotes with separate sexes. This is because the genomes of two individuals are combined, so

individual somatic mutations that occur in one individual will always be heterozygous in offspring (Long

et al. 2015). A more effective approach would be to make self-fertilizations in a hermaphroditic species,

so a proportion of offspring would be homozygous for novel somatic mutations. In this study, we make

crosses within individual clones of hermaphroditic plants to assess the fitness effects of inherited

mutations that accumulated during somatic growth.

Hermaphroditic plants have several advantages for the study of somatic mutations. Separate

stems on the same plant contain multiple germ cell lineages that are derived from the same zygote, but

they may differ for somatic mutations that have accumulated during stem elongation. To produce

recombinant progeny segregating for somatic mutations unique to each stem, crosses can be made

either within the same flower (autogamy), or between flowers on separate stems of the same plant

(inter-ramet geitonogamy – hereafter referred to simply as geitonogamy; Fig. 1). These crosses are both

self-fertilizations, but the offspring will differ in the complement of somatic mutations that they inherit.

Mutations unique to a stem will be homozygous in 25% of autogamous progeny, while none of the

progeny from geitonogamous crosses will be homozygous, and half will be heterozygous for unique

somatic mutations. Thus, the average fitness effects of somatic mutations unique to each stem can be

evaluated by comparing progeny generated by autogamous and geitonogamous crosses of the same

stem. The potential for somatic mutations to accumulate during vegetative growth, and the ability to

study their effects among recombinant progeny, make plants an attractive model for the study of

fundamental processes of clonal evolution.

Plants grow from the division of a population of undifferentiated meristem cells within the stem

tip that is known as the central zone. These germ cell lineages go on to produce future stem, leaf, and

reproductive tissues (flowers). As plants grow, individual ramets of the same genet (separate stems or

vegetatively propagated plants) can continue to become differentiated as they accumulate somatic

mutations (Fig. 2). Therefore, when we consider the potential for meiotic and somatic mutations to

contribute to the total mutational load of plant populations – particularly for long-lived plants – it

becomes evident that not all of the mutations occurring during a plant's lifespan are passed to the next

generation (Cruzan 2018 pages 86-98). Indeed, plants have mechanisms of "developmental selection"

(Buchholz 1922) that occur during vegetative growth and reproduction to filter the set of mutations that

are inherited by progeny. For example, because new somatic mutations occur as a single copy within the

diploid genome, their immediate fitness effects will depend on the combination of their fitness effects

and their expression in the heterozygous state. Mathematical models have demonstrated that multiple

unexpressed mutations (i.e. neutral and recessive deleterious mutations) are likely to accumulate as

germ cells divide, so any selective cell deaths that occur during somatic growth will reduce the total

mutational load and alter the composition of mutations carried by the germ cell population (Otto and

Orive 1995; Otto and Hastings 1998). Thus, lineages that carry expressed deleterious mutations causing

slower rates of division can be eliminated as they are replaced by cell lineages displaying more rapid

growth (Otto and Orive 1995; Otto and Hastings 1998; Orive 2001; Elena and Lenski 2003; Greaves and

Maley 2012; Long et al. 2015). (Fig. 2A). More rarely, expressed beneficial mutations will arise. Similar to

the process of clonal evolution in mammalian cancer, any mutations that elevate the rate of cell division

will tend to increase in frequency until they have replaced the entire germ cell population (clonal

selective sweep; Nowell 1976; Lang et al. 2013; Fig. 2B green arrows). This process – referred to as cell

lineage selection – will lead to some ramets containing a prevalence of deleterious somatic mutations,

while others could possess beneficial mutations. The potential for these clonal selective sweeps to occur

during vegetative growth has been confirmed by the observation of somatic mutations occurring at

frequencies near 50% (heterozygosity) within the stem tissue (Yu et al. 2020; Schwoch et al. unpublished

data). Hence, germ cell populations in plant apical meristems have the potential to undergo clonal

evolution during vegetative growth.

In addition to cell lineage selection, some proportion of recessive deleterious mutations may be

eliminated during the haploid life stage due to pollen tube attrition and pollen competition

(Gemetophytic Selection; Mulcahy 1979; Cruzan 1989; Mable and Otto 1998; Armbruster and Rogers

2004; Arunkumar et al. 2013; Harder et al. 2016). Finally, a portion of deleterious mutations will be

homozygous in zygotes, which can lead to higher rates of seed and fruit abortion (Selective Embryo

Abortion; Husband and Schemske 1995; Korbecka et al. 2002), thereby increasing the average fitness of

surviving offspring (Cruzan and Thomson 1997; Mena-Alĺ and Rocha 2005). The sum effects of these

processes of developmental selection, which includes cell lineage selection, gametophytic selection, and

selective embryo abortion, may filter the set of mutations that enters the next generation. Indeed, the

effects of deleterious somatic mutations often are apparent as higher rates of embryo abortion after

autogamous compared to geitonogamous pollinations, which is referred to as autogamy depression

(Schultz and Scofield 2009). While autogamy depression for seed and fruit abortion has been observed

in several species (reviewed in Bobiwash et al. 2013), no previous study has evaluated the fitness effects

of somatic mutations inherited by progeny. The question we address here is whether somatic mutations

that characterize individual ramets have measurable fitness effects in the offspring generation.

In this study, we use autogamous and geitonogamous self-pollinations to estimate the fitness

effects of somatic mutations segregating in offspring of *Mimulus guttatus* DC (*Erythranthe guttata* G.L.

Nesom; Phrymaceae). Populations of *M. guttatus* display a wide range of life histories – from annuals to

herbaceous perennials that outcross to varying degrees (Wu et al. 2008). We use perennial *M. guttatus*

plants that produce substantial vegetative growth prior to initiation of flowering. These plants are easily

propagated from rosettes, and selfing produces substantial numbers of seeds to allow for statistical

comparisons. In two separate experiments, we characterized the fitness effects of somatic mutations in

perennial *M. guttatus* plants in controlled environments.

We begin by describing the analytical approaches we use, followed by providing specifics of the

two experiments that we conducted. We show that somatic mutations accumulate during stem growth,

as progeny from autogamous crosses have greater variance in mean fitness compared to progeny from

geitonogamous crosses. We further reveal evidence for the presence of beneficial mutations in some

stems, which indicates that developmental selection may be responsible for filtering mutations and

modifying the distribution of fitness effects transmitted to progeny. The results from these experiments

provide the first evidence that inherited somatic mutations can have fitness consequences for plants in

the next generation, which suggests that somatic mutations may be an additional source of genetic

variation that can be used for adaptation.

Approach

We made autogamous and geitonogamous self-pollinations across multiple stems and estimated the

fitness effects of somatic mutations segregating in progeny. To increase the chances of detecting

somatic mutations that impacted fitness, we exposed plants to novel environments. According to

Fisher's geometric model, displacement from a fitness optimum increases the chances that any

mutation would move a genotype in a direction that increases fitness (Fisher 1930; Stearns and Fenster

2016). Furthermore, we evaluated fitness of seedlings under novel selection regimes in the greenhouse.

Exposing parental plants and seedlings to the same controlled environments is common in experimental

designs that test for the fitness effects of mutations accumulating across generations (Shaw et al. 2002;

Baer et al. 2007; Halligan and Keightley 2009), or in our case, for observing the fitness effects of somatic

mutations arising within a single generation.

For a diploid plant, we can assume that somatic mutations ($a \rightarrow a'$) will be in the heterozygous

state when they first appear. For progeny arising by autogamy, a mutation will segregate as 25%

homozygous ($a'a'$), 50% heterozygous (aa'), and 25% the original homozygote (aa). By contrast, because

progeny from geitonogamous crosses will be segregating for mutations that are unique to each stem,

half of them will be carrying mutations in the heterozygous state, and none of the progeny will be

homozygous for mutations that arose in a single stem. The fitness effects of somatic mutations

accumulating during vegetative growth have been quantified in terms of "autogamy depression" (δ_{AD})

using the formula; $\delta_{AD} = 1 - w_A/w_G$ (Schultz and Scofield 2009; Bobiwash et al. 2013), where w_A and w_G

are the relative fitnesses of progeny derived from autogamous and geitonogamous crosses, respectively.

Tests of autogamy depression have been applied previously only to estimates of fitness based

on fruit and seed set, and it generally assumes that fitness from autogamous crosses will be lower than

from geitonogamy. However, somatic mutations that are transmitted to progeny can affect fitness in the

offspring generation across different life stages, and beneficial mutations are possible. Thus, for each

stem, we examined the fitness effects of somatic mutations unique to stem k as: $\delta_{AD(k)} = \overline{w}_{k(A)} - \overline{w}_{k(G)}$,

where $\overline{w}_{k(A)}$ and $\overline{w}_{k(G)}$ are the average fitnesses of the progeny from autogamous and geitonogamous

crosses from stem k, respectively. We evaluate δ_{AD} for each stem separately, because we expect that

individual stems will have unique complements of somatic mutations and thus differ in their fitness

effects. By controlling for environmental effects with a common garden design, the parameter $\delta_{AD(k)}$ and

its sign describe the average magnitude and overall direction of the fitness effects of all expressed

somatic mutations that have been transmitted to the next generation. To evaluate the reliability of $\delta_{AD(k)}$

as an estimate of the fitness effects of somatic mutations, we simulated variation in the strength of

selection and dominance of somatic mutations across multiple pairs of stems (ramets). We determined

that values of $\delta_{AD(k)}$ were accurate across a range of values and assumptions for the effects of multiple

mutations (Appendix 1).

To provide additional confirmation of the $\delta_{AD(k)}$ estimates, we developed an alternative approach

for estimating the fitness effects of somatic mutations that was based entirely on the variance in mean

fitness among autogamous progeny. Individual germ cell lineages within the meristem may possess

unique sets of somatic mutations (Yu et al. 2020; Schwoch et al. in preparation), and flowers on the

same stem can develop from only a portion of the germ cells present in the meristem (McManus and

Veit 2002). Hence, no pair of flowers within a stem is likely to contain exactly the same complement of

mutations. Therefore, we generated an additional estimate of the fitness effects of somatic mutations

unique to a stem that is based on the variance in fitness of offspring from autogamous crosses. If

somatic mutations affect offspring fitness, the variance in fitness should be greater for progeny groups

from autogamy than from geitonogamy from the same stem, as long as mutations do not have complete

expression in heterozygotes (i.e. $h < 1.0$; Appendix 2). Somatic mutations that are unique to stems will

have different effects on the variance in fitness among progeny from autogamy depending on their

fitness (w) and their expression in the heterozygous state (h). Our analytical results demonstrated that

the large majority of variance in fitness among progeny from autogamy (as measured by the standard

deviation, SD) could be explained by the fitness effects of a mutation rather than the dominance level

(Appendix 2; Fig. S1). Moreover, the derived relationship allows us to estimate the fitness effects of

mutations based on the standard deviation in fitness according to the equation: $w_{SD} = cSD$, where c is

the slope of the linear relationship between w_{SD} and SD. For dominance levels of $h = 0.0$ and 1.0, $c =$

2.31, and the slope reaches a maximum of 3.01 when $h = 0.5$ (Appendix 2; Fig. S1). Thus, the variance in

fitness among progeny from autogamous crosses provides a good approximation of the magnitude of

the fitness effects (w) for somatic mutations accumulating in each stem. These two approaches are

independent of each other, and therefore provide an important means to confirm our estimates of the

average fitness effects of somatic mutations present in individual stems (Appendix 1 and 2).

203   From the considerations above, we can make three specific predictions that allow us to test for

the fitness effects of somatic mutations that occur during stem growth and are passed to offspring. First,

there should be greater variation among progeny groups from autogamy compared to those from

geitonogamy (i.e. as indicated by the stem x cross type interaction for progeny fitness). Second, the

average fitness of progeny from autogamy should be different from the progeny of geitonogamous

crosses from the same stem. The difference in the fitness of progeny from autogamous vs.

geitonogamous crosses provides an approximate estimate of the sign and magnitude of the average

fitness effects of somatic mutations for individual stems. In addition to influencing the mean fitness,

somatic mutations will increase the variance in fitness among progeny from autogamy. Consequently,

the third prediction we can make is that the mean fitness effects of somatic mutations will scale linearly

with the within-family standard deviation in fitness among progeny from autogamous crosses (Appendix

2). In general, we predict that if somatic mutations with fitness effects accumulate during vegetative

growth, then the mean progeny fitness should differ between autogamous and geitonogamous crosses,

and the variance in fitness of progeny from autogamous crosses would be greater than for progeny from

geitonogamous crosses. Below, we test these predictions using two different experiments with *Mimulus*

guttatus.

First Experiment

221   *Methods*–This experiment originally was designed to test the effects of somatic mutations on

seed set and ovule abortion (autogamy depression; Schultz and Scofield 2009; Bobiwash et al. 2013), so

we used multiple plants from several populations to obtain a large number of fruits. We grew plants of

M. guttatus in the Research Greenhouse facility at Portland State University from seed collected in July

2013 from three different populations in northern Oregon (Jackson Bottom Wetlands–JB: 45.501794 N, -

122.98776 W; and two from Saddle Mountain–SMB: 45.9861 N, -123.6859 W, and SMC: 45.9634 N, -

123.6837 W). We assumed that greenhouse conditions were different enough from field environments

to provide a novel selection regime. In August 2013, seeds were cold stratified on moist paper towels at

2°C for 30 days prior to being sown in soil. Seedlings were transplanted to pots (approximately 10 x 10 x

12 cm) and grown for seven months before the application of pollination treatments. Temperature was

maintained between 21–26°C during the day, and 15–21°C at night. Supplemental HID lights ran for 12

hours a day when the seedlings first emerged, and 14 hours a day during adult growth.

After plants became established and began producing multiple stems, we conducted

autogamous and geitonogamous self-pollinations using flowers on several stems (15 to 20 cm in length)

from two plants from each of four maternal families representing each of the three populations. Flowers

from pairs of stems on individual plants were reciprocally crossed (geitonogamy), or individual flowers

from these same stems were self-pollinated (autogamy). A total of 139 pollinations were conducted

across two treatments: limited (pollen was applied to stigmas with one touch from a plastic pipette tip)

or excess (where the stigma surface was coated with pollen). Pollinations were conducted on 12

different days (pollination date) over several weeks in July 2014. Mature fruits were collected and

placed individually into paper envelopes, and their contents were examined under a Leica MZ-16

stereoscope. The first 100 ovules from each fruit were categorized as filled seeds (brown, almond-

shaped), unfertilized ovules (small, flattened and light-colored), or aborted (larger than unfertilized,

dark-colored, shriveled). Ovules that were flattened and appeared to lack endosperm were assumed to

be unfertilized or aborted and were not used in germination tests. Seed set and ovule abortion were

analyzed using ANOVA models with the GLM procedure of SAS (SAS 2008), with population, maternal

plant nested within population, and pollination date as random effects, and cross type (autogamous or

geitonogamous) and pollination treatment as fixed effects. Seed set and ovule abortion data were

approximately normal so were not transformed prior to analysis.

We assessed the fitness of progeny arising by autogamy and geitonogamy in the same

greenhouse environment that was used to grow the parental plants. Seedlings from a subset of ten

maternal plants that had fruits from both cross types and at least 20 filled seeds were sown in soil and

transplanted to 36-cell trays (blocks) in September in a randomized incomplete block design. After three

months of growth, the progeny were scored for survival, and above ground biomass was measured after

drying at 60°C for at least 24 hours. Since all seedlings germinated within a few days of each other, the

final biomass is an estimate of growth rate. The fitness of each progeny was estimated as its final

biomass, which was log transformed and weighted by the survival frequency of progeny from the same

cross. Growth rate is considered to be an appropriate estimate of fitness for perennials (Younginger et

al. 2017). Furthermore, we evaluated fitness of seedlings under novel selection regimes in the

greenhouse rather than under field conditions, which allowed us to control for environmental

differences by having parental plants and seedlings exposed to the same conditions. These estimates

were rescaled relative to the maximum value from all crosses, so that $\bar{w}$ ranged from 0 to 1 across

progeny from all crosses. Data were analyzed to estimate the mean fitness for progeny from autogamy

($\bar{w}_{k(A)}$) and geitonogamy ($\bar{w}_{k(G)}$) from each stem, and the variance in fitness (as measured by the

standard deviation; SD) for each group of progeny produced by autogamy and geitonogamy from single

stems.

Results–Levels of seed set and germination were similar between cross types, but ovule abortion

was greater in autogamous crosses. The mean number of seeds produced after autogamous (56.72

±2.24, N = 87) and geitonogamous (57.52 ±3.42, N = 52) pollinations was similar (Cross in Table S2), and

there was little variation among populations (Pop in Table S2), maternal plants nested within population

(Mat(Pop) in Table S2), and pollination dates (Date in Table S2). Limited pollinations produced fewer

seeds than excess pollinations (Pollen in Table S2), but the effect of pollen dosage in the overall model

was not significant. However, the mean number of aborted ovules was greater for autogamous (17.36

±1.27, N = 87) than for geitonogamous (13.90 ±1.70, N = 52) pollinations (Cross in Table S3), which is

consistent with the predicted effects of autogamy depression (Schultz and Scofield 2009; Bobiwash et al.

2013). There also were significant differences among populations in the level of ovule abortion (Pop in

Table S3), which perhaps reflects differences in historical inbreeding and levels of genetic load. There

were significant differences in ovule abortion among maternal plants nested within population (Mat

(Pop) in Table S3), but not for pollination treatments (Pollen in Table S3) or pollination dates (Date in

Table S3). The overall model for ovule abortion was significant (Table S3). Seed germination was similar

between autogamous (mean = 13.03 out of 20 planted per fruit) and geitonogamous crosses (mean =

13.28; $F_{1,101}$ = 0.16, P = 0.694). Of the 354 seedlings that germinated, 202 survived, and interestingly,

survival was higher for progeny from autogamy (67%) compared to geitonogamy (50%; chi-square =

12.03, P = 0.0005). For the surviving plants that flowered by the end of the experiment, flower

production was correlated with growth rate (r = 0.58, P < 0.001, N = 282).

The average fitness of progeny from autogamy and geitonogamy varied widely among individual

stems. Consistent with our first prediction for the effects of somatic mutations on offspring fitness,

there was a significant interaction between cross type and stem identity (Table S4). Little of the variation

in progeny fitness was explained by differences among stems (Mat in Table S4), but larger amounts were

explained by cross type (Cross in Table S4) and among blocks (Tray in Table S4). When data were

analyzed separately for each cross type, we found that mean fitness of progeny from autogamy varied

among stems to a much greater degree ($F_{8,130} = 4.44$, $P < 0.0001$) than progeny from geitonogamous

crosses ($F_{8,70} = 1.47$, $P = 0.1866$); a result that supports our first prediction for the effects of somatic

mutations on progeny fitness. These data are analyzed further along with data from the second

experiment described below.

Second Experiment

The observation of greater variation in fitness among stems for progeny from autogamous

crosses in the first experiment is consistent with the hypothesis that somatic mutations arising during

vegetative growth are transmitted and can have substantial fitness effects in the next generation.

Furthermore, the results from the first experiment do not support the view that somatic mutations

during vegetative growth are rare (Burian et al. 2016; Groot and Laux 2016; Watson et al. 2016;

Kuhlemeier 2017; Schmid-Siegert et al. 2017), but they are consistent with several studies

demonstrating that somatic mutations can be inherited by offspring in the next generation (Klekowski

and Godfrey 1989; Schultz and Scofield 2009; Bobiwash et al. 2013). However, the first experiment was

not designed specifically to compare the effects of autogamous and geitonogamous pollinations on

offspring fitness. Thus, to confirm results from the first experiment, we performed a second experiment

to provide a more robust assessment of the effects of autogamous and geitonogamous self-pollination

on the fitness of offspring. The second experiment used a single clone of *M. guttatus* to make

comparisons between autogamous and geitonogamous pollinations paired at the same node. The plant

chosen was a self-compatible perennial that displayed vigorous vegetative growth. We exposed plants

to novel conditions using high salinity under hydroponics cultivation, which increased the potential that

mutations would have fitness consequences. Moreover, our recent genomic results confirm more low-

frequency somatic mutation among plants exposed to high salt stress compared to the control

hydroponics treatment without added salt (Schwoch et al. in preparation). This experimental design

provided a more refined test of the hypothesis that the fitness effects of somatic mutations differed in

the offspring of autogamous and geitonogamous crosses.

Methods–To assess the fitness effects of mutations that accumulated during vegetative growth,

a single plant (genet BV, obtained from Willamette Gardens native plant nursery, Corvallis, OR) was

vegetatively propagated to generate 12 plants (ramets) that were exposed to high salinity and control

conditions. Plants were grown in pea gravel (4–8 mm) in pots placed in four 53 L tubs using a flood and

drain hydroponics system (flooding at 15 min intervals). Two tubs had no added salt and were used as

controls, and two tubs had high salinity. The initial salt concentration in the high salinity treatment was

5 mM but increased weekly to 25 mM after plants became established. Salt concentrations were

monitored using a conductivity meter to ensure stable concentrations. To provide nutrients, 30 ml of

hydroponics fertilizer (FloraGrow, Planet Natural, Bozeman, MT) was added per tub. During the course

of the experiment, some plants grew substantially. We imposed selection to favor the fastest growing

ramets over the next three months by repeatedly removing single rosettes and transplanting them back

into the hydroponics system up to four times.

To promote stress recovery, plants were transplanted to soil for six months, which included a

two-month vernalization period in a growth chamber (4°C and 8 h light; Conviron E8, Controlled

Environments Ltd., Winnipeg, Manitoba, Canada). After vernalization, plants were returned to the

greenhouse to induce flowering. Autogamous and geitonogamous pollinations were made to pairs of

flowers at single nodes or consecutive nodes (seven nodes and 14 pollinations total) on the largest

ramets in each of the control and high salt treatments. To account for somatic mutation turnover that

may occur due to the effects of cell lineage selection during stem growth, we compared progeny from

autogamous and geitonogamous pollinations at pairs of flowers from the same node. Without *a priori*

knowledge of the expression of somatic mutations in heterozygotes, it is difficult to determine the best

pollen donor for geitonogamous crosses. Consequently, we opted to generate progeny from

geitonogamy by pollinating flowers with pollen from a potentially more genetically divergent ramet

from the other treatment (i.e. salt pollinated with control pollen, and control pollinated with salt

pollen). The fruits were collected, and the total number of aborted and mature seeds and unfertilized

ovules were counted under a dissecting microscope. Seeds were planted in soil in trays with three seeds

per cell. Seeds were cold stratified in moist soil for three weeks before they germinated in the

greenhouse.

To determine whether seedlings from autogamous crosses from ramets exposed to salt stress

showed improved performance under the same conditions, all progeny were exposed to high salinity.

After germination and establishment in soil, seedlings from autogamous and geitonogamous crosses

from control and salt stress ramets were transplanted into pots filled with pea gravel and subjected to

high salt in the hydroponics system, as described above. A total of 239 seedlings from 11 fruits (five

autogamous and six geitonogamous) were randomly and evenly distributed among 12 hydroponic tubs

to ensure equal representation across blocks (tubs). Plant size was measured as the product of the

length and width of vegetative spread after two months of growth and was used as a proxy for overall

plant performance. Since plants germinated within a few days of each other, plant size represents a

good estimate of growth rate. Salt concentration increased from 10 mM - 37.5 mM over the course of

the experiment to induce mortality (~57% across all progeny groups). Fitness was estimated as plant size

(log transformed to improve normality) weighted by the survival frequency for progeny from the same

cross.

Results–Similar to experiment 1 and consistent with our first prediction for the effects of

somatic mutations, there was greater variation in the fitness of progeny from autogamous compared to

geitonogamous crosses, as indicated by a significant interaction between cross type and stem/node

identity (Stem/Node*Cross in Table S5). There was no consistent difference in progeny fitness between

autogamous and geitonogamous pollinations when averaged across stems (Cross in Table S5), but there

were larger differences among individual stems/nodes (Stem/Node in Table S5). Also consistent with our

first prediction, there was significant variation in progeny fitness among stems ($F_{2,89}$ = 3.76, P = 0.028)

and nodes (nested within stems; $F_{3,89}$ = 2.59, P = 0.0596) for autogamous, but not geitonogamous

crosses ($F_{1,136}$ = 0.63, P = 0.431 and $F_{4,136}$ = 1.19, P = 0.317 for stems and nodes, respectively). There were

significant differences in average size among the blocks (Tray in Table S5), but there was no consistent

difference in the performance of progeny based on the treatment history of their maternal ramets ($F_{1,126}$

= 0.02, P = 0.888), and there was no interaction between cross type and historical treatment ($F_{1,136}$ =

0.64, P = 0.423). Three nodes were dropped from further analysis, because one of the paired fruits

aborted or produced fewer than five seedlings. However, similar to experiment 1, and consistent with

the hypothesis that somatic mutations accumulate during stem growth, the variance in fitness among

progeny from autogamy among stems was greater than for geitonogamy. Since results from the two

experiments were similar, the data for four nodes from experiment 2 were analyzed together with the

first experiment, as described in the next section.

Fitness Estimates

Results from the two experiments described above are consistent with the hypothesis that somatic

mutations unique to individual stems can have demonstrable effects on the fitness of progeny when a

proportion is made homozygous by autogamous self-pollination. In both experiments, the variance in

fitness among progeny groups from autogamy was significantly greater than among progeny from

geitonogamy from the same set of stems. As the results from these two experiments are qualitatively

similar, we estimated the fitness effects of somatic mutations in each stem (ten comparisons of

autogamy and geitonogamy from Experiment 1 and four from Experiment 2, for a total of 14

comparisons). Specifically, we tested the second and third predictions described above. Since 25% of

somatic mutations unique to a stem will be homozygous in autogamous progeny, we expect that their

fitness effects will be greater than for progeny from geitonogamous crosses on the same stem

(providing an estimate of $\delta_{AD(k)}$). In addition, the within-family variance in fitness among progeny should

be greater after autogamy compared to geitonogamy (providing an estimate of w_{SD}). These estimates

of the fitness effects of somatic mutations should be similar, so we assessed whether there was a

positive relationship between the variance in fitness among progeny from single fruits produced by

autogamous pollination (w_{SD}) and the fitness effects of somatic mutations unique to each stem

estimated from the average fitness of progeny from autogamous and geitonogamous crosses to the

same stem ($\delta_{AD(k)}$).

Estimates of the average fitness effects of somatic mutations unique to each stem were made

based on the fitness of progeny from autogamous ($\overline{w}_{k(A)}$) and geitonogamous ($\overline{w}_{k(G)}$) crosses (Table S1

in Appendix 3). In both experiments, fitness was estimated as growth rate weighted by the survival of

seedlings from the same fruit and scaled to a maximum value of 1.0. We have demonstrated that

growth rate is correlated with flower production, and it has been shown to be a good predictor of

fitness among perennial plants (Younginger et al. 2017). The average fitness effects of mutations ($\delta_{AD(k)}$)

for each stem (first experiment) or stem/node combination (second experiment) were calculated as the

difference in fitness between progeny from autogamous and geitonogamous crosses, as described

above ($\delta_{AD(k)} = \overline{w}_{k(A)} - \overline{w}_{k(G)}$). We observed extensive variation in $\delta_{AD(k)}$ among stems, with nine stems

that were significantly different from zero (Fig. 3). Four of the stems had average fitness effects of

somatic mutations that were positive, suggesting a net beneficial effect of somatic mutations

transmitted to offspring. In addition, the average fitness of progeny from autogamous and

geitonogamous crosses on the same stems or nodes was positively correlated (Fig. S2; Table S6). Note

also that the $\delta_{AD(k)}$ estimates that deviated the most from zero generally had the highest (or lowest)

$\overline{w}_{k(A)}$ values relative to all of the stems (Fig. S3). However, this was not always the case, as one stem

(stem 9) with a value of $\delta_{AD(k)}$ close to zero produced progeny with relatively high fitness after both

autogamous and geitonogamous pollination.

To make an independent estimate of the average fitness effects for mutations unique to each

stem or node, we used the standard deviation in progeny fitness from autogamous crosses based on the

relationship $w_{SD} = cSD$, where $c = 3$, which assumes dominance close to $h = 0.5$ (qualitatively similar

results are obtained if we choose other values of c between 3 and 2.31; Appendix 2). Because the sign of

w_{SD} could not be inferred directly from this approach, we used the sign estimated from the $\delta_{AD(k)}$ method

(Fig. 4; Appendix 2). Our simulations show that the sign of $\delta_{AD(k)}$ is almost always estimated correctly

(Appendix 1). For stems with values of $\delta_{AD(k)}$ greater than zero, there was a strong positive relationship

between $\delta_{AD(k)}$ and estimates of w_{SD} made from the within-family variation among progeny from

autogamy (Fig. 4). In contrast, the relationship for negative values of $\delta_{AD(k)}$ appeared to be driven largely

by a single observation. Note that this observation was not supported by a similarly high value of w_{SD},

and the remaining negative fitness effects were more modest based on both estimates. This one highly

negative estimate of $\delta_{AD(k)}$, may be due to a large influence of mutations from the second stem on the

fitness of geitonogamous progeny (note that this stem had the highest estimate of $\overline{w}_{k(G)}$; Fig. S2). It is

also notable that the variance in fitness for progeny from autogamy did not decline to zero for values of

$\delta_{AD(k)}$ close to zero, which could be due to the presence of both beneficial and detrimental mutations,

and possibly genetic background effects (i.e. epistasis), but it may also reflect uncontrolled

environmental variation. Overall, the results from both approaches reveal consistent estimates of the

average fitness consequences associated with the accumulation of somatic mutations in individual

stems.

Discussion

In this study, we have shown that somatic mutations can accumulate during vegetative growth and have

substantial fitness effects on plants in the next generation. Consistent with previous work (Bobiwash et

al. 2013), we observed significant autogamy depression in the form of greater ovule abortion in

autogamous relative to geitonogamous crosses. However, we also detected more variation in survival

and growth rate for progeny from autogamous compared to geitonogamous self-pollinations, which

provides the first evidence that somatic mutations that accumulated during vegetative growth can have

demonstrable effects on the fitness of plants in the next generation. Both the differences in the mean

fitness between progeny from autogamy and geitonogamy ($\delta_{AD(k)}$) and variation in fitness of autogamous

progeny (w_{SD}) provided consistent estimates of the average fitness effects of somatic mutations

segregating in progeny. We found evidence for the effects of beneficial mutations in progeny from some

stems, with estimates of $\delta_{AD(k)}$ and w_{SD} exceeding 0.1 in four cases, while estimates for negative fitness

effects were more modest (mostly > -0.15). Furthermore, the observation that progeny from

autogamous crosses occasionally had higher fitness than progeny from geitonogamous crosses argues

for the accumulation of beneficial mutations and implies that many deleterious mutations are culled by

the various types of developmental selection prior to seed dispersal. These results imply that even

though numerous somatic mutations are generated during vegetative growth, many of them are filtered

due to different forms of developmental selection, which results in a shift in the distribution of fitness

effects for the mutations that are passed to the next generation.

Somatic mutations accumulating during vegetative growth had an overall positive effect for four

of the stems tested. This result is contrary to widely held views that the accumulation of beneficial

somatic mutations should be exceedingly rare (Crow 1993; Charlesworth and Willis 2009). However, an

explanation for these findings is afforded by the unique biology of plants; somatic mutations in germ

cells in the central zone of the apical meristem could contribute to higher rates of division for some cell

lineages over others (Otto and Orive 1995; Otto and Hastings 1998). This hypothesis is supported by the

observation that a minority of somatic mutations occur at high frequencies in meristem tissue (Yu et al.

2020; Schwoch et al. unpublished data), suggesting that clonal selective sweeps have occurred during

stem elongation. Thus, cell lineage selection has the potential to retain beneficial somatic mutations, as

individual cell lineages divide at faster rates resulting in clonal evolution during vegetative growth.

Although there may be few opportunities for beneficial changes to alter basic cellular

metabolism, it is becoming apparent from experimental evolution studies with microbes that even basic

aspects of cellular metabolism can be sensitive to environmental conditions, which can increase the

chances that mutations in clonal populations are beneficial (e.g., Lang et al. 2013; Lee and Marx 2013;

Maharjan et al. 2015). In this regard, cell lineage selection in a plant meristem represents a potentially

powerful forum for the removal of deleterious somatic mutations while favoring the retention of

beneficial ones. In addition, gametophytic selection and selective embryo abortion can act as prominent

additional filters (Mulcahy 1979; Cruzan 1989; Mable and Otto 1998; Armbruster and Rogers 2004;

Arunkumar et al. 2013; Harder et al. 2016), but they are most likely to have effects by culling deleterious

mutations. Regardless, the combined effects of these different forms of developmental selection appear

to have had a considerable effect on filtering of somatic mutations under controlled conditions in the

greenhouse, such that the distribution of fitness effects among stems has shifted to include more

beneficial mutations than expected. Future studies should test whether similar findings are found under

field conditions, which could indicate a prominent role for somatic mutation in local adaptation.

An alternative explanation for the observed fitness differences is that exposure to

environmental stress has induced heritable epigenetic modifications (Quadrana and Colot 2016).

However, epigenetic modifications are considered to be consistent in direction and predictable, as they

are hypothesized to represent an adaptive response to historic exposure to similar stressors (Baulcombe

and Dean 2014; Itabashi et al. 2018). The results of the experiments described here are unlikely to

support a role for epigenetics, because fitness responses in the next generation were inconsistent in

direction and magnitude, and they were not predictable based on environmental exposure of the parent

stem. Indeed, among stems and nodes, the mean fitness of progeny from autogamy displayed both

increases and decreases compared to the geitonogamy controls, which is consistent with the hypothesis

that individual ramets are accumulating unique complements of somatic mutations. Furthermore, these

conclusions are supported by the observation of numerous somatic variants in the transcriptomes of

meristem tissue from multiple ramets derived from a single genet (Yu et al. 2020; Schwoch et al.

unpublished data). Thus, the results of the current study appear to indicate that somatic mutations

accumulating during stem growth are responsible for the observed fitness effects.

The potential for the acquisition of mutations during vegetative growth is a well-known aspect

of plant biology (Klekowski 2003; Schultz and Scofield 2009; Bobiwash et al. 2013), but no previous study

has demonstrated the effects of somatic mutations on the fitness of progeny in the next generation.

Moreover, most studies have focused on the detrimental effects of somatic mutations; chloroplast

mutants have been observed in a number of species (Klekowski 2003), declines in pollen fertility were

found in older clones of quaking aspen (Ally et al. 2010), and higher rates of seed and fruit abortion after

autogamous pollinations were found in several studies (reviewed in Bobiwash et al. 2013). In contrast,

agriculturalists have taken advantage of beneficial somatic mutations to improve economically

important plants (Aradhya et al. 2003; McKey et al. 2010; Miller and Gross 2011; Vezzulli et al. 2012;

Jarni et al. 2015; Pelsy et al. 2015), and a handful of studies report phenotypic responses to selection in

asexual lineages. For example, Breese et al. (1965) succeeded in selecting for increased tillering ability

(production of new grass stems) within genets of perennial ryegrass (*Lolium perenne*). Similarly, artificial

selection on clonal lineages effectively improved branching in the red seaweed, *Asparagopsis armata*

(Monro and Poore 2009). The current study on *M. guttatus* contributes to this literature by highlighting

the potential for plants to exhibit significant levels of clonal variation within a single generation.

The transmission of beneficial mutations in autogamous crosses may explain some heretofore

difficult to understand results from mutation accumulation studies. Our results suggest that beneficial

somatic mutations are likely to be partially dominant (i.e. not completely recessive), because they would

have to be expressed in the heterozygous state to be favored by cell lineage selection. Mutations

accumulating during vegetative growth could contribute to standing genetic variation, but autogamy

may be more effective for the accumulation of beneficial somatic mutations in populations than

geitonogamy or outcrossing, because beneficial mutations could be made homozygous in progeny in a

single generation. Depending on the crossing design, outcrossing would take at least two generations for

beneficial somatic mutations to be made homozygous, and under geitonogamy they would remain

heterozygous. It is striking that high rates of beneficial mutation accumulation have been observed in at

least some mutation accumulation studies in the autogamous plant *Arabidopsis thaliana* (Shaw et al.

2002; Rutter et al. 2010; Rutter et al. 2012), but not in outcrossing and partially-selfing species of

Amsinckia (Schoen 2005). With a few exceptions (e.g., Baer et al. 2005; Denver et al. 2010), nearly all

mutation accumulation studies on animals consistently show a prevalence of deleterious mutations

(Baer et al. 2007; Halligan and Keightley 2009). Similarly, our results suggest that the adaptive potential

of autogamous plants may be greater than previously thought, which may help explain the wider

geographic ranges of selfing compared to closely-related outcrossing species (Grossenbacher et al. 2015;

Grant and Kalisz 2020). Although developmental selection has the potential to contribute to adaptation

in all plants, its effects may be enhanced in autogamous lineages, because beneficial mutations arising

during vegetative growth have a greater chance of becoming homozygous in offspring and being

retained across generations.

As stems grow, mutations can be generated during every mitotic cell division, so the potential

for somatic mutation accumulation in plants appears substantial. Thus, understanding how long-lived

plants, such as trees, avoid mutational meltdown from the accumulation of deleterious somatic

mutations remains a longstanding question. Paradoxically, however, the rate of mutation accumulation

observed across generations in plant and animal genomes is similar (Gaut et al. 2011). One hypothesis

for this pattern is that, similar to animal germlines, a subset of cell lines in the apical meristem

undergoes fewer mitotic divisions, which would protect lineages from the negative effects of mutation

accumulation during development of the soma (Plant Germline Hypothesis; Burian et al. 2016; Cruzan

2018; page 90). An alternative hypothesis posits that somatic mutations are generated in apical

meristems during plant growth, but these mutations are filtered by developmental selection prior to the

establishment of offspring (Somatic Mutation Accumulation Hypothesis). Because plants have retained

the capacity to undergo clonal evolution from their algal ancestors, the ability to filter mutations during

growth and reproduction has existed for some time, and thus developmental selection has the potential

to skew the distribution of fitness effects of transmitted mutations to include a larger proportion of

beneficial mutations than would be expected through random processes. In addition, this provides a

reasonable explanation for why longer-lived plants appear to have slower rates of mutation

accumulation across generations (Yue et al. 2010; Gaut et al. 2011). This could be due to the fact that

longer generation time leads to more time between recombination events, which can lead to more

background selection in non-recombining cell lineages during vegetative growth (Cruzan 2018; pages 94-

95). Recent studies in oaks, spruce, and eel grass (Plomion et al. 2018; Hanlon et al. 2019; Yu et al.

2020), as well as our unpublished data from *M. guttatus*, confirm that multiple somatic variants are

likely, even in plants of very different size. Moreover, results from the current study identify the

presence of beneficial mutations that were passed to offspring in the next generation, likely a

consequence of developmental selection. Future work that combines information from experiments

evaluating the genomic consequences of somatic variation with anatomical estimates of stem cell

population dynamics will allow for the development of new models that provide insights into the extent

and limitations of somatic evolution in plants.

In conclusion, despite the potential for somatic mutation accumulation to generate novel

genetic variation in plant populations, its role in evolution remains almost entirely unexplored. Our

estimates of the fitness effects of somatic mutations were consistent across two different methods and

indicate that some stems possessed a prevalence of deleterious mutations while others produced

autogamous progeny with high fitness, which indicates the presence of beneficial mutations. Even

though high levels of mutation accumulation are often believed to be detrimental, the basic biology of

plants suggests that the role of somatic mutations in plant evolution should be considered carefully in

the future. Moving forward, our results indicate that we must keep in mind that all eukaryotes are not

necessarily equivalent, and specific features of the organism's biology may have unexpected

consequences for the evolutionary phenomena that we observe. Future lines of investigation will

improve our understanding of these fundamental aspects of plant development and evolution that may

have contributed to the remarkable diversification of plants, and may help to account for some of the

extensive variation in mutation rates detected among lineages.

Acknowledgements

We thank J. Thompson, E. Perez, and our research greenhouse manager Linda Taylor for plant

maintenance and assistance with this research in the greenhouse and in the lab. Several people made

helpful suggestions on this manuscript including N. Diaz, M. Grasty, and J. Persinger. This research was

supported by an NIH NIGMS BUILD EXITO grant (5TL4GM118965-03, 5UL1GM118964-03, and

5RL5GM118963-03) to Portland State University.

Data Archiving

Data and all scripts for data analysis will be submitted to DataDryad upon publication.

Literature Cited

Ally, D., K. Ritland, and S. P. Otto. 2010. Aging in a Long-Lived Clonal Tree. PLoS Biol. 8. (http://doi.org/10.1371/journal.pbio.1000454).

Antolin, M. F. and C. Strobeck. 1985. The population genetics of somatic mutation in plants. The American Naturalist 126:52-62. (http://doi.org/10.1086/284395).

Aradhya, M. K., G. S. Dangl, B. H. Prins, J. M. Boursiquot, M. A. Walker, C. P. Meredith, and C. J. Simon. 2003. Genetic structure and differentiation in cultivated grape, *Vitis vinifera* L. Genet. Res. 81:179-192. (http://doi.org/10.1017/s0016672303006177).

Armbruster, W. S. and D. G. Rogers. 2004. Does pollen competition reduce the cost of inbreeding? Am. J. Bot. 91:1939-1943. (http://doi.org/10.3732/ajb.91.11.1939).

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not sperm-specific, genes show stronger purifying selection and higher rates of positive selection than sporophytic genes in *Capsella grandiflora*. Mol. Biol. Evol. 30:2475-2486. (http://doi.org/10.1093/molbev/mst149).

Baer, C. F., M. M. Miyamoto, and D. R. Denver. 2007. Mutation rate variation in multicellular eukaryotes: causes and consequences. Nature Reviews Genetics 8:619-631.

Baer, C. F., F. Shaw, C. Steding, M. Baumgartner, A. Hawkins, A. Houppert, N. Mason, M. Reed, K. Simonelic, W. Woodard, and M. Lynch. 2005. Comparative evolutionary genetics of spontaneous mutations affecting fitness in rhabditid nematodes. Proceedings of the National Academy of Sciences of the United States of America 102:5785-5790. (http://doi.org/10.1073/pnas.0406056102).

Barfield, S., G. V. Aglyamova, and M. V. Matz. 2016. Evolutionary origins of germline segregation in Metazoa: evidence for a germ stem cell lineage in the coral *Orbicella faveolata* (Cnidaria, Anthozoa). Proceedings of the Royal Society B: Biological Sciences 283:20152128. (http://doi.org/doi:10.1098/rspb.2015.2128).

Baulcombe, D. C. and C. Dean. 2014. Epigenetic regulation in plant responses to the environment. Cold Spring Harbor Perspectives in Biology 6:a019471. (http://doi.org/10.1101/cshperspect.a019471).

Bobiwash, K., S. T. Schultz, and D. J. Schoen. 2013. Somatic deleterious mutation rate in a woody plant: estimation from phenotypic data. Heredity 111:338-344. (http://doi.org/10.1038/hdy.2013.57).

Breese, E. L., M. D. Hayward, and A. C. Thomas. 1965. Somatic selection in perennial ryegrass. Heredity 20:367-&. (http://doi.org/10.1038/hdy.1965.50).

Buchholz, J. T. 1922. Developmental selection in vascular plants. Botanical Gazette 73:249-286.

Burian, A., P. B. de Reuille, and C. Kuhlemeier. 2016. Patterns of stem cell divisions contribute to plant longevity. Curr. Biol. 26:1385-1394. (http://doi.org/10.1016/j.cub.2016.03.067).

Buss, L. W. 1983. Evolution, development, and the units of selection. Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences 80:1387-1391. (http://doi.org/10.1073/pnas.80.5.1387).

Charlesworth, D. and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev. Genet. 10. (http://doi.org/10.1038/nrg2664).

Crow, J. F. 1993. Mutation, mean fitness, and genetic load. Pp. 3-42. Oxford Surveys in Evolutionary Biology. Oxford University Press.

Cruzan, M. and J. Thomson. 1997. Effects of pre-dispersal selection on offspring growth and survival in *Erythronium grandiflorum*. J. Evol. Biol. 10:295-314.

Cruzan, M. B. 1989. Pollen tube attrition in *Erythronium grandiflorum*. Amer. J. Bot. 76:562-570.

Cruzan, M. B. 2018. Evolutionary Biology - A Plant Perspective. Oxford University Press, New York.

Denver, D. R., D. K. Howe, L. J. Wilhelm, C. A. Palmer, J. L. Anderson, K. C. Stein, P. C. Phillips, and S. Estes. 2010. Selective sweeps and parallel mutation in the adaptive recovery from deleterious mutation in Caenorhabditis elegans. Genome Res. 20:1663-1671. (http://doi.org/10.1101/gr.108191.110).

Dubrovina, A. S. and K. V. Kiselev. 2016. Age-associated alterations in the somatic mutation and DNA methylation levels in plants. Plant Biol. 18:185-196. (http://doi.org/10.1111/plb.12375).

Elena, S. F. and R. E. Lenski. 2003. Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. Nat. Rev. Genet. 4:457-469.

Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.

Fogle, C. A., J. L. Nagle, and M. M. Desai. 2008. Clonal interference, multiple mutations and adaptation in large asexual populations. Genetics 180:2163-2173. (http://doi.org/10.1534/genetics.108.090019).

Gaut, B., L. Yang, S. Takuno, and L. E. Eguiarte. 2011. The patterns and causes of variation in plant nucleotide substitution rates. Pp. 245-266 in D. J. Futuyma, H. B. Shaffer, and D. Simberloff, eds. Annual Review of Ecology, Evolution, and Systematics, Vol 42.

Grant, A.-G. and S. Kalisz. 2020. Do selfing species have greater niche breadth? Support from ecological niche modeling. Evolution 74:73-88. (http://doi.org/10.1111/evo.13870).

Greaves, M. and C. C. Maley. 2012. Clonal evolution in cancer. Nature 481:306-313. (http://doi.org/http://www.nature.com/nature/journal/v481/n7381/abs/nature10762.html#supplementary-information).

Groot, Edwin P. and T. Laux. 2016. Ageing: how do long-lived plants escape mutational meltdown? Curr. Biol. 26:R530-R532. (http://doi.org/https://doi.org/10.1016/j.cub.2016.05.049).

Grossenbacher, D., R. Briscoe Runquist, E. E. Goldberg, and Y. Brandvain. 2015. Geographic range size is predicted by plant mating system. Ecol. Lett. 18:706-713. (http://doi.org/10.1111/ele.12449).

Halligan, D. L. and P. D. Keightley. 2009. Spontaneous Mutation Accumulation Studies in Evolutionary Genetics. Pp. 151-172. Annual Review of Ecology Evolution and Systematics.

Hanlon, V. C. T., S. P. Otto, and S. N. Aitken. 2019. Somatic mutations substantially increase the per-generation mutation rate in the conifer Picea sitchensis. Evolution Letters 3:348-358. (http://doi.org/doi:10.1002/evl3.121).

Harder, L. D., M. A. Aizen, and S. A. Richards. 2016. The population ecology of male gametophytes: the link between pollination and seed production. Ecol. Lett. 19:497-509. (http://doi.org/10.1111/ele.12596).

Hill, W. G. and A. Robertson. 1966. The effect of linkage on limits to artificial selection. Genetics Research 8:269-294.

Husband, B. and D. W. Schemske. 1995. Evolution of the magnitude and timing of inbreeding depression in plants. Evolution 50:54-70.

Itabashi, E., K. Osabe, R. Fujimoto, and T. Kakizaki. 2018. Epigenetic regulation of agronomical traits in Brassicaceae. Plant Cell Rep. 37:87-101. (http://doi.org/10.1007/s00299-017-2223-z).

Jarni, K., J. Jakse, and R. Brus. 2015. Vegetative propagation: linear barriers and somatic mutation affect the genetic structure of a Prunus avium L. stand. Forestry 88:612-621. (http://doi.org/10.1093/forestry/cpv029).

Klekowski, E. J. 2003. Plant clonality, mutation, diplontic selection and mutational meltdown. Biol. J. Linn. Soc. 79:61-67. (http://doi.org/10.1046/j.1095-8312.2003.00183.x).

Klekowski, E. J. and P. J. Godfrey. 1989. Ageing and mutation in plants. Nature 340:389-391. (http://doi.org/10.1038/340389a0).

Korbecka, G., P. G. L. Klinkhamer, and K. Vrieling. 2002. Selective embryo abortion hypothesis revisited - A molecular approach. Plant Biol. 4:298-310. (http://doi.org/10.1055/s-2002-32331).

Kuhlemeier, C. 2017. How to get old without aging. Nature Plants 3:916-917.
(http://doi.org/10.1038/s41477-017-0076-7).
Lang, G. I., D. P. Rice, M. J. Hickman, E. Sodergren, G. M. Weinstock, D. Botstein, and M. M. Desai. 2013.
Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. Nature
500:571-574. (http://doi.org/10.1038/nature12344).
Lee, M.-C. and C. J. Marx. 2013. Synchronous waves of failed soft sweeps in the laboratory: remarkably
rampant clonal Interference of alleles at a single locus. Genetics 193:943-+.
(http://doi.org/10.1534/genetics.112.148502).
Long, A., G. Liti, A. Luptak, and O. Tenaillon. 2015. Elucidating the molecular architecture of adaptation
via evolve and resequence experiments. Nature Reviews Genetics 16:567-582.
(http://doi.org/10.1038/nrg3937).
Lynch, M., J. Blanchard, D. Houle, T. Kibota, S. Schultz, L. Vassilieva, and J. Willis. 1999. Perspective:
Spontaneous deleterious mutation. Evolution 53:645-663. (http://doi.org/10.2307/2640707).
Mable, B. K. and S. P. Otto. 1998. The evolution of life cycles with haploid and diploid phases. Bioessays
20:453-462. (http://doi.org/10.1002/(sici)1521-1878(199806)20:6<453::aid-bies3>3.0.co;2-n).
Maharjan, R. P., B. Liu, L. Feng, T. Ferenci, and L. Wang. 2015. Simple phenotypic sweeps hide complex
genetic changes in populations. Genome Biology and Evolution 7:531-544.
(http://doi.org/10.1093/gbe/evv004).
McKey, D., M. Elias, B. Pujol, and A. Duputie. 2010. The evolutionary ecology of clonally propagated
domesticated plants. New Phytol. 186:318-332. (http://doi.org/10.1111/j.1469-
8137.2010.03210.x).
McKnight, T. D., K. Riha, and D. E. Shippen. 2002. Telomeres, telomerase, and stability of the plant
genome. Plant Mol. Biol. 48:331-337. (http://doi.org/10.1023/a:1014091032750).
McManus, M. T. and B. E. Veit, eds. 2002. Meristematic Tissues in Plant Growth and Development. CRC
Press LLC, Boca Raton, FL.
Mena-Alí, J. I. and O. J. Rocha. 2005. Selective seed abortion affects the performance of the offspring in
Bauhinia ungulata. Ann. Bot. 95:1017-1023. (http://doi.org/10.1093/aob/mci109).
Miller, A. J. and B. L. Gross. 2011. From forest to field: perennial fruit crop domestication. Am. J. Bot.
98:1389-1414. (http://doi.org/10.3732/ajb.1000522).
Monro, K. and A. G. B. Poore. 2009. The potential for evolutionary responses to cell-lineage selection on
growth form and its plasticity in a red seaweed. Am. Nat. 173:151-163.
(http://doi.org/10.1086/595758).
Mulcahy, D. L. 1979. The rise of the angiosperms: A genecological factor. Science 206:20-23.
Nowell, P. C. 1976. Clonal evolution of tumor-cell populations. Science 194:23-28.
(http://doi.org/10.1126/science.959840).
Orive, M. E. 2001. Somatic mutations in organisms with complex life histories. Theor. Popul. Biol.
59:235-249. (http://doi.org/10.1006/tpbi.2001.1515).
Otto, S. P. and I. M. Hastings. 1998. Mutation and selection within the individual. Genetica 102-103:507-
524. (http://doi.org/10.1023/a:1017074823337).
Otto, S. P. and M. E. Orive. 1995. Evolutionary consequences of mutation and selection within an
individual. Genetics 141:1173-1187.
Pelsy, F., V. Dumas, L. Bevilacqua, S. Hocquigny, and D. Merdinoglu. 2015. Chromosome Replacement
and Deletion Lead to Clonal Polymorphism of Berry Color in Grapevine. PLoS Genet. 11.
(http://doi.org/10.1371/journal.pgen.1005081).
Plomion, C., J.-M. Aury, J. Amselem, T. Leroy, F. Murat, S. Duplessis, S. Faye, N. Francillonne, K. Labadie,
G. Le Provost, I. Lesur, J. Bartholomé, P. Faivre-Rampant, A. Kohler, J.-C. Leplé, N. Chantret, J.
Chen, A. Diévart, T. Alaeitabar, V. Barbe, C. Belser, H. Bergès, C. Bodénès, M.-B. Bogeat-
Triboulot, M.-L. Bouffaud, B. Brachi, E. Chancerel, D. Cohen, A. Couloux, C. Da Silva, C. Dossat, F.

Ehrenmann, C. Gaspin, J. Grima-Pettenati, E. Guichoux, A. Hecker, S. Herrmann, P. Hugueney, I. Hummel, C. Klopp, C. Lalanne, M. Lascoux, E. Lasserre, A. Lemainque, M.-L. Desprez-Loustau, I. Luyten, M.-A. Madoui, S. Mangenot, C. Marchal, F. Maumus, J. Mercier, C. Michotey, O. Panaud, N. Picault, N. Rouhier, O. Rué, C. Rustenholz, F. Salin, M. Soler, M. Tarkka, A. Velt, A. E. Zanne, F. Martin, P. Wincker, H. Quesneville, A. Kremer, and J. Salse. 2018. Oak genome reveals facets of long lifespan. Nature Plants 4:440-452. (http://doi.org/10.1038/s41477-018-0172-3).

Quadrana, L. and V. Colot. 2016. Plant Transgenerational Epigenetics. Pp. 467-491 *in* N. M. Bonini, ed. Annual Review of Genetics, Vol 50.

Reusch, T. B. H. and C. Bostrom. 2011. Widespread genetic mosaicism in the marine angiosperm Zostera marina is correlated with clonal reproduction. Evol. Ecol. 25:899-913. (http://doi.org/10.1007/s10682-010-9436-8).

Rutter, M. T., A. Roles, J. K. Conner, R. G. Shaw, F. H. Shaw, K. Schneeberger, S. Ossowski, D. Weigel, and C. B. Fenster. 2012. Fitness of *Arabidopsis thaliana* mutation accumulation lines whose spontaneous mutations are known. Evolution 66:2335-2339. (http://doi.org/10.1111/j.1558-5646.2012.01583.x).

Rutter, M. T., F. H. Shaw, and C. B. Fenster. 2010. Spontaneous mutation parameters for *Arabidopsis thaliana* measured in the wild. Evolution 64:1825-1835. (http://doi.org/10.1111/j.1558-5646.2009.00928.x).

SAS. 2008. SAS/STAT 9.2 user's guide. SAS Institute Inc., Cary, North Carolina, USA.

Schmid-Siegert, E., N. Sarkar, C. Iseli, S. Calderon, C. Gouhier-Darimont, J. Chrast, P. Cattaneo, F. Schütz, L. Farinelli, M. Pagni, M. Schneider, J. Voumard, M. Jaboyedoff, C. Fankhauser, C. S. Hardtke, L. Keller, J. R. Pannell, A. Reymond, M. Robinson-Rechavi, I. Xenarios, and P. Reymond. 2017. Low number of fixed somatic mutations in a long-lived oak tree. Nature Plants 3:926-929. (http://doi.org/10.1038/s41477-017-0066-9).

Schoen, D. J. 2005. Deleterious mutation in related species of the plant genus *Amsinckia* with contrasting mating systems. Evolution 59:2370-2377.

Schultz, S. T. and D. G. Scofield. 2009. Mutation accumulation in real branches: fitness assays for genomic deleterious mutation rate and effect in large-statured plants. Am. Nat. 174:163-175. (http://doi.org/10.1086/600100).

Schweinsberg, M., R. A. González Pech, R. Tollrian, and K. P. Lampert. 2014. Transfer of intracolonial genetic variability through gametes in Acropora hyacinthus corals. Coral Reefs 33:77-87. (http://doi.org/10.1007/s00338-013-1102-5).

Shaw, F. H., C. J. Geyer, and R. G. Shaw. 2002. A comprehensive model of mutations affecting fitness and inferences for *Arabidopsis thaliana*. Evolution 56:453-463.

Stearns, F. W. and C. B. Fenster. 2016. Fisher's geometric model predicts the effects of random mutations when tested in the wild. Evolution 70:495-501. (http://doi.org/10.1111/evo.12858).

Vezzulli, S., L. Leonardelli, U. Malossini, M. Stefanini, R. Velasco, and C. Moser. 2012. Pinot blanc and Pinot gris arose as independent somatic mutations of Pinot noir. J. Exp. Bot. 63:6359-6369. (http://doi.org/10.1093/jxb/ers290).

Watson, J. M., A. Platzer, A. Kazda, S. Akimcheva, S. Valuchova, V. Nizhynska, M. Nordborg, and K. Riha. 2016. Germline replications and somatic mutation accumulation are independent of vegetative life span in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America 113:12226-12231. (http://doi.org/10.1073/pnas.1609686113).

Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008. Mimulus is an emerging model system for the integration of ecological and genomic studies. Heredity 100:220-230. (http://doi.org/10.1038/sj.hdy.6801018).

Younginger, B. S., D. Sirová, M. B. Cruzan, and D. J. Ballhorn. 2017. Is biomass a reliable estimate of plant fitness? Applications in Plant Sciences 5:apps.1600094. (http://doi.org/10.3732/apps.1600094).

Yu, L., C. Boström, S. Franzenburg, T. Bayer, T. Dagan, and T. B. H. Reusch. 2020. Somatic genetic drift
and multilevel selection in a clonal seagrass. Nature Ecology & Evolution.
(http://doi.org/10.1038/s41559-020-1196-4).
Yue, J.-X., J. Li, D. Wang, H. Araki, D. Tian, and S. Yang. 2010. Genome-wide investigation reveals high
evolutionary rates in annual model plants. BMC Plant Biol. 10:1-12.
(http://doi.org/10.1186/1471-2229-10-242).

Figure Captions

Fig. 1. Experimental design to test for the average fitness effects of somatic mutations accumulating in
stems during vegetative growth. See text for further explanation.

Fig. 2. An illustration of cell lineage selection as it acts on a number of cell lines that differ in growth rate
within the apical meristem of a plant. A) Expressed deleterious mutations are predicted to lead to the
demise of a cell line (brown cells) that is replaced by neighboring cell lines. The appearance of a
mutation that increases cell growth rate in the current environment results in a selective sweep of the
meristem (light green cells). B) Multiple ramets of the same genet become differentiated by unique
somatic mutations during vegetative growth. Most ramets carry deleterious mutations (orange, red,
purple, and pink arrows), but expressed beneficial mutations cause clonal selective sweeps of the
meristem in some ramets (green arrows).

Fig. 3. Estimates of $\delta_{AD(k)}$ for fourteen different stems (ramets) from two separate experiments
(Experiment 1–blue bars; Experiment 2–orange bars) based on mean progeny fitness after autogamous
$\overline{w}_{k(A)}$ and geitonogamous $\overline{w}_{k(G)}$ self-pollinations ($\delta_{AD(k)}$ = $\overline{w}_{k(A)}$ - $\overline{w}_{k(G)}$). Horizontal lines represent
standard errors. Asterisks indicate values of $\delta_{AD(k)}$ that are significantly different from zero based on the t
value, calculated as $t = \delta_{AD(k)}/\sqrt{SE}$ with n-1 df, where n is the mean of sample sizes for progeny from
autogamy and geitonogamy. The relationship between $\overline{w}_{k(A)}$ and $\overline{w}_{k(G)}$ across stems is shown in Fig. S2.
Means and sample sizes for progeny groups are available in Table S6 and Fig. S3.

Fig. 4. Relationships between estimates of fitness effects of somatic mutations, based either on the
difference in fitness of progeny from autogamy and geitonogamy ($\delta_{AD(k)}$), or the standard deviation in
fitness within progeny groups from autogamy for each stem (w_{SD}). Estimates of w_{SD} corresponding to
negative values of $\delta_{AD(k)}$ were transformed to negative values. Estimates from Experiment 1 are indicated
by blue circles and from Experiment 2 are orange squares. Dashed lines indicate the separate
relationships for positive and negative values of fitness estimates.

Fig. 1

Fig. 2

Cell Generations

A. Cell Lineage Selection

B. Differentiation among Ramets

Fig. 3

Average Fitness Effects ($\delta_{AD(k)}$)

Fig. 4

823
824

Appendix 1. Validity of the $\delta_{AD(k)}$ Estimate

For a pair of stems from the same genet, we assume that w_G represents the fitness of geitonogamous
progeny after a cross between the pair of stems, and that w_A represents the fitness of autogamous
progeny after selfing a single flower on one stem:

$w_G = 1 + 0.5(h_1s_1+h_2s_2)$

$w_A = 0.25(1) + 0.5(1+ h_1s_1) + 0.25(1+s_1)$

Where, s_1 and s_2 represent the selective effects of mutations, and h_1 and h_2 represent the dominance of
single somatic mutations in stem 1 and stem 2, respectively.

For simulations, we randomly chose values of s ranging from -0.2 to 0.2 for 200 pairs of stems. Values of
h for $s > 0$ were allowed to range from 0 to 0.7 for each stem. Because deleterious alleles tend to be
recessive, values of h were constrained to range between 0 and 0.1 for $s < 0$. Dominance values were
chosen assuming that deleterious somatic mutations with high levels of expression would be eliminated
by cell lineage selection during vegetative growth, and beneficial mutations with high dominance would
not display strong fitness effect differences between autogamous and geitonogamous progeny. To
match our experimental design in *Mimulus guttatus*, we calculated the fitness of autogamous progeny
(w_A) for one of the stems in each pair. We then calculated w_G using information on s and h for both
stems in each pair. Then we calculated the difference in fitness between each estimate as a measure of
autogamy depression ($\delta_{AD(k)} = w_A - w_G$). This simulation was repeated 20 times (4,000 pairs of stems
total) and average values were calculated.

Results:

1.　Estimates of $\delta_{AD(k)}$ and s_1 are strongly correlated, with R^2 values ranging from 0.76 to 0.82 (one
example of 200 random estimates):

2. The frequency of $\delta_{AD(k)}$ estimates having the opposite sign of s_1 was very low. Mean number of
correct signs was 178.7 (+/- 1.04 standard error) and was 21.30 (+/-1.04; or 12%) for incorrect
signs.

3. Estimates of $\delta_{AD(k)}$ with incorrect signs were closer to zero. Mean absolute values of s for $\delta_{AD(k)}$
estimates with incorrect signs was 0.054 (+/- 0.0021) and was 0.106 (+/-0.0011) for correct
signs. Hence, estimates of $\delta_{AD(k)}$ that are further from zero are more reliable estimates of s_1.

We note that the formulation used above for w_G could result in estimates that are out of range when
large numbers of mutations are involved (i.e., $w_G = 1 + 0.5 \sum h_i\, s_i$ for a large number of loci segregating
for deleterious alleles). A better estimate may be the average effect of n loci segregating for deleterious
alleles (i.e. $\sum h_i\, s_i\, /n$). If we apply this approach for two loci (one mutation per stem) we obtain,

$w_G = 0.25 + 0.25(1 + h_1s_1) + 0.25(1 + h_2s_2) + 0.25(1 + (h_1s_1+h_2s_2)/2)$

$w_G = 1 + 0.375(h_1s_1+h_2s_2)$

Using this formula, the relationship between $\delta_{AD(k)}$ and s_1 is stronger (R^2 ranging from 0.86 to 0.90), with
only an average of 13 out of 200 estimates with opposite signs (7%). The average of correct signs was
similar to the one above (0.105, +/- 0.0007), but the mean for incorrectly assigned values was much
closer to zero (0.038, +/- 0.0022).

Another model for the effects of alleles at multiple loci on fitness assumes the interaction is
multiplicative. In this case the estimate of w_G becomes,

$w_G = 0.25 + 0.25(1 + h_1s_1) + 0.25(1 + h_2s_2) + 0.25(1 + (h_1s_1h_2s_2))$

$w_G = 1 + 0.25(h_1s_1+h_2s_2) + 0.25(h_1s_1h_2s_2)$

Using this formula, the relationship between $\delta_{AD(k)}$ and s_1 is stronger (R^2 ranging from 0.93 to 0.95), with
only an average of 7.1 out of 200 estimates with opposite signs (4%). The average of correct signs was
similar to the ones above (0.103, +/- 0.0010), but the mean for incorrectly assigned values was closer to
zero (0.023, +/- 0.0020).

Interpretation:

These results indicate that our estimate of $\delta_{AD(k)}$ is a valid estimate of s for stem 1. However, it does
appear that we are underestimating s (i.e., the slope of the line is less than 1), as these estimates do not
consider the effects of multiple somatic mutations or possible genetic background effects (i.e. epistasis).

It is also notable that our simulation assumes a wider range of h values than are generally observed for
deleterious mutations. Estimates indicate that the product of hs for deleterious alleles is generally
around 0.02, and that the relationship between s and h is generally negative (hyperbolic; Lynch et al.
1999). Minor effects of deleterious mutations are expected for our data, because somatic mutations
having strong effects in the heterozygous condition would be eliminated by cell lineage selection.
Adding the constraint of a negative relationship between h and s to our simulation model would
marginally improve the predictive power of $\delta_{AD(k)}$ for estimates of s.

These simulations also support our use of the correlation between w_A and w_G to confirm that our
estimates are similar to the simulation estimates.

From the simulations:

From our data:

Appendix 2. Estimating the selection coefficient due to somatic mutations from the variance in fitness
among progeny.
897
898 In this Appendix, we demonstrate that the standard deviation (SD) in fitness among progeny from
899 autogamous crosses can be used to estimate the selection coefficient for the average effects of somatic
900 mutations occurring within a stem. The variance in fitness among progeny from autogamy can be
901 affected by the selection coefficient (s) and the dominance effects (h) of somatic mutations unique to a
902 stem.
903
904 To evaluate the effects of a single mutation on progeny fitness, we assume fitnesses of;
905   $w_{aa} = 1$
906   $w_{aa'} = 1 + hs$
907   $w_{a'a'} = 1 + s$
908
909 The mean progeny fitness is then calculated as
910
911 $\overline{w} = ¼(1) + ½(1+hs) + ¼(1+s)$
912
913 Which simplifies to
914
915 $\overline{w} = (1 + ½\ hs + ¼\ s)$
916
917 The variance in progeny fitness can be calculated as
918
919 $V(w) = ¼(w_{aa} - \overline{w})^2 + ½(w_{aa'} - \overline{w})^2 + ¼(w_{a'a'} - \overline{w})^2$
920
921 After substituting, we get
922
923 $V(w) = [¼\ (-½\ hs - ¼\ s)^2] + [½\ (½\ hs - ¼\ s)^2] + [¼\ (¾\ s - ½\ hs\ )^2]$
924
925 For simplification, we separate the effects of homozygous and heterozygous genotypes (terms within
926 the three sets of square brackets)
927
928 $V(w) = [1] + [2] + [3]$
929
930 and evaluate each one;
931
932 $[1]: ¼\ (-½\ hs - ¼\ s)^2 = ¼\ (-½\ hs\ (-½\ hs - ¼\ s)- ¼\ s(-½\ hs - ¼\ s)) = (\ 1/16\ h^2s^2 + 1/16\ hs^2 + 1/64\ s^2\ )$
933
934 $[2]: ½\ (½\ hs - ¼\ s)^2 = ½\ (½\ hs\ (½\ hs - ¼\ s) - ¼\ s(½\ hs - ¼\ s)) = ½\ (1/4\ h^2s^2 - 1/8\ hs^2 - 1/8\ hs^2 + 1/16\ s^2\ ) = ($
935 $1/8\ h^2s^2 - 1/8\ hs^2 + 1/32\ s^2\ )$
936
937 $[3]: ¼\ (¾\ s - ½\ hs\ )^2 = ¼\ (¾\ s\ (¾\ s - ½\ hs\ ) - ½\ hs\ (¾\ s - ½\ hs\ )) = ¼\ (9/16\ s^2 - 3/8\ hs^2 - 3/8\ hs^2 + ¼\ h^2s^2\ ) =$
938 $(9/64\ s^2 - 3/16\ hs^2 + 1/16\ h^2s^2\ )$
939
By combining and then simplifying these three results, we get

$V(w) = 3/16\ s^2 - \tfrac{1}{4}\ hs^2 + \tfrac{1}{4}\ h^2 s^2$

And the standard deviation can be obtained by taking

the square root of V(w)

$SD(w) = \sqrt{V(w)} = 0.433\ s - \tfrac{1}{2}\ s\sqrt{h} + \tfrac{1}{2}\ hs$

We note that this relationship holds regardless of the

sign (positive or negative values of s), so for this

relationship we can substitute w for s. The standard

deviation in progeny fitness scales linearly with the

selective effects of somatic mutations, and the effects

of dominance on variation among progeny is minor

(Fig. S1; which shows values of s or w predicted from

Fig. S1. The effects of selection (s) or fitness (w) and dominance (h) on the standard deviation (SD) in fitness among progeny. The effect of dominance on variation in fitness is minimized when h = 0.5 and maximized when h = 0.0 and h = 1.0.

values of SD for fitness ranging from 0 to 0.5). We note that the standard deviation is highest when

mutations are completely recessive or dominant (h = 0 or 1) and SD(w) = 0.433s. When h = 0.5, the

standard deviation is at its minimal value and SD(w) = 0.332s. By rearranging, we can obtain the

regression equations to estimate the selective effects of somatic mutations from the standard deviation

in progeny fitness as $s_{SD} = cSD$, where c is the slope of the regression line in Fig. S1. From this analysis, it

is clear that the effects of h on the within-family SD in fitness is minor compared to the effects of s. We

obtain estimates of the regression slopes from the standard deviation in progeny fitness to produce the

relationships in Fig. S1;

$w = 2.31*SD(w)$ for h = 0 or 1, and

$w = 3.01*SD(w)$ for h = 0.5

From these relationships we can estimate the selective effects of somatic mutations from the standard

deviation in progeny fitness as $w_{SD} = cSD$, where c is the slope of the regression lines in Fig. S1.

Appendix 3. Supplemental Tables and Figures

975 Table S1.

	Autogamy			Geitonogamy				
Stem	$\overline{w}_A$	**N**	**Stdev**	$\overline{w}_G$	**N**	**Stdev**	$\delta_{AD(k)}$	$\overline{w}_A - \overline{\overline{w}}_G$
JB51	0.407	33	0.171	0.144	7	0.080	0.263	0.078
JB52	0.314	12	0.161	0.295	9	0.068	0.019	-0.015
JB54	0.258	8	0.066	0.298	11	0.103	-0.040	-0.071
SMB11	0.286	22	0.072	0.260	9	0.077	0.009	-0.043
SMB51	0.256	11	0.054	0.323	4	0.035	-0.067	-0.073
SMB52	0.273	7	0.072	0.280	6	0.051	-0.007	-0.056
SMB53	0.245	4	0.021	0.327	6	0.108	-0.081	-0.083
SMB54	0.289	10	0.023	0.245	5	0.056	0.044	-0.040
SMC11	0.303	16	0.071	0.238	8	0.085	0.065	-0.026
SMC12	0.356	8	0.071	0.239	6	0.059	0.117	0.027
C12	0.423	36	0.170	0.707	5	0.153	-0.283	0.094
C13	0.626	9	0.195	0.322	7	0.237	0.090	0.297
D17	0.692	5	0.170	0.485	4	0.314	0.207	0.363
D18	0.419	12	0.130	0.443	13	0.211	-0.025	0.090

980 Table S2. Analysis of variance results for seed set after autogamous and geitonogamous pollinations
981 (Cross) with limited and excess pollen (Pollen). Population (Pop), maternal stem nested within
982 population (Mat(Pop)), and pollination date (Date) were declared as random effects in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	21623.26022	655.25031	1.43	0.0888
Cross	1	100.647606	100.647606	0.22	0.6400
Pop	2	552.228601	276.114301	0.60	0.5487
Pollen	1	2183.436724	2183.436724	4.78	0.0312
Mat(Pop)	18	6490.364191	360.575788	0.79	0.7089
Date	11	8000.908005	727.355273	1.59	0.1127
Error	101	46182.07311	457.24825		

985 Table S3. Analysis of variance results for aborted ovules after autogamous and geitonogamous
986 pollinations (Cross) with limited and excess pollen (Pollen). Population (Pop), maternal stem nested
987 within population (Mat(Pop)), and pollination date (Date) were declared as random effects in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	9337.61468	282.95802	2.76	<.0001
Cross	1	449.176948	449.176948	4.38	0.0389
Pop	2	2174.689160	1087.344580	10.60	<.0001
Pollen	1	0.597410	0.597410	0.01	0.9393
Mat(Pop)	18	2217.350757	123.186153	1.20	0.2751
Date	11	1995.681210	181.425565	1.77	0.0695
Error	101	10363.24458	102.60638		

988
989 Table S4. Analysis of variance for progeny fitness (biomass weighted by survival of seedlings) after
990 autogamous and geitonogamous pollinations (Cross) for ten maternal stems (Mat) and their interaction
991 (Mat*Cross). Planting block (Tray) was declared as a random effect in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	1.42418594	0.04315715	4.18	<.0001
Tray	14	0.45040769	0.03217198	3.11	0.0002
Mat	9	0.06845581	0.00760620	0.74	0.6756
Cross	1	0.03196532	0.03196532	3.09	0.0805
Mat*Cross	9	0.51373985	0.05708221	5.52	<.0001
Error	168	1.73638043	0.01033560		

992
993
994 Table S5. Analysis of variance for progeny fitness (biomass weighted by survival of seedlings) after
995 autogamous and geitonogamous pollinations (Cross) for seven paired pollinations at individual nodes
996 (Stem/Node) and their interaction (Stem/Node*Cross). Hydroponic planting block (Tray) was declared as
997 a random effect in the model and initial size of seedlings (Initial Size) was entered as a covariate.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	23	1.31384069	0.05712351	6.05	<.0001
Tray	11	0.46856600	0.04259691	4.51	<.0001
Initial Size	1	0.46861414	0.46861414	49.61	<.0001
Stem/Node	6	0.12071536	0.02011923	2.13	0.0514
Cross	1	0.00555294	0.00555294	0.59	0.4441
Stem/Node*Cross	4	0.13005812	0.03251453	3.44	0.0095
Error	203	1.91760656	0.00944634		

 Table S6. Means and standard deviations (SD) for progeny groups from autogamous and geitonogamous
 pollinations in experiments 1 and 2. Experiment 1 included ten different genets. Experiment 2 was
 conducted with two ramets (C1 and D1) from the same genet. The fitness of progeny was estimated as
 its final biomass (log transformed to improve normality), weighted by the survival frequency of progeny
 from the same cross.

Exp	ID	N	Fitness Autogamy	SD	N	Fitness Geitonogamy	SD
2	C13	9	0.626	0.195	7	0.322	0.237
1	JB51	33	0.316	0.193	7	0.114	0.041
2	D17	7	0.662	0.170	60	0.465	0.314
1	SMC12	8	0.222	0.080	6	0.103	0.046
1	SMC11	16	0.175	0.065	8	0.099	0.071
1	JB52	12	0.195	0.158	9	0.162	0.071
1	SMB11	22	0.155	0.082	9	0.152	0.096
1	SMB54	10	0.128	0.020	5	0.129	0.072
2	D18	12	0.419	0.130	13	0.443	0.211
1	JB54	8	0.118	0.063	11	0.17	0.109
1	SMB52	7	0.117	0.069	6	0.173	0.057
1	SMB51	11	0.11	0.040	4	0.171	0.048
1	SMB53	4	0.069	0.021	6	0.21	0.108
2	C12	36	0.423	0.188	5	0.706	0.153
		13.93			11.14		

1006
1007
1008

1009
1010
1011   Fig. S2. The relationship between the autogamous and geitonogamous progeny from the 14
1012   comparisons from experiments 1 (blue circles) and 2 (orange squares) as listed in Table S6. The solid line
1013   represents a 1:1 relationship and the dashed line represents the best-fit to the data. Stems falling above
1014   the solid line apparently had a prevalence of deleterious somatic mutations whereas those falling below
1015   the line displayed a prevalence of beneficial somatic mutations.
1016
1017
1018

1019

1020

1021

1022

1023   Fig S3. Means and standard errors (lines) for fitness estimates of autogamous (upper panel) and
1024   geitonogamous (lower panel) progeny groups from experiment 1 (blue bars) and 2 (orange bars) listed in
1025   Table S6. Progeny groups are ordered as in Fig. 4.

1026

1027

1028