1

Phenotypic Effects of Somatic Mutations Accumulating during Vegetative Growth

Mitchell B. Cruzan[1, 2], Matthew A. Streisfeld[3], and Jaime A. Schwoch[1]

[1]Department of Biology, Portland State University, Portland, OR 97207

[3]Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403

[2]Author for correspondence: cruzan@pdx.edu

Submitted: _______________; Electronically published: __________________

Online enhancements: Three appendices. *Dryad Data:* __________________

Abstract

The unique life form of plants promotes the accumulation of large numbers of somatic mutations that can be passed to offspring in the next generation. However, rates of mutation accumulation in plants remain similar to rates in animals. The explanation for this paradox may be that plants have the potential to filter somatic mutations by eliminating deleterious mutations prior to seed dispersal. We present experimental tests of this hypothesis using crosses in plants of *Mimulus guttatus*. The fitness effects of somatic mutations are expected to differ between progeny from self-pollinations made within the same flower (autogamy) relative to those made between stems on the same plant (geitonogamy), as autogamy leads to homozygosity of a proportion of somatic mutations, but progeny from geitonogamy remain heterozygous. Using two different approaches, we find that somatic mutations have consistent fitness effects on offspring. Surprisingly, several progeny groups from autogamous crosses displayed increases in fitness compared to progeny from geitonogamy controls, indicating that beneficial somatic mutations were prevalent in some stems. These results support the hypothesis that somatic mutations accumulate during vegetative growth, but they are filtered by natural selection, resulting in the culling of expressed deleterious mutations and the retention of beneficial mutations.

Key words: Acquired mutations, autogamy, autogamy depression, cell lineage selection, dominance, *Erythranthe guttata*, geitonogamy, *Mimulus guttatus*, somatic mutations.

Introduction

Mutation is the source of variation for evolution and adaptation, but organisms differ in whether

mutations originating during gamete formation (meiosis) or somatic growth (mitosis) contribute to

heritable variation. For the vast majority of organisms, including viruses, unicellular microbes, and some

multicellular eukaryotes, sexual reproduction is rare or absent. In these organisms, mutations occur

during cell replication, and their primary process for adaptation and diversification is by clonal evolution.

By contrast, genetic modification occurring during somatic growth of animals is usually limited to the

appearance of acquired mutations and the development of cancer (Nowell 1976; Beerenwinkel et al.

2015), neither of which contributes to heritable variation in populations. This is due to the fact that in

metazoans, the germline is determined early in development and relatively few cell divisions occur

before the formation of gametes (the Weismann Barrier; Buss 1983). This may be true even for some

highly clonal animals, such as cnidarians, but it remains unclear whether clonal variation in corals is

transferred to the next generation remains (e.g., Schweinsberg et al. 2014), but not others (e.g., Barfield

et al. 2016).

Plants differ from animals and microbes, because mutations contributing to heritable variation

can arise both during gamete formation and somatic growth (Antolin and Strobeck 1985; Klekowski and

Godfrey 1989). This is due to the fact that plants lack separate germ lines; the same cells within apical

meristems subtend both vegetative growth and gamete development, leading to sexual reproduction.

Consequently, plants have the capacity for clonal evolution, as somatic mutations that occur in plant

meristems can accumulate and be passed to the next generation (McKnight et al. 2002; Klekowski 2003;

Schultz and Scofield 2009; Bobiwash et al. 2013; Dubrovina and Kiselev 2016; Watson et al. 2016;

Schmid-Siegert et al. 2017). This aspect of plant biology is well known (Monro and Poore 2009; Ally et al.

2010; Reusch and Bostrom 2011), and somatic mutation accumulation has been important in

agriculture, where the origin of many clonally-derived varieties of fruits, including citrus, apples, and

wine grapes, have been cultivated by grafting from genetically differentiated bud tips (Aradhya et al.

2003; McKey et al. 2010; Miller and Gross 2011; Vezzulli et al. 2012; Jarni et al. 2015; Pelsy et al. 2015).

However, there is disagreement over the degree and biological importance of somatic mutation

accumulation in plants (Schultz and Scofield 2009; Burian et al. 2016; Watson et al. 2016; Kuhlemeier

2017; Schmid-Siegert et al. 2017; Plomion et al. 2018; Hanlon et al. 2019).

Clonal evolution occurs as individual cell lineages accumulate mutations during mitotic division

(Otto and Orive 1995; Otto and Hastings 1998; Orive 2001; Elena and Lenski 2003; Greaves and Maley

2012; Long et al. 2015). Like mutations occurring during meiosis, somatic mutations are expected to be

deleterious more often than beneficial, but the actual distribution of fitness effects is generally unknown

(Orr 2010). In diploids, new somatic mutations occur as a single copy within the genome, so their

immediate fitness effects will depend on the combination of their selection coefficient (s) and their

expression in the heterozygous state (dominance; h). We expect expressed mutations that reduce rates

of cell division and growth will tend to be eliminated. Similarly, any mutations that elevate the rate of

cell division will tend to increase in frequency in the germ cell population. In general, it is difficult to

assess the effects of somatic mutations, because clonal lineages lack recombination, and the phenotypic

effects of beneficial mutations can be muted by the co-occurrence of detrimental genetic variants (Hill-

Robertson Effect; Selective Interference; Hill and Robertson 1966; Fogle et al. 2008). Although sexual

reproduction does allow mutations to segregate among progeny, it is difficult to detect the phenotypic

effects of novel mutations in eukaryotes with separate sexes, because the genomes of two individuals

are combined in offspring (Long et al. 2015). In this study, we make crosses within individual clones of

hermaphroditic plants to assess the phenotypic effects of novel mutations accumulating during somatic

growth.

78   Plants have several advantages over other organisms for the study of clonal evolution. Separate

stems on the same plant consist of multiple clonal cell lineages that are derived from the same zygote

but may differ for mutations that have accumulated during stem elongation. To produce recombinant

progeny segregating for somatic mutations unique to each stem, crosses can be made either within the

same flower (autogamy) or between flowers on separate stems of the same plant (geitonogamy; Fig. 1).

These crosses are both self-fertilizations, but they differ in the complement of somatic mutations that

are unique to each stem. Progeny from autogamous crosses will segregate for the mutations that

accumulated within a single stem, while the progeny from geitonogamous crosses will segregate for

mutations from both stems. After autogamy, somatic mutations unique to a stem will be homozygous in

approximately 25% of progeny, but they will be either heterozygous or absent in progeny from

geitonogamous crosses. Thus, the average effects of mutations accumulating during stem growth can be

evaluated by comparing the average fitness of progeny generated by autogamous and geitonogamous

crosses. The potential for somatic mutations to accumulate during vegetative growth, and the ability to

study their effects among recombinant progeny, make plants an attractive model for the study of

fundamental processes of clonal evolution.

93   Plants grow from the division of a population of meristem cells within the stem tip that is known

as the central zone. These germ cell lineages go on to produce future stem, leaf, and reproductive

tissues (flowers). As plants grow, individual ramets of the same genet (separate stems or vegetatively

propagated plants) can continue to become differentiated as they accumulate somatic mutations (Fig.

2). Mathematical models have demonstrated that multiple unexpressed mutations (i.e. neutral and

recessive deleterious mutations) are likely to accumulate as germ cells divide, so selective cell deaths

during somatic growth will reduce the total mutational load and alter the composition of mutations

carried by the germ cell population (Otto and Orive 1995; Otto and Hastings 1998). Lineages that carry

expressed deleterious mutations may have slower rates of division and can be eliminated as they are

replaced by cell lineages displaying more rapid growth (Fig. 2A). Moreover, theoretical treatments

predict constant turnover, as expressed deleterious mutations cause the loss of a cell lineage along with

all of its somatic mutations (Otto and Orive 1995; Otto and Hastings 1998; Orive 2001). This process –

referred to as cell lineage selection – has the potential to maintain moderate numbers of somatic

mutations within each meristem over the life of each ramet. More rarely, expressed beneficial

mutations will arise. These variants will be retained and accumulate during stem growth as these cell

lineages outcompete others, which leads to a process analogous to a selective sweep (Otto and Orive

1995; Fig. 2B green arrows; Otto and Hastings 1998; Orive 2001). The end result is that some ramets will

be characterized by a prevalence of deleterious somatic mutations, while others could possess beneficial

mutations. The question we address here is whether somatic mutations that characterize individual

ramets have measurable effects on fitness in the offspring generation.

When we consider the potential for meiotic and somatic mutations to contribute to the total

mutational load of plant populations–particularly for long-lived plants–it becomes evident that not all of

the mutations occurring during a plant's lifespan are passed on to the next generation (Cruzan 2018

pages 86-98). As described above, the process of cell lineage selection will tend to filter deleterious

mutations and favor the retention of beneficial mutations. In addition, some proportion of recessive

deleterious mutations may be eliminated during the haploid life stage due to pollen tube attrition and

pollen competition (Gemetophytic Selection; Mulcahy 1979; Cruzan 1989; Mable and Otto 1998;

Armbruster and Rogers 2004; Arunkumar et al. 2013; Harder et al. 2016). A portion of deleterious

mutations also will be homozygous in zygotes, which can lead to higher rates of seed and fruit abortion

(Selective Embryo Abortion; Husband and Schemske 1995; Korbecka et al. 2002), thereby increasing the

average fitness of surviving offspring (Cruzan and Thomson 1997; Mena-Alĺ and Rocha 2005). Indeed,

the effects of deleterious somatic mutations often are apparent as higher rates of embryo abortion after

autogamous compared to geitonogamous pollinations, which is referred to as autogamy depression

(Schultz and Scofield 2009; Bobiwash et al. 2013). In contrast, mutations that result in faster growth of

stem cells will be favored by cell lineage selection and will have an elevated chance of being inherited.

The sum effects of this intra-generation selection, which includes cell lineage selection, gametophytic

selection, and selective embryo abortion, may result in non-random complements of mutations entering

the next generation.

In this study, we use autogamous and geitonogamous self-pollinations to estimate the effects

of somatic mutations segregating in offspring of *Mimulus guttatus* DC (*Erythranthe guttata* G.L. Nesom;

Phrymaceae). Populations of *M. guttatus* display a wide range of life histories – from annuals to

herbaceous perennials that outcross to varying degrees (Wu et al. 2008). We use perennial *M. guttatus*

plants that produce substantial vegetative growth prior to initiation of flowering. These plants are easily

propagated from rosettes, and selfing produces substantial numbers of seeds to allow for statistical

comparisons. In two separate experiments, we characterized the fitness effects of somatic mutations

that accumulated during vegetative growth of perennial *M. guttatus* plants that were subjected to novel

environments. We made autogamous and geitonogamous self-pollinations across multiple stems and

asked whether there was evidence for autogamy depression for seed set and embryo abortion. In

addition, we estimated the fitness effects of somatic mutations segregating in progeny. We begin by

describing our analytical approach in greater detail, followed by providing specifics of the two

experiments. We show that somatic mutations accumulate during stem growth, as progeny from

autogamous crosses have greater variance in mean fitness compared to progeny from geitonogamous

crosses. We further reveal evidence for the presence of beneficial mutations in some stems, which

indicates that intra-generation selection may be filtering mutations to modify the distribution of fitness

effects transmitted to progeny. The results from these experiments challenge widely accepted notions

concerning mutation accumulation during vegetative growth and the distribution of fitness effects of

mutations in plant populations.

Approach

While previous studies have detected autogamy depression expressed as seed and fruit abortion

(reviewed in Bobiwash et al. 2013), no earlier work has evaluated the effects of somatic mutations on

the fitness of seedlings in the next generation. We use a comparative approach to evaluate the fitness

consequences of somatic mutations that are unique to each stem.

For a diploid plant, we can assume that somatic mutations ($a \rightarrow a'$) will be in the heterozygous

state when they first appear, so in progeny arising by autogamy, they will segregate as approximately

25% homozygous ($a'a'$), 50% heterozygous (aa'), and 25% the original homozygote (aa). By contrast,

because progeny from geitonogamous crosses will be segregating for mutations that are unique to each

stem, about half of them will be carrying mutations in the heterozygous state, and none of the progeny

will be homozygous for mutations that arose in a single stem. The fitness effects of somatic mutations

accumulating during vegetative growth have been quantified in terms of "autogamy depression" (δ_{AD})

using the formula; $\delta_{AD} = 1 - w_A/w_G$ (Schultz and Scofield 2009; Bobiwash et al. 2013), where w_A and w_G

are the relative fitnesses of progeny derived from autogamous and geitonogamous crosses, respectively.

This equation is analogous to the measure of inbreeding depression based on differences in the relative

fitness of offspring derived from selfing vs. outcrossing.

Autogamy depression has been applied previously only to estimates of fitness based on fruit and

seed set, and it generally assumes that the fitness of progeny from autogamous crosses will be lower

than from geitonogamy. Thus, for each stem, we estimated the fitness effects of somatic mutations

unique to that stem ($\bar{s}_k$) as the difference in mean fitness between the progeny from autogamous and

geitonogamous crosses from a single stem: $\bar{s}_k = \overline{w}_{k(A)} - \overline{w}_{k(G)}$, where $\overline{w}_{k(A)}$ and $\overline{w}_{k(G)}$ are the average

fitnesses of the progeny from autogamous and geitonogamous crosses from stem k, respectively.

Provided that environmental effects are controlled for with a common garden design, the parameter $\bar{s}_k$

represents the combined effects of one or more mutations that have different magnitudes and/or

directions of effects on fitness, which provides an estimate of the average fitness effects of all expressed

somatic mutations that have accumulated during vegetative growth. To evaluate the reliability of $\bar{s}_k$ as

an estimate of s we conducted simulations by randomizing the selective effects and dominance of

somatic mutations in pairs of stems. We determined that values of $\bar{s}_k$ were accurate estimates of s

across a range of values and assumptions for the effects of multiple mutations (Appendix 1).

While simulations indicate that our estimates of $\bar{s}_k$ should provide reliable information on the

sign and magnitude of the average fitness effects of somatic mutations unique to a stem, we also

developed an alternative approach for further validation. Individual germ cell lineages within the

meristem may possess unique sets of somatic mutations (Schwoch et al. in preparation), and flowers on

the same stem can develop from only a portion of the germ cells present in the meristem (McManus

and Veit 2002). Hence, no pair of flowers within a stem are likely to contain exactly the same

complement of mutations. Therefore, we generated an additional estimate of s that is based only on the

variance in fitness of offspring from autogamous crosses. If somatic mutations affect offspring fitness,

the variance in fitness should be greater for progeny groups from autogamy than from geitonogamy

from the same stem, as long as mutations do not have complete expression in heterozygotes (i.e. $h <$

1.0; Appendix 2). Somatic mutations that are unique to stems will have different effects on the variance

in fitness among progeny from autogamy depending on the strength of selection (s) and their expression

in the heterozygous state (*h*). To evaluate these effects, we considered a single somatic mutation

occurring in a stem with genotype *aa'* that segregated in offspring in a 1:2:1 ratio following autogamy.

We assumed that the fitness of the *aa*, *aa'*, and *a'a'* genotypes was 1, 1+*hs*, and 1+*s*, respectively (where

s ranges from -1 to 1, and *h* ranges from 0 to 1). Our analytical results demonstrated that the large

majority of variance in fitness among progeny from autogamy (as measured by the standard deviation,

SD) could be explained by the selection coefficient rather than the dominance level (Appendix 2; Fig. S1).

Moreover, the derived relationship allows us to estimate the selective effects of mutations based on the

standard deviation in fitness according to the equation: $s_{SD} = cSD$, where *c* is the slope of the linear

relationship between s_{SD} and *SD*. For dominance levels of *h* = 0.0 and 1.0, *c* = 2.31, and the slope reaches

a maximum of 3.01 when *h* = 0.5 (Appendix 2; Fig. S1). Thus, the variance in fitness among progeny from

autogamous crosses provides a good approximation of the magnitude of the selection coefficient (*s*) for

somatic mutations accumulating in each stem. However, because *s* is estimated from the standard

deviation, only the magnitude of selection but not its direction can be inferred. Thus, although each

approach has potential shortcomings (but for different reasons), identifying consistent results using

both approaches would provide a robust estimate of the average selective effects of somatic mutations

present in individual stems (Appendix 1 and 2).

From the considerations above, we can make three specific predictions that allow us to test for

the fitness effects of somatic mutations that occur during stem growth and are passed on to offspring.

First, there should be greater variation among progeny groups from autogamy compared to those from

geitonogamy (i.e. as indicated by the stem x cross type interaction for progeny fitness). Second, the

average fitness of progeny from autogamy should be different from the progeny of geitonogamous

crosses from the same stem. The difference in the fitness of progeny from autogamous vs.

geitonogamous crosses allows us to make an approximate estimate of the sign and magnitude of

selection coefficients for individual stems. In addition to influencing the mean fitness, somatic mutations

will increase the variance in fitness among progeny from autogamy. Consequently, the third prediction

we can make is that the selective effects of somatic mutations will scale linearly with the within-family

standard deviation in fitness among progeny from autogamous crosses (Appendix 2). In general, we

predict that if somatic mutations with fitness effects accumulate during vegetative growth, then the

mean fitness should differ between progeny from autogamy and geitonogamy, and the variance in

fitness of progeny from autogamous crosses would be greater than for progeny from geitonogamous

crosses. Below, we test these predictions using two different experiments with *Mimulus guttatus*.

First Experiment

225   *Methods*–This experiment originally was designed to test the effects of somatic mutations on

seed set and ovule abortion (autogamy depression; Schultz and Scofield 2009; Bobiwash et al. 2013), so

we used multiple plants from several populations to obtain a large number of fruits. We grew plants of

M. guttatus in the Research Greenhouse facility at Portland State University from seed collected in July

2013 from three different populations in northern Oregon (Jackson Bottom Wetlands–JB: 45.501794 N, -

122.98776 W; and two from Saddle Mountain–SMB: 45.9861 N, -123.6859 W, and SMC: 45.9634 N, -

123.6837 W). We assumed that greenhouse conditions were different enough from field environments

to provide a novel selection regime that would increase the chances mutations would enhance fitness

because genotypes were further from the new fitness optimum (Fisher 1930; Bull and Otto 2005;

Stearns and Fenster 2016). In August 2013, seeds were cold stratified on moist paper towels at 2°C for

30 days prior to being sown in soil. Seedlings were transplanted to pots (approximately 10 x 10 x 12 cm)

and grown for seven months before the application of pollination treatments. Temperature was

maintained between 21–26°C during the day, and 15–21°C at night. Supplemental HID lights ran for 12

hours a day when the seedlings first emerged, and 14 hours a day during adult growth.

239   After plants became established and began producing multiple stems, we conducted

autogamous and geitonogamous self-pollinations using flowers on several stems (15 to 20 cm in length)

from two plants from each of four maternal families representing each of the three populations. Flowers

from pairs of stems on individual plants were reciprocally crossed (geitonogamy), or individual flowers

from these same stems were self-pollinated (autogamy). A total of 139 pollinations were conducted

across two treatments: limited (pollen was applied to stigmas with one touch from a plastic pipette tip)

or excess (where the stigma surface was coated with pollen). Pollinations were conducted on 12

different days (pollination date) over several weeks in July 2014. Mature fruits were collected and

placed individually into paper envelopes, and their contents were examined under a Leica MZ-16

stereoscope. The first 100 ovules from each fruit were categorized as filled seeds (brown, almond-

shaped), unfertilized ovules (small, flattened and light-colored), or aborted (larger than unfertilized,

dark-colored, shriveled). Ovules that were flattened and appeared to lack endosperm were assumed to

be unfertilized or aborted and were not used in germination tests. Seed set and ovule abortion were

analyzed using ANOVA models with the GLM procedure of SAS (SAS 2008), with population, maternal

plant nested within population, and pollination date as random effects, and cross type (autogamous or

geitonogamous) and pollination treatment as fixed effects. Seed set and ovule abortion data were

approximately normal so were not transformed prior to analysis.

We assessed the fitness of progeny arising by autogamy and geitonogamy in the same
greenhouse environment that was used to grow the parental plants. Seedlings from a subset of ten
maternal plants that had fruits from both cross types and at least 20 filled seeds were sown in soil and
transplanted to 36-cell trays (blocks) in September in a randomized incomplete block design. After three
months of growth, the progeny were scored for survival, and above ground biomass was measured after
drying at 60°C for at least 24 hours. The fitness of each progeny was estimated as its final biomass,
which was log transformed and weighted by the survival frequency of progeny from the same cross.
Biomass is considered to be an appropriate estimate of fitness for perennials (Younginger et al. 2017).
These estimates were rescaled relative to the maximum value from all crosses, so that $\overline{w}$ ranged from 0
to 1 across progeny from all crosses. Data were analyzed to estimate the mean fitness for progeny from
autogamy ($\overline{w}_{k(A)}$) and geitonogamy ($\overline{w}_{k(G)}$) progeny from each stem, and the variance in fitness (as
measured by the standard deviation; SD) for each group of progeny produced by autogamy and
geitonogamy from a single stem.
Results–Levels of seed set and germination were similar between cross types, but embryo
abortion was greater in autogamous crosses. The mean number of seeds produced after autogamous
(56.72 ±2.24, N = 87) and geitonogamous (57.52 ±3.42, N = 52) pollinations was similar (Cross in Table
S2), and there was little variation among populations (Pop in Table S2), maternal plants nested within
population (Mat(Pop) in Table S2), and pollination dates (Date in Table S2). Limited pollinations
produced fewer seeds than excess pollinations (Pollen in Table S2), but the effect of pollen dosage in the
overall model was not significant. However, the mean number of aborted ovules was greater for
autogamous (17.36 ±1.27, N = 87) than for geitonogamous (13.90 ±1.70, N = 52) pollinations (Cross in
Table S3), which is consistent with the predicted effects of autogamy depression (Schultz and Scofield
2009; Bobiwash et al. 2013). There also were significant differences among populations in the level of
ovule abortion (Pop in Table S3), which perhaps reflects differences in historical inbreeding and levels of
genetic load. There were significant differences in ovule abortion among maternal plants nested within
population (Mat (Pop) in Table S3), but not for pollination treatments (Pollen in Table S3) or pollination
dates (Date in Table S3). The overall model for ovule abortion was significant (Table S3). Seed
germination was similar between autogamous (mean = 13.03 out of 20 planted per fruit) and
geitonogamous crosses (mean = 13.28; $F_{1,101}$ = 0.16, P = 0.694). Of the 354 seedlings that germinated,
202 survived, and survival was higher for progeny from autogamy (67%) compared to geitonogamy
(50%; chi-square = 12.03, P = 0.0005). Nearly all surviving plants flowered by the end of the experiment,
and flower production was correlated with above ground biomass (r = 0.58, P < 0.001, N = 282).

The average fitness of progeny from autogamy and geitonogamy varied widely among individual

stems. Consistent with our first prediction for the effects of somatic mutations on offspring fitness,

there was a significant interaction between cross type and stem identity (Table S4). Little of the variation

in progeny fitness was explained by differences among stems (Mat in Table S4), but larger amounts were

explained by cross type (Cross in Table S4) and among blocks (Tray in Table S4). When data were

analyzed separately for each cross type, we found that mean fitness of progeny from autogamy varied

among stems to a much greater degree ($F_{8,130}$ = 4.44, P < 0.0001) than progeny from geitonogamous

crosses ($F_{8,70}$ = 1.47, P = 0.1866); a result that supports our first prediction for the effects of somatic

mutations on progeny fitness. These data are analyzed further along with data from the second

experiment described below.

Second Experiment

The observation of greater variation in fitness among stems for progeny from autogamous

crosses in the first experiment is consistent with the hypothesis that somatic mutations arising during

vegetative growth are transmitted and can have substantial phenotypic effects in the next generation.

Furthermore, the results from the first experiment do not support the view that somatic mutations

during vegetative growth are rare (Burian et al. 2016; Groot and Laux 2016; Watson et al. 2016;

Kuhlemeier 2017; Schmid-Siegert et al. 2017), but they are consistent with several studies

demonstrating that somatic mutations can affect offspring fitness (Klekowski and Godfrey 1989; Schultz

and Scofield 2009; Bobiwash et al. 2013). However, the first experiment was not designed specifically to

compare the effects of autogamous and geitonogamous pollinations on offspring fitness. Thus, to

confirm results from the first experiment, we performed a second experiment to provide a more robust

assessment of the effects of autogamous and geitonogamous self-pollination on the fitness of offspring.

The second experiment used a single clone of *M. guttatus* to make comparisons between autogamous

and geitonogamous pollinations paired at the same node. The plant chosen was a self-compatible

perennial that displayed vigorous vegetative growth. We attempted to increase the potential for the

appearance of beneficial mutation by exposing plants to novel conditions provided by high salinity under

hydroponics cultivation. A control hydroponics treatment without added salt was included with an

expectation that fewer mutations would appear. Our recent genomic results confirm higher numbers of

low-frequency somatic mutation for plants exposed to high salt stress (Schwoch et al. in preparation).

This design provided a more refined test of the hypothesis that the fitness effects of somatic mutations

differed in the offspring of autogamous and geitonogamous crosses.

Methods–To assess the fitness effects of mutations that accumulated during vegetative growth,

a single plant (genet BV, obtained from Willamette Gardens native plant nursery, Corvallis, OR) was

vegetatively propagated to generate 12 plants (ramets) that were exposed to high salinity and control

conditions. Plants were grown in pea gravel (4–8 mm) in pots placed in four 53 L tubs using a flood and

drain hydroponics system (flooding at 15 min intervals). Two tubs had no added salt and were used as

controls, and two tubs had high salinity. The initial salt concentration in the high salinity treatment was

5 mM but increased weekly to 25 mM after plants became established. Salt concentrations were

monitored using a conductivity meter to ensure stable concentrations. To provide nutrients, 30 ml of

hydroponics fertilizer (FloraGrow, Planet Natural, Bozeman, MT) was added per tub. During the course

of the experiment, some plants grew substantially. We imposed selection to favor the fastest growing

ramets over the next three months by repeatedly removing single rosettes and transplanting them back

into the hydroponics system up to four times.

To promote stress recovery, plants were transplanted to soil for six months, which included a

two-month vernalization period in a growth chamber (4°C and 8 h light; Conviron E8, Controlled

Environments Ltd., Winnipeg, Manitoba, Canada). After vernalization, plants were returned to the

greenhouse to induce flowering. Autogamous and geitonogamous pollinations were made to pairs of

flowers at single nodes or consecutive nodes (seven nodes and 14 pollinations total) on the largest

ramets in each of the control and high salt treatments. To account for somatic mutation turnover that

may occur due to the effects of cell lineage selection during stem growth, we compared progeny from

autogamous and geitonogamous pollinations at pairs of flowers from the same node. Without *a priori*

knowledge of the expression of somatic mutations in heterozygotes, it is difficult to determine the best

pollen donor for geitonogamous crosses. Consequently, we opted to generate progeny from

geitonogamy by pollinating flowers with pollen from a potentially more genetically divergent ramet

from the other treatment (i.e. salt pollinated with control pollen, and control pollinated with salt

pollen). The fruits were collected, and the total number of aborted and mature seeds and unfertilized

ovules were counted under a dissecting microscope. Seeds were planted in soil in trays with three seeds

per cell. Seeds were cold stratified in moist soil for three weeks before they germinated in the

greenhouse.

To determine whether seedlings from autogamous crosses from ramets exposed to salt stress

showed improved performance under the same conditions, all progeny were exposed to high salinity.

After germination and establishment in soil, seedlings from autogamous and geitonogamous crosses

from control and salt stress ramets were transplanted into pots filled with pea gravel and subjected to

high salt in the hydroponics system, as described above. A total of 239 seedlings from 11 fruits (five

autogamous and six geitonogamous) were randomly and evenly distributed among 12 hydroponic tubs

to ensure equal representation across blocks (tubs). Plant size was measured as the product of the

length and width of vegetative spread after two months of growth and was used as a proxy for biomass.

Salt concentration increased from 10 mM - 37.5 mM over the course of the experiment to induce

mortality (~57% across all progeny groups). Fitness was estimated as plant size (log transformed to

improve normality) weighted by the survival frequency for progeny from the same cross.

359   *Results*–Similar to experiment 1 and consistent with our first prediction for the effects of

somatic mutations, there was greater variation in the fitness of progeny from autogamous compared to

geitonogamous crosses, as indicated by a significant interaction between cross type and stem/node

identity (Stem/Node*Cross in Table S5). There was no consistent difference in progeny fitness between

autogamous and geitonogamous pollinations when averaged across stems (Cross in Table S5), but there

were larger differences among individual stems/nodes (Stem/Node in Table S5). Also consistent with our

first prediction, there was significant variation in progeny fitness among stems ($F_{2,89}$ = 3.76, P = 0.028)

and nodes (nested within stems; $F_{3,89}$ = 2.59, P = 0.0596) for autogamous, but not geitonogamous

crosses ($F_{1,136}$ = 0.63, P = 0.431 and $F_{4,136}$ = 1.19, P = 0.317 for stems and nodes, respectively). There were

significant differences in average size among the blocks (Tray in Table S5), but there was no consistent

difference in the performance of progeny based on the treatment history of their maternal ramets ($F_{1,126}$

= 0.02, P = 0.888), and there was no interaction between cross type and historical treatment ($F_{1,136}$ =

0.64, P = 0.423). Three nodes were dropped from further analysis, because one of the paired fruits

aborted or produced fewer than five seedlings. However, similar to experiment 1, and consistent with

the hypothesis that somatic mutations accumulate during stem growth, the variance in fitness among

progeny from autogamy among stems was greater than for geitonogamy. Since results from the two

experiments were similar, the data for four nodes from experiment 2 were analyzed together with the

first experiment, as described in the next section.

Selection Coefficient Estimates

Results from the two experiments described above are consistent with the hypothesis that somatic

mutations unique to individual stems can have demonstrable effects on the fitness of progeny when a

proportion is made homozygous by autogamous self-pollination. In both experiments, the variance in

fitness among progeny groups from autogamy was significantly greater than among progeny from

geitonogamy from the same set of stems. Since the results from these two experiments are qualitatively

similar, we combined them to estimate the selection coefficients of somatic mutations in each stem (ten

comparisons of autogamy and geitonogamy from Experiment 1 and four from Experiment 2, for a total

of 14 comparisons). Specifically, we tested the second and third predictions described above. We expect

that the effect of somatic mutations will be greater for progeny from autogamy, so their mean fitness

should be different from progeny of geitonogamous crosses on the same stem (providing an estimate of

$\bar{s}_k$). Moreover, the within-family variance in fitness among progeny should be greater after autogamy

compared to geitonogamy (providing an estimate of s_{SD}). These estimates of the selective effects of

somatic mutations should be similar, so we assessed whether there was a positive relationship between

the variance in fitness among progeny from single fruits produced by autogamous pollination (s_{SD}) and

the selection coefficient estimated from the average fitness of progeny from autogamous and

geitonogamous crosses to the same stem ($\bar{s}_k$).

Estimates of the average selective effects of somatic mutations unique to each stem were made

based on the fitness of progeny from autogamous ($\overline{w}_{k(A)}$) and geitonogamous ($\overline{w}_{k(G)}$) crosses (Table S1

in Appendix 3). In both experiments, fitness was estimated as the above ground biomass weighted by

the survival of seedlings from the same fruit and scaled to a maximum value of 1.0. The average

selective value of mutations ($\bar{s}_k$) for each stem (first experiment) or stem/node combination (second

experiment) was calculated as the difference in fitness between progeny from autogamous and

geitonogamous crosses, as described above ($\bar{s}_k = \overline{w}_{k(A)} - \overline{w}_{k(G)}$). We observed extensive variation in $\bar{s}_k$

among stems, with nine stems that were significantly different from zero (Fig. 3). Four of the stems had

average selective effects of somatic mutations that were positive, suggesting a net beneficial effect of

somatic mutations transmitted to offspring. In addition, the average fitness of progeny from

autogamous and geitonogamous crosses on the same stems or nodes was positively correlated (Fig. S2;

Table S6). Note also that the $\bar{s}_k$ estimates that deviated the most from zero generally had the highest (or

lowest) $\overline{w}_A$ values relative to all of the stems (Fig. S3). However, this was not always the case, as one

stem (stem 9) with a value of $\bar{s}_k$ close to zero produced progeny with relatively high fitness after both

autogamous and geitonogamous pollination.

To make additional estimates of the average selection coefficients for mutations unique to each

stem or node, by using the standard deviation in progeny fitness from autogamous crosses based on the

relationship $s_{SD} = cSD$, where $c = 3$, which assumes dominance close to $h = 0.5$ (qualitatively similar

results are obtained if we choose other values of c between 3 and 2.31; Appendix 2). Because the sign of

s_{SD} could not be inferred directly from this approach, we used the sign estimated from the $\bar{s}_k$ method

(Fig. 4; Appendix 2). Our simulations show that the sign of $\bar{s}_k$ is almost always estimated correctly

(Appendix 1). For stems with values of $\bar{s}_k$ greater than zero, there was a strong positive relationship

between $\bar{s}_k$ and estimates of s_{SD} made from the within-family variation among progeny from autogamy

(Fig. 4). In contrast, the relationship for negative values of $\bar{s}_k$ appeared to be driven largely by a single

observation. Note that this observation was not supported by a similarly high value of s_{SD}, and the

remaining negative selection coefficients were more modest based on both estimates. This one highly

negative, and apparently inaccurate estimate of $\bar{s}_k$, may be due to a large influence of mutations from

the second stem on the fitness of geitonogamous progeny (note that this stem had the highest estimate

of $\bar{w}_{k(G)}$; Fig. S2). It is also notable that the variance in fitness for progeny from autogamy did not

decline to zero for values of $\bar{s}_k$ close to zero, which could be due to the presence of both beneficial and

detrimental mutations, and possibly genetic background effects (i.e. epistasis), but it may also reflect

uncontrolled environmental variation. Overall, the results from both approaches reveal consistent

estimates of the average fitness consequences associated with the accumulation of somatic mutations

in individual stems.

Discussion

The observation of higher variation in fitness among stems for progeny from autogamous compared to

geitonogamous self-pollinations provides strong evidence for the accumulation of somatic mutations

during vegetative growth in *Mimulus guttatus*. Two estimates of the effects of selection due to somatic

mutations provided consistent results on offspring fitness: 1) Differences in the mean fitness between

progeny from autogamy and geitonogamy ($\bar{s}_k$), and 2) variation in fitness of autogamous progeny (s_{SD}).

We found evidence for the prevalent effects of beneficial mutations in progeny, with estimates of $\bar{s}_k$

and s_{SD} exceeding 0.1 in four cases, while estimates for negative selection coefficients were more

modest (mostly > -0.15). These results are consistent with the hypothesis that somatic mutation

accumulation during vegetative growth is substantial enough to have demonstrable effects on progeny

fitness and to contribute to standing genetic variation. Furthermore, the observation that progeny from

autogamous crosses occasionally had higher fitness than progeny from geitonogamous crosses argues

for the accumulation of beneficial mutations and implies that many deleterious mutations are culled by

the various types of intra-generation selection prior to seed dispersal. These results suggest that somatic

mutations are generated during vegetative growth, but many of them are filtered due to intra-

generation selection, which results in a shift in the distribution of fitness effects for the mutations that

are passed on to the next generation.

Somatic mutations accumulating during vegetative growth had an overall positive effect for four

of the stems tested. This result is contrary to widely held views that the accumulation of beneficial

somatic mutations should be exceedingly rare (Crow 1993; Charlesworth and Willis 2009). However, a

potential explanation for these findings is afforded by the unique biology of plants; somatic mutations in

meristem cells could contribute to higher rates of division for some cell lineages over others (Otto and

Orive 1995; Otto and Hastings 1998). Thus, cell lineage selection has the potential to retain beneficial

somatic mutations, as individual cell lineages divide at faster rates. Although there may be few

opportunities for beneficial changes to alter basic cellular metabolism, it is becoming apparent from

experimental evolution studies with microbes that even basic aspects of cellular metabolism can be

sensitive to environmental conditions, which can increase the chances that mutations in clonal

populations are beneficial (e.g., Lee and Marx 2013; Maharjan et al. 2015). In this regard, cell lineage

selection in a plant meristem is analogous to clonal evolution in microbial populations, and it represents

a potentially powerful forum for the removal of deleterious somatic mutations while favoring the

retention of beneficial ones. In addition, gametophytic selection and selective embryo abortion can act

as prominent filters (Mulcahy 1979; Cruzan 1989; Mable and Otto 1998; Armbruster and Rogers 2004;

Arunkumar et al. 2013; Harder et al. 2016), but they are most likely to have effects by culling deleterious

mutations. Regardless, the combined effects of these different forms of intra-generation selection

appear to have had a considerable effect on filtering of somatic mutations, such that the distribution of

fitness effects among stems has shifted to include more beneficial mutations than expected.

An alternative explanation for the observed phenotypic changes is that exposure to

environmental stress has induced heritable epigenetic modifications (Quadrana and Colot 2016).

However, epigenetic modifications are often considered to be consistent in direction and predictable, as

they are hypothesized to represent an adaptive response to historic exposure to similar stressors

(Baulcombe and Dean 2014; Itabashi et al. 2018). The results of the experiments described here are

unlikely to support a role for epigenetics, because phenotypic responses in the next generation were

inconsistent in direction and magnitude, and they were not predictable based on environmental

exposure of the parent stem. Indeed, among stems and nodes, the mean fitness of progeny from

autogamy displayed both increases and decreases compared to the geitonogamy controls, which is

consistent with the hypothesis that individual ramets are accumulating unique complements of somatic

mutations. Furthermore, these conclusions are supported by the observation of numerous somatic

variants in the transcriptomes of meristem tissue from multiple ramets derived from a single genet

(Schwoch et al. in preparation). Thus, the results of the current study appear to indicate that somatic

mutations accumulating during stem growth are responsible for the observed phenotypic effects.

The potential for the acquisition of mutations during vegetative growth is a well-known aspect

of plant biology (Klekowski 2003; Schultz and Scofield 2009; Bobiwash et al. 2013), but no previous study

has demonstrated the effects of somatic mutations on the fitness of progeny in the next generation.

Most studies have focused on the detrimental effects of somatic mutations; chloroplast mutants have

been observed in a number of species (Klekowski 2003), declines in pollen fertility were found in older

clones of quaking aspen (Ally et al. 2010), and higher rates of seed and fruit abortion after autogamous

pollinations were found in several studies (reviewed in Bobiwash et al. 2013). In contrast, agriculturalists

have taken advantage of beneficial somatic mutations by conducting clonal selection to improve

economically important plants (Aradhya et al. 2003; McKey et al. 2010; Miller and Gross 2011; Vezzulli

et al. 2012; Jarni et al. 2015; Pelsy et al. 2015), and a handful of studies report phenotypic responses to

selection in asexual lineages. For example, Breese et al. (1965) succeeded in selecting for increased

tillering ability (production of new grass stems) within genets of perennial ryegrass (*Lolium perenne*).

Similarly, artificial selection on clonal lineages effectively improved branching in the red seaweed,

Asparagopsis armata (Monro and Poore 2009). The current study on *M. guttatus* contributes to this

literature by highlighting the potential for plants to exhibit significant levels of clonal variation within a

single generation.

The transmission of beneficial mutations in autogamous crosses may explain some heretofore

difficult to understand results from mutation accumulation studies. Our results suggest that beneficial

somatic mutations are likely to be partially dominant (i.e. not completely recessive), because they would

have to be expressed in the heterozygous state to be favored by cell lineage selection. Mutations

accumulating during vegetative growth could contribute to standing genetic variation, but autogamy

may be more effective for the accumulation of beneficial somatic mutations in populations than

geitonogamy or outcrossing, because beneficial mutations could be made homozygous in progeny in a

single generation. Depending on the crossing design, outcrossing would take at least two generations for

beneficial somatic mutations to be made homozygous, and under geitonogamy they would remain

heterozygous. It is striking that high rates of beneficial mutation accumulation have been observed in at

least some mutation accumulation studies in the autogamous plant *Arabidopsis thaliana* (Shaw et al.

2002; Rutter et al. 2010; Rutter et al. 2012), but not in outcrossing and partially-selfing species of

Amsinckia (Schoen 2005). With a few exceptions (e.g., Baer et al. 2005; Denver et al. 2010), nearly all

mutation accumulation studies on animals consistently show a prevalence of deleterious mutations

(Baer et al. 2007; Halligan and Keightley 2009). Similarly, our results suggest that the adaptive potential

of autogamous plants may be greater than previously thought, which may help explain the wider

geographic ranges of selfing compared to closely-related outcrossing species (Grossenbacher et al.

2015). Although intra-generation selection has the potential to contribute to adaptation in all plants, its

effects may be enhanced in autogamous lineages, because beneficial mutations arising during

vegetative growth have a greater chance of becoming homozygous in offspring and being retained

across generations.

As stems grow, mutations are generated during every mitotic cell division, so the potential for

somatic mutation accumulation in plants appears substantial. Thus, understanding how long-lived

plants, such as trees, avoid mutational meltdown from the accumulation of deleterious somatic

mutations remains a longstanding question. Paradoxically, however, the rate of mutation accumulation

observed across generations in plant and animal genomes is similar (Gaut et al. 2011). One explanation

for this pattern is that somatic mutations in plants are repressed during vegetative growth, similar to

animal germlines, which would protect lineages from the negative effects of mutation accumulation

during development of the soma (Plant Germline Hypothesis; Burian et al. 2016; Cruzan 2018; page 90).

An alternative hypothesis posits that somatic mutations are generated in apical meristems during plant

growth, but these mutations are filtered by intra-generation selection occurring prior to the

establishment of offspring (Somatic Mutation Accumulation Hypothesis). Because plants have retained

the capacity to undergo clonal evolution from their algal ancestors, the ability to filter mutations during

growth and reproduction has existed for some time, and thus intra-generation selection has the

potential to skew the distribution of fitness effects of transmitted mutations to include a larger

proportion of beneficial mutations than would be expected through random processes. In addition, this

provides a reasonable explanation for why longer-lived plants appear to have slower rates of mutation

accumulation across generations (Yue et al. 2010; Gaut et al. 2011). This could be due to the fact that

longer generation time leads to more time between recombination events, which can lead to more

background selection in non-recombining cell lineages during vegetative growth (Cruzan 2018; pages 94-

95). Recent studies in oaks and spruce (Plomion et al. 2018; Hanlon et al. 2019), as well as our

unpublished data from *M. guttatus*, confirm that multiple somatic variants are likely, even in plants of

very different size. Moreover, results from the current study identify the presence of beneficial

mutations that may be passed on to offspring in the next generation, likely a consequence of intra-

generation selection. Future work that combines information from experiments evaluating the genomic

consequences of somatic variation with anatomical estimates of stem cell population dynamics will

allow for the development of new models that provide insights into the extent and limitations of

somatic evolution in plants.

In conclusion, despite the potential for somatic mutation accumulation to generate novel

genetic variation in plant populations, its role in evolution remains almost entirely unexplored. Our

estimates of the selective effects of somatic mutations were consistent across several methods and

indicate that some stems possessed a prevalence of deleterious mutations while others produced

autogamous progeny with high fitness, which indicates a prevalence of beneficial mutations. Even

though high levels of mutation accumulation are often believed to be detrimental, the basic biology of

plants suggests that the role of somatic mutations in plant evolution should be considered carefully in

the future. Moving forward, our results indicate that we must keep in mind that all eukaryotes are not

necessarily equivalent, and features of the organism's biology may have unexpected consequences for

the evolutionary phenomena that we observe. Future lines of investigation will improve our

understanding of these fundamental aspects of plant development and evolution that may have

contributed to the remarkable diversification of plants, and may help to account for some of the

extensive variation in mutation rates detected among lineages.

Acknowledgements

We thank J. Thompson, E. Perez, and our research greenhouse manager Linda Taylor for plant

maintenance and assistance with this research in the greenhouse and in the lab. Several people made

helpful suggestions on this manuscript including N. Diaz, M. Grasty, and J. Persinger. This research was

supported by an NIH NIGMS BUILD EXITO grant (5TL4GM118965-03, 5UL1GM118964-03, and

5RL5GM118963-03) to Portland State University.

Data Archiving

Data and all scripts for data analysis will be submitted to DataDryad upon publication.

Literature Cited

Ally, D., K. Ritland, and S. P. Otto. 2010. Aging in a Long-Lived Clonal Tree. PLoS Biol. 8.
(http://doi.org/10.1371/journal.pbio.1000454).

Antolin, M. F. and C. Strobeck. 1985. The population genetics of somatic mutation in plants. The
American Naturalist 126:52-62. (http://doi.org/10.1086/284395).

Aradhya, M. K., G. S. Dangl, B. H. Prins, J. M. Boursiquot, M. A. Walker, C. P. Meredith, and C. J. Simon.
2003. Genetic structure and differentiation in cultivated grape, *Vitis vinifera* L. Genet. Res.
81:179-192. (http://doi.org/10.1017/s0016672303006177).

Armbruster, W. S. and D. G. Rogers. 2004. Does pollen competition reduce the cost of inbreeding? Am. J.
Bot. 91:1939-1943. (http://doi.org/10.3732/ajb.91.11.1939).

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not sperm-
specific, genes show stronger purifying selection and higher rates of positive selection than
sporophytic genes in *Capsella grandiflora*. Mol. Biol. Evol. 30:2475-2486.
(http://doi.org/10.1093/molbev/mst149).

Baer, C. F., M. M. Miyamoto, and D. R. Denver. 2007. Mutation rate variation in multicellular eukaryotes:
causes and consequences. Nature Reviews Genetics 8:619-631.

Baer, C. F., F. Shaw, C. Steding, M. Baumgartner, A. Hawkins, A. Houppert, N. Mason, M. Reed, K.
Simonelic, W. Woodard, and M. Lynch. 2005. Comparative evolutionary genetics of spontaneous
mutations affecting fitness in rhabditid nematodes. Proceedings of the National Academy of
Sciences of the United States of America 102:5785-5790.
(http://doi.org/10.1073/pnas.0406056102).

Barfield, S., G. V. Aglyamova, and M. V. Matz. 2016. Evolutionary origins of germline segregation in
Metazoa: evidence for a germ stem cell lineage in the coral <i>Orbicella faveolata</i> (Cnidaria,
Anthozoa). Proceedings of the Royal Society B: Biological Sciences 283:20152128.
(http://doi.org/doi:10.1098/rspb.2015.2128).

Baulcombe, D. C. and C. Dean. 2014. Epigenetic regulation in plant responses to the environment. Cold
Spring Harbor Perspectives in Biology 6:a019471.
(http://doi.org/10.1101/cshperspect.a019471).

Beerenwinkel, N., R. F. Schwarz, M. Gerstung, and F. Markowetz. 2015. Cancer evolution: mathematical
models and computational inference. Syst. Biol. 64:e1-25.
(http://doi.org/10.1093/sysbio/syu081).

Bobiwash, K., S. T. Schultz, and D. J. Schoen. 2013. Somatic deleterious mutation rate in a woody plant:
estimation from phenotypic data. Heredity 111:338-344. (http://doi.org/10.1038/hdy.2013.57).

Breese, E. L., M. D. Hayward, and A. C. Thomas. 1965. Somatic selection in perennial ryegrass. Heredity
20:367-&. (http://doi.org/10.1038/hdy.1965.50).

Bull, J. J. and S. P. Otto. 2005. The first steps in adaptive evolution. Nat. Genet. 37:342-343.
(http://doi.org/10.1038/ng0405-342).

Burian, A., P. B. de Reuille, and C. Kuhlemeier. 2016. Patterns of stem cell divisions contribute to plant
longevity. Curr. Biol. 26:1385-1394. (http://doi.org/10.1016/j.cub.2016.03.067).

Buss, L. W. 1983. Evolution, development, and the units of selection. Proceedings of the National
Academy of Sciences of the United States of America-Biological Sciences 80:1387-1391.
(http://doi.org/10.1073/pnas.80.5.1387).

Charlesworth, D. and J. H. Willis. 2009. The genetics of inbreeding depression. Nat. Rev. Genet. 10.
(http://doi.org/10.1038/nrg2664).

Crow, J. F. 1993. Mutation, mean fitness, and genetic load. Pp. 3-42. Oxford Surveys in Evolutionary
Biology. Oxford University Press.

Cruzan, M. and J. Thomson. 1997. Effects of pre-dispersal selection on offspring growth and survival in
Erythronium grandiflorum. J. Evol. Biol. 10:295-314.
Cruzan, M. B. 1989. Pollen tube attrition in *Erythronium grandiflorum*. Amer. J. Bot. 76:562-570.
Cruzan, M. B. 2018. Evolutionary Biology - A Plant Perspective. Oxford University Press, New York.
Denver, D. R., D. K. Howe, L. J. Wilhelm, C. A. Palmer, J. L. Anderson, K. C. Stein, P. C. Phillips, and S.
Estes. 2010. Selective sweeps and parallel mutation in the adaptive recovery from deleterious
mutation in Caenorhabditis elegans. Genome Res. 20:1663-1671.
(http://doi.org/10.1101/gr.108191.110).
Dubrovina, A. S. and K. V. Kiselev. 2016. Age-associated alterations in the somatic mutation and DNA
methylation levels in plants. Plant Biol. 18:185-196. (http://doi.org/10.1111/plb.12375).
Elena, S. F. and R. E. Lenski. 2003. Evolution experiments with microorganisms: the dynamics and
genetic bases of adaptation. Nat. Rev. Genet. 4:457-469.
Fisher, R. A. 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.
Fogle, C. A., J. L. Nagle, and M. M. Desai. 2008. Clonal interference, multiple mutations and adaptation in
large asexual populations. Genetics 180:2163-2173.
(http://doi.org/10.1534/genetics.108.090019).
Gaut, B., L. Yang, S. Takuno, and L. E. Eguiarte. 2011. The patterns and causes of variation in plant
nucleotide substitution rates. Pp. 245-266 *in* D. J. Futuyma, H. B. Shaffer, and D. Simberloff, eds.
Annual Review of Ecology, Evolution, and Systematics, Vol 42.
Greaves, M. and C. C. Maley. 2012. Clonal evolution in cancer. Nature 481:306-313.
(http://doi.org/http://www.nature.com/nature/journal/v481/n7381/abs/nature10762.html#su
pplementary-information).
Groot, Edwin P. and T. Laux. 2016. Ageing: how do long-lived plants escape mutational meltdown? Curr.
Biol. 26:R530-R532. (http://doi.org/https://doi.org/10.1016/j.cub.2016.05.049).
Grossenbacher, D., R. Briscoe Runquist, E. E. Goldberg, and Y. Brandvain. 2015. Geographic range size is
predicted by plant mating system. Ecol. Lett. 18:706-713. (http://doi.org/10.1111/ele.12449).
Halligan, D. L. and P. D. Keightley. 2009. Spontaneous Mutation Accumulation Studies in Evolutionary
Genetics. Pp. 151-172. Annual Review of Ecology Evolution and Systematics.
Hanlon, V. C. T., S. P. Otto, and S. N. Aitken. 2019. Somatic mutations substantially increase the per-
generation mutation rate in the conifer Picea sitchensis. Evolution Letters 3:348-358.
(http://doi.org/doi:10.1002/evl3.121).
Harder, L. D., M. A. Aizen, and S. A. Richards. 2016. The population ecology of male gametophytes: the
link between pollination and seed production. Ecol. Lett. 19:497-509.
(http://doi.org/10.1111/ele.12596).
Hill, W. G. and A. Robertson. 1966. The effect of linkage on limits to artificial selection. Genetics
Research 8:269-294.
Husband, B. and D. W. Schemske. 1995. Evolution of the magnitude and timing of inbreeding depression
in plants. Evolution 50:54-70.
Itabashi, E., K. Osabe, R. Fujimoto, and T. Kakizaki. 2018. Epigenetic regulation of agronomical traits in
Brassicaceae. Plant Cell Rep. 37:87-101. (http://doi.org/10.1007/s00299-017-2223-z).
Jarni, K., J. Jakse, and R. Brus. 2015. Vegetative propagation: linear barriers and somatic mutation affect
the genetic structure of a Prunus avium L. stand. Forestry 88:612-621.
(http://doi.org/10.1093/forestry/cpv029).
Klekowski, E. J. 2003. Plant clonality, mutation, diplontic selection and mutational meltdown. Biol. J.
Linn. Soc. 79:61-67. (http://doi.org/10.1046/j.1095-8312.2003.00183.x).
Klekowski, E. J. and P. J. Godfrey. 1989. Ageing and mutation in plants. Nature 340:389-391.
(http://doi.org/10.1038/340389a0).

Korbecka, G., P. G. L. Klinkhamer, and K. Vrieling. 2002. Selective embryo abortion hypothesis revisited - A molecular approach. Plant Biol. 4:298-310. (http://doi.org/10.1055/s-2002-32331).

Kuhlemeier, C. 2017. How to get old without aging. Nature Plants 3:916-917. (http://doi.org/10.1038/s41477-017-0076-7).

Lee, M.-C. and C. J. Marx. 2013. Synchronous waves of failed soft sweeps in the laboratory: remarkably rampant clonal Interference of alleles at a single locus. Genetics 193:943-+. (http://doi.org/10.1534/genetics.112.148502).

Long, A., G. Liti, A. Luptak, and O. Tenaillon. 2015. Elucidating the molecular architecture of adaptation via evolve and resequence experiments. Nature Reviews Genetics 16:567-582. (http://doi.org/10.1038/nrg3937).

Mable, B. K. and S. P. Otto. 1998. The evolution of life cycles with haploid and diploid phases. Bioessays 20:453-462. (http://doi.org/10.1002/(sici)1521-1878(199806)20:6<453::aid-bies3>3.0.co;2-n).

Maharjan, R. P., B. Liu, L. Feng, T. Ferenci, and L. Wang. 2015. Simple phenotypic sweeps hide complex genetic changes in populations. Genome Biology and Evolution 7:531-544. (http://doi.org/10.1093/gbe/evv004).

McKey, D., M. Elias, B. Pujol, and A. Duputie. 2010. The evolutionary ecology of clonally propagated domesticated plants. New Phytol. 186:318-332. (http://doi.org/10.1111/j.1469-8137.2010.03210.x).

McKnight, T. D., K. Riha, and D. E. Shippen. 2002. Telomeres, telomerase, and stability of the plant genome. Plant Mol. Biol. 48:331-337. (http://doi.org/10.1023/a:1014091032750).

McManus, M. T. and B. E. Veit, eds. 2002. Meristematic Tissues in Plant Growth and Development. CRC Press LLC, Boca Raton, FL.

Mena-Alí, J. I. and O. J. Rocha. 2005. Selective seed abortion affects the performance of the offspring in *Bauhinia ungulata*. Ann. Bot. 95:1017-1023. (http://doi.org/10.1093/aob/mci109).

Miller, A. J. and B. L. Gross. 2011. From forest to field: perennial fruit crop domestication. Am. J. Bot. 98:1389-1414. (http://doi.org/10.3732/ajb.1000522).

Monro, K. and A. G. B. Poore. 2009. The potential for evolutionary responses to cell-lineage selection on growth form and its plasticity in a red seaweed. Am. Nat. 173:151-163. (http://doi.org/10.1086/595758).

Mulcahy, D. L. 1979. The rise of the angiosperms: A genecological factor. Science 206:20-23.

Nowell, P. C. 1976. Clonal evolution of tumor-cell populations. Science 194:23-28. (http://doi.org/10.1126/science.959840).

Orive, M. E. 2001. Somatic mutations in organisms with complex life histories. Theor. Popul. Biol. 59:235-249. (http://doi.org/10.1006/tpbi.2001.1515).

Orr, H. A. 2010. The population genetics of beneficial mutations. Philos Trans R Soc Lond B Biol Sci 365:1195-1201. (http://doi.org/10.1098/rstb.2009.0282).

Otto, S. P. and I. M. Hastings. 1998. Mutation and selection within the individual. Genetica 102-103:507-524. (http://doi.org/10.1023/a:1017074823337).

Otto, S. P. and M. E. Orive. 1995. Evolutionary consequences of mutation and selection within an individual. Genetics 141:1173-1187.

Pelsy, F., V. Dumas, L. Bevilacqua, S. Hocquigny, and D. Merdinoglu. 2015. Chromosome Replacement and Deletion Lead to Clonal Polymorphism of Berry Color in Grapevine. PLoS Genet. 11. (http://doi.org/10.1371/journal.pgen.1005081).

Plomion, C., J.-M. Aury, J. Amselem, T. Leroy, F. Murat, S. Duplessis, S. Faye, N. Francillonne, K. Labadie, G. Le Provost, I. Lesur, J. Bartholomé, P. Faivre-Rampant, A. Kohler, J.-C. Leplé, N. Chantret, J. Chen, A. Diévart, T. Alaeitabar, V. Barbe, C. Belser, H. Bergès, C. Bodénès, M.-B. Bogeat-Triboulot, M.-L. Bouffaud, B. Brachi, E. Chancerel, D. Cohen, A. Couloux, C. Da Silva, C. Dossat, F. Ehrenmann, C. Gaspin, J. Grima-Pettenati, E. Guichoux, A. Hecker, S. Herrmann, P. Hugueney, I.

Hummel, C. Klopp, C. Lalanne, M. Lascoux, E. Lasserre, A. Lemainque, M.-L. Desprez-Loustau, I. Luyten, M.-A. Madoui, S. Mangenot, C. Marchal, F. Maumus, J. Mercier, C. Michotey, O. Panaud, N. Picault, N. Rouhier, O. Rué, C. Rustenholz, F. Salin, M. Soler, M. Tarkka, A. Velt, A. E. Zanne, F. Martin, P. Wincker, H. Quesneville, A. Kremer, and J. Salse. 2018. Oak genome reveals facets of long lifespan. Nature Plants 4:440-452. (http://doi.org/10.1038/s41477-018-0172-3).

Quadrana, L. and V. Colot. 2016. Plant Transgenerational Epigenetics. Pp. 467-491 *in* N. M. Bonini, ed. Annual Review of Genetics, Vol 50.

Reusch, T. B. H. and C. Bostrom. 2011. Widespread genetic mosaicism in the marine angiosperm Zostera marina is correlated with clonal reproduction. Evol. Ecol. 25:899-913. (http://doi.org/10.1007/s10682-010-9436-8).

Rutter, M. T., A. Roles, J. K. Conner, R. G. Shaw, F. H. Shaw, K. Schneeberger, S. Ossowski, D. Weigel, and C. B. Fenster. 2012. Fitness of *Arabidopsis thaliana* mutation accumulation lines whose spontaneous mutations are known. Evolution 66:2335-2339. (http://doi.org/10.1111/j.1558-5646.2012.01583.x).

Rutter, M. T., F. H. Shaw, and C. B. Fenster. 2010. Spontaneous mutation parameters for *Arabidopsis thaliana* measured in the wild. Evolution 64:1825-1835. (http://doi.org/10.1111/j.1558-5646.2009.00928.x).

SAS. 2008. SAS/STAT 9.2 user's guide. SAS Institute Inc., Cary, North Carolina, USA.

Schmid-Siegert, E., N. Sarkar, C. Iseli, S. Calderon, C. Gouhier-Darimont, J. Chrast, P. Cattaneo, F. Schütz, L. Farinelli, M. Pagni, M. Schneider, J. Voumard, M. Jaboyedoff, C. Fankhauser, C. S. Hardtke, L. Keller, J. R. Pannell, A. Reymond, M. Robinson-Rechavi, I. Xenarios, and P. Reymond. 2017. Low number of fixed somatic mutations in a long-lived oak tree. Nature Plants 3:926-929. (http://doi.org/10.1038/s41477-017-0066-9).

Schoen, D. J. 2005. Deleterious mutation in related species of the plant genus *Amsinckia* with contrasting mating systems. Evolution 59:2370-2377.

Schultz, S. T. and D. G. Scofield. 2009. Mutation accumulation in real branches: fitness assays for genomic deleterious mutation rate and effect in large-statured plants. Am. Nat. 174:163-175. (http://doi.org/10.1086/600100).

Schweinsberg, M., R. A. González Pech, R. Tollrian, and K. P. Lampert. 2014. Transfer of intracolonial genetic variability through gametes in Acropora hyacinthus corals. Coral Reefs 33:77-87. (http://doi.org/10.1007/s00338-013-1102-5).

Shaw, F. H., C. J. Geyer, and R. G. Shaw. 2002. A comprehensive model of mutations affecting fitness and inferences for *Arabidopsis thaliana*. Evolution 56:453-463.

Stearns, F. W. and C. B. Fenster. 2016. Fisher's geometric model predicts the effects of random mutations when tested in the wild. Evolution 70:495-501. (http://doi.org/10.1111/evo.12858).

Vezzulli, S., L. Leonardelli, U. Malossini, M. Stefanini, R. Velasco, and C. Moser. 2012. Pinot blanc and Pinot gris arose as independent somatic mutations of Pinot noir. J. Exp. Bot. 63:6359-6369. (http://doi.org/10.1093/jxb/ers290).

Watson, J. M., A. Platzer, A. Kazda, S. Akimcheva, S. Valuchova, V. Nizhynska, M. Nordborg, and K. Riha. 2016. Germline replications and somatic mutation accumulation are independent of vegetative life span in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America 113:12226-12231. (http://doi.org/10.1073/pnas.1609686113).

Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, and J. H. Willis. 2008. Mimulus is an emerging model system for the integration of ecological and genomic studies. Heredity 100:220-230. (http://doi.org/10.1038/sj.hdy.6801018).

Younginger, B. S., D. Sirová, M. B. Cruzan, and D. J. Ballhorn. 2017. Is biomass a reliable estimate of plant fitness? Applications in Plant Sciences 5:apps.1600094. (http://doi.org/10.3732/apps.1600094).

Yue, J.-X., J. Li, D. Wang, H. Araki, D. Tian, and S. Yang. 2010. Genome-wide investigation reveals high
evolutionary rates in annual model plants. BMC Plant Biol. 10:1-12.
(http://doi.org/10.1186/1471-2229-10-242).

Figure Captions

Fig. 1. Experimental design to test for the average fitness effects of somatic mutations accumulating in
stems during vegetative growth. See text for further explanation.

Fig. 2. An illustration of cell lineage selection as it acts on a number of cell lines that differ in growth rate
within the apical meristem of a plant. A) Expressed deleterious mutations are predicted to lead to the
demise of a cell line (brown cells) that is replaced by neighboring cell lines. The appearance of a
mutation that increases cell growth rate in the current environment results in a selective sweep of the
meristem (light green cells). B) Multiple ramets of the same genet become differentiated by unique
somatic mutations during vegetative growth. Most ramets carry deleterious mutations (orange, red,
purple, and pink arrows), but expressed beneficial mutations cause selective sweeps of the meristem in
some ramets (green arrows).

Fig. 3. Estimates of $\bar{s}_k$ for fourteen different stems (ramets) from two separate experiments (Experiment
1–blue bars; Experiment 2–orange bars) based on mean progeny fitness after autogamous $\bar{w}_{k(A)}$ and
geitonogamous $\bar{w}_{k(G)}$ self-pollinations ($\bar{s}_k = \bar{w}_{k(A)} - \bar{w}_{k(G)}$). Horizontal lines represent standard errors.
Asterisks indicate values of $\bar{s}_k$ that are significantly different from zero based on the *t* value, calculated
as $t = \bar{s}_k/\sqrt{SE}$ with *n-1* df, where *n* is the mean of sample sizes for progeny from autogamy and
geitonogamy. The relationship between $\bar{w}_{k(A)}$ and $\bar{w}_{k(G)}$ across stems is shown in Fig. S2. Means and
sample sizes for progeny groups are available in Table S6 and Fig. S3.

Fig. 4. Relationships between estimates of selection coefficients, based either on the difference in
fitness of progeny from autogamy and geitonogamy ($\bar{s}_k$), or the standard deviation in fitness within
progeny groups from autogamy for each stem (s_{SD}). Estimates of s_{SD} corresponding to negative values of
$\bar{s}_k$ were transformed to negative values. Estimates from Experiment 1 are indicated by blue circles and
from Experiment 2 are orange squares. Dashed lines indicate the separate relationships for positive and
negative values of the selection coefficients.

Fig. 1

Fig. 2

Cell Generations

A. Cell Lineage Selection

B. Differentiation among Ramets

Fig. 3

Average Selective Value ($\bar{S}_k$)

Fig. 4

811
812

Appendix 1. Validity of the $\bar{s}_k$ Estimate

For a pair of stems from the same genet, we assume that w_G represents the fitness of geitonogamous
progeny after a cross between the pair of stems, and that w_A represents the fitness of autogamous
progeny after selfing a single flower on one stem:

$w_G = 1 + 0.5(h_1s_1 + h_2s_2)$

$w_A = 0.25(1) + 0.5(1 + h_1s_1) + 0.25(1 + s_1)$

Where, s_1 and s_2 represent the selective effects of mutations, and h_1 and h_2 represent the dominance of
single somatic mutations in stem 1 and stem 2, respectively.

For simulations, we randomly chose values of s ranging from -0.2 to 0.2 for 200 pairs of stems. Values of
h for $s > 0$ were allowed to range from 0 to 0.7 for each stem. Because deleterious alleles tend to be
recessive, values of h were constrained to range between 0 and 0.1 for $s < 0$. To match our experimental
design in *Mimulus guttatus*, we calculated the fitness of autogamous progeny (w_A) for one of the stems
in each pair. We then calculated w_G using information on s and h for both stems in each pair. Then we
calculated the difference in fitness between each estimate as a measure of autogamy depression ($s_k =$
$w_A - w_G$). This simulation was repeated 20 times (4,000 pairs of stems total) and average values were
calculated.

Results:

1. Estimates of s_k and s_1 are strongly correlated, with R^2 values ranging from 0.76 to 0.82 (one
831   example of 200 random estimates):

834   2. The frequency of s_k estimates having the opposite sign of s_1 was very low. Mean number of
835   correct signs was 178.7 (+/- 1.04 standard error) and was 21.30 (+/-1.04; or 12%) for incorrect
836   signs.

837   3.   Estimates of s_k with incorrect signs were closer to zero. Mean absolute values of s for s_k
838   estimates with incorrect signs was 0.054 (+/- 0.0021) and was 0.106 (+/-0.0011) for correct
839   signs. Hence, estimates of s_k that are further from zero are more reliable estimates of s_1.
840
We note that the formulation used above for w_G could result in estimates that are out of range when
large numbers of mutations are involved (i.e., $w_G = 1 + 0.5 \sum h_i \, s_i$ for a large number of loci segregating
for deleterious alleles). A better estimate may be the average effect of n loci segregating for deleterious
alleles (i.e. $\sum h_i \, s_i \, /n$). If we apply this approach for two loci (one mutation per stem) we obtain,

$w_G = 0.25 + 0.25(1 + h_1 s_1) + 0.25(1 + h_2 s_2) + 0.25(1 + (h_1 s_1 + h_2 s_2)/2)$

$w_G = 1 + 0.375(h_1 s_1 + h_2 s_2)$

Using this formula, the relationship between s_k and s_1 is stronger (R^2 ranging from 0.86 to 0.90), with
only an average of 13 out of 200 estimates with opposite signs (7%). The average of correct signs was
similar to the one above (0.105, +/- 0.0007), but the mean for incorrectly assigned values was much
closer to zero (0.038, +/- 0.0022).

Another model for the effects of alleles at multiple loci on fitness assumes the interaction is
multiplicative. In this case the estimate of w_G becomes,

$w_G = 0.25 + 0.25(1 + h_1 s_1) + 0.25(1 + h_2 s_2) + 0.25(1 + (h_1 s_1 h_2 s_2))$

$w_G = 1 + 0.25(h_1 s_1 + h_2 s_2) + 0.25(h_1 s_1 h_2 s_2)$

Using this formula, the relationship between s_k and s_1 is stronger (R^2 ranging from 0.93 to 0.95), with
only an average of 7.1 out of 200 estimates with opposite signs (4%). The average of correct signs was
similar to the ones above (0.103, +/- 0.0010), but the mean for incorrectly assigned values was closer to
zero (0.023, +/- 0.0020).

Interpretation:

These results indicate that our estimate of s_k is a valid estimate of s for stem 1. However, it does appear
that we are underestimating s (i.e., the slope of the line is less than 1), as these estimates do not
consider the effects of multiple somatic mutations or possible genetic background effects (i.e. epistasis).

It is also notable that our simulation assumes a wider range of h values than are generally observed for
deleterious mutations. Estimates indicate that the product of hs for deleterious alleles is generally
around 0.02, and that the relationship between s and h is generally negative (hyperbolic). Minor effects
of deleterious mutations are expected for our data, because somatic mutations having strong effects in
the heterozygous condition would be eliminated by cell lineage selection. Adding the constraint of a
negative relationship between h and s to our simulation model would marginally improve the predictive
power of s_k for estimates of s.

These simulations also support our use of the correlation between w_A and w_G to confirm that our
estimates are similar to the simulation estimates.

From the simulations:

From our data:

Appendix 2. Estimating the selection coefficient due to somatic mutations from the variance in fitness
among progeny.
882
883 In this Appendix, we demonstrate that the standard deviation (SD) in fitness among progeny from
884 autogamous crosses can be used to estimate the selection coefficient for the average effects of somatic
885 mutations occurring within a stem. The variance in fitness among progeny from autogamy can be
886 affected by the selection coefficient (s) and the dominance effects (h) of somatic mutations unique to a
887 stem.
888
889 To evaluate the effects of a single mutation on progeny fitness, we assume fitnesses of;
890 $w_{aa} = 1$
891 $w_{aa'} = 1 + hs$
892 $w_{a'a'} = 1 + s$
893
894 The mean progeny fitness is then calculated as
895
896 $\overline{w} = \frac{1}{4}(1) + \frac{1}{2}(1+hs) + \frac{1}{4}(1+s)$
897
898 Which simplifies to
899
900 $\overline{w} = (1 + \frac{1}{2} hs + \frac{1}{4} s)$
901
902 The variance in progeny fitness can be calculated as
903
904 $V(w) = \frac{1}{4}(w_{aa} - \overline{w})^2 + \frac{1}{2}(w_{aa'} - \overline{w})^2 + \frac{1}{4}(w_{a'a'} - \overline{w})^2$
905
906 After substituting, we get
907
908 $V(w) = [\frac{1}{4} (- \frac{1}{2} hs - \frac{1}{4} s)^2] + [\frac{1}{2} (\frac{1}{2} hs - \frac{1}{4} s)^2] + [\frac{1}{4} (\frac{3}{4} s - \frac{1}{2} hs)^2]$
909
910 For simplification, we separate the effects of homozygous and heterozygous genotypes (terms within
911 the three sets of square brackets)
912
913 $V(w) = [1] + [2] + [3]$
914
915 and evaluate each one;
916

917 [1]: $\frac{1}{4} (- \frac{1}{2} hs - \frac{1}{4} s)^2 = \frac{1}{4} (- \frac{1}{2} hs (- \frac{1}{2} hs - \frac{1}{4} s)- \frac{1}{4} s(- \frac{1}{2} hs - \frac{1}{4} s)) = (\frac{1}{16} h^2s^2 + \frac{1}{16} hs^2 + \frac{1}{64} s^2)$
918

919 [2]: $\frac{1}{2} (\frac{1}{2} hs - \frac{1}{4} s)^2 = \frac{1}{2} (\frac{1}{2} hs (\frac{1}{2} hs - \frac{1}{4} s) - \frac{1}{4} s(\frac{1}{2} hs - \frac{1}{4} s)) = \frac{1}{2} (\frac{1}{4} h^2s^2 - \frac{1}{8} hs^2 - \frac{1}{8} hs^2 + \frac{1}{16} s^2) = ($
920 $\frac{1}{8} h^2s^2 - \frac{1}{8} hs^2 + \frac{1}{32} s^2)$
921

922 [3]: $\frac{1}{4} (\frac{3}{4} s - \frac{1}{2} hs)^2 = \frac{1}{4} (\frac{3}{4} s (\frac{3}{4} s - \frac{1}{2} hs) - \frac{1}{2} hs (\frac{3}{4} s - \frac{1}{2} hs)) = \frac{1}{4} (\frac{9}{16} s^2 - \frac{3}{8} hs^2 - \frac{3}{8} hs^2 + \frac{1}{4} h^2s^2) =$
923 $(\frac{9}{64} s^2 - \frac{3}{16} hs^2 + \frac{1}{16} h^2s^2)$
924
By combining and then simplifying these three results, we get

$V(w) = 3/16\ s^2 - \frac{1}{4}\ hs^2 + \frac{1}{4}\ h^2 s^2$

And the standard deviation can be obtained by taking

the square root of V(w)

$SD(w) = \sqrt{V(w)} = 0.433\ s - \frac{1}{2}\ s\sqrt{h} + \frac{1}{2}\ hs$

The standard deviation in progeny fitness scales

linearly with the selective effects of somatic

mutations, and the effects of dominance on variation

among progeny is minor (Fig. S1; which shows values

of *s* predicted from values of SD for fitness ranging

from 0 to 0.5). We note that the standard deviation is

Fig. S1. The effects of selection (*s*) and dominance (*h*) on the standard deviation (SD) in fitness among progeny. The effect of dominance on variation in fitness is minimized when *h = 0.5* and maximized when *h = 0.0* and *h = 1.0*.

highest when mutations are completely recessive or dominant (h = 0 or 1) and SD(w) = 0.433s. When h =

0.5, the standard deviation is at its minimal value and SD(w) = 0.332s. By rearranging, we can obtain the

regression equations to estimate the selective effects of somatic mutations from the standard deviation

in progeny fitness as $s_{SD} = cSD$, where *c* is the slope of the regression line in Fig. S1. From this analysis, it

is clear that the effects of *h* on the within-family SD in fitness is minor compared to the effects of *s*. We

obtain estimates of the regression slopes from the standard deviation in progeny fitness to produce the

relationships in Fig. S1;

s = 2.31*SD(w) for *h* = 0 or 1, and

s = 3.01*SD(w) for *h* = 0.5

From these relationships we can estimate the selective effects of somatic mutations from the standard

deviation in progeny fitness as $s_{SD} = cSD$, where *c* is the slope of the regression lines in Fig. S1.

Appendix 3. Supplemental Tables and Figures

Table S1.

Stem	Autogamy $\overline{w}_A$	N	Stdev	Geitonogamy $\overline{w}_G$	N	Stdev	$\overline{s}_k$	$\overline{w}_A - \overline{\overline{w}}_G$
JB51	0.407	33	0.171	0.144	7	0.080	0.263	0.078
JB52	0.314	12	0.161	0.295	9	0.068	0.019	-0.015
JB54	0.258	8	0.066	0.298	11	0.103	-0.040	-0.071
SMB11	0.286	22	0.072	0.260	9	0.077	0.009	-0.043
SMB51	0.256	11	0.054	0.323	4	0.035	-0.067	-0.073
SMB52	0.273	7	0.072	0.280	6	0.051	-0.007	-0.056
SMB53	0.245	4	0.021	0.327	6	0.108	-0.081	-0.083
SMB54	0.289	10	0.023	0.245	5	0.056	0.044	-0.040
SMC11	0.303	16	0.071	0.238	8	0.085	0.065	-0.026
SMC12	0.356	8	0.071	0.239	6	0.059	0.117	0.027
C12	0.423	36	0.170	0.707	5	0.153	-0.283	0.094
C13	0.626	9	0.195	0.322	7	0.237	0.090	0.297
D17	0.692	5	0.170	0.485	4	0.314	0.207	0.363
D18	0.419	12	0.130	0.443	13	0.211	-0.025	0.090

Table S2. Analysis of variance results for seed set after autogamous and geitonogamous pollinations (Cross) with limited and excess pollen (Pollen). Population (Pop), maternal stem nested within population (Mat(Pop)), and pollination date (Date) were declared as random effects in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	21623.26022	655.25031	1.43	0.0888
Cross	1	100.647606	100.647606	0.22	0.6400
Pop	2	552.228601	276.114301	0.60	0.5487
Pollen	1	2183.436724	2183.436724	4.78	0.0312
Mat(Pop)	18	6490.364191	360.575788	0.79	0.7089
Date	11	8000.908005	727.355273	1.59	0.1127
Error	101	46182.07311	457.24825		

968 Table S3. Analysis of variance results for aborted ovules after autogamous and geitonogamous
969 pollinations (Cross) with limited and excess pollen (Pollen). Population (Pop), maternal stem nested
970 within population (Mat(Pop)), and pollination date (Date) were declared as random effects in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	9337.61468	282.95802	2.76	<.0001
Cross	1	449.176948	449.176948	4.38	0.0389
Pop	2	2174.689160	1087.344580	10.60	<.0001
Pollen	1	0.597410	0.597410	0.01	0.9393
Mat(Pop)	18	2217.350757	123.186153	1.20	0.2751
Date	11	1995.681210	181.425565	1.77	0.0695
Error	101	10363.24458	102.60638		

971
972 Table S4. Analysis of variance for progeny fitness (biomass weighted by survival of seedlings) after
973 autogamous and geitonogamous pollinations (Cross) for ten maternal stems (Mat) and their interaction
974 (Mat*Cross). Planting block (Tray) was declared as a random effect in the model.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	33	1.42418594	0.04315715	4.18	<.0001
Tray	14	0.45040769	0.03217198	3.11	0.0002
Mat	9	0.06845581	0.00760620	0.74	0.6756
Cross	1	0.03196532	0.03196532	3.09	0.0805
Mat*Cross	9	0.51373985	0.05708221	5.52	<.0001
Error	168	1.73638043	0.01033560		

975
976
977 Table S5. Analysis of variance for progeny fitness (biomass weighted by survival of seedlings) after
978 autogamous and geitonogamous pollinations (Cross) for seven paired pollinations at individual nodes
979 (Stem/Node) and their interaction (Stem/Node*Cross). Hydroponic planting block (Tray) was declared as
980 a random effect in the model and initial size of seedlings (Initial Size) was entered as a covariate.

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	23	1.31384069	0.05712351	6.05	<.0001
Tray	11	0.46856600	0.04259691	4.51	<.0001
Initial Size	1	0.46861414	0.46861414	49.61	<.0001
Stem/Node	6	0.12071536	0.02011923	2.13	0.0514
Cross	1	0.00555294	0.00555294	0.59	0.4441
Stem/Node*Cross	4	0.13005812	0.03251453	3.44	0.0095
Error	203	1.91760656	0.00944634		

  Table S6. Means and standard deviations (SD) for progeny groups from autogamous and geitonogamous
  pollinations in experiments 1 and 2. Experiment 1 included ten different genets. Experiment 2 was
  conducted with two ramets (C1 and D1) from the same genet. The fitness of progeny was estimated as
  its final biomass (log transformed to improve normality), weighted by the survival frequency of progeny
  from the same cross.

Exp	ID	N	Fitness Autogamy	SD	N	Fitness Geitonogamy	SD
2	C13	9	0.626	0.195	7	0.322	0.237
1	JB51	33	0.316	0.193	7	0.114	0.041
2	D17	7	0.662	0.170	60	0.465	0.314
1	SMC12	8	0.222	0.080	6	0.103	0.046
1	SMC11	16	0.175	0.065	8	0.099	0.071
1	JB52	12	0.195	0.158	9	0.162	0.071
1	SMB11	22	0.155	0.082	9	0.152	0.096
1	SMB54	10	0.128	0.020	5	0.129	0.072
2	D18	12	0.419	0.130	13	0.443	0.211
1	JB54	8	0.118	0.063	11	0.17	0.109
1	SMB52	7	0.117	0.069	6	0.173	0.057
1	SMB51	11	0.11	0.040	4	0.171	0.048
1	SMB53	4	0.069	0.021	6	0.21	0.108
2	C12	36	0.423	0.188	5	0.706	0.153
		13.93			11.14		

987
988

989
990
991

992
993
994   Fig. S2. The relationship between the autogamous and geitonogamous progeny from the 14
995   comparisons from experiments 1 (blue circles) and 2 (orange squares) as listed in Table S6. The solid line
996   represents a 1:1 relationship and the dashed line represents the best-fit to the data. Stems falling above
997   the solid line apparently had a prevalence of deleterious somatic mutations whereas those falling below
998   the line displayed a prevalence of beneficial somatic mutations.
999
1000
1001

1002

1003

1004

1005

1006 Fig S3. Means and standard errors (lines) for fitness estimates of autogamous (upper panel) and
1007 geitonogamous (lower panel) progeny groups from experiment 1 (blue bars) and 2 (orange bars) listed in
1008 Table S6. Progeny groups are ordered as in Fig. 4.

1009

1010

1011