Regional Striatal Cholinergic Involvement in Human Behavioural Flexibility

Role of human striatal choline in reversal learning

Tiffany Bell[1,2], Michael Lindner[1], Angela Langdon[3], Paul Gerald Mullins[4], Anastasia Christakou[1]

1. School of Psychology and Clinical Language Sciences, and Centre for Integrative Neuroscience and Neurodynamics, University of Reading, RG6 6AL, UK
2. Department of Radiology, Cumming School of Medicine, University of Calgary, AB, T2N 1N4 Canada
3. Princeton Neuroscience Institute, Princeton University, NJ 08544,USA
4. School of Psychology, Bangor University, LL57 2DG UK

Corresponding author: Dr Anastasia Christakou
                         School of Psychology and Clinical Language Sciences
                         University of Reading
                         Reading RG6 6AL
                         anastasia.christakou@reading.ac.uk

Number of Figures: 5

Number of Tables: 6

Number of Words: Abstract   243

                    Introduction   588

                    Discussion   1440

Conflict of Interest   The authors declare no competing interests.

Acknowledgments: This study was supported by a Human Frontier Science Program (HFSP) grant (RGP0048/2012), and an Engineering and Physical Sciences Research Council (EPSRC) doctoral training grant (EP/L505043/1). The funders had no involvement in study design, in the collection, analysis, and interpretation of data, in writing the report, or in the decision to submit the paper for publication. The authors would like to thank Rosie Gillespie and Emma Davis for assistance with data collection.

ABSTRACT

Animal studies have shown that the striatal cholinergic system plays a role in behavioural flexibility but, until recently, this system could not be studied in humans due to a lack of appropriate non-invasive techniques. Using proton magnetic resonance spectroscopy (^{1}H-MRS) we recently showed that the concentration of dorsal striatal choline (an acetylcholine precursor) changes during reversal learning (a measure of behavioural flexibility) in humans. The aim of the present study was to examine whether regional average striatal choline was associated with reversal learning. We measured choline at rest in both the dorsal and ventral striatum using ^{1}H-MRS and examined its relationship with performance on a probabilistic learning task with a reversal component. Task performance was described using a simple reinforcement learning model that dissociates the contributions of positive and negative prediction errors to learning. Average levels of choline in the dorsal striatum were associated with performance during reversal, but not during initial learning. Specifically, lower levels of choline in the dorsal striatum were associated with a lower number of perseverative trials. Moreover, choline levels explained inter-individual variance in perseveration over and above that explained by learning from negative prediction errors. These findings suggest that the dorsal striatal cholinergic system plays an important role in behavioural flexibility, in line with evidence from the animal literature and our previous work in humans. Additionally, this work provides further support for the idea of measuring choline with ^{1}H-MRS as a non-invasive way of studying human cholinergic neurochemistry.

SIGNIFICANCE STATEMENT

Behavioural flexibility is a crucial component of adaptation and survival, as well as everyday living. Evidence from the animal literature shows the striatal cholinergic system is fundamental to reversal learning, a key paradigm for studying behavioural flexibility. However, this system remains understudied in humans, largely due to a lack of non-invasive techniques. Using proton magnetic resonance spectroscopy (^{1}H-MRS), we showed that choline levels at rest in the dorsal striatum are associated with individual differences in performance specifically during reversal learning. These novel findings help to bridge the gap between animal studies of the striatal cholinergic system and human studies of basal ganglia function, by demonstrating the importance of cholinergic function in the dorsal striatum in human behavioural flexibility. Importantly, the methods described here can not only be applied to furthering our understanding of healthy human neurochemistry, but can also be applied to extending our understanding of cholinergic disorders.

INTRODUCTION

Acetylcholine (ACh) plays an important role in adaptive behaviour, and has been implicated in disorders of cognitive flexibility, such as Parkinson's disease (Tanimura et al., 2018; Zucca, Zucca, & Wickens, 2018). Studies in rodents have repeatedly demonstrated that ACh transmission, determined by the activity and regulation of cholinergic interneurons in the dorsal striatum (DS), is involved in reversal learning and similar forms of behavioural flexibility (Aoki et al., 2018; Bradfield, Bertran-Gonzalez, Chieng, & Balleine, 2013; H. D. Brown, Baker, & Ragozzino, 2010; McCool, Patel, Talati, & Ragozzino, 2008; Okada, Nishizawa, Setogawa, Hashimoto, & Kobayashi, 2018; Ragozzino, Jih, & Tzavos, 2002; Ragozzino, Mohler, Prior, Palencia, & Rozman, 2009; Tzavos, Jih, & Ragozzino, 2004). Further, ACh efflux has been shown to increase specifically during reversal learning (but not during initial learning), and this effect is specific to the dorsomedial striatum (with no changes in ACh levels in either the dorsolateral striatum or the ventral striatum) (Ragozzino et al., 2009). It is clear then that cholinergic activity in the DS plays an important role in reversal learning but, despite the importance of understanding this system, there remain important challenges in probing ACh function in humans due to a lack of appropriate non-invasive techniques.

Proton magnetic resonance spectroscopy ([1]H-MRS) is a non-invasive method for measuring brain metabolites *in vivo* (Puts & Edden, 2012). Although it cannot be used to study ACh directly due to its low concentration (Hoover, Muth, & Jacobowitz, 1978), [1]H-MRS can be used to measure levels of certain choline containing compounds (CCCs) involved in the ACh cycle, including choline (CHO). CHO is the product of ACh hydrolysis, and its uptake in cholinergic terminals is the rate-limiting step in ACh biosynthesis (Lockman & Allen, 2002). Using functional [1]H-MRS we previously demonstrated task-driven changes in the concentration of CHO in the human DS during reversal learning (Bell, Lindner, Mullins, & Christakou, 2018). Although [1]H-MRS studies typically model CCCs as a single peak due to their proximity on the spectrum, we showed that using this

method may mask CHO-specific effects. Therefore, in the context of studying ACh function, it is

necessary to separate the metabolites when measuring individual differences in CHO levels (Bell et

al., 2018; Lindner, Bell, Iqbal, Mullins, & Christakou, 2017).

Among the many open questions around this approach is the nature of the relationship between

baseline levels of CHO availability and function-relevant ACh activity. Animal studies have shown

that ACh synthesis is tightly coupled to CHO availability. For example, depletion of CHO has been

shown to reduce ACh synthesis (Jope, 1979) and administration of CHO has been shown to increase

it (Koshimura et al., 1990). Further, overexpression (Holmstrand et al., 2013) and under-expression

(Parikh, St. Peters, Blakely, & Sarter, 2013) of presynaptic CHO up-take transporters has been

shown to increase and decrease ACh levels respectively. It is possible, therefore, that baseline CHO

availability may modulate ACh activity, leading to effects on behavioural flexibility. In this study,

we used ^{1}H-MRS to test whether baseline levels of regional striatal CHO are related to individual

differences in performance during a probabilistic reversal learning task. To do this, we obtained

average measures of CHO from the dorsal and ventral regions of the striatum (DS and VS,

respectively). Additionally, CHO levels from the cerebellum were used as a control to demonstrate

specificity. In line with the animal literature and our previous findings in humans (Bell et al., 2018),

we predicted that average levels of CHO in the dorsal, but not the ventral, striatum would be

associated with performance during reversal, but not initial, learning.

METHODS

Participants

The study was approved by the University of Reading Research Ethics Committee. 36 volunteers (20 female) between the ages of 18.3 and 32.8 (mean = 24.8, SD = 3.5) were recruited by opportunity sampling. All participants were healthy, right handed non-smokers and written informed consent was taken prior to participation. Two participants were excluded from analyses due to a high proportion of missed responses (participant 14: 35% during initial learning and 39% during reversal learning; participant 31: 27% during initial learning, 54% during reversal learning).

Behavioural Data

Learning Task

The task used was a probabilistic multi-alternative learning task previously described (Bell et al., 2018), and was programmed using MATLAB (2014a, The Mathworks, Inc., Natick, MA, United States) and Psychtoolbox (Brainard, 1997).

First, participants were presented with a fixation cross displayed in the centre of the visual display. Participants were then presented with four decks of cards. Each deck contained a mixture of winning and losing cards, corresponding respectively to a gain or loss of 50 points. The probability of getting a winning card differed for each deck (75%, 60%, 40%, and 25%) and the probabilities were randomly assigned across the four decks for each participant. Participants indicated their choice of deck using a computer keyboard. Outcomes were pseudo-randomised so that the assigned probability was true over every 20 times that deck was selected. Additionally, no more than 4 cards of the same result (win/loose) were presented consecutively in the 75% and 25% decks and no more than 3 cards of the same result in the 60% and 40% decks. A cumulative points total was displayed

in the bottom right-hand corner throughout the session and in the centre of the visual display at the

end of each trial (Figure 1). Participants were instructed that some decks may be better than others,

they are free to switch between decks as often as they wish, and they should aim to win as many

points as possible.

The learning criterion was set at selection of either of the two highest decks (60% or 75%) on at

least 80% of the time over ten consecutive trials. Though the optimal strategy is to repeatedly

choose the 75% deck, pilot testing revealed the participants were not always able to distinguish

between the 75% and 60% decks. Therefore, as both decks generate an overall gain in points and

choice of either deck could be considered a good strategy, both decks are included in the learning

criterion.

The initial learning phase (round 1) was completed when either the learning criterion was reached,

or the participant completed 100 trials. The deck probabilities were then reversed so that the high

probability decks became low probability and vice versa. Participants were not informed of the

reversal. The task ended either after the learning criterion was reached following the reversal (round

2), or after another 100 trials (Figure 2).

Figure 1: General outline of learning task trials. Participants were instructed to choose between four decks of

cards. Each deck had a different probability of generating wins:losses (75:25, 60:40, 40:60, 25:75). Once the

learning criterion had been reached, the deck probabilities were reversed so that high probability decks became low probability decks and vice versa. Participants were not informed of this in advance and were simply instructed to gain as many points as possible. Each screen was shown for 2.5s, RT = reaction time.

Impulsivity

Previous research has shown that trait levels of impulsivity can influence decision making (Bayard, Raffard, & Gely-Nargeot, 2011). Individuals with higher levels of impulsivity have been shown to demonstrate sub-optimal performance on decision making tasks, displaying a decreased ability to learn reward and punishment associations and implement these to make appropriate decisions. For instance, individuals with high levels of impulsivity were relatively impaired in adapting their behaviour during a reversal learning task (Franken, van Strien, Nijs, & Muris, 2008). Other tasks of cognitive flexibility have also been shown to be influenced by trait impulsivity levels (e.g. Müller, Langner, Cieslik, Rottschy, & Eickhoff, 2014). Therefore all participants completed the Barratt Impulsiveness Scale (BIS-11; Patton, Stanford, & Barratt, 1995) and their total score was used as a trait measure of impulsivity. This was included in the analysis to account for effects driven by individual differences in impulsivity.

Data Analysis

Participants were split into two groups based on performance. Those who learnt both rounds (i.e. reached criterion both during initial learning and after reversal) were classified as learners and those who did not learn both rounds were classified as non-learners.

Behaviour was analysed for learners only. Those who did not reach criterion during round 1 will not have realised there was a change in contingencies and will not have experienced a reversal, therefore their behaviour during the reversal stage is likely to be different to those who did experience the reversal. Additionally, because the task stops at 100 trials per round if the criterion is not met, there is a ceiling effect for those who did not reach criterion. Consequently, there will be a ceiling effect for those who did not reach criterion in both rounds, and for those who did not reach

criterion in either round 1 or round 2 only. Therefore, participants who did not reach criteria in

either one round or both rounds were excluded from behaviour analysis.

Performance was measured using the number of trials taken to reach criterion in round 1 (initial

learning) and in round 2 (reversal learning). Round 2 was subdivided into perseverative trials and

post-reversal learning (Figure 2). The number of perseverative trials was defined as the number of

trials after reversal until the probability of selecting the previously favoured deck reached chance

level (0.25), i.e. the number of trials taken to identify the reversal and switch behaviour. Post-

reversal learning was defined as the number of trials taken to reach criterion in round 2, minus the

number of perseverative trials, i.e. the number of trials to reach criterion after the reversal had been

detected. In other words, post-reversal leaning is measured by the number of trials the participant

took to learn the contingencies once they had realised the deck probabilities had reversed.

Additionally, the post-reversal learning period included a measure of regressive errors. The number

of regressive errors was defined as the number of times the previously favoured deck was selected

during the post-reversal learning period (i.e. after the perseverative period had ended).

Figure 2: General overview of learning task structure. Participants completed the initial learning phase
(round 1) by reaching the predefined accuracy criterion or after 100 trials. Upon completion of the initial
learning phase, the deck probabilities were reversed. Participants then completed a reversal learning phase
(round 2). For behavioural analysis, this was subdivided into perseverative trials (PER) and a post-reversal
learning period. The number of perseverative trials was defined as the number of trials after reversal until the
probability of selecting the previously favoured card reached chance level (0.25). The post-reversal learning
period was the number of trials to reach criterion in round 2, minus the number of perseverative trials. The
number of regressive errors was defined as the number of times the previously favoured deck was selected

during the post-reversal learning period. The task ended once participants either reached the same accuracy
criterion in round 2 or after 100 round 2 trials.

Temporal Difference Reinforcement Learning Model

196 We modelled participants' choice behaviour as a function of their previous choices and rewards

197 using a temporal difference reinforcement learning algorithm (Sutton & Barto, 1998). This allows

198 us to track trial-and-error learning for each participant, during each task stage, in terms of a

199 subjective expected value for each deck. On each trial t, the probability that deck c was chosen was

200 given by a soft-max probability distribution,

$$P(c_t = c) = e^{m_t(c)} \Big/ \sum_j e^{m_t(j)} \tag{1}$$

201 where $m_t(c)$ is the preference for the chosen deck and j indexes the four possible decks. The

202 preference for the chosen deck was comprised of the participant's expected value of that deck on

203 that trial, $V_t(c)$, multiplied by the participant's individual value impact parameter β (equivalent to

204 the inverse temperature):

$$m_t(c) = \beta V_t(c). \tag{2}$$

The parameter β describes the extent to which trial-by-trial choices follow the distribution of the

expected values of the decks: a low β indicates choices are not strongly modulated by expected

value, being effectively random with respect to this quantity (i.e. participants are not choosing

based exclusively on value, and are effectively exploring all options); conversely, a high β indicates

choices largely follow expected value (i.e. participants choose the deck with the highest expected

value; exploitation).

To update the subjective value of each deck, a prediction error was generated on each trial, pe_t

based on whether participants experienced a reward or a loss (*reward$_t$*, = +1 or -1 respectively). The

expected value of the chosen deck was subtracted from the actual trial reward to give the prediction

error:

$$pe_t = reward_t - V_t(c) \qquad\qquad (3)$$

Studies have shown that individuals differ in the degree to which they learn from better than

expected outcomes (positive prediction errors) and worse than expected outcomes (negative

prediction errors) (Bull, Tippett, & Addis, 2015; Christakou et al., 2013; Gray, 1970; Niv, Edlund,

Dayan, & O'Doherty, 2012). To account for this, two learning rate parameters were used to model

sensitivity to prediction errors in updating the expected values: the weight of learning from better

than expected outcomes (learning rate from positive prediction errors: η^+) and the weight of

learning from worse than expected outcomes (learning rate from negative prediction errors: η^-). For

example, individuals who are reward seeking will place a high weight on the former, whereas those

who are loss-aversive will place a high weight on the latter. The prediction error on each trial was

multiplied by either the positive (η^+) or negative (η^-) learning rate and used to update the value of

the chosen deck.

$$\delta_t = \eta^+ \times pe_t \quad if\ pe_t > 0 \qquad\qquad (4)$$

$$\delta_t = \eta^- \times pe_t \quad if\ pe_t < 0 \qquad\qquad (5)$$

$$V(chosen_t) = V(chosen_{t-1}) + \delta_t \qquad\qquad (6)$$

226　Thus, the model has three parameters of interest (β, η^+ and η^-). In psychological terms, β captures

227　the degree to which the subjective value of the chosen deck influenced decisions, while the learning

228　rates capture the individual's preference for learning from positive (η^+) or negative (η^-) prediction

229　errors to guide choice behaviour during this task.

Model Fitting

The model was fit per participant to provide parameters that maximised the likelihood of the

observed choices given the model (individual maximum likelihood fit; Daw, 2011). The reward

value was updated as 1 (win) or -1 (loss). Subjective value was initialised at zero for all decks and

the initial parameter values were randomised. To ensure the model produced consistent,

interpretable parameter estimates, η was limited to between 0 and 1 and β was assumed positive.

The parameters were constrained by the following distributions based on Christakou et al (2013):

$$\beta \sim Gamma\ (2,1)$$

$$\eta \sim Beta\ (1.2, 1.2)$$

The model was fit separately over the trials encompassing round 1 (R1, initial learning) and round 2

(R2, perseverative trials and post-reversal learning, denoted as reversal learning). This was done to

capture the change in influence of the model parameters from initial learning to reversal learning.

The model was not fit over the perseverative trials separately as the average number of

perseverative trials was too small to generate a stable model fit.

Traditionally, to investigate the fit of a temporal difference reinforcement learning model the

Bayesian information criterion (BIC) is used. The BIC is a post hoc fit criterion which looks at the

adequacy of a model whilst penalising the number of parameters used. A lower number indicates a

better fit (Steingroever, Wetzels, & Wagenmakers, 2016). However, the BIC is generally used to

compare different models, rather than model fits over different sets of data, and will penalise

different sized data sets. Alternatively, the corrected likelihood per trial (CLPT) can be used. The

CLPT is a more intuitive measure of fit that takes into account the number of trials completed

without penalising different sized data sets. The CLPT varies between 0 and 1, with higher values

indicating a better fit (Leong & Niv, 2013; Niv et al., 2015).

Wilcoxon signed-rank tests showed there was no significant difference between the CLPT values

for the model fit over round 1 (Mdn = 0.23) and round 2 (Mdn = 0.23; Z= -1.308, p = 0.191).

Additionally, there was no significant difference between the BIC values for the model fit over

round 1 (M = 75.7, SD = 45.5) and round 2 (M = 90.9, SD = 43.6; t(33) = -1.533, p = 0.135, r =

0.26).

To summarise, the model fit equally well across rounds. Therefore, differences in parameter

estimates across the task can be examined.

Magnetic Resonance Spectroscopy

Data Acquisition

Data was collected at the University of Reading on a Siemens Trio 3T MRI scanner using a transmit-receive head coil. A high-resolution whole-brain T1 structural image was acquired for voxel placement using an MPRAGE sequence parallel to the anterior-posterior commissure line (176 x 1mm slices; TR = 2020ms; TE = 2.9ms; FOV = 250mm).

Voxels were placed in either the left or right dorsal striatum (DS), ventral striatum (VS) and the cerebellum, with hemisphere placement and order of measurements counterbalanced across participants. Anatomy was used to guide voxel positioning. The top of the DS was identified by slice-by-slice examination of the structural scan. The slice below the slice where the top of the striatum was no longer visible was selected and the top of the voxel was aligned with this slice. The same technique was applied for the VS voxel. The slice above the slice where the bottom of the striatum could no longer be seen was selected and used for alignment of the VS voxel. The cerebellum voxel was placed as high in the superior cerebellar vermis as possible whilst ensuring only cerebellar tissue was contained in the voxel. The superior cerebellar vermis was chosen as it has been shown to have the lowest variability in both inter and intra subject metabolite ratios as measured with [1]H-MRS at rest (Currie, Hadjivassiliou, Wilkinson, Griffiths, & Hoggard, 2013). All voxels were visually inspected to ensure minimal cerebrospinal fluid was included in the voxels.

A PRESS sequence was used to acquire data from the three separate voxel positions (voxel size = 10mm x 15mm x 15mm; TR = 2000ms; TE = 30ms). 128 spectra were collected and averaged for each area. A water-unsuppressed spectrum was also obtained from each area for data processing, which consisted of an average of 15 spectra. Spectra were obtained for all participants from the cerebellum and the DS. The spectrum obtained from the VS was only usable from 34 participants due to noise levels. The SIEMENS Auto Align Scout was used in between each scan to adjust the

voxel position based on the actual head position of the participant. This was used to correct for

participant motion and minimize the variability of the voxel position.

Figure 3: Location of voxels and example spectra. Heat maps showing the sum of the [1]H-MRS voxels over

all subjects in MNI space, along with a representative spectrum from a single subject (A = Dorsal Striatum,

MNI coordinates: -3.41, 2.37, 11.16; B = Ventral Striatum, MNI coordinates: -2.99, 5.92, -3.93; C =

Cerebellum, MNI coordinates: -2.10, -61.03, 19.20).

Structural Segmentation

Structural scans were processed using FSL version 5.0.8 (Jenkinson, Beckmann, Behrens,

Woolrich, & Smith, 2012; Smith et al., 2004). First, the skull was removed using the brain

extraction tool (BET) (Smith, 2002). Images were segmented into three separate tissue types: grey

matter (GM), white matter (WM) and cerebrospinal fluid (CSF) using the FAST tool (Zhang,

Brady, & Smith, 2001). The coordinates and dimensions of the voxel were then superimposed on

these images and the proportion of each of the three tissue types contained within the voxel was

calculated.

Quantitation

Data was processed in the time domain using Java-Based Magnetic Resonance User Interface

(jMRUI software version 5.0 (http://www.mrui.uab.es/mrui); Naressi et al., 2001). Phase correction

was performed using the corresponding water spectrum from each area. Each spectrum was then

apodized using a Gaussian filter of 3Hz to improve signal quality, reduce noise and reduce effects

of signal truncation (Jiru, 2008). The residual water peak was removed using the Hankel-Lanczos

Singular Value Decomposition (HLSVD) filter tool.

Metabolite models were generated using the software Versatile Simulation, Pulses and Analysis

(VEsPA (https://scion.duhs.duke.edu/vespa/project); Soher, Semanchuk, Todd, Steinberg, &

Young., 2010). 14 typical brain metabolites (Acetate, Aspartate, CHO, Creatine, Gamma-

Aminobutyric Acid (GABA), Glucose, Glutamate, Glutamine, Lactate, Myo-inositol, N-acetyl

Aspartate (NAA), Phosphocreatine, PC & GPC, Scyllo-inositol, Succinate, Taurine) were simulated

at a field strength of 3T using a PRESS pulse sequence (TE1 = 20ms, TE2 = 10ms, main field =

123.25MHz). For initial analyses, CHO was modelled separately from PC+GPC based on the

method described in Bell et al., 2018. Additionally, the sum of the three peaks (total choline,

tCHO) was included in the analyses for comparison. If tCHO produced similar results to CHO, it

would potentially suggest that there may not be a need to separate the three peaks, or that the

quantitation method is not separating CHO effectively.

The jMRUI tool Accurate Quantification of Short Echo time domain Signals (AQSES) was used for

automatic quantification of spectra signals. AQSES was applied using the method described in

Minati, Aquino, Bruzzone, & Erbetta, 2010. To correct for any chemical shift displacement, the

spectrum was shifted so that the peak for n-acetyl-aspartate (NAA) was at 2.02ppm. The frequency

range selected for processing was limited to 0-8.6ppm (equal phase for all metabolites, begin time fixed, delta damping (-10 to 25Hz), delta frequency (-5 to 5Hz), no background handling, 0 truncated points, 2048 points in AQSES and normalisation on). Based on common practice in the field, values with a CRB higher than 30% were excluded on a case by case basis.

Metabolite concentrations were calculated for CHO, PC+GPC, tCHO, NAA and total creatine (tCR, creatine + phosphocreatine), correcting for partial-volume and relaxation effects, using the formula described in (Gasparovic et al., 2006).

Experimental Design and Statistical Analysis

Statistical analysis was performed using SPSS (IBM Corp. Released 2013. IBM SPSS Statistics for Windows, Version 22.0. Armonk, NY: IBM Corp).

The relationships between model parameters and behaviour, along with model parameters and metabolite levels and behaviour and metabolite levels was assessed using correlation analysis. The distribution of the data was analysed using measures of skewness and kurtosis, along with the Shapiro-Wilk test. When the assumptions of normality and homogeneity were met, Pearson's correlation (r) was used to assess correlations. When assumptions of normality were not met, Kendall's Tau ($r_\square$) was used to assess correlations, as it provides a better estimation of the correlation in a small sample size compared to other non-parametric methods (Field, 2009). Where appropriate, hierarchical multiple regression analysis was used to assess the variance explained by metabolite levels in behaviour, after the model parameters were accounted for.

Confounding Variables

There were no significant differences in metabolite levels between hemispheres, therefore the results were combined across hemisphere of acquisition.

To examine if variations in the metabolite values might be caused by differing proportions of tissue

composition, correlations were performed between CCC levels and proportion of grey and white

matter present in the voxel. Additionally, metabolite values were checked against the water signal

for the same reason. No significant correlations were found between CCCs and grey/white matter

content, indicating any variance seen is generated by differing metabolite levels. The water signal

significantly correlated with DS tCHO ($r_□$ (35) = -0.348, p = 0.003) and VS PC+GPC ($r_□$ (32) = -

0.270, p = 0.001). Therefore, analyses involving DS tCHO or VS PC+GPC were corrected for this

source of variance using partial correlations. No other significant correlations were seen between

the water signal and metabolite levels of interest.

There is evidence that metabolite levels in the brain can vary based on time of day (Soreni et al.,

2006) and age (Pfefferbaum, Adalsteinsson, Spielman, Sullivan, & Lim, 1999; Reyngoudt et al.,

2012). Therefore, all metabolites were checked against these two variables to ensure this was not a

source of variance. Time of day significantly correlated with DS tCHO ($r_□$ (35) = 0.249, p = 0.038)

and cerebellum tCHO ($r_□$ (31) = 0.285, p = 0.026). Therefore, analyses involving DS tCHO or

cerebellum tCHO were corrected for this source of variance using partial correlations. No other

significant correlations were seen between metabolite levels and time of day or age of participant.

Controls

The cerebellum was used as a control to demonstrate the regional specificity of results. None of the

effects were present in the cerebellum and therefore these results are not reported further. NAA and

tCR were used as controls to demonstrate the neurochemical specificity of the results (i.e. that the

relevant individual differences were specific to choline and not to spectrum-wide inter-individual

differences). None of the effects were present in either NAA or tCR and therefore these results are

not reported further. Furthermore, none of the reported effects were found when using tCHO as a

measure of cholinergic availability and therefore these results are not reported further.

 # RESULTS

 # **Behavioural Results**

Twenty-two (22) participants reached criterion during both rounds (i.e. they reached criterion both

during initial learning and after the reversal) and were included in the analysis.

Model parameters and performance

A reinforcement-learning model was used to disentangle components of learning that contribute to

overall behaviour. We looked at three parameters of interest, the learning rates from positive (η^{+})

and negative (η^{-}) prediction errors, and the overall impact of subjective value of the deck on the

participants choice (value impact parameter, β). To test the contribution of the model parameters to

behaviour, we looked at correlations between behaviour (as measured by trials to criterion, number

of perseverative trials and number of regressive errors) and the corresponding model parameters,

i.e. behaviour during initial learning was correlated with model parameters fit over the initial

learning period, and likewise for the reversal learning period.

Table 1: Performance variables

	Average Number of Trials	*SD*
Initial Learning	44	28
Reversal Learning		
Perseveration Period	12	8
Post Reversal Learning	35	22
Regressive Errors	7	6
Total	47	

381   Table 2: Estimates of model parameters

	η^+	η^-	β
Initial Learning	0.37	0.42	1.44
	(SD = 0.30)	(SD = 0.31)	(SD = 0.56)
Reversal Learning	0.24	0.31	1.37
	(SD = 0.35)	(SD = 0.27)	(SD = 0.97)

Note: η^+ = learning rate from positive prediction errors; η^- = learning rate from negative prediction errors; β =
impact of subjective value on choice.

Table 3 shows the correlation coefficients for the relationships between model parameters and

behaviour. Faster initial learning (low number of trials to criterion) was associated with a higher

learning rate from positive prediction errors ($r(22)$ = -0.439, p = 0.041) and a higher value impact

parameter ($r(21)$ = -0.536, p = 0.012). A lower number of perseverative trials was associated with a

higher learning rate from negative prediction errors ($r(22)$ = -0.527, p = 0.012). As was the case

during initial learning, during post-reversal learning (after the reversal has been identified) a lower

number of trials taken to reach criterion was associated with a higher learning rate from positive

prediction errors (r (22) = -0.335, p = 0.03), and a higher value impact parameter (r (22) = -

0.352, p = 0.022). Additionally, during post-reversal learning, a lower number of regressive errors

was associated with a higher learning rate from positive prediction errors (r (22) = -0.355, p =

0.023) and a higher value impact parameter (r (22) = -0.337, p = 0.031).

Table 3: Correlation coefficients for relationships between model parameters and behaviour

	η^+	η^-	β
Initial Learning (TTC)	-0.439*	-0.218	-0.536*
Reversal Learning			
Perseverative Errors	-0.176	-0.527*	0.132
Post Reversal Learning (TTC)	-0.335*	0.322	-0.352*
Regressive Errors	-0.355*	0.292	-0.337*

396   Note: η^+ = learning rate from positive prediction errors; η^- = learning rate from negative prediction errors; β
397   = value impact parameter; * p<0.05.

Effects of trait impulsivity on performance

To investigate the influence of impulsivity on decision making, we looked at correlations between impulsivity (total BIS-11 score) and measures of behaviour (including model parameters) in learners. Higher impulsivity levels were associated with a lower number of perseverative errors ($r(22) = -0.470$, p = 0.027). No other measures of behaviour correlated with impulsivity.

Summary

Faster initial learning was indexed by both higher learning rates from positive prediction errors (R1η^+) and higher value impact parameters (R1β). Reduced numbers of perseverative trials were associated with higher learning rates from negative prediction errors (R2η^-) and higher impulsivity levels. Similar to initial learning, faster post-reversal learning was associated with higher learning rates from positive prediction errors (R2η^+) and higher value impact parameters (R2β). Additionally, during post-reversal learning, lower numbers of regressive errors were associated with higher learning rates from positive prediction errors (R2η^+) and higher value impact parameters (R2β).

Spectroscopy Results

One participant was excluded from spectroscopy analysis due to issues with segmentation of the structural scan.

Table 4: Average metabolite levels in the DS

	CHO	*PC+GPC*	*tCHO*	*NAA*	*tCR*
Learners	0.15	0.27	0.42	8.73	11.58
	(SD = 0.20)	(SD = 0.10)	(SD = 0.12)	(SD = 0.77)	(SD = 1.74)
Non-Learners	0.11	0.36	0.46	8.83	11.80
	(SD = 0.16)	(SD = 0.14)	(SD = 0.10)	(SD = 2.37)	(SD = 2.31)

Note: CHO = choline, PC+GPC = phosphocholine and glycerophosphocholine, tCHO = total choline, NAA = n-acetyl aspartate, tCR = total creatine.

418 Table 5: Average metabolite levels in the VS

	CHO	PC+GPC	tCHO	NAA	tCR
Learners	0.24	0.27	0.5	5.39	12.02
	(SD = 0.17)	(SD = 0.12)	(SD = 0.17)	(SD = 1.97)	(SD = 2.26)
Non-Learners	0.23	0.25	0.48	5.45	11.13
	(SD = 0.17)	(SD = 0.14)	(SD = 0.16)	(SD = 1.54)	(SD = 3.95)

419 Note: CHO = choline, PC+GPC = phosphocholine and glycerophosphocholine, tCHO = total choline, NAA
420 = n-acetyl aspartate, tCR = total creatine.

Group Comparisons

422 To investigate whether average levels of CHO in the striatum relate to task performance, the
423 average levels were compared between learners and non-learners. There was no significant
424 difference in CHO levels between learners and non-learners in either the DS or the VS.

Correlations with Behaviour

426 To further investigate the relationship between average metabolite levels and task performance,
427 correlations were performed between metabolite levels and measures of performance in learners
428 (numbers of trials to criterion and model parameters).

Initial Learning

No significant correlations were seen with measures of performance in round 1 (trials to criterion,
$R1\eta^+$ or $R1\beta$) and average levels of CHO in the DS.

VS CHO did not correlate with trials to criterion in round 1. However, low levels of CHO in the VS
were associated with higher learning rates from positive prediction errors ($r(20) = -0.625$, p =
0.003; Figure 4A) and lower value impact parameters ($r(19) = 0.555$, p = 0.014; Figure *4*B).

Figure 4: Correlations between VS CHO levels and performance during initial learning **A:** Negative

correlation between learning rate based on positive prediction errors derived from round 1 (R1η+) and levels

of CHO in the VS (r(20) = -0.625, p = 0.003). **B:** Negative correlation between impact of participant's

subjective value on their future choice derived from round 1 (R1β) and levels of CHO) in the VS (r(19) =

0.555, p = 0.014). VS: Ventral Striatum; CHO: Choline.

Perseverative Trials

A lower number of perseverative trials was associated with lower levels of DS CHO ($r_\square$ (21) =

0.367, p = 0.021; Figure 5A). The opposite effect was seen with DS PC+GPC (r(21) = -0.447, p =

0.042. Additionally, higher learning rates from negative prediction errors were associated with

lower DS CHO levels ($r_\square$ (21) = -0.371, p = 0.019; Figure 5B). This result is specific to DS CHO,

with no other DS metabolites found to correlate with learning rates from negative prediction errors.

Additionally, VS CHO was not found to correlate with either the number of perseverative trials or

learning rates from negative prediction errors.

Figure 5: Correlations between DS CHO levels and performance during reversal **A:** Positive correlation
between the number of perseverative trials and levels of CHO in the DS (r_s (21) = 0.367, p = 0.021). **B:**
Negative correlation between the learning rate based on negative prediction errors derived from round 2
(R2η-) and levels of CHO in the DS (r_s (21) = -0.371, p = 0.019). DS: Dorsal Striatum; CHO: Choline.

After establishing an association between CHO levels and reversal performance, we wanted to

examine whether CHO contributed to reversal efficiency over and above behavioural and

personality variables. Using a hierarchical multiple regression, we first modelled the contribution of

variance from learning rates from negative prediction errors and total BIS scores to the variance in

the number of perseverative trials (Model 1; $F(2,18) = 9.460$ p = 0.002, $R^2 = 0.512$; Table 6). The

second model looked at whether the addition of DS CHO would explain significantly more

variance, over and above that explained by learning rates from negative prediction errors and total

BIS score (Model 2; $F(3,17) = 9.574$ p = 0.001, $R^2 = 0.628$; Table 6).

The amount of variance in the number of perseverative trials explained by learning rates from

negative prediction errors was significant in both Model 1 (β = -0.493, t(18) = -2.980, p = 0.008;

Table 6) and Model 2 (β = -0.430, t(17) = -2.843, p = 0.011; Table 6). Additionally, total BIS score

also explained a significant amount of variance in both Model 1 (β = -0.472, t(18) = -2.855, p =

0.011; Table 6) and Model 2 (β = -0.419, t(17) = -2.787, p = 0.013; Table 6).

In Model 2, DS CHO also explained a significant amount of variance in the number of

perseverative trials ($\beta = 0.351$, $t(17) = 2.300$, $p = 0.034$; Table 6). The addition of DS CHO to the

model increased R^2 by 0.116 and this increase was statistically significant ($F(1,23) = 5.291$, $p =$

0.034; Table 6).

To assess the specificity of this result, DS PC+GPC was also included in the model. However,

analysis of multicollinearity diagnostics showed a tolerance of 0.175, which is below the acceptable

value of 0.2. This is due to the strong significant correlation between DS CHO and DS PC+GPC (r

(21) = -0.667 $p < 0.001$). As a result, including the two variables in the same regression model

would violate the assumption of multicollinearity and the regression model would not be able to

provide unique estimates of the regression coefficients, as each will account for overlapping

variance (Field, 2009). Therefore, we instead repeated the hierarchical regression with DS PC+GPC

in place of DS CHO. The amount of variance explained by DS PC+GPC was not significant ($\beta = -$

0.301, $t(17) = -1.900$, $p = 0.075$). The addition of DS PC+GPC to the model increased R^2 by 0.085

and this increase was not statistically significant ($F(1,23) = 3.611$, $p = 0.075$). This indicates that DS

CHO levels can explain part of the variance in the number of perseverative trials, however DS

PC+GPC levels cannot.

Table 6: Summary of hierarchical regression analyses for variables predicting perseveration

		B	SE B	β	R^2	ΔR^2	p
Model 1					0.512		0.002
	R2η⁻	-14.476	4.858	-0.493			0.008
	BIS Total	-0.504	0.176	-0.472			0.011
Model 2					0.628	0.116	0.034
	R2η⁻	-12.619	4.439	-0.430			0.011
	BIS Total	-0.447	0.160	-0.419			0.013
	DS CHO	5.306	2.307	0.351			0.034

Note, for $\Delta R^2 = 0.139$, p = 0.037

B = unstandardized coefficient, SE = standard error, β = standardised coefficient

Post Reversal Learning

No significant correlations were seen with either DS or VS CHO levels and measures of

performance during post reversal learning (trials to criterion, $R2\eta^{+}$ or $R2\beta$). Additionally, there were

no significant correlations between DS or VS CHO levels and the number of regressive errors.

Summary
492

In the DS, average CHO levels were associated with performance during reversal, but not during

initial learning. There was a significant positive correlation between DS CHO levels and the

number of perseverative trials, and a significant negative correlation between DS CHO levels and

learning rates from negative prediction errors ($R2\eta^{-}$). Additionally, DS CHO levels explained

variance in the number of perseverative trials over and above that explained by learning rates from

negative prediction errors.

In the VS, average CHO levels were not associated with performance during reversal learning.

Although VS CHO levels were not associated with the speed of initial learning, there was a

significant positive correlation between VS CHO levels and learning rates from positive prediction

errors, and a significant negative correlation between VS CHO levels and the value impact

parameter during initial learning.

DISCUSSION

We used [1]H-MRS to investigate the relationship between average CHO levels in the human striatum at rest and performance during probabilistic reversal learning. Here we show that baseline levels of CHO in the human DS are associated specifically with individual differences in reversal learning efficiency, but not in initial learning, and that this effect is specific to the dorsal, but not the ventral striatum.

Behaviourally, we show that faster initial learning is indexed by a higher learning rate from positive prediction errors (η^+) and a higher value impact parameter (β). Therefore, during this period, participants are using wins and expected value to guide their choices. This is also seen during the post-reversal learning period, in which faster post-reversal learning is indexed by higher learning rates from positive prediction errors (η^+) and higher value impact parameters (β). Faster reversal (less perseveration), however, was indexed by higher learning rates from negative prediction errors (η^-) only. During this period, participants must now pay increased attention to worse than expected outcomes in order to identify the change in contingencies. Therefore, to adapt to changes in task structure, participants adapt their strategy by altering the weight of learning from prediction errors based on reward history.

The learning rate for negative prediction errors, accounting for trait impulsivity, explained a significant amount of variance during perseveration, providing a simple mechanism to explain reversal efficiency. However, average DS CHO levels explained variance in the number of perseverative trials over and above this original model. This suggests a more complex mechanism in which perseveration is influenced, in part, by the learning rate from negative prediction errors (which can change due to task demand) and by resting levels of DS CHO. Indeed, Franklin & Frank, 2015 showed that a model which takes into account cholinergic activity performs better on a reversal learning task than a model based solely on dopamine prediction error signalling.

Our results indicate that participants who were quicker to reverse had lower average levels of DS

CHO, suggesting that low trait levels of DS CHO are beneficial for reversal learning. Based on

evidence that ACh efflux increases during reversal learning (H. D. Brown et al., 2010; Ragozzino et

al., 2009), this suggests two potential mechanisms. Firstly, lower levels of DS CHO at rest could

reflect lower levels of ACh at rest. This is also supported by evidence from the animal literature,

which has shown a positive correlation between ACh levels at rest as measured by microdialysis

and average CCCs as measured by [1]H-MRS (Wang, Du, Tian, & Wang, 2008). Additionally, higher

levels of CHO availability have been shown to lead to higher levels of ACh release, implying a

positive correlation between the two metabolites (Koshimura et al., 1990). Based on this notion, the

findings here suggest that lower levels of ACh at rest may be beneficial for reversal learning

because they enable a higher contrast between ACh levels at rest and during reversal learning.

However, it is important to note that Wang et al. (2008) modelled all three CCCs as a single peak. It

is likely that the relationship between CHO levels as measured by spectroscopy and ACh levels in

the brain is not straightforward, and this interpretation should be considered with caution. Indeed,

animal studies have shown the relationship between CHO and ACh can change based on neuronal

firing and ACh requirement (Klein, Weichel, Ruhr, Dvorak, & Loffelholz, 2002; Löffelholz, 1998).

Furthermore, we have previously demonstrated a drop in CHO levels in the human DS during

reversal learning, thought to reflect the sustained increase in ACh release seen in animal studies

(e.g. Ragozzino et al., 2009). This drop is thought to be due to an increase in translocation of CHO

uptake receptors in response to sustained neural firing (Bell et al., 2018). Though we have described

the measurements in this study as "at rest", cholinergic interneurons are tonically active, and

therefore the relationship between CHO and ACh levels in the striatum will likely reflect a more

complex dynamical relationship between the two.

The second potential mechanism supported by our findings is that lower levels of DS CHO at rest

may result from a more efficient CHO uptake system. Mice carrying mutations in the gene coding

for CHO uptake transporters have reduced neuronal capacity to both clear CHO and release ACh.

Moreover, performance on an attention task was impaired in these mice (Parikh et al., 2013).

Additionally, in a study of frontal cortex cholinergic modulation during attention, humans with a

gene polymorphism which reduces CHO transport capacity showed reduced activation in the

prefrontal cortex during an attentional task. Furthermore, the pattern of activation predicted CHO

genotype (Berry, Blakely, Sarter, & Lustig, 2015). Further work is needed to determine the

relationship between CHO uptake, ACh release and reversal learning.

Disruption of cholinergic signalling in rodents typically results in an increase in regressive errors

(Bradfield et al., 2013; H. D. Brown et al., 2010). However, here we found no association between

DS CHO levels and the number of regressive errors. In humans, measures of individual differences

in perseverative and regressive errors are likely to be confounded by individual differences in

representation of the task structure. Rather than making perseverative and regressive errors based

solely on feedback, the ability to flexibly alter response depends in part on a higher level

representation of the task, which is thought to be maintained in frontal areas of the cortex

(Armbruster, Ueltzhöffer, Basten, & Fiebach, 2012). It should be noted that the basal ganglia-

thalamo-cortical system has been shown to be modulated by the maintenance of task rules, with

those with stronger representation of the task structure showing higher activation in the caudate and

thalamus during a behaviour switch (Ueltzhöffer, Armbruster-Genç, & Fiebach, 2015), indicating

that representation of task structure likely modulates DS activity in response to the need for

behavioural flexibility. Inevitably, caution is needed when translating evidence from rodent studies

of learning to human studies. This emphasises the need to further develop non-invasive techniques

for studying human neurochemistry *in vivo*

As predicted, and in line with evidence from the animal literature (Ragozzino et al., 2009), levels of

CHO in the VS were not associated with reversal learning. However, VS CHO levels were

associated with model parameters which contributed to initial learning. Though Ragozzino et al.

demonstrated that ACh levels in the VS did not change during reversal learning, they did not test if

they changed during initial learning. Successful learning requires the ability to learn from feedback,

which is signalled by dopaminergic prediction error signalling in the VS (Schultz, Dayan, &

Montague, 1997). The rodent VS has a higher density of cholinergic interneurons than the DS

(Matamales, Götz, & Bertran-Gonzalez, 2016) and changes in cholinergic activity are time locked

to changes in dopaminergic activity, which is thought to enhance the contrast of prediction error

signalling (Aosaki, Miura, Suzuki, Nishimura, & Masuda, 2010). Indeed, cholinergic activity in the

VS has been linked with effective learning of a stimulus-outcome association (M. T. C. Brown et

al., 2012), therefore, it is likely that cholinergic activity in the VS is involved in some aspect with

goal-directed learning, and further studies should explore this contribution.

We used several controls to demonstrate that these effects are specific to CHO levels in the

striatum, not least because our MRS application method is novel. We acquired data from a voxel in

the cerebellum, geometrically identical to the striatal voxels. No learning effects were present in the

cerebellum, demonstrating that our findings are specific to the striatum. Additionally, we also

quantified two control metabolites (NAA and tCR) to ensure that the results were specific to the

metabolite of interest, rather than a general measurement or region effect. None of the effects were

seen in levels of NAA and tCR in the DS or VS. Importantly, none of the effects were seen when

modelling all three peaks together (tCHO), highlighting once more the importance of separating

CHO when using [1]H-MRS to investigate individual differences in CCC levels.

In conclusion, [1]H-MRS was used to demonstrate that average levels of CHO in the human DS are

associated with performance during probabilistic reversal, but not during initial learning. This is in

line with evidence from the animal literature and out own prior work with humans which suggests a

specific role for cholinergic activity in the DS during reversal learning. These results provide

evidence for the role of the human cholinergic striatum in reversal learning and behavioural

flexibility more generally. Additionally, these findings further support the idea of using CHO levels

as measured by [1]H-MRS as a tool for non-invasive *in vivo* monitoring of both healthy human

neurochemistry, as well as disorders of the human cholinergic system.

REFERENCES

Aoki, S., Liu, A. W., Akamine, Y., Zucca, A., Zucca, S., & Wickens, J. R. (2018). Cholinergic interneurons in the rat striatum modulate substitution of habits. *European Journal of Neuroscience*, *47*(10), 1194–1205. http://doi.org/10.1111/ejn.13820

Aoki, S., Liu, A. W., Zucca, A., Zucca, S., & Wickens, J. R. (2015). Role of Striatal Cholinergic Interneurons in Set-Shifting in the Rat. *Journal of Neuroscience*, *35*(25), 9424–9431. http://doi.org/10.1523/JNEUROSCI.0490-15.2015

Aosaki, T., Miura, M., Suzuki, T., Nishimura, K., & Masuda, M. (2010). Acetylcholine-dopamine balance hypothesis in the striatum: An update. *Geriatrics and Gerontology International*, *10*(SUPPL. 1), S148–S157. http://doi.org/10.1111/j.1447-0594.2010.00588.x

Armbruster, D. J. N., Ueltzhöffer, K., Basten, U., & Fiebach, C. J. (2012). Prefrontal cortical mechanisms underlying individual differences in cognitive flexibility and stability. *Journal of Cognitive Neuroscience*, *24*(12), 2385–2399. http://doi.org/10.1162/jocn_a_00286

Bayard, S., Raffard, S., & Gely-Nargeot, M.-C. (2011). Do facets of self-reported impulsivity predict decision-making under ambiguity and risk? Evidence from a community sample. *Psychiatry Research*, *190*(2–3), 322–326. http://doi.org/10.1016/j.psychres.2011.06.013

Bell, T., Lindner, M., Mullins, P. G., & Christakou, A. (2018). Functional neurochemical imaging of the human striatal cholinergic system in reversal learning. *European Journal of Neuroscience*, *47*(10), 1184–1193. http://doi.org/10.1111/ejn.13803

Berry, A. S., Blakely, R. D., Sarter, M., & Lustig, C. (2015). Cholinergic capacity mediates prefrontal engagement during challenges to attention: Evidence from imaging genetics. *NeuroImage*, *108*, 386–395. http://doi.org/10.1016/j.neuroimage.2014.12.036

Bradfield, L. A., Bertran-Gonzalez, J., Chieng, B., & Balleine, B. W. (2013). The thalamostriatal

pathway and cholinergic control of goal-directed action: interlacing new with existing learning

in the striatum. *Neuron*, *79*(1), 153–166. http://doi.org/10.1016/j.neuron.2013.04.039

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433–436.

Brown, H. D., Baker, P. M., & Ragozzino, M. E. (2010). The parafascicular thalamic nucleus

concomitantly influences behavioral flexibility and dorsomedial striatal acetylcholine output in

rats.   *The*   *Journal*   *of*   *Neuroscience*,   *30*(43),   14390–8.

http://doi.org/10.1523/JNEUROSCI.2167-10.2010

Brown, M. T. C., Tan, K. R., O'Connor, E. C., Nikonenko, I., Muller, D., & Lüscher, C. (2012).

Ventral tegmental area GABA projections pause accumbal cholinergic interneurons to enhance

associative learning. *Nature*, *492*(7429), 452–6. http://doi.org/10.1038/nature11657

Bull, P. N., Tippett, L. J., & Addis, D. R. (2015). Decision making in healthy participants on the

Iowa Gambling Task: new insights from an operant approach. *Frontiers in Psychology*,

6(APR). http://doi.org/10.3389/fpsyg.2015.00391

Christakou, A., Gershman, S. J., Niv, Y., Simmons, A., Brammer, M., & Rubia, K. (2013). Neural

and Psychological Maturation of Decision-making in Adolescence and Young Adulthood.

Journal of Cognitive Neuroscience, *25*(11), 1807–1823. http://doi.org/10.1162/jocn_a_00447

Currie, S., Hadjivassiliou, M., Wilkinson, I. D., Griffiths, P. D., & Hoggard, N. (2013). Magnetic

resonance spectroscopy of the normal cerebellum: What degree of variability can be expected?

Cerebellum, *12*(2), 205–211. http://doi.org/10.1007/s12311-012-0415-1

Daw, N. D. (2011). Trial-by-trial data analysis using computational models. *Attention and*

Performance XXIII, *23*, 1.

Field, A. (2009). Discovering Statistics using SPSS. In *Third Edition*.

Franken, I. H. A., van Strien, J. W., Nijs, I., & Muris, P. (2008). Impulsivity is associated with

behavioral decision-making deficits. *Psychiatry Research*, *158*(2), 155–163. http://doi.org/10.1016/j.psychres.2007.06.002

Franklin, N. T., & Frank, M. J. (2015). A cholinergic feedback circuit to regulate striatal population uncertainty and optimize reinforcement learning. *ELife*, *4*(DECEMBER2015). http://doi.org/10.7554/eLife.12029

Gasparovic, C., Chen, H., & Mullins, P. G. (2018). Errors in 1 H-MRS estimates of brain metabolite concentrations caused by failing to take into account tissue-specific signal relaxation. *NMR in Biomedicine*, (November 2017), e3914. http://doi.org/10.1002/nbm.3914

Gasparovic, C., Song, T., Devier, D., Bockholt, H. J., Caprihan, A., Mullins, P. G., … Morrison, L. A. (2006). Use of tissue water as a concentration reference for proton spectroscopic imaging. *Magnetic Resonance in Medicine*, *55*(6), 1219–1226. http://doi.org/10.1002/mrm.20901

Gray, J. A. (1970). The psychophysiological basis of introversion-extraversion. *Behaviour Research and Therapy*, *8*(3), 249–266. http://doi.org/10.1016/0005-7967(70)90069-0

Holmstrand, E. C., Lund, D., Cherian, A. K., Wright, J., Martin, R. F., Ennis, E. A., … Blakely, R. D. (2013). Transgenic overexpression of the presynaptic choline transporter elevates acetylcholine levels and augments motor endurance. *Neurochemistry International*, *73*, 217–28. http://doi.org/10.1016/j.neuint.2013.11.008

Hoover, D. B., Muth, E. A., & Jacobowitz, D. M. (1978). A mapping of the distribution of acetycholine, choline acetyltransferase and acetylcholinesterase in discrete areas of rat brain. *Brain Research*, *153*(2), 295–306. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/687983

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). FSL. *NeuroImage*, (62), 782–790.

Jiru, F. (2008). Introduction to post-processing techniques. *European Journal of Radiology*, *67*(2),

202–17. http://doi.org/10.1016/j.ejrad.2008.03.005

Jope, R. S. (1979). High Affinity Choline Transport and AcetylCoA Production in Brain and their roles in the Regulation of Acetyl-Choline Synthesis. *Brain Research Reviews*, *1*, 313–344.

Klein, J., Weichel, O., Ruhr, J., Dvorak, C., & Loffelholz, K. (2002). A homeostatic mechanism counteracting K+-evoked choline release in adult brain. *Journal of Neurochemistry*, *80*(5), 843–849. http://doi.org/10.1046/j.0022-3042.2001.00754.x

Koshimura, K., Miwa, S., Lee, K., Hayashi, Y., Hasegawa, H., Hamahata, K., … Itokawa, Y. (1990). Effects of choline administration on in vivo release and biosynthesis of acetylcholine in the rat striatum as studied by in vivo brain microdialysis. *Journal of Neurochemistry*, *54*(2), 533–9.

Leong, Y. C., & Niv, Y. (2013). Human reinforcement learning processes act on learned attentionally-filtered representations of the world. *Reinforcement Learning and Decision-Making Conference*, *43*, 43–47.

Lindner, M., Bell, T., Iqbal, S., Mullins, P. G., & Christakou, A. (2017). In vivo functional neurochemistry of human Cortical cholinergic function during visuospatial attention. *PLoS ONE*, *12*(2), e0171338. http://doi.org/10.1371/journal.pone.0171338

Lockman, P. R., & Allen, D. D. (2002). The transport of choline. *Drug Development and Industrial Pharmacy*, *28*(7), 749–71. http://doi.org/10.1081/DDC-120005622

Löffelholz, K. (1998). Brain choline has a typical precursor profile. *Journal of Physiology-Paris*, *92*(3–4), 235–239. http://doi.org/10.1016/S0928-4257(98)80025-9

Matamales, M., Götz, J., & Bertran-Gonzalez, J. (2016). Quantitative imaging of cholinergic interneurons reveals a distinctive spatial organization and a functional gradient across the mouse striatum. *PLoS ONE*, *11*(6), e0157682. http://doi.org/10.1371/journal.pone.0157682

McCool, M. F., Patel, S., Talati, R., & Ragozzino, M. E. (2008). Differential involvement of M1-

type and M4-type muscarinic cholinergic receptors in the dorsomedial striatum in task

switching. *Neurobiology of Learning and Memory*, *89*(2), 114–124.

http://doi.org/10.1016/j.nlm.2007.06.005

Minati, L., Aquino, D., Bruzzone, M., & Erbetta, A. (2010). Quantitation of normal metabolite

concentrations in six brain regions by in-vivo 1 H-MR spectroscopy. *Journal of Medical*

Physics, *35*(3), 154. http://doi.org/10.4103/0971-6203.62128

Müller, V. I., Langner, R., Cieslik, E. C., Rottschy, C., & Eickhoff, S. B. (2015). Interindividual

differences in cognitive flexibility: influence of gray matter volume, functional connectivity

and trait impulsivity. *Brain Structure and Function*, *220*(4), 2401–2414.

http://doi.org/10.1007/s00429-014-0797-6

Naressi, A., Couturier, C., Devos, J. M., Janssen, M., Mangeat, C., de Beer, R., & Graveron-

Demilly, D. (2001). Java-based graphical user interface for the MRUI quantitation package.

Magnetic Resonance Materials in Physics, Biology and Medicine, *12*(2–3), 141–52. Retrieved

from http://www.ncbi.nlm.nih.gov/pubmed/11390270

Niv, Y., Daniel, R., Geana, A., Gershman, S. J., Leong, Y. C., Radulescu, A., & Wilson, R. C.

(2015). Reinforcement Learning in Multidimensional Environments Relies on Attention

Mechanisms. *Journal of Neuroscience*, *35*(21), 8145–8157.

http://doi.org/10.1523/JNEUROSCI.2978-14.2015

Niv, Y., Edlund, J. A., Dayan, P., & O'Doherty, J. P. (2012). Neural Prediction Errors Reveal a

Risk-Sensitive Reinforcement-Learning Process in the Human Brain. *Journal of Neuroscience*,

32(2), 551–562. http://doi.org/10.1523/JNEUROSCI.5498-10.2012

Okada, K., Nishizawa, K., Setogawa, S., Hashimoto, K., & Kobayashi, K. (2018). Task-dependent

function of striatal cholinergic interneurons in behavioural flexibility. *European Journal of*

Neuroscience, *47*(10), 1174–1183. http://doi.org/10.1111/ejn.13768

Parikh, V., St. Peters, M., Blakely, R. D., & Sarter, M. (2013). The Presynaptic Choline Transporter Imposes Limits on Sustained Cortical Acetylcholine Release and Attention. *Journal of Neuroscience*, *33*(6), 2326–2337. http://doi.org/10.1523/JNEUROSCI.4993-12.2013

Patton, J. H., Stanford, M. S., & Barratt, E. S. (1995). Factor structure of the barratt impulsiveness scale. *Journal of Clinical Psychology*, *51*(6), 768–774. http://doi.org/10.1002/1097-4679(199511)51:6<768::AID-JCLP2270510607>3.0.CO;2-1

Pfefferbaum, A., Adalsteinsson, E., Spielman, D., Sullivan, E. V, & Lim, K. O. (1999). In vivo spectroscopic quantification of the N-acetyl moiety, creatine, and choline from large volumes of brain gray and white matter: effects of normal aging. *Magnetic Resonance in Medicine*, *41*(2), 276–84. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/10080274

Puts, N. A. J., & Edden, R. A. E. (2012). In vivo magnetic resonance spectroscopy of GABA: A methodological review. *Progress in Nuclear Magnetic Resonance Spectroscopy*, *60*, 29–41. http://doi.org/10.1016/J.PNMRS.2011.06.001

Ragozzino, M. E., Jih, J., & Tzavos, A. (2002). Involvement of the dorsomedial striatum in behavioral flexibility: role of muscarinic cholinergic receptors. *Brain Research*, *953*(1–2), 205–214. http://doi.org/10.1016/S0006-8993(02)03287-0

Ragozzino, M. E., Mohler, E. G., Prior, M., Palencia, C. A., & Rozman, S. (2009). Acetylcholine activity in selective striatal regions supports behavioral flexibility. *Neurobiology of Learning and Memory*, *91*(1), 13–22. http://doi.org/10.1016/j.nlm.2008.09.008

Reyngoudt, H., Claeys, T., Vlerick, L., Verleden, S., Acou, M., Deblaere, K., … Achten, E. (2012). Age-related differences in metabolites in the posterior cingulate cortex and hippocampus of normal ageing brain: a 1H-MRS study. *European Journal of Radiology*, *81*(3), e223-31. http://doi.org/10.1016/j.ejrad.2011.01.106

Schultz, W., Dayan, P., & Montague, P. R. (1997). A Neural Substrate of Prediction and Reward.

Science, *275*(5306), 1593–1599. http://doi.org/10.1126/science.275.5306.1593

Smith, S. M. (2002). Fast robust automated brain extraction. *Human Brain Mapping*, *17*(3), 143–

155.

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg,

H., … Matthews, P. M. (2004). Advances in functional and structural MR image analysis and

implementation as FSL. *NeuroImage*, *23*(S1), 208–219.

Soher, B. J., Semanchuk, P., Todd, D., Steinberg, J., & Young., K. (2010). Vespa: Integrated

applications for RF pulse design, spectral simulation and MRS data analysis. In *ISMRM*

Stockholm, Sweden (p. 3169).

Soreni, N., Noseworthy, M. D., Cormier, T., Oakden, W. K., Bells, S., & Schachar, R. (2006).

Intraindividual variability of striatal (1)H-MRS brain metabolite measurements at 3 T.

Magnetic Resonance Imaging, *24*(2), 187–94. http://doi.org/10.1016/j.mri.2005.10.027

Steingroever, H., Wetzels, R., & Wagenmakers, E. (2016). Bayes factors for reinforcement-learning

models   of   the   Iowa   gambling   task.   *Decision*,   *3*(2),   115–131.

http://doi.org/10.1037/dec0000040

Sutton, R. S., & Barto, A. G. (1998). *Reinforcement Learning: An Introduction*. (P.-B. Books, Ed.),

IEEE   Transactions   on   Neural   Networks   (Vol.   9).   MIT   Press.

http://doi.org/10.1109/TNN.1998.712192

Tanimura, A., Pancani, T., Lim, S. A. O., Tubert, C., Melendez, A. E., Shen, W., & Surmeier, D. J.

(2018). Striatal cholinergic interneurons and Parkinson's disease. *European Journal of*

Neuroscience, *47*(10), 1148–1158. http://doi.org/10.1111/ejn.13638

Tzavos, A., Jih, J., & Ragozzino, M. E. (2004). Differential effects of M1 muscarinic receptor

blockade and nicotinic receptor blockade in the dorsomedial striatum on response reversal

learning. *Behavioural Brain Research*, *154*(1), 245–53.

http://doi.org/10.1016/j.bbr.2004.02.011

Ueltzhöffer, K., Armbruster-Genç, D. J. N., & Fiebach, C. J. (2015). Stochastic dynamics

underlying cognitive stability and flexibility. *PLoS Computational Biology*, *11*(6), e1004331.

http://doi.org/10.1371/journal.pcbi.1004331

Wang, X.-C., Du, X.-X., Tian, Q., & Wang, J.-Z. (2008). Correlation between choline signal

intensity and acetylcholine level in different brain regions of rat. *Neurochemical Research*,

33(5), 814–9. http://doi.org/10.1007/s11064-007-9509-4

Zhang, Y., Brady, M., & Smith, S. M. (2001). Segmentation of brain MR images through a hidden

Markov random field model and the expectation-maximization algorithm. *IEEE Trans Med*

Imaging, *20*(1), 45–57.

Zucca, A., Zucca, S., & Wickens, J. (2018). Cholinergic mechanisms in adaptive behaviour.

European Journal of Neuroscience, *47*(10), 1146–1147. http://doi.org/10.1111/ejn.13926