1   A genus definition for *Bacteria* and *Archaea* based on genome relatedness and taxonomic
2   affiliation.
3

4   Barco, R.A.[1,5*], Garrity, G.M.[3], Scott, J.J.[4], Amend, J.P.[1,2], Nealson, K.H.[1,2], and Emerson, D.[5]
5
[1]Department of Earth Sciences, University of Southern California, Los Angeles, CA, USA.
[2]Department of Biological Sciences, University of Southern California, Los Angeles, CA,
USA.
[3]Department of Microbiology and Molecular Genetics, Michigan State University, East
Lansing, MI, USA.
[4]Smithsonian Tropical Research Institute, Panama, Republic of Panama.
[5]Bigelow Laboratory for Ocean Sciences, East Boothbay, ME, USA.

13
14
15 Abstract
16

Genus assignment is fundamental in the characterization of microbes, yet there is
currently no unambiguous way to demarcate genera solely using standard genomic relatedness
indices. Here, we propose an approach to demarcate genera that relies on the combined use of
the average nucleotide identity, genome alignment fraction, and the distinction between type
species and non-type species. More than 750 genomes representing type strains of species from
10 different phyla, and 19 different taxonomic orders/families in Gram-positive/negative,
bacterial and archaeal lineages were tested. Overall, all 19 analyzed taxa conserved significant
genomic differences between members of a genus and type species of other genera in the same
taxonomic family. *Bacillus*, *Flavobacterium, Hydrogenovibrio*, *Lactococcu*s, *Methanosarcina*,
Thiomicrorhabdus, *Thiomicrospira*, *Shewanella*, and *Vibrio* are discussed in detail. Less than
1% of the type strains analyzed need reclassification, highlighting that the adoption of the 16S
rRNA gene as a taxonomic marker has provided consistency to the classification of
microorganisms in recent decades. One exception to this is the genus *Bacillus* with 61% of type
strains needing reclassification, including the human pathogens *B. cereus* and *B. anthracis*. The
results provide a first line of evidence that the combination of genomic indices provides
appropriate resolution to effectively demarcate genera within the current taxonomic framework
that is based on the 16S rRNA gene. We also identify the emergence of natural breakpoints at
the genome level that can further help in the circumscription of genera. Altogether, these results
show that a distinct difference between distant relatives and close relatives at the genome level
(i.e., genomic coherence) is an emergent property of genera in *Bacteria* and *Archaea*.

37
38 Introduction
39
At the time of writing, 17,300 bacterial and archaeal species and over 3,000 genera
with validly published names have been described in the taxonomic literature (Garrity, 2016;
Namesforlife Database); however, based on simple phylogenetic definition there are over
200,000 bacterial and archaeal species and 60,000 genera so far detected in the SILVA
database (Quast *et al.*, 2013; Yarza *et al.*, 2014). Sequence data stored in the Joint Genome
Institute (JGI) database, which includes data from other databases, has increased
exponentially over the past decade (Markowitz *et al.*, 2015) with >130,000 genomes of

bacterial and archaeal isolates, >6,000 metagenome-assembled genomes, and >2,000 single-
cell amplified genomes currently available. Because the number of available datasets is
constantly increasing (Mukherjee *et al.*, 2017), it is becoming easier to access data that
represent taxa of interest. Consequently, this added layer of available information could aid in
the formal characterization of microorganisms. An essential aspect of this characterization
process is the proper assignment of genus and species.
Historically, DNA-DNA hybridization (DDH) has been the 'gold standard' for species
delineation with a DDH value of ≥70% being recognized as the species boundary between
two strains (Tindall *et al.*, 2010 and other references therein). Stackebrandt and Goebel (1994)
conducted a correlation analysis between DDH values and 16S rRNA gene sequence
identities, and based on this, proposed a boundary of 16S rRNA gene sequence similarity of
97% for species delineation. This value, still largely used today for operational-taxonomic-
unit (OTU) based analysis of microbial communities, has been updated by Stackebrandt and
Ebers (2006) to a value between 98.7-99.0%, based on a greater amount of available sequence
data. Subsequently, as genome sequencing has become common, whole genome comparisons
became possible leading to the advent of genome relatedness indices such as the average
nucleotide identity (ANI), amino acid identity (AAI), and digital DDH. More recently, Kim *et*
al. (2014) proposed a 16S rRNA gene sequence similarity threshold value of 98.65% for
species delineation as it equated ANI values of 95-96%, which in turn have been equated to
the classical standard species delineation threshold DDH value of 70% (Goris *et al.*, 2007;
Richter and Rosselló-Móra, 2009). A method that only relies on protein-coding genes (i.e.
neither rRNA nor tRNA are included in analysis) is the Microbial Species Identifier (MiSI),
which employs both alignment fractions (AF) and ANI for demarcation of species,
recommending threshold AF:ANI values of 0.6:96.5% using complete or nearly complete
genomes (Varghese *et al.*, 2015).
Despite these advancements in resolving species delineation, practical guidelines that
incorporate genomic properties to demarcate genera have been lacking even though genus
assignment is key to performing meaningful comparisons regarding the physiology,
metabolism and genomic potential of microbes. Methods to demarcate genera have been
proposed that are based on either the amino acid identity (AAI) or the percentage of
conserved proteins (Konstantinidis and Tiedje, 2007; Qin *et al.*, 2014). However, these
methods directly rely on the 16S rRNA gene sequence, which is in some cases, insensitive to
evolutionary changes in the rest of the genome of a given organism, as revealed by different
species sharing >99% identity over the length of this gene. Additionally, the generally-used
arbitrary genus threshold of 95% 16S rRNA gene identity has been recently revisited to a
lower minimum value of 94.5% with confidence interval of 94.55-95.05 and median sequence
identity of 96.4% (Yarza *et al.*, 2014). In borderline cases, interpretation of results may be
unclear if there are no alternative ways to confirm genus assignment. This is also the case for
microorganisms with multiple, highly-divergent 16S rRNA genes. Here, we propose a novel
approach that is based on the application of the MiSI method (Varghese *et al.*, 2015) and
provide an objective and reproducible method of delimiting genera. We implement this
approach and method by testing a variety of taxonomic groups of *Bacteria* and *Archaea*.
Furthermore, supporting evidence is presented to support the reclassification of
Hydrogenovibrio halophilus and the extensive rearrangement of species within *Bacillus,*
including the human pathogens *B. anthracis* and *B. cereus*.
92

93   Materials and Methods
94
ANI and AF
ANI and alignment fraction (AF) values were obtained by the Microbial Species
Identifier (MiSI) method (https://ani.jgi-psf.org/html/; Varghese *et al.*, 2015), which is also
implemented in the JGI-Integrated Microbial Genomes (IMG) system
(https://img.jgi.doe.gov/cgi-bin/m/main.cgi) via its Pairwise ANI tool. ANI, as defined by
Varghese *et al.* (2015), is calculated for a pair of genomes by averaging the nucleotide
identity of orthologous genes identified as bidirectional best hits (BBHs), which are the genes
that show ≥70% sequence identity and ≥70% alignment of the shorter gene. AF, as defined by
Varghese *et al.* (2015), is calculated as a fraction of the sum of the lengths of BBH genes
divided by the sum of the lengths of all genes in a genome.
105
Strains
The Namesforlife Database (http://www.namesforlife.com; NamesforLife, LLC, East
Lansing, MI; Garrity, 2010) was primarily used to retrieve information about type strains and
type species associated with different genera. Complementary to this, equivalent strain
numbers assigned by different biological resource centers (e.g., DSMZ - German Collection
of Microorganism and Cell Cultures) were searched in their respective online catalogues. In
addition to the above, primary literature sources were used to confirm some of these
designations. All strain designations were cross-referenced in at least two databases.
114
Genomes and 16S rRNA gene sequences
Genomes were obtained from IMG or the National Center for Biotechnology
Information (NCBI; https://www.ncbi.nlm.nih.gov/). Generally, phyla with taxonomic
families/orders that have available genomes of ≥4 type species of genera and ≥4 non-type
species in a given genus were considered. Most of the genomes used in this study were
sequenced directly by the JGI (47%), National Institute of Technology and Evaluation
(https://www.nite.go.jp/; 5%), J. Craig Venter Institute (3%), and Broad Institute (3%). About
38% of the genomes included in this study were sequenced as part of the JGI-Genomic
Encyclopedia of *Bacteria* and *Archaea*, a project that focuses on sequencing the genomes of
type strains (Wu et al., 2009; Mukherjee et al., 2017). The rest of the genomes were
sequenced by >140 different institutions. Sequences of the 16S rRNA gene were obtained
from either IMG, NCBI, or EZBioCloud (http://www.ezbiocloud.net/). Alignments were
separately performed in the SILVA Incremental Aligner version 1.2.11 (https://www.arb-
silva.de/aligner/) and ClustalW via the Geneious platform (version R6; Biomatters, Auckland,
New Zealand). Genetic distances were calculated in Geneious. PhyML (Guindon and
Gascuel, 2003) was used via the Geneious platform to generate the maximum likelihood
phylogenetic tree with the following settings: Hasegawa-Kishino-Yano (HKY85) substitution
model, 1000 bootstraps, estimated transition/transversion ratio, estimated proportion of
invariable sites, estimated gamma distribution, branch lengths and substitution rate optimized.
134
Pangenome Analysis
Pangenome analysis of genomes from *Bacillus* and other genera in *Bacillaceae* were
processed in anvi'o (version 4)(Eren et al., 2015) following the workflow for microbial
pangenomics (merenlab.org/2016/11/08/pangenomics-v2/, last accessed 13 April, 2018;

Delmont and Eren, 2018). In brief, we generated contig databases for each genome contig file
using the command 'anvi-gen-contigs-database'. Prodigal (Hyatt et al., 2010) was used to
identify open reading frames and subsequently each database was populated with HMM
profiles by comparison to several single-copy gene collections (Campbell et al., 2011; Rinke
et al., 2013) using HMMER (Eddy, 2011). Once contig databases were generated for all
genomes, we used 'anvi-gen-genomes-storage' to generate a master genome storage database
to use in the pangenome analysis. We used the ncbi-blast option in 'anvi-pan-genome' to
calculate gene similarity and MCL (van Dongen and Abreu-Goodger, 2012) for clustering
under the following settings: minbit, 0.5; mcl inflation, 2; minimum occurrence, 2. Finally, to
organize both gene clusters and genomes, we used Euclidian distance and Ward linkage.
149
Statistical Analysis
All bidirectional, matched pair values of ANI (ANI1-2; ANI2-1) and AF (AF1-2;
AF2-1) were reported as single averaged ANI and AF values in this study. The nonparametric
Wilcoxon test was performed separately for each ANI and AF values to determine significant
($P<0.05$) differences between type- and non-type species. A discriminant multivariate analysis
was performed using quadratic fitting methods, which resulted in probabilities of group
assignment (i.e., type vs. non-type species). All statistical analyses were performed using the
statistical software JMP Pro, version 12 (SAS Institute Inc.).
158
Rationale
The nomenclatural type or "type" is defined by the International Code of
Nomenclature of Prokaryotes (ICNP) as "that element of the taxon with which the name is
permanently associated" (Parker et al., 2015). A type strain acts as the single reference for a
given species. Each validly published species has a type strain. A type species, which is
represented by its type strain, acts as the single reference for a given genus. Type strains and
type species do not necessarily have to be the most representative of the species or genus,
respectively. The rationale used in this study is that sister species in a given genus should be
relatively similar to the type species of the cognate genus indicating high similarity at the
genome level. Therefore, when a type species of a genus is compared to a non-type, sister
species of the same genus, the AF:ANI values should be relatively high. In contrast to this,
type species of different sister genera (i.e., in the same taxonomic order or family) should be
relatively dissimilar at the genome level. This dissimilarity should be reflected in relatively
low AF:ANI values.
173
Guidelines
In addition to the genome(s) under consideration (e.g., of a cultured strain that needs
classification), at least two genomes of type species in the same order or family are needed: a)
the primary reference which is the phylogenetically closest relative that is a type species (i.e.,
the type species that all other microbes will be compared to), b) all other type species (≥1) in
the same order or family. All available genomes of non-type, sister species that belong to the
same genus of the primary reference are needed. In order for the results to be meaningful, a
species must only be represented by the type strain unless its genome is not available, in
which case a non-type strain of the same species that is ≥99 % identical in 16S rRNA gene
sequence(s) (i.e., all copies of 16S rRNA gene in the genome) may be used. Accession
numbers for these genomes should be reported.

185
Interpretation
Every data point in the AF:ANI plot is calculated in relation to the primary reference
which is defined above as the type strain of the type species of the specific genus to be
analyzed; therefore, the plot represents genomic similarity to the reference genome. If the
primary reference is compared against an identical genome (e.g., or a direct comparison
against itself), an AF value of 1 and ANI value of 100 would result. If the primary reference is
compared against relatively dissimilar genomes, the AF and ANI values will reflect relatively
lower values. Thus, the plots that are generated effectively illustrate how distant other data
points are to the primary reference in terms of AF and ANI values, or in other words, how
distant other data points are to the upper-right corner (AF=1 and ANI=100).
When all comparisons are made in relation to the primary reference, a type species
should always cluster with other type species in the same taxonomic order or family,
reflecting relatively low AF and ANI values. A non-type, sister species should always cluster
with other non-type, sister species of the same genus, reflecting relatively high AF and ANI
values. The distinct clustering of type species and non-type, sister species should be
conserved without any overlap. If a non-type, sister species is positioned within the cluster of
type species in the same taxonomic order or family, then this indicates that the non-type, sister
species could potentially be assigned to a new genus, as it is as genomically different to the
primary reference as are other type species in the same taxonomic order or family. If a type
species is positioned within the cluster of non-type, sister species, then there is support for
potential reclassification of that species within the same genus of the primary reference, given
the accumulation of other supporting evidence (e.g., 16S rRNA gene identity and phenotypic
properties). It is emphasized that any proposed taxonomic rearrangement or reclassification
based on AF:ANI results will necessarily affect nomenclature; therefore, any corresponding
changes in nomenclature resulting from such proposal must be in alignment with the ICNP
(Parker et al., 2015).
212
Results and Discussion
Table 1 summarizes the 19 separate taxonomic groups that were investigated in this
study along with results of statistical tests for both AF and ANI. A total of 790 genomes
including 302 type species and 488 non-type species, representing 705 type strains and 10
phyla were used. For each taxonomic group, the combination of AF and ANI genomic indices
resulted in distinct clustering of type and non-type species. The means of these clusters were
significantly different in all tested cases with AF as a variable. In the majority of the cases
(17/19), the means of these clusters were also significantly different with ANI as a variable.
Discriminatory multivariate analyses, which give a sense of how separated the clusters are,
supported the distinct clustering of type and non-type species in 18 out of 19 cases with 91-
100% of the species correctly classified. A few representative cases will be discussed in more
detail below.
To initially test our approach with Gram-positive bacteria, the order *Lactobacillales*
was chosen, since it contains a number of well-characterized genera and species. A continuum
of AF:ANI values characterizes a typical result when intergeneric and intrageneric species of
the order *Lactobacillales* (e.g., all against all) are compared without distinguishing between
type and non-type strains or species (Figure 1A). Because there is no primary taxonomic
reference genome to compare to, differentiation between different taxonomic groups is not

possible. However, when the type species of *Lactococcus, L. lactis* subsp. *lactis* ATCC
19435[T], is used as the primary reference, a clear distinction can be made between type species
and non-type species (Figure 1B). The type species of genera in *Lactobacillales* form a
distinct cluster toward lower AF:ANI values while the non-type species of *Lactococcus*
cluster toward higher AF:ANI values (Table 1; Figure 1C-D). There is no overlap between
these two clusters. Discriminatory multivariate analysis supported the distinct clustering of
type and non-type species successfully classifying 98% of the samples.

To test our approach with Gram-negative microorganisms, type species of genera in
the order *Alteromonadales* and non-type *Shewanella* spp. were pairwise-compared to the type
species *S. putrefaciens* JCM 20190[T], which is the primary reference (Figure 2A; Table 1). A
distinct clustering is seen between the type species and non-type species. Discriminatory
multivariate analysis also supported the distinct clustering of type and non-type species
successfully classifying 100% of the samples.

Additional examples show that this trend is also seen with taxonomic families (Figure
2B; Table 1). The *Flavobacteriaceae* was investigated as an example of a diverse family with
one of the largest number of validly published genera (currently >160). Despite this, the
separate clustering of type species and non-type species was conserved. When the type
species of the genus *Flavobacterium, F. aquatile* LMG 4008[T] is used as the primary
reference, all type species (i.e. with available genomes) in the *Flavobacteriaceae* clustered
towards lower AF:ANI values, while the vast majority of the non-type species of
Flavobacterium clustered toward higher AF:ANI values. Discriminatory multivariate analysis
successfully classified 96% of the samples. The five non-type *Flavobacterium* spp. that were
outside the AF:ANI genus boundary had 16S rRNA gene sequence identities of <94% to the
primary reference. In a separate example, the type species *Vibrio cholerae* ATCC 14035[T] was
used as the primary reference and compared to non-type species in the genus *Vibrio* and other
type species of genera in the family *Vibrionaceae* (Figure 2C; Table 1). A distinct clustering,
without overlap, is seen between the type species and non-type species. Discriminatory
multivariate analysis also supported the distinct clustering of type and non-type species with
>91% of the samples successfully classified, even though the clusters were in much closer
proximity to each other than in previous examples.

This approach was also tested on *Methanosarcinaceae*, an archaeal taxonomic family
in the phylum *Euryarchaeota* (Figure 2D; Table 1). The primary reference was the type
species *Methanosarcina barkeri* JCM 10043[T]. The results were consistent with previous
examples on bacterial taxonomic groups displaying clustering of non-type species separately
from type species. Discriminatory multivariate analysis successfully classified 100% of the
samples. Similar results were obtained in the family *Haloferacaceae*. Additionally, the family
Thermoproteaceae in the phylum *Crenarchaeota* was tested with *Pyrobaculum islandicum*
DSM 4184[T] as a primary reference (Figure S1). A significant distinction between type and
non-type species was seen with AF (P=0.0200) but not with ANI (P=0.713). Despite this,
there was no overlap of the type/non-type clusters and discriminatory multivariate analysis
successfully classified 100% of the samples.

In a similar manner, genus assignment in the recently rearranged *Hydrogenovibrio-*
Thioalkalimicrobium-Thiomicrospira cluster was tested as an example of a bacterial
taxonomic group with historical taxonomic issues (Takai *et al*., 2004; Tourova *et al*., 2006;
Figure S2). Boden *et al.* (2017) provided a detailed evaluation of the taxonomy of this cluster
which belongs to the family *Piscirickettsiaceae* and proposed to place many of the species

into different genera. We tested these newly proposed assignments using the approach
described above and the MiSI method. When testing genus assignment to *Hydrogenovibrio*,
the type species *H. marinus* DSM 11271[T] was used as a primary reference (Figure 3). The
single data point that crossed the boundaries set by the AF:ANI values between type species
of genera in the family *Piscirickettsiaceae* and the non-type species of the genus
Hydrogenovibrio belonged to *Hydrogenovibrio halophilus* DSM 15072[T]. Aside from this data
point, there was no overlap of AF:ANI values between the type- and non-type species,
indicating support for the assignment of *H. crunogenus*, *H. thermophilus*, and *H. kuenenii* to
the genus *Hydrogenovibrio*. Similarly, the recently proposed new genus *Thiomicrorhabdus*
(Boden *et al*., 2017) was also tested (Figure S3). The results indicate that the new
classification is supported by AF and ANI values as the type species of genera in the family
Piscirickettsiaceae and *Thiomicrorhabdus* spp. remain well separated. The reclassification of
Thioalkalimicrobium spp. as *Thiomicrospira* spp. (Boden *et al*., 2017) is also supported by
ANI and AF values (Figure S4), as the *Thioalkalimicrobium* spp. (now non-type
Thiomicrospira spp.) cluster away from the type species of genera in the family
Piscirickettsiaceae.
Our approach has identified a potential misclassification of *H. halophilus* (basonym
Thiomicrospira halophila) because it clusters with the type species of genera in the family
Piscirickettsiaceae, other than *Hydrogenovibrio*. Below we show additional evidence that
could support a potential reclassification of *H. halophilus*. A closer look at the multiple 16S
rRNA gene sequences of *H. halophilus* revealed that they have 94.0-94.2% pairwise-identities
to the type species, *H. marinus* DSM 11271[T]. Our results differ from those of Boden *et al*.,
who reported 95.7 % sequence identity (2017) between *H. halophilus* and *H. marinus*. The
difference in identity appears to be due to differences in the 16S rRNA gene sequence
between the near-full-length version of Sorokin *et al*. (2006; 1420 bp; accession number
DQ390450) and the full-length versions of this gene (1439 bp each; IMG ID 2518265101 and
2518265550) originating from a draft genome (Scott *et al.,* 2018) and used in this study. The
phylogenetic placement of this organism also supports the idea that *H. halophilus* is distinctly
positioned in its own clade as it branches away from the cluster of other *Hydrogenovibrio* spp.
(Figure 3B). One differentiating characteristic of *H. halophilus* is the DNA G+C content of
56.6%, much higher than the ca. 44% of the other newly classified *Hydrogenovibrio* spp.
Another major differentiating aspect is the higher NaCl optimum/maximum of 1.5/3.5 M for
H. halophilus versus the lower 0.2-0.5/0.6-1.2 M for other *Hydrogenovibrio* spp. Therefore,
based on AF and ANI results, molecular, phylogenetic, and physiological evidence, *H.*
halophilus could potentially be assigned to a new genus.
The *Bacillaceae* was also investigated as an example of taxonomic family that is
medically relevant (Figure 4A). Generally, there were significant differences between non-
type *Bacillus* species and type species of genera in the family *Bacillaceae* when *B. subtilis*
6051[T] (Kabisch et al., 2013) was used as a primary reference (Table 1; Figure S5). However,
a large number of non-type *Bacillus* species clustered with type species (50% misclassified by
multivariate, discriminatory method), which strongly suggests that the genus *Bacillus* is in
need of an extensive and thorough rearrangement. Of note, it is highlighted that human
pathogens *B. anthracis* and *B. cereus* cluster with type species of genera in *Bacillaceae*,
indicating that they are at least as genomically different to *B. subtilis* as are other type species
in this family, warranting a taxonomic rearrangement. As shown above with *H. halophilus*, a
similar case could be made for the taxonomic rearrangement of each one of these species

(e.g., <94.5% identity in 16S rRNA gene sequence to the type species *B. subtilis* 6051[T]).
Compared to our analyses of other taxonomic groups, these results are atypical in the sense
that the non-type species cluster overlaps considerably with the type species cluster. We
highlight the fact that a large number of these specimens were designated non-type species of
Bacillus prior to the advent of the universal use of 16S rRNA gene as a taxonomic marker in
the 1990s (i.e., >20% of all type strains of non-type *Bacillus* with available genomes in our
dataset). For example, *B. anthracis* was first described in 1872 (Cohn). When the primary
reference is changed to *Anoxybacillus pushchinoensis* K1[T], a type species of a genus
established in 2000 (Pikuta *et al.*) using the taxonomic framework of the 16S rRNA gene, the
clustering of type species of genera in *Bacillaceae* and non-type species of *Anoxybacillus* is
distinct and without overlap (Table 1).
The AF:ANI plots reveal different rates of differentiation at the DNA level. In the few
cases where there are sufficient samples to analyze, and values are evenly distributed, the
AF:ANI plots reveal a polynomial shape approximating a quartic function, with R^2 values
>0.97. A second derivative of this function results in a quadratic function that can be used to
determine inflection points with precision. Estimates based on several examples in this study
indicate two inflection points. The first inflection point at 71-74% ANI, generally
approximates the genus demarcation line in all cases herein presented based on taxonomic
affiliation. However, the distribution of non-type, sister species a given genus generally
includes values that fall below this inflection point at lower AF:ANI values, suggesting a
potential underestimation of genera diversity within a taxonomic order or family. Estimation
of this genus inflection point via Logistic and Gompertz functions generally agree with the
estimations by quartic function and their use is recommended for accuracy. The second
inflection point at 89-90% ANI, is consistently 5-7% lower than current ANI species
thresholds (95-96.5% ANI), and is only seen with the quartic function. Because the second
inflection point potentially deals with species delineation, detailed exploration (e.g., using
subspecies type strains) and discussion of this topic are outside of the scope of this
manuscript. However, it is noted that ANI values of 89-90% largely correspond to 16S rRNA
gene identity values of ≥98.65 (Figure 3 in Kim et al., 2014).
The genus inflection point serves as practical guide for maximum AF:ANI values of
genus boundary and for the identification of a genus transition zone. Ideally, and assuming
abundance of type strains genomes and consequently data points, the estimated genus
inflection point should closely match the genus boundary. As an example, three different
functions were used to estimate the genus inflection point in *Bacillus* (Figure 4B). The
obtained values give an estimation of this inflection point at AF=0.39-0.45 and ANI=72.71-
73.80. An analysis of the pangenome of *Bacillus* using the most conservative estimated genus
inflection point as a guide shows that only non-type *Bacillus* spp. above the genus inflection
point cluster with the type species *B. subtilis* (the primary reference) while *B. anthracis* and *B.*
cereus cluster with type species of other genera in *Bacillaceae* (Figure 5), corroborating, as
pointed out above, that these two pathogens warrant reclassification and renaming. However,
doing so would require considerable care so as not to raise objections over safety issues and a
push for conservation of the current names over newly proposed names (Parker *et al.*, 2015).
These results were further corroborated in phylogenomic analyses of *Bacillaceae* taking into
account either 138 single copy genes or only ribosomal genes (Figures S13-S14).
It is important to note that the genus inflection point is *estimated* based on nonlinear
regression and only serves as a guide for future taxonomic designation by providing

maximum AF:ANI values. This is particularly important in the case of a new taxonomic
designation with a genome displaying AF:ANI values that are right above the effective genus
boundary, in a transition zone; is it a novel type species or a novel non-type species? Having
an estimated inflection point helps in making this decision. Thus far, this could be properly
calculated only for a few taxonomic groups due to lack of genomes of type strains and/or
actual isolates. *B. pumilus*, *B. safensis*, *B. invictae*, *B. altitudinis*, *B. zhangzhouensis*, *B.*
xiamenensis, and *B. australimaris* are borderline species positioned within the limits of the
most and least conservative estimated genus inflection points. Their 16S rRNA gene identities
in relation to the primary reference range from 96.8% to 97.4%, suggesting that the genus
inflection point of *Bacillus* approximates a value that is much higher than the current genus
threshold based on 16S rRNA gene identity (94.5%).
The approach we have used to demarcate genera is complementary and does not
replace the conventional polyphasic method of designating a genus, which includes thorough
analyses of the full-length 16S rRNA gene, phylogeny, physiology and metabolism, among
other aspects. The AF:ANI boundary values for genus demarcation in a given order or family
can be refined as more type strains and type species are genomically sequenced and formally
described. This approach emphasizes the importance of type strains and type species in the
evaluation of taxonomy with genomic indices (e.g., ANI or AF), as these specimens function
as reference points for their respective taxonomic groups. In this respect, our approach is in
line with recommendations regarding taxonomy, including relevant comparisons to type
strains and type species for the characterization of novel *Bacteria* and *Archaea* (Tindall *et al.*,
2010). It is reiterated that any rearrangement or reclassification of taxa should be in alignment
with the ICNP (Parker *et al.*, 2015). In particular, a potential rearrangement of type species
must be carefully analyzed as a genus can only contain one type species, and rules and
principles of nomenclature must be followed in order to properly do that in accordance with
the ICNP.
Uncultivated *Bacteria* and *Archaea* represented by metagenomic assembled genomes
(MAGs) and/or single cell amplified genomes (SAGs) could potentially be used in the
implementation of this approach; however, it is noted that type strains and type species should
be used as references for such analyses, if genomes are available (see *Guidelines*). The
designation of type strains and type species is governed by the ICNP and require a thorough
characterization of the microorganism and the deposit of the type strain in at least two
different culture collections. Naturally, the proper characterization, and therefore the type
designation is, in most cases, impossible for uncultivated *Bacteria* and *Archaea* given the
current guidelines set by the ICNP. This could change if genomes are subsequently
considered type material, as recently proposed (Whitman, 2015) and discussed
(Konstantinidis *et al.*, 2017). In lieu of type strains and type species, proper and consistent
genomic references must be established in order to analyze taxonomic groups without
cultivated representatives. Caution is advised in the interpretation of data resulting from
MAGs since they can represent composite genomes of different strains. In theory, SAGs
could serve as good taxonomic frames of reference for uncultivated genera; however, genome
incompleteness in SAGs is an issue that would need to be addressed.
The MiSI method explicitly excludes tRNA and rRNA genes in order to avoid
inflation of AF or ANI values. However, the current taxonomic framework is largely based on
the use of the 16S rRNA gene for classification. Therefore, the approach herein investigated is
not completely independent of the 16S rRNA gene. The fact that in all the groups tested, with

the exception of the *Bacillaceae* (note: although significant differences between type and non-type species were still detected for this group), there was strong concordance of type species of genera being delineated from non-type species of the primary reference provides evidence to the success of the use of the 16S rRNA gene within the taxonomic framework. The 16S rRNA gene has served and is continuing to serve the scientific community well, especially in maintaining consistency in classifications. However, our estimation of inflection points indicates that the 16S RNA gene identity minimum threshold value of 94.5% could underestimate genera diversity in some taxa [e.g. *Bacillaceae (Bacillus)*] and overestimate it in others [e.g. *Enterobacterales (Xenorhabdus)*], especially if multiple non-identical 16S rRNA genes are present in the genomes. It is clear that different taxonomic groups are characterized by different AF:ANI threshold values for genus demarcation. However, even within established taxonomic groups, the effective AF:ANI threshold values will still be "moving targets", dependent on the assignment and re-assignment of new type species. Thus, a single, universal, stationary threshold for genus delineation will not be able to sensitively resolve genus assignments for all taxa.

The reader is made aware that variations in bidirectional, matched pair AF values (AF1-2; AF2-1) exist. These variations are usually small, but could be relevant in the interpretation of data, especially in borderline cases. Bidirectional, matched pair ANI values (ANI1-2; ANI2-1) differ by <0.1 % in the majority of the cases, with a few cases differing by <0.3%, which would make the error bars invisible to the naked eye. Figures with corresponding standard deviation values for AF are included in a supplemental file (Figure S6-S12).

The use of ANI has been previously stated as not applicable for the demarcation of genera (Qin *et al*, 2014; Kim *et al*., 2014). Our results indicate that the MiSI method, and therefore AF and ANI, can be used to visualize natural breakpoints that can be used to circumscribe genera, if the guidelines presented in this study are followed. Adaptation of this method to demarcate higher taxonomic ranks, including phyla, has not been tested as it is beyond the scope of this study. Nonetheless, such an approach would be warranted. Finally, we highlight that the results demonstrate a conserved genomic coherence at the genus level for numerous different taxa, shedding light on a fundamental emergent property of *Bacteria* and *Archaea*, a property that has proven largely elusive at the species level.

Table 1. Summary of taxonomic groups and genomes included in this study. Gray fill-color indicates if either the taxonomic family or order was investigated. For a detailed list including accession numbers, see Table S1.

Domain	Phylum	Class	Order*	Family*	Genus	Primary reference	# of Type Species of Genera§	# of Non-Type Species	Wilcoxon Test (T vs. NT) AF (P-value)	ANI (P-value)	Discriminatory Test % Correctly Classified
Archaea	Crenarchaeota	Thermoprotei	Thermoproteales	Thermoproteaceae	Pyrobaculum	P. islandicum DSM 4184^T	5	5	0.0200	0.7130	100
Archaea	Euryarchaeota	Halobacteria	Haloferacales	Haloferacaceae	Haloferax	H. volcanii DS2^T	9	10	0.0004	0.0004	100
Archaea	Euryarchaeota	Methanococci	Methanosarcinales	Methanosarcinaceae	Methanosarcina	M. barkeri JCM 10043^T	7	9	0.0016	0.0018	100
Bacteria	Actinobacteria	Actinomycetes	Microbacteriales	Curtobacteriaceae	Curtobacterium	C. citreum NS330$^#$	23	4	0.0020	0.0020	100
Bacteria	Bacteroidetes	Flavobacteria	Flavobacteriales	Flavobacteriaceae	Flavobacterium	F. aquatile LMG 4008^T	70	83	<0.0001	<0.0001	96
Bacteria	Bacteroidetes	Flavobacteria	Flavobacteriales	Flavobacteriaceae	Maribacter	M. sedimenticola DSM 19840^T	70	12	<0.0001	<0.0001	100
Bacteria	Deinococcus-Thermus	Deinococci	Thermales	Thermaceae	Thermus	T. aquaticus YT-1^T	4	12	0.0113	0.0115	100
Bacteria	Firmicutes	Bacilli	Bacillales	Bacillaceae	Anoxybacillus	A. pushchinoensis K1^T	27	8	<0.0001	<0.0001	97
Bacteria	Firmicutes	Bacilli	Bacillales	Bacillaceae	Bacillus	B. subtilis 6051-HGW^T	"	110	<0.0001	<0.0001	50
Bacteria	Firmicutes	Bacilli	Lactobacillales	Streptococcaceae	Lactococcus	L. lactis ATCC 19435^T	38	6	0.0001	0.0001	97
Bacteria	Fusobacteria	Fusobacteria	Fusobacteriales	Leptotrichiaceae	Streptobacillus	S. moniliformis DSM 12112^T	9	3	0.0184	0.0189	100
Bacteria	Fusobacteria	Fusobacteria	Fusobacteriales	Fusobacteriaceae	Fusobacterium	F. nucleatum ATCC 25586^T	"	9	0.0081	0.0081	100
Bacteria	Planctomycetes	Planctomycetia	Planctomycetales	Planctomycetaceae	Rhodopirellula	R. baltica SH 1^T	13	2	0.0358	0.0354	100
Bacteria	Proteobacteria	Gammaproteobacteria	Alteromonadales	Shewanellaceae	Shewanella	S. putrefaciens JCM 20190^T	20	23	<0.0001	<0.0001	100
Bacteria	Proteobacteria	Deltaproteobacteria	Desulfuromonadales	Geobacteraceae	Geobacter	G. metallireducens GS-15^T	6	12	0.0019	0.0019	100
Bacteria	Proteobacteria	Gammaproteobacteria	Enterobacteriales	Morganellaceae	Xenorhabdus	X. nematophila ATCC 19061^T	46	10	<0.0001	<0.0001	100
Bacteria	Proteobacteria	Gammaproteobacteria	Vibrionales	Vibrionaceae	Vibrio	V. cholerae ATCC 14035^T	6	65	0.0005	0.0005	91
Bacteria	Proteobacteria	Gammaproteobacteria	Thiotrichales	Piscirickettsiaceae	Hydrogenovibrio	H. marinus DSM 11271^T	11	4	0.013	0.0089	100
Bacteria	Spirochaetes	Spirochaetia	Spirochaetales	Spirochaetaceae	Leptospira	L. interrogans ATCC 43642^T	6	19	0.0008	0.1768	100

* = shading indicates the taxonomic resolution at which the group was analyzed.

§ = including the primary reference.

= >99% identity to 16S rRNA gene sequence of C. citreum DSM 20528^T, which does not have a sequenced genome.

Figure 1. AF:ANI pairwise genome comparisons in the order *Lactobacillales*. A. Type and non-type strains/species of 15 different genera in the order *Lactobacillales* were pairwise-compared (n=6090 comparisons). B. Type species (n=37) of genera within the order *Lactobacillales* (circles) and non-type species (n=6) of the genus *Lactococcus* (in order *Lactobacillales*; triangles) were pairwise-compared *only* to the primary reference, type species *Lactococcus lactis* subsp. *lactis* ATCC 19435[T]. The bottom-left quadrant demarcates the boundary between type and non-type species. Boxplot diagrams of AF (C) and ANI (D) values as related to non-type species (NT) of the genus *Lactococcus* and type species (T) of genera within the order *Lactobacillales*. Means are shown in green. Standard deviations are shown in blue.

478
Figure 2. Pairwise genome comparisons to the following primary references: a) type species
of the genus *Shewanella*: *S. putrefaciens* JCM 20190[T]. In circles: type species (n=19) of
genera within the order *Alteromonadales*. In triangles: non-type species (n=23) of the genus
Shewanella; b) type species of the genus *Flavobacterium*: *F. aquatile* LMG 4008[T]. In circles:
type species (n=69) of genera within the family *Flavobacteriaceae*. In triangles: non-type
species (n=83) of the genus *Flavobacterium*. Inset: zoomed-in boxed area; c) type species of
the genus *Vibrio*: *V. cholerae* ATCC 14035[T]. Circles: type species (n=5) of genera within the
family *Vibrionaceae*. Triangles: non-type *Vibrio* spp. (n=65). Inset: zoomed-in boxed area; d)
type species of the genus *Methanosarcina*: *M. barkeri* JCM 10043[T]. Circles: type species
(n=6) of genera within the family *Methanosarcinaceae*. Triangles: non-type *Methanosarcina*

spp. (n=9). In all cases, the bottom-left quadrant demarcates the boundary between type and
non-type species. Boxplots of AF (center panel) and ANI (right panel) values as related to
non-type (NT) species of genus analyzed and type species (T) of genera within the order or
family analyzed. Means are shown in green. Standard deviations are shown in blue.
493
494

495
496
Figure 3. Analysis of *Hydrogenovibrio*, as rearranged by Boden *et al.* (2007). A. Pairwise
genome comparisons to the primary reference *Hydrogenovibrio marinus* DSM 11271[T]. Non-
type species within the genus *Hydrogenovibrio* are shown in triangles. The type strain *H.*
thermophilus I78[T] does not have a sequenced genome; therefore, *H.* sp. MA2-6 (>99%
pairwise identity in 16S rRNA gene sequence) was used instead. Similarly, *H. crunogenus*
XCL-2 (>99% pairwise identity in 16S rRNA gene sequence) is used instead of the type strain
H. crunogenus TH-55[T], as the latter does not have a sequenced genome. Type species of
genera within the family *Piscirickettsiaceae* are shown in circles. The bottom-left quadrant
demarcates the boundary between type and non-type species. B. Maximum-likelihood
phylogenetic tree based on an alignment of 16S rRNA genes indicating the phylogenetic
positioning of *H. halophilus* DSM 15072[T]. Type species of genera are shown in bold. The
scale bar indicates 20% sequence divergence.

522
523
Figure 4. Pairwise genome comparisons to the type species of the genus *Bacillus*: *B. subtilis*
ATCC 6051[T]. Circles: type species (n=26) of genera within the family *Bacillaceae*. Triangles:
non-type *Bacillus* spp. (n=110). a) The bottom-left quadrant (black dashed-line) demarcates
the boundary between type and non-type species; inset: zoomed-in boxed area. b) Estimated
inflection points (colored dashed-lines) for *Bacillus* based on quartic (blue), logistic 4P (red),
and Gompertz 4P (green) functions. The quartic function is forced to go through the AF:ANI
value of 1:100.

557
558
Figure 5. Pangenome of *Bacillus* (in blue). Type species of genera in the family *Bacillaceae*
(i.e., other than *B. subtilis* ATCC 6051[T]) that are relatively close to the genus demarcation
AF:ANI threshold (see Figure 4A) are shown in orange. For this pangenomic view, only
Bacillus spp. with AF:ANI values above an estimated genus inflection point (i.e.,
>0.39:72.71; Figure 4B) are shown, with exception of *B. anthracis* (0.18:69.8) and *B. cereus*
(0.17:70.0). ANI and AF values were obtained via pairwise genome comparisons to the
primary reference, *B. subtilis* ATCC 6051[T] (in purple). Hierarchical clustering was performed
on the presence-absence of gene clusters using Euclidean distance and Ward linkage. * =
Bacillus spp. above the estimated genus inflection point; SCG = single copy genes.

568
Acknowledgements

We would like to thank Anna Ratner at JGI for updating the interface of the Pairwise ANI
tool on the IMG platform. We would also like to thank Trudy Wassenaar and Nikos Kyrpides

for comments and suggestions on this manuscript. RAB was supported by the National
Science Foundation postdoctoral research fellowship in Biology (award #1523639). JJS was
supported by the Gordon and Betty Moore Foundation through a grant to the Smithsonian
Tropical Research Institute (GBMF5603).

577
578
579   References:
580

581   Boden, R., Scott, K.M., Williams, J., Russel, S., Antonen, K., Rae, A.W., and Hutt, L.P.
582   (2017) An evaluation of *Thiomicrospira*, *Hydrogenovibrio* and *Thioalkalimicrobium*:
583   reclassification of four species of *Thiomicrospira* to each *Thiomicrorhabdus* gen. nov. and
584   *Hydrogenovibrio*, and reclassification of all four species of *Thioalkalimicrobium* to
585   *Thiomicrospira*. *Int. J. Syst. Evol. Microbiol*. 67:1140-1151. doi:10.1099/ijsem.0.001855
586
587   Campbell, B.J., Yu, L., Heidelberg, J.F., and Kirchman, D.L. (2011) Activity of abundant and
588   rare bacteria in a coastal ocean. *Proc. Natl. Acad. Sci. USA*. 108(31):12776-12781.
589   doi:10.1073/pnas.1101405108
590
591   Cohn, F. (1872) Untersuchungen über Bakterien. *Beitr. Biol. Pflanz*. 1:127-224.
592
593   Eren, A.M., Esen, Ö.C., Quince, C., Vineis, J.H., Morrison, H.G., Sogin, M.L., Delmont, T.O.
594   (2015) Anvi'o: an advanced analysis and visualization platform for 'omics' data. *PeerJ*.
595   3:e1319.
596
597   Eddy, S.R. (2011) Accelerated profile HMM searches. *PLoS Comp. Biol*. 7(10):e1002195.
598   doi:10.1371/journal.pcbi.1002195.
599
600   Delmont, T.O. and Eren, A.M. (2018) Linking pangenomes and metagenomes: the
601   Prochlorococcus metapangenome. *PeerJ*. 6:e4320.
602
603   Garrity, G.M. (2010) NamesforLife: BrowserTool takes expertise out of the database and puts
604   it right in the browser. *Microbiology Today*. 2(2):9.
605
606   Garrity, G.M. (2016) A new genomics-driven taxonomy of Bacteria and Archaea: are we
607   there yet? *J. Clin. Microbiol*. 54:1956-1963. doi:10.1128/JCM.00200-16
608
609   Goris, J., Konstantinidis, K.T., Klappenbach, J.A., Coenye, T., Vandamme, P. and Tiedje,
610   J.M. (2007) DNA-DNA hybridization values and their relationship to whole-genome
611   sequence similarities. *Int. J. Syst. Evol. Microbiol*. 57:81-91. doi:10.1099/ijs.0.64483-0
612
613   Guindon, S. and Gascuel, O. (2003) A simple, fast, and accurate algorithm to estimate large
614   phylogenies by maximum likelihood. *Systematic Biology*. 52(5):696-704.
615
Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010)
Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*
Bioinformatics. 11:119. doi:10.1186/1471-2105-11-119

Kabisch, J., Thürmer, A., Hübel, T., Popper, L., Daniel, R., Schweder, T. (2013) Journal of
Biotechnology. 163(2):91-04. doi:10.1016/j.jbiotec.2012.06.034

Kim, M., Oh, H-S., Park, S-C., and Chun, J. (2014) Towards a taxonomic coherence between
average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation
of prokaryotes. *Int. J. Syst. Evol. Microbiol*. 64:346-351. doi:10.1099/ijs.0.059774-0

Konstantinidis, K.T. and Tiedje, J.M. (2005) Towards a genome-based taxonomy for
prokaryotes. *J. Bacteriol.* 187(18):6258-6264. doi:10.1128/JB.187.18.6258-6264.2005

Konstantinidis, K.T. and Tiedje, J.M. (2007) Prokaryotic taxonomy and phylogeny in the
genomic era: advancements and challenges ahead. *Current Opinion in Microbiology*. 10:504-
509.

Konstantinidis, K.T., Rosselló-Móra, R., and Amann, R. (2017) Uncultivated microbes in
need of their own taxonomy. *ISME J.* 11:2399-2406.

Markowitz, V.M., Chen, I.M., Chu, K., Pati, A., Ivanova, N.N., and Kyrpides, N.C. (2015)
Ten years of maintaining and expanding a microbial genome and metagenome analysis
system. *Trends in Microbiology*. 23(11):730-741. doi:10.1016/j.tim.2015.07.012

Mukherjee, S., Seshadri, R., Varghese, N.J., Eloe-Fadrosh, E.A., Meier-Kolthoff, J.P., Goker,
M., Coates, R.C., Hadjithomas, M., Pavlopoulos, G.A., Paez-Espino, D., Yoshikuni, Y.,
Visel, A., Whitman, W.B., Garrity, G.M., Eisen, J.A., Hugenholtz, P., Pati, A., Ivanova, N.N.,
Woyke, T., Klenk, H.-P., and Kyrpides, N.C. (2017) 1,003 reference genomes of bacterial and
archaeal isolates expand coverage of the tree of life. *Nature Biotechnology*. 35:676-683.
doi:10.1038/nbt.3886

Parte, A.C. (2014) LPSN — list of prokaryotic names with standing nomenclature. *Nucleic
Acids Research*. 42:D613-D616. doi:10.1093/nar/gkt1111

Parker, C.T., Tindall, B.J., Garrity, G.M. (2015) International Code of Nomenclature of
Prokaryotes. *Int. J. Syst. Evol. Microbiol*. doi:10.1099/ijsem.0.000778

Pikuta, E., Lysenko, A., Chuvilskaya, N., Mendrock, U., Hippe, H., Suzina, N., Nikitin, D.,
Osipov, G., Laurinavichius, K. (2000) *Anoxybacillus pushchinensis* gen. nov., sp. nov., a
novel anaerobic, alkaliphilic, moderately thermophilic bacterium from manure, and
description of *Anoxybacillus flavithermus* comb. nov. *Int. J. Syst. Evol. Microbiol.* 50:2109-
2117. doi:10.1099/00207713-50-6-2109

Pruesse, E., Peplies, J. and Glöckner, F.O. (2012) SINA: accurate high-throughput multiple
sequence alignment of ribosomal RNA genes. *Bioinformatics.* 28:1823-1829.

Qin, Q., Xie, B., Zhang, X., Chen, X., Zhou, B., Zhou, J., Oren, A., Zhang, Y. (2014) A
proposed genus boundary for the Prokaryotes based on genomic insights. *Journal of*
Bacteriology. 196(12): 2210-2215. doi:10.1128/JB.01688-14
666

667   Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and
668   Glöckner, F.O. (2013) The SILVA ribosomal RNA gene database project: improved data
669   processing and web-based tools. *Nucleic Acids Res*. 41:D590-D596.
670

671   Richter, M. and Rosselló-Móra, R. (2009) Shifting the genomic gold standard for the
672   prokaryotic species definition. *Proc. Natl. Acad. Sci. USA*. 106(45):19126-19131.
673   doi:10.1073/pnas.0906412106
674

675   Rinke, C., Schwientek, P., Sczyrba, A., Ivanova, N.N., Anderson, I.J., Cheng, J.-F., Darling,
676   A., Malfatti, S., Swan, B.K., Gies, E.A., Dodsworth, J.A., Hedlund, B.P., Tsiamis, G., Sievert,
677   S.M., Liu, W.-T., Eisen, J.A., Hallam, S.J., Kyrpides, N.C., Stepanauskas, R., Rubin, E.M.,
678   Hugenholtz, P., and Woyke, T. (2013) Insights into the phylogeny and coding potential of
679   microbial dark matter. *Nature*. 499:431-437. doi:10.1038/nature12352
680

Scott, K.M., Williams, J., Porter, C.M.B., Russel, S., Harmer, T.L., Paul, J.H., Antonen,
K.M., Bridges, M.K., Camper, G.J., Campla, C.K., Casella, L.G., Chase, E., Conrad, J.W.,
Cruz, M.C., Dunlap, D.S., Duran, L., Fahsbender, E.M., Goldsmith, D.B., Keeley, R.F.,
Kondoff, M.R., Kussy, B.I., Lane, M.K., Lawler, S., Leigh, B.A., Lewis, C., Lostal, L.M.,
Marking, D., Mancera, P.A., McClenthan, E.C., McIntyre, E.A., Mine, J.A., Modi, S., Moore,
B.D., Morgan, W.A., Nelson, K.M., Nguyen, K.N., Ogburn, N., Parrino, D.G., Pedapudi,
A.D., Pelham, R.P., Preece, A.M., Rampersad, E.A., Richardson, J.C., Rodgers, C.M.,
Schaffer, B.L., Sheridan, N.E., Solone, M.R., Staley, Z.R., Tabuchi, M., Waide, R.J.,
Wanjugi, P.W., Young, S., Clum, A., Daum, C., Huntemann, M., Ivanova, N., Kyrpides, N.,
Mikhailova, N., Palaniappan, K., Pillay, M., Reddy, T.B.K., Shapiro, N., Stamatis, D.,
Varghese, N., Woyke, T., Boden, R., Freyermuth, S.K., Kerfeld, C.A. (2018) Genomes of
ubiquitous marine and hypersaline *Hydrogenovibrio*, *Thiomicrorhabdus* and *Thiomicrospira*
spp. encode a diversity of mechanisms to sustain chemolithoautotrophy in heterogeneous
environments. *Environ. Microbiol*. doi:10.1111/1462-2920.14090
695

696   Stackebrandt, E. and Goebel, B.M. (1994) Taxonomic note: a place for DNA-DNA
697   reassociation and 16S rRNA sequence analysis in the present species definition in
698   Bacteriology. *Int. J. Syst. Bacteriol*. 44(4):846-849.
699

700   Stackebrandt, E. and Ebers, J. (2006) Taxonomic parameters revisited: tarnished gold
701   standards. *Microbiol. Today*. 33:152-155.
702

703   Sorokin, D.Y., Tourova, T.P., Kolganova, T.V., Spiridonova, E.M., Berg, I.A., and Muyzer,
704   G. (2006) *Thiomicrospira halophila* sp. nov., a moderately halophilic, obligately
705   chemolithoautotrophic, sulfur-oxidizing bacterium from hypersaline lakes. *Int. J. Syst. Evol.*
706   *Microbiol*. 56:2375-2380. doi:10.1099/ijs.0.64445-0
707

708 Takai, K., Hirayama, H., Nakagawa, T., Suzuki, Y., Nealson, K.H., and Horikoshi, K. (2004)
709 *Thiomicrospira thermophila* sp. nov., a novel microaerobi, thermotolerant, sulfur-oxidizing
710 chemolithomixotroph isolated from a deep-sea hydrothermal fumarole in the TOTO caldera,
711 Mariana Arc, Western Pacific. *Int. J. Syst. Evol. Microbiol.* 54:2325-2333.
712 doi:10.1099/ijs.0.63284-0
713
714 Tindall, B.J., Rosselló-Móra, R., Busse, H.-J., Ludwig, W. and Kämpfer, P. (2010) Notes on
715 the characterization of prokaryote strains for taxonomic purposes. *Int. J. Syst. Evol.*
716 *Microbiol.* 60:249-266.
717
718 Tourova, T.P., Spiridonova, E.M., Berg, I.A., Kuznetsov, B.B., and Sorokin, D.Y. (2006)
719 Occurrence, phylogeny and evolution of ribulose-1,5-bisphosphate carboxylase/oxygenase
720 genes in obligately chemolithoautotrophic sulfur-oxidizing bacteria of the genera
721 *Thiomicrospira* and *Thioalkalimicrobium*. *Microbiology*. 152:2159-2169.
722 doi:10.1099/mic.0.28699-0
723
724 van Dongen, S. and Abreu-Goodger, C. (2012) Using MCL to extract clusters from networks.
725 *Methods Mol. Biol.* 804:281-295. doi:10.1007/978-1-61779-361-5_15
726
727 Varghese, N.J., Mukherjee, S., Ivanova, N., Konstantinidis, K.T., Mavrommatis, K.,
728 Kyrpides, N.C., and Pati, A. (2015) Microbial species delineation using whole genome
729 sequences. *Nucleic Acids Research*. 43(14): 6761-6771. doi:10.1093/nar/gkv657
730
731 Verslyppe, B., De Smet, W., De Baets, B., De Vos, P., and Dawyndt, P. (2014) StrainInfo
732 introduces electronic passports for microorganisms. *Syst. Appl. Microbiol.* 37(1):42-50.
733 doi:10.1016/j.syapm.2013.11.002
734
735 Whitman, W.B. (2015) Genome sequences as the type material for taxonomic descriptions of
736 prokaryotes. *Syst. Appl. Microbiol*. 38:217-222.
737
Wu, D., Hugenholtz, P., Mavromatis, K., Pukall, R., Dalin, E., Ivanova, N.N., Kunin, V.,
Goodwin, L., Wu, M., Tindall, B.J., Hooper, S.D., Pati, A., Lykidis, A., Spring, S., Anderson,
I.J., D'haeseleer, P., Zemla, A., Singer, M., Lapidus, A., Nolan, M., Copeland, A., Han, C.,
Chen, F., Cheng, J.-F., Lucas, S., Kerfeld, C., Lang, E., Gronow, S., Chain, P., Bruce, D.,
Rubin, E.M., Kyrpides, N.C., Klenk, H.-P., and Eisen, J.A. (2009) A phylogeny-driven
genomic encyclopaedia of Bacteria and Archaea. *Nature*. 462(7276):1056-1060.
doi:10.1038/nature08656
745
746 Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F.O., Ludwig, W., Schleifer, K., Whitman,
747 W.B., Euzéby, J., Amann, R., and Móra, R.R. (2014). Uniting the classification of cultured
748 and uncultured bacteria and archaea using 16S rRNA gene sequences. *Nature Reviews*
749 *Microbiology*. 12: 635-645. doi:10.1038/nrmicro.3330
750
751 Yoon, S. H., Ha, S. M., Kwon, S., Lim, J., Kim, Y., Seo, H. and Chun, J. (2017). Introducing
752 EzBioCloud: A taxonomically united database of 16S rRNA and whole genome
753 assemblies. *Int. J. Syst. Evol. Microbiol*. 67(5):1613-1617. doi:10.1099/ijsem.0.001755.

755   Supplemental Section
756

757
758
759   Figure S1. Pairwise genome comparisons to the type species of the genus *Pyrobaculum*: *P.*
760   *islandicum* DSM 4184[T]. Circles: type species (n=4) of genera within the family
761   *Thermoproteaceae*. Triangles: non-type *Pyrobaculum* spp. (n=5).
762

763
764
Figure S2. Pairwise genome comparisons to the type species *Thiomicrospira pelophia* DSM
1534[T] using the old classification scheme (*i.e.*, prior to rearrangement by Boden *et al.*, 2017)
show clear taxonomic issues and no distinct clustering between type and non-type species.
Non-type species within the genus *Thiomicrospira* are shown in triangles. Type species of
genera within the family *Piscirickettsiaceae* are shown in circles. Inset: zoomed-in boxed
area.

Figure S3. Pairwise genome comparisons to *Thiomicrorhabdus* sp. KP2 (> 99% pairwise identity to the 16S rRNA gene sequence of the type species *Tmr. frisia* JB-A2[T], which does not have a sequenced genome) using the new re-classification scheme by Boden *et al.* (2017). Non-type species within the genus *Thiomicrorhabdus* are shown in triangles. Type species of genera within the family *Piscirickettsiaceae* are shown in circles. The bottom-left quadrant demarcates the boundary between type and non-type species.

Figure S4. Pairwise genome comparisons to *Thiomicrospira pelophila* DSM 1534[T] using the new re-classification scheme by Boden *et al.* (2017). Non-type species within the genus *Thiomicrospira* are shown in triangles. Type species of genera within the family

788 *Piscirickettsiaceae* are shown in circles, with the exception of *Thiomicrorhabdus frisia* JB-
789 A2^T which does not have a sequenced genome; *Thiomicrorhabdus* sp. KP2 (> 99% pairwise
790 identity to the 16S rRNA gene sequence) was used instead. The bottom-left quadrant
791 demarcates the boundaries between type and non-type species.
792
793

794
795 Figure S5. Boxplot diagrams of AF (left panel) and ANI (right panel) values as related to non-
796 type species (NT; n=120) of the genus *Bacillus* and type species (T, n=26) of genera within
797 the family *Bacillaceae*. Means are shown in green. Standard deviations are shown in blue.
798
799

800
Figure S6. AF:ANI pairwise genome comparisons in the order *Lactobacillales* (Figure 1-
right). Type species (n=37) of genera within the order *Lactobacillales* are shown in circles.
Non-type species (n=6) of the genus *Lactococcus* are shown in triangles. The primary
reference is the type species *Lactococcus lactis* subsp. *lactis* ATCC 19435^T. The bottom-left
quadrant demarcates the boundary between type and non-type species. Error bars indicate one
standard deviation from the mean.
807
808
809
810

811
812

813
814
815 Figure S7. Pairwise genome comparisons to the type species of the genus *Shewanella*: *S.*
816 *putrefaciens* JCM 20190[T] (Figure 2A). In circles: type species (n=19) of genera within the
817 order *Alteromonadales*. In triangles: non-type species (n=23) of the genus *Shewanella*. The
818 bottom-left quadrant demarcates the boundary between type and non-type species. Error bars
819 indicate one standard deviation from the mean.
820
821

822
823 Figure S8. Pairwise genome comparisons to the type species of the genus *Flavobacterium*: *F.*
824 *aquatile* LMG 4008[T] (Figure 2B). In circles: type species (n=69) of genera within the family
825 *Flavobacteriaceae*. In triangles: non-type species (n=83) of the genus *Flavobacterium*. The
826 bottom-left quadrant demarcates the boundary between type and non-type species. Error bars
827 indicate one standard deviation from the mean.

828
829

830
831 Figure S9. Pairwise genome comparisons to the type species of the genus *Vibrio*: *V. cholerae*
832 ATCC 14035^T (Figure 2C). Circles: type species (n=5) of genera within the family
833 *Vibrionaceae*. Triangles: non-type *Vibrio* spp. (n=65). The bottom-left quadrant demarcates
834 the boundary between type and non-type species. Error bars indicate one standard deviation
835 from the mean.
836
837

838
839 Figure S10. Pairwise genome comparisons to the type species of the genus *Methanosarcina*:
840 *M. barkeri* JCM 10043^T (Figure 2D). Circles: type species (n=6) of genera within the family
841 *Methanosarcinaceae*. Triangles: non-type *Methanosarcina* spp. (n=9). Error bars indicate one
842 standard deviation from the mean.
843
844

848 Figure S11. Pairwise genome comparisons to *Hydrogenovibrio marinus* DSM 11271[T] using

849 the new re-classification scheme by Boden *et al.* (2017)(Figure 3A). Non-type species within

850 the genus *Hydrogenovibrio* are shown in triangles. Type species of genera within the family

851 *Piscirickettsiaceae* are shown in circles. The bottom-left quadrant demarcates the boundary

852 between type and non-type species. Error bars indicate one standard deviation from the mean.

857 Figure S12. Pairwise genome comparisons to the type species of the genus *Bacillus*: *B.*

858 *subtilis* ATCC 6051[T]. Circles: type species (n=26) of genera within the family *Bacillaceae*.

859 Triangles: non-type *Bacillus* spp. (n=110). The bottom-left quadrant demarcates the boundary

860 between type and non-type species. Error bars indicate one standard deviation from the mean.

866 Figure S13. Genome clustering based on either pangenomic or phylogenomic analysis. All
867 138 single copy genes were included in the phylogenomic analysis.

889
890 Figure S14. Genome clustering based on either pangenomic or phylogenomic analysis. Only
891 the ribosomal genes were included in the phylogenomic analysis.