Demography and behavior of polygyne nests of the supercolonial
***Cataglyphis niger*:**

Does kinship matter?

Tali Reiner Brodetzki[1], Ofer Feinerman[2], Guy Brodetzki and Abraham Hefetz[1]

[1] School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University, Tel
Aviv, 6997801, Israel.

[2] Department of Physics of Complex Systems, Weizmann Institute of Science,
Rehovot, 7610001, Israel.

Corresponding author: Tali Reiner Brodetzki

School of Zoology, George S. Wise Faculty of Life Sciences, Tel Aviv University

Tel Aviv, 69978, Israel

Phone: + 972 3 6408766

e-mail: talireiner@gmail.com

Running headline: *C. niger* polygyne nest demography and behavior

Abstract

The basic ant colony structure is presumed to have evolved trough kin selection.

However, ants show a remarkable diversity in social organizations from a

monogynous and monoandrous queen to the more derived states of polygyny with

polyandrous queens. It is not clear whether kin selection can still explain these

derived societies. The existence of polygyny is still an evolutionary enigma since kin

selection theory predicts that queens should thrive for reproductive monopoly on

the one hand, and workers are predicted to favor their own matriline in rearing

gynes, on the other hand. Using a Quick Response (QR) - barcoding system that

allows tracking individual queens and workers interactions, along with polymorphic

DNA microsatellite markers indicating the matriline and patriline of each worker in

the nest, we demonstrate the complex social interactions in polygyne nests of

Cataglyphis niger. *C. niger* is not only polygyne but constitute a supercolony in the

study site. Our findings that both the residing queens and the workers in the nest are

not necessarily related to each other, are first for this species, support the

supercolony structure of the population. Also in line with such a social structure, we

demonstrate that they contribute equally to the nest production and rearing of the

queens. Unlike invasive supercolonial species, *C. niger* is native in Israel, raising

questions about the driving forces that stabilize this society, apart from kin selection.

Keywords: nepotism, social structure, supercolony, QR tracking, kin selection,
polygyne

Introduction

In the middle ages, the imposed sterility of catholic popes has driven an inclusive fitness behavior of appointing their nephews as cardinals who will later on succeed them as popes, this is known as nepotism. Similarly, monogymous ant colonies, the postulated basic colony structure, presumably evolved via kin selection (Hamilton, 1964; Hölldobler and Wilson, 1990; Hughes *et al.*, 2008; Boomsma, 2009). Notwithstanding, more derived states where the colony is composed of multiple queens (polygyny) that may also be multiply inseminated, are very common and pose a conflict for inclusive fitness theory. Polygyny poses two problems: from the queens' perspective, queens are predicted to compete for monopoly in reproduction; from the workers' perspective, workers are predicted to favor their own matriline in rearing gynes (young unmated queens; (Keller, 1993). The social consequences of polygyny in ant colonies are variable, ranging from equal reproductive share among queens to functional monogyny (i.e., only one queen reproduces while all other queens behave as workers), and from complete openness to incoming queens to limited acceptance of only related queens (Hölldobler and Wilson, 1977). Social polymorphism is mostly believed to be the result of multiple selective pressures such as ecological constraints on nesting success, predation, increased genetic diversity, and is usually associated with profound changes in life history strategies and dispersal behavior (Bourke and Franks, 1995; Keller, 1995; Pamilo and Crozier, 1996; Ross, 2001; Chapuisat, Bocherens and Rosset, 2004). Inclusive fitness theory also assumes favorable discrimination of closely related individuals (e.g., nepotism), unless between-matriline competition hampers total colony reproductive output. This discrimination is likely mediated through kin recognition cues, similar to nestmate recognition cues. Although, recognition cues exist in ants (Soroker *et al.*, 1994; Lahav *et al.*, 1998) they seem to play a role in within-colony kin discrimination only in rare cases (El-Showk *et al.*, 2010). Additionally, in most cases relatedness in these polygyne nests is low thus questioning the benefits of inclusive fitness theory at all in these cases. In fact, so far nepotism in eusocial insects, specifically in ants has been mostly rejected ((DeHeer and Ross, 1997; Zinck, Châline and Jaisson, 2009; Friend and Bourke, 2012), except

for *Formica fusca* (Hannonen and Sundström, 2003) which was criticized for not

having an actual behavioral assay and rather looked into the outcomes of behaviors

(Holzer *et al.*, 2006; e.g. kin composition of brood), and *Leptothorax acervorum* ants

(Gill and Hammond, 2011). Therefore, the extent of nepotism in eusocial insects is

still unclear, and the need for actual behavioral assays exists. Boomsma and

d'Ettorre (2013) suggested that within-colony kin discrimination might be

maintained in a secondary polygyny where there is a high queen turnover and

fluctuating relatedness in the colony.

The ant genus of *Cataglyphis* comprises over one hundred described species that are

distributed over the palaerctic realm in arid environments (Agosti, 1990).

Cataglyphis exhibits high social polymorphism from strictl monogyny (*C. sabulosa*,

C.bicolor , C. emmae and *C. hispanica*) through facultative polygyny (*C. velox* and *C.*

livida) to polygyny (*C. niger* and *C. mauritanica*; (Lenoir *et al.*, 2009; Boulay *et al.*,

2017). The genus also shows a diversity of reproduction modes (sexual and asexual),

which allows us to investigate the evolution of sociality. Thus, the importance of

understanding the mode of social organization in each species is high. Specifically, it

is important to understand interactions within a polygyny nest where there are

multiple possibilities of nest foundation and relatedness asymmetries (Keller and

Reeve, 1995).

Here, we focus on a population of *C. niger* that has polygyne nests that are organized

as a supercolony (Leniaud *et al.*, 2011). In this population structure, brood and

workers are often translocated to other nests in the population according to the

changing needs of the widespread colony. This complex social system encompasses

multiple thoughts about the mechanisms leading to this evolutionary stable status.

Nepotism has been investigated in the imported fire ant which has a similar

population and social structure in its invasive range, refuting the existence of

nepotism (DeHeer and Ross, 1997). However, *C. niger* is a native species, and

therefore could behave differently and possibly show within-colony kin

discrimination more than an invasive species.

Using a barcode system for tracking individual movement, we monitored worker-

worker and workers-queen interactions in polygyne nests. We also genotyped the

workers and queens in order to assess the within-nest relatedness. For the first time,

to the best of our knowledge, we obtained extensive data of both individual

interactions and relatedness between these individuals, showing the complex

relationships present in polygyne nests, and dismissing the possibility of nepotism.

Methods

<u>Ant collection and maintenance</u>

Three complete polygyne nests (nests 677, 868 and 869 having 3, 3 and 4 queens,

respectively) of *C. niger* (niger haplotype, as described in Eyer 2017) were excavated

from the Tel-Baruch population in the semi-stabilized sand dunes of Tel Aviv, Israel.

Prior to any manipulation, nests were kept in the lab for about a month under

constant temperature and humidity and were given water and a diet according to

(Bhatkar and Whitcomb, 1970), supplemented with tuna fish. For observations, a

subset of randomly selected workers (n=200 for each of the 3 nests) and the queens

were housed in artificial nests (20cm x 30cm) closed with an IR filtered roof and

given water and food *ad libitum*. Workers and queens were each marked with

individual 2mm QR barcodes using skin adhesive (SAUER-HAUTKLEBER, Manfred

sauer GMBH). Using a Canon T3i we took snap shots of the nest every 2 seconds, for

10,000 continuous frames (approximately 5.5 hours). The following observation

protocol was done for each of the 3 nests. Three days after tagging the first session

began, constituting free roaming queens and workers, and lasted for 3 days. In the

following session the queens were tethered at the petiole with fine nylon string (2-3

cm long), restricting each to a limited space in different corners of the nest, while

allowing free movement of the workers throughout the nest. At the end of this

session the queens were freed for three days, after which a second session of

tethered queens began, in which the queens were placed at corners different from

previous.

 Statistical analysis

 Behavioral analyses were performed using BugTag, a barcode analyzing system
 (developed by *robiotec*). The tags allow for constant monitoring of the location of
 each individual ant. The direction of the tags was corrected to match the position of
 the head of the ants, indicative of the direction at which the ant faced. Workers with
 low detection rates (their identity was certain in less than 30% of the 10000 frames)
 were excluded from the analyses. In cases where missed detections occurred in
 segments that were either fewer than five frames (less than 10 second apart), or if
 the change in the individual's position before and after the detection gap was lower
 than 2cm in distance, the position was interpolated using linear regression. Thus, the
 numbers of workers in each analysis were 137 and 122 for nest 677, 149 and 161 for
 nest 868, and 161 and 159 for nest 869.

 Using MATLAB, we defined a circular virtual arena (2cm radius) around each queen,
 hence termed "retinue space". Workers were considered to engaged in retinue
 around the queen if they were positioned in the "retinue space" while facing the
 queen for over 10% of the observation time. Ants that only occasionally interacted
 with a queen or sporadically shared her roaming space were not considered as
 involved in retinue behavior. The constancy of these groups over time, in terms of
 ant identity, and the fidelity of each worker to a specific queen or task were also
 measured. We compared our results with random permutations of worker-queen
 sets, allowing random positioning of workers next to the queens.

 Genetic analysis

 At the end of the experiment, all the queens and workers were sacrificed for genetic
 analyses. For nest 868 we analyzed the 3 queens and all the 165 workers in the
 experimental nest. For the remaining 2 nests (677and 869) we analyzed the queens
 and a sub set of workers (18 and 28 respectively), which included the top three
 "retinue" workers for each queen (the minimal number of retinue workers for all of
 the queens in all of the sessions) along with randomly selected nestmates. Queens'
 ovaries were dissected for assessing their degree of activation. Under binocular we
 compared the number of reproductively active ovarioles as well as the number and

size of oocytes in them, creating an activation index as follows: **0** - non-activated

ovarioles; **1** - 2 to 4 activated ovarioles with developed oocytes; **2** - 4 activated

ovarioles with small oocytes; **3** - all ovarioles with large, ready to lay eggs). We also

kept the spermathecal content of each queen for genetic analyses in ringer's buffer

at -80°c until DNA was extracted. DNA was extracted with 5% CHELEX (*BIO-RAD*) and

amplified with 7 microsatellite markers that had been designed previously for *C.*

hispanica (Ch23 ;Darras, Kuhn and Aron, 2014), *C. cursor* (Cc51,Cc89, and

Cc99;Pearcy *et al.*, 2004), and *C. niger* (Cn02, Cn04, and Cn08 ;Saar *et al.*, 2014)

using Type-it PCR mix (QIAGEN). PCR products were sequenced by ABI3500 genetic

analyzer and genotype analyses were performed with GeneMarker. Relatedness was

evaluated with Coancectry (Wang, 2011) and ML Relate (Kalinowski, Wagner and

Taper, 2006). We checked for possible correlation of relatedness to the mean

distances between all the individuals in the nest (workers and queens) using

SOCPROG 2.8. Network analysis was done using UCI6 and NetDraw.

Results

<u>Nests compositions and within nest relatedness</u>

The experimental nests had 3-4 queens (667- 3; 868- 3; 869-4), all of which were

multiply inseminated by 3-7 males. All the queens in the experiment (n=10) had well

developed ovaries with ready to lay eggs (level 3), except for one queen in nest 868

(Q3- tag # 552) and one queen in nest 869 (Q4- tag # 1189) who had medium size

oocytes (level 2). Table 1 presents the level of relatedness among the queens in the

experimental nests. In nest 869 three of the four queens were highly related (x>0.5;

full sisters or daughters), whereas the 4[th] queen was completely unrelated (x<0; not

different from relatedness=0). In nest 677 two of the queens were slightly related

(0.2181; half sisters or cousins), while the 3[rd] was not related to any of the other two

(x<0.1). All three queens in nest 868 were completely unrelated (x<0.1).

To assess nest demography, we genotyped all the workers and queens from nest 868

that were engaged in the tagging experiment (three queens and 165 workers). Since

these workers were randomly selected out of the approximately 1,000 workers that

populated the nest, we assume that they reflect a good assessment of the genotype

distribution within the entire nest. The genotype analysis revealed that 9.6% of the

workers were highly related (x>=0.5), 14.8% were moderately related (0.25<=x<0.5),

19.3% had low relatedness (0.125<=x<0.25) and 56.3% were unrelated (x<0.125).

Figure 1 shows a network analysis of relatedness values with ties indicating

relatedness greater than 0.25 and nodes are sized according to centrality measures.

Relatedness distribution of the workers to the different queens was as follows:

Highly related workers (x >=0.5, possible daughters or full sisters) Q1=17, Q2=15 (of

which 1 worker was similarly related to both Q1 and Q2), and Q3= 10. Moderately

related workers (0.25<=x<0.5, possible sisters) to the different queens was Q1=24,

Q2=26 (of which 4 workers were similarly related to both Q1 and Q2), and Q3= 4.

The remaining 63 workers were not related to any of the three queens, (X< 0.25).

Overall, nest relatedness for nest 868 was 0.058, and average relatedness of the

workers to each of the queens was similar (0.089±0.004). These results are slightly

lower than the within nest relatedness of the whole population of 0.112±0.009

(based on 129 workers from 6 nests; data not presented).

For nests 677 and 869 we genotyped a subset of workers as described in table 2. In

nest 677, five of 18 workers were highly related to one of the queens (relatedness

>=0.5): Q1=1, Q2= 4, and Q3=0. Seven workers were moderately related

(0.25<=x<0.5, possible sisters): Q1=5, Q2=1 (that was equally related to Q1), and Q3=

2. The remaining seven workers were not related to any of the queens (x<0.25). In

nest 869, eight of the workers were highly related to one of the queens (relatedness

>=0.5): Q1=2, Q2=1 (similarly related to Q1 and Q4), Q3=2 (one worker was similarly

related to Q4), Q4=5. Moderaltly related workers to queens were as follows: Q1= 4

(one similar to Q4), Q2= 6 (one similar to Q3 and one to Q4), Q3= 2, Q4=5. The

remaining 12 workers were not related to any of the queens.

<u>Identity and genotype of "retinue workers"</u>

The first set of experiments was done in an "open-range" arena where the queens

and workers could roam freely. Figure 2A depicts network analyses of the mean

distances between the individuals in the experimental nest 868. The network

presents only the ties between individuals that were in high proximity to each other over time (3 days; x<mean+2SD). The three queens as well as a small number of workers were highly central in the network, that is, they had close interactions with many individuals. These workers can be considered as 'retinue workers'. Queens also tended to stay very close to each other. It is also evident that, over time, many of the workers did not retain high proximity to each other (had few or no ties). Figure 2B shows a network of relatedness of the 'retinue' workers only and the queens, depicting only ties higher than 0.25 (moderately to highly related). Queens were not very central with respect to relatedness values, having had only a few daughters among the retinue workers. Furthermore, the central retinue workers were not very centrally related to others in the nest. Aligning relatedness results and mean proximity locations of all the workers and queens showed no clear correlation (i.e., workers or queens do not interact more with highly related individuals; Mantel Z test= -0.005, p=0.75).

In the "open-range" arena queens tended to be near each other, which made it difficult to precisely determine which worker tended which queen. Therefore, to assess whether workers do prefer specific queens we spaced out the queens within the nest by confining each to a specific location in the nest while allowing workers free movement, and thus giving them the opportunity to selectively retinue specific queens. Figure 3 shows the number of workers (averaged over 500 frames) that occupied the retinue space around each queen. Queens' retinue consisted of 3.86±0.15 (±SE) workers (range of averages 2.8-4.9 of workers attending a queen through the whole session). In none of the nests there was apparent preference to specific queens. Workers varied in the time they spent interacting with the different queens (Fig. 4a+b), but most did not have prolonged interactions. Approximately a quarter of the workers spent more than 10% of their time in one of the queens' retinue space. The maximal proportion of time that workers spent next to a specific queen was 30%. The relatedness of the top three "retinue" workers (which spent more than 10% of their time in retinue) to the attended queens ranged between -0.29 to 0.54 (Fig. 5), demonstrating that the queens' attendants were not necessarily related to the attended queen.

Workers were rarely loyal to the queens they initially attended to, but directed their

retinue behavior also towards the other queens. Worker switches between queens

occurred both during the same session as well as between the two replicate sessions

(supplementary material 1). Finally, worker preference in retinue of specific queen

was not different than random permutations of the workers' preference for the

different queens.

Discussion

Interactions between nestmates in an ant colony are the main focus of many recent

studies, all attempting to underline the mechanisms of 'self- organization' a hallmark

in the evolution of social insects (Pinter-Wollman *et al.*, 2011; Mersch, Crespi and

Keller, 2013; Heyman *et al.*, 2017). Here, for the first time, we combined individual

tracking with genetic data, in order to elucidate the possible effects of relatedness

and kin selection to worker in-nest behavior.

Polygyny is still an evolutionary enigma with respect to its adaptive value. Several

explanations have been proposed for the mutual benefits of this social dynamic,

from high relatedness of the co-inhabiting queens (the inclusive fitness explanation;

Giraud *et al.*, 2001) to robustness of the colony (higher genetic diversity, and colony

size explanations; Keller, 1995; Hughes, Ratnieks and Oldroyd, 2008; Heinze and

Rueppell, 2014). In such complexed societies where relatedness varies highly

between individuals, often reaching that of non-eusocial non-viscous population, we

expect to see altruistic/nepotistic interactions, like those commonly found in many

vertebrate societies (Eberhard, 1975; Clutton-Brock, 2002; Hatchwell, 2010).

Theoretically nepotistic interactions should be favored unless they negatively affect

the society's total fitness.

Additionally, in the present study, we assessed whether relatedness influences the

dynamics of the nest (in terms of worker behavior) in this species case where

polygyne nests also comprise a supercolony. As expected from a unicolonial system,

we found that in *C. niger* inter- as well as intra-colony relatedness is very low; only

about a third of nestmate workers are related to the level of sisters or half sisters. It

was argued by Helanterä *et al.*(2009), that in order to maintain such a unicolonial

structures it is necessary to have nepotism. Otherwise there is a chance for the

degradation of the social structure over evolutionary time scale. We could not detect

in our system any nepotism.

Relatedness between the queens was also rather variable, ranging from sisters to un-

related co-nesting queens. This suggests that this species exercises dependent

colony foundation, conforming to the usual case in polygyne nests (Cronin *et al.*,

2013). Some of the queens return to their natal nests after mating or are transferred

by workers to a new nest (budding; in the case of queens who are highly related)

while other queens (with low relatedness) disperse and join existing nests (this could

also be explained by colony fission). From our observations at the collection site, we

observed only a few newly mated queens that attempted independent colony

foundation, and very few nests with a single queen (unpublished data), thus

dependent colony foundation seems to be a more preferred strategy in this

population.

Our results also verify for the first time that this is a functional polygynous nest.

Assessment of the ovarian status as well as the relatedness between the queens and

workers indicate that all the queens contribute similarly to the nest population, thus

is a true functional polygyny. Further contributing to the low relatedness is our

finding that the queens are multiply inseminated. Finally, we observed frequent

movement between nests of workers and brood. While we cannot exclude that this

is simply nest relocation, we interpret this as free movement of workers brood and

queens between the nests of the supercolony, according to changes in available

resources (Robinson, 2014).

Queens in our experimental nests, were attended by the same number of workers

on average, respectively they also showed similar oocyte development. These

retinue workers, comprised only a small proportion of all the workers in the nest

suggesting there is a retinue caste, as seen in honey bees (Seely, 1979), and in *C.*

fellah as nurses' caste (Mersch, Crespi and Keller, 2013). These retinue workers were

not closely related to the queens. In fact, it seems as the retinue workers attended

the different queens at random, rather than being loyal to a specific queen. Indeed,

worker locations were not significantly difference in comparisons to random

permutations. The same is true when we examine the relatedness between

attendants and attendees. Nestmates do not associate more with highly related

individuals, irrespective of whether they are workers or queens. All this emphasizes

a stochastic intra-colony interactions, regarding both worker-queen and between

workers interactions (i.e. the interactions are not different than expected at random)

and thus, refutes any possibility of nepotistic behavior.

But is nepotism such a parsimony theory in this case? Under nepotism, we presume

an unequal distribution of resources in favor of specific matrilines, which perturbs

normal colony development, rendering polygyny evolutionary unstable. On the other

hand, if nepotism is maladaptive, what is the driving force for workers to keep

tending non-kin, and what are the mechanisms (if any) that prevent the downgrade

of non-kin-biased worker behavior. This evolutionary conflict may explain why

polygynous supercolonies are so rare, the benefits are mostly short lived and die out

through the evolutionary timescale. For this kind of social structure to be stable,

there should be other 'hard wired' mechanisms involved. An example is the 'social

chromosome', a non combining chromosome that exists only in polygynous

populations (Wang *et al.*, 2013; Purcell *et al.*, 2014). Future work should attempt to

elucidate possible mechanisms that keep these populations stable, such as worker

fixed sterility.

Acknowledgements

This project was funded by the Israel Science foundation (ISF grant # 844/13 to AH).
We would like to thank Oded Schor and Yael Heyman for help with designing and
analyzing the tracking methodology. We would also like to thank P.A. Eyer, Margarita
Orlova, and Tovit Simon for help with sample collection and technical assistance.

Authors' contributions

TRB, OF and AH conceived the ideas and designed the methodology; TRB collected
the data; TRB analyzed the genetic data; TRB and GB analyzed the tracking data; TRB
and AH led the writing of the manuscript. All authors contributed critically to the
drafts and gave final approval for publication.

Bibliography

Agosti, D. (1990) 'Review and reclassification of Cataglyphis (Hymenoptera,
Formicidae)', *Journal of Natural history*. Taylor & Francis Group, 24(6), pp. 1457–
1505.

Bhatkar, A. and Whitcomb, W. H. (1970) 'Artificial diet for rearing various species of
ants', *Florida Entomologist*. JSTOR, pp. 229–232.

Boomsma, J. J. (2009) 'Lifetime monogamy and the evolution of eusociality',
Philosophical Transactions of the Royal Society B: Biological Sciences. The Royal
Society, 364(1533), pp. 3191–3207.

Boomsma, J. J. and d'Ettorre, P. (2013) 'Nice to kin and nasty to non-kin: revisiting
Hamilton's early insights on eusociality', *Biology letters*. The Royal Society, 9(6), p.
20130444.

Boulay, R. *et al.* (2017) 'Social Life in Arid Environments: The Case Study of
Cataglyphis Ants', *Annual Review of Entomology*, 62(1), pp. 305–321. doi:
10.1146/annurev-ento-031616-034941.

Bourke, A. F. G. and Franks, N. R. (1995) *Social evolution in ants*. Princeton University
Press.

Chapuisat, M., Bocherens, S. and Rosset, H. (2004) 'Variable queen number in ant
colonies: no impact on queen turnover, inbreeding, and population genetic
differentiation in the ant Formica selysi', *Evolution*, 58(5), pp. 1064–1072.

Clutton-Brock, T. (2002) 'Breeding together: kin selection and mutualism in
cooperative vertebrates', *Science*. American Association for the Advancement of
Science, 296(5565), pp. 69–72.

Cronin, A. L. *et al.* (2013) 'Recurrent evolution of dependent colony foundation
across eusocial insects', *Annual Review of Entomology*, 58.

Darras, H., Kuhn, A. and Aron, S. (2014) 'Genetic determination of female castes in a
hybridogenetic desert ant', *Journal of Evolutionary Biology*. Wiley Online Library,
27(10), pp. 2265–2271. doi: 10.1111/jeb.12470.

DeHeer, C. J. and Ross, K. G. (1997) 'Lack of detectable nepotism in multiple-queen
colonies of the fire ant Solenopsis invicta (Hymenoptera: Formicidae)', *Behavioral
Ecology and Sociobiology*. Springer-Verlag, 40(1), pp. 27–33.

Eberhard, M. J. W. (1975) 'The evolution of social behavior by kin selection', *The
Quarterly Review of Biology*. Stony Brook Foundation, Inc., 50(1), pp. 1–33.

El-Showk, S. *et al.* (2010) 'Are you my mother? Kin recognition in the ant Formica
fusca', *Journal of evolutionary biology*. Wiley Online Library, 23(2), pp. 397–406.

Friend, L. A. and Bourke, A. F. G. (2012) 'Absence of Within-Colony Kin Discrimination
in a Multiple-Queen Ant, Leptothorax acervorum', *Ethology*. Wiley Online Library,

118(12), pp. 1182–1190.

Gill, R. J. and Hammond, R. L. (2011) 'Workers influence royal reproduction', *Proceedings of the Royal Society B: Biological Sciences*, 278(1711), pp. 1524–1531. doi: 10.1098/rspb.2010.1774.

Giraud, T. *et al.* (2001) 'High genetic relatedness among nestmate queens in the polygynous ponerine ant Gnamptogenys striatula in Brazil', *Behavioral Ecology and Sociobiology*, 49(2–3), pp. 128–134. doi: 10.1007/s002650000284.

Hamilton, W. D. (1964) 'The genetical evolution of social behaviour. II', *Journal of theoretical biology*. Elsevier, 7(1), pp. 17–52.

Hannonen, M. and Sundström, L. (2003) 'Sociobiology: Worker nepotism among polygynous ants', *Nature*. Nature Publishing Group, 421(6926), p. 910. doi: 10.1038/421910a.

Hatchwell, B. J. (2010) 'Cryptic kin selection: kin structure in vertebrate populations and opportunities for kin-directed cooperation', *Ethology*. Wiley Online Library, 116(3), pp. 203–216.

Heinze, J. and Rueppell, O. (2014) 'The frequency of multi-queen colonies increases with altitude in a Nearctic ant', *Ecological Entomology*, 39(4), pp. 527–529. doi: 10.1111/een.12119.

Helanterä, H. *et al.* (2009) 'Unicolonial ants: where do they come from, what are they and where are they going?', *Trends in Ecology and Evolution*, 24(6), pp. 341–349. doi: 10.1016/j.tree.2009.01.013.

Heyman, Y. *et al.* (2017) 'Ants regulate colony spatial organization using multiple chemical road-signs', *Nature communications*. Nature Publishing Group, 8, p. 15414.

Hölldobler, B. and Wilson, E. O. (1977) 'The number of queens: An important trait in ant evolution', *Naturwissenschaften*, 64(1), pp. 8–15. doi: 10.1007/BF00439886.

Hölldobler, B. and Wilson, E. O. (1990) *The ants*. Harvard University Press.

Holzer, B. *et al.* (2006) 'Sham nepotism as a result of intrinsic differences in brood viability in ants', *Proceedings of the Royal Society B: Biological Sciences*, 273(1597), p. 2049 LP-2052. doi: 10.1098/rspb.2006.3553.

Hughes, A. R. *et al.* (2008) 'Ecological consequences of genetic diversity', *Ecology Letters*, 11(6), pp. 609–623. doi: 10.1111/j.1461-0248.2008.01179.x.

Hughes, W. O. H. H., Ratnieks, F. L. W. W. and Oldroyd, B. P. (2008) 'Multiple paternity or multiple queens: two routes to greater intracolonial genetic diversity in the eusocial Hymenoptera', *Journal of evolutionary biology*. Wiley Online Library, 21(4), pp. 1090–1095. doi: 10.1111/j.1420-9101.2008.01532.x.

Kalinowski, S. T., Wagner, A. P. and Taper, M. L. (2006) 'ML-Relate: a computer program for maximum likelihood estimation of relatedness and relationship', *Molecular Ecology Resources*. Wiley Online Library, 6(2), pp. 576–579.

Keller, L. (1993) 'Queen number and sociality in insects'. Oxford University Press
Oxford.

Keller, L. (1995) 'Social life: the paradox of multiple-queen colonies', *Trends in*
Ecology & Evolution. Elsevier Current Trends, 10(9), pp. 355–360. doi:
10.1016/S0169-5347(00)89133-8.

Keller, L. and Reeve, H. K. (1995) 'Why do females mate with multiple males? The
sexually selected sperm hypothesis', *Advances in the Study of Behaviour*. New York:
Academic Press, 1965-, 24, pp. 291–316.

Lahav, S. *et al.* (1998) 'Nestmate recognition in the ant Cataglyphis niger: Do queens
matter?', *Behavioral Ecology and Sociobiology*, 43(3), pp. 203–212. doi:
10.1007/s002650050482.

Leniaud, L. *et al.* (2011) 'Multiple mating and supercoloniality in *Cataglyphis* desert
ants', *Biological Journal of the Linnean Society*, 104(4), pp. 866–876. doi:
10.1111/j.1095-8312.2011.01772.x.

Lenoir, A. *et al.* (2009) 'Cataglyphis desert ants: a good model for evolutionary
biology in Darwin's anniversary year-A review', *Israel Journal of Entomology*, 39, pp.
1–32. Available at: http://digital.csic.es/handle/10261/65135.

Mersch, D. P., Crespi, a. and Keller, L. (2013) 'Tracking Individuals Shows Spatial
Fidelity Is a Key Regulator of Ant Social Organization', *Science*, 340(6136), pp. 1090–
1093. doi: 10.1126/science.1234316.

Pamilo, P. and Crozier, R. H. (1996) 'Reproductive skew simplified', *Oikos*. JSTOR, pp.
533–535.

Pearcy, M. *et al.* (2004) 'Characterization of nuclear DNA microsatellite markers in
the ant Cataglyphis cursor', *Molecular Ecology Resources*. Wiley Online Library, 4(4),
pp. 642–644.

Pinter-Wollman, N. *et al.* (2011) 'The effect of individual variation on the structure
and function of interaction networks in harvester ants', *Journal of The Royal Society*
Interface, 8(64), pp. 1562–1573. doi: 10.1098/rsif.2011.0059.

Purcell, J. *et al.* (2014) 'Convergent Genetic Architecture Underlies Social
Organization in Ants', *Current Biology*, 24(22), pp. 2728–2732. doi:
https://doi.org/10.1016/j.cub.2014.09.071.

Robinson, E. J. H. (2014) 'Polydomy: the organisation and adaptive function of
complex nest systems in ants', *Current Opinion in Insect Science*. Elsevier, 5, pp. 37–
43.

Ross, K. G. (2001) 'Molecular ecology of social behaviour: analyses of breeding
systems and genetic structure', *Molecular Ecology*. Blackwell Science Ltd, 10(2), pp.
265–284.

Saar, M. *et al.* (2014) 'At the brink of supercoloniality: genetic, behavioral, and

chemical assessments of population structure of the desert ant Cataglyphis niger',
Frontiers in Ecology and Evolution, 2(May), pp. 1–10. doi: 10.3389/fevo.2014.00013.

Seely, T. D. (1979) 'Queen substance dispersal by messenger workers in honeybee
colonies', *Behavioral Ecology and Sociobiology*. Springer, 5(4), pp. 391–415.

Soroker, V. *et al.* (1994) 'The postpharyngeal gland as a "Gestalt" organ for nestmate
recognition in the antCataglyphis niger', *Naturwissenschaften*, 81(11), pp. 510–513.
doi: 10.1007/BF01132686.

Wang, J. (2011) 'COANCESTRY: a program for simulating, estimating and analysing
relatedness and inbreeding coefficients', *Molecular ecology resources*. Blackwell
Publishing Ltd, 11(1), pp. 141–145.

Wang, J. *et al.* (2013) 'A Y-like social chromosome causes alternative colony
organization in fire ants', *Nature*. Nature Publishing Group, 493(7434), pp. 664–668.
doi: 10.1038/nature11832.

Zinck, L., Châline, N. and Jaisson, P. (2009) 'Absence of nepotism in worker–queen
care in polygynous colonies of the ant Ectatomma tuberculatum', *Journal of insect
behavior*. Springer US, 22(3), pp. 196–204.

Table 1: Relatedness values between queens

Nest - Queen	**869 Q2**	**869 Q3**	**869 Q4**
869 Q1	0.5304	-0.1901	0.5362
869 Q2	--	0.0626	0.6122
869 Q3	--	--	-0.0165
	667 Q2	**667 Q3**	--
667 Q1	0.2181	0.047	--
667 Q2	--	0.0794	--
	868 Q2	**868 Q3**	--
868 Q1	-0.1573	0.0873	--
868 Q2	--	-0.3619	--

NEST_QUEEN	RELATEDNESS LEVEL	
	0.25	0.5
677_Q1	6^{12}	1
677_Q2	5^{2}	4
677_Q3	2^{1}	0
869_Q1	$6^{12*\circ}$	2^{*}
869_Q2	$7^{1*\circ^{wy}}$	1^{*}
869_Q3	$4^{1^{w}}$	2^{1}
869_Q4	10^{12wy}	5^{1}

Table 2: Number of workers who have relatedness values of 0.25/0.5 to any of the

queens. N= 18 for nest 677 and 28 for nest 869. $^{12*\circ^{wy}}$ indicate individuals that are

related to both queens.

Figure legends:

Figure 1: Network analysis of relatedness values of the ants in nest 868 (with a

threshold of 0.25). Queens colored in red. The size of the circle indicates its

centrality (how many relationships he holds). The thickness of the line indicates the

strength of the tie = relatedness value between individuals.

Figure 2: A) Network analysis of the averaged minimal distance (proximity) between

the ants in nest 868 (threshold of x< 2*SD). Queens colored in red. The size of the

square indicates its centrality (how many relationships he holds). The thickness of

the lines indicate the strength of the tie= average proximity between individuals. B)

Network analysis of relatedness values of the retinue ants in nest 868 showing ties

larger than 0.25). Queens colored in red. The size of the circle indicates its centrality

(how many relationships he holds). The thickness of the line indicates the strength of

the tie = relatedness value between individuals.

Figure 3: Assessment of queens' retinue, number of workers around each queen

averaged for 500 frames. Average retinue was 3.86 (SE=0.15; range:2.8-4.9) workers

for each queen. (a)nest 868; queen 552 in blue had smaller oocytes (b) nest 677 (c)

nest 869; queen 1189 in purple had smaller oocytes.

505   Figure 4: The proportion of time a worker interacted with a specific queen,

506   normalized to each of the workers detection success in nest 868. a) First session

507   where queens were tied. b) The second session of tied queens, after an intermediate

508   time where queens were free.

Figure 5: The level of Relatedness between individual ants, line thickness and length

represents the level of relatedness, color and shape represent the different retinue

groups of the queens (W=worker; Q= queen). Queens and their workers attendants

(for each queen three of the workers who spent the most time next to the queen) in

nest 677,868 (relatedness under zero was emitted from the network). A) nest 677 b)

nest 868 c) nest 869

 <u>Figure Supplementary 1</u>: a) Individual workers who spent more than X% out of 20

 minute segments in a specific queen's retinue space. Each queen's retinue space is

 defined by a different color, other locations are black. b) Workers who attended the

 queens for more than 10% of the time. In dark blue the workers did not attend the

 specific queen, in light blue they attended the queen for only one repeat of the

 experiment, in yellow they attended to the queen in both repeats.

23