Plasticity in striatal dopamine release is governed by release-independent

depression and the dopamine transporter

Mark D. Condon[1], Nicola J. Platt[1], Bradley M. Roberts[1], Min-Yee Tseu[1], Stefania Vietti-Michelina[1],

Sarah Threlfell[1,2] and Stephanie J. Cragg[1,2]*

[1]Department of Physiology, Anatomy and Genetics, Parks Road, University of Oxford, OX1 3PT, UK

[2]Oxford Parkinson's Disease Centre, Oxford, OX1 3PT

*Author for Correspondence: Stephanie.cragg@dpag.ox.ac.uk, +44 1865 282513

Short Title: Short-term plasticity in dopamine release

Keywords: Dopamine, striatum, release probability, dopamine release, short-term plasticity, short-term

depression, short-term facilitation, dopamine transporter, release-independent depression, axonal

excitability, voltammetry

Key Findings:

• Short-term plasticity in dopamine release is only weakly governed by Ca^{2+} and initial release

• Short-term depression is strongly dependent on regulation of axonal excitability

• The DAT determines short-term plasticity and drives short-term depression

• DAT governs the balance between release-dependent and release-independent mechanisms

Abstract

Action potentials in mesostriatal dopamine (DA) neurons encode the motivational value of salient stimuli. DA neurons possess extensively branched axonal arbours, and how action potentials are converted to DA output will be influenced dynamically and critically by axonal mechanisms that are poorly understood. We explored the role for mechanisms governing release probability and axonal excitability in determining short-term plasticity (STP) of DA release, using fast-scan cyclic voltammetry in *ex vivo* mouse striatum. The brief short-term facilitation (STF) and longer short-term depression (STD) seen were only weakly dependent on initial release. Rather, STP was strongly governed by release-independent mechanisms that involved K^+-dependent axonal excitability and surprisingly, the dopamine transporter (DAT), particularly in dorsal striatum. The DAT acts as a master regulator of DA STP, governing the balance between release-dependent and release-independent mechanisms that also show region-specific gating.

Introduction

Dopamine (DA) release in the striatum plays key roles in action selection and behavioural reinforcement, and is dysregulated in diverse disorders including Parkinson's disease and addictions. DA neurons generate action potentials (APs) at a range of frequencies, firing tonically below 10 Hz, and intermittently in short bursts with instantaneous frequencies of up to approximately 40 Hz (1-6). Bursts occur in response to salient environmental stimuli predicting reward value or action signals (6-8). However, the extent to which AP activity is conveyed into striatal DA release remains unclear: DA axons are important sites for local regulation of DA output (9-12) which shows strong short-term plasticity (STP) and distorts the relationship between activity and DA output (11, 13-16).

Despite classical accounts of axons as simple high-fidelity cables, neurotransmitter release from CNS axons is shaped by factors that regulate AP propagation and axonal excitability such as axonal morphology, and the expression of ion channels, presynaptic receptors and other diverse mechanisms that regulate neurotransmitter release probability and STP (17). Axons of DA neurons are remarkable; they comprise vast,

extensively branched arbours (18, 19), and therefore axonal properties are likely to be particularly important

in governing striatal DA output. One major axonal influence is input from striatal cholinergic interneurons

onto nicotinic receptors (nAChRs) which influence the pattern of DA release and promote short-term

depression (STD) (11, 14, 20-22). In the absence of nAChR activation, DA release shows an underlying

spectrum of STP ranging from short-term facilitation (STF) to STD (11, 14, 21, 23, 24) but the underlying

mechanisms are poorly understood. There is evidence that STP might be only weakly related to initial levels

of release (16, 23, 24).

Here, we delineate the roles of three types of intrinsic mechanisms on DA axons that could underlie

STP of DA release. We examine, firstly, the role for initial release, a proxy for probability of release (P_r);

secondly, the role for K^+-dependent processes such as axonal membrane excitability and repolarization; and

thirdly, whether STP can be influenced by the dopamine transporter (DAT). Besides mediating DA uptake,

the DAT generates a depolarising transport-coupled conductance in midbrain DA neurons (25, 26), and limits

vesicular release through suggested interactions with synaptic vesicle proteins (27, 28). Thus, the DAT has

the potential to govern STP of DA release via regulation of axonal excitability and/or P_r.

We reveal that initial P_r plays a limited role in STP which is largely release-independent, except on

the shortest of timescales when STF operates. We identify that STD is strongly determined by membrane

excitability, particularly in dorsal striatum, and that the DAT drives STF and STD, effectively clamping release

at physiological inter-pulse intervals. We propose a region-specific hierarchy of interacting drivers of STP,

overseen by the DAT, with the DAT promoting release-insensitive over release-dependent mechanisms.

Results

Patterns of short-term plasticity at DA release sites are region-specific and weakly release-dependent

We assessed STP of DA release in acute coronal striatal slices evoked using pairs of local electrical pulses. All experiments were conducted without the confounding effects of evoked striatal ACh by inclusion of nAChR antagonist DHβE (1 μM) throughout. Inclusion of nAChR antagonists during local stimulation also makes evoked DA release in *ex vivo* slice resemble DA release evoked *in vivo* by midbrain or medial forebrain bundle stimulation: sensitivity to frequency as seen *in vivo* (29, 30) is unmasked in slices by nAChR inhibition (14, 31). The ratio of DA release evoked by the second pulse in a pair of pulses compared to a single pulse (paired-pulse ratio, PPR) varied with inter-pulse intervals (IPI) of 10-200 ms in dorsolateral striatum (CPu) (**Fig. 1A,C**) and in nucleus accumbens core (NAc) (**Fig. 1B,C**). In both regions, PPR decreased with increasing IPI. Short-term facilitation (STF, PPR>1) was observed at the shortest IPIs (10 ms in CPu, 10-25 ms in NAc), whereas short-term depression (STD) (PPR<1) was observed at longer IPIs (≥ 25 ms in CPu, >40 ms in NAc), as shown previously (14, 32). In CPu, STD at IPIs ≥ 25 ms was significantly more pronounced than in NAc (**Fig. 1C,** Two-way ANOVA, region x IPI interaction, $F_{4,25}$ = 4.195, *P* = 0.010, *n* = 7). STP of DA release therefore shows strong STD in both CPu and NAc, and to a greater extent in CPu, which suggests region-specific gating of STP.

PPR is usually inversely related to initial release probability (P_r) at central synapses, resulting in STF when P_r is low, and STD when P_r is high (33-36). STF at these synapses consists of several temporally-overlapping processes with a range of time constants from 30 – 500 ms (37). An inverse relationship between PPR and single pulse release has been seen in CPu for very short IPI of 10 ms but not at 100 ms (23), suggesting that STP of DA release might reflect initial P_r at only very short intervals. We probed the relationship between PPR and the DA release evoked by a single pulse (as a proxy of initial P_r) in CPu and in NAc across a population of release sites that exhibited a 10-fold range in $[DA]_o$ evoked by a single pulse (1p

[DA]$_o$). We found an inverse relationship between 1p [DA]$_o$ and PPR for IPI of 10 ms in both CPu and NAc,

and for 40 ms in NAc only, but in neither region at intervals of 100 ms (**Fig. 1D,** linear regressions, 10 ms,

CPu: β = -0.79 ± 0.14 [95% CI: -1.08 to -0.50], $F_{1,33}$ = 30.9, *P* = 3.56 x 10^{-6}, *n* = 35; NAc: β = -0.52 ± 0.19 [95% CI:

-0.93 to -0.11], $F_{1,17}$ = 7.136, *P* = 0.016, *n* = 19; 40 ms, CPu, *P* = 0.152, *n* = 35; NAc 95% CI: -0.38 to -0.001], $F_{1,}$

$_{17}$ = 4.489, *P* = 0.049, *n* = 19; 100 ms, CPu: *P* = 0.122, n = 35; NAc: *P* = 0.917, n = 19). These data suggest that

DA PPR is a function of P$_r$ at only very short intervals and specifically, only at those intervals at which we

observe a STF. These data also confirm that STP of DA release is dominated by STD at all but the shortest

intervals (≤25 ms) and indicate that PPR at physiological IPIs (≥40 ms) is not strongly influenced by P$_r$. Taken

together these findings suggest that STP might be governed by release-dependent mechanisms at short

intervals, but by mechanisms unrelated to initial release at longer physiological intervals, across striatum.

We tested whether this expression of STD could be explained by activation of inhibitory D2-type

autoreceptors (D2Rs). We prevented activation of D2Rs with the D2/D3-receptor antagonist L741,626.

L741,626 (1 µM) did not appear to influence peak [DA]$_o$ (**Fig. 1E**) and did not significantly alter PPR at IPIs up

to 200 ms in either CPu or NAc (**Fig. 1F**, Two-way ANOVA, CPu: *P* = 0.420, n= 5, NAc: *P* = 0.2197, n = 6),

consistent with previous findings that D2Rs modify release at only longer IPI (>200 ms) (13, 38-40). These

findings show that STD at these intervals is not explained by D2R activation.

STP is shaped by release-dependent facilitation and release-independent depression

To test directly whether STP is governed by P$_r$, we varied P$_r$ at a given release site by varying the extracellular

concentration of Ca^{2+} ([Ca^{2+}]$_o$) between 1.2 and 3.6 mM. [DA]$_o$ evoked by 1p was strongly dependent on

[Ca^{2+}]$_o$, in both CPu and NAc (**Fig. 2A,B**) as expected, and varying [Ca^{2+}]$_o$ had a stronger outcome on 1p [DA]$_o$

in CPu than in NAc (**Fig. 2C**, Two-way ANOVA, [Ca^{2+}]$_o$ x region interaction, $F_{2,20}$ = 17.25, *P* = 4.43 x10^{-5}, *n* = 6),

consistent with previous findings (24). However, despite large changes in 1p-evoked DA release (2.5 to 4-

fold), [Ca^{2+}]$_o$ had only modest effects on PPR. In CPu, a trend towards an inverse relationship at 10 ms IPI did

not reach significance (**Fig. 2D,F**, Two-way ANOVA, *P* = 0.604, n = 4), while in NAc, PPR varied inversely with

[Ca^{2+}]$_o$ at short intervals, reaching significance at only 10 ms IPI (**Fig. 2E,F,** Two-way ANOVA, IPI x [Ca^{2+}]$_o$

interaction, $F_{8,30}$ = 4.497, *P* = 0.001, *n* = 3). Therefore, the effects of varying P_r on PPR are subtle and limited

to very short IPIs, revealing that STP of DA release is only weakly governed by classic presynaptic release-

dependent mechanisms, and suggesting that other mechanisms might be important.

We noted also that the greater range in PPR seen in NAc than in CPu across different $[Ca^{2+}]_o$ occurred

over a smaller range of 1p $[DA]_o$ than in CPu: there was an apparently stronger relationship between initial

release and STP in NAc than CPu (**Fig. 2F**). The difference in STP between CPu and NAc cannot therefore be

attributed to a difference in the sensitivity of initial P_r to $[Ca^{2+}]_o$ (see Fig. 2C). Thus, STP of DA release can be

governed only partly by $[Ca^{2+}]_o$-dependent processes, and must involve additional participating mechanisms

that are not modulated by P_r which, by definition, can be referred to as release-insensitive (or release-

independent).

To confirm that STD and insensitivity to $[Ca^{2+}]$ are intrinsic properties of DA axons, and are not a

result of co-activation of an undefined input to DA axons, we used optogenetic stimulation of DA axons to

selectively evoke DA release. In striatum from heterozygote DAT[IRES-Cre] mice expressing ChR2-eYFP, a brief

light flash (2 ms, 473 nm wavelength) evoked $[DA]_o$ transients that varied with varying $[Ca^{2+}]_o$ (**Fig. 2G**) and

with a steeper $[Ca^{2+}]_o$ concentration-response curve in CPu than in NAc (**Fig. 2H**, nonlinear regression, CPu:

R^2 = 0.91, n = 3, NAc: R^2 = 0.94, n = 3, comparison of fits: $F_{4,118}$ = 5.94, *P* = 2 x 10^{-4}), as seen for electrically

evoked DA release. Furthermore, at an interval of 40 ms, at which ChR2 can reliably drive repeat spiking in

DA neurons (1) and can be followed by the release of DA (11), PPR was greater in NAc than in CPu as

observed using electrical stimulation (see Fig. 1C) and moreover, there was no inverse relationship between

PPR and 1p $[DA]_o$ (**Fig. 2I,** linear regression, CPu: *P* = 0.874, n = 3, NAc: *P* = 0.286, n = 3). These data confirm

that STD is not attributable to concurrent activation of other inputs. For subsequent experiments, we chose

to electrical over optogenetic stimulation (with nAChR inhibitors included throughout) to avoid confounding

effects of ChR2 conductances on Ca^{2+} entry.

132 A release-insensitive mechanism of short-term depression is controlled by K$^+$-dependent gating

We next tested whether STD is driven by mechanisms related to membrane excitability by varying $[K^+]_o$.

Varying $[K^+]_o$ can alter membrane potential and/or rate of repolarisation through the Nernstian driving forces

underlying K$^+$-mediated currents (41) and/or through K$^+$-dependent inhibition of K$^+$-channel inactivation (42).

Reduced $[K^+]_o$ can favour membrane hyperpolarisation, and faster repolarisation between pulses promoting

de-inactivation of Na$^+$ channels, but can also promote inactivation of K$^+$-channels, leading to use-dependent

depolarisation and enhanced Na$^+$-channel recruitment. In either case, recruitment of voltage-gated Na$^+$

channels at a subsequent depolarising event will be altered.

We found that varying $[K^+]_o$ (1.25 - 7.5 mM) did not change 1p $[DA]_o$ in either CPu (**Fig. 3A**) or NAc

(**Fig. 3B**) but significantly modulated STP (**Fig. 3A-D**). In CPu, PPR varied inversely with $[K^+]_o$ at all IPIs, with

STD minimized at lowest $[K^+]_o$ (**Fig. 3C**, Two-way ANOVA, $[K^+]_o$ x IPI interaction, $F_{8,30}$ = 7.66, P = 1.53 x 10^{-5}, n =

3). In NAc, the effect of $[K^+]_o$ on PPR was less evident than in CPu, but showed an overall significance (**Fig. 3D**,

Two-way ANOVA, main effect of $[K^+]_o$, $F_{2,30}$ = 15.15, P = 2.83 x 10^{-5}, n = 3). Since we observed no effect of $[K^+]_o$

on 1p $[DA]_o$, these effects on PPR were unrelated to initial P_r at either short (10 ms) or longer (100 ms) IPI

(**Fig. 3E,** vertical slopes). These findings demonstrate directly that STP of DA release can be dissociated from

initial P_r, i.e. there is a release-independent STP. Furthermore, they suggest that STD is governed by K$^+$-

dependent membrane polarisation/excitability particularly in CPu.

149 K$^+$-dependent gating of short-term depression requires axonal K$_v$ channels

To identify whether the control of STP by $[K^+]_o$ is accounted for by voltage-gated K channels (K$_v$) channels we

tested whether the effects of $[K^+]_o$ on PPR could be substituted for, and prevented by, the broad-spectrum

blocker 4-aminopyridine (4-AP). We hypothesised that by reducing K$_v$ conductances, 4-AP should promote

STD and prevent effects of further changes in $[K^+]_o$. In CPu, 4-AP (100 μM), unlike changes in $[K^+]_o$ alone,

profoundly increased 1p $[DA]_o$ (**Fig. 4A, B**, Two-way ANOVA, main effect of 4-AP: $F_{1,3}$ = 33.63, P = 0.010, n =

4). This effect is consistent with the widening of the action potential by 4-AP seen in DA neurons (43) which

will increase Ca^{2+} entry and neurotransmitter release for a single stimulus. In addition, 4-AP, similar to the

effects of high $[K^+]_o$ (see Fig. 3), but unlike high $[Ca^{2+}]_o$ (see Fig. 2), reduced PPR at all IPIs, and also prevented

the effects of changing $[K^+]_o$ on PPR (**Fig. 4C, 4D,** Two-way ANOVA, 4-AP x $[K^+]_o$ interaction, $F_{1, 12} = 6.003$, $P =$

0.031, *n* = 4). These findings indicate that inhibition of K_v channels promotes STD, and suggest conversely

that enhanced K_v currents relieve STD, and suggest that hyperpolarizing/repolarizing conditions limit STD and

promote subsequent release.

In NAc, we observed no significant effect of 4-AP on 1p $[DA]_o$ (**Fig. 4E,F,** Two-way ANOVA, $P = 0.266$,

n = 3). 4-AP did decrease PPR at all IPIs, although there was no effect of varying $[K^+]_o$ in control conditions or

in the presence of 4-AP (**Fig. 4G, H**). These findings verify further that STD can be distinct from the effects on

initial P_r and that STD is underpinned by K_v-gated regulation of axonal excitability.

K^+-dependent gating does not alter release-dependence of STP

We investigated whether $[K^+]_o$-dependent STD could be a limiting factor that prevents STP of DA release

from responding to changes in P_r. We explored whether a relief of STD in low $[K^+]_o$ could unmask a greater

sensitivity of STP to changes in P_r driven by variation in $[Ca^{2+}]_o$. In CPu, 1p-evoked $[DA]_o$ varied with $[Ca^{2+}]_o$ in

a similar manner in 1 mM and 5 mM $[K^+]_o$ (**Fig. 5A-C**) and although PPRs were higher when $[K^+]_o$ was lower

(**Fig. 5D**) (as seen in Figs. 3C, 4C), there was no significant interaction between $[K^+]_o$ and $[Ca^{2+}]_o$ at IPIs of

either 10 ms or 100 ms (**Fig. 5D**, Two-way ANOVA, 10 ms IPI: $P = 0.086$, 100 ms IPI: $P = 0.654$, n = 10). By

limiting STD, we did not enhance Ca^{2+}-dependent modulation of STF or reveal a relationship between STD

and P_r in CPu.

In NAc, 1p $[DA]_o$ varied with $[Ca^{2+}]_o$ in a manner that did not depend on $[K^+]_o$ (**Fig. 5E-G**).

Furthermore, there was no significant interaction between the effects of $[K^+]_o$ and $[Ca^{2+}]_o$ on PPR at either 10

ms or 100 ms IPI (**Fig. 5H**, Two-way ANOVA, 10 ms IPI: $P = 0.963$, 100 ms IPI: $P = 0.883$, n = 7). The release-

independent mechanism through which $[K^+]_o$ modulates STP does therefore not apparently mask classical

roles for Ca^{2+} in STP, in either CPu or NAc. This suggests that an additional mechanism may operate to limit

the Ca^{2+}-dependence of STP besides K_v-regulation of axonal excitability and polarization.

The dopamine transporter regulates STP of dopamine release

The dopamine transporter (DAT) has previously been shown to limit DA release (27, 28) by a mechanism that

is independent of DA transport (44-46), suggesting that the DAT might play a key role in regulating P_r

alongside its role in the uptake of DA. The DAT is also known to mediate electrogenic currents during DA

binding and transport (25, 47) that have been shown to modulate the membrane potential of DA neurons *in*

vitro (26). We therefore investigated whether DAT in striatum regulates STP.

We explored PPR in the presence of a range of uptake inhibitors. We found that the monoamine

uptake inhibitors cocaine and methylphenidate (MPH) and the DAT-selective inhibitor nomifensine

significantly altered the pattern of STP in a common manner. Cocaine (5 µM), MPH (5 µM) or nomifensine

(10 µM) increased the peak amplitude and decreased the rate of decay of $[DA]_o$ transients in CPu and NAc in

keeping with DA re-uptake inhibition (**Fig. 6A,C,G,I**). Furthermore, they significantly altered STP. In CPu, DAT

inhibition prevented STF and significantly relieved STD (Cocaine: **Fig. 6A,B**, Two-way ANOVA, cocaine x IPI

interaction, $F_{4, 30} = 11.97$, *P* = 6.24 x 10^{-6}, *n* = 4, MPH: **Fig. 6C,D**, Two-way ANOVA, MPH x IPI interaction, $F_{4, 40}$

= 5.316, *P* = 0.002, *n* = 5; Nomifensine: **Fig. 6E,F**, Two-way ANOVA, nomifensine x IPI interaction, $F_{4, 80}$ =

21.12, *P* = 6.5 x 10^{-12}, *n* = 3). These findings indicate that the DAT is a key regulator of STP of DA release. At

shortest IPI, DAT promotes STF, whereas at longer IPI, DAT clamps release to promote STD in CPu.

We ruled out potential local anaesthetic actions of cocaine on voltage-gated Na^+ channels (VGSCs),

since lignocaine (10 µM), an inhibitor of VGSCs, did not significantly alter STP (**Fig. S1**), consistent with

reports that cocaine does not inhibit VGSCs at the concentration used in this study (48).

In NAc, cocaine and MPH (we did not test nomifensine in this region) prevented STF at 10 ms IPI (**Fig.**

6G,H, Two-way ANOVA, cocaine x IPI interaction, $F_{4, 30} = 7.884$, *P* = 1.82 x 10^{-4}, *n* = 4; MPH: **Fig 6I,J,** Two-way

ANOVA, MPH x IPI interaction, $F_{4, 20} = 3.986$, *P* = 0.016, *n* = 3) but in contrast to CPu, STD at longer intervals

was not relieved. In NAc, the DAT regulated STP at only short interpulse intervals. Since it is at these intervals

that STP is related to P_r (see Fig 1), these data suggest that it is the effect of DAT inhibition on promoting DA

release that underlies its effects on STP at short intervals in NAc.

We tested whether the effects of DAT inhibition on STP were dependent on synapsin III, since it has

been suggested that this protein might mediate the role of the DAT in vesicle segregation and in limiting

release (27). However, we found that the effects of cocaine persisted in mice lacking synapsin III (**Fig. S2**),

indicating that the DAT regulates STP via a synapsin III-independent or a redundant mechanism.

210   The DAT limits release-dependence of STP in CPu

Since the DAT operates a clamp on DA release in CPu that drives STD, we tested whether the DAT might

prevent STP from being release-dependent. We explored whether inhibition of the DAT could expose a

stronger relationship between $[Ca^{2+}]_o$ and PPR in CPu. We found that whereas there was no effect on PPR of

changing $[Ca^{2+}]_o$ in the absence of cocaine (**Fig. 7A,B**, Two-way ANOVA, P = 0.209, n = 4), in the presence of

cocaine by contrast there was a significant effect of $[Ca^{2+}]_o$: PPR at IPIs of 10-40 ms were significantly

elevated in 1.2 versus 2.4 mM $[Ca^{2+}]_o$ (**Fig. 7A,C-E**, Two-way ANOVA, $[Ca^{2+}]_o$ x IPI interaction, $F_{3, 24}$ = 6.137, P =

0.003, n = 4). There was a significantly stronger effect of $[Ca^{2+}]_o$ on PPR in the presence of cocaine than in

control conditions at 25 ms (**Fig. 7D;** Two-way ANOVA, cocaine x $[Ca^{2+}]_o$ interaction; $F_{1, 12}$ = 7.86; P = 0.016; n

= 4) and 40 ms IPI (**Fig. 7D**; Two-way ANOVA, cocaine x $[Ca^{2+}]_o$ interaction; $F_{1, 12}$ = 4.72, P = 0.050, n = 4), but

not at 10 ms or 100 ms IPI. These effects were not due to a potential electrochemical change in dopamine

adsorption/desorption kinetics at the electrode that might occur with a change in divalent cations, since

these effects of cocaine were maintained when the reduction in $[Ca^{2+}]_o$ was compensated by equimolar

substitution with Mg^{2+} (**Fig. S3**)**.** Inhibition of the DAT therefore relieves the clamp on a relationship between

PPR and initial release, suggesting that the DAT drives STD and limits the release-dependence of STP.

The DAT maintains K$^+$-dependent gating of STP

We noted that the effect of DAT inhibitors on relieving STD resembled the effect of low [K$^+$]$_o$. Since inhibition

of the DAT might prevent DAT-mediated depolarising conductances and lead to axonal hyperpolarisation, we

tested whether cocaine and low [K$^+$]$_o$ relieve STD via an overlapping mechanism i.e. promoting membrane

hyperpolarisation/repolarisation. To test this, we identified whether cocaine precluded the effect of varying

[K$^+$]$_o$ (1 vs 5 mM) on STP, in CPu and NAc.

In CPu, in the absence of cocaine, PPR were elevated in lower [K$^+$]$_o$ (**Fig. 8A,B**, Two-way ANOVA, main

effect [K$^+$]$_o$, $F_{1, 18}$ = 20.47, P = 2.62 x 10^{-4}, n = 4) as seen in Fig. 3. Cocaine increased 1p-evoked peak [DA]$_o$ to a

similar extent in 5 mM versus 1 mM [K$^+$]$_o$ (**Fig. 8C**) but prevented the effect on PPR of varying [K$^+$]$_o$ (**Fig. 8D**,

Two-way ANOVA, [K$^+$]$_o$ x cocaine interaction, $F_{1, 12}$ = 7.686, P = 0.017, n = 8). The limited ability of low [K$^+$]$_o$ to

increase PPR was not due simply to some ceiling effect such as depletion of DA due to increased P1 release

in cocaine, since cocaine also prevented the opposite effects of a raise in [K$^+$]$_o$ (to 7.5 mM) from decreasing

PPR (**Fig. S4**).

In NAc, PPR were slightly elevated in lower [K$^+$]$_o$ (**Fig. 8E,F**, Two-way ANOVA, main effect [K$^+$]$_o$, $F_{1,12}$ =

6.840, P = 0.023, n = 3). Cocaine increased 1p-evoked peak [DA]$_o$ to a similar extent in 5 mM versus 1 mM

[K$^+$]$_o$ (**Fig. 8G**), and, as in CPu, prevented the effect of varying [K$^+$]$_o$ on PPR (**Fig. 8F,H**, Two-way ANOVA, [K$^+$]$_o$ x

cocaine interaction, $F_{1,8}$ = 6.756, P = 0.032, n = 3). The effects of [K$^+$]$_o$ on STP are therefore abolished by

cocaine, suggesting that STP is regulated by a pathway common to [K$^+$]$_o$ and DAT function. DAT function

appears to regulate STP in a manner that supports [K$^+$]$_o$-dependent modulation.

Discussion

We explored the major intrinsic mechanisms operating on DA axons that govern the dynamic probability of

DA release, in dorsal and ventral striatum. We show that STP is governed in only a limited manner by Ca^{2+}-

dependent regulation of release probability, which participates subsequently in determining STF but not

STD, and to greater extent in ventral than dorsal striatum. We reveal that release-independent mechanisms

drive strong STD of release, particularly in dorsal striatum, where axonal excitability and the DAT are major

players and dominate over Ca^{2+}-dependent gating. The DAT appears to be a master regulator that sets DA P_r

and its resulting STF, and clamps release leading to release-independent STD particularly in dorsal striatum.

The differences seen between regions could in turn lead to different DA outputs in dorsal and ventral

striatum in response to changes in AP firing, and could underpin different outcomes of striatal modulatory

inputs.

Calcium-dependent facilitation

Studies of STP at central synapses commonly demonstrate an inverse relationship to P_r; facilitation is

typically observed when P_r is low, and depression when P_r is high (49). Here, we show that, in NAc, an

inverse relationship between STF and P_r could be observed for short IPI corresponding to instantaneous

frequencies seen during fast burst firing (~25 Hz), but the strength of STD at lower frequencies was not

altered. By contrast, in CPu, changes to DA P_r do not strongly influence STP. There seems to be a limited

relationship between factors that influence Ca^{2+} entry, initial P_r as defined here, and the dynamic plasticity of

DA release, especially in CPu, where other release-independent mechanisms are particularly influential.

These findings also suggest that mechanisms governing STF might operate differently between NAc

and CPu. STF at other synapses is often attributed to summation of residual Ca^{2+} (50, 51), or Ca^{2+}-dependent

ultrafast recruitment of vesicles (52), and Ca^{2+} dynamics also vary with local Ca^{2+} buffering mechanisms (53).

In NAc, we saw that Ca^{2+}-dependent STF plays a stronger role in regulating STP than in CPu, despite a weaker

relationship between $[Ca^{2+}]_o$ and P_r than in CPu. These differences in Ca^{2+}-dependence between NAc and

CPu could correspond in part to differences in their Ca^{2+} binding proteins. For example, the high-affinity fast

Ca^{2+} buffer calbindin-D_{28k} is expressed at 2-3-fold greater levels in DA neurons of ventral tegmental area

(VTA) than substantia nigra pars compacta (SNc), which predominantly project to NAc and CPu respectively

(54-56). Saturation of calbindin following initial Ca^{2+} entry could increase P_r at subsequent stimulations

resulting in enhanced STF, as seen in other neurons (57, 58).

Release-insensitive depression

At longer interpulse intervals (40-200 ms) corresponding to a large range of physiological firing frequencies

(5-25 Hz), STP of DA release is dominated by powerful STD. This was not due to D_2 autoreceptor (D2R)

activation. STD of release of other neurotransmitters can result from depletion of readily-releasable vesicles

(59), but the STD seen here for DA does not seem to be due to depletion. Not only is a low fraction of

presynaptic DA estimated to be released after stimulation (13, 16, 60, 61), but moreover STD in the present

study was not relieved by reducing initial release (by $[Ca^{2+}]_o$), inconsistent with a depletion mechanism. VGCC

inactivation by a Ca^{2+} sensor has been proposed to drive STD at some central synapses (62-64) but since low

$[Ca^{2+}]_o$ did not relieve STD for DA release, this mechanism is unlikely to be a critical factor for DA STD.

We show that STD of DA release is not sensitive to the magnitude of initial release (a measure of P_r),

and so can be considered a release-insensitive depression. We found, particularly in CPu, that the strength of

STD was inversely correlated with $[K^+]_o$, consistent with findings at some other central synapses (65). We

found further that we could dissociate mechanisms governing STD from those governing initial release: STD

varied with $[K^+]_o$ despite no change in initial P_r. The $[K^+]_o$-dependent modulation of STP was dependent on K_v

channels, which regulate terminal excitability and rates of repolarisation in several axon types (66-69)

indicating that membrane polarisation is a critical factor in determining STD for DA release.

These findings together suggest that mechanisms regulating Ca^{2+} entry into DA axons will change the

overall amplitude of DA signals, but will not change the dynamic contrast in DA signals when DA neurons

change their firing frequency, except at very highest frequencies in NAc. Conversely, mechanisms that

modulate K^+ conductances or other mechanisms underlying axonal membrane potential and repolarisation,

will particularly influence the dynamic contrast in DA signals. Varying $[K^+]_o$ changes the Nernstian driving

force for K^+ currents (41), altering the rate of repolarisation at DA axons and therefore the degree of Na^+

channel inactivation, but also alters the rate of K^+ channel inactivation (42) leading to use-dependent

changes in membrane potential that will influence Na^+ channel recruitment at the second pulse. We cannot

distinguish here which of these opposing mechanisms dominates to govern STP but in either scenario, Na^+-

channel recruitment will be modified and will govern subsequent release events, leading to varying dynamic

summation of $[DA]_o$. Other mechanisms that influence Na^+ channel availability might also regulate STP; it is

noteworthy in this regard that nAChRs can significantly promote DA release and in turn drive a strong STD

that limits the frequency sensitivity in DA signals (11, 14).

The critical role for axonal excitability in DA STD is particularly pertinent given DA axon morphology.

Midbrain DA neurons form small diameter, unmyelinated, extensively arborised axons each containing on

the order of $\sim 10^4$ branch points and $\sim 10^5$ varicosities (19, 70). These morphological features will not readily

favour reliable conduction of action potentials (71). Changes in $[K^+]_o$ that influence membrane

potential/repolarisation are likely to change axonal conduction failure (65, 72) and may therefore influence

the fidelity and the extent of action potential propagation through the DA axonal tree. AP propagation or

failure could therefore be a key determinant of the STD observed in this study as seen elsewhere (73), and

furthermore could then be a mechanism by which the size of the DA axonal arbour invaded and the

consequent size of the striatal field modulated by DA could be dynamically adapted by presynaptic activity or

neuromodulatory inputs.

Changes in propagation fidelity would also be expected to have a greater limiting effect on release

where axonal arbours are more extensive. The axonal arbours formed by DA neurons projecting to CPu are

more extensive than those projecting to NAc (18, 19, 74), and in parallel, STP showed stronger dependence

on K^+-dependent gating in CPu than in NAc. This association suggests that action potentials in VTA DA

neurons might more readily invade a fuller extent of their axonal tree than action potentials in SNc DA

neurons, a speculation which could have significant implications for the signal processing performed by

these different neurons.

Role of the DAT in STP

Besides curtailing the amplitude and lifetime of $[DA]_o$ through re-uptake (75, 76), DAT function has also been

shown to limit DA release (44, 45). The DAT has been suggested to inhibit vesicle mobilisation via

interactions with SNARES such as synapsins, with synapsin III indicated as a potential candidate (27, 28).

Consistent with a role of the DAT in limiting P_r, our findings reveal that DAT function promotes STF of DA

release seen at very short interpulse intervals (up to 25 ms). At longer intervals, corresponding to typical

firing frequencies (5-25 Hz), when STP is dominated by release-independent depression, we found that the

DAT clamps release which promotes STD in CPu. Our data do not, however, support a key role for the

synapsin III isoform in these DAT functions. Through a different mechanism, the DAT supports both initial P_r,

STF and STD, in contrast to the effects of changing either $[Ca^{2+}]_o$ or $[K^+]_o$ alone, indicating that the DAT limits

initial DA release probability and also ongoing release.

The effects of DAT inhibition on STF were surprisingly large given that, under control conditions, the

relationship between P_r and STF was weak. This disparity is reconciled by our finding that DAT inhibition

unmasked an enhanced Ca^{2+}-dependent modulation of STP, suggesting that the DAT is a critical player that

limits the surprisingly poor relationship between P_r and STP. The mechanism for this interplay is unclear, but

since the DAT is an electrogenic transporter (77, 78) that mediates a depolarising current (25, 26, 47) and

that it can also regulate VGCCs (79), the DAT is likely to influence both axonal excitability and intracellular

Ca^{2+} dynamics in DA axons.

Both the DAT and $[K^+]_o$ acted to clamp release, and apparently operated through a common

mechanism. The effects of the DAT and $[K^+]_o$ in driving STD of DA release could be indicative of depolarising

influences that attenuate conduction propagation through the DA axon arbour, since axonal

hyperpolarisation and low $[K^+]_o$ can actually promote AP conduction in some neurons (80). It is also possible

that the DAT might interact directly with voltage-gated K^+ channels, as suggested from DAT proteome

analyses (81), or indirectly through an interaction with D2Rs (82, 83) which regulate DA release via K^+

channel activity (84).

Thus overall, our findings provide support for the DAT acting to limit vesicular release, and promote

STD, but allowing limited STF at short intervals. DAT action explains the limited expected effects of Ca^{2+}-

dependent release probability on STP. Our data do not support a synapsin3-dependent inhibition of vesicle

recruitment by the DAT, but do not exclude other synapsins that might be involved in DA vesicle recruitment

(27), but might point to a direct action on VGCC function. The overlap with DAT effects and those of K^+ and K_v

channels on STD indicates that the DAT can also act through mechanisms governing excitability that could

include DAT-dependent depolarising conductances and/or Na^{2+} entry that appear to limited Na^+-channel

recruitment and promote STD. It is of note that DATs are widely distributed throughout the length of DA

axons (85), and are found at locations thought to correspond to both release-active zones and inactive zones

(16). Thus DATs could be ideally positioned to act to govern both fiber recruitment for AP propagation in

axons as well as vesicle/VGCC recruitment at active zones. Now defined, these hypotheses should be tested

directly in future studies.

Furthermore, it is well known that DAT function can alter the frequency-dependence of DA release

during ongoing activity (23, 30, 32, 86). These effects are consistent with the effects of uptake on

extracellular summation between stimuli and also with the role of DAT in the dynamic probability of release

we have identified here. The DAT might in turn also be expected to determine the spatial range of striatal DA

transmission not only through its established role in limiting the diffusion of extracellular DA away from

release sites but also by limiting AP conduction within the axonal arbour and/or the probability of DA

release. It is of further note that the apparently greater role for DAT in driving STD in CPu and NAc correlates

with the greater striatal density of DAT at more dorsolateral co-ordinates (87), and with greater DAT

transcript levels in SNc than VTA neurons (56).

In conclusion, we propose a hierarchy of intrinsic mechanisms that control STP of DA release. Within

this hierarchy, the DAT represents a 'master regulator' of STP, governing the balance between release-

dependent and release-independent mechanisms that differently dominate in dorsal versus ventral striatum.

In turn, DAT inhibitors such as cocaine will have profound effects on DA signalling through promoting P_r,

relieving STD, and altering the timecourse and spatial field of DA signals.

Materials and Methods

Animals

Experiments were carried out using adult male C57Bl6/J mice (Jackson Laboratories), heterozygote DAT$^{IRES\text{-}Cre}$ mice conditionally expressing ChR2, or synapsin-III knockout (S3KO) mice on a C57Bl6/J background. For experiments using optical stimulation of DA axons, male heterozygote DAT-internal ribosome entry site (IRES)-Cre mice were bred from homozygote DAT$^{IRES\text{-}Cre}$ mice on a C57Bl6/J background (B6.SJL-*Slc6a3$^{tm1.1(cre)Bkmn}$*/J, stock # 006660, Jackson Laboratories). S3KO mice were bred as described previously (Feng et al., 2002), and were kindly supplied by Professor HT Kao (Brown University). All procedures were performed in accordance with the Animals in Scientific Procedures Act 1986 (Amended 2012) with ethical approval from the University of Oxford, and under authority of a Project Licence granted by the UK Home Office.

Surgery

Heterozygote DAT$^{IRES\text{-}Cre}$ mice were injected intracerebrally with a Cre-inducible recombinant AAV serotype 5 vector containing an inverted gene for channelrhodopsin-2 fused in-frame with a gene encoding enhanced yellow fluorescent protein (pAAV-double floxed-hChR2(H134R)-EYFP-WPRE-pA). Mice were placed in a stereotaxic frame under isoflurane anaesthesia and a craniotomy was made above the injection site. Injections of 1 μL virus were given either unilaterally or bilaterally in either VTA (co-ordinates from Bregma in mm: AP -3.1, ML ±0.5, DV -4.4) or in the SNc (from Bregma in mm: AP -3.5, ML ±1.2, DV -4.0) using a 2.5 μL 33-gauge Hamilton syringe at 0.2 μL/min with a microinjector. Animals were maintained for at least 3 weeks following surgery to allow virus expression in striatum.

Slice Preparation

Striatal slices were prepared as described previously (31, 88). Mice were sacrificed by cervical dislocation and the brains removed and transferred to ice-cold HEPES-based buffer containing in mM: 120 NaCl, 20 NaHCO$_3$,

6.7 HEPES acid, 5 KCl, 3.3 HEPES salt, 2 $CaCl_2$, 2 $MgSO_4$, 1.2 KH_2PO_4, 10 glucose, saturated with 95%O_2/5%CO_2.

Acute 300 µm thick coronal striatal slices, containing both dorsal striatum (CPu) and nucleus accumbens core

(NAc) were prepared in ice-cold HEPES-based buffer and cut using a vibratome (VT1000S or VT1200S; Leica).

Slices were kept at room temperature in HEPES-based buffer for 1 hour before being transferred to the

recording chamber and superfused at 1.8-2.0 ml/min in bicarbonate buffer-based artificial CSF (aCSF)

containing in mM: 124.3 NaCl, 26 $NaHCO_3$, 3.8 KCl, 2.4 $CaCl_2$, 1.3 $MgSO_4$, 1.2 KH_2PO_4, 10 glucose, saturated

with 95% O_2/5% CO_2, at 31-33^{o}C. In all experiments, recordings were carried out in the presence of bath-

applied dihydro-β-erythroidine (DHβE, 1 µM), an antagonist at β2 subunit-containing nAChRs. Slices were

allowed to equilibrate for 30 minutes prior to recording.

Voltammetry

Evoked extracellular DA concentration ($[DA]_o$) was measured using FCV at carbon-fibre microelectrodes (fibre

diameter 7-10 µm, tip length 50-120 µm). A triangular voltage waveform was scanned across the

microelectrode (-700 to +1300 mV and back vs Ag/AgCl reference, scan rate 800 V/s) using a Millar

voltammeter (Julian Millar, Barts and the London School of Medicine and Dentistry), with a sweep frequency

of 8 Hz, as described previously (31). This sampling rate is sufficient to capture the rising and falling phase of

the DA transients; faster sampling rates do not change the data interpretations (not illustrated). Evoked

currents were confirmed as DA by comparison of the voltammogram with that produced during calibration

with applied DA in aCSF (oxidation peak +500-600 mV and reduction peak -200 mV). Currents at the oxidation

peak potential were measured from the baseline of each voltammogram and plotted against time to provide

profiles of $[DA]_o$ versus time. Electrodes were calibrated in 2 µM DA in each experimental solution used,

including all solutions where $[Ca^{2+}]$ was varied, since electrode sensitivity varies with divalent ion

concentration (89). Calibration solutions were made immediately before use from stock solution of 2.5 mM

DA in 0.1 M $HClO_4$ stored at 4^{o}C.

Stimulation and experimental design

For experiments using electrical stimulation, DA release was evoked using a surface bipolar concentric Pt/Ir electrode (25 µm diameter, FHC) placed approximately 100 µm from the recording electrode. Stimulation pulses of 200 µs duration were applied at 0.6 mA unless described otherwise. In experiments using tissue from DAT[IRES-Cre] mice conditionally expressing ChR2 in DA axons, DA release was evoked with 2 ms full-field illumination of the slice with an LED system emitting light at 470 nm wavelength (OptoLED, Cairn Research) as described previously (11). The LED system illuminated an area of 2.2 mm diameter. The current delivered by the LED power supply was set to produce a perimaximal light intensity (i.e. the minimum light intensity able to evoke maximum $[DA]_o$ following a single 2 ms pulse). Since the perimaximal light intensity is dependent on the level of ChR2 expression, which varies between animals, the appropriate current was determined at the beginning of each experiment. Electrical or optical stimulations were delivered every 2.5 minutes, after which dopamine release was fully recovered.

Release probability P_r for DA as defined here is a composite measure of synaptic P_r (a function of vesicular P_r and the size of the pool of vesicles) and the number of release sites and fibres recruited by the stimulus. Short-term plasticity (STP) in DA release was explored by applying alternating single pulses (1p) or paired pulses (2p) with inter-pulse intervals (IPIs) of 10 ms – 200 ms in pseudorandom order and in triplicate at each recording site. IPIs of 40 to 200 ms fall within the range commonly observed during burst firing in DA neurons *in vivo* (3, 4). IPIs of 10 to 25 ms IPIs have been observed during burst firing in DA neurons (Hyland et al., 2002) but are particularly useful to interrogate short-term facilitation which occurs on this timescale (14, 23). We calculated paired-pulse ratio (PPR) as a measure of STP. We define PPR as the ratio P2/P1, where P1 is peak $[DA]_o$ detected following 1p stimulus and P2 is the peak $[DA]_o$ attributable to the second stimulation. P2 was determined by subtracting the $[DA]_o$ transient including decay phase after a single pulse from the summated paired-pulse recording, and therefore DA decay kinetics are accounted for. We cannot control for variable fiber recruitment at different stimuli, but can control for variable modulation by other

local inputs: PPR on the timecourses explored here is not modified by D2 dopamine receptors (see Results),

or by GABA receptors (Lopes and Cragg, unpublished observations).

Calcium concentration ($[Ca^{2+}]_o$) in aCSF was varied by varying $[CaCl_2]$. In some experiments where

stated, a reduction in $[CaCl_2]$ was substituted with an equimolar quantity of $MgCl_2$. However, Ca^{2+}/Mg^{2+}

substitution was not routine because other effects of $[Mg^{2+}]_o$ can confound interpretation. Changes in $[Mg^{2+}]$

have been shown to alter Ca^{2+} currents through VGCCs (90, 91), alter G-protein-coupled receptor function,

including dopamine D2 receptors (92), change the affinity of calbindin binding to Ca^{2+} (93), trigger

complexing of Ca^{2+} by ATP (94), and alter the sensitivity of FCV electrodes to DA (89).

Where extracellular potassium concentration ($[K^+]_o$) was varied (1.0 - 7.5 mM), changes in osmolarity

were corrected by varying $[Na^+]$. The Na^+- and K^+-containing salts present in aCSF were varied as follows (in

mM): In experiments with 1 mM $[K^+]_o$, salts were 127.1 NaCl, 1.0 KCl, 0 KH_2PO_4, 1.2 NaH_2PO_4; For 1.25 mM

$[K^+]_o$ salts were 128.0 NaCl, 0.05 KCl, 1.2 KH_2PO_4; For 7.5 mM $[K^+]_o$, salts were 121.8 NaCl, 6.3 KCl and 1.2

KH_2PO_4. Recordings in control conditions were collected before and after each experimental manipulation at

each recording site. Slices were equilibrated in each solution for at least 20 minutes before data were

included for analysis. Experiments were carried out in either the dorsal half of the caudate-putamen (CPu) or

nucleus accumbens core (NAc), one site per slice. Release is TTX-sensitive.

Data analysis

Data were acquired and analysed using Axoscope 10.2, 10.5 (Molecular Devices) or Strathclyde Whole Cell

Program (University of Strathclyde, Glasgow, UK) and locally-written scripts in Visual Basic for Applications

(Microsoft). Data are expressed as mean ± standard error (SEM) and n = number of animals (the number of

biological replicates). Within a given experiment, technical replicates were usually obtained in at least

triplicate before averaging to obtain the value for that experiment. Statistical comparisons were carried out

using GraphPad Prism 6 (GraphPad Software Inc.).

Drugs

DHβE and lignocaine were obtained from Tocris Biosciences or Ascent Scientific. Cocaine hydrochloride and

nomifensine maleate were obtained from Sigma. Methylphenidate (Ritalin™) was obtained from Novartis. All

drugs were dissolved in either de-ionised water, aqueous acid (nomifensine maleate) or ethanol (lignocaine)

to make stock aliquots at 1000-10,000x final concentration and stored at -20^{o}C. The final concentration was

made up in aCSF immediately before use.

Acknowledgements: Funding sources: MRC, BBSRC Interdisciplinary Doctoral Training Program, Clarendon

Fund, Monument Trust Parkinson's UK Discovery Award, and Christ Church Oxford.

Author contributions: M.D.C. designed and performed experiments, analysed and interpreted the data, and

co-wrote the manuscript. N.J.P. designed and performed experiments, and analysed and interpreted the

data. B.M.R, S.V.M. and M.Y.T performed experiments and analysed data. S.T. and S.J.C. supervised the

work. S.J.C. designed the experiments, interpreted the data, and co-wrote the manuscript.

Declaration of Interests: The authors declare no competing interests.

References

1. Cohen JY, Haesler S, Vong L, Lowell BB, Uchida N. Neuron-type-specific signals for reward and punishment in the ventral tegmental area. Nature. 2012;482(7383):85-8.

2. Grace AA, Bunney BS. The control of firing pattern in nigral dopamine neurons: single spike firing. J Neurosci. 1984;4(11):2866-76.

3. Grace AA, Bunney BS. The control of firing pattern in nigral dopamine neurons: burst firing. J Neurosci. 1984;4(11):2877-90.

4. Hyland BI, Reynolds JNJ, Hay J, Perk CG, Miller R. Firing modes of midbrain dopamine cells in the freely moving rat. Neuroscience. 2002;114(2):475-92.

5. Pan WX, Schmidt R, Wickens JR, Hyland BI. Dopamine Cells Respond to Predicted Events during Classical Conditioning: Evidence for Eligibility Traces in the Reward-Learning Network. J Neurosci. 2005;25(26):6235-42.

6. Jin X, Costa RM. Start/stop signals emerge in nigrostriatal circuits during sequence learning. Nature. 2010;466(7305):457-62.

7. Matsumoto M, Hikosaka O. Two types of dopamine neuron distinctly convey positive and negative motivational signals. Nature. 2009;459(7248):837-41.

8. Schultz W. Responses of midbrain dopamine neurons to behavioral trigger stimuli in the monkey. Journal of neurophysiology. 1986;56(5):1439-61.

9. Sulzer D, Cragg SJ, Rice ME. Striatal dopamine neurotransmission: Regulation of release and uptake. Basal Ganglia. 2016;6(3):123-48.

10. Hamid AA, Pettibone JR, Mabrouk OS, Hetrick VL, Schmidt R, Vander Weele CM, et al. Mesolimbic dopamine signals the value of work. Nat Neurosci. 2016;19(1):117-26.

11. Threlfell S, Lalic T, Platt NJ, Jennings KA, Deisseroth K, Cragg SJ. Striatal dopamine release is triggered by synchronized activity in cholinergic interneurons. Neuron. 2012;75(1):58-64.

12. Cachope R, Mateo Y, Mathur BN, Irving J, Wang HL, Morales M, et al. Selective activation of cholinergic interneurons enhances accumbal phasic dopamine release: setting the tone for reward processing. Cell Rep. 2012;2(1):33-41.

13. Cragg SJ. Variable Dopamine Release Probability and Short-Term Plasticity between Functional Domains of the Primate Striatum. J Neurosci. 2003;23(10):4378-85.

14. Rice ME, Cragg SJ. Nicotine amplifies reward-related dopamine signals in striatum. Nat Neurosci. 2004;7(6):583-4.

15. Montague PR, McClure SM, Baldwin PR, Phillips PEM, Budygin EA, Stuber GD, et al. Dynamic gain control of dopamine delivery in freely moving animals. J Neurosci. 2004;24(7):1754-9.

16. Liu C, Kershberg L, Wang J, Schneeberger S, Kaeser PS. Dopamine Secretion Is Mediated by Sparse Active Zone-like Release Sites. Cell. 2018;172(4):706-18.e15.

17. Debanne D. Information processing in the axon. Nature Rev Neurosci. 2004;5(4):304-16.

18.   Aransay A, Rodríguez-López C, García-Amado M, Clascá F, Prensa L. Long-range projection neurons of
the mouse ventral tegmental area: a single-cell axon tracing analysis. Front Neuroanat. 2015;9:59-.

19.   Matsuda W, Furuta T, Nakamura KC, Hioki H, Fujiyama F, Arai R, et al. Single nigrostriatal
dopaminergic neurons form widely spread and highly dense axonal arborizations in the neostriatum. J
Neurosci. 2009;29(2):444-53.

20.   Collins AL, Aitken TJ, Greenfield VY, Ostlund SB, Wassum KM. Nucleus Accumbens Acetylcholine
Receptors Modulate Dopamine and Motivation. Neuropsychopharmacol. 2016;41:2830-8.

21.   Zhang H, Sulzer D. Frequency-dependent modulation of dopamine release by nicotine. Nat Neurosci.
2004;7(6):581-2.

22.   Wang L, Shang S, Kang X, Teng S, Zhu F, Liu B, et al. Modulation of dopamine release in the striatum
by physiologically relevant levels of nicotine. Nat Commun. 2014;5:3925.

23.   Jennings KA, Platt NJ, Cragg SJ. The impact of a parkinsonian lesion on dynamic striatal dopamine
transmission depends on nicotinic receptor activation. Neurobiol Dis. 2015;82:262-8.

24.   Brimblecombe KR, Gracie CJ, Platt NJ, Cragg SJ. Gating of dopamine transmission by calcium and
axonal N-, Q-, T- and L-type voltage-gated calcium channels differs between striatal domains. J Physiol.
2015;593(4):929-46.

25.   Carvelli L, McDonald PW, Blakely RD, Defelice LJ. Dopamine transporters depolarize neurons by a
channel mechanism. Proc Natl Acad Sci USA. 2004;101(45):16046-51.

26.   Ingram SL, Prasad BM, Amara SG. Dopamine transporter-mediated conductances increase
excitability of midbrain dopamine neurons. Nat Neurosci. 2002;5(10):971-8.

27.   Kile BM, Guillot TS, Venton BJ, Wetsel WC, Augustine GJ, Wightman RM. Synapsins differentially
control dopamine and serotonin release. J Neurosci. 2010;30(29):9762-70.

28.   Venton BJ, Seipel AT, Phillips PEM, Wetsel WC, Gitler D, Greengard P, et al. Cocaine Increases
Dopamine Release by Mobilization of a Synapsin-Dependent Reserve Pool. Journal of Neuroscience.
2006;26(12):3206-9.

29.   Gonon FG. Nonlinear relationship between impulse flow and dopamine released by rat midbrain
dopaminergic neurons as studied by in vivo electrochemistry. Neuroscience. 1988;24(1):19-28.

30.   Garris PA, Wightman RM. Different kinetics govern dopaminergic transmission in the amygdala,
prefrontal cortex, and striatum: an in vivo voltammetric study. J Neurosci. 1994;14(1):442-50.

31.   Threlfell S, Clements MA, Khodai T, Pienaar IS, Exley R, Wess J, et al. Striatal muscarinic receptors
promote activity dependence of dopamine transmission via distinct receptor subtypes on cholinergic
interneurons in ventral versus dorsal striatum. J Neurosci. 2010;30(9):3398-408.

32.   Zhang L, Doyon WM, Clark JJ, Phillips PEM, Dani JA. Controls of Tonic and Phasic Dopamine
Transmission in the Dorsal and Ventral Striatum. Molec Pharmacol. 2009;76(2):396-404.

33.   Debanne D, Guérineau NC, Gähwiler BH, Thompson SM. Paired-pulse facilitation and depression at
unitary synapses in rat hippocampus: quantal fluctuation affects subsequent release. J Physiol.
1996;491(1):163-76.

34.   Dobrunz LE, Stevens CF. Heterogeneity of Release Probability, Facilitation, and Depletion at Central
Synapses. Neuron. 1997;18(6):995-1008.

35.   Thomson AM. Molecular frequency filters at central synapses. ProgNeurobiol. 2000;62(2):159-96.

36.   Wilcox KS, Dichter MA. Paired pulse depression in cultured hippocampal neurons is due to a
presynaptic mechanism independent of GABAB autoreceptor activation. J Neurosci. 1994;14(3 Pt 2):1775-88.

37.   Fisher SA, Fischer TM, Carew TJ. Multiple overlapping processes underlying short-term synaptic
enhancement. Trends Neurosci. 1997;20(4):170-7.

38.   Benoit-Marand M, Borrelli E, Gonon F. Inhibition of dopamine release via presynaptic D2 receptors:
time course and functional characteristics in vivo. J Neurosci. 2001;21(23):9134-41.

39.   Phillips PEM, Hancock PJ, Stamford JA. Time window of autoreceptor-mediated inhibition of limbic
and striatal dopamine release. Synapse. 2002;44(1):15-22.

40.   Schmitz Y, Schmauss C, Sulzer D. Altered dopamine release and uptake kinetics in mice lacking D2
receptors. J Neurosci. 2002;22(18):8002-9.

41.   Balestrino M, Aitken PG, Somjen GG. The effects of moderate changes of extracellular K+ and Ca2+
on synaptic and neural function in the CA1 region of the hippocampal slice. Brain Res. 1986;377(2):229-39.

42.   Baukrowitz T, Yellen G. Modulation of K+ current by frequency and external [K+]: A tale of two
inactivation mechanisms. Neuron. 1995;15(4):951-60.

43.   Nedergaard S. Regulation of Action Potential Size and Excitability in Substantia Nigra Compacta
Neurons: Sensitivity to 4-Aminopyridine. Journal of Neurophysiology. 1999;82(6).

44.   Hoffman AF, Spivak CE, Lupica CR. Enhanced Dopamine Release by Dopamine Transport Inhibitors
Described by a Restricted Diffusion Model and Fast-Scan Cyclic Voltammetry. ACS Chem Neurosci.
2016;7:700-9.

45.   Lee TH, Balu R, Davidson C, Ellinwood EH. Differential time-course profiles of dopamine release and
uptake changes induced by three dopamine uptake inhibitors. Synapse. 2001;41(4):301-10.

46.   Stamford JA, Kruk ZL, Millar J. Dissociation of the actions of uptake blockers upon dopamine
overflow and uptake in the rat nucleus accumbens: In vivo voltammetric data. Neuropharmacology.
1989;28(12):1383-8.

47.   Sonders MS, Zhu S-J, Zahniser NR, Kavanaugh MP, Amara SG. Multiple Ionic Conductances of the
Human Dopamine Transporter: The Actions of Dopamine and Psychostimulants. J Neurosci. 1997;17(3):960-
74.

48.   Acevedo-Rodriguez A, Zhang L, Zhou F, Gong S, Gu H, De Biasi M, et al. Cocaine inhibition of nicotinic
acetylcholine receptors influences dopamine release. Front Synaptic Neurosci. 2014;6:19-.

49.   Zucker RS, Regehr WG. Short-term synaptic plasticity. Ann Review Physiol. 2002;64:355-405.

50.   Chen C, Regehr WG. Contributions of residual calcium to fast synaptic transmission. J Neurosci.
1999;19(15):6257-66.

51.   Katz B, Miledi R. The role of calcium in neuromuscular facilitation. J Physiol. 1968;195(2):481-92.

52.   Doussau F, Schmidt H, Dorgans K, Valera AM, Poulain B, Isope P. Frequency-dependent mobilization
of heterogeneous pools of synaptic vesicles shapes presynaptic plasticity. Elife. 2017;6.

53.   Regehr WG. Short-term presynaptic plasticity. Cold Spring Harbor Perspect Biol. 2012;4(7):a005702-
a.

54.   Chung CY, Seo H, Sonntag KC, Brooks A, Lin L, Isacson O. Cell type-specific gene expression of
midbrain dopaminergic neurons reveals molecules involved in their vulnerability and protection. Hum Molec
Genet. 2005;14(13):1709-25.

55.   Greene JG, Dingledine R, Greenamyre JT. Gene expression profiling of rat midbrain dopamine
neurons: implications for selective vulnerability in parkinsonism. Neurobiol Dis. 2005;18(1):19-31.

56.   Haber SN, Ryoo H, Cox C, Lu W. Subsets of midbrain dopaminergic neurons in monkeys are
distinguished by different levels of mRNA for the dopamine transporter: Comparison with the mRNA for the
D2 receptor, tyrosine hydroxylase and calbindin immunoreactivity. J Comp Neurol. 1995;362(3):400-10.

57.   Blatow M, Caputi A, Burnashev N, Monyer H, Rozov A. Ca2+ Buffer Saturation Underlies Paired Pulse
Facilitation in Calbindin-D28k-Containing Terminals. Neuron. 2003;38(1):79-88.

58.   Felmy F, Neher E, Schneggenburger R. Probing the Intracellular Calcium Sensitivity of Transmitter
Release during Synaptic Facilitation. Neuron. 2003;37(5):801-11.

59.   Stevens CF, Williams JH. Discharge of the Readily Releasable Pool With Action Potentials at
Hippocampal Synapses. J Neurophys. 2007;98(6):3221-9.

60.   Dreyer JK, Herrik KF, Berg RW, Hounsgaard JD. Influence of Phasic and Tonic Dopamine Release on
Receptor Activation. J Neurosci. 2010;30(42):14273-83.

61.   Rooney KE, Wallace LJ. Computational modeling of extracellular dopamine kinetics suggests low
probability of neurotransmitter release. Synapse. 2015;69(11):515-25.

62.   Nanou E, Sullivan JM, Scheuer T, Catterall WA. Calcium sensor regulation of the CaV2.1 Ca2+ channel
contributes to short-term synaptic plasticity in hippocampal neurons. Proc Natl Acad Sci USA.
2016;113(4):1062-7.

63.   Sheng J, He L, Zheng H, Xue L, Luo F, Shin W, et al. Calcium-channel number critically influences
synaptic strength and plasticity at the active zone. Nat Neurosci. 2012;15(7):998-1006.

64.   Xu J, Wu L-G. The decrease in the presynaptic calcium current is a major cause of short-term
depression at a calyx-type synapse. Neuron. 2005;46(4):633-45.

65.   He Y, Zorumski CF, Mennerick S. Contribution of Presynaptic Na+ Channel Inactivation to Paired-
Pulse Synaptic Depression in Cultured Hippocampal Neurons. J Neurophysiol. 2002;87(2):925-36.

66.   Geiger JRP, Jonas P. Dynamic Control of Presynaptic Ca2+ Inflow by Fast-Inactivating K+ Channels in
Hippocampal Mossy Fiber Boutons. Neuron. 2000;28(3):927-39.

67.   Ishikawa T, Nakamura Y, Saitoh N, Li W-B, Iwasaki S, Takahashi T. Distinct Roles of Kv1 and Kv3
Potassium Channels at the Calyx of Held Presynaptic Terminal. J Neurosci. 2003;23(32):10445-53.

68.   Kawaguchi S-y, Sakaba T. Control of inhibitory synaptic outputs by low excitability of axon terminals
revealed by direct recording. Neuron. 2015;85(6):1273-88.

69.   Ueda A, Wu C-F. Distinct Frequency-Dependent Regulation of Nerve Terminal Excitability and
Synaptic Transmission by IA and IK Potassium Channels Revealed by Drosophila Shaker and Shab Mutations. J
Neurosci. 2006;26(23):6238-48.

70.   Bolam JP, Pissadaki EK. Living on the edge with too many mouths to feed: Why dopamine neurons
die. Mov Disord. 2012;27(12):1478-83.

71.   Ofer N, Shefi O, Yaari G. Branching morphology determines signal propagation dynamics in neurons.
Scientific Reports. 2017;7(1):8877-.

72.   Brody DL, Yue DT. Release-Independent Short-Term Synaptic Depression in Cultured Hippocampal
Neurons. J Neurosci. 2000;20(7):2480-94.

73.   Debanne D, Campanac E, Bialowas A, Carlier E, Alcaraz G. Axon physiology. Physiol Rev.
2011;91(2):555-602.

74.   Pacelli C, Giguère N, Bourque M-J, Lévesque M, Slack Ruth S, Trudeau L-É. Elevated Mitochondrial
Bioenergetics and Axonal Arborization Size Are Key Contributors to the Vulnerability of Dopamine Neurons.
Current Biology. 2015;25(18):2349-60.

75.   Jones SR, Gainetdinov RR, Wightman RM, Caron MG. Mechanisms of amphetamine action revealed
in mice lacking the dopamine transporter. Journal of Neuroscience. 1998;18(6):1979-86.

76.   Mateo Y, Budygin EA, John CE, Banks ML, Jones SR. Voltammetric assessment of dopamine clearance
in the absence of the dopamine transporter: no contribution of other transporters in core or shell of nucleus
accumbens. J Neurosci Meth. 2004;140(1-2):183-7.

77.   Gu H, Wall SC, Rudnick G. Stable expression of biogenic amine transporters reveals differences in
inhibitor sensitivity, kinetics, and ion dependence. J Biol Chem. 1994;269(10):7124-30.

78.   McElvain JS, Schenk JO. A multisubstrate mechanism of striatal dopamine uptake and its inhibition
by cocaine. Biochem Pharmacol. 1992;43(10):2189-99.

79.   Cameron KN, Solis E, Ruchala I, De Felice LJ, Eltit JM. Amphetamine activates calcium channels
through dopamine transporter-mediated depolarization. Cell Calcium. 2015;58(5):457-66.

80.   Smith DO. Mechanisms of action potential propagation failure at sites of axon branching in the
crayfish. J Physiol. 1980;301:243-59.

81.   Maiya R, Ponomarev I, Linse KD, Harris RA, Mayfield RD. Defining the dopamine transporter
proteome by convergent biochemical and in silico analyses. Genes Brain Behav. 2007;6(1):97-106.

82.   Benoit-Marand M, Ballion B, Borrelli E, Boraud T, Gonon F. Inhibition of dopamine uptake by D2
antagonists: an in vivo study. J Neurochem. 2011;116(3):449-58.

83.   Lee FJS, Pei L, Moszczynska A, Vukusic B, Fletcher PJ, Liu F. Dopamine transporter cell surface
localization facilitated by a direct interaction with the dopamine D2 receptor. EMBO J. 2007;26(8):2127-36.

84.   Martel P, Leo D, Fulton S, Bérard M, Trudeau L-E. Role of Kv1 potassium channels in regulating
dopamine release and presynaptic D2 receptor function. PLoS One. 2011;6(5):e20402-e.

85. Nirenberg MJ, Vaughan RA, Uhl GR, Kuhar MJ, Pickel VM. The dopamine transporter is localized to
dendritic and axonal plasma membranes of nigrostriatal dopaminergic neurons. Journal of Neuroscience.
1996;16(2):436-47.

86. Suaud-Chagny MF. In vivo monitoring of dopamine overflow in the central nervous system by
amperometric techniques combined with carbon fibre electrodes. Methods. 2004;33(4):322-9.

87. Marshall JF, O'Dell SJ, Navarrete R, Rosenstein AJ. Dopamine high-affinity transport site topography
in rat brain: Major differences between dorsal and ventral striatum. Neuroscience. 1990;37(1):11-21.

88. Exley R, McIntosh JM, Marks MJ, Maskos U, Cragg SJ. Striatal alpha5 nicotinic receptor subunit
regulates dopamine transmission in dorsal striatum. J Neurosci. 2012;32(7):2352-6.

89. Kume-Kick J, Rice ME. Dependence of dopamine calibration factors on media Ca^{2+} and Mg^{2+} at
carbon-fiber microelectrodes used with fast-scan cyclic voltammetry. J Neurosci Meth. 1998;84(1-2):55-62.

90. Agus ZS, Kelepouris E, Dukes I, Morad M. Cytosolic magnesium modulates calcium channel activity in
mammalian ventricular cells. Am J Physiol - Cell Physiol. 1989;256(2).

91. Lansman JB, Hess P, Tsien RW. Blockade of current through single calcium channels by Cd^{2+}, Mg^{2+},
and Ca^{2+}. Voltage and concentration dependence of calcium entry into the pore. J Gen Physiol.
1986;88(3):321-47.

92. van der Westhuizen ET, Valant C, Sexton PM, Christopoulos A. Endogenous allosteric modulators of
G protein-coupled receptors. J Pharmacol Exp Therapeut. 2015;353(2):246-60.

93. Berggård T, Miron S, Onnerfjord P, Thulin E, Åkerfeldt KS, Enghild JJ, et al. Calbindin D28k Exhibits
Properties Characteristic of a Ca^{2+} Sensor. J Biol Chem. 2002;277(19):16662-72.

94. Michailova A, McCulloch AD. Model Study of ATP and ADP Buffering, Transport of Ca^{2+} and Mg^{2+},
and Regulation of Ion Pumps in Ventricular Myocyte. Biophys J. 2001;81:614-29.

Figure Legends

Figure 1. Short-term plasticity at DA release sites is region-specific and only weakly release-dependent.

(A, B) Mean profiles of $[DA]_o$ transients ($\pm$ SEM) elicited by single or paired electrical pulses at IPI of 10-200 ms in CPu (A, n = 4) and NAc (B, n = 3). Dotted traces, $[DA]_o$ attributable to the second pulse, P2, in a pair (P1+P2 – P1). **(C)** Mean P2/P1 ($\pm$ SEM) vs. IPIs in CPu (*filled circles*) and NAc (*open circles*). **(D)** P2/P1 vs peak 1p-evoked $[DA]_o$ for IPIs of 10 ms (*left*), 40 ms (*centre*) and 100 ms (*right*). Each data point is 1 site in CPu (*filled circles*, n = 35) or NAc (*open circles*, n = 19). Linear fits to data show *P*-values (F-test) to indicate whether the slope is significantly > 0. **(E,F)** (*Left)* Mean profiles of $[DA]_o$ transients in CPu (E, n = 5) and NAc (F, n = 6) in control conditions vs L741,626 (1 μM, *green*). (*Right*) Mean P2/P1 vs IPIs in control conditions and L741,626 in CPu (E) or NAc (F). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons: *$P < 0.05$, **$P < 0.01$.

Figure 2. STP is shaped by release-dependent facilitation and release-independent depression. (A, B) Mean profiles of $[DA]_o$ transients ($\pm$ SEM) evoked by single or paired electrical pulses in 1.2 mM (*light grey*), 2.4 mM (*dark grey*) and 3.6 mM (*black*) $[Ca^{2+}]_o$ in CPu (A, n = 3) and NAc (B, n = 3). **(C)** Mean 1p evoked $[DA]_o$ ($\pm$ SEM, normalised to 2.4 mM $[Ca^{2+}]_o$) vs $[Ca^{2+}]_o$ in CPu (*filled*) and NAc (*open circles*). **(D, E)** Mean P2/P1 ($\pm$ SEM) vs. IPI in 1.2 mM (*light grey*), 2.4 mM (*dark grey*) and 3.6 mM (*black*) $[Ca^{2+}]_o$ in CPu (D, *filled*) and NAc (D, *open circles*). **(F)** Mean P2/P1 ($\pm$ SEM) vs mean peak 1p $[DA]_o$ ($\pm$ SEM) at 10 ms IPI in 1.2 mM (*light grey*), 2.4 mM (*dark grey*) and 3.6 mM (*black*) $[Ca^{2+}]_o$ in CPu (*filled*) and NAc (*open circles*). **(G)** Mean profiles of $[DA]_o$ transients ($\pm$ SEM) elicited by single or paired light pulses at 40 ms IPI (473 nm, 2 ms) in 1.0 mM (*light grey*), 2.0 mM (*dark grey*) and 3.0 mM (*black*) $[Ca^{2+}]_o$ in CPu (*left*, n = 3) and NAc (*right*, n = 3). **(H)** Mean peak 1p $[DA]_o$ ($\pm$ SEM, normalised to 2.4 mM $[Ca^{2+}]_o$) vs $[Ca^{2+}]_o$ evoked by light stimulation in CPu (*filled*) and NAc (*open circles*). Second-order polynomial curve fit; F-test of relative sum of squares shows curve fits are different. **(I)** Mean P2/P1 ($\pm$ SEM) at 40 ms IPI vs mean peak 1p $[DA]_o$ ($\pm$ SEM) evoked by optical stimulation in CPu (*filled*) and NAc (*open circles*). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons except where stated otherwise: *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 3. A release-insensitive mechanism of short-term depression is controlled by [K$^+$]-dependent gating.

(A, B) Mean profiles of [DA]$_o$ (± SEM) elicited by single or paired electrical pulses in 1.25 mM (*light grey*), 5 mM

(*dark grey*) and 7.5 mM [K$^+$]$_o$ (*black*) in CPu (A, *n* = 3) and NAc (B, *n* = 3). **(C, D)** Mean P2/P1 (± SEM) vs IPIs in

1.25 mM (*light grey*), 5 mM (*dark grey*) and 7.5 mM [K$^+$]$_o$ (*black*) in CPu (C) and NAc (D). **(E)** Mean P2/P1 (±

SEM) vs. mean peak 1p [DA]$_o$ (± SEM) in 1.25 mM (*light grey*), 5 mM (*dark grey*) and 7.5 mM (*black*) [K$^+$]$_o$ in

CPu (*filled*) and NAc (*open circles*) at 10 ms IPI (*left*) and 100 ms IPI (*right*). Two-way ANOVA with Bonferroni's

test for *post-hoc* comparisons; *$P < 0.05$, ***$P < 0.001$.

Figure 4. K$^+$-dependent gating of short-term depression requires axonal Kv channels. (A, E) Mean profiles of

[DA]$_o$ transients (± SEM) in CPu elicited by single or paired electrical pulses in control conditions (*black*) or with

100 μM 4-AP (*grey*) in CPu (A, *n* = 4) and NAc (E, *n* = 3). Vertical scale is normalised to peak 1p [DA]$_o$ in control

conditions. **(B, F)** Mean 1p [DA]$_o$ (± SEM) in CPu in 1 mM and 5 mM [K$^+$]$_o$ in control conditions (*black*) or 4-AP

(*grey*) in CPu (B) or in NAc (F). **(C, G)** Mean P2/P1 (± SEM) in CPu vs IPI in 5 mM (*black*) and 1 mM (*grey*) [K$^+$]$_o$

in control conditions (*left*) and 4-AP (*right*) in CPu (C) and in NAc (G). **(D, H)** Mean P2/P1 (± SEM) in CPu at 25

ms IPI in 1 mM and 5 mM [K$^+$]$_o$ in control conditions (*black*) or 4-AP (*grey*) in CPu (D) and in NAc (H). (D),

asterisks indicate interaction between [K$^+$]$_o$ and 4-AP. Two-way ANOVA with Bonferroni's test for *post-hoc*

comparisons; *$P < 0.05$, **$P < 0.01$.

Figure 5. K$^+$-dependent gating does not alter release-dependence of STP. (A,B,E,F) Mean profiles of [DA]$_o$

transients (± SEM) elicited by single or paired pulses in 1.2 mM [Ca^{2+}] (*light grey*), 2.4 mM [Ca^{2+}]$_o$ (*dark grey*)

and 3.6 mM [Ca^{2+}]$_o$ (*black*) in either 5 mM [K$^+$]$_o$ (*A,E*) or 1 mM [K$^+$]$_o$ (*B,F*) in CPu (A,B, *n* = 11) and in NAc (E,F, *n*

= 7). **(C, G)** Mean peak 1p [DA]$_o$ (± SEM, normalised to 2.4 mM [Ca^{2+}]$_o$) in 5 mM [K$^+$]$_o$ (*black*) and 1 mM [K$^+$]$_o$

(*grey*) in CPu (C) and in NAc (G). **(D, H)** Mean P2/P1 (± SEM) at 10 ms IPI (*left*) and 100 ms IPI (*right*) in 5 mM

[K$^+$]$_o$ (*black*) and 1 mM [K$^+$]$_o$ (*grey*) in CPu (D) and in NAc (H). Two-way ANOVA with Bonferroni's test for *post-*

hoc comparisons.

Figure 6. The dopamine transporter regulates STP of dopamine release. (A, G) Mean profiles of $[DA]_o$

transients (± SEM) elicited by single or paired electrical pulses in control conditions (*black*) or with 5 µM

cocaine (*blue*) in CPu (A, $n = 4$) or NAc (G, $n = 4$). **(B, H)** Mean P2/P1 (± SEM) against IPI in control conditions

(*black*) and with cocaine (*blue*) in CPu (B) or NAc (H). **(C, I)** Mean profiles of $[DA]_o$ transients (± SEM) elicited by

single or paired electrical pulses in control conditions (*black*) or with 5 µM methylphenidate (*MPH, grey*) in

CPu (C, $n = 5$) or NAc (G, $n = 3$). **(D, J)** Mean P2/P1 (± SEM) vs IPI in control conditions (*black*) and with MPH

(*green*) in CPu (D) or NAc (J). **(E)** Mean profiles of $[DA]_o$ transients (± SEM) elicited by single or paired electrical

pulses in control conditions (*black*) or with 10 µM nomifensine (*red*) in CPu ($n = 3$). **(F)** Mean P2/P1 (± SEM) vs

IPI in control conditions (*black*) and with nomifensine (*red*) in CPu. Two-way ANOVA with Bonferroni's test for

post-hoc comparisons; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 7. The DAT limits release-dependence of STP in CPu. (A) Mean profiles of $[DA]_o$ transients (± SEM,

vertical scale normalised to 1p in 2.4 mM $[Ca^{2+}]_o$) ($n = 4$) elicited by single or paired electrical pulses in 2.4 mM

(*upper*) and 1.2 mM $[Ca^{2+}]_o$ (*lower*) in control conditions (*black*) or 5 µM cocaine (*blue*). **(B, C)** Mean P2/P1 (±

SEM) vs. IPI in 1.2 mM (*light*) and 2.4 mM $[Ca^{2+}]_o$ (*dark*) in control conditions (B) or cocaine (C). **(D)** Mean P2/P1

(± SEM) in 1.2 mM and 2.4 mM $[Ca^{2+}]_o$ in control conditions (*black-grey*) or cocaine (*blue*) at 25 ms IPI (*left*) or

40 ms IPI (*right*). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons; *$P < 0.05$, **$P < 0.01$, ***P

< 0.001.

Figure 8. The DAT and $[K^+]_o$ gate STP through overlapping mechanisms. (A, E) Mean profiles of $[DA]_o$

transients (± SEM) elicited by single or paired electrical pulses in control conditions or cocaine in 5 mM $[K^+]_o$

(*upper*) and 1 mM $[K^+]_o$ (*lower*) in CPu (A, $n = 4$) and in NAc (E, $n = 3$). **(B, F)** Mean P2/P1 (± SEM) vs. IPI in 5 mM

$[K^+]_o$ (*black*) and 1 mM $[K^+]_o$ (*grey*) in control conditions (*left*) or cocaine (*right*) in CPu (B) and in NAc (F). **(C, G)**

Mean 1p $[DA]_o$ (± SEM) in control conditions (*black-grey*) or cocaine (*blue*) in CPu (C) and in NAc (G). **(D, H)**

Mean P2/P1 (± SEM) in CPu at 25 ms IPI in 1 mM and 5 mM $[K^+]_o$ in control conditions or with cocaine in CPu

(D) and in NAc (H). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons; *$P < 0.05$, **$P < 0.01$.

Figure 1
Condon et al

Figure 2
Condon et al

Figure 3
Condon et al

Figure 4
Condon et al

Figure 5
Condon et al

Figure 6
Condon et al

Figure 7
Condon et al

Figure 8
Condon et al

Supporting Data

Figures S1-S4 and Legends

Figure S1. Voltage-gated Na$^+$ channel inhibitor lignocaine does not replicate the effects of cocaine on STP in CPu

(A) Mean profiles of [DA]$_o$ transients (± SEM) evoked by electrical stimulation in control conditions (*black*) or in the presence of 5 μM lignocaine (*grey*) in CPu (n = 3). Lignocaine significantly decreased 1p [DA]$_o$ ($p <$ 0.001, n = 3). **(B)** Mean P2/P1 ±SEM vs. IPI in control conditions (*black*) or in the presence of lignocaine (*grey*).

Figure S2. The effects of cocaine on STP are not dependent on synapsin III

(A, E) Mean profiles of [DA]$_o$ transients (± SEM) evoked by single pulses in slices from wild-type (*black*) and S3KO (*grey*) animals in CPu (A, *n* = 6) and in NAc (E, *n* = 6). **(B, F)** Mean peak [DA]$_o$ (± SEM) following single pulses in wild-type (*black*) or S3KO (*grey*) animals in CPu (B) and in NAc (F). **(C, G)** Mean profiles of [DA]$_o$ transients (± SEM) evoked by single or paired pulses in control conditions (*black*) or cocaine (*blue*) in S3KO animals in CPu (C) or in NAc (G). **(D, H)** Mean P2/P1 (± SEM) vs. IPIs in control conditions (*black*) or in cocaine (*blue*) in CPu (D) and in NAc (H). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons; *$P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Figure S3. The response to $[Ca^{2+}]_o$ revealed after DAT inhibition (see Fig. 7) is not caused by a change in the extracellular concentration of divalent cations

(A) Mean profiles of $[DA]_o$ transients (± SEM, vertical scale normalised to 1p in 2.4 mM $[Ca^{2+}]_o$) (n = 4) elicited by single or paired electrical pulses in 2.4 mM $[Ca^{2+}]_o$ (*upper*) and 1.2 mM $[Ca^{2+}]_o$ with 1.2 mM $[Mg^{2+}]_o$ (*lower*) in control conditions (*black*) or 5 µM cocaine (*blue*). **(B, C)** Mean P2/P1 (± SEM) vs. IPI in 1.2 mM (*light*) and 2.4 mM $[Ca^{2+}]_o$ (*dark*) in control conditions (B) or cocaine (C). *** $P<0.001$.

Figure S4. Cocaine prevents enhancement of STD by increased [K⁺]ₒ

(**A**) Mean profiles (± SEM) of [DA]ₒ transients in 5 mM (*upper*) and 7.5 mM [K⁺]ₒ (*lower*) in control conditions (*black-grey*) and in cocaine (*blue*). (**B**) Mean P2/P1 (± SEM) at IPIs of 10 ms and 100 ms in control conditions (*left*) and in cocaine (*right*) in 5 mM (*dark*) and 7.5 mM (*light*) [K⁺]ₒ. (**C**) Mean P2/P1 (± SEM) in CPu at 10 ms IPI (*left*) and 100 ms IPI (*right*) in 5 mM and 7.5 mM [K⁺]ₒ in control conditions (*black-grey*) or with cocaine (*blue*). Two-way ANOVA with Bonferroni's test for *post-hoc* comparisons; * $P < 0.05$, *** $P < 0.001$.